Genomic changes are varied across congeneric species pairs

Francis, Warren R.[1]; Vargas, Sergio[1]; Wörheide, Gert[1,2,3,*]

[1] Department of Earth and Environmental Sciences, Paleontology and Geobiology, Ludwig-Maximilians-Universität München, Munich, Germany.
[2] GeoBio-Center, Ludwig-Maximilians-Universität München, Munich, Germany
[3] Staatliche Naturwissenschaftliche Sammlungen Bayerns (SNSB)–Bayerische Staatssammlung für Paläontologie und Geologie, Munich, Germany

* Corresponding author: woerheide@lmu.de

Synteny, the shared arrangement of genes on chromosomes between related species, is a marker of shared ancestry, and synteny-breaking events can result in genomic incompatibilities between populations and ultimately lead to speciation events. Despite its pivotal role as a driver of speciation, the role of synteny breaks on speciation is poorly studied due to a lack of chromosome-level genome assemblies for a taxonomically broad sample of organisms. Here, using 22 congeneric animal genome pairs, we find a link between protein identity, microsynteny, and macrosynteny, but no evidence for a universal path of genomic change during divergence. We observed varied trajectories of synteny conservation relative to protein identity in non-model organisms, with many species pairs showing no karyotypic changes and others displaying large genomic rearrangements. This contrasts with previous studies on model organisms and indicates that the genomic changes preceding or resulting from speciation are likely very contextual between clades.

Introduction

The "species" is a fundamental concept in biology, and many different concepts – at least 26 – exist of what constitutes a species and how species should be defined and delineated (De Queiroz 2007; Hausdorf 2011; Zachos 2016). Delimiting species and correctly assessing species numbers in an ecosystem is essential to assess, for example, the resilience of species assemblages and ecosystems to environmental challenges, with consequences for biodiversity conservation and management (Hey et al. 2003; Frankham et al. 2012). Probably the most important species concept proposed is the "biological species" of Ernst Mayr (Mayr 1942). Mayr defined species as "groups of actually or potentially interbreeding natural populations, which are reproductively isolated from other such groups". This condition is only testable for (a few) species that can be bred under controlled conditions, for example, domesticated species or organisms that can be easily kept in the laboratory (rodents, flowers, etc.). However, it may be very difficult or impossible to directly test for most organisms, especially marine species. Hence, other "more practical" species concepts, such as the phylogenetic species concept, the evolutionary species concept, and the morphological species concept, are frequently used (reviewed in Ghiselin 2010; Frankham et al. 2012).

Under the biological species concept, reproductive isolation is key. This is because even if the exact mechanisms leading to the reproductive isolation of two or more populations remain unknown, it is predicted that reproductively isolated populations will become genetically incompatible with time through the accumulation of random mutations (i.e., Dobzhansky-Müller incompatibilities) in their genomes (Orr 1996). Importantly, this process can occur if the populations are under selection or can proceed under pure genetic drift. Thus, the accumulation of Dobzhansky-Müller incompatibilities provides a universal mechanism whereby species could arise in nature. For instance, genetic incompatibilities leading to hybrid inviability have been detected in mice (Zechner et al. 1996; Vrana et al. 2000), and in shrimp sperm/eggs recognition incompatibility has been described (Ulate and Alfaro-Montoya 2010). However, the genomic causes leading to genetic incompatibility and, eventually, hybrid inviability remain to be assessed for most species.

Although the Dobzahnsky-Müller model typically considers mutations as the causal agent behind genetic incompatibilities, synteny-breaking genome rearrangements could have similar effects on hybrid viability. In this regard, changes in both macro-synteny (i.e., the shared gene content on a chromosome, Figure 1A) after chromosomal translocations, for instance, and micro-synteny (i.e., the preserved order of several genes in a chromosome, Figure 1B) can lead to chromosome pairing problems during mitosis in hybrid zygotes, resulting in hybrid inviability and, eventually, in speciation. Several studies have observed that macro-synteny is well-conserved across animal phyla (Srivastava et al. 2008; Lv et al. 2011; Kenny et al. 2020; Simakov et al. 2022) suggesting that

chromosome gene content is constrained and that translocations are either infrequent or selected against across long time scales. One proposed mechanism underlying this constraint is that gene dosage of multiple genes on a chromosome must be maintained (Lv et al. 2011). Since chromosome translocations may result in a zygote/cell with either 1x or 3x copies of a gene, it is likely that the loss of viability resulting from these chromosome rearrangements is selected negatively, as proposed by Ohno (1970). Thus, macro-synteny breaks likely play a minor role in speciation as their effects are too deleterious to spread in a population.

In contrast, changes in micro-synteny resulting from chromosome fusions and inversions appear to be more widespread, possibly because they do not result in gene dosage imbalances. This is expected to some extent because eukaryotic genes typically have their own promoters so that genes "moving" together with their linked regulatory regions should preserve their expression patterns. Supporting this idea, the expression of nested genes appears to be uncorrelated, and gene proximity and dosage typically are also uncorrelated (Assis et al. 2008). There are some exceptions: genes controlled by bidirectional promotors or highly-conserved non-coding elements (HCNEs) controlling a group of neighbouring genes (Engström et al. 2007; Kikuta et al. 2007); or genes with regulatory elements inside the introns of neighbouring genes (e.g., the human lactase, which enhancer is located inside its upstream gene mcmc6 (Enattah et al. 2002)). Thus, it is likely that micro-synteny changes are either neutral or only slightly deleterious for most eukaryotic genes allowing these changes to spread in populations after their appearance and eventually become fixed (Ohno 1970; Slijepcevic 1998).

Despite the potential role of micro-synteny in speciation, the degree of micro-synteny is variable within the animal lineages examined to date. For instance, extensive syntenic blocks, of tens to hundreds of genes, have been found in octocorals (Jiang et al. 2019) but not medusozoans (Jiang et al. 2019; Khalturin et al. 2019), or in glass sponges (Francis et al. 2023) but not demosponges (Kenny et al. 2020). Even in two smaller phyla, Placozoa and Ctenophora, microsynteny was extensive in placozoans (Eitel et al. 2018), but was almost completely absent in ctenophores (Schultz et al. 2021). The effects of conserved synteny have only been examined in model organisms and their close relatives, making it difficult to generalise the role of micro-synteny breaking in speciation across the animal kingdom. To evaluate this, we examined macro- and micro-synteny in 22 con-generic pairs with chromosome-level genome assemblies representing six animal phyla. We find varied trajectories of synteny conservation, suggesting that no universal path of genomic change during speciation exists in animals.

Material & Methods
Datasets from NCBI

A total of 17 animal genera accounting for 22 pairs of congeneric species were downloaded from NCBI (see Table 1, and Supplemental Figure 1). The bulk of the 55 available congeneric genome pairs on NCBI mostly consisted of arthropods (e.g., 18 *Drosophila* species) and chordates (mostly mammals, e.g., cow, pig, mouse). We limited this selection to include only chromosome-level genome assemblies for which NCBI annotations for that specific version are also available and excluded redundant pairs within genera from the analyses e.g. we only kept 4 *Drosophila* species (see Table 1). For additional comparisons to remake the analyses of previous studies, we also included a set of seven genomes of additional mammal genera, namely human, mouse, chimpanzee, gorilla, orangutan, gibbon, and opossum.

Addition of representatives from a coral genus

For the coral genus *Acropora*, both *A. hyacinthus* and *A. millepora* had chromosome-level assemblies, but only *A. millepora* had an annotation file on NCBI. Thus, we used the annotation file from the previous version of *A. hyacinthus*, downloaded from the Marine Genomics Group at OIST (https://marinegenomics.oist.jp/ahya/viewer/download?project_id=91).
From the v1 annotation (Shinzato et al. 2021), we generated transcripts using gffread v0.12.7 (Pertea and Pertea 2020), which were then filtered to remove splice variants, and mapped to the chromosome-level assembly using minimap2 v2.23 (Li 2018).

Genus	N spp.	Representing taxonomic group	Phylum	Genome size (Mb)	N Chrs
Acropora	2	Staghorn corals	Cnidaria	446-475	14
Anastrepha	2	Fruit flies	Arthropoda	820-836	6
Bombus	2	Bumble-bees	Arthropoda	254-392	18 (24[a])
Bufo	2	Toads	Chordata	4545-5044	11
Cervus	2	Deer	Chordata	2526-2886	34
Crassostrea	3	Oysters	Mollusca	624-684	10
Culex	2	Mosquitoes	Arthropoda	566-573	3
Daphnia	2	Water fleas	Arthropoda	133-161	10,12
Drosophila	4	Fruit flies	Arthropoda	143-191	4-7
Epinephelus	3	Groupers (fish)	Chordata	1030-1087	24
Lytechinus	2	Sea urchins	Echinodermata	869-999	19
Mauremys	2	Turtles	Chordata	2367-2484	27
Octopus	2	Octopus	Mollusca	2342-2719	30

Genus	N spp.	Representing taxonomic group	Phylum	Genome size (Mb)	N Chrs
Oreochromis	2	Tilapia (fish)	Chordata	1005	22
Perca	2	Perch (fish)	Chordata	877-951	24
Thunnus	2	Tuna (fish)	Chordata	782-792	24
Vespa	2	Hornets	Arthropoda	193-229	25 (26 [b])

1
Table 1: The 17 genera used in analysis.
[a] several unassigned scaffolds in both *Bombus* species appear roughly
the size of chromosomes and would account for an extra 4-6 chromo-
somes.
[b] 6Mb of unassigned scaffolds are given as a single scaffold in *V. crabro*,
with mostly masked bases and effectively no annotated genes, hence
no synteny was detected with *V. velutina*.

9
Removal of additional protein isoforms

Because the NCBI annotation pipeline will include splice variants where
evidence is available, several pre-processing steps were applied for each spe-
cies to avoid isoforms from interfering with nearly all steps of the analysis
(e.g., counting duplicate orthologs). For our analyses, only the longest iso-
form was retained for each gene using a custom Python script "get_gen-
bank_longest_isoforms.py". This script simultaneously removes the shorter
protein isoforms for each gene and the annotation entries from the respec-
tive GFF files. It should be noted that the longest isoform is not necessarily
the "primary" or "correct" one, however, the role of splice variation in evo-
lution or phylogenetics is not well-studied, and the main focus of our anal-
yses was the maintenance or break of synteny blocks that can be identified
with any of the alternative isoforms. The filtered versions of the genomes
were used for all subsequent analyses, including the microsynteny and pro-
tein sequence identity. Scripts are available online at
https://github.com/PalMuc/congeneric_synteny/tree/main.

26
Generation of ortholog pairs

For each species, homologous protein clusters were generated using the fol-
lowing strategy. Proteins were aligned using DIAMOND v2.0.13 (Buchfink
et al. 2015) with default parameters. Ortholog pairs were generated using
the custom script `makehomologs.py`, which first filters the DIAMOND hits
by removing self-matches, removing matches with a low bitscore to length
ratio, and finally leverages `MCL` for the actual clustering (Enright et al.
2002).
Several filtering steps were applied (see Supplemental Figure 1), and only
one-to-one protein clusters were retained for subsequent analysis (i.e., pa-
ralogs were removed). This means that fragmented proteins (where one
gene is fragmented in either genome and results in two proteins that align
to a single protein in the other genome) were excluded. This also means
that genes duplicated in either genome (1-to-many and many-to-many
orthologs) are also excluded, even if they were identical. It should be noted
that this method does not directly identify pairs by synteny. That is, clus-
tered proteins or identical proteins may still be omitted if that cluster is not
single-copy in both genomes, even if the synteny potentially could be de-
tected by a stricter approach. For instance, the three copies of human cal-
modulin are 100% identical to each other and to the four mouse homologs,
but this would result in a cluster of seven proteins, and then get excluded
from the analysis regardless of their chromosomal arrangement. Each
ortholog pair was then subsequently aligned with MAFFT v7.487 (Katoh
and Standley 2013) to account for the entire protein in calculations of pro-
tein identity. Amino acid sites were defined as identical, different, or gaps
for each species using the custom Python script "alignment_con-
served_site_to_dots.py". Protein identity was calculated as the number of
identities divided by the number of identities plus differences, meaning that

gap sites were ignored. The frequency of gaps by species is examined in
Supplemental Figure 5 but does not appear to cause any obvious bias.
57
Quantification of microsynteny in genome pairs

We then quantified the number of genes that could be found in syntenic
blocks between each pair of congeneric species using the custom Python
script "microsynteny.py", with various options depending on the species
pair (see https://github.com/wrf/genomeGTFtools/blob/master/microsyn-
teny.py). Here, we define microsynteny as blocks of three or more consec-
utive genes in the same order in both species, which are then assumed to
be homologous, or "identical by descent".
As intergenic length is linearly related to total genome size (Francis and
Wörheide 2017), programs measuring microsynteny are very sensitive to
any parameters controlling the allowed distance to the next gene. We cali-
brated the allowed distance to the next gene across several species, finding
the optimal length of ca. 1/5000 of the total length of the genome (Supple-
mental Figure 2). For instance, for the tuna, this length accounted for 99.4%
of intergenic distances (Supplemental Figure 3), meaning that a value was
used to account for nearly all genes. Identifying syntenic blocks can also be
affected by gene insertions or deletions, so the minimum block length and
number of "intervening genes" (i.e., those inserted/deleted) were also eval-
uated. Based on this, we required a minimum of three genes per block and
allowed up to four intervening genes (Supplemental Figure 4). When the
minimum was set to two genes per block, as was used by Zdobnov and Bork
(2007), the results were not appreciably different. Since spurious synteny
blocks were frequent in randomized datasets when the minimum number
of genes per block was set to two (Supplemental Figure 4 A, B) and were
greatly reduced (10-20 fold reduction) if the minimum number of syntenic
genes was set to three (Supplemental Figure 4 A) we used three genes per
block as a conservative approach to synteny determination.

85
Quantification of macrosynteny

For macrosynteny, we counted the number of genes that could be found on
homologous chromosomes between each pair of congeneric species using
custom scripts. All proteins were aligned using DIAMOND v2.0.13
(Buchfink et al. 2015), and the resulting alignments were used to generate
dot plots with the Python script "scaffold_synteny.py". Macrosynteny was
then determined using the R script "calculate_main_block_genes.R" by
counting the number of genes for each scaffold that have DIAMOND
matches to the most common scaffold in the other species; these genes were
counted as "main" or "macrosyntenic" genes. All genes matching any other
scaffold were classified as "non-macrosyntenic". To remove genes from un-
assigned scaffolds, we only considered scaffolds greater than 1% of the total
assembly size. All code is available at https://github.com/PalMuc/conge-
neric_synteny/tree/main

Results and Discussion
Microsynteny broadly correlates with protein identity
across animals

We found a loose correlation between pairwise protein
identity and the number of genes in syntenic blocks, with
most species pairs differing in synteny but showing rela-
tively high sequence identity (Figure 1C). For instance, the
fraction of syntenic genes in the coral genus *Acropora* was
50.2%, but its average protein identity was 95.8%. The re-
duced synteny at high protein identity levels indicates that
in many species, microsynteny is not as selectively con-
strained as amino acid sequence.

On the other hand, in some genera (notably, *Drosophila*
melanogaster against *D. grimshawi*) pairs show low protein
identity but are relatively conserved gene order. In this case,
sequence identity decreases across the genus without sub-
stantial synteny breaks, indicating that synteny may be con-
strained. Because insertion of new genes (duplications,

(retro)transpositions) or deletions would also disrupt synteny, these processes may also be uncommon in *Drosophila.*

To compare to analyses from other studies, we added two sets of species pairs, that included species not in the same genus, one from insects (gray) and one from mammals (yellow-green) (Figure 1C). While there is not any obvious connection between insects and mammals, these additional sets showed a different pattern of protein sequence identity and microsynteny. This suggests a different genomic history, whereby two events occur: one is the overall complete loss of synteny; the other is a decreasing protein identity, such that orthologs can no longer be identified by BLAST (see Supplemental Figure 6). This may be accompanied further by an overall decrease in the total number of ortholog pairs, because our pipeline excludes recently duplicated genes. In general, this agrees with earlier findings in insects (Zdobnov and Bork 2007), though that study was focused on insect species between orders.

Microsynteny changes more frequently than macrosynteny

Given that many inversions (of three or more genes) do not disrupt microsynteny, many inverted blocks would still be detected as syntenic, and the slow accumulation of

inversions would not change the fraction of genes in blocks or affect the overall microsynteny estimates. Translocations, however, can change macrosynteny but not microsynteny, as genes move as a block to a different chromosome. We examined this for each congeneric species pair as a function of the number of genes on homologous chromosomes in both species compared to the number of translocated genes or inserted novel genes.

Like the relationship with protein identity, we found that both macrosynteny and microsynteny together tend to decrease with speciation (Figure 1D), but can be decoupled in many species pairs. In most species pairs (other than the mammal set) microsynteny appears to be less constrained than macrosynteny. For instance, although microsynteny is almost absent in the insects analysed, macrosynteny remains broadly detectable in these organisms. Mammals, however, display a different pattern of chromosomal evolution, with several species pairs showing chromosomal rearrangements while preserving their microsynteny.

While there are examples of changes in karyotype between species (such as the fusion resulting in Chr 2 in humans, Yunis and Prakash 1982), only a few of the species in our study had obvious karyotype differences. Macrosynteny is relatively conserved among Drosophila species with very conserved microsynteny and relatively low protein sequence identity. Between *D. melanogaster* and *D. erecta* (the closest pair in our study), only one translocation is evident in the genome (Figure 2A). In Octopus, which possesses the largest invertebrate genome in our study, the karyotype appears to be the same between the two species analysed, but with frequent small inversions in most chromosomes (Figure 2B). Gene arrangement in Octopus is known to be variable, such as the rearrangement of the Hox cluster (Albertin et al. 2015). Other species have karyotype variations. The water flea *Daphnia pulex* has 12 chromosomes, while *D. magna* has only ten. These *Daphnia* species show conserved macrosynteny but rapidly decaying microsynteny (Figure 2C). Also, the karyotype change in *Daphnia* appears to have been caused by recent Robertsonian fusion events in *D. magna*, as signs of mixing after fusion were detected. For *Crassostea* (oysters), a chromosomal fusion and a split were evident (Figure 2D), despite their similar karyotype.

Distribution of protein identity values is similar across species pairs

We examined genomic signatures of speciation across 23 pairs of congeneric species across the animal kingdom. Species are not typically defined by molecular distances, so all of the species pairs in our study, though formally described as species, may all be at different genetic stages of divergence. If all pairs show the same pattern of genetic changes, then they can be thought of as being at different stages along the same underlying process. However, if they show different patterns between species, then multiple underlying processes are possible.

Figure 1: (A) Schematic dot plot of microsynteny. Arrows along the axes represent enes, with red genes indicating inversions in one of the two genomes. Red boxes indicate microsyntenic blocks. (B) Schematic dot plot of macrosynteny. Bars along the axes represent chromosomes and points represent positions of homologous genes between the two species. Blue points would be considered macrosyntenic, red points are microsyntenic but with a translocation, and gray points are any other hits, e.g. genes sitting on very different positions in the compared species pairs. (C) Pairwise protein sequence identity for all orthologs plotted against fraction of genes in synteny blocks. Gray (insects) and yellow-green (mammals) indicate additional comparisons between species not in the same genus. The dotted line indicates the calculated regression line from Zdobnov and Bork (2007) of the same two axes. (D) Relationship between microsynteny and macrosynteny.

Figure 2: Example dotplots between congeneric species, showing a range of karyotypic states. All axes indicate positions in the genome in megabases. (A) *Drosophila*, showing two translocations/splits in blue boxes, one of which is unassigned to a chromosome in the assembly. (B) Octopus, with no obvious translocations but many small inversions. (C) *Daphnia*, red boxes indicate two chromosomal fusions with mixing in *D. magna*. (D) *Crassostrea*, with an evident chromosomal fusion and split between the two species, as red and blue boxes, respectively. See online repository data for higher resolution versions of all species pairs.

Among congeneric species, most one-to-one orthologous
proteins are nearly identical for many pairs (Figure 3A,D,
Supplemental Figure 7), and only few proteins were highly
divergent (Supplemental Figure 7). As mutations accumu-
late (Figure 3B,C,E,F), very few proteins remain completely
identical. The more divergent proteins become difficult to
detect by BLAST, effectively "dropping off the graph," and
are neither counted for protein sequence identity nor de-
tected in synteny blocks. The drop-off threshold appears to
be at roughly 40% protein identity (Figure 3, see Supple-
mental Figure 8 for histograms of all species).

Lack of genomic signatures of speciation

The taxonomic hierarchy can be considered an abstraction
of the true divergence within a clade. However, taxonomic
ranks are arbitrary and do not reflect divergence similarly in
different groups. Yet, as ranks can be used as proxies for phy-
logenetic diversity, taxonomic ranks remain meaningful in
contexts such as conservation biology (Hendricks et al.
2014). In our study, as there are multiple trajectories of ge-
netic change among congeneric species (Figure 1), we could
not identify a clear genomic distinction based on macro-
and/or microsyntenic patterns or percent protein identity

Figure 3: Histograms of protein percent identity for *Crassostrea* (A-C) and *Drosophila* (D-F). Protein identity is binned at intervals of 0.1% identity.

that could possibly be useful to delimit different animal species, and generally, any other taxonomic level. Within some clades, patterns related to specific chromosomal features, such as centromere position, may emerge. In our data, some species pairs (e.g., in genera *Crassostrea*, *Daphnia*) show low levels of microsynteny and relatively low protein sequence identity indicating that the taxonomic rank 'genus' in these pairs point to different stages of the evolutionary process (see, for example, also Avise and Liu 2011). For instance, the two Crassostrea species, *C. angulata* and *C. virginica*, represent a pair where substantial rearrangements are visible in the genome. These two species are also separated geographically, as *C. angulata* was originally from the Pacific, while *C. virginica* is from the Atlantic.

As in most other similar studies, the synteny assignments in our method are ultimately dependent upon local sequence similarity (BLAST, DIAMOND, etc). Hence, one of two effects could result in the exclusion of a gene from our analysis: duplication (no longer a one-to-one ortholog) or mutation beyond the detection of BLAST. This second cause may explain the regression line found by Zdobnov and Bork (2007) and the apparent lower bound for protein identity to microsynteny (Figure 1C). In this regard, as protein pairs diverge beyond the detection limit imposed by sequence similarity searches, e.g., in more distantly related species or genes with low evolutionary pressure, they cannot be detected as syntenic. Thus, it is very unlikely to find a species pair with very high microsynteny and low protein identity but not the opposite.

Difficulty in quantification of syntenic blocks and microsynteny

Meaningful measurements of microsynteny are challenging to get with current data. For instance, assembly quality could affect synteny assessments, as lower quality assemblies would be expected to increase the number of synteny blocks but not necessarily affect the number of syntenic genes. Due to the high sensitivity of the analysis to different parameters, real biological features such as gene insertions, duplications, or long intergenic regions can result in synteny blocks being split in two. Thus, the number of synteny blocks themselves is less important than the number of genes contained in blocks. A further confounding factor is the presence of unassigned scaffolds (i.e., the number of scaffolds is far above the number of chromosomes) in all assemblies analyzed. These may be unmerged haplotypes and would affect the measurement by increasing the number of

genes not found in matching synteny blocks as they appear as duplicates.

One attempt to model the syntenic change between mouse and human (Nadeau and Taylor 1984) had also assumed that reciprocal translocations and not inversions typically caused the synteny breaks. Under that model, it was assumed that the length of synteny blocks follows a negative exponential distribution and that synteny breaks disrupt both micro- and macro-synteny, something not typically observed. Nadeau's model was disputed by Zdobnov and Bork (2007), who argued that the distribution of block length followed a power law, not an exponential distribution. We found similar correlations in block length in our dataset, though R2 ranged between 0.713 and 0.945 (Supplemental Figure 9). Thus, the correlation between block length and number of blocks is not as strong as proposed by Zdobnov and Bork (2007), suggesting that other biological factors affect chromosomal breakpoints (Pevzner and Tesler 2003).

Challenges in quantification of rearrangement time

A prominent unanswered question of this study is whether chromosome rearrangements occur at the same rate across a clade. Such a study would be of great interest in evolutionary biology, even if difficult to design correctly. Firstly, such a study would require enough sequenced high-quality chromosome-scale species-pair genomes to identify which lineage had the inversion or translocation based on genomic triangulation. We are currently far from this objective. In our study, there were only 18 genera on NCBI with more than two species with chromosome-level assemblies, and several of those still did not include an annotation.

Secondly, one must infer divergence times between taxa in a way that is not reliant on protein identity. Divergence time estimates are often based on molecular clock inferences. While fossil calibrations are necessary for this analysis, the time estimates are then shifted based on molecular changes, i.e, the inferred divergence time depends on the protein identity. Unsurprisingly, one would likely find a strong correlation between protein identity and divergence time (as in Figure 1), but that is a problem of circular logic; the divergence time was a function of protein identity in the first place, so it cannot be applied as an independent measurement for rearrangements. This is extended with the correlation of protein identity to the number of genes in synteny blocks. Consequently, protein identity, microsynteny, and molecular clock-based divergence estimates are likely correlated. Unfortunately, this means that molecular clock divergence time should not be used to test the rearrangement rate in lineages. One possibility for timing narrow-scale speciation could be the usage of codon evolution models. For instance, in Placozoa, it was observed that the nucleotide identity differed from protein identity such that highly conserved genes had many synonymous nucleotide changes instead of simply having fewer total mutations (Eitel et al. 2018). Thus,

codon evolution models could provide a different age estimate than the proteins.

Conclusions

Defining and delimiting species is one of the important and basic requirements for studying biodiversity and its evolution, and it is pivotal for the management and conservation of natural resources. However, even in the times of DNA barcoding and genome sequencing, no universal threshold that applies, for example, across all animals, has been identified, and different species concepts are applied in different taxa. Synteny, i.e., the shared arrangement of genes on chromosomes between species, has been used as a marker of shared ancestry, with synteny-breaking events resulting in genomic incompatibilities between populations, ultimately leading to speciation. However, in our study of the genomes of 22 con-generic pairs, we found that genomic changes inferred from speciation are variable and contextual and that synteny should not be used at present as a measure of divergence or speciation. Once more high-quality, chromosome-scale genomes of species-pairs are available, a clear pattern of genome evolution may emerge.

Data availability

Additional data, figures, scripts and analytical resources can be accessed in a GitHub repository https://github.com/Pal-Muc/congeneric_synteny, archived with a DOI at Zenodo (https://zenodo.org/doi/10.5281/zenodo.10069099).

Acknowledgements

G.W. acknowledges funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 764840 (ITN IGNITE). WRF would like to thank M. Dohrmann and D. Schultz for helpful discussions. We would like to thank M. Eitel for helpful discussions and comments. G.W. dedicates this work to his late wife Connie Wörheide, who left us too early during the preparation of this manuscript.

References

Albertin CB, Simakov O, Mitros T, Wang ZY, Pungor JR, Edsinger-Gonzales E, Brenner S, Ragsdale CW, Rokhsar DS. 2015. The octopus genome and the evolution of cephalopod neural and morphological novelties. *Nature* 524:220–224.

Assis R, Kondrashov AS, Koonin EV, Kondrashov FA. 2008. Nested genes and increasing organizational complexity of metazoan genomes. *Trends Genet.* 24:475–478.

Avise JC, Liu J-X. 2011. On the temporal inconsistencies of Linnean taxonomic ranks. *Biol. J. Linn. Soc. Lond.* 102:707–714.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND. *Nat. Methods* 12:59–60.

De Queiroz K. 2007. Species concepts and species delimitation. *Syst. Biol.* 56:879–886.

Eitel M, Francis WR, Varoqueaux F, Daraspe J, Osigus H-J, Krebs S, Vargas S, Blum H, Williams GA, Schierwater B, et al. 2018. Comparative genomics and the nature of placozoan species. *PLoS Biol.* 16:e2005359.

Enattah NS, Sahi T, Savilahti E, Terwilliger JD, Peltonen L, Järvelä I. 2002. Identification of a variant associated with adult-type hypolactasia. *Nat. Genet.* 30:233–237.

Engström PG, Ho Sui SJ, Drivenes O, Becker TS, Lenhard B. 2007. Genomic regulatory blocks underlie extensive microsynteny conservation in insects. *Genome Res.* 17:1898–1908.

Enright AJ, Van Dongen S, Ouzounis CA. 2002. An efficient algorithm for large-scale detection of protein families. *Nucleic Acids Res.* 30:1575–1584.

Francis WR, Eitel M, Vargas S, Garcia-Escudero CA, Conci N, Deister F, Mah JL, Guiglielmoni N, Krebs S, Blum H, et al. 2023. The genome of the reef-building glass sponge *Aphrocallistes vastus* provides insights into silica biomineralization. *R Soc Open Sci* 10:230423.

Francis WR, Wörheide G. 2017. Similar Ratios of Introns to Intergenic Sequence across Animal Genomes. *Genome Biol. Evol.* 9:1582–1598.

Frankham R, Ballou JD, Dudash MR, Eldridge MDB, Fenster CB, Lacy RC, Mendelson JR, Porton IJ, Ralls K, Ryder OA. 2012. Implications of different species concepts for conserving biodiversity. *Biol. Conserv.* 153:25–31.

Ghiselin MT. 2010. Species Concepts. *eLS* [Internet]. Available from: https://onlinelibrary.wiley.com/doi/10.1002/9780470015902. a0001744.pub2

Hausdorf B. 2011. Progress toward a general species concept. *Evolution* 65:923–931.

Hendricks JR, Saupe EE, Myers CE, Hermsen EJ, Allmon WD. 2014. The Generification of the Fossil Record. *Paleobiology* 40:511–528.

Hey J, Waples RS, Arnold ML, Butlin RK, Harrison RG. 2003. Understanding and confronting species uncertainty in biology and conservation. *Trends Ecol. Evol.* 18:597–603.

Jiang JB, Quattrini AM, Francis WR, Ryan JF, Rodríguez E, McFadden CS. 2019. A hybrid de novo assembly of the sea pansy (*Renilla muelleri*) genome. *Gigascience* 8:giz026.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30:772–780.

Kenny NJ, Francis WR, Rivera-Vicéns RE, Juravel K, de Mendoza A, Díez-Vives C, Lister R, Bezares-Calderón LA, Grombacher L, Roller M, et al. 2020. Tracing animal genomic evolution with the chromosomal-level assembly of the freshwater sponge *Ephydatia muelleri*. *Nat. Commun.* 11:1–11.

Khalturin K, Shinzato C, Khalturina M, Hamada M, Fujie M, Koyanagi R, Kanda M, Goto H, Anton-Erxleben F, Toyokawa M, et al. 2019. Medusozoan genomes inform the evolution of the jellyfish body plan. *Nat Ecol Evol* 3:811–822.

Kikuta H, Laplante M, Navratilova P, Komisarczuk AZ, Engström PG, Fredman D, Akalin A, Caccamo M, Sealy I, Howe K, et al. 2007. Genomic regulatory blocks encompass multiple neighboring genes and maintain conserved synteny in vertebrates. *Genome Res.* 17:545–555.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100.

Lv J, Havlak P, Putnam NH. 2011. Constraints on genes shape long-term conservation of macro-synteny in metazoan genomes. *BMC Bioinformatics* 12 Suppl 9:S11.

Mayr E. 1942. Systematic and the origin of species. New York: Columbia University Press

Nadeau JH, Taylor BA. 1984. Lengths of chromosomal segments conserved since divergence of man and mouse. *Proc. Natl. Acad. Sci. U. S. A.* 81:814–818.

Ohno S. 1970. Gene Duplication as an Evolutionary Force. In: Peeters H, editor. Protides of the Biological Fluids. Vol. 17. Elsevier. p. 39–42.

Orr HA. 1996. Dobzhansky, Bateson, and the genetics of speciation. *Genetics* 144:1331–1335.

Pertea G, Pertea M. 2020. GFF Utilities: GffRead and GffCompare. *F1000Res.* 9. Available from: http://dx.doi.org/10.12688/f1000research.23297.2

Pevzner P, Tesler G. 2003. Human and mouse genomic sequences reveal extensive breakpoint reuse in mammalian evolution. *Proc. Natl. Acad. Sci. U. S. A.* 100:7672–7677.

Schultz DT, Francis WR, McBroome JD, Christianson LM, Haddock SHD, Green RE. 2021. A chromosome-scale genome assembly and karyotype of the ctenophore *Hormiphora californensis*. *G3* 11:jkab302.

Shinzato C, Khalturin K, Inoue J, Zayasu Y, Kanda M, Kawamitsu M, Yoshioka Y, Yamashita H, Suzuki G, Satoh N. 2021. Eighteen coral genomes reveal the evolutionary origin of *Acropora* strategies to accommodate environmental changes. *Mol. Biol. Evol.* 38:16–30.

Simakov O, Bredeson J, Berkoff K, Marletaz F, Mitros T, Schultz DT, O'Connell BL, Dear P, Martinez DE, Steele RE, et al. 2022. Deeply conserved synteny and the evolution of metazoan chromosomes. *Sci Adv* 8:eabi5884.

Slijepcevic P. 1998. Telomeres and mechanisms of Robertsonian fusion. *Chromosoma* 107:136–140.

Srivastava M, Begovic E, Chapman J, Putnam NH, Hellsten U, Kawashima T, Kuo A, Mitros T, Salamov A, Carpenter ML, et al. 2008. The *Trichoplax* genome and the nature of placozoans. *Nature* 454:955–960.

Ulate K, Alfaro-Montoya J. 2010. Sperm–egg incompatibility in inter-specific inseminations of *Penaeus (Litopenaeus) occidentalis*, *P. (Litopenaeus) stylirostris* and *P. (Litopenaeus) vannamei*. *Aquaculture* 309:290–292.

Vrana PB, Fossella JA, Matteson P, del Rio T, O'Neill MJ, Tilghman SM. 2000. Genetic and epigenetic incompatibilities underlie hybrid dysgenesis in *Peromyscus*. *Nat. Genet.* 25:120–124.

Yunis JJ, Prakash O. 1982. The origin of man: a chromosomal pictorial legacy. *Science* 215:1525–1530.

Zachos FE. 2016. Species Concepts in Biology. Springer International Publishing.

Zdobnov EM, Bork P. 2007. Quantification of insect genome divergence. *Trends Genet.* 23:16–20.

Zechner U, Reule M, Orth A, Bonhomme F, Strack B, Guénet, Hameister H, Fundele R. 1996. An X-chromosome linked locus contributes to abnormal placental development in mouse interspecific hybrid. *Nat. Genet.* 12:398–403.