Genomic changes are varied across congeneric species pairs of animals

Francis, Warren R.[1]; Vargas, Sergio[1]; Wörheide, Gert[1,2,3,*]

[1] Department of Earth and Environmental Sciences, Paleontology and Geobiology, Ludwig-Maximilians-

Universität München, Munich, Germany.

[2] GeoBio-Center, Ludwig-Maximilians-Universität München, Munich, Germany

[3] Staatliche Naturwissenschaftliche Sammlungen Bayerns (SNSB)–Bayerische Staatssammlung für

Paläontologie und Geologie, Munich, Germany

* Corresponding author: woerheide@lmu.de

Abstract

Synteny, the shared arrangement of genes on chromosomes between related species, is a marker of

shared ancestry, and synteny-breaking events can result in genomic incompatibilities between

populations and ultimately lead to speciation events. Despite its pivotal role as a driver of speciation,

the role of synteny breaks on speciation is poorly studied due to a lack of chromosome-level genome

assemblies for a taxonomically broad sample of organisms. Here, using 22 congeneric animal genome

pairs, we find a link between protein identity, microsynteny, and macrosynteny, but no evidence for a

universal path of genomic change during divergence. We observed varied trajectories of synteny

conservation relative to protein identity in non-model organisms, with many species pairs showing no

karyotypic changes and others displaying large genomic rearrangements. This contrasts with previous

studies on model organisms and indicates that the genomic changes preceding or resulting from

speciation are likely very contextual between clades.

Introduction

The "species" is a fundamental concept in biology, and many different concepts – at least 26 – exist of

what constitutes a species and how species should be defined and delineated (De Queiroz, 2007;

Hausdorf, 2011; Zachos, 2016). Delimiting species and correctly assessing species numbers in an

ecosystem is essential to assess, for example, the resilience of species assemblages and ecosystems to

environmental challenges, with consequences for biodiversity conservation and management

(Frankham et al., 2012; Hey et al., 2003). Probably the most important species concept proposed is the

"biological species" of Ernst Mayr (Mayr, 1942). Mayr defined species as "*groups of actually or*

potentially interbreeding natural populations, which are reproductively isolated from other such
groups". This condition is only testable for (a few) species that can be bred under controlled
conditions, for example, domesticated species or organisms that can be easily kept in the laboratory
(rodents, flowers, etc.). However, it may be very difficult or impossible to directly test for most
organisms, especially marine species. Hence, other "more practical" species concepts, such as the
phylogenetic species concept, the evolutionary species concept, and the morphological species
concept, are frequently used (Frankham et al., 2012; reviewed in Ghiselin, 2010).

Under the biological species concept, reproductive isolation is key. This is because even if the exact
mechanisms leading to the reproductive isolation of two or more populations remain unknown, it is
predicted that reproductively isolated populations will become genetically incompatible with time
through the accumulation of random mutations (i.e., Dobzhansky-Müller incompatibilities) in their
genomes (Orr, 1996). Importantly, this process can occur if the populations are under selection or can
proceed under pure genetic drift. Thus, the accumulation of Dobzhansky-Müller incompatibilities
provides a universal mechanism whereby species could arise in nature. For instance, genetic
incompatibilities leading to hybrid inviability have been detected in mice (Vrana et al., 2000; Zechner
et al., 1996), and in shrimp sperm/eggs recognition incompatibility has been described (Ulate &
Alfaro-Montoya, 2010). However, the genomic causes leading to genetic incompatibility and,
eventually, hybrid inviability remain to be assessed for most species.

Although the Dobzahnsky-Müller model typically considers mutations as the causal agent behind
genetic incompatibilities, synteny-breaking genome rearrangements could have similar effects on
hybrid viability. In this regard, changes in both macro-synteny (i.e., the shared gene content on a
chromosome, Figure 1A) after chromosomal translocations, for instance, and micro-synteny (i.e., the
preserved order of several genes in a chromosome, Figure 1B) can lead to chromosome pairing
problems during mitosis in hybrid zygotes, resulting in hybrid inviability and, eventually, in
speciation. Several studies have observed that macro-synteny is well-conserved across animal phyla
(Kenny et al., 2020; Lv et al., 2011; Simakov et al., 2022; Srivastava et al., 2008) suggesting that
chromosome gene content is constrained and that translocations are either infrequent or selected
against across long time scales. One proposed mechanism underlying this constraint is that gene
dosage of multiple genes on a chromosome must be maintained (Lv et al., 2011). Since chromosome
translocations may result in a zygote/cell with either 1x or 3x copies of a gene, it is likely that the loss
of viability resulting from these chromosome rearrangements is selected negatively, as proposed by
Ohno (1970). Thus, translocations may only play a minor role in speciation as their effects are too
deleterious to spread in a population while other macro-synteny changes, such as Robertsonian
fusions, may still have effects on speciation.

In contrast, changes in micro-synteny resulting from chromosome fusions and inversions appear to be
more widespread, possibly because they do not result in gene dosage imbalances. This is expected to
some extent because eukaryotic genes typically have their own promoters so that genes "moving"
together with their linked regulatory regions should preserve their expression patterns. Supporting this
idea, the expression of nested genes appears to be uncorrelated, and gene proximity and dosage
typically are also uncorrelated (Assis et al., 2008). There are some exceptions: genes controlled by
bidirectional promotors or highly-conserved non-coding elements (HCNEs) controlling a group of
neighbouring genes (Engström et al., 2007; Kikuta et al., 2007); or genes with regulatory elements
inside the introns of neighbouring genes (e.g., the human lactase, which enhancer is located inside its
upstream gene *mcmc6* (Enattah et al., 2002)). Thus, it is likely that micro-synteny changes are either
neutral or only slightly deleterious for most eukaryotic genes allowing these changes to spread in
populations after their appearance and eventually become fixed (Ohno, 1970; Slijepcevic, 1998).
13

Despite the potential role of micro-synteny in speciation, the degree of micro-synteny is variable
within the animal lineages examined to date. For instance, extensive syntenic blocks, of tens to
hundreds of genes, have been found in octocorals (Jiang et al., 2019) but not medusozoans (Jiang et
al., 2019; Khalturin et al., 2019), or in glass sponges (Francis et al., 2023) but not demosponges
(Kenny et al., 2020). Even in two smaller phyla, Placozoa and Ctenophora, microsynteny was
extensive in placozoans (Eitel et al., 2018), but was almost completely absent in ctenophores (Schultz
et al., 2021). The effects of conserved synteny have only been examined in model organisms and their
close relatives, making it difficult to generalise the role of micro-synteny breaking in speciation across
the animal kingdom. To evaluate this, we examined macro- and micro-synteny in 22 con-generic pairs
with chromosome-level genome assemblies representing five animal phyla. We find varied
trajectories of synteny conservation, suggesting that no universal path of genomic change during
speciation exists in animals.

Material & Methods

Datasets from NCBI

A total of 17 animal genera accounting for 22 pairs of congeneric species were downloaded from
NCBI (see Table 1, and Supplemental Figure 1). The bulk of the 55 available congeneric genome
pairs on NCBI mostly consisted of arthropods (e.g., 18 *Drosophila* species) and chordates (mostly
mammals, e.g., cow, pig, mouse). We limited this selection to include only chromosome-level
genome assemblies for which NCBI annotations for that specific version were also available, and also
excluded redundant pairs within genera from the analyses e.g. we only kept 4 *Drosophila* species (see
Table 1). In one of the genera, *Crassostrea*, two of the species were reassigned into another genus

Magallana (Salvi & Mariottini, 2017) on NCBI during the review of this manuscript, though the

original comparisons are retained below. For additional comparisons to remake the analyses of

previous studies, we analyzed the insects against each other, using *Drosophila melanogaster* as one

species, and then one of *Anastrepha*, *Culex*, *Bombus*, and *Vespa* for each pair (matching what was

done in (Zdobnov & Bork, 2007). We also included a set of seven genomes of additional mammal

genera, namely human, *Mus musculus* (mouse), *Pan troglodytes* (chimpanzee), *Gorilla gorilla*

(gorilla), *Pongo abelii* (orangutan), *Nomascus leucogenys* (gibbon), and *Monodelphis domestica*

(opossum).

9   Addition of representatives from a coral genus

For the coral genus *Acropora*, both *A. hyacinthus* and *A. millepora* had chromosome-level assemblies,

but only *A. millepora* had an annotation file on NCBI. Thus, we used the annotation file from the

previous version of *A. hyacinthus*, downloaded from the Marine Genomics Group at OIST

(https://marinegenomics.oist.jp/ahya/viewer/download?project_id=91). From the v1 annotation

(Shinzato et al., 2021), we generated transcripts using gffread v0.12.7 (Pertea & Pertea, 2020), which

were then filtered to remove splice variants, and mapped to the chromosome-level assembly using

minimap2 v2.23 (Li, 2018).

Genus	N spp.	Representing taxonomic group	Phylum	Genome size (Mb)	N Chrs
Acropora	2	Staghorn corals	Cnidaria	446-475	14
Anastrepha	2	Fruit flies	Arthropoda	820-836	6
Bombus	2	Bumblebees	Arthropoda	254-392	18 (24 [a])
Bufo	2	Toads	Chordata	4545-5044	11
Cervus	2	Deer	Chordata	2526-2886	34
Crassostrea/ Magallana ([b])	3	Oysters	Mollusca	624-684	10
Culex	2	Mosquitoes	Arthropoda	566-573	3
Daphnia	2	Water fleas	Arthropoda	133-161	10,12
Drosophila	4	Fruit flies	Arthropoda	143-191	4-7
Epinephelus	3	Groupers (fish)	Chordata	1030-1087	24
Lytechinus	2	Sea urchins	Echinodermata	869-999	19
Mauremys	2	Turtles	Chordata	2367-2484	27
Octopus	2	Octopus	Mollusca	2342-2719	30
Oreochromis	2	Tilapia (fish)	Chordata	1005	22
Perca	2	Perch (fish)	Chordata	877-951	24
Thunnus	2	Tuna (fish)	Chordata	782-792	24
Vespa	2	Hornets	Arthropoda	193-229	25 (26 [c])

Table 1: The 17 genera used in analysis.

^a several unassigned scaffolds in both *Bombus* species appear roughly the size of chromosomes, and would
account for an extra 4-6 chromosomes.

b at the time of downloading the data from NCBI, all species used were of the genus *Crassostrea*, though now
some species in this genus have been reassigned to *Magellana* (see Salvi & Mariottini, 2017, 2021)

^c 6Mb of unassigned scaffolds are given as a single scaffold in *V. crabro*, with mostly masked bases and
effectively no annotated genes, hence no synteny was detected with *V. velutina*.

8 Removal of additional protein isoforms

Because the NCBI annotation pipeline will include splice variants where evidence is available, several

pre-processing steps were applied for each species to avoid isoforms from interfering with nearly all

steps of the analysis (e.g., counting duplicate orthologs). For our analyses, only the longest isoform

was retained for each gene using a custom Python script "get_genbank_longest_isoforms.py". This

script simultaneously removes the shorter protein isoforms for each gene and the annotation entries

from the respective GFF files. It should be noted that the longest isoform is not necessarily the

"primary" or "correct" one, however, the role of splice variation in evolution or phylogenetics is not

well-studied, and the main focus of our analyses was the maintenance or break of synteny blocks that

can be identified with any of the alternative isoforms. The filtered versions of the genomes were used

for all subsequent analyses, including the microsynteny and protein sequence identity. Scripts are

available online at https://github.com/PalMuc/congeneric_synteny/tree/main.

20 Generation of ortholog pairs

For each species pair, homologous protein clusters were generated using the following strategy.

Proteins were aligned using DIAMOND v2.0.13 (Buchfink et al., 2015) with default parameters.

Ortholog pairs were generated using the custom script `makehomologs.py`, which first filters the

DIAMOND hits by removing self-matches, removing matches with a low bitscore to length ratio, and

finally leverages `MCL` for the actual clustering (Enright et al., 2002).

Several filtering steps were applied (see Supplemental Figure 1), and only one-to-one protein clusters

were retained for subsequent analysis (i.e., paralogs were removed). This means that fragmented

proteins (where one gene is fragmented in either genome and results in two proteins that align to a

single protein in the other genome) were excluded. This also means that genes duplicated in either

genome (1-to-many and many-to-many orthologs) are also excluded, even if they were identical. It

should be noted that this method does not directly identify pairs by synteny. That is, clustered proteins

or identical proteins may still be omitted if that cluster is not single-copy in both genomes, even if the

synteny potentially could be detected by a stricter approach. For instance, the three copies of human

calmodulin are 100% identical to each other and to the four mouse homologs, but this would result in

a cluster of seven proteins, and then get excluded from the analysis regardless of their chromosomal

arrangement. Each ortholog pair was then subsequently aligned with MAFFT v7.487 (Katoh &

Standley, 2013) to account for the entire protein in calculations of protein identity. Amino acid sites

were defined as identical, different, or gaps for each species using the custom Python script

"alignment_conserved_site_to_dots.py". Protein identity was calculated as the number of identities

divided by the number of identities plus differences, meaning that gap sites were ignored. The

frequency of gaps by species is examined in Supplemental Figure 2, but does not appear to cause any

obvious bias, such as if highly-divergent regions were systematically missed in many protein pairs.

Quantification of microsynteny in genome pairs

We then quantified the number of genes that could be found in syntenic blocks between each pair of

congeneric species using the custom Python script "microsynteny.py", with various options depending

on the species pair (see https://github.com/wrf/genomeGTFtools/blob/master/microsynteny.py). Here,

we define microsynteny as blocks of three or more consecutive genes in the same order in both

species, which are then assumed to be homologous, or "identical by descent".

As intergenic length is linearly related to total genome size (Francis & Wörheide, 2017), programs

measuring microsynteny are very sensitive to any parameters controlling the allowed distance to the

next gene. We calibrated the allowed distance to the next gene across several species, finding the

optimal length of ca. 1/5000 of the total length of the genome (Supplemental Figure 3). For instance,

for the tuna, this length accounted for 99.4% of intergenic distances (Supplemental Figure 4),

meaning that a value was used to account for nearly all genes. Identifying syntenic blocks can also be

affected by gene insertions or deletions, so the minimum block length and number of "intervening

genes" (i.e., those inserted/deleted) were also evaluated. Based on this, we required a minimum of

three genes per block and allowed up to four intervening genes (Supplemental Figure 5). When the

minimum was set to two genes per block, as was used by (Zdobnov & Bork, 2007), the results were

not appreciably different. Since spurious synteny blocks were frequent in randomised datasets when

the minimum number of genes per block was set to two (Supplemental Figure 5 A, B) and were

greatly reduced (10-20 fold reduction) if the minimum number of syntenic genes was set to three

(Supplemental Figure 5 A) we used three genes per block as a conservative approach to synteny

determination.

Quantification of macrosynteny

31 For macrosynteny, we counted the number of genes that could be found on homologous chromosomes

32 between each pair of congeneric species using custom scripts. All proteins were aligned using

33 DIAMOND v2.0.13 (Buchfink et al., 2015), and the resulting alignments were used to generate dot

34 plots with the Python script "scaffold_synteny.py". Macrosynteny was then determined using the R

script "calculate_main_block_genes.R" by counting the number of genes for each scaffold that have DIAMOND matches to the most common scaffold in the other species; these genes were counted as "main" or "macrosyntenic" genes. All genes matching any other scaffold were classified as "non-macrosyntenic". To remove genes from unassigned scaffolds, we only considered scaffolds greater than 1% of the total assembly size. All code is available at:

https://github.com/PalMuc/congeneric_synteny/tree/main

Results and Discussion

Microsynteny broadly correlates with protein identity across animals

We found a loose correlation between pairwise protein identity and the number of genes in syntenic blocks, with most species pairs differing in synteny but showing relatively high sequence identity (Figure 1C). For instance, the fraction of syntenic genes in the coral genus *Acropora* was 50.2%, but its average protein identity was 95.8%. The reduced synteny at high protein identity levels indicates that in many species, microsynteny is not as selectively constrained as amino acid sequence.

On the other hand, in some genera (notably, *Drosophila melanogaster* against *D. grimshawi*) pairs show low protein identity but are relatively conserved gene order. In this case, sequence identity decreases across the genus without substantial synteny breaks, indicating that synteny may be constrained. Because insertion of new genes (duplications, (retro)transpositions) or deletions would also disrupt synteny, these processes may also be uncommon in *Drosophila*.

Figure 1: (A) Schematic dot plot of microsynteny. Arrows along the axes represent genes, with red
genes indicating inversions in one of the two genomes. Red boxes indicate microsyntenic blocks. (B)
Schematic dot plot of macrosynteny. Bars along the axes represent chromosomes and points
represent positions of homologous genes between the two species. Blue points would be considered
macrosyntenic, red points are microsyntenic but with a translocation, and gray points are any other
hits, e.g. genes sitting on very different positions in the compared species pairs. (C) Pairwise protein
sequence identity for all orthologs plotted against fraction of genes in synteny blocks. Gray (insects)
and yellow-green (mammals) indicate additional comparisons between species not in the same
genus. The dotted line indicates the calculated regression line from (Zdobnov & Bork, 2007) of the
same two axes. (D) Relationship between microsynteny and macrosynteny.

To compare to analyses from other studies, we added two sets of species pairs, that included species

not in the same genus, one from insects (gray) and one from mammals (yellow-green) (Figure 1C).

While there is not any obvious connection between insects and mammals, these additional sets

showed a different pattern of protein sequence identity and microsynteny. This suggests a different

genomic history, whereby two events occur: one is the overall complete loss of synteny; the other is a

decreasing protein identity, such that orthologs can no longer be identified by BLAST (see

Supplemental Figure 6). This may be accompanied further by an overall decrease in the total number

1 of ortholog pairs, because our pipeline excludes recently duplicated genes. In general, this agrees with

2 earlier findings in insects (Zdobnov & Bork, 2007), though that study was focused on insect species

3 between orders.

Microsynteny changes more frequently than macrosynteny

Given that many inversions (of three or more genes) do not disrupt microsynteny, many inverted

blocks would still be detected as syntenic, and the slow accumulation of inversions would not change

the fraction of genes in blocks or affect the overall microsynteny estimates. Translocations, however,

can change macrosynteny but not microsynteny, as genes move as a block to a different chromosome.

We examined this for each congeneric species pair as a function of the number of genes on

homologous chromosomes in both species compared to the number of translocated genes or inserted

novel genes.

Like the relationship with protein identity, we found that macrosynteny and microsynteny broadly

tend to correlate together, (Figure 1D), but can be decoupled in many species pairs. In most species

pairs (other than the mammal set) microsynteny appears to be less constrained than macrosynteny. For

instance, although microsynteny is almost absent in the insects analysed, macrosynteny remains

broadly detectable in these organisms. Mammals, however, display a different pattern of

chromosomal evolution, with several species pairs showing chromosomal rearrangements while

preserving their microsynteny.

While there are examples of changes in karyotype between species (such as the fusion resulting in Chr

2 in humans (Yunis & Prakash, 1982)), only a few of the species in our study had obvious karyotype

differences. Macrosynteny is relatively conserved among *Drosophila* species with very conserved

microsynteny and relatively low protein sequence identity. Between *D. melanogaster* and *D. erecta*

(the closest pair in our study), only one translocation is evident in the genome (Figure 2A). In

Octopus, which possesses the largest invertebrate genome in our study, the karyotype appears to be

the same between the two species analysed, but with frequent small inversions in most chromosomes

(Figure 2B). Gene arrangement in *Octopus* is known to be variable, such as the rearrangement of the

Hox cluster (Albertin et al., 2015). Other species have karyotype variations. The water flea *Daphnia*

pulex has 12 chromosomes, while *D. magna* has only ten. These *Daphnia* species show conserved

macrosynteny but rapidly decaying microsynteny (Figure 2C). Also, the karyotype change in *Daphnia*

appears to have been caused by recent Robertsonian fission/fusion events in one of the species. For

Crassostrea / *Magallana* (oysters), two opposite chromosomal fusion/fission events were evident

(Figure 2D), producing the same number of chromosomes in the two species.

Figure 2: Example dotplots between congeneric species, showing a range of karyotypic states. All axes indicate positions in the genome in megabases. (A) *Drosophila*, showing two translocations/splits in blue boxes, one of which is unassigned to a chromosome in the assembly. (B) *Octopus*, with no obvious translocations but many small inversions. (C) *Daphnia*, red boxes indicate two chromosomal fusions/fissions with mixing in *D. magna*. (D) *Crassostrea/Magallana*, with two evident chromosomal fusions/fissions and split between the two species, as red and blue boxes, respectively. See online repository data for higher resolution versions of all species pairs.

Distribution of protein identity values is varied across species pairs

We then examined genomic signatures of speciation across the pairs of congeneric species across the animal kingdom. Species are not typically defined by molecular distances, so all of the species pairs in our study, though formally described as species, may all be at different genetic stages of divergence, or varied amounts of time of divergence. If all pairs show the same broader pattern of genetic changes, then they can be thought of as being at different stages along the same underlying process. However, if they show very different patterns between species, then multiple underlying processes are possible.

Among congeneric species, many one-to-one orthologous proteins are nearly identical for several pairs (Figure 3A,D, Supplemental Figure 7), and only few proteins were highly divergent (Supplemental Figure 7). Over time, as mutations accumulate throughout the genomes of the two separate species (Figure 3B,C,E,F), the number of completely identical proteins decreases, in a sense turning Figure 3A into 3B. Additionally, the total number of protein pairs being compared also decreases because the more-divergent proteins become difficult to detect by BLAST, effectively "dropping off the graph," and are neither counted for protein sequence identity nor detected in synteny blocks. The drop-off threshold appears to be at roughly 40% protein identity (Figure 3, see Supplemental Figure 8 for histograms of all species). In the case of *Crassostrea*, where two of the species are now assigned to the genus *Magallana*, the observed distributions of protein identity in Figure 3B,C are potentially are a sign of long-past separations, and that what used to be several species in a single genus indeed may be better described as separate genera. Similar distributions of protein identity are seen below in *Drosophila* (Figure 3E,F), and in *Daphnia* (Supplement Figure 8F).

Figure 3: Histograms of protein percent identity for *Crassostrea / Magallana* (A-C) and *Drosophila* (D-F). Protein identity is binned at intervals of 0.1% identity. Note the large differences in Y-axis scale, driven by a large number of 100% identical proteins in A and D.

Lack of genomic signatures of speciation

The taxonomic hierarchy can be considered an abstraction of the true divergence within a clade.

However, taxonomic ranks are arbitrary and do not reflect divergence similarly in different groups.

Yet, as ranks can be used as proxies for phylogenetic diversity, taxonomic ranks remain meaningful in

contexts such as conservation biology (Hendricks et al., 2014). In our study, as there are multiple

trajectories of genetic change among congeneric species (Figure 1), we could not identify a clear

genomic distinction based on macro- and/or microsyntenic patterns or percent protein identity that

could possibly be useful to delimit different animal species, and generally, any other taxonomic level.

Within some clades, patterns related to specific chromosomal features, such as centromere position,

may emerge. In our data, some species pairs (e.g., in genera *Crassostrea, Daphnia*) show low levels

of microsynteny and relatively low protein sequence identity indicating that the taxonomic rank

'genus' in these pairs point to different stages of the evolutionary process (see, for example, also

Avise and Liu 2011). For instance, the two oyster species, now *M. angulata* and *C. virginica*,

represent a pair where substantial rearrangements are visible in the genome. These two species are

also separated geographically, as *M. angulata* was originally from the Pacific, while *C. virginica* is

from the Atlantic. The degree of rearrangement in *Daphnia* (Figure 2C) is similar, and potentially this genus should be re-evaluated to see whether it is better described as two genera.

As in most other similar studies, the synteny assignments in our method are ultimately dependent upon local sequence similarity (BLAST, DIAMOND, etc). Hence, one of two effects could result in the exclusion of a gene from our analysis: duplication (no longer a one-to-one ortholog) or mutation beyond the detection of BLAST. This second cause may explain the regression line found by (Zdobnov & Bork, 2007) and the apparent lower bound for protein identity to microsynteny (Figure 1C). In this regard, as protein pairs diverge beyond the detection limit imposed by sequence similarity searches, e.g., in more distantly related species or genes with low evolutionary pressure, they cannot be detected as syntenic. Thus, it is very unlikely to find a species pair with very high microsynteny and low protein identity but not the opposite.

Difficulty in quantification of syntenic blocks and microsynteny

Meaningful measurements of microsynteny are challenging to get with current data. For instance, assembly quality could affect synteny assessments, as lower quality assemblies would be expected to increase the number of synteny blocks but not necessarily affect the number of syntenic genes. Due to the high sensitivity of the analysis to different parameters, real biological features such as gene insertions, duplications, or long intergenic regions can result in synteny blocks being split in two. Thus, the number of synteny blocks themselves is less important than the number of genes contained in blocks. A further confounding factor is the presence of unassigned scaffolds (i.e., the number of scaffolds is far above the number of chromosomes) in all assemblies analyzed. These may be unmerged haplotypes and would affect the measurement by increasing the number of genes not found in matching synteny blocks as they appear as duplicates.

One attempt to model the syntenic change between mouse and human (Nadeau & Taylor, 1984) had also assumed that reciprocal translocations and not inversions typically caused the synteny breaks. Under that model, it was assumed that the length of synteny blocks follows a negative exponential distribution and that synteny breaks disrupt both micro- and macro-synteny, something not typically observed. Nadeau's model was disputed by Zdobnov and Bork (2007), who argued that the distribution of block length followed a power law, not an exponential distribution. We found similar correlations in block length in our dataset, though R^2 ranged between 0.713 and 0.945 (Supplemental Figure 9). Thus, the correlation between block length and number of blocks is not as strong as proposed by Zdobnov and Bork (2007), suggesting that other biological factors affect chromosomal breakpoints (Pevzner & Tesler, 2003).

Challenges in quantification of rearrangement time

A prominent unanswered question of this study is whether chromosome rearrangements occur at the same rate across a clade. Such a study would be of great interest in evolutionary biology, even if difficult to design correctly. Firstly, such a study would require enough sequenced high-quality chromosome-scale species-pair genomes to identify which lineage had the inversion or translocation based on genomic triangulation. We are currently far from this objective. In our study, there were only 18 genera on NCBI with more than two species with chromosome-level assemblies, and several of those still did not include an annotation.

Secondly, one must infer divergence times between taxa in a way that is not reliant on protein identity. Divergence time estimates are often based on molecular clock inferences. While fossil calibrations are necessary for this analysis, the time estimates are then shifted based on molecular changes, i.e, the inferred divergence time depends on the protein identity. Unsurprisingly, one would likely find a strong correlation between protein identity and divergence time (as in Supplemental Figure 10), but that is a problem of circular logic; the divergence time was a function of protein identity in the first place, so it cannot be applied as an independent measurement for rearrangements. This is extended with the correlation of protein identity to the number of genes in synteny blocks. Consequently, protein identity, microsynteny, and molecular clock-based divergence estimates are likely correlated. Unfortunately, this means that molecular clock divergence time should not be used to test the rearrangement rate in lineages. One possibility for timing narrow-scale speciation could be the usage of codon evolution models. For instance, in Placozoa, it was observed that the nucleotide identity differed from protein identity such that highly conserved genes had many synonymous nucleotide changes instead of simply having fewer total mutations (Eitel et al., 2018). Thus, codon evolution models could provide a different age estimate than the proteins.

Conclusions

Defining and delimiting species is one of the important and basic requirements for studying biodiversity and its evolution, and it is pivotal for the management and conservation of natural resources. However, even in the times of DNA barcoding and genome sequencing, no universal threshold that applies, for example, across all animals, has been identified, and different species concepts are applied in different taxa. Synteny, i.e., the shared arrangement of genes on chromosomes between species, has been used as a marker of shared ancestry, with synteny-breaking events resulting in genomic incompatibilities between populations, ultimately leading to speciation. However, in our study of the genomes of congeneric pairs, we found that genomic changes inferred from speciation are variable and contextual and that synteny should not be used at present as a measure of divergence or

speciation. Once more high-quality, chromosome-scale genomes of sister species-pairs are available, a clear pattern of genome evolution may emerge..

Data, script, and code availability

Additional data, figures, scripts and analytical resources can be accessed in a GitHub repository https://github.com/PalMuc/congeneric_synteny, archived with a DOI at Zenodo (https://zenodo.org/doi/10.5281/zenodo.10069099).

Supplementary Information

Supplementary information is available from https://github.com/PalMuc/congeneric_synteny/tree/main/supplements_for_paper

Funding

G.W. acknowledges funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 764840 (ITN IGNITE).

Acknowledgements

WRF would like to thank M. Dohrmann and D. Schultz for helpful discussions. We would like to thank M. Eitel for helpful discussions and comments. G.W. dedicates this work to his late wife Connie Wörheide, who left us too early during the preparation of this manuscript.

References

Albertin, C. B., Simakov, O., Mitros, T., Wang, Z. Y., Pungor, J. R., Edsinger-Gonzales, E., Brenner, S., Ragsdale, C. W., & Rokhsar, D. S. (2015). The octopus genome and the evolution of cephalopod neural and morphological novelties. *Nature*, *524*(7564), 220–224. http://www.nature.com/doifinder/10.1038/nature14668

Assis, R., Kondrashov, A. S., Koonin, E. V., & Kondrashov, F. A. (2008). Nested genes and increasing organizational complexity of metazoan genomes. *Trends in Genetics*, *24*(10), 475– 478. https://doi.org/10.1016/j.tig.2008.08.003

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND.

Nature Methods, *12*(1), 59–60. https://doi.org/10.1038/nmeth.3176

De Queiroz, K. (2007). Species concepts and species delimitation. *Systematic Biology*, *56*(6), 879–886. https://doi.org/10.1080/10635150701701083

Eitel, M., Francis, W. R., Varoqueaux, F., Daraspe, J., Osigus, H.-J., Krebs, S., Vargas, S., Blum, H., Williams, G. A., Schierwater, B., & Wörheide, G. (2018). Comparative genomics and the nature of placozoan species. *PLoS Biology*, *16*(7), e2005359. https://doi.org/10.1371/journal.pbio.2005359

Enattah, N. S., Sahi, T., Savilahti, E., Terwilliger, J. D., Peltonen, L., & Järvelä, I. (2002). Identification of a variant associated with adult-type hypolactasia. *Nature Genetics*, *30*(2), 233–237. https://doi.org/10.1038/ng826

Engström, P. G., Ho Sui, S. J., Drivenes, O., Becker, T. S., & Lenhard, B. (2007). Genomic regulatory blocks underlie extensive microsynteny conservation in insects. *Genome Research*, *17*(12), 1898–1908. https://doi.org/10.1101/gr.6669607

Enright, A. J., Van Dongen, S., & Ouzounis, C. A. (2002). An efficient algorithm for large-scale detection of protein families. *Nucleic Acids Research*, *30*(7), 1575–1584. https://doi.org/10.1093/nar/30.7.1575

Francis, W. R., Eitel, M., Vargas, S., Garcia-Escudero, C. A., Conci, N., Deister, F., Mah, J. L., Guiglielmoni, N., Krebs, S., Blum, H., Leys, S. P., & Wörheide, G. (2023). The genome of the reef-building glass sponge *Aphrocallistes vastus* provides insights into silica biomineralization. *Royal Society Open Science*, *10*(6), 230423. https://doi.org/10.1098/rsos.230423

Francis, W. R., & Wörheide, G. (2017). Similar ratios of introns to intergenic sequence across animal genomes. *Genome Biology and Evolution*, *9*(6), 1582–1598. https://doi.org/10.1093/gbe/evx103

Frankham, R., Ballou, J. D., Dudash, M. R., Eldridge, M. D. B., Fenster, C. B., Lacy, R. C., Mendelson, J. R., Porton, I. J., Ralls, K., & Ryder, O. A. (2012). Implications of different species concepts for conserving biodiversity. *Biological Conservation*, *153*, 25–31. https://doi.org/10.1016/j.biocon.2012.04.034

Ghiselin, M. T. (2010). Species Concepts. In *eLS*. John Wiley & Sons, Ltd. https://doi.org/10.1002/9780470015902.a0001744.pub2

Hausdorf, B. (2011). Progress toward a general species concept. *Evolution; International Journal of Organic Evolution*, *65*(4), 923–931. https://doi.org/10.1111/j.1558-5646.2011.01231.x

Hendricks, J. R., Saupe, E. E., Myers, C. E., Hermsen, E. J., & Allmon, W. D. (2014). The generification of the fossil record. *Paleobiology*, *40*(4), 511–528. https://doi.org/10.1666/13076

Hey, J., Waples, R. S., Arnold, M. L., Butlin, R. K., & Harrison, R. G. (2003). Understanding and confronting species uncertainty in biology and conservation. *Trends in Ecology & Evolution*, *18*(11), 597–603. https://doi.org/10.1016/j.tree.2003.08.014

Jiang, J. B., Quattrini, A. M., Francis, W. R., Ryan, J. F., Rodríguez, E., & McFadden, C. S. (2019). A hybrid de novo assembly of the sea pansy (*Renilla muelleri*) genome. *GigaScience*, *8*(4), giz026. https://doi.org/10.1093/gigascience/giz026

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–780. https://doi.org/10.1093/molbev/mst010

Kenny, N. J., Francis, W. R., Rivera-Vicéns, R. E., Juravel, K., de Mendoza, A., Díez-Vives, C., Lister, R., Bezares-Calderón, L. A., Grombacher, L., Roller, M., Barlow, L. D., Camilli, S., Ryan, J. F., Wörheide, G., Hill, A. L., Riesgo, A., & Leys, S. P. (2020). Tracing animal genomic evolution with the chromosomal-level assembly of the freshwater sponge *Ephydatia muelleri*. *Nature Communications*, *11*(1), 1–11. https://doi.org/10.1038/s41467-020-17397-w

Khalturin, K., Shinzato, C., Khalturina, M., Hamada, M., Fujie, M., Koyanagi, R., Kanda, M., Goto, H., Anton-Erxleben, F., Toyokawa, M., Toshino, S., & Satoh, N. (2019). Medusozoan genomes inform the evolution of the jellyfish body plan. *Nature Ecology & Evolution*, *3*(5), 811–822. https://doi.org/10.1038/s41559-019-0853-y

Kikuta, H., Laplante, M., Navratilova, P., Komisarczuk, A. Z., Engström, P. G., Fredman, D., Akalin, A., Caccamo, M., Sealy, I., Howe, K., Ghislain, J., Pezeron, G., Mourrain, P., Ellingsen, S., Oates, A. C., Thisse, C., Thisse, B., Foucher, I., Adolf, B., … Becker, T. S. (2007). Genomic regulatory blocks encompass multiple neighboring genes and maintain conserved synteny in vertebrates. *Genome Research*, *17*(5), 545–555. https://doi.org/10.1101/gr.6086307

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* , *34*(18),

3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Lv, J., Havlak, P., & Putnam, N. H. (2011). Constraints on genes shape long-term conservation of macro-synteny in metazoan genomes. *BMC Bioinformatics*, *12 Suppl 9*(Suppl 9), S11. https://doi.org/10.1186/1471-2105-12-S9-S11

Mayr, E. (1942). *Systematic and the origin of species*. Columbia University Press.

Nadeau, J. H., & Taylor, B. A. (1984). Lengths of chromosomal segments conserved since divergence of man and mouse. *Proceedings of the National Academy of Sciences of the United States of America*, *81*(3), 814–818. https://doi.org/10.1073/pnas.81.3.814

Ohno, S. (1970). Gene duplication as an evolutionary force. In H. Peeters (Ed.), *Protides of the Biological Fluids* (Vol. 17, pp. 39–42). Elsevier. https://doi.org/10.1016/B978-0-08-015566-1.50009-5

Orr, H. A. (1996). Dobzhansky, Bateson, and the genetics of speciation. *Genetics*, *144*(4), 1331–1335. https://doi.org/10.1093/genetics/144.4.1331

Pertea, G., & Pertea, M. (2020). GFF Utilities: GffRead and GffCompare. *F1000Research*, *9*. https://doi.org/10.12688/f1000research.23297.2

Pevzner, P., & Tesler, G. (2003). Human and mouse genomic sequences reveal extensive breakpoint reuse in mammalian evolution. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(13), 7672–7677. https://doi.org/10.1073/pnas.1330369100

Salvi, D., & Mariottini, P. (2017). Molecular taxonomy in 2D: a novel ITS2 rRNA sequence-structure approach guides the description of the oysters' subfamily Saccostreinae and the genus *Magallana* (Bivalvia: Ostreidae). *Zoological Journal of the Linnean Society*, *179*(2), 263–276. https://doi.org/10.1111/zoj.12455

Salvi, D., & Mariottini, P. (2021). Revision shock in pacific oysters taxonomy: The genus *Magallana* (formerly *Crassostrea* in part) is well-founded and necessary. *Zoological Journal of the Linnean Society*, *192*(1), 43–58. https://doi.org/10.1093/zoolinnean/zlaa112

Schultz, D. T., Francis, W. R., McBroome, J. D., Christianson, L. M., Haddock, S. H. D., & Green, R. E. (2021). A chromosome-scale genome assembly and karyotype of the ctenophore *Hormiphora californensis*. *G3* , *11*(11), jkab302. https://doi.org/10.1093/g3journal/jkab302

Shinzato, C., Khalturin, K., Inoue, J., Zayasu, Y., Kanda, M., Kawamitsu, M., Yoshioka, Y., Yamashita, H., Suzuki, G., & Satoh, N. (2021). Eighteen coral genomes reveal the evolutionary origin of Acropora strategies to accommodate environmental changes. *Molecular Biology and Evolution*, *38*(1), 16–30. https://doi.org/10.1093/molbev/msaa216

Simakov, O., Bredeson, J., Berkoff, K., Marletaz, F., Mitros, T., Schultz, D. T., O'Connell, B. L., Dear, P., Martinez, D. E., Steele, R. E., Green, R. E., David, C. N., & Rokhsar, D. S. (2022). Deeply conserved synteny and the evolution of metazoan chromosomes. *Science Advances*, *8*(5), eabi5884. https://doi.org/10.1126/sciadv.abi5884

Slijepcevic, P. (1998). Telomeres and mechanisms of Robertsonian fusion. *Chromosoma*, *107*(2), 136–140. https://doi.org/10.1007/s004120050289

Srivastava, M., Begovic, E., Chapman, J., Putnam, N. H., Hellsten, U., Kawashima, T., Kuo, A., Mitros, T., Salamov, A., Carpenter, M. L., Signorovitch, A. Y., Moreno, M. A., Kamm, K., Grimwood, J., Schmutz, J., Shapiro, H., Grigoriev, I. V., Buss, L. W., Schierwater, B., … Rokhsar, D. S. (2008). The *Trichoplax* genome and the nature of placozoans. *Nature*, *454*(7207), 955–960. https://doi.org/10.1038/nature07191

Ulate, K., & Alfaro-Montoya, J. (2010). Sperm–egg incompatibility in inter-specific inseminations of *Penaeus (Litopenaeus) occidentalis*, *P. (Litopenaeus) stylirostris* and *P. (Litopenaeus) vannamei*. *Aquaculture* , *309*(1), 290–292. https://doi.org/10.1016/j.aquaculture.2010.08.029

Vrana, P. B., Fossella, J. A., Matteson, P., del Rio, T., O'Neill, M. J., & Tilghman, S. M. (2000). Genetic and epigenetic incompatibilities underlie hybrid dysgenesis in Peromyscus. *Nature Genetics*, *25*(1), 120–124. https://doi.org/10.1038/75518

Yunis, J. J., & Prakash, O. (1982). The origin of man: a chromosomal pictorial legacy. *Science*, *215*(4539), 1525–1530. https://doi.org/10.1126/science.7063861

Zachos, F. E. (2016). *Species Concepts in Biology*. Springer International Publishing. https://doi.org/10.1007/978-3-319-44966-1

Zdobnov, E. M., & Bork, P. (2007). Quantification of insect genome divergence. *Trends in Genetics*, *23*(1), 16–20. https://doi.org/10.1016/j.tig.2006.10.004

Zechner, U., Reule, M., Orth, A., Bonhomme, F., Strack, B., Guénet, Hameister, H., & Fundele, R.

1   (1996). An X-chromosome linked locus contributes to abnormal placental development in mouse

2   interspecific hybrid. *Nature Genetics*, *12*(4), 398–403. https://doi.org/10.1038/ng0496-398