Dynamic estimation of the attentional field from visual cortical activity

Ilona M. Bloem[1,2]*, Leah Bakst[1]*, Joseph T. McGuire[1], Sam Ling[1]

[1] Department of Psychological & Brain Sciences, Boston University, Boston, USA
[2] Department of Computational Cognitive Neuroscience and Neuroimaging, Netherlands Institute for Neuroscience, Amsterdam, Netherlands & Spinoza Centre for Neuroimaging, Amsterdam, Netherlands

* These authors provided equal contribution

Correspondence should be addressed to Sam Ling (samling@bu.edu).

Abbreviated title (50-character max): Dynamical estimation of the attentional field

Number of pages: 25
Number of figures: 6
Number of words for:
  Abstract: 247
  Introduction: 649
  Discussion: 1149

The authors declare no competing financial interests.

Acknowledgements: This work was supported by National Science Foundation grants SMA-1809071, BCS-1625552, and BCS-1755757, National Institutes of Health grants F32-EY029134, R01-EY028163, and R01-MH126971, Office of Naval Research grant N00014-17-1-2304, and the Center for Systems Neuroscience Postdoctoral Fellowship at Boston University. The content of this paper does not necessarily represent the official views of the funding agencies.

Author contributions: Conceptualization, Methodology & Writing – Review & Editing I.M.B., L.B., J.T.M., S.L.; Investigation, Analysis, Visualization, & Writing – Original Draft, I.M.B. & L.B.; Resources & Supervision, J.T.M & S.L.; Funding Acquisition L.B., J.T.M., S.L.

Abstract

Navigating around the world, we must adaptively allocate attention to our surroundings based on anticipated future stimuli and events. This allocation of spatial attention boosts visuocortical representations at attended locations and locally enhances perception. Indeed, spatial attention has often been analogized to a "spotlight" shining on the item of relevance. Although the neural underpinnings of the locus of this attentional spotlight have been relatively well studied, less is known about the size of the spotlight: to what extent can the attentional field be broadened and narrowed in accordance with behavioral demands? In this study, we developed a paradigm for dynamically estimating the locus and spread of covert spatial attention, inferred from visuocortical activity using fMRI in humans. We measured BOLD activity in response to an annulus while participants (4 female, 4 male) used covert visual attention to determine whether more numbers or letters were present in a cued region of the annulus. Importantly, the width of the cued area was systematically varied, calling for different sizes of the attentional spotlight. The deployment of attention was associated with an increase in BOLD activity in corresponding retinotopic regions of visual areas V1—V3. By modeling the visuocortical attentional modulation, we could reliably recover the cued location, as well as a broadening of the attentional enhancement with wider attentional cues. This modeling approach offers a useful window into the dynamics of attention and spatial uncertainty.

Significance Statement

This study explores whether spatial attention can dynamically adapt by shifting and broadening the attentional field. While previous research has focused on the modulation of neural responses at attended locations, less is known about how the size of the attentional field is represented within visual cortex. Using fMRI, we developed a novel paradigm to estimate the spatial tuning of the attentional field and demonstrate that we were able to recover both the location as well as the width of the attentional field. Our findings offer new insights into the neural mechanisms underlying the deployment of spatial attention, contributing to a deeper understanding of how spatial attention supports visual perception.

Introduction

We bounce attention around all the time. Take, for instance, when we're monitoring oncoming traffic while driving. It isn't sufficient to attend to the single most likely source of traffic. Instead, attention adaptively broadens and narrows to cover the anticipated spatial distribution of relevant events. The need to spread attention across different swaths of the visual field is driven, to a large degree, by spatial uncertainty: statistical regularities give us a general sense as to where something useful might happen, and this evolves from moment to moment. We navigate this uncertainty by dynamically deploying spatial attention.

Covert spatial attention improves behavioral performance at attended locations at the cost of performance at unattended locations (Posner, 1980), leading to a common metaphor that spatial selective attention acts as a 'spotlight' or 'zoom lens' (Shaw and Shaw, 1977; Posner, 1980; Eriksen and St. James, 1986; Carrasco, 2011). This attentional 'spotlight' is characterized by a specific size and location and traverses the visual field based on behavioral demands (Eriksen and St. James, 1986; Castiello and Umiltà, 1990), selectively boosting information at the attended location within the visual system while suppressing information elsewhere. Animal studies have observed multiplicative increases in visuocortical neural responses at attended locations (McAdams and Maunsell, 1999; Maunsell, 2015) and human neuroimaging studies have found similar focal enhancements of population responses (Kastner *et al.*, 1998; Brefczynski and DeYoe, 1999; McMains and Somers, 2004; Datta and DeYoe, 2009; Sprague and Serences, 2013; Puckett and DeYoe, 2015; Samaha, Sprague and Postle, 2016; Shioiri *et al.*, 2016; Bloem and Ling, 2019).

While neural modulation at the locus of attention has been relatively well studied, less is known regarding the neural signatures of the size of the attentional field (Yeshurun, 2019). Spreading attention over a larger region of visual space can decrease behavioral performance, but only a handful of studies have interrogated associated effects within visual cortex (Müller *et al.*, 2003; Herrmann *et al.*, 2010; Itthipuripat *et al.*, 2014; Feldmann-Wüstefeld and Awh, 2020). This is surprising, as the spatial distribution of the attentional field is a key feature in an influential theoretical model of attention (Reynolds and Heeger, 2009). The model assumes that the size of the attentional field can be adjusted based on task demands and that the interaction between attentional field size and stimulus-related factors can predict observed attentional gain effects.

While the studies that have experimentally manipulated the attentional field size found evidence congruent with this prominent theory (Herrmann *et al.*, 2010; Itthipuripat *et al.*, 2014; Kınıklıoğlu and Boyaci, 2022), few studies have directly investigated the spatial extent of the attentional window and its concomitant neural representation. One neuroimaging study revealed that the attentional field expanded in the face of greater task-related uncertainty (Herrmann *et al.*, 2010), while other studies showed that the responsive area of visual cortex increased in size, coupled with a decrease of the overall population response (Müller *et al.*, 2003; Feldmann-Wüstefeld and Awh, 2020). While these studies are consistent with the notion that the attentional field size can be detected

in visual cortex, methods for dynamically recovering location and field size from moment
to moment are lacking.
In this study, we developed a paradigm that allowed us to dynamically
characterize the spatial tuning of spatial attention across the visual field. Using fMRI in
humans, we examined whether attentional modulation of the BOLD response spanned
a larger area of visual cortex when participants were cued to perform attend to a larger
region of space. Behavioral performance confirmed that participants could successfully
allocate their attention to different-sized swaths of the visual field. This deployment of
attention was associated with a boost in cortical activity in the corresponding retinotopic
areas of visual cortex. By modeling the location and spread of the visuocortical
modulation, we dynamically recovered the cued location from the attentional activity with
a high degree of fidelity, together with a broadening of the attentional enhancement for
wider attentional cues.

Materials and Methods

2 **Participants.** Eight healthy adults (4 female, 4 male, mean age = 30) participated in the
3 main attention experiment, five of whom also participated in a second experiment
4 featuring a contrast manipulation. All participants had normal or corrected-to-normal
5 vision. All procedures were approved by the Boston University Institutional Review
6 Board, and informed consent was obtained from all participants.
7
Apparatus and stimuli. Participants were presented with stimuli generated using
PsychoPy (v1.85.1; Peirce, 2007) on a MacBook Pro. The visual stimuli were displayed
on a rear-projection screen (subtending ~20°x16° visual angle) using a VPixx
Technologies PROPixx DLP LED projector (maximum luminance 306 cd/m^2).
Participants viewed the screen through a front surface mirror. Participants were placed
comfortably in the scanner with padding to minimize head motion.
14
Procedure.
Attentional width manipulation. Participants were instructed to fixate a central point
(radius 0.08° visual angle) while dynamic pixelwise white noise (flickering at 10 Hz, 50%
contrast) was presented in the periphery (annulus spanning 4.6° to 7.4° visual angle).
The annulus was segmented into 20 bins (18° polar angle per bin) by white grid lines
radiating from a white circle at the center of the screen (radius 0.25°), passing behind
the annulus, and terminating at 8.5° eccentricity. In the middle of each bin, a number or
letter (height: 2.1°) was superimposed on the white noise annulus (see **Figure 1a**). For a
subset of the participants (3 out of 8) the screen distance inside the scanner was
changed, therefore for those participants the letter size was 1.86° visual angle, and the
white noise annulus spanned 4.1° to 6.5° visual angle. The set of possible letters included
all lowercase letters of the Latin alphabet except a, b, e, g, i, o, and u. The set of numbers
included 2, 3, 4, 5, 7, and 8.
Participants were cued to attend covertly to a contiguous subset of the bins and
their task was to report, via button press, whether there were more *numbers* or *letters*
present within the cued region. The cue was a bold red segment on the central white
circle, which corresponded to 1, 3, 5, or 9 bins (18°, 54°, 90°, or 162° polar angle; see
Figure 1a). The true proportion of letters versus numbers was controlled within each cue
width condition. For cued regions of 1 bin, there was either a single number or letter in
the bin. For cued regions of 3 bins, the ratio was always 2:1 (either two numbers and one
letter or vice versa). For cued regions of 5 bins, the ratio was 3:2, and for cued regions
of 9 bins, the ratio was 6:3. Cues could be centered on any of the 20 bins.
Participants completed 8 to 12 runs of the task (mean = 10.4), with each run
lasting 341 s and containing 100 trials. Each cue remained constant for a block of five
trials (lasting 15.5 s, 10 TRs), although the letters and numbers within the cued region
changed on every trial. Thus, each participant saw 20 unique cues (combinations of cue
location and width) per run. Each run began and ended with 15.5 s of the dynamic noise
annulus.

During each trial, the cue and white noise annulus were presented alone for 1.35
s. The numbers and letters were then displayed for 0.5 s. Thereafter, the cue and white
noise remained visible while the participant had 1.25 s to indicate whether there had
been more digits or letters within the cued region, resulting in a total trial duration of 3.1
s (2 TRs). No accuracy feedback was provided during the main experiment. However, all
participants completed three training runs with trial-by-trial feedback prior to the scan
session. During training runs, the response window was shortened to 1 s and the
remaining 0.25 s presented feedback in the form of a change in color of the fixation point
(blue for correct responses and orange for incorrect responses).
52
Physical contrast manipulation. A subset of participants (n=5) also participated in an
experiment that enhanced the physical contrast intensity of the dynamic visual noise in
segments of the annulus. This additional experiment was carried out during the same
scan session and allowed for benchmarking the detectability of stimulus-evoked
modulation in visual cortex using our analyses. The stimuli and trial structure were similar
to the attentional manipulation. The task differed in the following ways: (1) the contrast
of the white noise annulus was increased to 100% for segments of the annulus
corresponding to 1, 3, 5, 7 or 9 bins (18°, 54°, 90°, 126°, or 162° polar angle), with a
Gaussian rolloff ($\sigma = 15°$) that spanned 25% of the furthest included bins and 25% of the
adjacent excluded bins; (2) the enhanced segments were always centered on the
cardinal directions (0°, 90°, 180°, and 270° polar angle); (3) the contrast increase
remained constant for 15.5 seconds (10 TRs); (4) participants performed a color change
detection task at fixation. Each unique combination of 4 locations and 5 widths of the
contrast enhancement was shown once per run, with the order randomized. To estimate
a baseline response, each run started and ended with 15.5 seconds without contrast
modulation. Participants completed two runs total, each lasting 341 seconds (220 TRs).
Throughout the *physical contrast* runs, participants were instructed to fixate on a
central point (radius 0.08° visual angle) and to press a button when the fixation point
switched color (alternating white and red). The fixation point remained a color for at least
one second and then had a 10% probability of switching every 100 ms. No cue was
presented associated with the regions of increased contrast. Additionally, no letters or
numbers were superimposed on the white noise annulus.
75
Population receptive field mapping. Population receptive field (pRF) estimates were
obtained for each participant in a separate scan session. We used the experimental
procedure as described in the Human Connectome Project 7T Retinotopy dataset
(Benson *et al.*, 2018). Stimuli were composed of a pink noise background with colorful
objects and faces at various spatial scales, displayed on a mean luminance gray
background. Stimuli were updated at a rate of 15 Hz while participants performed a color
change detection task at fixation. Participants viewed two types of mapping stimuli: (1)
contracting/expanding rings and rotating wedges; (2) moving bar stimuli (Dumoulin and
Wandell, 2008; Kay *et al.*, 2013). A total of 4-6 scans (300 TRs) were collected for each
participant (2-3 scans per stimulus type).
86

MRI data acquisition. All MRI data were acquired at Boston University's Cognitive
Neuroimaging Center (Boston, Massachusetts) on a research-dedicated Siemens
Prisma 3T scanner using a 64-channel head coil. A scanning session lasted 2 hours.
All functional neuroimaging data were acquired using a simultaneous multislice (SMS)
gradient echo echoplanar acquisition protocol (Moeller *et al.*, 2010; Setsompop *et al.*,
2012): 2 mm isotropic voxels; FoV = 212 x 212 mm; 72 axial slices; TR = 1.55 s; TE =
35.60 ms; flip angle = 72°; multiband acceleration factor 4. We computed distortion field
maps by using a spin echo echoplanar protocol with opposite y-axis phase encoding
directions (2 mm isotropic voxels; FOV = 212 x 212 mm; TR = 8850 ms; TE = 70.80 ms;
flip angle = 90°). During a separate scan session, we acquired a whole-brain anatomical
scan using a T1-weighted multi-echo MPRAGE 3d sequence (1 mm isotropic; FoV = 256
x 256 mm; 176 sagittal slices; TR = 2530 ms; TE = 1.69 ms; flip angle = 7°), and the pRF
scans (occipital coverage only; right-left phase encoding; 2 mm isotropic voxels; FoV =
136 x 136 mm; 36 slices; TR = 1 s; TE = 35.4 ms; flip angle = 64°; multiband acceleration
factor 3).
102
103 ***MRI data analysis.***
104 *Structural data preprocessing.* Whole brain T1-weighted anatomical data were analyzed
105 using the standard 'recon-all' pipeline provided by Freesurfer software (Freesurfer
106 version 5.3, (Fischl, 2012)), generating cortical surface models, whole-brain
107 segmentation, and cortical parcellations.
108
Functional data preprocessing. All analyses were performed in the native space for each
participant. First, EPI distortion correction was applied to all fMRI BOLD time-series data
using a reverse phase-encode method (Andersson, Skare and Ashburner, 2003)
implemented in FSL (Smith *et al.*, 2004). All functional data were then preprocessed using
FS-FAST (Fischl, 2012), including standard motion-correction procedures, Siemens slice
timing correction, and boundary-based registration between anatomical and functional
volumetric spaces (Greve and Fischl, 2009). To facilitate voxel-wise analysis, no
volumetric smoothing was performed and across-run within-modality robust rigid
registration was applied (Reuter, Rosas and Fischl, 2010), with the middle time-point of
the first run serving as the target volume, and the middle time-point of each subsequent
run used as a movable volume for alignment. Lastly, data were detrended (0.005 Hz
high-pass filter) and converted to percent signal change for each voxel independently
using custom code written in MATLAB (version 2020b).
122
Population receptive field mapping and voxel selection. The time series were analyzed
using the analyzePRF toolbox in MATLAB, implementing a compressive spatial
summation pRF model (Kay *et al.*, 2013). The results of the pRF analysis were used to
manually draw boundaries between early visual regions (V1, V2, and V3), which served
as our regions of interest (ROIs).
Within each ROI, pRF modeling results were used to constrain voxel selection
used in the main experiment. We excluded voxels with a preferred eccentricity outside
the bounds of the pRF stimulus (<0.7° and >9.1°), with a pRF size smaller than 0.01°, or
with poor spatial selectivity as indicated by the pRF model fit ($R^2 < 10\%$). Following our

2D visualizations (see below), we further constrained voxel selection by only including
voxels whose pRF overlapped with the white noise annulus.

2D visualizations of attentional modulation. To visualize the topography of attentional
modulation under different cue widths, we projected the average BOLD responses for a
given block (10 TRs with a consistent cue location and width, shifted by 3 TRs [4.65 s]
to compensate for the hemodynamic delay) into the visual field using each voxel's pRF
location. This method is similar to that described in (Favila, Kuhl and Winawer, 2022).
First, we computed the Cartesian (x,y) coordinates from the pRF eccentricity and polar
angle estimates for each voxel. Then, within a given ROI, we interpolated the BOLD
responses over (x,y) space to produce a full-field representation. Each representation
was then z-scored to allow for comparison across blocks, cue conditions, and
participants. Finally, the representation was rotated so that the center of the cue was
aligned to the right horizontal meridian (see **Figure 2a**).

1D spatial profile of attentional modulation. We also examined the spatial profile of
attentional modulation as a function of polar angle. Voxels with pRFs overlapping the
white noise annulus were grouped into 60 bins according to their pRF polar angle
estimate (6° polar angle bin width). We computed a median BOLD response within each
bin. To improve the signal-to-noise ratio, the resulting profile was smoothed with a
moving average filter (width 18° polar angle; see **Figure 2b**).

Model fitting. We quantified the spatial profile of attentional modulation with a
generalized Gaussian model (Nadarajah, 2005). The generalized Gaussian function (G)
combines Gaussian and Laplace distributions:

$$G = exp\left\{-\left|\frac{x-\mu}{\sigma}\right|^{\beta}\right\} \tag{1}$$

The function has free parameters for location (μ), scale (σ), and shape (β). The shape
parameter enables the tails of the distribution to become heavier than Gaussian (when
$\beta < 2$), or lighter than Gaussian (when $\beta > 2$); as $\beta \to \infty$, the model approaches a
uniform distribution.

Next, G was normalized to range between 0 and 1, and vertically scaled and shifted by
two additional free parameters for amplitude (a) and baseline offset (b):

$$\hat{y} = a \cdot G + b \tag{2}$$

We fit the five free parameters (μ, σ, β, a, b) using the MATLAB optimization tool *fmincon,*
minimizing the squared error between the model prediction and the 1D profile described
above. To avoid local minima, we first ran a grid search to find the initialization values
with the lowest SSE (6 possible values for μ, equally spaced between 0 and 360°,

crossed with 6 possible values for σ, equally spaced between 9° and 162° polar angle;
$\beta = 4$; $a = 1$; $b = 0$). We imposed the following parameter bounds on the search: σ: [6°,
180° polar angle], β: [1.8, 50], and a: [0, 20]. μ was unbounded, but was wrapped to
remain within [0°, 360°].
From the model fits we computed the following summary metrics: 1) angular error,
defined as the polar-angle distance between the true and estimated location; 2) the full
width at half-maximum (FWHM) of the best-fitting generalized Gaussian function, which
served as our measure of the width of attentional modulation. The FWHM was controlled
mainly by the scale parameter (σ) but also to a lesser degree by the shape parameter (β;
see **Figure 3a**); 3) the gain modulation of the spatial profile (a); 4) the model's goodness
of fit quantified as the percentage of explained variance (R^2) in the spatial response
profile:

$$R^2 = 1 - \frac{(y - \hat{y})^2}{(y - \bar{y})^2} \tag{3}$$

180 ***Statistical testing***. To assess how the attentional cue width manipulation influenced the
181 1D spatial profile of BOLD modulation, we tested whether the computed summary
182 metrics (absolute angular error, FWHM, and amplitude) varied as a function of cue width.
183 Specifically, we performed a linear regression for each metric and tested whether the
184 slopes differed from zero. This was done independently for each ROI.
185
Eye-position monitoring. Gaze data were collected for all participants using an MR-
compatible SR Research EyeLink 1000+ eye tracker sampling at 1 kHz. Data from blink
periods were excluded from analysis. Participants maintained fixation throughout the
task, with average gaze eccentricity below 0.5° for all participants. Gaze eccentricity did
not significantly vary by cued width (pairwise comparison of width conditions using a
paired t-test, all $p >= 0.205$ with Bonferroni correction for multiple comparisons) nor
location (pairwise comparison, all $p >= 0.522$ with Bonferroni correction for multiple
comparisons).

1 Results

2 Behavioral performance indicates effective deployment of covert spatial attention

We set out to investigate the spatial distribution of attentional modulation within
visual cortex. To do so, we first ensured that participants (*n*=8) could successfully
allocate covert spatial attention to cued portions of the visual field. During the
experiment, participants' task was to fixate the center of the screen and report whether
there were more *numbers* or *letters* in a cued peripheral region (**Figure 1a**). The cued
region varied in location and width: it could be centered on any of 20 polar angles and
could span any of four widths (18°, 54°, 90°, and 162° of polar angle). Task performance
indicated that participants used the cue effectively, as the proportion of correct
responses was significantly above chance for all width conditions (**Figure 1b**; t-test, all
p<.001). We verified, with eye tracking, that participants performed the task using
peripheral vision while maintaining central fixation. The upper bound of the 95% CI for
each participant's average gaze eccentricity ranged from 0.29° (degrees of visual angle)
to 0.64° (mean = 0.48°; **Figure 1c**), suggesting that gaze did not exceed the cue annulus
at fixation and that participants used covert spatial attention to perform the task.

a. Attention task trial sequence

Figure 1. a. Task schematic. Participants' were instructed to maintain central fixation and use covert spatial attention to determine whether there were more numbers or letters present within a cued region of a white noise annulus. On each trial, the red cue was displayed alone for 1.35 s and remained present throughout the trial. Twenty digits and letters were then presented for 0.5 s, equally spaced and overlaid on the annulus. Participants had 1.25 s to indicate via button press whether more digits or letters were present in the cued region. The cue remained stable for 5 trials (10 TRs, 15.5 s), had a width of 1, 3, 5, or 9 segments (18°, 54°, 90°, or 162°), and was centered on any of the 20 digit/letter slots. **b.** Behavioral task performance: Group mean accuracy for each cue width. Error bars are SEM; gray circles show individual participants. **c.** Group mean gaze eccentricity (in degrees of visual angle) for each cue width, conventions as in **b**.

b. Behavioral task performance   *c. Gaze distance from fixation*

Attentional modulation of BOLD responses broadens with cue width

We assessed the spatial distribution of attention by visualizing how the BOLD
response was modulated by the location and width of the cue. To do so, we used each
voxel's population receptive field (pRF) to project BOLD responses for each attentional
cue into the visual field. The resulting 2D visual field maps were averaged across trials
for each cue width by rotating the maps so the attentional cue aligned to 0° polar angle
(right horizontal meridian). The reconstructed visual field maps revealed that increasing
cue width led to a concomitant broadening of attentional modulation in cortex (**Figure
2a**). While this pattern was evident in all three early visual regions (V1–V3), the effect
appeared to strengthen when ascending the visuocortical hierarchy.

Next, we computed the one-dimensional profile of attentional modulation at a
fixed eccentricity. We were able to do this because we manipulated the location of the
attentional field only as a function of polar angle, so all cues directed the attentional field
to iso-eccentric locations. We selected voxels with pRFs that overlapped the white noise
annulus and sorted them according to their polar angle preference.

a. Attention: 2D BOLD activity reconstruction

b. Spatial profile attentional modulation

Figure 2. a. BOLD response projected into the visual field for each attentional cue width. Heatmaps represent the group mean BOLD activity using each voxel's population receptive field (pRF) location within the visual field, shown separately for V1, V2, and V3. Maps were rotated to align all attentional cue locations to 0° polar angle (rightward). Concentric circles indicated by black dashed lines represent the location of the white noise annulus. **b.** Average spatial modulation profiles at the eccentricity of the annulus. The spatial profiles were recentered to 0° polar angle based on the cue location. Solid lines represent the group mean BOLD activity and shaded regions the SEM across participants.

For visualization purposes, the spatial response modulations were recentered to
align all cues at 0° polar angle and averaged across trials for each cue width separately.
Much like in the visual field reconstructions, there was a clear attentional enhancement
centered on 0°, which broadened and decreased in amplitude with cue width – a pattern
that was particularly evident in area V3 (**Figure 2b**).

Dynamic model-based recovery of the attentional field

We next applied a modeling approach to estimate the location and width of
attentional modulation, allowing us to further investigate the spread of attention in visual
cortex. To do this, we averaged the spatial response profiles across TRs within each 10-
TR block, in which the cue maintained a consistent location and width, yielding between
27 and 53 averaged spatial response profiles per participant for each width condition.
We fit a generalized Gaussian function to each of these spatial profiles to estimate the
location and width of attentional modulation per spatial profile (see **Figure 3a**). The width
of attentional modulation was quantified in terms of the full width at half maximum
(FWHM) of the best fitting model prediction (see **Figure 3b**).

a. Generalized Gaussian model   b. Quantify attentional field

Figure 3. a. Modeling approach. The generalized Gaussian model is characterized by parameters for location (μ), scale (σ), and shape (β). **b.** Example model fits for two spatial profiles. Dots indicate BOLD response amplitudes for two attentional cues differing in position and width. Solid lines indicate the best fitting model estimate. To quantify the attentional field, we extracted the location and gain (dashed arrows), as well as the width (FWHM; solid arrows).

Can we dynamically recover the attentional field from activity within visual cortex?
Model fits explained a substantial proportion of variance in the spatial profiles of BOLD
activity (**V1**: for 18° cues, mean [standard deviation] of R^2 = 0.42 [0.03]; for 54° cues, 0.43
[0.03]; for 90° cues, 0.44 [0.03]; for 162° cues, 0.42 [0.03]; **V2**: for 18° cues, 0.51 [0.05];
for 54° cues, 0.54 [0.05]; for 90° cues, 0.54 [0.04]; for 162° cues, 0.55 [0.04]; **V3**: for 18°
cues, 0.50 [0.03]; for 54° cues, 0.56 [0.04]; for 90° cues, 0.55 [0.03]; for 162° cues, 0.51
[0.02]). To interpret the estimated model parameters, we excluded the bottom 20% of
fits based on a pooled R^2 across V1, V2, and V3, leaving roughly equal proportions of
included blocks across cue width conditions (18°: mean [standard deviation] = 0.78
[0.04], 54°: 0.83 [0.05], 90°: 0.83 [0.04], 162°: 0.77 [0.07]).

To assess how well the model-estimated attentional field matched the cued
location, we first calculated the angular error between the cue center and the model's
estimated location parameter. The angular error distribution across blocks, separated by
width condition, is shown in **Figure 4a** for one example participant to display block-to-
block variation. The model reliably captured the location of the attentional field with low
angular error. This result was consistent across participants. The group mean absolute
angular error in V1 was 41.9° (SEM=2.86°), in V2 was 32.2° (2.31°), and in V3 was 24.7°
(1.54°). Additionally, the magnitude of the absolute error did not vary linearly with the
width of the cue in V1 or V2 (regression slope tested against zero; both $p>=.468$; **Figure**
4b). In V3, we observed a small but statistically significant increase in absolute error
magnitude associated with greater cue widths ($t(1,30)=2.86$, $p=.008$).

a. Example observer

b. Group results

Figure 4. Attentional field parameter estimates. **a.** The full parameter estimate distributions across blocks for location, width, and amplitude are shown for one example participant in V1, V2, and V3. Median parameter estimates are shown by the white points, with the box plot representing the 25th to 75th percentile, and whiskers extending to all non-outlier points. **b.** Group results for location, width, and amplitude estimates. Overall group mean and standard error are shown, separated by cue width and brain region.

Next, we evaluated the width of the attentional field by visualizing the distribution
of FWHM for the same example participant (**Figure 4a**), and at the group level (**Figure**
4b). Confirming the broadening of the attentional field observed in the visual field
reconstruction maps, we found that the estimated FWHM increased with greater cue
widths in V2 and V3 (V2 t(1,30)=4.60, p<.001; V3 t(1,30)=6.71, p<.001). The effect was
not statistically significant in V1 (t(1,30)=1.61, p=.118).

Finally, we assessed the gain of the attentional modulation in the model (**Figure**
4a and **4b** for the example participant and group data, respectively). We observed no
significant relationship between amplitude and cue width (V1 t(1,30)=-.18, p=.861; V2
t(1,30)=-.42, p=.677; V3 t(1,30)=-1.00, p=.325). We also found that the overall gain was
greater in V2 and V3 compared to V1 (paired t-test, both p<=.01).

Temporal interval analysis

In the previous analyses, we leveraged the fact that the attentional cue remained
constant for 5-trial blocks (spatial profiles were computed by averaging BOLD
measurements across a block of 10 TRs). We next examined the degree to which we
were able to recover the attentional field on a moment-by-moment (TR-by-TR) basis. To
examine the consistency of the attentional field over a varying number of TRs with an
identical cue, we systematically adjusted the number of TRs that contributed to the
averaged spatial response profile. To maintain a constant number of observations across
the temporal interval conditions, we randomly sampled a subset of TRs from each block.

When we systematically varied the number of TRs included for each model fit (1,
2, 3, 5, or 10 TRs), we found a significantly positive linear relationship between cue width
and recovered FWHM when averaging two or more TRs in V3 (all p<=.008), and five or
more TRs in V2 (both p<=.034; **Figure 5a**). As described above, V1 did not reliably show
a monotonic relationship between cue width and FWHM, even when averaging ten TRs.
We did not find that varying the number of TRs systematically altered the FWHM
estimates in V1 or V3 (V1, t(1,18)=1.70, p=.106; V3, t(1,18)=.85, p=.405), although a
significant effect was observed in V2 such that larger numbers of TRs were associated
with greater width estimates (t(1,18)=2.39, p=.028).

The number of TRs significantly affected the absolute angular error associated
with the estimated location of the attentional field **(Figure 5b)**. Error magnitude
decreased with TRs in all three visual regions (V1, t(1,18)=-4.40, p<.001; V2, t(1,18)=-
5.15, p<.001; V3, t(1,18)=-4.10, p<.001), suggesting that more data yielded more
accurate estimates, though absolute angular error remained consistently below chance
(90°) even when fitting the model to single-TR BOLD responses. Location error remained
stable across width conditions (V1, t(1,18)=-.38, p=.706; V2, t(1,18)=.66, p=.520; V3,
t(1,18)=1.29, p=.215).

Figure 5. Effect of number of TRs. Model fits were computed using BOLD data averaged across different temporal intervals (1, 2, 3, 5, or 10 TRs). Group means (with SEM) are plotted for FWHM, absolute angular error, amplitude estimates, and R^2, separated by cue width, brain region, and the number of TRs used for each model fit.

The estimated gain of the attentional enhancement showed a dependence on
number of TRs, with more TRs associated with lower gain estimates (V1, $t(1,18)=-6.67$,
$p<.001$; V2, $t(1,18)=-6.03$, $p<.001$; V3, $t(1,18)=-5.74$, $p<.001$), with no evident
dependence on cue width (V1 $t(1,18)=-.03$, $p=.976$; V2 $t(1,18)=-.24$, $p=.810$; V3 $t(1,18)=-$
$.28$, $p=.782$; **Figure 5c**).
Finally, the model's goodness of fit improved with more data, with larger R^2
associated with greater numbers of TRs included in the average profiles (V1,
$t(1,18)=5.82$, $p<.001$; V2, $t(1,18)=12.89$, $p<.001$; V3, $t(1,18)=8.30$, $p<.001$), though all R^2
were above 0.3 across all visual regions even for single-TR model fits. We did not

observe a dependence of R² on cue width (V1, $t(1,18)=.23$, $p=.823$; V2, $t(1,18)=.46$,
$p=.653$; V3, $t(1,18)=.05$, $p=.959$; **Figure 5d**).

Width of the attentional field mimics perceptual modulation

While the attentional field broadened as expected when participants were cued
to attend to a larger portion of the white noise annulus, the size of the estimated
attentional modulation was greater than the true size of the cued region. The cue width
varied between 18° and 162°, whereas the width estimate derived from spatial profiles
of BOLD modulation varied between 103° and 179° (**Figure 4b**). We wondered what the
underlying cause of this disparity might be. One possibility is that the BOLD-derived
FWHM might tend to overestimate the retinotopic extent of the modulation. If this were
the case, we would expect to obtain overestimates of FWHM when applying the same
modeling approach to perceptual modulations as well. Alternatively, the true subjective
attentional field might be consistently broader than cued, despite the presence of nearby
distractors. If this were the case, modulation driven by perceptual differences should *not*
result in the same large FWHM estimates.

To address this, we compared our estimates of the attentional field with
equivalent estimates for spatial profiles induced by a perceptual manipulation. In this
additional experiment, we varied the contrast intensity of sections of the white noise
annulus. Participants were not asked to deploy spatial attention to the stimulus and were
instead instructed to perform a color change detection task at fixation. The regions of
increased noise contrast matched the attentional cue widths (18°, 54°, 90°, and 162°,
plus an additional intermediate width of 126º), and were centered on one of the four
cardinal locations (0°, 90°, 180°, 270° polar angle).

As expected, we observed a broadening of the spatial profile of BOLD modulation
in all three visual areas as the region of increased contrast widened (**Figure 6a**). Using
an identical modeling procedure, we estimated the spatial profile of the *perceptual* BOLD
modulation. The group results for model estimates revealed that: 1) we were highly
accurate in estimating the location of the contrast increment; 2) FWHM of the spatial
profiles broadened across contrast widths, and 3) the amplitude remained stable across
contrast widths (**Figure 6b**).

Mirroring the results from the attentional manipulation, FWHM estimates
systematically exceeded the nominal size of the perceptually modulated region of the
visual field. Comparing the estimated FWHMs of the perceptual and attentional spatial
profiles (**Figure 6c**) revealed that the estimated widths were highly comparable (Pearson
correlation r=0.664 across width conditions and visual regions). This finding implies that
the BOLD-derived generative Gaussian model may have had a general tendency to
return upwardly biased width estimates, but that it recovered relative differences in a
similar manner for attentional and perceptual forms of modulation.

For the perceptual contrast manipulation, the increase in the recovered FWHM
with contrast width was observed in both V1 and V3 (**Figure 6b**; V1, $t(1,23)=4.59$, $p<.001$;
V3, $t(1,23)=4.96$, $p<.001$), though this effect was not clearly observed in V2 ($t(1,23)=1.22$,
$p=.236$). The mean magnitude of angular error between the model-estimated location
and the center of the contrast stimulus did not depend linearly on contrast width in any
of the three visual areas (all $p>=.584$). The estimated amplitude of modulation also did

not show a relationship to contrast width in any of the visual areas (V1, $t(1,23)=1.19$,
$p=.245$; V2, $t(1,23)=-.32$, $p=.749$; V3, $t(1,23)=.28$, $p=.783$).
160

a. Spatial profile contrast modulation

b. Group results

c. FWHM comparison

Figure 6. a. Spatial profiles of perceptual modulation. Solid lines represent the group mean BOLD activity and shaded regions the SEM. **b.** Group level parameter estimates. Overall group mean and standard error are shown for the absolute angular error, FWHM, and amplitude, separated by contrast width and brain region. **c.** Comparison of FWHM estimates obtained from the attentional manipulation and the physical contrast manipulation. Dot color indicates brain region; each point represents the mean FWHM for a given width condition across participants.

1 Discussion

We investigated the topographic spread of spatial attention in human visual cortex
by characterizing the spatial profile of BOLD responses while participants attended to
different portions of the visual field. Behavioral performance confirmed that participants
used the fixation cue to dynamically allocate attention to different swaths of the visual
field. Attention allocation was associated with a boost in the BOLD response in
corresponding retinotopic areas of visual cortex. To characterize the topography of that
boost, our approach involved selecting voxels with pRF preferred eccentricities that
overlapped our white noise annulus, and organizing those voxels into one-dimensional
profiles of attentional modulation as a function of preferred polar angle. This allowed us
to model the location and spread of the attentional field and test how well it tracked the
nominal location and width of the cue presented at fixation. Using a generalized
Gaussian model, the cued location could be recovered with high fidelity. Furthermore,
we observed a broadening of the estimated attentional field in areas V2 and V3 with the
cue width, suggesting our method was capable of dynamically recovering the location
and size of the attentional field from moment to moment.
This work builds on the concept of an attentional 'spotlight' or 'zoom lens' that
has long been theorized to aid in spatial attention (Shaw and Shaw, 1977; Posner, 1980;
Eriksen and St. James, 1986; Carrasco, 2011). By flexibly adjusting and shifting the focus
of the spotlight, visual representations are selectivity enhanced within a specific region
of the visual field. However, the empirical evidence demonstrating that attention can
change its *spread* across the visual field by modulating brain responses is surprisingly
lacking (Yeshurun, 2019). Our understanding of how the attentional window interacts
with spatial representations is mainly based on behavioral reports (Gobell, Tseng and
Sperling, 2004; Palmer and Moore, 2009; Herrmann *et al.*, 2010; Beilen *et al.*, 2011;
Taylor *et al.*, 2015; Huang *et al.*, 2017; Kınıklıoğlu and Boyaci, 2022), but see (Hopf *et al.*,
2006; Itthipuripat *et al.*, 2014; Tkacz-Domb and Yeshurun, 2018; Feldmann-Wüstefeld
and Awh, 2020), despite it forming a crucial component in an influential theoretical model
of attention (Reynolds and Heeger, 2009). This model proposes that the interaction
between stimulus properties (such as its size and specific features) and the attentional
field can explain a wide variety of attentional effects reported in behavioral and
neurophysiological studies (Herrmann *et al.*, 2010; Itthipuripat *et al.*, 2014; Bloem and
Ling, 2019; Jigo, Heeger and Carrasco, 2021). The present study sought to address this
gap, with our results showing that the visuocortical attentional field broadened as we
increased the cue width (**Figure 4**). This provides compelling evidence that the attention-
related cortical response can, in fact, flexibly vary in its position and spatial distribution.
In this study, we modeled the attentional field using a one-dimensional
distribution. This approach aligned with our experimental design, as the attentional cue
was manipulated only as a function of polar angle. However, we know that spatial
processing varies substantially as a function of eccentricity. Spatial resolution is highest
at the fovea and rapidly drops in the periphery (Anton-Erxleben and Carrasco, 2013). The
spatial distribution of attention will presumably also vary with eccentricity and will likely

take on different functional properties close to the fovea, where spatial resolution is high,
compared to the far periphery where spatial resolution is low (Intriligator and Cavanagh,
2001; Jigo, Heeger and Carrasco, 2021). Future work can help provide a better
understanding of the contribution of spatial attention by considering how the attentional
field interacts with these well described spatial variations across the visual field.
Measuring the full spatial distribution of the attentional field (across both eccentricity and
polar angle) will shed light on how spatial attention guides perception by interacting with
the non-uniformity of spatial representations.

The spread of the attentional field likely influences the degree to which spatial
resolution at the attended location is transformed, leading to enhanced behavioral
performance. In our experiment, we cued participant to varying swaths of an iso-
eccentric annulus of white noise and participants had to discriminate whether more
numbers or more letters were presented within the cued region. Spatial attention was
vital for this task, as enhanced spatial perception allowed the participants to better
discriminate all stimuli within the cued region (Anton-Erxleben and Carrasco, 2013).
However, the estimated spatial spread of the attentional modulation (as indicated by the
recovered FWHM) was consistently wider than the cued region itself. We therefore
compared the spread of the attention field with the spatial profile of a *perceptually*
induced width manipulation. Our model overestimated the retinotopic extent of the cued
region in both the attentional and perceptual versions of the task (**Figure 6c**), suggesting
that the BOLD-derived FWHM systematically overestimated the extent of modulation.
Future work could unpack the degree to which the size of the attentional field influences
the spatial resolution of visual cortical representations (Klein, Harvey and Dumoulin,
2014; Vo, Sprague and Serences, 2017; Tünçok, Carrasco and Winawer, 2024), and how
this influences spatial perception.

Beyond addressing core questions related to the function of spatial attention, this
method also lays groundwork for addressing questions about spatial predictive
uncertainty and belief updating. Prior work on these topics has relied almost entirely on
inferring participants' predictions from their behavior, often requiring participants to
report overt point predictions (Nassar *et al.*, 2010; McGuire *et al.*, 2014; D'Acremont and
Bossaerts, 2016; Nassar, Bruckner and Frank, 2019), or inferring participants'
predictions from their sequences of decisions (Daw *et al.*, 2006; Behrens *et al.*, 2007;
Payzan-LeNestour and Bossaerts, 2011; Payzan-LeNestour *et al.*, 2013). These
approaches have shed light on how we dynamically adapt our learning and belief
updating processes over time in differently structured contexts. However, methods for
recovering information about full predictive belief distributions have been limited, relying
on indirect measurements such as eye movements (O'Reilly *et al.*, 2013; Bakst and
McGuire, 2021, 2023), and physiological measures of uncertainty and surprise in EEG
and pupillometry (Preuschoff, 't Hart and Einhauser, 2011; Nassar *et al.*, 2012; Nassar,
Bruckner and Frank, 2019). The methods developed here offer a potential way to recover
the location and width of a spatial predictive distribution via the attentional field in
contexts in which it is unknown *a priori* and might be dependent on how a given
participant has integrated previous sequential evidence. Future work could extend this
method to more directly interrogate how predictive uncertainty is represented
throughout the brain on a moment-by-moment basis.

In summary, we found evidence that people could dynamically adapt the spread
of spatial attention, and that the retinotopic extent of attentional enhancement of the
BOLD response reflected this dynamic adaptation. These findings address a gap in our
understanding of spatial attentional control, supporting core theoretical models of
attention. Our modeling approach also lays the groundwork to address further questions
related to how the attentional field interacts with the non-uniformity of spatial
representations and how uncertainty in spatial contexts is represented in the human
brain.

References

1) Andersson, J.L.R., Skare, S. and Ashburner, J. (2003) 'How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging', *NeuroImage*, 20(2), pp. 870–888. Available at: https://doi.org/10.1016/S1053-8119(03)00336-7.

2) Anton-Erxleben, K. and Carrasco, M. (2013) 'Attentional enhancement of spatial resolution: linking behavioural and neurophysiological evidence', *Nature Reviews Neuroscience*, 14(3), pp. 188–200. Available at: https://doi.org/10.1038/nrn3443.

3) Bakst, L. and McGuire, J.T. (2021) 'Eye movements reflect adaptive predictions and predictive precision', *Journal of Experimental Psychology: General*, 150(5), pp. 915–929. Available at: https://doi.org/10.1037/xge0000977.

4) Bakst, L. and McGuire, J.T. (2023) 'Experience-driven recalibration of learning from surprising events', *Cognition*, 232, p. 105343. Available at: https://doi.org/10.1016/j.cognition.2022.105343.

5) Behrens, T.E.J. *et al.* (2007) 'Learning the value of information in an uncertain world', *Nature Neuroscience*, 10(9), pp. 1214–1221. Available at: https://doi.org/10.1038/nn1954.

6) Beilen, M. van *et al.* (2011) 'Attentional Window Set by Expected Relevance of Environmental Signals', *PLOS ONE*, 6(6), p. e21262. Available at: https://doi.org/10.1371/journal.pone.0021262.

7) Benson, N.C. *et al.* (2018) 'The Human Connectome Project 7 Tesla retinotopy dataset: Description and population receptive field analysis', *Journal of Vision*, 18(13), p. 23. Available at: https://doi.org/10.1167/18.13.23.

8) Bloem, I.M. and Ling, S. (2019) 'Normalization governs attentional modulation within human visual cortex', *Nature Communications*, 10(1), p. 5660. Available at: https://doi.org/10.1038/s41467-019-13597-1.

9) Brefczynski, J.A. and DeYoe, E.A. (1999) 'A physiological correlate of the "spotlight" of visual attention', *Nature Neuroscience*, 2(4), pp. 370–374. Available at: https://doi.org/10.1038/7280.

10) Carrasco, M. (2011) 'Visual attention: The past 25 years', *Vision Research*, 51(13), pp. 1484–1525. Available at: https://doi.org/10.1016/j.visres.2011.04.012.

11) Castiello, U. and Umiltà, C. (1990) 'Size of the attentional focus and efficiency of processing', *Acta Psychologica*, 73(3), pp. 195–209. Available at: https://doi.org/10.1016/0001-6918(90)90022-8.

12) D'Acremont, M. and Bossaerts, P. (2016) 'Neural Mechanisms Behind Identification of Leptokurtic Noise and Adaptive Behavioral Response', *Cerebral Cortex*, 26(4), pp. 1818–1830. Available at: https://doi.org/10.1093/cercor/bhw013.

13) Datta, R. and DeYoe, E.A. (2009) 'I know where you are secretly attending! The
topography of human visual attention revealed with fMRI', *Vision Research*, 49(10),
pp. 1037–1044. Available at: https://doi.org/10.1016/j.visres.2009.01.014.

14) Daw, N.D. *et al.* (2006) 'Cortical substrates for exploratory decisions in humans',
Nature, 441(7095), pp. 876–879. Available at: https://doi.org/10.1038/nature04766.

15) Dumoulin, S.O. and Wandell, B.A. (2008) 'Population receptive field estimates in
human visual cortex', *NeuroImage*, 39(2), pp. 647–660. Available at:
https://doi.org/10.1016/j.neuroimage.2007.09.034.

16) Eriksen, C.W. and St. James, J.D. (1986) 'Visual attention within and around the field
of focal attention: A zoom lens model', *Perception & Psychophysics*, 40(4), pp. 225–
240. Available at: https://doi.org/10.3758/BF03211502.

17) Favila, S.E., Kuhl, B.A. and Winawer, J. (2022) 'Perception and memory have distinct
spatial tuning properties in human visual cortex', *Nature Communications*, 13(1), p.
5864. Available at: https://doi.org/10.1038/s41467-022-33161-8.

18) Feldmann-Wüstefeld, T. and Awh, E. (2020) 'Alpha-band Activity Tracks the Zoom
Lens of Attention', *Journal of Cognitive Neuroscience*, 32(2), pp. 272–282. Available
at: https://doi.org/10.1162/jocn_a_01484.

19) Fischl, B. (2012) 'FreeSurfer', *NeuroImage*, 62(2), pp. 774–781. Available at:
https://doi.org/10.1016/j.neuroimage.2012.01.021.

20) Gobell, J.L., Tseng, C. and Sperling, G. (2004) 'The spatial distribution of visual
attention', *Vision Research*, 44(12), pp. 1273–1296. Available at:
https://doi.org/10.1016/j.visres.2004.01.012.

21) Greve, D.N. and Fischl, B. (2009) 'Accurate and robust brain image alignment using
boundary-based registration', *NeuroImage*, 48(1), pp. 63–72. Available at:
https://doi.org/10.1016/j.neuroimage.2009.06.060.

22) Herrmann, K. *et al.* (2010) 'When size matters: attention affects performance by
contrast or response gain', *Nature Neuroscience*, 13(12), pp. 1554–1559. Available
at: https://doi.org/10.1038/nn.2669.

23) Hopf, J.-M. *et al.* (2006) 'The Neural Site of Attention Matches the Spatial Scale of
Perception', *Journal of Neuroscience*, 26(13), pp. 3532–3540. Available at:
https://doi.org/10.1523/JNEUROSCI.4510-05.2006.

24) Huang, D. *et al.* (2017) 'The time course of attention modulation elicited by spatial
uncertainty', *Vision Research*, 138, pp. 50–58. Available at:
https://doi.org/10.1016/j.visres.2017.06.008.

25) Intriligator, J. and Cavanagh, P. (2001) 'The Spatial Resolution of Visual Attention',
Cognitive Psychology, 43(3), pp. 171–216. Available at:
https://doi.org/10.1006/cogp.2001.0755.

26) Itthipuripat, S. *et al.* (2014) 'Changing the Spatial Scope of Attention Alters Patterns
of Neural Gain in Human Cortex', *Journal of Neuroscience*, 34(1), pp. 112–123.
Available at: https://doi.org/10.1523/JNEUROSCI.3943-13.2014.

27) Jigo, M., Heeger, D.J. and Carrasco, M. (2021) 'An image-computable model of how
endogenous and exogenous attention differentially alter visual perception',
Proceedings of the National Academy of Sciences, 118(33), p. e2106436118.
Available at: https://doi.org/10.1073/pnas.2106436118.

28) Kastner, S. *et al.* (1998) 'Mechanisms of Directed Attention in the Human Extrastriate
Cortex as Revealed by Functional MRI', *Science*, 282(5386), pp. 108–111. Available
at: https://doi.org/10.1126/science.282.5386.108.

29) Kay, K.N. *et al.* (2013) 'Compressive spatial summation in human visual cortex',
Journal of Neurophysiology, 110(2), pp. 481–494. Available at:
https://doi.org/10.1152/jn.00105.2013.

30) Kınıklıoğlu, M. and Boyaci, H. (2022) 'Increasing the spatial extent of attention
strengthens surround suppression', *Vision Research*, 199, p. 108074. Available at:
https://doi.org/10.1016/j.visres.2022.108074.

31) Klein, B.P., Harvey, B.M. and Dumoulin, S.O. (2014) 'Attraction of Position Preference
by Spatial Attention throughout Human Visual Cortex', *Neuron*, 84(1), pp. 227–237.
Available at: https://doi.org/10.1016/j.neuron.2014.08.047.

32) Maunsell, J.H.R. (2015) 'Neuronal Mechanisms of Visual Attention', *Annual Review of
Vision Science*, 1(Volume 1, 2015), pp. 373–391. Available at:
https://doi.org/10.1146/annurev-vision-082114-035431.

33) McAdams, C.J. and Maunsell, J.H. (1999) 'Effects of attention on orientation-tuning
functions of single neurons in macaque cortical area V4', *The Journal of
Neuroscience: The Official Journal of the Society for Neuroscience*, 19(1), pp. 431–
441. Available at: https://doi.org/10.1523/JNEUROSCI.19-01-00431.1999.

34) McGuire, J.T. *et al.* (2014) 'Functionally Dissociable Influences on Learning Rate in a
Dynamic Environment', *Neuron*, 84(4), pp. 870–881. Available at:
https://doi.org/10.1016/j.neuron.2014.10.013.

35) McMains, S.A. and Somers, D.C. (2004) 'Multiple Spotlights of Attentional Selection
in Human Visual Cortex', *Neuron*, 42(4), pp. 677–686. Available at:
https://doi.org/10.1016/S0896-6273(04)00263-6.

36) Moeller, S. *et al.* (2010) 'Multiband multislice GE-EPI at 7 tesla, with 16-fold
acceleration using partial parallel imaging with application to high spatial and
temporal whole-brain fMRI', *Magnetic Resonance in Medicine*, 63(5), pp. 1144–1153.
Available at: https://doi.org/10.1002/mrm.22361.

37) Müller, N.G. *et al.* (2003) 'A Physiological Correlate of the "Zoom Lens" of Visual
Attention', *Journal of Neuroscience*, 23(9), pp. 3561–3565. Available at:
https://doi.org/10.1523/JNEUROSCI.23-09-03561.2003.

38) Nadarajah, S. (2005) 'A generalized normal distribution', *Journal of Applied Statistics*,
32(7), pp. 685–694. Available at: https://doi.org/10.1080/02664760500079464.

39) Nassar, M.R. *et al.* (2010) 'An Approximately Bayesian Delta-Rule Model Explains the
Dynamics of Belief Updating in a Changing Environment', *Journal of Neuroscience*,
30(37), pp. 12366–12378. Available at: https://doi.org/10.1523/JNEUROSCI.0822-
10.2010.

40) Nassar, M.R. *et al.* (2012) 'Rational regulation of learning dynamics by pupil-linked
arousal systems', *Nature Neuroscience*, 15(7), pp. 1040–1046. Available at:
https://doi.org/10.1038/nn.3130.

41) Nassar, M.R., Bruckner, R. and Frank, M.J. (2019) 'Statistical context dictates the
relationship between feedback-related EEG signals and learning', *eLife*. Edited by
T.H. Donner et al., 8, p. e46975. Available at: https://doi.org/10.7554/eLife.46975.

42) O'Reilly, J.X. *et al.* (2013) 'Dissociable effects of surprise and model update in parietal
and anterior cingulate cortex', *Proceedings of the National Academy of Sciences*,
110(38), pp. E3660–E3669. Available at: https://doi.org/10.1073/pnas.1305373110.

43) Palmer, J. and Moore, C.M. (2009) 'Using a filtering task to measure the spatial extent
of selective attention', *Vision Research*, 49(10), pp. 1045–1064. Available at:
https://doi.org/10.1016/j.visres.2008.02.022.

44) Payzan-LeNestour, E. *et al.* (2013) 'The Neural Representation of Unexpected
Uncertainty during Value-Based Decision Making', *Neuron*, 79(1), pp. 191–201.
Available at: https://doi.org/10.1016/j.neuron.2013.04.037.

45) Payzan-LeNestour, E. and Bossaerts, P. (2011) 'Risk, Unexpected Uncertainty, and
Estimation Uncertainty: Bayesian Learning in Unstable Settings', *PLOS*
Computational Biology, 7(1), p. e1001048. Available at:
https://doi.org/10.1371/journal.pcbi.1001048.

46) Posner, M.I. (1980) 'Orienting of Attention', *Quarterly Journal of Experimental*
Psychology, 32(1), pp. 3–25. Available at:
https://doi.org/10.1080/00335558008248231.

47) Preuschoff, K., 't Hart, B.M. and Einhauser, W. (2011) 'Pupil Dilation Signals Surprise:
Evidence for Noradrenaline's Role in Decision Making', *Frontiers in Neuroscience*, 5.
Available at: https://doi.org/10.3389/fnins.2011.00115.

48) Puckett, A.M. and DeYoe, E.A. (2015) 'The Attentional Field Revealed by Single-Voxel
Modeling of fMRI Time Courses', *Journal of Neuroscience*, 35(12), pp. 5030–5042.
Available at: https://doi.org/10.1523/JNEUROSCI.3754-14.2015.

49) Reuter, M., Rosas, H.D. and Fischl, B. (2010) 'Highly accurate inverse consistent
registration: A robust approach', *NeuroImage*, 53(4), pp. 1181–1196. Available at:
https://doi.org/10.1016/j.neuroimage.2010.07.020.

50) Reynolds, J.H. and Heeger, D.J. (2009) 'The Normalization Model of Attention',
Neuron, 61(2), pp. 168–185. Available at:
https://doi.org/10.1016/j.neuron.2009.01.002.

51) Samaha, J., Sprague, T.C. and Postle, B.R. (2016) 'Decoding and Reconstructing the
Focus of Spatial Attention from the Topography of Alpha-band Oscillations', *Journal*
of cognitive neuroscience, 28(8), pp. 1090–1097. Available at:
https://doi.org/10.1162/jocn_a_00955.

52) Setsompop, K. *et al.* (2012) 'Blipped-controlled aliasing in parallel imaging for
simultaneous multislice echo planar imaging with reduced g-factor penalty',
Magnetic Resonance in Medicine, 67(5), pp. 1210–1224. Available at:
https://doi.org/10.1002/mrm.23097.

53) Shaw, M.L. and Shaw, P. (1977) 'Optimal allocation of cognitive resources to spatial
locations', *Journal of Experimental Psychology: Human Perception and Performance*,
3(2), pp. 201–211. Available at: https://doi.org/10.1037/0096-1523.3.2.201.

54) Shioiri, S. *et al.* (2016) 'Visual attention spreads broadly but selects information
locally', *Scientific Reports*, 6(1), p. 35513. Available at:
https://doi.org/10.1038/srep35513.

55) Smith, S.M. *et al.* (2004) 'Advances in functional and structural MR image analysis
and implementation as FSL', *NeuroImage*, 23, pp. S208–S219. Available at:
https://doi.org/10.1016/j.neuroimage.2004.07.051.

56) Sprague, T.C. and Serences, J.T. (2013) 'Attention modulates spatial priority maps in
the human occipital, parietal and frontal cortices', *Nature Neuroscience*, 16(12), pp.
1879–1887. Available at: https://doi.org/10.1038/nn.3574.

57) Taylor, J.E.T. *et al.* (2015) 'Attentional cartography: mapping the distribution of
attention across time and space', *Attention, Perception, & Psychophysics*, 77(7), pp.
2240–2246. Available at: https://doi.org/10.3758/s13414-015-0943-0.

58) Tkacz-Domb, S. and Yeshurun, Y. (2018) 'The size of the attentional window when
measured by the pupillary response to light', *Scientific Reports*, 8(1), p. 11878.
Available at: https://doi.org/10.1038/s41598-018-30343-7.

59) Tünçok, E., Carrasco, M. and Winawer, J. (2024) 'Spatial attention alters visual
cortical representation during target anticipation'. bioRxiv, p. 2024.03.02.583127.
Available at: https://doi.org/10.1101/2024.03.02.583127.

60) Vo, V.A., Sprague, T.C. and Serences, J.T. (2017) 'Spatial Tuning Shifts Increase the
Discriminability and Fidelity of Population Codes in Visual Cortex', *Journal of*
Neuroscience, 37(12), pp. 3386–3401. Available at:
https://doi.org/10.1523/JNEUROSCI.3484-16.2017.

61) Yeshurun, Y. (2019) 'The spatial distribution of attention', *Current Opinion in*
Psychology, 29, pp. 76–81. Available at:
https://doi.org/10.1016/j.copsyc.2018.12.008.

a. Anisotropic 2D BOLD activity reconstruction

b. Spatial profile attentional modulation

a. Generalized Gaussian model   *b.* Quantify attentional field

BOLD response (a.u.)

Cued polar angle (°)

BOLD response (a.u.)

Cued polar angle (°)

a. Example observer

Location — Width — Amplitude

b. Group results

Location — Width — Amplitude

V1 **V2** **V3**

a. Attentional field width

FWHM (°)

b. Attentional field angular error

Absolute
angular error (°)

c. Attentional gain

Amplitude
(%SC)

d. Goodness of fit

R²

BOLD response
(% signal change)

V1 V2 V3

Polar angle relative to center location (°)

b. *Group results*

Location

Angular error (°)

Contrast width

Width

FWHM (°)

Contrast width

Amplitude

BOLD (%SD)

18° 54° 90°126°162°

Contrast width

c. *FWHM comparison*

FWHM
Contrast modulation (°)

FWHM
Attention field (°)