1 **Genomic characterization of foodborne *Salmonella enterica* and *Escherichia coli* isolates from**
2 **Saboba district and Bolgatanga Municipality Ghana**
3

4 Gabriel Temitope Sunmonu [1], Frederick Adzitey [2], Erkison Ewomazino Odih [1], Boniface Awini Tibile [2],
5 Rejoice Ekli [2], Martin Aduah [2], Anderson O Oaikhena [1], Olabisi C Akinlabi [1], Akebe Luther King Abia [3],
6 Daniel Gyamfi Amoako [3, 4], and Iruka N Okeke [1]
7
8 1. Department of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Ibadan, Nigeria.
9 2. Faculty of Agriculture, Food and Consumer Sciences, University for Development Studies, Tamale, Ghana
10 3. Antimicrobial Research Unit, College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa
11 4. Department of Integrative Biology and Bioinformatics, University of Guelph, Ontario, Canada
12

Abstract

Salmonella enterica and *Escherichia coli* are well-known bacteria commonly associated with
foodborne illnesses in humans and animals. Genomic characterization of these pathogens
provides valuable insights into their evolution, virulence factors and resistance determinants.
This study aimed to characterized previously isolated *Salmonella* (n = 14) and *E. coli* (n = 19)
from milk, meat and its associated utensils in Ghana using whole-genome sequencing. Most of
the *Salmonella* serovars (Fresno, Plymouth, Infantis, Give and Orleans) identified in this study
are yet to be reported in Ghana. Most *Salmonella* isolates were pan-sensitive, but genes
conferring resistance to fosfomycin (*fosA7.2*) and tetracycline (*tet(A)*) were detected in one and
three isolates, respectively. Seven of the *Salmonella* isolates carry the IncI1-I(Gamma) plasmid
replicon. Although antimicrobial resistance was not common among *Salmonella* strains, most
(11/19) of the *E. coli* strains had at least one resistance gene, with nearly half (8/19) being
multidrug resistant and carrying mobile elements. Three of the 19 *E. coli* strains belong to
serovars commonly associated with enteroaggregative *E. coli* (EAEC) pathotype. While strains
belonging to virulence-associated lineages lacked key plasmid-encoded virulence plasmids,
several plasmid replicons were detected in most of the *E. coli* (14/19) strains. Food
contaminated with these pathogens can serve as a vehicle for disease transmission, posing a
significant public health risk and necessitating stringent food safety and hygiene practices to
prevent outbreaks. Hence, there is need for continuous surveillance and preventive measures
to stop the spread of foodborne pathogens and reduce the risk of associated illnesses in Ghana.
33

INTRODUCTION
Milk and meat are essential protein sources that constitute a significant and nutrient-rich
component of human diets. However, their consumption is often associated with foodborne
infections (1), particularly those caused by *Salmonella* and *Escherichia coli* (2-5). Salmonellosis
is under-reported in Ghana, and only a few studies have investigated the plausible role of
contaminated milk (6), meat, meat products, handlers' hands and associated surfaces such as
knives, tables and aprons (7-9) in facilitating their transmission. Knowledge of food safety
practices by key food handlers in Ghana has recently been reported to be suboptimal, as have
food safety infrastructure and regulatory enforcement (10, 11). There have been a few reports
of meat samples contaminated with *E. coli* in Ghana (11-14). Similarly, *E. coli* has been
recovered from milk, milking utensils, faeces of lactating cow and milkers' hands (15). Some of

these *E. coli* strains harbour both virulence and antimicrobial resistance genes (ARGs), raising
public health concerns. However, few of these strains have been thoroughly characterized.
47
There are public health and food safety implications of finding *Salmonella* and *E. coli* in food
because they are invariably of faecal origin. Globally, in addition to contamination risks,
Salmonella and *E. coli* sourced from milk and meat increasingly exhibit resistance to different
classes of antibiotics that is commonly mediated by mobile elements (13, 16-18). The
prevalence of multidrug-resistant (MDR) *Salmonella* and *E. coli* is also on the rise in clinical
infections (19, 20). Notably, resistance to extended spectrum beta-lactams,
trimethoprim/sulfamethoxazole, chloramphenicol and ciprofloxacin has been reported, often
associated with plasmids that could mediate their spread, in both *Salmonella* and *E. coli* isolates
from milk and retail meats in Ghana (6, 12, 14).
57
Various methods, including serotyping, antibiotic profiling, pulsed-field gel electrophoresis and
whole genome sequencing, have been employed to elucidate the phenotypic and genotypic
attributes of foodborne pathogens and to determine their interrelationships and connections to
pandemic clones of interest (13, 21). Next-generation sequencing (NGS) technology, the most
versatile and informative approach, has gained recent prominence (22, 23). NGS is now used by
PulseNet to categorize foodborne diseases, enabling nuanced epidemiological investigations
(24). Identification of virulence factors, antibiotic resistance genes, and serotypes are all
possible by genomic analysis, which can also provide enhanced information on strain inter-
relatedness, and therefore enable source attribution. Data on the genomic characterization of
Salmonella and *E. coli* from milk and meat are few from low- and middle-income countries,
including Ghana (7, 25). In light of this gap, this study aims to characterize the resistance,
virulence and plasmid profile of previously isolated *Salmonella* and *E. coli* isolated from fresh
different retail meats, milk, and associated samples (handler's hand swab, table, knife and
faecal samples) in Saboba district and Bolgatanga Municipality of Ghana.
72
METHODS
Strains
A total of 33 bacterial isolates (14 *Salmonella* and 19 *E. coli* species) previously isolated from
various fresh and ready-to-eat meats, meat sellers' tables, milk, milk-collecting utensils, milkers'
hands and faeces of lactating cows were characterized for this study (Table 1) (8, 15, 26, 27).
The isolates originated from markets and farms in Bolgatanga Municipality and Saboba District
in Northern Ghana and were cryopreserved in 50% glycerol in Luria broth at -80^{o}C.
80
***Salmonella* and *E. coli* identification**
Salmonella isolates were initially confirmed using a latex agglutination kit for *Salmonella* (Oxoid
Limited, Basingstoke, UK) and by PCR targeting the *invA* gene as described by Rahn et al. (1992)
(28), using PCR oligonucleotides *invA139f* GTGAAATTATCGCCACGTTCGGGCAA and *invA141r*
TCATCGCACCGTCAAGGAACC. PCR was performed using PuRe Taq Ready-To-Go PCR Beads
(illustra^{TM}). The PCR cycle used an initial denaturation temperature of 95^{0}C for two minutes,
followed by 35 cycles of denaturation at 95^{0}C for 30 seconds, annealing at 55^{0}C for 30 seconds
and extension at 72^{0}C for two minutes, then a terminal extension at 72^{0}C for five minutes.

Visualization of the 284 bp amplicon was accomplished after electrophoresis on 1.5% (w/v)
agarose gels stained with Gel red (biotium), using a UVP GelMax transilluminator and imager.
Salmonella isolates positive for *invA* and *E. coli* isolates were biotyped with the Gram-negative
(GN) test kit (Ref: 21341) on VITEK 2 systems (version 2.0, Marcy-l'Etoile, France, Biomérieux)
according to manufacturer's instructions
94
95 Table 1: *E. coli* and *Salmonella* isolates characterized in this study
96

Id	Code	Species	Sample	Reference
GH-FA-M23_S31	M2 3	*Salmonella enterica*	Milk	(27)
GH-FA-M87_S6	M8 1	*Salmonella enterica*	Milk	(27)
GH-FA-FS24_S23	FS24	*Salmonella enterica*	Faecal Sample (Milking cow)	(27)
GH-FA-M25_S2	M25	*Salmonella enterica*	Milk	(27)
GH-FA-US15_S32	US15-2	*Salmonella enterica*	Utensil sample	(27)
GH-FA-FK4_S10	FK4	*Salmonella enterica*	Knife (Fresh meat)	(8)
GH-FA-FK4d_S7	FK4d	*Salmonella enterica*	Knife (Fresh meat)	(8)
GH-FA-RCH2_S11	Rch2 extra	*Salmonella enterica*	RTE Chicken	(8)
GH-FA-RG2_S29	RG2	*Salmonella enterica*	RTE Guinea fowl	(8)
GH-FA-RK4_S19	RK4	*Salmonella enterica*	Knife (RTE utensil)	(8)
GH-FA-RM4_S20	RM4	*Salmonella enterica*	RTE Mutton	(8)
GH-FA-RP5_S5	RP5	*Salmonella enterica*	RTE Pork	(8)
GH-FA-RP5D_S11	RP5D	*Salmonella enterica*	RTE Pork	(8)
GH-FA-RPSD_S30	RPSD	*Salmonella enterica*	RTE Pork	(8)
GH-FA-FS23_S22	FS23	*Escherichia coli*	Faecal Sample (milking cow)	(27)
GH-FA-US24_S14	US24	*Escherichia coli*	Utensil Sample	(15)
GH-FA-HS11_S26	HS11	*Escherichia coli*	Hand swab	(15)
GH-FA-HS3_S27	HS3	*Escherichia coli*	Hand swab	(15)
GH-FA-M9_S16	M9	*Escherichia coli*	Milk	(15)
GH-FA-US3_S21	US3	*Escherichia coli*	Utensil sample	(15)
GH-FA-HS12_S21	HS12	*Escherichia coli*	Hand Swab	(15)
GH-FA-M25D_S22	M25	*Escherichia coli*	Milk	(15)
GH-FA-FB4_S24	FB4	*Escherichia coli*	Fresh Beef	(26)
GH-FA-FC1_S23	FC1	*Escherichia coli*	Fresh Chicken	(26)
GH-FA-FH2_S15	FH2	*Escherichia coli*	Hand Swab (Fresh meat vendor)	(26)
GH-FA-FGS_S16	FG5	*Escherichia coli*	Fresh Guinea fowl	(26)
GH-FA-FM4_S34	FM4	*Escherichia coli*	Fresh Mutton	(26)
GH-FA-FT1_S17	FT1	*Escherichia coli*	Table swab (Fresh meat)	(26)
GH-FA-RB4_S24	RB4	*Escherichia coli*	RTE Beef	(26)
GH-FA-RH4_S3	RH4	*Escherichia coli*	Hand swab (RTE meat Vendor)	(26)
GH-FA-RT3_S6	RT3	*Escherichia coli*	Table swab (RTE meat)	(26)

GH-FA-RT3_S29	RT3d	*Escherichia coli*	Table swab (RTE meat)	(26)
GH-FA-RCLI_S18	Rch1	*Escherichia coli*	RTE Chevon	(26)

Key: RTE: Ready-to-eat – meats sampled in prepared form

DNA Extraction, Library Preparation and Whole Genome Sequencing
Genomic DNA of the isolates was extracted using Wizard DNA extraction kit (Promega;
Wisconsin, USA) in accordance with manufacturer's protocol. Using a dsDNA Broad Range
quantification assay, the extracted DNA was quantified on a Qubit fluorometer (Invitrogen;
California, USA). dsDNA libraries were prepared using NEBNext Ultra II FS DNA library kit for
Illumina with 96-unique indexes (New England Biolabs, Massachusetts, USA; Cat. No: E6609L).
DNA libraries was quantified using dsDNA High Sensitivity quantification assay on a Qubit
fluorometer (Invitrogen; California, USA) and fragment length analysed with the Bioanalyzer
(Agilent). Denatured libraries were sequenced on an Illumina MiSeq (Illumina, California, USA).
The raw sequence reads were *de novo* assembled using SPAdes v3.15.3 (29) according to GHRU
protocols (https://gitlab.com/cgps/ghru/pipelines/dsl2/pipelines/assembly).

Sequence Typing of *Salmonella* and *E. coli* Genomes
Sequence reads were deposited in the *Salmonella* and *E. coli* database for *Salmonella* and *E. coli*
respectively on EnteroBase (30) and analyzed using publicly available tools that we have
previously validated (31, 32). Multi-locus sequence types (MLST) for the isolates were
determined using ARIBA (33). Novel ST strains were assigned ST using EnteroBase (30). The
Salmonella genome assemblies were analysed using the *Salmonella* In-Silico Typing Resource
(SISTR) for the prediction of serovars and serogroups (34) (https://github.com/phac-
nml/sistr_cmd), while the serotyping of the *E. coli* genome was done using ECtyper (35).

Identification of AMR, Plasmids and Virulence genes
PlasmidFinder (36) was utilized to identify plasmid replicons that were present in the
assembled genomes. AMRFinderPlus v3.10.24 (37) and its associated database (version 2022-
04-04.1) were used to predict the antimicrobial resistance genes carried by the isolates and the
drug classes to which they probably conferred resistance. Using ARIBA (33) and the virulence
factor database (VFDB), we were also able to identify the virulence genes that were present in
the isolates.

Single Nucleotide Polymorphism (SNP) Calling and Phylogenetic analysis
For phylogenetic analysis, reference sequences for the *Salmonella* and *E. coli* genomes were
objectively selected from the National Center for Biotechnology Information Reference
Sequence (RefSeq) database (https://www.ncbi.nlm.nih.gov/refseq/) using BactinspectorMax
v0.1.3 (https://gitlab.com/antunderwood/bactinspector). The selected references were the *S.*
enterica subsp. enterica serovar Fresno strain (assembly accession: GCF_003590695.1) and the
E. coli O25b:H4-ST131 strain (assembly accession: GCF_000285655.3). The sequence reads for
each species were then mapped to the chromosome of the reference using BWA (v0.7.17) (38)
and variants were called and filtered using bcftools (v1.9) (39) as implemented in the GHRU SNP

phylogeny pipeline (https://gitlab.com/cgps/ghru/pipelines/snp_phylogeny). Variant positions
were concatenated into a pseudoalignment and used to generate a maximum likelihood tree
using iqtree (v1.6.8) (40). SNP distances between the genome pairs were calculated using snp-
dists v.0.8.2 (https://github.com/tseemann/snp-dists).
142
143   **Data availability**
144   The sequence reads were deposited in the European Nucleotide Archive with the study
145   accession PRJEB58695 (Supplementary table 1).
146
RESULTS
Salmonella **Serotypes, Sequence Types (STs), Virulence Factors and Phylogeny**
We used SISTR software to predict the serovars of the 14 *Salmonella* strains characterized in
this study from whole genome sequence. The most common serotype was Fresno (n=6)
followed by Give (n=3), Orleans (n=2), Plymouth (n=1), Agona (n=1) and Infantis (n=1). The *S.*
Fresno and *S.* Orleans isolates were from previously unreported sequence types, now
designated ST10742 and ST10465 respectively (Supplementary table 2). As shown in Figure 1,
all the isolates from ready-to-eat pork, mutton and chicken belong to serovar Fresno. *S.* Fresno
isolates were also isolated from a meat vendor's knife, as was *S.* Orleans. The three milk
isolates, which were all from Saboba, belong to the serovars Plymouth (ST565), Give (ST516)
and Agona (ST13) (Figure 1). Two more ST516 *S.* Give isolates were recovered from the faeces
of milking cow and from a milking utensil.
159
160   Phylogenetic analysis of the 14 *Salmonella* isolates from this study showed that all *S.* Fresno
161   isolates, irrespective of source, clustered together and differed by < 3 SNPs. The two *S.* Orleans
162   isolates were identical (0 SNPs) and the three *S.* Give isolates were also identical, with the
163   isolates originating from milk, faeces of milking cow and from a milking utensil, also in Saboba.
164

Figure 1: Core genome SNP-based Maximum likelihood tree showing phylogenetic relationships among strains sequenced in this study. The matrix below the tree shows the sequence types, serotypes, location of isolation (Bolgatanga municipality or Saboba district) as well as the source/host sample and presence or absence of the *tetA* and *fosA* resistance genes.

All *Salmonella* isolates harboured curli (*csg*) genes as well as *bcf*, *fim* and *ste* fimbrial operons and ten of them, representing all serovars except *S.* Give and Plymouth, carried long polar fimbriae (*lpf)* genes. The *S.* Infantis and *S.* Agona strains carried *ratB* and *shdA*. Type III secretion system effector genes, such as: *inv, org, prg, sif, spa, sse, ssa* and *sop* were detected in all the isolates while *avr* was present in 57.1% (8/14) of the isolates and only one isolate harbours *gogB*. Four of the isolates also encode the cytholethal distending toxin gene, *cdtB* (Supplementary table 3).

Plasmid Replicons and ARG Profiles of *Salmonella*
The *Salmonella* isolates were largely pan-sensitive but genes conferring resistance to fosfomycin (*fosA7.2*) and tetracycline (*tet(A)*) were detected in one and three isolates respectively. Both *S.* Orleans isolates and one of the *S.* Fresno, from ready-to-eat mutton carried *tetA*, along with an IncI1-I(Gamma) plasmid replicon, which was also seen in six other, *tetA*-negative strains. Interestingly, the IncI1-I(Gamma) plasmid replicon was detected in all isolates from Bolgatanga municipality, irrespective of serovar, and no isolate from Saboba district harboured this plasmid. The fosfomycin resistance gene was found in the *S.* Agona genome, in which no plasmid replicons were detected (Figure 1).

***E. coli* Serotypes, Virulence factors and Phylogeny**

A total of 19 *E. coli* isolates were identified. *E. coli* serotyping using the ECtyper revealed that
the most common serotypes among the *E. coli* isolates were -:H7 (n=2), O138:H48 (n=2),
O6:H16 (n=2), and O8/O160:H16 (n=2). A number of these serovars and STs are associated with
pathogenicity, notably O6:H16 (41), as well as O8/O160:H16 and O77/O17/O44/O106/O73:H18
(ST394; (42-44)). The strains belonging to these lineages lacked the defining virulence genes of
the respective pathotypes but did contain accessory virulence genes, as shown in
supplementary table 4.
198
Irrespective of whether they belonged to a lineage commonly associated with virulence, most
of the isolates contained a range of adhesins and iron utilization genes. *E. coli* extracellular
protein (ECP) export pathway (*ecp/yag*) and *ompA* and type I fimbriae-encoding operon, *fim*,
encoding genes seen in most *E. coli* genomes, were present in 94.7% (18/19) of *E. coli* isolates.
Fimbriae encoding gene, *f17d*, often seen in enterotoxigenic *E. coli* recovered from animals,
was present in the two O8/O160:H16 isolates and an O-:H7 isolate.
205
The phylogenetic analysis of the 19 *E. coli* isolates from this study and a reference genome
(NZ_HG941718.1) based on SNP is presented in Figure 2. The range of isolates was broader
than with *Salmonella* but closely related pairs of isolates belonging to the same serovar, and ST
were found in three instances. Very similar (2347 SNPs) O6:H16 isolates were recovered from
different food preparation table samples in Bolgatanga. One of the two isolates from milk in
Saboba belonged to ST2165 and was identical (0 SNPs) to a Saboba ST2165 utensil isolate. The
two ST4 isolates from fresh beef and a cow milker's hand differed by 2347 SNPs and are
unlikely to be connected.
214

215   (a)

216
217   (b)

218
219 Figure 2: Phylogeny, plasmid replicons and resistance genes of *E. coli* isolates. (a). SNP-based
220 phylogenetic tree of the *E. coli* isolates showing the range of sequence types, serovars, plasmid
221 replicons and resistance genes detected (b) AMR determinants detected in *E. coli* and the
222 resistance they confer

Plasmid Replicons and ARG Profiles of *E. coli*
Antimicrobial resistance determinants present in the *E. coli* isolates include those encoding
resistance to aminoglycosides (*aph(3")-Ib, aph(6)-Id, aph(6)-Id, aph(3")-Ib*), beta-lactams (*bla*$_{LAP-}$
$_2$, *bla*$_{TEM-1}$), fosfomycin (*fosA7.5*), quinolones (*qnrB19, qnrS1*), sulfonamide (*sul2*), tetracycline
(*tet(A), tet(B)*) and trimethoprim (*dfrA14*) (Figure 2). At least 3 antimicrobial resistance genes
(ARGs) which confer resistance to different classes of antibiotics were present in 8 isolates.
Three isolates carried one ARG each while 8 isolates had no ARGs. Six strains carried the genes
aph(3'')-Ib, aph(6)-Id, blaTEM-1, dfrA14, sul2, qnrS1 and *tet(A)*. The *dfrA14-qnrS1-tet(A)*
resistance gene combination has previously been reported from Nigeria, being part of a
transposon transmitted in an IncX plasmid (45). In this study however, IncX replicons were not
detected. The most common plasmid replicon type detected among the *E. coli* isolates was
pO111 (n=6), originally described in an *E. coli* virulence plasmid and found in the
aforementioned strains belonging to virulence-associated lineages. The other plasmid replicon
types detected were IncY (n=5), IncFII (n=2), IncFIA(HI1) (n=2), Col(pHAD28) (n=2),
IncFIB(pB171) (n=1), IncR (n=1), IncHI1A (n=1) and IncI1-I(Gamma), which was common among
the *Salmonella*, (n=1) (Figure 2). All multidrug resistant *E. coli* strains in this study encode
pO111 or IncY replicons.

DISCUSSION
Salmonella and *E. coli* are the main causes of bacterial foodborne illnesses in Ghana (46). Retail
meat, along with milk and their products are recognized as primary sources of foodborne

Salmonellosis (12) and *E. coli* infection (7). Post-cooking handling practices, exposure during the
points of sale, and suboptimal meat storage conditions collectively contribute to an increased
presence of both pathogenic and spoilage bacteria in ready-to-eat (RTE) meat (47). Within
Ghana, food safety has been inadequately studied in the northern region (10). In this study, we
characterized the genomes of 14 *Salmonella* and 19 *E. coli* previously isolated from Saboba
district and Bolgatanga Municipality in northern Ghana.

The *Salmonella* serovars (Fresno, Plymouth, Infantis, Give and Orleans) identified in this study
are yet to be reported in Ghana, but Guinee et al. (1961) (48) isolated *S.* Agona from cattle in
Ghana and *S.* Give (but ST524, different from ST516 in this study) has been reported from beef
in Nigeria (49). Isolation of *S.* Infantis from retail poultry meat has also been reported in
Ecuador (50), Belgium (51) and Italy (52) and all the serovars detected in this study have been
implicated in human infections. The identification of *Salmonella* serovars not previously
documented in the country in meat and milk products highlights the need for heightened
surveillance and preventive measures to curb the spread of foodborne pathogens and reduce
the risk of associated illnesses.

Unlike *Salmonella*, not all *E. coli* are potential pathogens. However, *E. coli* serve as markers for
faecal contamination and therefore the potential that other pathogens are present. The
predominant *E. coli* STs (ST4, ST10, ST219, ST2522) detected in our study have been previously
isolated from food animals and have been associated with pathogenicity (53-55).

While none of the *E. coli* isolates carried genes encoding ETEC heat-sensitive or heat-labile
enterotoxins, f17d fimbrial genes present in four of the *E. coli* genomes encode ETEC
colonization factors commonly associated with colonization of cattle and other ruminant
isolates (56). *E. coli,* F17 fimbriae are associated with pathogenic *E. coli* recovered from
diarrhoea and septicaemia outbreaks in calves, lambs, and humans, including from outbreaks.

Additionally, two of the ST4 *E. coli* isolates from this study not harbouring f17d fimbrial genes
belong to the serovar O6:H16, one of the most widely disseminated lineages of human
enterotoxigenic *E. coli* (ETEC). O6:H16 ETEC cause outbreaks, often associated with food and/or
inadequate handwashing (57-59). ETEC, by definition, produce plasmid-encoded heat-labile
and/or heat stable toxins not present in the genomes of the isolates from this study.

However, the serovars O8/O160:H16 and O77/O17/O44/O106/O73:H18 belong to a previously
described enteroaggregative *E. coli* (EAEC) lineage (42, 44). Isolates from this study belonging to
the serovars O8/O160:H16 and O77/O17/O44/O106/O73:H18 possessed no EAEC accessory
genes. These strains are from virulent lineages but lack key virulence genes that are plasmid-
encoded, which could mean that these plasmids were lost in the food chain or during isolation
but could be reacquired. Nevertheless, the presence of these strains in food could increase the
risk of foodborne illness.

While strains belonging to virulence-associated lineages lacked key plasmid-encoded virulence
plasmids, several plasmid replicons were detected in the isolate genomes. ColE, Incl1, IncF, and

IncX were commonly detected plasmid replicons in *Salmonella* from food animals in the US
according to McMillan et al. (2019) (60). In this study, the Incl1 replicon was predominant, and
nine of the thirteen *Salmonella* strains harboured Incl1 plasmid replicon, three harboured the
tetA. This is likely to be an instance of a successful mobile element with extraordinary local
reach, a few of which have been reported from West Africa, including Ghana, in the past (45,
61-63). The Incl1-I(Gamma) plasmid replicon observed among *Salmonella* isolates was detected
in all *Salmonella* isolates from Bolgatanga municipality - three different serovars - and none of
the Saboba district isolates. However, one *E. coli* isolate (from a recently reported ST, ST8274)
from Saboba did have this replicon. As it is an Incl1 plasmid replicon, its plasmid should be
better characterized and, it should remain under surveillance because numerous articles have
reported associations with multiple ARGs, such as *tetB*, *tetAR*, bla_{CMY-2}, bla_{TEM-1}, *aac3VIa*, *aphA*,
aadA and *sul1* (60), bla_{CTXM-1} (64), *strA*, *strB*, *cmlA*, *floR*, bla_{SHV-12}, bla_{OXA-2} and *FosA3* (65) in Incl1
plasmids in *Salmonella*. As our own sequence was generated by short read only, the first step
would be to generate long read sequence that could fully assemble the plasmid and make it
possible to identify genetic factors supporting its success.
304
Among the *E. coli* isolates, the plasmid replicons pO111 was the most common replicon. A
previous study by Balbuena-Alonso et al. (2022) (66), revealed that pO111 is usually associated
with extended spectrum beta lactamases gene and is very common in food and clinical isolates.
In this study, all isolates carrying pO111 harbour at least one beta-lactamase gene. Likewise, all
the pO111 plasmid bearing isolates in this study carry ARGs that confer resistance to at least 4
classes of antibiotics. Altogether, these data demonstrate a concerning reservoir of resistance
genes in these foodborne bacteria.
312
Conclusion
This study has characterized the genomes of *Salmonella* and *E. coli* in milk, meat and their
associated utensils. The diverse serovars and virulence genes detected in *Salmonella* strains
indicate potential pathogenicity. Although not all *E. coli* strains are pathogenic, their presence
serves as an indicator of faecal contamination, suggesting the potential presence of other
harmful pathogens. The presence of EAEC strains in food is concerning as EAEC is a well-known
cause of diarrhoeal diseases, particularly in children and immunocompromised individuals,
making its presence in food a serious concern. While antimicrobial resistance was not common
among *Salmonella* strains, most of the *E. coli* strain had at least one resistance gene and almost
half were multidrug resistance and carry mobile elements. Moreover, there have been recent
reports of resistant *Salmonella* and *E. coli* from meat and milk elsewhere in West Africa (49).
324
A recent scoping review reported weak enforcement of food safety regulations as well as a lack
of infrastructure, knowledge and skills to implement them (10, 11, 46). Food contaminated with
and *Salmonella* and *E. coli* can serve as a vehicle for their transmission, posing a significant
public health risk. We recommend that food safety regulation be strengthened in northern
Ghana and elsewhere in West Africa. It is also important to increase awareness among
consumers so that food is handled in such a way to prevent pathogen transmission. There is an
additional need for continuous surveillance and preventive measures to stop the spread of
foodborne pathogens and reduce the risk of associated illnesses in Ghana.

Supplementary Materials

Supplementary Table 1: Accession numbers for genomes generated in this study

Supplementary Table 2: Novel *Salmonella* allelic profile and assigned ST.

Supplementary Table 3: *Salmonella* strain metadata including serotype, ST, plasmid replicon, AMR and virulence profile.

Supplementary Table 4: *E. coli* strain metadata including serotype, ST, plasmid replicon, AMR and virulence profile.

Acknowledgements

We thank Ayorinde Afolayan, Faith I. Oni, and Abeeb Adeniyi for technical assistance and Jola-Ade Ajiboye, Kesiena Akpede and Pernille Nilsson for logistic support.

Whole genome sequencing and analysis for this project was supported through SEQAFRICA. The SEQAFRICA project is funded by the Department of Health and Social Care's Fleming Fund using UK aid. The views expressed in this publication are those of the authors and not necessarily those of the UK Department of Health and Social Care or its Management Agent, Mott MacDonald. INO is a Calestous Juma Fellow supported by the Bill and Melinda Gates Foundation.

REFERENCES:

1.   Grace D, Wu F, Havelaar AH. MILK Symposium review: Foodborne diseases from milk and milk products in developing countries-Review of causes and health and economic implications. J Dairy Sci. 2020;103(11):9715-29.

2.   Abraham S, O'Dea M, Sahibzada S, Hewson K, Pavic A, Veltman T, et al. Escherichia coli and Salmonella spp. isolated from Australian meat chickens remain susceptible to critically important antimicrobial agents. PLoS One. 2019;14(10):e0224281.

3.   Elzhraa F, Al-Ashmawy M, El-Sherbini M, Abdelkhalek A. Critical occurrence of verotoxgenic E.coli and non-typhoidal salmonella in some heat treated dairy products. Ital J Food Saf. 2021;10(2):9318.

4.   Gutema FD, Abdi RD, Agga GE, Firew S, Rasschaert G, Mattheus W, et al. Assessment of beef carcass contamination with Salmonella and E. coli O 157 in slaughterhouses in Bishoftu, Ethiopia. International Journal of Food Contamination. 2021;8:1-9.

5.   Lay KK, Jeamsripong S, Sunn KP, Angkititrakul S, Prathan R, Srisanga S, et al. Colistin Resistance and ESBL Production in Salmonella and Escherichia coli from Pigs and Pork in the Thailand, Cambodia, Lao PDR, and Myanmar Border Area. Antibiotics (Basel). 2021;10(6).

6.   Parry-Hanson Kunadu A, Holmes M, Miller EL, Grant AJ. Microbiological quality and antimicrobial resistance characterization of Salmonella spp. in fresh milk value chains in Ghana. Int J Food Microbiol. 2018;277:41-9.

7.   Adzitey F, Asante J, Kumalo HM, Khan RB, Somboro AM, Amoako DG. Genomic Investigation into the Virulome, Pathogenicity, Stress Response Factors, Clonal Lineages, and Phylogenetic Relationship of Escherichia coli Strains Isolated from Meat Sources in Ghana. Genes (Basel). 2020;11(12).

8.   Aduah M, Adzitey F, Amoako DG, Abia ALK, Ekli R, Teye GA, et al. Not All Street Food Is Bad: Low Prevalence of Antibiotic-Resistant Salmonella enterica in Ready-to-Eat (RTE) Meats in Ghana Is Associated with Good Vendors' Knowledge of Meat Safety. Foods. 2021;10(5).

9.   Tay MYF, Adzitey F, Sultan SA, Tati JM, Seow KLG, Schlundt J. Whole-Genome Sequencing of Nontyphoidal Salmonella enterica Isolates Obtained from Various Meat Types in Ghana. Microbiol Resour Announc. 2019;8(15).

10.   Botha NN, Ansah EW, Segbedzi CE, Darkwa S. Public health concerns for food contamination in Ghana: A scoping review. PLoS One. 2023;18(8):e0288685.

11.   Asati DA, Abdulai PM, Boateng KS, Appau AAA, Ofori LA, Agyekum TP. Food safety knowledge and practices among raw meat handlers and the microbial content of raw meat sold at Kumasi Abattoir Butchery Shops in Kumasi, Ghana. BMC Public Health. 2024;24(1):975.

12.   Adzitey F, Assoah-Peprah P, Teye GA. Whole-genome sequencing of Escherichia coli isolated from contaminated meat samples collected from the Northern Region of Ghana reveals the presence of multiple antimicrobial resistance genes. J Glob Antimicrob Resist. 2019;18:179-82.

13.   Adzitey F, Teye GA, Amoako DG. Prevalence, phylogenomic insights, and phenotypic characterization of Salmonella enterica isolated from meats in the Tamale metropolis of Ghana. Food Sci Nutr. 2020;8(7):3647-55.

14.   Falgenhauer L, Imirzalioglu C, Oppong K, Akenten CW, Hogan B, Krumkamp R, et al. Detection and Characterization of ESBL-Producing Escherichia coli From Humans and Poultry in Ghana. Front Microbiol. 2018;9:3358.

15.   Adzitey F, Yussif S, Ayamga R, Zuberu S, Addy F, Adu-Bonsu G, et al. Antimicrobial Susceptibility and Molecular Characterization of Escherichia coli Recovered from Milk and Related Samples. Microorganisms. 2022;10(7).

16.   Moawad AA, Hotzel H, Awad O, Tomaso H, Neubauer H, Hafez HM, et al. Occurrence of Salmonella enterica and Escherichia coli in raw chicken and beef meat in northern Egypt and dissemination of their antibiotic resistance markers. Gut Pathog. 2017;9:57.

17.   Geletu US, Usmael MA, Ibrahim AM. Isolation, Identification, and Susceptibility Profile of E. coli, Salmonella, and S. aureus in Dairy Farm and Their Public Health Implication in Central Ethiopia. Vet Med Int. 2022;2022:1887977.

18.   Worku W, Desta M, Menjetta T. High prevalence and antimicrobial susceptibility pattern of salmonella species and extended-spectrum beta-lactamase producing Escherichia coli from raw cattle meat at butcher houses in Hawassa city, Sidama regional state, Ethiopia. PLoS One. 2022;17(1):e0262308.

19.   Lyu N, Feng Y, Pan Y, Huang H, Liu Y, Xue C, et al. Genomic Characterization of Salmonella enterica Isolates From Retail Meat in Beijing, China. Front Microbiol. 2021;12:636332.

20.   Zhan Z, Xu X, Gu Z, Meng J, Wufuer X, Wang M, et al. Molecular epidemiology and antimicrobial resistance of invasive non-typhoidal Salmonella in China, 2007-2016. Infect Drug Resist. 2019;12:2885-97.

21.   Jaja IF, Bhembe NL, Green E, Oguttu J, Muchenje V. Molecular characterisation of antibiotic-resistant Salmonella enterica isolates recovered from meat in South Africa. Acta Trop. 2019;190:129-36.

22.   Rantsiou K, Kathariou S, Winkler A, Skandamis P, Saint-Cyr MJ, Rouzeau-Szynalski K, et al. Next generation microbiological risk assessment: opportunities of whole genome sequencing (WGS) for foodborne pathogen surveillance, source tracking and risk assessment. Int J Food Microbiol. 2018;287:3-9.

23.   Hilt EE, Ferrieri P. Next Generation and Other Sequencing Technologies in Diagnostic Microbiology and Infectious Diseases. Genes (Basel). 2022;13(9).

24.   Nadon C, Van Walle I, Gerner-Smidt P, Campos J, Chinen I, Concepcion-Acevedo J, et al. PulseNet International: Vision for the implementation of whole genome sequencing (WGS) for global food-borne disease surveillance. Euro Surveill. 2017;22(23).

25.   Saba CK, Escudero JA, Herrera-Leon S, Porrero MC, Suarez M, Dominguez L, et al. First identification of Salmonella Urbana and Salmonella Ouakam in humans in Africa. J Infect Dev Ctries. 2013;7(10):691-5.

26.   Abass A, Adzitey F, Huda N. Escherichia coli of Ready-to-Eat (RTE) Meats Origin Showed Resistance to Antibiotics Used by Farmers. Antibiotics (Basel). 2020;9(12).

27.   Tibile AB. Incidence of Salmonella enterica and total bacterial count obtained from cow milk and its related samples from Saboba district: One health approach [A dissertation]: University for Development Studies, Ghana; 2022.

28.   Rahn K, De Grandis SA, Clarke RC, McEwen SA, Galan JE, Ginocchio C, et al. Amplification of an invA gene sequence of Salmonella typhimurium by polymerase chain reaction as a specific method of detection of Salmonella. Mol Cell Probes. 1992;6(4):271-9.

29.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De Novo Assembler. Curr Protoc Bioinformatics. 2020;70(1):e102.

30.   Zhou Z, Alikhan NF, Mohamed K, Fan Y, Agama Study G, Achtman M. The EnteroBase user's guide, with case studies on Salmonella transmissions, Yersinia pestis phylogeny, and Escherichia core genomic diversity. Genome Res. 2020;30(1):138-52.

31.   Ikhimiukor OO, Oaikhena AO, Afolayan AO, Fadeyi A, Kehinde A, Ogunleye VO, et al. Genomic characterization of invasive typhoidal and non-typhoidal Salmonella in southwestern Nigeria. PLoS Negl Trop Dis. 2022;16(8):e0010716.

32.   Akinlabi OC, Nwoko EQ, Dada RA, Ekpo S, Omotuyi A, Nwimo CC, et al. Epidemiology and Risk Factors for Diarrheagenic Escherichia coli Carriage among Children in Northern Ibadan, Nigeria. Am J Trop Med Hyg. 2023;109(6):1223-32.

33.   Hunt M, Mather AE, Sanchez-Buso L, Page AJ, Parkhill J, Keane JA, et al. ARIBA: rapid antimicrobial resistance genotyping directly from sequencing reads. Microb Genom. 2017;3(10):e000131.

34.   Yoshida CE, Kruczkiewicz P, Laing CR, Lingohr EJ, Gannon VP, Nash JH, et al. The Salmonella In Silico Typing Resource (SISTR): An Open Web-Accessible Tool for Rapidly Typing and Subtyping Draft Salmonella Genome Assemblies. PLoS One. 2016;11(1):e0147101.

35.   Bessonov K, Laing C, Robertson J, Yong I, Ziebell K, Gannon VPJ, et al. ECTyper: in silico Escherichia coli serotype and species prediction from raw and assembled whole-genome sequence data. Microb Genom. 2021;7(12).

36.   Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. Antimicrob Agents Chemother. 2014;58(7):3895-903.

37.   Feldgarden M, Brover V, Gonzalez-Escalona N, Frye JG, Haendiges J, Haft DH, et al. AMRFinderPlus and the Reference Gene Catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence. Sci Rep. 2021;11(1):12728.

38.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25(14):1754-60.

39.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of SAMtools and BCFtools. Gigascience. 2021;10(2).

40.   Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015;32(1):268-74.

41.   Scotland SM, Gross RJ, Rowe B. Serotype-related enterotoxigenicity in Escherichia coli O6.H16 and O148.H28. J Hyg (Lond). 1977;79(3):395-403.

42.   Okeke IN, Wallace-Gadsden F, Simons HR, Matthews N, Labar AS, Hwang J, et al. Multi-locus sequence typing of enteroaggregative Escherichia coli isolates from Nigerian children uncovers multiple lineages. PLoS One. 2010;5(11):e14093.

43.   Chattaway MA, Day M, Mtwale J, White E, Rogers J, Day M, et al. Clonality, virulence and antimicrobial resistance of enteroaggregative Escherichia coli from Mirzapur, Bangladesh. J Med Microbiol. 2017;66(10):1429-35.

44.   Chen Y, Chen X, Zheng S, Yu F, Kong H, Yang Q, et al. Serotypes, genotypes and antimicrobial resistance patterns of human diarrhoeagenic Escherichia coli isolates circulating in southeastern China. Clin Microbiol Infect. 2014;20(1):52-8.

45.   Sumrall ET, Gallo EB, Aboderin AO, Lamikanra A, Okeke IN. Dissemination of the transmissible quinolone-resistance gene qnrS1 by IncX plasmids in Nigeria. PLoS One. 2014;9(10):e110279.

46.   Christiana Cudjoe D, Balali GI, Titus OO, Osafo R, Taufiq M. Food Safety in Sub-Sahara Africa, An insight into Ghana and Nigeria. Environ Health Insights. 2022;16:11786302221142484.

47.   Henriques A, Fraqueza M. Listeria monocytogenes and ready-to-eat meat-based food products: Incidence and control. Listeria monocytogenes: incidence, growth behavior and control. 2015:71-103.

48.   Guinee PA, Kampelmacher EH, Willems HM. Six new Salmonella types, isolated in Ghana (S. volta, S. agona, S. wa, S. techimani, S. mampong and S. tafo). Antonie Van Leeuwenhoek. 1961;27:469-72.

49.   Aworh MK, Nilsson P, Egyir B, Owusu FA, Hendriksen RS. Rare serovars of non-typhoidal Salmonella enterica isolated from humans, beef cattle and abattoir environments in Nigeria. PLoS One. 2024;19(1):e0296971.

50.   Mejia L, Vela G, Zapata S. High Occurrence of Multiresistant Salmonella Infantis in Retail Meat in Ecuador. Foodborne Pathog Dis. 2021;18(1):41-8.

51.   Zeng H, De Reu K, Gabriel S, Mattheus W, De Zutter L, Rasschaert G. Salmonella prevalence and persistence in industrialized poultry slaughterhouses. Poult Sci. 2021;100(4):100991.

52.   Casagrande Proietti P, Musa L, Stefanetti V, Orsini M, Toppi V, Branciari R, et al. mcr-1-Mediated Colistin Resistance and Genomic Characterization of Antimicrobial Resistance in ESBL-Producing Salmonella Infantis Strains from a Broiler Meat Production Chain in Italy. Antibiotics (Basel). 2022;11(6).

53.   Araujo S, I ATS, Tacao M, Patinha C, Alves A, Henriques I. Characterization of antibiotic resistant and pathogenic Escherichia coli in irrigation water and vegetables in household farms. Int J Food Microbiol. 2017;257:192-200.

54.   Silva A, Silva V, Pereira JE, Maltez L, Igrejas G, Valentao P, et al. Antimicrobial Resistance and Clonal Lineages of Escherichia coli from Food-Producing Animals. Antibiotics (Basel). 2023;12(6).

55.   Zhao X, Miao Y, Adam FEA, Zhao H, Zhou Z, Su M, et al. ESBLs-producing Escherichia coli from sheep-origin: Genome-wide virulence genes identification and in vivo virulence assessment in mice and Galleria mellonella. Transbound Emerg Dis. 2022;69(6):3606-17.

56.   Bihannic M, Ghanbarpour R, Auvray F, Cavalie L, Chatre P, Boury M, et al. Identification and detection of three new F17 fimbrial variants in Escherichia coli strains isolated from cattle. Vet Res. 2014;45(1):76.

57.   Pacheco AB, Soares KC, de Almeida DF, Viboud GI, Binsztein N, Ferreira LC. Clonal nature of enterotoxigenic Escherichia coli serotype O6: H16 revealed by

randomly amplified polymorphic DNA analysis. Journal of clinical microbiology. 1998;36(7):2099-102.

58.   Beatty ME, Adcock PM, Smith SW, Quinlan K, Kamimoto LA, Rowe SY, et al. Epidemic diarrhea due to enterotoxigenic Escherichia coli. Clin Infect Dis. 2006;42(3):329-34.

59.   Willshaw GA, Cheasty T, Rowe B, Smith HR, Faithfull-Davies DN, Brooks TG. Isolation of enterotoxigenic Escherichia coli from British troops in Saudi Arabia. Epidemiol Infect. 1995;115(3):455-63.

60.   McMillan EA, Gupta SK, Williams LE, Jove T, Hiott LM, Woodley TA, et al. Antimicrobial Resistance Genes, Cassettes, and Plasmids Present in Salmonella enterica Associated With United States Food Animals. Front Microbiol. 2019;10:832.

61.   Dekker D, Eibach D, Boahen KG, Akenten CW, Pfeifer Y, Zautner AE, et al. Fluoroquinolone-Resistant Salmonella enterica, Campylobacter spp., and Arcobacter butzleri from Local and Imported Poultry Meat in Kumasi, Ghana. Foodborne Pathog Dis. 2019;16(5):352-8.

62.   Labar AS, Millman JS, Ruebush E, Opintan JA, Bishar RA, Aboderin AO, et al. Regional dissemination of a trimethoprim-resistance gene cassette via a successful transposable element. PLoS One. 2012;7(5):e38142.

63.   Monarrez R, Braun M, Coburn-Flynn O, Botelho J, Odetoyin BW, Otero-Vera JI, et al. A large self-transmissible resistance plasmid from Nigeria contains genes that ameliorate a carrying cost. Sci Rep. 2019;9(1):19624.

64.   Abraham S, Kirkwood RN, Laird T, Saputra S, Mitchell T, Singh M, et al. Dissemination and persistence of extended-spectrum cephalosporin-resistance encoding IncI1-bla(CTXM-1) plasmid among Escherichia coli in pigs. ISME J. 2018;12(10):2352-62.

65.   Foley SL, Kaldhone PR, Ricke SC, Han J. Incompatibility Group I1 (IncI1) Plasmids: Their Genetics, Biology, and Public Health Relevance. Microbiol Mol Biol Rev. 2021;85(2).

66.   Balbuena-Alonso MG, Cortes-Cortes G, Kim JW, Lozano-Zarain P, Camps M, Del Carmen Rocha-Gracia R. Genomic analysis of plasmid content in food isolates of E. coli strongly supports its role as a reservoir for the horizontal transfer of virulence and antibiotic resistance genes. Plasmid. 2022;123-124:102650.