1 **Genomic characterization of foodborne *Salmonella enterica* and *Escherichia coli* isolates from**
2 **Saboba district and Bolgatanga Municipality Ghana**
3

4 Gabriel Temitope Sunmonu [1], Frederick Adzitey [2], Erkison Ewomazino Odih [1], Boniface Awini Tibile [2],
5 Rejoice Ekli [2], Martin Aduah [2], Anderson O Oaikhena [1], Olabisi C Akinlabi [1], Akebe Luther King Abia [3], Daniel
6 Gyamfi Amoako [3, 4], and Iruka N Okeke [1]
7
8 1. Department of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Ibadan, Nigeria.
9 2. Faculty of Agriculture, Food and Consumer Sciences, University for Development Studies, Tamale, Ghana
10 3. Antimicrobial Research Unit, College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa
11 4. Department of Integrative Biology and Bioinformatics, University of Guelph, Ontario, Canada
12

Abstract
Salmonella and *Escherichia coli* are well-known bacteria commonly associated with foodborne
illnesses in humans and animals. Genomic characterization of these pathogens provides valuable
insights into their evolution, virulence factors and resistance determinants. This study
characterized *Salmonella* (n = 14) and *E. coli* (n = 19) from milk, meat and its associated utensils
in Ghana using whole-genome sequencing. Most of the *Salmonella* serovars (Fresno, Plymouth,
Infantis, Give and Orleans) identified in this study are yet to be reported in Ghana. The *Salmonella*
enterica isolates were largely pan-sensitive but genes conferring resistance to fosfomycin
(*fosA7.2*) and tetracycline (*tet(A)*) were detected in one and three isolates respectively. While 7of
the *Salmonella* isolates carry the IncI1-I(Gamma) plasmid replicon. While antimicrobial resistance
was not common among *Salmonella* strains, most (11/19) of the *E. coli* strains had at least one
resistance gene and almost half (8/19) were multidrug resistance and carry mobile elements.
Three of the 19 *E. coli* strains belong to serovars commonly associated with enteroaggregative *E.*
coli (EAEC) lineage. While strains belonging to virulence-associated lineages lacked key plasmid-
encoded virulence plasmids, several plasmid replicons were detected in most of the *E. coli*
(14/19) strains. Food contaminated with these pathogens can serve as a vehicle for transmission,
posing a significant public health risk and necessitating stringent food safety and hygiene
practices to prevent outbreaks. Hence, there is need for continuous surveillance and preventive
measures to stop the spread of foodborne pathogens and reduce the risk of associated illnesses
in Ghana.
33

INTRODUCTION
Milk and meat are essential protein sources that constitute a significant and nutrient-rich
component of human diets. However, their consumption is frequently associated with foodborne
infections (1), particularly those caused by *Escherichia coli* and *Salmonella* (2-5). Salmonellosis is
under-reported in Ghana, and only a few studies have investigated the plausible role of
contaminated milk, meat or meat products in facilitating their transmission (6-8). There have
been a few reports of meat samples contaminated with *E. coli* in Ghana (9-11). Some of these *E.*
coli strains harbour virulence and antimicrobial resistance genes (ARGs), which is of public health
concern however few have been characterized in detail.
43
There are public health and food safety implications of finding E. coli and Salmonella in food
because they are invariably of faecal origin. Globally, *Salmonella* and *Escherichia coli* sourced

from milk and meat increasingly exhibit resistance to different classes of antibiotics that is
commonly mediated by mobile elements (10, 12-14). The prevalence of multidrug-resistant
(MDR) *S. enterica* and *E. coli* is also on the rise in clinical infections (15, 16). Notably, resistance
to extended spectrum beta-lactams, trimethoprim/sulfamethoxazole, chloramphenicol and
ciprofloxacin has been reported, often associated with plasmids that could mediate their spread,
in both *S. enterica* and *E. coli* isolates from milk and retail meats in Ghana (7, 9, 11).
52
Various methods, including serotyping, antibiotic profiling, pulsed-field gel electrophoresis and
whole genome sequencing, have been employed to elucidate the phenotypic and genotypic
attributes of foodborne pathogens and determine their interrelationships and connections to
pandemic clones of interest (10, 17). Next-generation sequencing (NGS) technology, the most
versatile and informative approach, has gained recent prominence (18, 19). NGS is now used by
PulseNet to categorize foodborne diseases, enabling nuanced epidemiological investigations
(20). Identification of virulence factors, antibiotic resistance genes, and serotypes are all possible
by genomic analysis, which can also provide enhanced information on strain inter-relatedness,
and therefore enable source attribution. Data on the genomic characterization of *Salmonella* and
E. coli from milk and meat are few from low- and middle-income countries, including Ghana (6,
21). In light of this gap, this study aims to characterize the resistance, virulence and plasmid
profile of *Salmonella* and *E. coli* isolated from fresh different retail meats, milk, and associated
samples (handler's hand swab, table, knife and faecal samples) in in Saboba district and
Bolgatanga Municipality of Ghana.
67
METHODS
Strains
A total of 33 bacterial isolates (14 *Salmonella* and 19 *E. coli* species) previously isolated from
various fresh and ready-to-eat meats, meat sellers' tables, milk, milk-collecting utensils, milkers'
hands and faeces of lactating cows were characterized for this study (22, 23). The isolates
originated from markets and farms in Bolgatanga Municipality and Saboba District in Northern
Ghana and were cryopreserved in 50% glycerol in Luria broth at -80°C.
75
***Salmonella* and *E. coli* identification**
Salmonella and *E. coli* were biotyped with the Gram-negative (GN) test kit (Ref: 21341) on VITEK
2 systems (version 2.0, Marcy-l'Etoile, France, Biomérieux) according to manufacturer's
instructions. *Salmonella* isolates were additionally confirmed using a latex agglutination kit for
Salmonella (Oxoid Limited, Basingstoke, UK) and by PCR targeting the *invA* gene as described by
Rahn et al. (1992) (24), using PCR oligonucleotides *invA139f* GTGAAATTATCGCCACGTTCGGGCAA
and *invA141r* TCATCGCACCGTCAAGGAACC. PCR was performed using PuRe Taq Ready-To-Go PCR
Beads (illustra™). The PCR cycle used an initial denaturation temperature of 95^0C for two
minutes, followed by 35 cycles of denaturation at 95^0C for 30 seconds, annealing at 55^0C for 30
seconds, extension at 72^0C for two minutes and then a terminal extension at 72^0C for five
minutes. Visualization of the 284 bp amplicon was accomplished after electrophoresis on 1.5%
(w/v) agarose gels stained with Gel red (biotium), using a UVP GelMax transilluminator and
imager.
89

DNA Extraction, Library Preparation and Whole Genome Sequencing

Genomic DNA of the isolates was extracted using Wizard DNA extraction kit (Promega; Wisconsin, USA) in accordance with manufacturer's protocol. Using a dsDNA Broad Range quantification assay, the extracted DNA was quantified on a Qubit fluorometer (Invitrogen; California, USA). dsDNA libraries were prepared using NEBNext Ultra II FS DNA library kit for Illumina with 96-unique indexes (New England Biolabs, Massachusetts, USA; Cat. No: E6609L). DNA libraries was quantified using dsDNA High Sensitivity quantification assay on a Qubit fluorometer (Invitrogen; California, USA) and fragment length analysed with the Bioanalyzer (Agilent). Denatured libraries were sequenced on an Illumina MiSeq (Illumina, California, USA). The raw sequence reads were *de novo* assembled using SAdes v3.15.3 (25) according to GHRU protocols (https://gitlab.com/cgps/ghru/pipelines/dsl2/pipelines/assembly).

Sequence Typing of *Salmonella* and *E. coli* Genomes

Sequence reads were deposited in the *Salmonella* and *E. coli* database for *Salmonella* and *E. coli* respectively on EnteroBase (26). Multi-locus sequence types (MLST) for the isolates were determined and core-genome MLST calculated. The *Salmonella* genome assemblies were analysed using the *Salmonella* In-Silico Typing Resource (SISTR) for the prediction of serovars and serogroups (27) (https://github.com/phac-nml/sistr_cmd), while the serotyping of the *E. coli* genome was done using ECtyper (28).

Identification of AMR, Plasmids and Virulence genes

PlasmidFinder (29) was utilized to identify plasmid replicons that were present in the assembled genomes. AMRFinderPlus v3.10.24 (30) and its associated database (version 2022-04-04.1) were used to predict the antimicrobial resistance genes carried by the isolates and the drug classes to which they probably conferred resistance. Using ARIBA (31) and the virulence factor database (VFDB), we were also able to identify the virulence genes that were present in the isolates.

Single Nucleotide Polymorphism (SNP) Calling and Phylogenetic analysis

For phylogenetic analysis, reference sequences for the *S. enterica* and *E. coli* genomes were objectively selected from the National Center for Biotechnology Information Reference Sequence (RefSeq) database (https://www.ncbi.nlm.nih.gov/refseq/) using BactinspectorMax v0.1.3 (https://gitlab.com/antunderwood/bactinspector). The selected references were the *S. enterica* subsp. enterica serovar Fresno strain (assembly accession: GCF_003590695.1) and the *E. coli* O25b:H4-ST131 strain (assembly accession: GCF_000285655.3). The sequence reads for each species were then mapped to the chromosome of the reference using BWA (v0.7.17) (32) and variants were called and filtered using bcftools (v1.9) (Danecek et al., 2021) as implemented in the GHRU SNP phylogeny pipeline (https://gitlab.com/cgps/ghru/pipelines/snp_phylogeny). Variant positions were concatenated into a pseudoalignment and used to generate a maximum likelihood tree using iqtree (v1.6.8) (33). SNP distances between the genome pairs were calculated using snp-dists v.0.8.2 (https://github.com/tseemann/snp-dists).

Data availability

The sequence reads were deposited in the European Nucleotide Archive with the study accession PRJEB58695 (Supplementary table 1).

134
RESULTS
Salmonella **Serotypes, Sequence Types (STs), Virulence Factors and Phylogeny**
We used SISTR software to predict the serovars of the 14 Salmonella strains characterized in this
study from whole genome sequence. The most common serotype was Fresno (n=6) followed by
Give (n=3), Orleans (n=2), Plymouth (n=1), Agona (n=1) and Infantis (n=1). The *S.* Fresno and *S.*
Orleans isolates were from previously unreported sequence types, now designated ST 10742 and
ST 10465 respectively (Supplementary table 2). As shown in Figure 1, all the isolates from ready-
to-eat pork, mutton and chicken belong to serovar Fresno. *S.* Fresno isolates were also isolated
from a meat vendor's knife, as was *S.* Orleans. The three milk isolates, which were all from
Saboba, belong to the serovars Plymouth (ST 565), Give (ST 516) and Agona (ST13) (Figure 1).
Two more ST 516 *S.* Give isolates were recovered from the faeces of milking cow and from a
milking utensil.
147
148 Phylogenetic analysis of the 14 *Salmonella enterica* isolates from this study showed that all *S.*
149 Fresno isolates, irrespective of source, clustered together and differed by < 3 SNPs. The two *S.*
150 Orleans isolates were identical (0 SNPs) and the three *S.* Give isolates were also identical, with
151 the isolates originating from milk, faeces of milking cow and from a milking utensil, also in
152 Saboba.
153

154

156 Figure 1: Core genome SNP-based Maximum likelihood tree showing phylogenetic relationships
157 among strains sequenced in this study. The matrix below the tree shows the sequence types,
158 serotypes, location of isolation (Bolgatanga municipality or Saboba district) as well as the
159 source/host sample and presence or absence of the tetA and fosA resistance genes.

160
All *Salmonella* isolates harboured curli (*csg*) genes as well as *bcf*, *fim* and *ste* fimbrial operons and
ten of them, representing all serovars except *S*. Give and Plymouth, carried long polar fimbriae
(*lpf*) genes. The *S*. Infantis and *S*. Agona strains carried *ratB* and *shdA*. Type III secretion system
effector genes, such as: *inv*, *org*, *prg*, *sif*, *spa*, *sse*, *ssa* and *sop* were detected in all the isolates
while *avr* was present in 57.1% (8/14) of the isolates and only one isolate harbours *gogB*. Four of
the isolates also encode the cytholethal distending toxin gene, *cdtB* (Supplementary table 3).
167
Plasmid Replicons and ARG Profiles of *Salmonella*
The *Salmonella enterica* isolates were largely pan-sensitive but genes conferring resistance to
fosfomycin (*fosA7.2*) and tetracycline (*tet(A)*) were detected in one and three isolates
respectively. Both *S*. Orleans isolates and one of the *S*. Fresno, from ready-to-eat mutton carried
tetA, along with an IncI1-I(Gamma) plasmid replicon, which was also seen in six other, *tetA*-
negative strains. Interestingly, the IncI1-I(Gamma) plasmid replicon was detected in all isolates
from Bolgatanga municipality, irrespective of serovar, and no isolate from Saboba district
harboured this plasmid. The fosfomycin resistance gene was found in the *S*. Agona genome, in
which no plasmid replicons were detected (Figure 1).
177
***E. coli* Serotypes, Virulence factors and Phylogeny**
A total of 19 *E. coli* isolates were identified. *E. coli* serotyping using the ECtyper revealed that the
most common serotypes among the *E. coli* isolates were -:H7 (n=2), O138:H48 (n=2), O6:H16
(n=2), and O8/O160:H16 (n=2). A number of these serovars and STs are associated with
pathogenicity, notably O6:H16 (34), as well as O8/O160:H16 and O77/O17/O44/O106/O73:H18
(ST394; (35-37)). The strains belonging to these lineages lacked the defining virulence genes of
the respective pathotypes but did contain accessory virulence genes, as shown in supplementary
table 4.
186
Irrespective of whether they belonged to a lineage commonly associated with virulence, most of
the isolates contained a range of adhesins and iron utilization genes. *E. coli* extracellular protein
(ECP) export pathway (*ecp/yag*) and *ompA* and type I fimbriae-encoding operon, *fim*, encoding
genes seen in most *E. coli* genomes, were present in 94.7% (18/19) of *E. coli* isolates. Fimbriae
encoding gene, *f17d*, often seen in enterotoxigenic *E. coli* recovered from animals, was present
in the two O8/O160:H16 isolates and an O-:H7 isolate.
193
The phylogenetic analysis of the 19 *E. coli* isolates from this study and a reference genome
(NZ_HG941718.1) based on SNP is presented in Figure 2. The range of isolates was broader than
with *Salmonella* but closely related pairs of isolates belonging to the same serovar, and ST were
found in three instances. Very similar (2347 SNPs) O6:H16 isolates were recovered from different
food preparation table samples in Bolgatanga. One of the two isolates from milk in Saboba
belonged to ST2165 and was identical (0 SNPs) to a Saboba ST2165 utensil isolate. The two ST4
isolates from fresh beef and a cow milker's hand differed by 2347 SNPs and are unlikely to be
connected.
202

203 (a)

204
205 (b)

Figure 2: Phylogeny, plasmid replicons and resistance genes of *E. coli* isolates. A. SNP-based phylogenetic tree of the *E. coli* isolates showing the range of sequence types, serovars, plasmid replicons and resistance genes detected (b) AMR determinants detected in *E. coli* and the resistance they confer

Plasmid Replicons and ARG Profiles of *E. coli*

Antimicrobial resistance determinants present in the *E. coli* isolates include those encoding resistance to aminoglycosides (*aph(3")-Ib*, *aph(6)-Id*, *aph(6)-Id*, *aph(3")-Ib*), beta-lactams (bla_{LAP-2}, bla_{TEM-1}), fosfomycin (*fosA7.5*), quinolones (*qnrB19*, *qnrS1*), sulfonamide (*sul2*), tetracycline (*tet(A)*, *tet(B)*) and trimethoprim (*dfrA14*) (Figure 2). At least 3 antimicrobial resistant genes (ARGs) which confer resistance to different classes of antibiotics were present in 8 isolates. Three isolates carried one ARG each while 8 isolates had no ARGs. Six strains carried the genes *aph(3'')-Ib, aph(6)-Id, blaTEM-1, dfrA14, sul2, qnrS1* and *tet(A). The dfrA14-qnrS1-tet(A) resistance* gene combination has previously been reported from Nigeria, being part of a transposon transmitted in an IncX plasmid (38). In this study however, IncX replicons were not detected. The most common plasmid replicon type detected among the *E. coli* isolates was pO111 (n=6), originally described in an *E. coli* virulence plasmid and found in the aforementioned strains belonging to virulence-associated lineages. The other plasmid replicon types detected were IncY (n=5), IncFII (n=2), IncFIA(HI1) (n=2), Col(pHAD28) (n=2), IncFIB(pB171) (n=1), IncR (n=1), IncHI1A (n=1) and IncI1-I(Gamma), which was common among the *Salmonella*, (n=1) (Figure 2). All multidrug resistant strains encode pO111 or IncY replicons.

DISCUSSION

Salmonella and *E. coli* are the main causes of bacterial foodborne illnesses in Ghana (39). Retail meat, along with milk and their products are recognized as primary sources of foodborne Salmonellosis (9) and *E. coli* infection (6). Post-cooking handling practices, exposure during the

points of sale, and suboptimal meat storage conditions collectively contribute to an increased
presence of both pathogenic and spoilage bacteria in ready-to-eat (RTE) meat (40). In this study,
we characterized the genomes of 14 *Salmonella* and 19 *E. coli* previously isolated from Saboba
district and Bolgatanga Municipality in northern Ghana.

The *Salmonella* serovars (Fresno, Plymouth, Infantis, Give and Orleans) identified in this study
are yet to be reported in Ghana, but Guinee et al. (1961) (41) isolated *S.* Agona from cattle in
Ghana. Isolation of *S.* Infantis from retail poultry meat has also been reported in Ecuador (42),
Belgium (43) and Italy (44) and all the serovars detected in this study have been implicated in
human infections. The identification of Salmonella serovars not previously documented in the
country in meat and milk products highlights the need for heightened surveillance and preventive
measures to curb the spread of foodborne pathogens and reduce the risk of associated illnesses.

Unlike *Salmonella*, not all *E. coli* are potential pathogens. However, *E. coli* serve as markers for
faecal contamination and therefore the potential that other pathogens are present. The
predominant *E. coli* STs (ST4, ST10, ST219, ST2522) detected in our study have been previously
isolated from food animals and have been associated with pathogenicity (45-47). While none of
the isolates carried genes encoding ETEC heat-sensitive or heat-labile enterotoxins, f17d fimbrial
genes present in four of the *E. coli* genomes encode ETEC colonization factors commonly
associated with colonization of cattle and other ruminant isolates (48). *E. coli,* F17 fimbriae are
associated with pathogenic *E. coli* recovered from diarrhoea and septicaemia outbreaks in calves,
lambs, and humans, including from outbreaks. Additionally, two of the ST4 *E. coli* isolates from
this study not harbouring f17d fimbrial genes belong to the serovar O6:H16, one of the most
widely disseminated lineages of human enterotoxigenic *E. coli* (ETEC). O6:H16 ETEC cause
outbreaks, often associated with food and/or inadequate handwashing (49-51). ETEC, by
definition, produce plasmid-encoded heat-labile and/or heat stable toxins not present in the
genomes of the isolates from this study. However, the serovars O8/O160:H16 and
O77/O17/O44/O106/O73:H18 belong to a previously described enteroaggregative *E. coli* (EAEC)
lineage (35, 37). Isolates from this study belonging to the serovars O8/O160:H16 and
O77/O17/O44/O106/O73:H18 possess no EAEC accessory genes. These strains are from virulent
lineages but lack key virulence genes that are plasmid-encoded, which could mean that these
plasmids were lost in the food chain or during isolation but could be reacquired. Nevertheless,
the presence of these strains in food could increase the risk of foodborne illness.

While strains belonging to virulence-associated lineages lacked key plasmid-encoded virulence
plasmids, several plasmid replicons were detected in the isolate genomes. ColE, IncI1, IncF, and
IncX were commonly detected plasmid replicons in *Salmonella* from food animals in the US
according to McMillan et al. (2019) (52). In this study, the IncI1 replicon was predominant, and
nine of the thirteen *Salmonella* strains harboured IncI1 plasmid replicon, three harboured the
tetA. This is likely to be an instance of a successful mobile element with extraordinary local reach,
a few of which have been reported from West Africa, including Ghana, in the past (38, 53-55).
The IncI1-I(Gamma) plasmid replicon observed among *S. enterica* strains was detected in all
Salmonella isolates from Bolgatanga municipality - three different serovars - and none of the
Saboba district isolates. However, one *E. coli* isolate (from a recently reported ST, ST8274) from

Saboba did have this replicon. As it is an IncI1 plasmid replicon, its plasmid should be better
characterized and, it should remain under surveillance because numerous articles have reported
associations with multiple ARGs, such as *tetB*, *tetAR*, *bla*$_{CMY-2}$, *bla*$_{TEM-1}$, *aac3VIa*, *aphA*, *aadA* and
sul1 (52), *bla*$_{CTXM-1}$ (56), *strA*, *strB*, *cmlA*, *floR*, *bla*$_{SHV-12}$, *bla*$_{OXA-2}$ and *FosA3* (57) in IncI1 plasmids
in *Salmonella*. As our own sequence was generated by short read only, the first step would be to
generate long read sequence that could fully assemble the plasmid and make it possible to
identify genetic factors supporting its success.

Among the *E. coli* isolates, the plasmid replicons pO111 was the most common replicon. A
previous study by Balbuena-Alonso et al. (2022) (58), revealed that pO111 is usually associated
with extended spectrum beta lactamases gene and is very common in food and clinical isolates.
In this study, all isolates carrying pO111 harbour at least one beta-lactamase. Likewise, all the
pO111 plasmid bearing isolates in this study carry ARGs that confer resistance to at least 4 classes
of antibiotics. Altogether, these data demonstrate a concerning reservoir of resistance genes in
these foodborne bacteria.

Conclusion
This study has characterized the genomes of *Salmonella* and *E. coli* in milk, meat and their
associated utensils. The diverse serovars and virulence genes detected in *Salmonella* strains
indicate potential pathogenicity. All *E. coli* are not always pathogenic, but they serve as indicators
of faecal contamination, suggesting a potential risk for the presence of other pathogens. The
presence of EAEC lineages in food is concerning as these bacteria are known to cause diarrhoea,
especially in children and immunocompromised individuals. While antimicrobial resistance was
not common among *Salmonella* strains, most of the *E. coli* strain had at least one resistance gene
and almost half were multidrug resistance and carry mobile elements. Food contaminated with
these pathogens can serve as a vehicle for transmission, posing a significant public health risk
and necessitating stringent food safety and hygiene practices to prevent outbreaks. Hence, there
is need for continuous surveillance and preventive measures to stop the spread of foodborne
pathogens and reduce the risk of associated illnesses in Ghana.

Supplementary Materials
Supplementary Table 1: Accession numbers for genomes generated in this study
Supplementary Table 2: Novel *Salmonella* allelic profile and assigned ST.
Supplementary Table 3: *Salmonella* strain metadata including serotype, ST, plasmid replicon,
AMR and virulence profile.
Supplementary Table 4: *E. coli* strain metadata including serotype, ST, plasmid replicon, AMR and
virulence profile.

Acknowledgements
We thank Ayorinde Afolayan, Faith I. Oni, and Abeeb Adeniyi for technical assistance and Jola-
Ade Ajiboye, Kesiena Akpede and Pernille Nilsson for logistic support.

Whole genome sequencing and analysis for this project was supported through SEQAFRICA. The
SEQAFRICA project is funded by the Department of Health and Social Care's Fleming Fund using

UK aid. The views expressed in this publication are those of the authors and not necessarily those
of the UK Department of Health and Social Care or its Management Agent, Mott MacDonald. INO
is a Calestous Juma Fellow supported by the Bill and Melinda Gates Foundation.
324

REFERENCES:

1. Grace D, Wu F, Havelaar AH. MILK Symposium review: Foodborne diseases from milk and milk products in developing countries-Review of causes and health and economic implications. J Dairy Sci. 2020;103(11):9715-29.

2. Abraham S, O'Dea M, Sahibzada S, Hewson K, Pavic A, Veltman T, et al. Escherichia coli and Salmonella spp. isolated from Australian meat chickens remain susceptible to critically important antimicrobial agents. PLoS One. 2019;14(10):e0224281.

3. Elzhraa F, Al-Ashmawy M, El-Sherbini M, Abdelkhalek A. Critical occurrence of verotoxgenic E.coli and non-typhoidal salmonella in some heat treated dairy products. Ital J Food Saf. 2021;10(2):9318.

4. Gutema FD, Abdi RD, Agga GE, Firew S, Rasschaert G, Mattheus W, et al. Assessment of beef carcass contamination with Salmonella and E. coli O 157 in slaughterhouses in Bishoftu, Ethiopia. International Journal of Food Contamination. 2021;8:1-9.

5. Lay KK, Jeamsripong S, Sunn KP, Angkititrakul S, Prathan R, Srisanga S, et al. Colistin Resistance and ESBL Production in Salmonella and Escherichia coli from Pigs and Pork in the Thailand, Cambodia, Lao PDR, and Myanmar Border Area. Antibiotics (Basel). 2021;10(6).

6. Adzitey F, Asante J, Kumalo HM, Khan RB, Somboro AM, Amoako DG. Genomic Investigation into the Virulome, Pathogenicity, Stress Response Factors, Clonal Lineages, and Phylogenetic Relationship of Escherichia coli Strains Isolated from Meat Sources in Ghana. Genes (Basel). 2020;11(12).

7. Parry-Hanson Kunadu A, Holmes M, Miller EL, Grant AJ. Microbiological quality and antimicrobial resistance characterization of Salmonella spp. in fresh milk value chains in Ghana. Int J Food Microbiol. 2018;277:41-9.

8. Tay MYF, Adzitey F, Sultan SA, Tati JM, Seow KLG, Schlundt J. Whole-Genome Sequencing of Nontyphoidal Salmonella enterica Isolates Obtained from Various Meat Types in Ghana. Microbiol Resour Announc. 2019;8(15).

9. Adzitey F, Assoah-Peprah P, Teye GA. Whole-genome sequencing of Escherichia coli isolated from contaminated meat samples collected from the Northern Region of Ghana reveals the presence of multiple antimicrobial resistance genes. J Glob Antimicrob Resist. 2019;18:179-82.

10. Adzitey F, Teye GA, Amoako DG. Prevalence, phylogenomic insights, and phenotypic characterization of Salmonella enterica isolated from meats in the Tamale metropolis of Ghana. Food Sci Nutr. 2020;8(7):3647-55.

11. Falgenhauer L, Imirzalioglu C, Oppong K, Akenten CW, Hogan B, Krumkamp R, et al. Detection and Characterization of ESBL-Producing Escherichia coli From Humans and Poultry in Ghana. Front Microbiol. 2018;9:3358.

12. Moawad AA, Hotzel H, Awad O, Tomaso H, Neubauer H, Hafez HM, et al. Occurrence of Salmonella enterica and Escherichia coli in raw chicken and beef meat in northern Egypt and dissemination of their antibiotic resistance markers. Gut Pathog. 2017;9:57.

13. Geletu US, Usmael MA, Ibrahim AM. Isolation, Identification, and Susceptibility Profile of E. coli, Salmonella, and S. aureus in Dairy Farm and Their Public Health Implication in Central Ethiopia. Vet Med Int. 2022;2022:1887977.

14.   Worku W, Desta M, Menjetta T. High prevalence and antimicrobial susceptibility pattern of salmonella species and extended-spectrum beta-lactamase producing Escherichia coli from raw cattle meat at butcher houses in Hawassa city, Sidama regional state, Ethiopia. PLoS One. 2022;17(1):e0262308.

15.   Lyu N, Feng Y, Pan Y, Huang H, Liu Y, Xue C, et al. Genomic Characterization of Salmonella enterica Isolates From Retail Meat in Beijing, China. Front Microbiol. 2021;12:636332.

16.   Zhan Z, Xu X, Gu Z, Meng J, Wufuer X, Wang M, et al. Molecular epidemiology and antimicrobial resistance of invasive non-typhoidal Salmonella in China, 2007-2016. Infect Drug Resist. 2019;12:2885-97.

17.   Jaja IF, Bhembe NL, Green E, Oguttu J, Muchenje V. Molecular characterisation of antibiotic-resistant Salmonella enterica isolates recovered from meat in South Africa. Acta Trop. 2019;190:129-36.

18.   Rantsiou K, Kathariou S, Winkler A, Skandamis P, Saint-Cyr MJ, Rouzeau-Szynalski K, et al. Next generation microbiological risk assessment: opportunities of whole genome sequencing (WGS) for foodborne pathogen surveillance, source tracking and risk assessment. Int J Food Microbiol. 2018;287:3-9.

19.   Hilt EE, Ferrieri P. Next Generation and Other Sequencing Technologies in Diagnostic Microbiology and Infectious Diseases. Genes (Basel). 2022;13(9).

20.   Nadon C, Van Walle I, Gerner-Smidt P, Campos J, Chinen I, Concepcion-Acevedo J, et al. PulseNet International: Vision for the implementation of whole genome sequencing (WGS) for global food-borne disease surveillance. Euro Surveill. 2017;22(23).

21.   Saba CK, Escudero JA, Herrera-Leon S, Porrero MC, Suarez M, Dominguez L, et al. First identification of Salmonella Urbana and Salmonella Ouakam in humans in Africa. J Infect Dev Ctries. 2013;7(10):691-5.

22.   Abass A, Adzitey F, Huda N. Escherichia coli of Ready-to-Eat (RTE) Meats Origin Showed Resistance to Antibiotics Used by Farmers. Antibiotics (Basel). 2020;9(12).

23.   Aduah M, Adzitey F, Amoako DG, Abia ALK, Ekli R, Teye GA, et al. Not All Street Food Is Bad: Low Prevalence of Antibiotic-Resistant Salmonella enterica in Ready-to-Eat (RTE) Meats in Ghana Is Associated with Good Vendors' Knowledge of Meat Safety. Foods. 2021;10(5).

24.   Rahn K, De Grandis SA, Clarke RC, McEwen SA, Galan JE, Ginocchio C, et al. Amplification of an invA gene sequence of Salmonella typhimurium by polymerase chain reaction as a specific method of detection of Salmonella. Mol Cell Probes. 1992;6(4):271-9.

25.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De Novo Assembler. Curr Protoc Bioinformatics. 2020;70(1):e102.

26.   Zhou Z, Alikhan NF, Mohamed K, Fan Y, Agama Study G, Achtman M. The EnteroBase user's guide, with case studies on Salmonella transmissions, Yersinia pestis phylogeny, and Escherichia core genomic diversity. Genome Res. 2020;30(1):138-52.

27.   Yoshida CE, Kruczkiewicz P, Laing CR, Lingohr EJ, Gannon VP, Nash JH, et al. The Salmonella In Silico Typing Resource (SISTR): An Open Web-Accessible Tool for Rapidly Typing and Subtyping Draft Salmonella Genome Assemblies. PLoS One. 2016;11(1):e0147101.

28.   Bessonov K, Laing C, Robertson J, Yong I, Ziebell K, Gannon VPJ, et al. ECTyper: in silico Escherichia coli serotype and species prediction from raw and assembled whole-genome sequence data. Microb Genom. 2021;7(12).

29.   Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. Antimicrob Agents Chemother. 2014;58(7):3895-903.

30.   Feldgarden M, Brover V, Gonzalez-Escalona N, Frye JG, Haendiges J, Haft DH, et al. AMRFinderPlus and the Reference Gene Catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence. Sci Rep. 2021;11(1):12728.

31.   Hunt M, Mather AE, Sanchez-Buso L, Page AJ, Parkhill J, Keane JA, et al. ARIBA: rapid antimicrobial resistance genotyping directly from sequencing reads. Microb Genom. 2017;3(10):e000131.

32.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25(14):1754-60.

33.   Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015;32(1):268-74.

34.   Scotland SM, Gross RJ, Rowe B. Serotype-related enterotoxigenicity in Escherichia coli O6.H16 and O148.H28. J Hyg (Lond). 1977;79(3):395-403.

35.   Okeke IN, Wallace-Gadsden F, Simons HR, Matthews N, Labar AS, Hwang J, et al. Multi-locus sequence typing of enteroaggregative Escherichia coli isolates from Nigerian children uncovers multiple lineages. PLoS One. 2010;5(11):e14093.

36.   Chattaway MA, Day M, Mtwale J, White E, Rogers J, Day M, et al. Clonality, virulence and antimicrobial resistance of enteroaggregative Escherichia coli from Mirzapur, Bangladesh. J Med Microbiol. 2017;66(10):1429-35.

37.   Chen Y, Chen X, Zheng S, Yu F, Kong H, Yang Q, et al. Serotypes, genotypes and antimicrobial resistance patterns of human diarrhoeagenic Escherichia coli isolates circulating in southeastern China. Clin Microbiol Infect. 2014;20(1):52-8.

38.   Sumrall ET, Gallo EB, Aboderin AO, Lamikanra A, Okeke IN. Dissemination of the transmissible quinolone-resistance gene qnrS1 by IncX plasmids in Nigeria. PLoS One. 2014;9(10):e110279.

39.   Christiana Cudjoe D, Balali GI, Titus OO, Osafo R, Taufiq M. Food Safety in Sub-Sahara Africa, An insight into Ghana and Nigeria. Environ Health Insights. 2022;16:11786302221142484.

40.   Henriques A, Fraqueza M. Listeria monocytogenes and ready-to-eat meat-based food products: Incidence and control. Listeria monocytogenes: incidence, growth behavior and control. 2015:71-103.

41.   Guinee PA, Kampelmacher EH, Willems HM. Six new Salmonella types, isolated in Ghana (S. volta, S. agona, S. wa, S. techimani, S. mampong and S. tafo). Antonie Van Leeuwenhoek. 1961;27:469-72.

42.   Mejia L, Vela G, Zapata S. High Occurrence of Multiresistant Salmonella Infantis in Retail Meat in Ecuador. Foodborne Pathog Dis. 2021;18(1):41-8.

43.   Zeng H, De Reu K, Gabriel S, Mattheus W, De Zutter L, Rasschaert G. Salmonella prevalence and persistence in industrialized poultry slaughterhouses. Poult Sci. 2021;100(4):100991.

44.   Casagrande Proietti P, Musa L, Stefanetti V, Orsini M, Toppi V, Branciari R, et al. mcr-1-Mediated Colistin Resistance and Genomic Characterization of Antimicrobial Resistance in ESBL-Producing Salmonella Infantis Strains from a Broiler Meat Production Chain in Italy. Antibiotics (Basel). 2022;11(6).

45.   Araujo S, I ATS, Tacao M, Patinha C, Alves A, Henriques I. Characterization of antibiotic resistant and pathogenic Escherichia coli in irrigation water and vegetables in household farms. Int J Food Microbiol. 2017;257:192-200.

46.   Silva A, Silva V, Pereira JE, Maltez L, Igrejas G, Valentao P, et al. Antimicrobial Resistance and Clonal Lineages of Escherichia coli from Food-Producing Animals. Antibiotics (Basel). 2023;12(6).

47.   Zhao X, Miao Y, Adam FEA, Zhao H, Zhou Z, Su M, et al. ESBLs-producing Escherichia coli from sheep-origin: Genome-wide virulence genes identification and in vivo virulence assessment in mice and Galleria mellonella. Transbound Emerg Dis. 2022;69(6):3606-17.

48.   Bihannic M, Ghanbarpour R, Auvray F, Cavalie L, Chatre P, Boury M, et al. Identification and detection of three new F17 fimbrial variants in Escherichia coli strains isolated from cattle. Vet Res. 2014;45(1):76.

49.   Pacheco AB, Soares KC, de Almeida DF, Viboud GI, Binsztein N, Ferreira LC. Clonal nature of enterotoxigenic Escherichia coli serotype O6: H16 revealed by randomly amplified polymorphic DNA analysis. Journal of clinical microbiology. 1998;36(7):2099-102.

50.   Beatty ME, Adcock PM, Smith SW, Quinlan K, Kamimoto LA, Rowe SY, et al. Epidemic diarrhea due to enterotoxigenic Escherichia coli. Clin Infect Dis. 2006;42(3):329-34.

51.   Willshaw GA, Cheasty T, Rowe B, Smith HR, Faithfull-Davies DN, Brooks TG. Isolation of enterotoxigenic Escherichia coli from British troops in Saudi Arabia. Epidemiol Infect. 1995;115(3):455-63.

52.   McMillan EA, Gupta SK, Williams LE, Jove T, Hiott LM, Woodley TA, et al. Antimicrobial Resistance Genes, Cassettes, and Plasmids Present in Salmonella enterica Associated With United States Food Animals. Front Microbiol. 2019;10:832.

53.   Dekker D, Eibach D, Boahen KG, Akenten CW, Pfeifer Y, Zautner AE, et al. Fluoroquinolone-Resistant Salmonella enterica, Campylobacter spp., and Arcobacter butzleri from Local and Imported Poultry Meat in Kumasi, Ghana. Foodborne Pathog Dis. 2019;16(5):352-8.

54.   Labar AS, Millman JS, Ruebush E, Opintan JA, Bishar RA, Aboderin AO, et al. Regional dissemination of a trimethoprim-resistance gene cassette via a successful transposable element. PLoS One. 2012;7(5):e38142.

55.   Monarrez R, Braun M, Coburn-Flynn O, Botelho J, Odetoyin BW, Otero-Vera JI, et al. A large self-transmissible resistance plasmid from Nigeria contains genes that ameliorate a carrying cost. Sci Rep. 2019;9(1):19624.

56.   Abraham S, Kirkwood RN, Laird T, Saputra S, Mitchell T, Singh M, et al. Dissemination and persistence of extended-spectrum cephalosporin-resistance encoding IncI1-bla(CTXM-1) plasmid among Escherichia coli in pigs. ISME J. 2018;12(10):2352-62.

57.   Foley SL, Kaldhone PR, Ricke SC, Han J. Incompatibility Group I1 (IncI1) Plasmids: Their Genetics, Biology, and Public Health Relevance. Microbiol Mol Biol Rev. 2021;85(2).

58.   Balbuena-Alonso MG, Cortes-Cortes G, Kim JW, Lozano-Zarain P, Camps M, Del Carmen Rocha-Gracia R. Genomic analysis of plasmid content in food isolates of E. coli strongly supports its role as a reservoir for the horizontal transfer of virulence and antibiotic resistance genes. Plasmid. 2022;123-124:102650.