Can reproductive interference be integrated into the Sterile Insect Technique for pest control? Insights from the spotted wing fly *Drosophila suzukii*

Flavia Cerasti [1], Massimo Cristofaro [2] Valentina Mastrantonio [1], Jessica Scifo[3], Adriano Verna[3], Daniele Canestrelli[4], Daniele Porretta[1*]

[1]Department of Environmental Biology, Sapienza University of Rome, Rome, Italy

[2]Biotechnology and Biological Control Agency (BBCA), Rome, Italy

[3]FSN-FISS-SNI Laboratory, Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Rome, Italy

[4]Department of Ecology and Biology, Tuscia University, Viterbo, Italy

[*]Correspondence: daniele.porretta@uniroma1.it

Abstract

BACKGROUND: Reproductive interference (i.e., sexual interaction between males of one species and females of another species that reduce the fitness of one or both the interacting individuals) is an important species interaction significantly affecting population dynamics and persistence. However, its exploitation in pest control remains overlooked. Here, we investigated the possible integration of reproductive interference into the Sterile Insect Technique (SIT) to develop a cross-species SIT. At this aim, we focused on the invasive pest species *Drosophila suzukii* and used *D. melanogaster* as the control species. First, we investigated the effect of irradiation on *D. melanogaster* male sterility and longevity Then, we tested the mating performance of irradiated males and their ability to reduce the *D. suzukii* fitness.

RESULTS: We found by microcosm experiments that: i) irradiation induced high levels of *D. melanogaster* male sterility without reducing longevity; ii) irradiated *D. melanogaster* males court *D. suzukii* females as much as *D. suzukii* males and they couple, mate and fecund heterospecific females; iii) irradiated *D. melanogaster* males significantly reduce the offspring of *D. suzukii* females under different species ratios.

CONCLUSIONS: Our results provide the first foundation to develop cross-species Sterile Insect Technique against *D. suzukii* and fuel to test this approach against other groups of pest species.

Keywords: Reproductive interference, pest control, Sterile Insect Technique, *Drosophila suzukii*, *Drosophila melanogaster*

1. Introduction

Species interactions, such as predation, parasitism, or competition, can significantly affect the population dynamics of the interacting species, affecting their abundance and persistence [1]. These interactions have been used for a long time to control pest species in agriculture and are the foundation of modern biological control approaches [2].

Along with the above species interactions, reproductive interference is now recognized as a major ecological process affecting population dynamics and species persistence [3-7] . Reproductive interference (or satyrisation in animals) consists of any sexual interaction between species that reduce the fitness of one or both the interacting individuals [8]. It occurs due to incomplete mating barriers between species and can occur at any stage of mate acquisition, from courtship to mating and hybridization [3]. Reproductive interference has been documented under laboratory and field conditions in a wide variety of sexually reproducing taxa [5,6]. Theoretical and empirical studies showed that reproductive interference, as competition, is density-dependent and can result in population or species exclusion [4,9].

The first attempts to exploit reproductive interference for pest control date back to the first mid of the XX century. During the 1930s – 40s, in pioneering works, F. L. Vanderplank and colleagues applied reproductive interference to suppress tsetse fly, *Glossina swynnertoni* Austen (reviewed in [10]). Laboratory experiments showed that offspring produced by cross-mating between *Glossina swynnertoni* (the vector species) and *G. morsitans* Westwood (a nonvector species) had low fertility, as all hybrid males and some hybrid females were sterile. Then, a large field experiment was carried out in the Shinyanga area, Tanzania, by releasing fertile *G. morsitans* pupae over seven months. After the mass release, the density of *G. swynnertoni* was

drastically reduced demonstrating the potential of hybrid sterility and sexual interference to
suppress tsetse flies [9,10].

Although the experience with tsetse flies showed the potential effectiveness of
reproductive interference for controlling a pest species, it did not lead to common exploitation of
this approach. A major concern about heterospecific sterility was that pre-mating isolation
mechanisms could constrain mating between released heterospecific males and wild females,
leading to failure to control the target species in the field [11,12]. A further concern has been the
risk of introducing non-native pest species. The mass release of fertile individuals of a closely
related species that is non-native or a pest itself, at least potentially, would lead to replacing one
pest species with another. These concerns and the success in the 1950s of Knipling and his team
in eradicating the screwworm *Cochliomyia hominivorax* (Coquerel) using ionizing radiation
[13], has led to the spread of homospecific Sterile Insect Technique (SIT) as the major approach
exploiting sterility in pest control during the last decades [14-18]. It consists of mass rearing,
sterilization by ionizing radiation, and massive release of conspecific individuals into the target
population. The unfertile mating between the released males and wild females leads to a
progressive decline of the target pest population [19,20].

Recently, however, a renewed interest has been in exploiting reproductive interference to
control pest species [8,21-24]. Mitchell et al. [8] reviewed the literature on reproductive
interference in natural populations. They highlighted the effects of these processes on population
decline in nature and proposed a framework for their use in pest control [8].

Interestingly, McInnis [21] tested the potential use of sterilized males of the oriental fruit
fly *Bactrocera dorsalis* (Hendel) against wild carambola fruit fly *B. carambolae*, suggesting that
the concern about using non-native or potential pest species to control another pest can be
overcome if the released heterospecific individuals are sterile, thus integrating reproductive
interference and SIT in a cross-species SIT approach. More recently, Honma et al. [22] have
proposed a framework for incorporating reproductive interference into a classic SIT program.
They argued that the sterile males released in a SIT program to suppress the wild-type population
of the same species could also lead to suppression of a closely related pest species through
reproductive interference (an approach that they called sterile interference).

In this paper, we aimed to explore the potential use of reproductive interference to develop
cross-species SIT against the spotted wing fly *Drosophila suzukii* Matsumura. This species is an

invasive pest that has spread in the last few decades from its native range in East Asia throughout

North America, Europe, and South America [25,26]. Unlike most Drosophilidae, *D. suzukii* can

lay eggs in unripe and healthy fruits, causing severe economic losses for fruit industries

worldwide [27,28]. We selected the fruit fly *D. melanogaster* as the species inducing

reproductive interference. Our previous study showed that post-mating isolation between *D.*

melanogaster and *D. suzukii* is complete, while there is an incomplete pre-mating isolation [24].

D. melanogaster males successfully courted *D. suzukii* females. Furthermore, they can

inseminate *D. suzukii* and reduce its offspring [24]. These results are a baseline for exploiting

sterilized *D. melanogaster* males in a cross-species SIT approach.

Here, we specifically aimed: *i*) to assess the effect of irradiation doses on the sterility degree

and longevity of *D. melanogaster* males. To this end, we irradiated virgin *D. melanogaster* males

with two gamma-ray doses (60 and 80 Gy) and assessed their fertility, through mating trials with

D. melanogaster females, and their longevity; *ii*) to analyze the mating performance of irradiated

D. melanogaster males in courting and mating with *D. suzukii* females; *iii*) to evaluate if

irradiated *D. melanogaster* males can reduce *D. suzukii* fitness with whom they mated. To this

end, we analyzed the effect of irradiated *D. melanogaster* males on the fertility of *D. suzukii*

females using different species ratios.

2.   Material and methods

2.1 Laboratory colonies

Drosophila suzukii and *D. melanogaster* individuals from the laboratory rearing facilities of the

Sapienza University of Rome were used in this study [24]. Both species were reared on an

artificial diet consisting of agar (7 g), table sugar (16 g), precooked ground maize (72 g), mother

yeast (18 g), soy flour (10 g), and methylparaben (2.5 g) [29]. The colonies were maintained in

entomological cages (30 x 30 x 30 cm), in a climate chamber at 25 ±1°C, 14:10 h light:dark

cycle.

2.2 Effect of irradiation dose on sterility and longevity of *D. melanogaster* males

To evaluate the effect of irradiation doses on the sterility degree of *D. melanogaster* males,

mating trials between irradiated males and fertile *D. melanogaster* females were carried out.

Virgin *D. melanogaster* adults were obtained by checking every thirty minutes the adult

emergence from the pupae confined in rearing falcons (50 ml), where they completed their full
larval development. As soon as new individuals emerged, we divided them into cages (15 x 15 x
15 cm) according to the gender. This procedure allows us to be sure to use only virgin males and
females. After selection, *D. melanogaster* males were sterilized using gamma radiation. Males
(48-72-96 hours old) were placed separately by age in 50 ml falcon tubes containing wet cotton
to avoid dehydration and transported to the Calliope Facility at ENEA Casaccia Research Centre
(Rome). The Calliope plant is a pool-type irradiation facility equipped with a ^{60}Co (mean energy
1.25 MeV) radio-isotopic source array in a high volume (7.0 m ´ 6.0 m ´ 3.9 m) shielded cell.
The irradiation cell can provide different dose rates by placing the samples in specific positions
and exposing them for varying periods [30]. We provided irradiation doses of 60 Gy and 80 Gy
to *D. melanogaster* males, according to Nelson [31] and Henneberry et al. [32].The dose rate was
175.03 Gy/h (2.92 Gy/min). After irradiation, five irradiated males were placed in falcon (50 ml)
with five non-irradiated *D. melanogaster* females. Each falcon contained food substrate to allow
females to lay eggs. In the control treatment, we used non-irradiated *D. melanogaster* males and
females. We performed twelve to sixteen replicates for each irradiation dose (60 and 80 Gy) and
the control treatment (0 Gy). The couples were left together for six days and then removed. New
individuals that emerged were removed and counted, checking oviposition substrates daily until
no newborn individuals were observed.

To evaluate the effect of irradiation doses on male longevity, we compared the average
lifespan between irradiated and non-irradiated *D. melanogaster* males. Virgin *D. melanogaster*
males were selected and irradiated as described above. A cage (30 x 30 x 30 cm) with twenty
males was set up for each dose treatment (60 and 80 Gy). Two more cages were set up as control,
one as "Control trip" and one as "Control home." In the "Control trip" cage, we placed twenty *D.*
melanogaster males who were previously transported to the Calliope facility without receiving
any irradiation dose. In this way, we evaluated the impact that the stress due to transport could
have on their longevity. Instead, in the "Control home" cage, we placed twenty non-irradiated *D.*
melanogaster males, which have not undergone transport stress but have always been kept in the
climate chamber conditions. Mortality was recorded every morning until all flies had died.

2.3 Mating performance of irradiated *D. melanogaster* males

All the following assays were carried out using only the *D. melanogaster* males irradiated with an 80 Gy irradiation dose, which led to higher sterility without affecting male longevity (see Results section).

First, mating performance was investigated by analyzing if the irradiated *D. melanogaster* males court *D. suzukii* females as much as *D. suzukii* males. At this aim, we compared the time the males of the two species spent courting *D. suzukii* females when they were alone or when the males of the other species co-occurred. Therefore, we set up three experimental setting by placing in 15 ml falcons: 1) one *D. suzukii* female and one *D. suzukii* male; 2) one *D. suzukii* female and one *D. melanogaster* male; 3) one *D. suzukii* female, one *D. suzukii* male, and one *D. melanogaster* male. The experiments were carried out with virgin individuals selected as described above. After 5 minutes of acclimatation, we recorded 10 minutes of the individual's behavior with an Olympus Tough TG-6 camera. The videos were used to analyze courtship behavior elements of *D. melanogaster* and *D. suzukii* males [33,34] using the BORIS software [35]. Twenty replicates were carried out.

Second, we investigated if irradiated *D. melanogaster* males were able to mate with *D. suzukii* females and fecund them. We selected virgin individuals of both species and irradiated the males at 48-72 hours old at 80 Gy as described previously. Then, we placed in 50 ml falcons with food substrates, one *D. suzukii* female and one irradiated *D. melanogaster* male. We left the couples inside the falcons for 6 days, to allow the females to oviposit. After that time, we removed the couples and, through a stereomicroscope Leica EZ4W at magnification 5x we checked daily the possible occurrence of eggs laid by *D. suzukii* females in food substrates. If present, eggs were photographed with a stereomicroscope digital camera and monitored for eclosion and for eventual larval development. Forty replicates were carried out.

2.4 Effect of irradiated *D. melanogaster* males on *D. suzukii* fitness

We evaluated whether the reproductive interactions between *D. melanogaster* males and *D. suzukii* individuals result in fitness costs for *D. suzukii* females. At this aim, we set up three experimental setting by placing in entomological cages (15 x 15 x 15 cm): 1) five pairs of no-irradiated virgin *D. suzukii* (selected as described above) without *D. melanogaster* males; 2) 40 irradiated *D. melanogaster* males and five pairs of no-irradiated virgin *D. suzukii* to test an 8:1 over-flooding ratio (sterile:wild males, OFR); 3) 60 irradiated *D. melanogaster* males with five

pairs of no-irradiated virgin *D. suzukii* to test an OFR of 12:1 [20]. Then, we compared the

number of newborn individuals that emerged from cages with only *D. suzukii* pairs and from

cages where *D. suzukii* pairs plus *D. melanogaster* irradiated males co-occurred.

2.5 Data analysis

To evaluate the effect of the irradiation doses on the sterility degree of *D. melanogaster* males, a

GLMM model (generalized mixed model with random effects; package lme4) [36] was applied.

We used the irradiation dose of *D. melanogaster* males as a fixed effect (0-60-80 Gy) on the

offspring produced by *D. melanogaster* females (the response variable), which was recorded as

an event with a continuous distribution. The age of experimental individuals (48, 72, and 96

hours old) was considered a random effect since they are a sampling of infinite possible

combinations, and we were not interested in studying them as such but in identifying whether

they constituted a source of significant variability. We analyzed the offspring produced with a

negative binomial distribution applied to the GLMM model. The model family was selected

comparing the AIC and BIC estimators and the likelihood ratio test. Multiple Comparisons of

Means Tukey Contrasts was performed as a post-hoc test using the *multcomp* package in R [37].

To evaluate the effect of the irradiation doses on the longevity of *D. melanogaster* males,

survival distributions of the four *D. melanogaster* groups ('Control cages', 'Control trip', '60

Gy', '80 Gy') were computed using the Kaplan–Meier method [38]. At this end, the *survival* and

survminer R packages were used [39]. The differences between survival distributions were

estimated using the Log-Rank Test.

To analyze the courtship behavior data, we used the Wilcoxon Mann–Whitney U test to

compare the time spent by *D. melanogaster* and *D. suzukii* males in courting *D. suzukii* females.

For what concern the analysis of the effect of *D. melanogaster on D. suzukii* fitness, a

GLMM analysis was performed using the *D. suzukii* offspring as response variable with a

continuous distribution. In the analysis, the number of *D. melanogaster* males (0, 40, or 60

individuals) in each replicate was considered a fixed effect (the explanatory variables). The age

of the experimental individuals (48, 72, and 96 hours old), as in the previous analysis on the *D.*

melanogaster sterility, was considered the random effect. Even in this case, we applied a

negative binomial distribution to the GLMM effect, comparing the models based on the optimal

parsimony principle (AIC and BIC estimators) and the likelihood ratio test. We performed the

Multiple Comparisons of Means Tukey Contrasts as a post-hoc test.

All analyses were performed using the software R version 3.6.2. [40].

3. Results

3.1 Effect of irradiation dose on sterility and longevity of *D. melanogaster* males

The mean number of offspring produced by *D. melanogaster* females was 128.66 ($\pm$ 10.67) in

the control and 35.81 ($\pm$ 4.08) (mean $\pm$ standard error) and 29.84 ($\pm$ 3.62) in the treatments at 60

and 80 Gy, respectively (Fig. 1). The GLMM model showed a significant effect of the irradiation

dose on the number of offspring produced by *D. melanogaster* females when the males were

irradiated at 60 Gy and 80 Gy (Table 1). The Tukey Multiple Comparison tests showed a

significant reduction in *D. melanogaster* offspring produced when *D. melanogaster* males were

irradiated at 60 Gy (z = -6.130, p = < 1e-04) and 80 Gy (z = - 9.057, p = < 1e-04) than when

there were no-irradiated *D. melanogaster* males (Fig. 1).

The Kaplan–Meier survival curves showed significant differences in the lifespan of *D.*

melanogaster males among treatments (60 Gy, 80 Gy, the Trip cage control and Home cage

control groups) (Mantel-Cox log-rank; $\chi 2$ = 17.8, d.f.= 3, P = 5e-04) (Fig. 2). The pairwise

comparisons test showed that the individuals irradiated with 60 and 80 Gy had higher survival

probability than control individuals. Significant differences were indeed observed between the

males irradiated at 60 Gy and those of the "Control home" (p= 0.0308) and the "Control trip"

(p=0.0151) groups, as well as between the males irradiated at 80 Gy and those of the two control

groups (control home p=0.0053; control trip p= 0.0028). No significant differences were

observed between the two control groups (p=0.7859) and between the two experimental

conditions (p=0.8541).

3.2 Mating performance of the irradiated *D. melanogaster* males

The mating performance of irradiated *D. melanogaster* males toward *D. suzukii* females was

investigated by courtship and mating trials.

First, we investigated if irradiated *D. melanogaster* males were able to court *D. suzukii* females

as much as *D. suzukii* males. Under all experimental conditions the typical behavior elements

during courtship were observed, including "orientation" (i.e., the male approaches the female,

quivering the abdominal and scissoring its wings), "tapping," (i.e., the male hits the female abdomen, or middle and hind legs by stretching his foreleg); "wing spreading", "wing scissoring" (i.e., the male is oriented toward the female front, quivers with the abdomen and scissors his wings keeping them at 180° for seconds to expose the upper side and wing spot toward to female) [33,34].

In the condition where *D. melanogaster* males were placed in 15 ml falcon only with *D. suzukii* females, the average courtship time spent by these males was 26.41% (± 5.90) (mean ± standard error). In the condition where *D. suzukii* males were placed in 15 ml falcon only with *D. suzukii* females, the average courtship time of males was 35.36% (± 8.17). No significant differences were observed between the two conditions (Wilcoxon Mann–Whitney test W = 118, p-value = 0.2673) (Figure 3a).

In the condition where *D. melanogaster* males and *D. suzukii* males were placed together in 15 ml falcon with *D. suzukii* females, the average courtship time spent by *D. melanogaster* males was 19.43% (± 6.16) and the average courtship time spent by *D. suzukii* males was 9.52% (± 2.29). There are no significant differences between the courtship time of *D. suzukii* and *D. melanogaster* males (Wilcoxon Mann–Whitney test W = 194.5, p-value = 0.9105) (Figure 3b).

Second, we investigated if irradiated *D. melanogaster* males were able to mate and fecund *D. suzukii* females. At this end, we analyzed the occurrence of oviposited eggs by *D. suzukii* females confined with irradiated *D. melanogaster* males. Monitoring daily the food substrates of every replicate, we found eggs in 12 out 40 falcons (30 %). We did not observe any subsequent larval development.

3.3 Effect of irradiated *D. melanogaster* males on *D. suzukii* fitness

The mean number of the individuals originating from five pairs of *D. suzukii* was 37.95 (± 3.62) (± standard error) in the control tests, while it was 16.11 (± 5.58) and 13.89 (±3.69), when 40 and 60 *D. melanogaster* males were added, respectively (Fig. 4). The GLMM model showed a significant effect on the *D. suzukii* offspring (Table 2). The Tukey Multiple Comparison tests showed a significant reduction of *D. suzukii* offspring when 40 (z = -2.758, p = 0.01601) and 60 (z = -3.218, p = 0.00376) *D. melanogaster* males were placed with *D. suzukii* couples (Fig. 4), while no significant differences were observed between treatments with 40 and 60 *D. melanogaster* irradiated males (z = -0.408, p = 0.91176).

4. **Discussion**

4.1 Irradiation dose, *D. melanogaster* male sterility and longevity

Cross-species SIT and SIT approaches can be developed under similar theoretical frameworks. Irradiation dose is an important factor that can affect the sterility degree, male longevity, and performance [41]. It is, therefore, necessary to find a high enough dose to induce sterility but that has a minimum impact on the biological quality of the irradiated males [42].

We found that 60 and 80 Gy irradiation doses led to high levels of *D. melanogaster* male sterility without reducing male longevity (Figures 1 and 2, respectively). Interestingly, irradiated males showed a higher lifespan than the control individuals. This effect of radiation has been observed also in other species, such as some *Anopheles* species [43,44] and the Stink bug Pest *Bagrada hilaris* [45], although the molecular basis underlying these physiological effects remain unclear.

As observed in previous studies [32], we also observed a residual fertility. Notably, the purpose of male sterilization is different in SIT and cross-species SIT approaches. In the former, male sterilization is the way to introduce sterility into the wild pest population. Residual fertility could hamper the SIT effectiveness as wild females could mate with released unsterilized males and produce fertile offspring. Conversely, in the cross-species SIT, the introduction of sterility in the target pest population is ensured by the reproductive post-zygotic isolation mechanisms between the released heterospecific males and the wild females of the target species. Male sterilization avoids potential adverse environmental effects if non-native or potentially pest species are used in the cross-species SIT. In the case of cross-species SIT using *D. melanogaster*, residual fertility would not reduce the effectiveness of the released males because also not irradiated male would induce sterility in *D. suzukii* females [24]. Furthermore, there are no concerns about the potential release of fertile *D. melanogaster* males. Indeed, *D. melanogaster* is not considered an agricultural pest as the female oviposits on rotten fruits [46].

4.2 Mating performance of the irradiated *D. melanogaster* males and effect on *D. suzukii* fitness

Cross-species SIT is based on the sterility between heterospecifics. The unfertile mating between
the released heterospecific males and wild females leads to a progressive decline of the target
pest population [19,20]. Therefore, the potential use of cross-species SIT as a control method
strictly depends on the mating ability of the released irradiated males [3]. Our results supported
that the *D. melanogaster* males irradiated at 80 Gy can be effective at different stages of mate
acquisition, from courtship to mating. The analysis of the courtship behavior showed that
irradiated *D. melanogaster* males courted *D. suzukii* females as much as *D. suzukii* males.
Furthermore, the results of mating trials showed that irradiated *D. melanogaster* males were able
to couple, mate and fecund *D. suzukii* females. Notably, we found that only 30% of *D. suzukii*
females oviposited eggs, and no larval development was observed in any tests, which confirms
that, contrary to pre-zygotic isolation, post-zygotic isolation between *D. suzukii* and *D.*
melanogaster is complete [24].

The results of our study clearly show that irradiated *D. melanogaster* males can lead to a
reproductive interference on *D. suzukii*. Indeed, *D. suzukii* had significantly reduced offspring in
the presence of irradiated *D. melanogaster* males, irrespective of the species ratio used (Fig. 4).
The reproductive interference on *D. suzukii* by irradiated *D. melanogaster* males is likely due to
interference during courtship and mating, as described above. However, other factors could
contribute to the offspring reduction of *D. suzukii* by *D. melanogaster* males. First, it has been
shown that *D. melanogaster*, during courtship, produces cis-vaccenyl acetate (cVA), which acts
as a repellent to *D. suzukii* female for laying eggs [47,48]. Furthermore, *D. melanogaster* males
release substances through seminal fluid that reduce female remating in homospecific matings
[49,50]. This latter phenomenon deserves future studies. Indeed, if it also occurs in heterospecific
matings between *D. suzukii* and *D. melanogaster*, it would be of great interest for cross-specific
SIT, preventing *D. suzukii* females from remating with co-specific wild males.

4.3 "The importance of being *melanogaster*"

334 The above arguments make us optimistic about the possible use of *D. melanogaster* in cross-
335 species SIT against *D. suzukii* and fuel to test this approach under well isolated confined-field
336 facilities.

337   However, let us highlight a further issue that makes cross-species SIT particularly
338 exciting to be explored: *D. melanogaster* is a model species. The huge amount of data available

on the genetics and biology of *D. melanogaster* can be exploited to address classical problems in developing SIT. Despite considerable research, for most species where SIT is used, we lack a deep understanding of the effects of radiation, rearing conditions (diet, light, density), as well as the effects of environmental conditions on individual biology and male performance in the field. Furthermore, some authors emphasized the need to build "a better male" [51] to overcome the prevailing approach based on overflooding ratios (sterile: wild males) in the field, which can hamper the economic feasibility of the SIT. In this context, *D. melanogaster*, whose biology has been dissected at the molecular, cellular, and physiological levels, could significantly contribute to unveiling these critical issues, making, in perspective, *D. melanogaster-suzukii* a model system for the development of cross-species SIT.

5. Conclusions

Reproductive interaction between heterospecific individuals can have multiple effects on pest management. When pre-mating barriers between interfertile species are weak, hybridization can occur and significantly threat pest control. It can favour pest outbreaks by improving population fitness and adaptation to environmental conditions, or by leading to control failures as consequence of introgression of pesticide resistance alleles from one species to another [52,53]. On the other hand, reproductive interaction can be beneficial for pest management [8,22]. Here, we highlight the potential use of reproductive interference in pest management by cross-species SIT. In last decades, the new sequencing technologies combined with classical reproductive incompatibility studies have improved our knowledge of pre- and post-zygotic isolation in sibling pest species allowing us to identify even more numerous potential systems for the development and application of cross-species SIT control approach [54,55].

Acknowledgments

We thank Elisa Michelangelo and Mark Eltenton for technical help. This study was carried out within the Agritech National Research Center and received funding from the European Union Next-Generation EU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4 – D.D. 1032 17 June 2022, CN00000022). This manuscript reflects only the authors' views and opinions, neither the European Union nor the European Commission can be considered responsible for them. FC was

funded by PhD resources within PON "RICERCA E INNOVAZIONE" 2014–2020", AZIONE IV.5 "DOTTORATI SU TEMATICHE GREEN, D.M. 1061 10 August 2021.

Data availability

Raw data will be permanently archived if the paper is accepted for publication.

Author contributions

DP and MC conceived the study. DP, MC, and DC designed the study. FC, VM, GS and AV collected the data. FC and VM analyzed the data. DP wrote the first draft of the manuscript. All authors contributed critically to the drafts and approved the final manuscript.

Conflict of Interest Statement

The authors declare no conflicts of interest.

References

1. Cain, ML, Bowman WD, Hacker SD. Ecology 2nd edition, ed. by Oxford University Press, Incorporate, Oxford, pp. 648 (2011).

2. Hoddle MS and Van Driesche RG, Biological control of insect pests. Encyclopedia of insects, ed. by Academic Press, Cambridge MA, pp. 91-101 (2009).

3. Gröning J, Hochkirch A, Reproductive Interference Between Animal Species. *Q Rev Biol* **83**:257–82 (2008).

4. Kishi S, Nishida T, Tsubaki Y, Reproductive interference determines persistence and exclusion in species interactions: Sexual interference governs competition. *J Anim Ecol* **78**:1043–9 (2009).

5. Kyogoku D, Reproductive interference: ecological and evolutionary consequences of interspecific promiscuity. *Popul Ecol* **57**:253–60 (2015).

6. Shuker DM, Burdfield-Steel ER, Reproductive interference in insects: Reproductive interference in insects. *Ecol Entomol* **42**:65–75 (2017).

7. Gómez-Llano M, Germain RM, Kyogoku D, McPeek MA, Siepielski AM, When Ecology Fails: How Reproductive Interactions Promote Species Coexistence. *Trends Ecol Evol* **36**:610–22 (2021).

8. Mitchell C, Leigh S, Alphey L, Haerty W, Chapman T, Reproductive interference and Satyrisation: mechanisms, outcomes and potential use for insect control. *J Pest Sci* **95**:1023–36 (2022).

9. Ribeiro JMC, Can Satyrs Control Pests and Vectors? *J Med Entomol* **25**:431–40 (1988).

10. Klassen W, Curtis CF, Hendrichs J, History of the sterile insect technique. Sterile insect technique, ed by CRC Press, Boca Raton, pp. 1-44 (2021).

11. Davidson G, The potential use of sterile hybrid males for the eradication of member species of the *Anopheles gambiae* complex. *Bull World Health Organ* **40**:221–8 (1969).

12. Davidson G, Odetoyinbo JA, Colussa B, Coz J, Field attempt to assess the mating competitiveness of sterile males produced by crossing 2 member species of the *Anopheles gambiae* complex. *Bull World Health Organ* **42**:55–67 (1970).

13. Knipling EF, Screwworm eradication: concepts and research leading to the sterile-male method. *In* Smithsonian Report for 1958. Smithsonian Inst., Washington DC, pp. 409–418 (1959).

14. Enkerlin WR, Impact of fruit fly control programmes using the sterile insect technique. Sterile insect technique, ed. by CRC Press, Boca Raton, pp. 979-1006 (2021).

15. Feldmann U, Dyck VA, Mattioli RC, Jannin J, Vreysen MJB, Impact of tsetse fly eradication programmes using the sterile insect technique. Sterile Insect Technique, ed. by CRC Press, Boca Raton, pp. 1051-1080 (2021).

16. Lees RS, Carvalho DO, Bouyer J, Potential impact of integrating the sterile insect technique into the fight against disease-transmitting mosquitoes». Sterile Insect Technique, ed by CRC Press, Boca Raton, pp. 1081-1118 (2021).

17. Simmons GS, Suckling DM, Carpenter JE, Addison MF, Dyck VA, Vreysen MJB, Improved quality management to enhance the efficacy of the sterile insect technique for lepidopteran pests. *J Appl Entomol* **134**:261–73 (2010).

18. Vargas-Terán M, Spradbery JP, Hofmann HC, Tweddle NE, Impact of screwworm eradication programmes using the sterile insect technique». Sterile insect technique, ed. by CRC Press, Boca Raton, pp. 949-978 (2021).

19. Bourtzis K, Vreysen MJB., Sterile Insect Technique (SIT) and Its Applications. *Insects*
12:638 (2021).

20. Dyck VA, Hendrichs J, Robinson AS, Sterile Insect Technique: Principles and Practice in
Area-Wide Integrated Pest Management, 2nd ed. by CRC Press, Boca Raton (2021).

21. McInnis DO, Rendon P, Jang E, Van Sauers-Muller A, Sugayama R, Malavasi A,
Interspecific Mating of Introduced, Sterile *Bactrocera dorsalis* with Wild *B. carambolae*
(Diptera: *Tephritidae*) in Suriname: a Potential Case for Cross-Species Sterile Insect
Technique. *Annal Entomol Soc Am* **92**:758–65 (1999).

22. Honma A, Kumano N, Noriyuki S, Killling two bugs with one stone: a perspective for
targeting multiple pest species by incorporating reproductive interference into sterile
insect technique: Reproductive interference and SIT. *Pest Manag Sci* **75**:571–7 (2019).

23. Tsurui-Sato K, Fujimoto S, Deki O, Suzuki T, Tatsuta H, Tsuji K, Reproductive
interference in live-bearing fish: the male guppy is a potential biological agent for
eradicating invasive mosquitofish. *Sci Rep* **9**:5439 (2019).

24. Cerasti F, Mastrantonio V, Dallai R, Cristofaro M, Porretta D, Applying Satyrization to
Insect Pest Control: The Case of the Spotted Wing Drosophila, *Drosophila suzukii*
Matsumura. *Insects* **14**: 569 (2023).

25. Fraimout A, Debat V, Fellous S, Hufbauer RA, Foucaud J, Pudlo P, et al., Deciphering
the routes of invasion of *Drosophila suzukii* by means of ABC random forest. *Mol Biol*
Evol **34**:980-996 (2017).

26. Tait G, Mermer S, Stockton D, Lee J, Avosani S, Abrieux A, et al., *Drosophila suzukii*
(Diptera: Drosophilidae): A Decade of Research Towards a Sustainable Integrated Pest
Management Program. Brewer M, ed. by *J Econ Entomol* **114**:1950–74 (2021).

27. Bolda M, Goodhue RE, Zalom FG, Spotted wing drosophila: potential economic impact
of a newly established pest. *Agric Resour Econ* **13**:5-8 (2010).

28. De Ros G, Conci S, Pantezzi T, Savini G, The economic impact of invasive pest
Drosophila suzukii on berry production in the Province of Trento, Italy. *J Berry Res*
5:89–96 (2015).

29. Silva-Soares NF, Nogueira-Alves A, Beldade P, Mirth CK, Adaptation to new nutritional
environments: larval performance, foraging decisions, and adult oviposition choices in
Drosophila suzukii. *BMC Ecol* **17**:21 (2017).

30. Baccaro S, Cemmi A, Gamma irradiation CALLIOPE facility at ENEA – Casaccia Research Centre, ed. by ENEA, Rome, pp. 48 (2019).

31. Nelson FRS, *Drosophila melanogaster*: Effects of Gamma Radiation on Fecundity and Longevity. *J Econ Entomol* **66**:257–8 (1973).

32. Henneberry TJ, Effects of Gamma Radiation on the Fertility and Longevity of *Drosophila melanogaster*. *J Econ Entomol* **56**: 279–81 (1963).

33. Lasbleiz C, Ferveur JF, Everaerts C, Courtship behaviour of *Drosophila melanogaster* revisited. *Anim Behav* **72**:1001–12 (2006).

34. Revadi S, Lebreton S, Witzgall P, Anfora G, Dekker T, Becher P, Sexual Behavior of *Drosophila suzukii*. *Insects* **6**:183–96 (2015).

35. Friard O, Gamba M, BORIS: a free, versatile open‐source event‐logging software for video/audio coding and live observations. *Methods Ecol Evol* **7**:1325–30 (2016).

36. Bates D, Mächler M, Bolker B, Walker S, Fitting Linear Mixed-Effects Models Using Lme4. *J Stat Softw* **67**:1–48 (2015).

37. Hothorn T, Bretz F, Westfall P, Simultaneous Inference in General Parametric Models. *Biometrical J* **50**:346–63 (2008).

38. Kaplan E and Meier P, Nonparametric estimation from incomplete observations. *J Amer Statist Assoc* **53**:457–481 (1958).

39. Kassambara A, Kosinski M, Biecek P, survminer: Drawing Survival Curves using 'ggplot2. R package version 0.4.9. (2021).

40. R Core Team, A Language and Environment for Statistical Computing. R-Foundation for Computer Statistics, (2019).

41. Ernawan B, Anggraeni T, Yusmalinar S, Ahmad I, Investigation of Developmental Stage/Age, Gamma Irradiation Dose, and Temperature in Sterilization of Male *Aedes aegypti* (Diptera: Culicidae) in a Sterile Insect Technique Program. *J Med Entomol* **59**:320–7 (2022).

42. Lance DR, McInnis DO, Biological basis of the sterile insect technique». Sterile Insect Technique, ed. by CRC Press, Boca Raton, pp.113-142 (2021).

43. Abdel-Malek AA, Tantawy AO, Wakid AM, Studies on the Eradication of Anopheles pharoensis Theobald by the Sterile-Male Technique Using Cobalt-60. I. Biological

Effects of Gamma Radiation on the Different Developmental Stages. *J Econ Entomol* **59**:672–8 (1966).

44. Helinski ME, Parker AG, Knols BG, Radiation-induced sterility for pupal and adult stages of the malaria mosquito *Anopheles arabiensis*. *Malar J* **5**:41 (2006).

45. Cristofaro M, Sforza RFH, Roselli G, Paolini A, Cemmi A, Musmeci S, et al., Effects of Gamma Irradiation on the Fecundity, Fertility, and Longevity of the Invasive Stink Bug Pest *Bagrada hilaris* (Burmeister) (Hemiptera: Pentatomidae). *Insects* **13**:787 (2022).

46. Markow TA, The secret lives of *Drosophila* flies. *eLife* **4**: e06793 (2015).

47. Dekker T, Revadi S, Mansourian S, Ramasamy S, Lebreton S, Becher PG, et al., Loss of *Drosophila* pheromone reverses its role in sexual communication in *Drosophila suzukii*. *Proc Roy Soc London Ser B* **282**:20143018 (2015).

48. Shaw B, Brain P, Wijnen H, Fountain MT, Reducing *Drosophila suzukii* emergence through inter-species competition: Reducing *D. suzukii* emergence through competition. *Pest Manag Sci* **74**:1466–71 (2018).

49. Manning A, A Sperm Factor Affecting the Receptivity of *Drosophila Melanogaster* Females. *Nature* **194**:252–3 (1962).

50. Gromko MH, Newport MEA, Kortier MG, Sperm Dependence of Female Receptivity to Remating in *Drosophila melanogaster*. *Evolution* **38**:1273 (1984).

51. Pérez‐Staples D, Shelly TE, Yuval B, Female mating failure and the failure of 'mating' in sterile insect programs. *Entomol Exp Appl* **146**:66–78 (2013).

52. Corrêa AS, Cordeiro EM, Omoto C, Agricultural insect hybridization and implications for pest management. *Pest Manag Sci* **75**:2857–64 (2019).

53. Porretta D, Canestrelli D, The ecological importance of hybridization. *Trends Ecol Evol* **38**:1097–108 (2023).

54. Ivey V, Hillier NK, Hybridization in heliothine moths: impacts on reproduction, pheromone communication, and pest management. *Front Ecol Evol* **11**:1208079 (2023).

55. Saveer AM, Becher PG, Birgersson G, Hansson BS, Witzgall P, Bengtsson M Mate recognition and reproductive isolation in the sibling species *Spodoptera littoralis* and *Spodoptera litura*. *Front Ecol Evol* (2014).

521 **Table 1. Effect of irradiation doses on sterility of *Drosophila melanogaster* males.** GLMM

522 model values are shown. Values in boldface indicate significant differences

Fixed Effects	Estimate	±SE	z Value	*p*-Value
(Intercept)	4.8674	0.1408	34,575	**< 2e-16**
IRR60	-1.1163	0.1821	-6.130	**8.8e-10**
IRR80	-1.7136	0.1892	-9.057	**< 2e-16**

Table 2. Effect of irradiated *Drosophila melanogaster* males on *D. suzukii* fitness.

GLMM model values are shown. Values in boldface indicate significant differences

Fixed Effects	Estimate	±SE	z Value	*p*-Value
(Intercept)	3.6361	0.1756	20,711	**< 2e-16**
40 *D. melanogaster* males	-0.8566	0.3105	-2.758	**-0.00581**
60 *D. melanogaster* males	-1.0050	0.3123	-3.218	**-0.00129**

Figure 1. Effect of irradiation on sterility of *Drosophila melanogaster* males. Comparison between the offspring originated from *D. melanogaster* females coupled with non- irradiated *D. melanogaster* males (green column) and irradiated at 60 Gy and 80 Gy (orange and blue column, respectively). Different letters mean significant differences by Tukey Multiple Comparison tests (p<0.05).

Figure 2. Effect of different treatments on the longevity of *Drosophila melanogaster* males.

Kaplan-Meier survival curves and confidence interval (CI 90%) for each condition are shown.

The p-value of the Log-Rank Test is also shown.

Figure 3. Time spent in courting *Drosophila suzukii* females by *D. suzukii* and irradiated *D.*

***melanogaster* males.** (a) Percentage of the total time spent courting *D. suzukii* females *by D.*

melanogaster and *D. suzukii* males when alone (green and orange column, respectively)

Wilcoxon Mann–Whitney test p-value > 0.05. (b) Percentage of the total time spent courting *D.*

suzukii females by *D. melanogaster* and males *D. suzukii* when placed together (green and

orange column, respectively) Wilcoxon Mann–Whitney test p-value > 0.05.

Figure 4. Effect of irradiated *Drosophila melanogaster* males on *D. suzukii* fitness. Offspring originated from five pairs of *D. suzukii* without irradiated *D. melanogaster* males (green column) and with 40 (orange column) or 60 (blue column) males. Different letters mean significant differences by Tukey Multiple Comparison tests.