Title

Image-Based Quantitative Single-Cell Method Showed Increase of Global Chromatin Accessibility in Tumor Compared to Normal Cells.

Authors

Mairead Commane[1], Vidula Jadhav[1], Katerina Leonova[1], Brian Buckley[2], Henry Withers[2, 3], Katerina Gurova[1, 3]

Affiliations

[1] – Department of Cell Stress Biology, Roswell Park Comprehensive Cancer Center, Elm and Carlton Str, Buffalo, NY, USA, 14263.

[2] – Drug Discovery Core Shared Resource, Roswell Park Comprehensive Cancer Center, Elm and Carlton Str, Buffalo, NY, USA, 14263.

[3] – Department of Bioinformatics and Biostatistics, Roswell Park Comprehensive Cancer Center, Elm and Carlton Str, Buffalo, NY, USA, 14263.

Corresponding author: Katerina Gurova, email: katerina.gurova@gmail.com

Competing interests: authors declare no competing financial interests in relation to the work described.

Abstract

The phenotypic plasticity of cancer cells has recently emerged as an important factor of treatment failure. The mechanisms of phenotypic plasticity are not fully understood. One of the hypotheses is that the degree of chromatin accessibility defines the easiness of cell transitions between different phenotypes. To test this, a method to compare overall chromatin accessibility between cells in a population or between cell populations is needed. We propose to measure chromatin accessibility by fluorescence signal from nuclei of cells stained with DNA binding fluorescent molecules. This method is based on the observations that small molecules bind nucleosome-free DNA more easily than nucleosomal DNA. Thus, nuclear fluorescence is proportional to the amount of nucleosome-free DNA, serving as a measure of chromatin accessibility. We optimized the method using several DNA intercalators and minor groove binders and known chromatin-modulating agents and demonstrated that chromatin accessibility is increased upon oncogene-induced transformation and further in tumor cells.

Introduction

Chromatin accessibility, or more precisely the accessibility of genomic DNA within chromatin for transcriptional machinery and other protein complexes, is a crucial determinant of a cell's transcriptional program and phenotype. Cells vary in their ability to change phenotype, known as phenotypic plasticity, which plays a significant role in various physiological and pathological processes such as differentiation, oncogenic transformation, tumor progression, inflammation, etc [1]. However, we lack a reliable quantitative measure to assess and compare the degree of phenotypic plasticity between different cells.

Theoretically, phenotypic plasticity refers to a cell's ability to switch between transcriptional programs. This can be measured by observing time-dependent changes in gene expression using techniques like bulk RNA-seq at different time points. While single-cell RNA sequencing (scRNA-seq) is a better approach for demonstrating the heterogeneity within a cell population, however it is not sufficiently quantitative. Additionally, both methods are expensive, time-consuming, and require a high degree of technical expertise and sophisticated data analyses.

Current methods to measure chromatin accessibility as a potential proxy for phenotypic plasticity are extremely laborious and not considered to be high throughput. Techniques like ATAC-seq or nuclease-seq are designed to identify differences in accessibility between specific genomic regions but fail to quantify altered genome-wide chromatin accessibility between cells [2]. This requires the use of spike-in controls to normalize signals between different samples [3, 4]. The extensive biochemical and bioinformatic manipulations needed often raise questions about the reliability of these comparisons.

Therefore, a simple and reliable method to measure general chromatin accessibility is highly demanded. Extensive data indicates that the degree of chromatin condensation not only

defines DNA accessibility to protein complexes, such as RNA polymerases, nucleases or transposases, but also to small molecules known as DNA ligands [5-8]. The two most specific types of DNA ligands are DNA intercalators, heterocyclic compounds that insert their planar bodies between nucleoside bases of DNA, and minor groove binders, crescent-shaped molecules that position themselves inside and along the minor groove of DNA. Among both classes, there are many well-known fluorescent molecules, and importantly, the fluorescent signal of many of these molecules increases many folds when they are bound to DNA [9].

Important property of both classes of molecules, discovered long ago, is that they bind free DNA more easily than DNA bound by proteins, especially histones within nucleosomes [5-8]. Thus, theoretically, the binding of these molecules to genomic DNA is proportional to DNA accessibility, and if they fluoresce only when bound, their fluorescence could be a measure of chromatin accessibility.

Several recent studies have proposed using DNA-binding small molecules to evaluate the chromatin state in cells (see Discussion) [10-13]. Building on these previous studies, we propose and optimize an approach to measure the intensity of total nuclear fluorescence of DNA-bound small molecules as a reporter of chromatin accessibility in individual cells. Using several DNA intercalators and minor groove binders, as well as established methods to manipulate chromatin condensation, we provide evidence that the nuclear fluorescence of DNA ligands can serve as an easy and quantifiable proxy for chromatin accessibility in cells and tissues. Importantly, this parameter is increased in cells with higher phenotypic plasticity, such as tumor and transformed cells.

Materials and Methods

Reagents

CBL0137 was provided by Incuron, Inc (Buffalo, NY). Propidium Iodide, RNAse A, Trichostatin A, Panobinostat, Valproic Acid, Vorinostat (SAHA), JQ1 were purchased from Sigma-Aldrich (St. Louis, MO), DAPI, Hoechst 33342 and Sybr Green were purchased from Invitrogen/TermoFisher (Grand Island, NY). EdU kit was from Click Chemistry Tools (Scottsdale, AZ).

Cells

HT1080, cells are from the American Type Culture Collection (ATCC). They were authenticated using short tandem repeat analysis (100% match). MCF10A cells are from ATCC. Primary human neonatal dermal fibroblasts (NDFs) were obtained from AllCells, LLC (Alameda, CA), as a pool of three separate donors. HT1080 and NDF cells were maintained in high glucose DMEM (Invitrogene) with 5% FBS (different vendors) and antibiotic solution in standard conditions. Composition of the medium for MCF10A cells is provided in Table S1.

Cell transformation

NDF and MCF10A cells were transduced with lentivirus harboring p53 dominant negative mutant GSE56 and HRasV12 oncogene connected via IRES [14]. Control cells were transduced with empty virus. Cells were transduced at MOI ~ 1. After that cells were split every 3-4 days for 2 weeks and then taken to the experiments.

MEFs were isolated from C57Bl/6 p53-heterozygous bred mice. Isolated MEFs were genotyped for the absence of p53, and p53-null MEFs were immortalized by stable transfection with SV40

large T antigen and then transformed with a vector expressing oncogenic mutant HRasV12 tagged with GFP [14]. GFP positive cells were sorted flow cytometry for further experiments as a model of transformed cells. To grow p53-null transformed MEFs (MEF p53KO-HRasV12) *in vivo*, 0.5×10^6 cells were implanted subcutaneously into wild-type C57BL/6 mice. When tumors reached 500-600mm^3, they were excised and underwent enzymatic digestion (1 mg/mL collagenase type IV + 0.02 mg/mL DNase) for 1 hour at 37°C. Digested tumors were filtered through 70 μm mesh filter and plated in DMEM supplemented with 10% FBS.

Cell treatment, fixation, and staining

For all imaging experiment cells were plated into 96 well black plates with clear bottom (Greiner Bio-One, Monroe, NC, cat # 655090), 5000 cells per well. For flow cytometry cells were plated in 6 well plate at 100,000 cells per well. Next day cells were treated with CBL0137 for 10-60 min or epigenetic drugs for 24 hours. After treatment medium with drugs was removed and cells were fixed with 4% paraformaldehyde (PFA) in PBS with 0.1% Triton X100 or 100% ice-cold methanol for 10 min at room temperature. For flow cytometry cells were first trypsinized, resuspended in medium with 5% FBS, washed from medium with PBS and fixed in 4% PFA. Then fixatives were removed, and cells were stained with DNA ligands with or without RNAse A (100 μg/ml). Unless otherwise stated, the following concentrations of DNA ligands were used Propidium Iodide (PI) – 1 μg/ml, DAPI and Hoechst 33342 – 1 μM, Sybr Green – 50 μM (1:100 dilution of solution provided by vendor). Plate was left in staining solution overnight at room temperature to allow RNA digestion.

Image acquisition and analyses

Plate imaging was done using Cytation 5 automated imager (Biotek/ Agilent Technologies, Santa

Clara, CA) using 4X objective in every well autofocus regimen. Each well image was montaged

from 4 photographs. Image processing and analysis was done using Gen5 Image Prime software

(Biotek/ Agilent). Data collection was done from ~300-5000 objects per well. Object masking was

done to include only nuclei. Before data analyses, objects from all wells were sorted by size and

object <15 m and > 40 μm were excluded to remove cell debris or cell clusters.

Microscopic imaging was done using Zeiss Axio Observer A1 inverted microscope, Zeiss MRC5

camera, and AxioVision Rel.4.8 software.

Statistical analyses

All experiments were repeated at least twice and included at least two replicate wells. The average

parameter for a well was used to calculate the mean of replicate wells. For object level analyses,

all objects from replicate wells were pooled together. The significance of difference between

conditions was calculated using t-test with SciPy function scipy.stats.ttest_ind [15].

Results

1. Theoretical assumptions

Multiple studies have shown that DNA ligands bind preferably naked DNA versus DNA

wrapped around nucleosomes [5-8]. Intercalator binding through the insertion of planar

heterocyclic moiety requires around a 2-fold increase of inter-base pair distance [16]. Spatial and

superhelical constraints of nucleosomal DNA limit this binding [12]. Some histone's amino acid

side chains intercalate between base pairs, thus competing with DNA ligands. The minor groove

of DNA facing histones is inaccessible to DNA minor groove ligands. Reduced binding of DNA

ligands to nucleosomal versus nucleosome-free DNA was firmly established in solution and cell-

based experiments [5-8].

However, high-affinity DNA ligands compete with histones for binding of nucleosomal DNA.

This causes DNA unwrapping from the histone core with loss of histones from chromatin, a

phenomenon known as chromatin damage [17]. Therefore, such DNA ligands themselves cause

an increase in chromatin accessibility, which makes measuring basal chromatin accessibility

difficult. Another problem is that many DNA ligands are a substrate of multidrug transporters and

their concentration in cells depends on the activity of multi-drug transporters, which is variable

between cells [18]. Both these problems can be resolved by cell fixation. The most suitable seems

fixation with short distance crosslinking agents, since they cause covalent links between

molecules, effectively gluing them together into an insoluble meshwork that preserves the

molecular anatomy of a cell as it existed at the moment of fixation. Thus, no further nucleosome

unwrapping is possible.

The next concern is the effect of the total amount of DNA in cells, which is different between

G1, S, and G2 cells and may be different between healthy and diseased cells (e.g., normal and

tumor cells due to aneuploidy, amplifications, and deletions). Thus, cells with longer total DNA

would bind more DNA ligands without having more accessible chromatin. Therefore,

normalization for the total DNA content per cell may be needed. However, the way to measure

nuclear fluorescence may mitigate this issue.

To access overall chromatin accessibility per cell nucleus, total amount of DNA-bound dye

needs to be quantified, what means collection of total fluorescent signal from a nucleus. Several

recently published methods based on high-resolution confocal microscopy measured fluorescence

from a section of a cell nucleus, not of the whole nucleus. Although images can be Z-stacked, they

are still the sum of sections, and the total signal depends on the number and thickness of sections

as well as thickness of a nucleus (Fig.1A).

Alternative methods are flow cytometry and non-confocal microscopy. Both methods allow

separation of G1, S, and G2 cells, so the problem of cell cycle-dependent DNA content may be

easily solved by assessing the cells in the same phase of the cell cycle, e.g., G1. However, flow

cytometry has poor distinction between cytoplasm and nucleus (unless using special equipment

like ImageStream) and ligand binding to mitochondrial DNA or double-stranded RNA in

cytoplasm may affect the signal.

Automated non-confocal microscopy like flow cytometry allows collection of fluorescent

signals from the whole volume of a cell (Fig.1A), however, the cytoplasm and the nucleus are

easier separated by masking since cells are attached to the matrix. It also allows significant process

miniaturization: as few as several cells can be detected in multi-well plates and it does not require

cell detachment from a matrix on which they normally grow. It can also be applied to tissues

without tissue disintegration (tissue slides). We tested effects of these theoretical considerations

on the performance of a method in conditions of controlled chromatin decondensation in normal

and tumor cells.

2. Optimization of an assay

2.1. Selection of a parameter of nuclear fluorescence

We used normal human diploid fibroblasts NDF and human fibrosarcoma cells HT1080.

The latter are near diploid (modal chromosome number = 46; range = 44 to 48) [19], thus

eliminating the potential effect of cell aneuploidy. Cells growing in a multiwell plate were fixed

and stained with intercalator propidium iodide (PI) or miner groove binder DAPI. Using non-

confocal automatic image acquisition, we collected the following parameters: nuclear area, mean

– the average intensity of all pixels per nucleus, and integral – the sum of all pixel intensities per

nucleus. Images were collected using a 4X objective, with pixel size equal to ~1.6 microns. These

data are presented in the form of histograms in Figure 1B and C. As we expected, the area and the

integral of cells had bimodal distributions reflecting the phase of a cell cycle similar to the

distribution of cell fluorescence intensity measured by flow cytometry. Therefore, average integral

values are significantly influence by the proportion of cycling cells in a population. However, the

distribution of mean values did not follow the same bimodal pattern, which is explained by the

cell-cycle dependent increase of the nuclear size. The normalization of integral by the area of a

nucleus produced a distribution very similar to the mean (Fig.1B, C). Thus, we selected mean

nuclear fluorescence as a potential measure of chromatin accessibility in cells due to less

dependence on the phase of cell cycle.

2.2.   Comparison of DNA ligand

We tested the performance of several intercalators, and minor groove binders known to be

highly fluorescent only when bound to nucleic acids. As an inducer of chromatin decondensation

we used curaxin CBL0137 which destabilizing effect on nucleosomes was shown in different

assays in solution and in cells [16, 20, 21]. We used a range of CBL0137 concentrations from 0.1

μM – no significant nucleosome unfolding, 0.3 μM – causing departure of H1 histone in some

cells [22], 1-3 μM - loss of outer histones H2A and H2B, and 10 μM – significant nucleosome

disassembly in cells ([23] and Fig. S1A). Importantly, these are approximate numbers, since these

effects do not happen equally genome-wide but depend on the pre-existing chromatin state and

cell type.

Cells were treated with CBL0137 for 30-60 min, since CBL0137 binds genomic DNA in cells

in a matter of seconds [16]. Preliminary experiments done using flow cytometry showed that a

shift in cell fluorescence is observed already after 10 minutes of incubation and is not significantly

increased anymore between 20 minutes and 1 hour (Fig. S1B and C). After treatment cells were

fixed and stained with Propidium Iodine (PI), Sybr Green, Hoechst 33342 or DAPI. Imaging was

done between 60 minutes and 24 hours with no difference between readings (data not shown).

The fluorescence of nuclei stained with different dyes increased with increasing doses of

CBL0137 (Fig. 2A, B). Though the increase in average fluorescent signal per well was very small

in case of minor groove binders (Fig.2A), it was highly significant when all individual cell values

were counted (Fig.2B). The increase was stronger with intercalators, PI and Sybr Green.

CBL0137-induced increase of Sybr Green fluorescence was the strongest when cells were

stained with 50 μM of Sybr Green. However, when we titrated the concentration of dyes in staining

solution, PI fluorescent intensity decreased linearly with dye concentration, while Sybr Green

fluorescent intensity was bell-shaped (Fig. S2A, B). We confirmed this non-linear change in

fluorescence by inspection of images taken with different exposures (Fig. S2 C, D). For both

intercalators, the CBL0137-induced increase of fluorescent signal became smaller with increase

in the intensity of fluorescence in basal conditions, suggesting a saturation effect, which was

especially strong in the case of Sybr Green (Fig. S2E, F). With this dye automatic quantitation

without visual confirmation may be misleading due to easy overexposure (compare visual effect

and quantitation in Fig. S2 D and F).

2.3.   Importance of RNAse A treatment

Although both intercalators and minor groove binders are known as specific DNA dyes, they

also bind to double-stranded RNA (dsRNA). Thus, we compared the performance of all DNA dyes

with and without treatment of fixed cells with RNAse A. Distribution of fluorescent signal was

changed upon RNAse A treatment in all conditions (Fig. 2B – compare blue and orange halves of

volcano plots), however the most significant change was seen for intercalators, especially for PI

(Fig. 2A, B and Fig. S3). This change was explained by stronger cytoplasmic signal in cells stained

with intercalators in the absence of RNAse A (Fig. S3). Cytoplasmic signal was gone upon RNAse

A treatment, suggesting that intercalators stain dsRNA stronger than minor groove binders.

Removal of RNA staining with RNAse A treatment made changes in chromatin accessibility

caused by CBL0137 more pronounced.

2.4.   The role of fixation method

Although we assumed that fixation with the short-range crosslinking agent would be the

best, we decided to see if other fixatives, such as methanol, which causes protein denaturation can

be used for the same purposes. Methanol fixation led to the increase in fluorescent signal in all

conditions compared with PFA fixed cells (Fig. 2C). There was similar increase in DAPI and PI

nuclear fluorescence upon CBL0137 treatment if cells were not treated with RNAse A. However,

RNAse A treatment led to further increase of the effect of CBL0137 if cells were fixed with PFA

and no additional increase if cells were fixed with methanol.

Thus, we defined the following conditions as the most sensitive and robust for detection of

changes in chromatin accessibility: (i) fixation of cells with 4% PFA, (ii) treatment of cells with

RNAse A, (iii) staining of cells with DNA intercalator dyes, with PI being less sensitive to changes

in dye concertation but more dependent on RNAse A treatment, and Sybr Green requiring more

accurate concentration optimization and monitoring of exposure time, due to easy overexposure.

3. Testing of the assay performance in cells treated with epigenetic drugs

After establishing optimal conditions, we tested assay performance upon treatment of cells

with several epigenetic drugs with known effects on chromatin accessibility. We used HDAC

inhibitors, trichostatin A (TSA), panobinostat (PNB), valproic acid (VA) and vorinostat

(SAHA) and inhibitor of BET domain proteins, JQ1, which does not directly increase chromatin

accessibility. Since most of these drugs inhibit enzymes, we treated cells for 24 hours and

monitored the toxicity by direct cell counting and microscopic imaging (Fig. 3A, B, E, F). As

expected, cells treated with all HDAC inhibitors accumulated more PI than untreated cells (Fig.

3C-F), in line with the known mechanism of action of these agents. This effect was obvious and

significant even when the drug caused growth arrest (reduced number of cells per well compared

with control in the absence of cell death) in the case of PNB and SAHA, or cell death in the case

of VA (Fig. 3A, B, E, F). Interestingly, all these drugs had a stronger effect on the viability and

chromatin accessibility of tumor HT1080 cells, than normal NDF cells (Fig. 3). JQ1 caused the

minimal changes in chromatin accessibility (Fig. 3 C, D) in line with its mechanism of action.

Thus, our method detected chromatin accessibility changes caused by several groups of

epigenetic drugs though neither of these drugs bind DNA.

4. Change of chromatin accessibility during cell cycle

The amount of DNA-bound small molecules increases with DNA length. Since DNA length

increases with cell cycle progression populations of cells with more cycling cells would bind more

DNA dye due to longer average DNA length per cell. We tried to distinguish the effect of increased

DNA length from increased DNA accessibility by comparison of PI incorporation at different

phases of the cell cycle and in growth-arrested and cycling cell populations.

First, we identified cells in the S phase by labeling them with EdU for 1 hour and then

measuring the fluorescent signal of DNA ligand in cells positive and negative for EdU. This

comparison showed that EdU-positive cells accumulate more dye molecules, which was not

surprising since they have more DNA than cells in G1, the predominant state of EdU-negative cells (~ 70 % of HT1080 cells in basal conditions) (Fig. 4A, B). To mitigate this factor to a certain extent, we used the integral of cell fluorescence to identify cells in G1 or G2/M phases of cell cycle (Fig. 4B). EdU positive cells withing these groups are cells which are early in S phase and therefore having DNA content close to G1 cells or very late in S phase, i.e., having DNA content close to G2/M cells. When we compared mean fluorescent signal of these cells, we did not see significant difference between EdU positive and negative cells (Fig. 4C, D), suggesting that DNA replication per se does not make chromatin more accessible at nuclei level.

Another approach was to compare cells arrested with combination of contact inhibition and serum starvation (48 hours in medium with 1% FBS) and normally growing cycling cells. Interestingly, there was or slight decrease in basal condition and no difference upon CBL0137 treatment in the dye accumulation in growing versus arrested normal NDF cells, while growing tumor HT1080 cells had more open chromatin. This shift was true even if we looked only at the position of G1 cells using the integral of fluorescence, and it was increased at high dose of CBL0137 (Fig. S4).

Thus, we observed more open chromatin state in cycling versus resting cells, especially in the case of tumor cells and this increase cannot be explained by increased DNA length.

5. Comparison of chromatin accessibility in normal and transformed or tumor cells

Existing data from literature suggest that chromatin may be overall more open in tumor than in normal cells (reviewed in [24]), though data comparing general chromatin accessibility between normal, transformed and tumor cells are missing. Thus, we decided to do this using our method.

First, we compared NDF and HT1080 cells using DAPI and PI. In both cases HT1080

accumulated more dye than NDF cells in basal conditions and upon CBL0137 treatment (Fig.5A,

B). This was true with both fixatives, PFA and methanol, and in the presence or absence of RNAse

A (Fig. S5A-D). Next, we compared syngeneic normal and transformed cells. NDF and

immortalized mammary epithelial cell line, MCF10A, were transduced with combination of p53

dominant negative inhibitor, GSE56 [25] and mutant HRasV12 oncogene (GR), or empty virus

(EV). Cells were passaged for 2 weeks to allow transformation phenotype to evolve and then they

were fixed and stained with DAPI. Though nuclear accumulation of this dye has been changed less

upon CBL0137 treatment than of PI, we observed significant difference between cells transduced

with empty vector and two independently generated transformed cell lines (GR1 and GR2). Both,

transformed fibroblasts and epithelial cells accumulated more dye than non-transformed cells (Fig.

5C, D), suggesting more open chromatin state in transformed versus normal cells.

Lastly, we compared chromatin accessibility of syngeneic mouse normal, transformed and

tumor cells. We used mouse embryo fibroblasts   from wild type and p53 knockout animals. The

latter were transduced with mutant HRasV12 to get transformed cells. These transformed cells

were implanted subcutaneously into C57Black/6 mice and after tumor reached the size of 500

mm^3, it was excised, disaggregated and established as cell line *in vitro*. Similar to human cells, we

saw increase in chromatin accessibility upon *in vitro* transformation of mouse cells and further

increase in tumor cells (Fig. 5E).

Since, as we saw in the previous section, mean fluorescence is higher in proliferating than

non-proliferating cells, we decided to confirm that the increase in chromatin accessibility is not

only due to the higher proportion of proliferating cells in transformed and tumor cell populations.

For this, we compared fluorescent intensities of G1 peaks obtained from the integral of

fluorescence. In all cases we saw shift in the position of G1 peaks corresponding to the shift of

mean fluorescence (Fig. 5F and S5 E-H), confirming increase in the chromatin accessibility during

oncogene induced transformation of human and mouse cells.

Discussion

The method which we described here is extremely simple and was used extensively for the

different purpose: to assess distribution of cells in populations along different phases of a cell cycle

via flow cytometry. In most cases laser settings were adjusted for every different cell type and

therefore shifts in the positions of histogram peaks between cell types, i.e., different fluorescent

intensity of different cells was largely ignored by cell and molecular biologists. At the same time

the fact that DNA ligands bind naked DNA better that DNA wrapped around histone core was very

firmly established long ago [5-8].

Differential accessibility of DNA in chromatin to different molecules is currently mostly

assessed using proteins, such as nucleases or transposases. These methods are expensive, time

consuming and require high level expertise in molecular biology and bioinformatics for data

processing. Most importantly they are not tuned for the comparison of global level of chromatin

accessibility between different samples.

Nucleosome dependent binding of small molecules to DNA in cells manifested with the

emission of light presents an easy opportunity to assess an abundance of nucleosome-free DNA in

different cells by comparing their nuclear fluorescence. This comparison can be done between cell

populations and between individual cells in population. Not surprisingly there were already several

attempts to use fluorescent DNA ligands for the assessment of chromatin organization in cells. The

closest approach was published by Rosevalentine Bosire et al [12]. They used different methods to decondense chromatin in fixed cells to demonstrate that intercalator incorporation into nuclear DNA in cells is constrained by superhelical stress due to DNA wrapping within nucleosome. Unlike us, they used laser scanning cytometry, i.e. they collected the signal from a section of a nucleus, and not from the whole nucleus. They observed that fluorescent signal of intercalators depends on the degree of chromatin condensation and negatively correlates with the number of histones per nucleus. Contrary to us they did not see the same dependence for minor groove binder DAPI and when they fixed cells with methanol, probably due to assessment of a signal from one section of a nucleus.

Differential binding of an intercalator to open and closed chromatin was used by Gali Bai et al to develop a different method, adduct sequencing (Add-seq), to probe chromatin accessibility by treating chromatin with the small molecule angelicin, which then was covalently bound to DNA. DNA regions with bound angelicin was detected by nanopore sequencing [10]. Though in general similar, this method requires nuclei isolation, DNA sequencing, and it is unclear whether it is good for measurement of differences in general chromatin compaction between cells.

Application of DNA ligands to discern chromatin structure between tumor and normal cells was proposed by Jianquan Xu et al [11]. They used minor groove binder Hoechst and stochastic optical reconstruction microscopy (STORM) [26] to evaluate chromatin organization. However, instead of Hoechst fluorescence, which is not optimal for super resolution, they attached another fluorophore, Cy5 to Hoechst. Cy5 does not bind DNA itself but provided necessary fluorescent parameters for STORM.   They observed significant differences between tumor and non-tumor cells, but their approach to image analysis was focused on the detection of different patterns of chromatin organization and not on the total cumulative fluorescent signal from a cell [11].

However, their description of the observed differences was in line with the quantitative differences which we observed here. They found that in normal cells chromatin domains are more compact, especially at the nuclear periphery. In precancerous cells, chromatin compaction was slightly disrupted, and in cancer cells chromatin at nuclear periphery was indistinguishable from the interior. This can be interpreted as general chromatin de-compaction in the process of tumorigenesis. This is an excellent method to detect and describe changes in the patterns of chromatin organization, however, it is much more complicated in data acquisition and analysis, than the method which we propose. Our method is simple enough that it can be done in a clinical lab. With additional optimization it can be applied to tissue slides, and data processing can be easily automated. We believe that the degree of chromatin decondensation may be an important measure of tumor aggressiveness [24]. Therefore, quantitative assessment of the chromatin decondensation in tumor cells and/or the proportion of cells in tumors with decondensed chromatin may be used as a prognostic marker.

Metastasis, invasion to the neighboring organs, and development of resistance to multiple therapies are factors responsible for poor prognosis of cancer patients. These traits are not often due to the accumulation of new mutations and selection of resistant clones, but to the adaptation of tumor cells to new conditions via changing of their phenotype, i.e., phenotypic plasticity [1, 27]. Although detailed mechanisms of phenotypic plasticity are still obscure, easiness of transitions between transcriptional programs is probably an obligatory factor of phenotypic plasticity. Transcriptional programs are controlled by chromatin organization at different genomic regions and therefore stable programs should be associated with stable chromatin with multiple constraints preventing easy activation and de-activation of genes. Easy transitions between active and inactive state of transcription may occur when silent and active states of genomic regions are not enforced

strongly enough by chromatin organization, which may include unstable nucleosomes at

regulatory region and gene bodies leading to easier access of transcriptional machinery to DNA,

mobile, flexible chromatin fibers and "poorly locked" chromatin loops making random collisions

of promoters and enhancer more probable and burst of transcription more stochastic. Thus, our

long-term hypothesis was that general destabilization of chromatin through reduced number of

histones per cell, prevalence of histone modifications or mutations making nucleosomes less

stable, loss of heterochromatin proteins or overexpression of proteins making nucleosomes more

open, e.g., HMG proteins, these all would result in more open chromatin state in tumor versus non-

tumor cells. Our quantitative method confirmed this hypothesis in human and mouse models. Our

next hypothesis is that the degree of chromatin decondensation correlates with tumor prognosis

can now be tested by staining of patient's tumor samples on slides and assessing correlation

between the degree of chromatin accessibility and patient outcomes. If proven correct, this may

become one of the simplest and universal prognostic biomarkers, which in case of development of

appropriate fluorescent probe can be even applied for *in vivo* imaging.

Although we believe that our method is simple, easy, quantitative and accurately revealing the

state of chromatin in cell populations and difference in chromatin condensation between individual

cells in a population, there are still limitations or uncertainties which need to be clarified in the

future studies. Ideally normalization for the total DNA length needs to be done. Better

understanding of the mechanisms of fluorescence as a result of interactions of molecules of DNA

ligand between themselves and with DNA is needed to understand which signals report accessible

chromatin and which is influenced by ligand-ligand interactions, as we see in the case of Sybr

Green. Another question is which DNA ligand better reflects actual chromatin state in different

conditions, minor groove binders or DNA intercalators. There is very limited data to understand

which properties of nucleosomal DNA define these molecules binding. Most probably their

binding would inform us about different complementary properties of nucleosomes. which can be

put together to better understand chromatin organization and dynamics. Although fluorescent

signal from the same cells in different experiments was in the same range, there were some

differences between experiments and between wells in the same experiments. For comparison of

different samples, especially in clinical conditions a set of controls for normalization of staining

and imaging and calibration of signal need to be developed.

Acknowledgments

We would like to acknowledge Dr. Razvan Chereji for helping with Fig. S1A

presentation, Drs. Andrei Gudkov and Subhamoy Dasgupta for the critical discussion of

the manuscript, Bruce Specht, Mary Morgan, Sarah Mercy and Dale Henry for the

administrative support.

Funding:

National Institutes of Health grant R01 CA266216 (KG)

National Institute of Health grant NCI NIH Core P30CA16056 to RPCCC, which

partially covers the costs Genomics, Bioinformatics, Flow Cytometry, Drug Discovery

Lab, Gene Modulation

Author contributions:

MC and VJ were responsible for running imaging and flow cytometry experiments. KL

generated mouse transformed and tumor cells. BB did imaging and image analyses, HW

performed analysis of some data, generated R script to define x-value of positions of G1

and G2 peaks on a histogram and editing of the manuscript. KG conceptualized the study,

analyzed all data, wrote the manuscript and prepared all figure.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: Transformed or tumor cells generated for this study are

available upon request. Raw imaging data are available upon request. R script to define x-

value of positions of G1 and G2 peaks on a histogram is available here:

https://github.com/HGWithers/cellphaseR

References

1   Gupta PB, Pastushenko I, Skibinski A, Blanpain C, Kuperwasser C. Phenotypic
     Plasticity: Driver of Cancer Initiation, Progression, and Therapy Resistance. *Cell Stem
     Cell* 2019; 24: 65-78.

2   Minnoye L, Marinov GK, Krausgruber T, Pan L, Marand AP, Secchia S *et al*. Chromatin
     accessibility profiling methods. *Nat Rev Methods Primers* 2021; 1.

3   Chen K, Hu Z, Xia Z, Zhao D, Li W, Tyler JK. The Overlooked Fact: Fundamental Need
     for Spike-In Control for Virtually All Genome-Wide Analyses. *Mol Cell Biol* 2015; 36:
     662-667.

4   Chereji RV, Bryson TD, Henikoff S. Quantitative MNase-seq accurately maps
     nucleosome occupancy levels. *Genome Biol* 2019; 20: 198.

5   Giangare MC, Prosperi E, Pedrali-Noy G, Bottiroli G. Flow cytometric evaluation of
     DNA stainability with propidium iodide after histone H1 extraction. *Cytometry* 1989; 10:
     726-730.

6   Gabbay EJ, Wilson WD. Intercalating agents as probes of chromatin structure. *Methods
     Cell Biol* 1978; 18: 351-384.

Lurquin PF. The use of intercalating dye molecules in the study of chromatin structure. *Chem Biol Interact* 1974; 8: 303-312.

Lawrence JJ, Louis M. Ethidium bromide as a probe of chromatin structure. *FEBS Lett* 1974; 40: 9-12.

Bag S, Bhowmik S. Fluorescence Spectroscopy: A Useful Method to Explore the Interactions of Small Molecule Ligands with DNA Structures. *Methods Mol Biol* 2024; 2719: 33-49.

Bai G, Dhillon N, Felton C, Meissner B, Saint-John B, Shelansky R *et al*. Probing chromatin accessibility with small molecule DNA intercalation and nanopore sequencing. *bioRxiv* 2024.

Xu J, Sun X, Kim K, Brand RM, Hartman D, Ma H *et al*. Ultrastructural visualization of chromatin in cancer pathogenesis using a simple small-molecule fluorescent probe. *Sci Adv* 2022; 8: eabm8293.

Bosire R, Nanasi P, Jr., Imre L, Dienes B, Szoor A, Mazlo A *et al*. Intercalation of small molecules into DNA in chromatin is primarily controlled by superhelical constraint. *PLoS One* 2019; 14: e0224936.

Imai R, Nozaki T, Tani T, Kaizu K, Hibino K, Ide S *et al*. Density imaging of heterochromatin in live cells using orientation-independent-DIC microscopy. *Mol Biol Cell* 2017; 28: 3349-3359.

Boiko AD, Porteous S, Razorenova OV, Krivokrysenko VI, Williams BR, Gudkov AV. A systematic search for downstream mediators of tumor suppressor function of p53 reveals a major role of BTG2 in suppression of Ras-induced transformation. *Genes Dev* 2006; 20: 236-252.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D *et al*. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* 2020; 17: 261-272.

Safina A, Cheney P, Pal M, Brodsky L, Ivanov A, Kirsanov K *et al*. FACT is a sensor of DNA torsional stress in eukaryotic cells. *Nucleic Acids Res* 2017; 45: 1925-1945.

Neefjes J, Gurova K, Sarthy J, Szabo G, Henikoff S. Chromatin as an old and new anticancer target. *Trends Cancer* 2024.

Szabo E, Kulin A, Jezso B, Kucsma N, Sarkadi B, Varady G. Selective Fluorescent Probes for High-Throughput Functional Diagnostics of the Human Multidrug Transporter P-Glycoprotein (ABCB1). *Int J Mol Sci* 2022; 23.

531   19   Chen TR, Hay RJ, Macy ML. Intercellular karyotypic similarity in near-diploid cell lines
532   of human tumor origins. *Cancer Genet Cytogenet* 1983; 10: 351-362.
533

534   20   Chang HW, Nizovtseva EV, Razin SV, Formosa T, Gurova KV, Studitsky VM. Histone
535   Chaperone FACT and Curaxins: Effects on Genome Structure and Function. *J Cancer
536   Metastasis Treat* 2019; 5.
537

538   21   Gasparian AV, Burkhart CA, Purmal AA, Brodsky L, Pal M, Saranadasa M *et al*.
539   Curaxins: anticancer compounds that simultaneously suppress NF-kappaB and activate
540   p53 by targeting FACT. *Sci Transl Med* 2011; 3: 95ra74.
541

542   22   Leonova K, Safina A, Nesher E, Sandlesh P, Pratt R, Burkhart C *et al*. TRAIN
543   (Transcription of Repeats Activates INterferon) in response to chromatin destabilization
544   induced by small molecules in mammalian cells. *Elife* 2018; 7.
545

546   23   Nesher E, Safina A, Aljahdali I, Portwood S, Wang ES, Koman I *et al*. Role of chromatin
547   damage and chromatin trapping of FACT in mediating the anticancer cytotoxicity of
548   DNA-binding small molecule drugs. *Cancer Res* 2018.
549

550   24   Gurova K. Can aggressive cancers be identified by the "aggressiveness" of their
551   chromatin? *Bioessays* 2022; 44: e2100212.
552

553   25   Ossovskaya VS, Mazo IA, Chernov MV, Chernova OB, Strezoska Z, Kondratov R *et al*.
554   Use of genetic suppressor elements to dissect distinct biological effects of separate p53
555   domains. *Proc Natl Acad Sci U S A* 1996; 93: 10309-10314.
556

557   26   Bates M, Jones SA, Zhuang X. Stochastic optical reconstruction microscopy (STORM): a
558   method for superresolution fluorescence imaging. *Cold Spring Harb Protoc* 2013; 2013:
559   498-520.
560

561   27   Shah S, Philipp LM, Giaimo S, Sebens S, Traulsen A, Raatz M. Understanding and
562   leveraging phenotypic plasticity during metastasis formation. *NPJ Syst Biol Appl* 2023; 9:
563   48.
564

566   **Figure Legends**

567   **Fig. 1. Measurement of nuclear fluorescence using non-confocal automated imaging. A**. Total

568   nuclear fluorescence is more accurately collected using non-confocal imaging. **B, C**. Distribution

569   of parameters of nuclear fluorescence in NDF and HT1080 cells in basal conditions. Cells were

570   fixed and stained with DAPI (**B**) or PI (**C**) 24 hours after plating.

Figure 2. Optimization of the assessment of chromatin accessibility using fluorescent DNA ligands. Nuclear fluorescence was increased with a short-term treatment of NDF and HT1080 cells with nucleosome-destabilizing compound CBL0137. **A, B**. Comparison of the performance of different DNA dyes in the presence and absence of RNAse **A**. Fold change in average nuclear fluorescence per well relative to untreated cells. Data from two replicate wells with error bars showing variability between replicates. A representative from > 3 similar experiments. **B**. Split violin plots with quartiles showing mean nuclear fluorescence of cells treated with different concentrations of CBL0137 for 60 min. Numbers above violins are p-values of the t-test comparing treated cells to the untreated control stained either without RNAse A (blue numbers) or with RNAse A (orange numbers). P-values were rounded to 4 decimal places. ns - p-value >0.05. **C**. Split violin plots with quartiles showing mean nuclear fluorescence per cell fixed with either 4% PFA or 100% methanol. Numbers on the plot are p-values of the t-test comparing treated cells to untreated control fixed either with PFA (blue numbers) or methanol (orange numbers). **D, E**. Fold change in average nuclear fluorescence per well of cells treated with CBL0137 relative to untreated cells and stained in the presence of absence of RNAse A. **D**. Cells were stained with DAPI. **E**. Cells were stained with PI. Data from two replicate wells with error bars showing variability between wells.

Figure 3. Effect of epigenetic drugs on chromatin accessibility measured with PI. NDF and HT 1080 cells were treated with CBL0137 (500 nM) , trichostatin A (TSA, 500 nM), panobinostat (PNB, 20 nM), valproic acid (VA, 0.5 mM), SAHA (1000 nM), or JQ1 (1000 nM) for 24 hours before fixation with PFA and staining with PI in the presence of RNAse A. **A, B**. Number of nuclei per well of NDF (**A**) or HT1080 (**B**) cells, untreated or treated with the drugs. Bars are mean value of 4 replicate wells, error bars are SDV. Number above the bar is p-value of t-test comparing

each drug with the control untreated cells, rounded to 2 decimal points. **C, D**. Violin plots with

quartiles of distribution of PI nuclear fluorescence of cells in all replicate wells of NDF (**C**) or

HT1080 (**D**) cells treated with epigenetic drugs. Number above violin is p-value of t-test

comparing each drug with the control untreated cells, rounded to 4 decimal points. Pink and blue

transparent squares show position of quartiles 0.5 and 0.75 (pink), and 0.5 and 0.25 (blue) in

control untreated samples. **E, F**. Microscopic images of PI-stained wells from the plate of cells

treated with epigenetic drugs. The same exposition time was used for all images of NDF (**E**) and

HT1080 (**F**) cells.

Figure 4. Dependence of chromatin accessibility of cell proliferation. **A-D**. HT1080 cells were

incubated in the presence of EdU for 1 hour. After that cells were fixed and stained for EdU and

DAPI. **A**. Violin plots with quartiles of nuclear fluorescence of of EdU positive and negative cells.

Bleu squares correspond to the positions of quartiles (0.25, 0.5 and 0.75) in EdU negative cells. **B**.

Histogram of distribution of total nuclear fluorescence (Integral) in EdU negative and positive

cells. Blue squares show values of nuclear fluorescence used to select cells with DNA content

close to G1 or G2/M cells. **C**, D. Violin plots with quartiles showing mean nuclear fluorescence

of EdU positive and negative cells with DNA content close to G1 (**C**) or G2 /M (**D**). ns – p-value

> 0.05. **E, F**. Comparison of mean nuclear fluorescence of NDF (**E**) or HT1080 (**F**) cells growing

in normal conditions or arrested with dense plating and medium with 1% FBS for 48 hours. Before

staining cells were treated with the indicated concentrations of CBL0137 for 60 minutes. Numbers

are p-values of t-test comparing growing and arrested cells at each concentration of CBL0137.

Figure 5. Comparison of chromatin accessibility between normal, transformed and tumor

cells. **A, B**. Split violin plots of mean nuclear fluorescence of NDF and HT1080 cells treated with

different concentrations of CBL0137 for 60 minutes and stained with DAPI (**A**) or PI (**B**). Numbers

are p-value of t-test comparing NDF and HT1080 cells. **C, D**. Violin plots with quartiles showing

mean nuclear fluorescence (DAPI) of two biological replicates of transformed NDF (**C**) or

MCF10A (**D**) cells (GR1, GR2) and control non transformed cells (EV). Numbers are p-values of

t-test comparing each of transformed variants (GR) with control non-transformed EV cells. **E, F**.

Comparison of nuclear fluorescence (PI) of mouse normal, transformed and tumor cells. **E**. Violin

plots with quartiles of mean nuclear fluorescence. Numbers above the lines show p-values of

ANOVA test comparing all groups. **F**. Distribution of total nuclear fluorescence. Numbers show

X-values of corresponding peaks on histograms. Dotted lines correspond to the positions of G1

peaks of normal MEF (red), transformed (blue) and tumor (green) cells.

 Figures

Fig. 1. Measurement of nuclear fluorescence using non-confocal automated imaging. A. Total nuclear fluorescence is more accurately collected using non-confocal imaging. **B, C**. Distribution of parameters of nuclear fluorescence in NDF and HT1080 cells in basal conditions. Cells were fixed and stained with DAPI (**B**) or PI (**C**) 24 hours after plating.

Figure 2. Optimization of the assessment of chromatin accessibility using fluorescent DNA ligands. Nuclear

fluorescence was increased with a short-term treatment of NDF and HT1080 cells with nucleosome-destabilizing

compound CBL0137. **A, B**. Comparison of the performance of different DNA dyes in the presence and absence of

RNAse **A**. Fold change in average nuclear fluorescence per well relative to untreated cells. Data from two replicate

wells with error bars showing variability between replicates. A representative from > 3 similar experiments. **B**. Split

violin plots with quartiles showing mean nuclear fluorescence of cells treated with different concentrations of

CBL0137 for 60 min. Numbers above violins are p-values of the t-test comparing treated cells to the untreated control

stained either without RNAse A (blue numbers) or with RNAse A (orange numbers). P-values were rounded to 4

decimal places. ns - p-value >0.05. **C**. Split violin plots with quartiles showing mean nuclear fluorescence per

cell fixed with either 4% PFA or 100% methanol. Numbers on the plot are p-values of the t-test comparing treated

cells to untreated control fixed either with PFA (blue numbers) or methanol (orange numbers). **D, E**. Fold change in

average nuclear fluorescence per well of cells treated with CBL0137 relative to untreated cells and stained in the

presence of absence of RNAse A. **D**. Cells were stained with DAPI. **E**. Cells were stained with PI. Data from two

replicate wells with error bars showing variability between wells.

Figure 3. Effect of epigenetic drugs on chromatin accessibility measured with PI. NDF and HT 1080 cells were

treated with CBL0137 (500 nM) , trichostatin A (TSA, 500 nM), panobinostat (PNB, 20 nM), valproic acid (VA, 0.5

mM), SAHA (1000 nM), or JQ1 (1000 nM) for 24 hours before fixation with PFA and staining with PI in the presence

of RNAse A. **A, B**. Number of nuclei per well of NDF (**A**) or HT1080 (**B**) cells, untreated or treated with the drugs.

Bars are mean value of 4 replicate wells, error bars are SDV. Number above the bar is p-value of t-test comparing

each drug with the control untreated cells, rounded to 2 decimal points. **C, D**. Violin plots with quartiles of distribution

of PI nuclear fluorescence of cells in all replicate wells of NDF (**C**) or HT1080 (**D**) cells treated with epigenetic drugs.

Number above violin is p-value of t-test comparing each drug with the control untreated cells, rounded to 4 decimal

points. Pink and blue transparent squares show position of quartiles 0.5 and 0.75 (pink), and 0.5 and 0.25 (blue) in

control untreated samples. **E, F**. Microscopic images of PI-stained wells from the plate of cells treated with epigenetic

drugs. The same exposition time was used for all images of NDF (**E**) and HT1080 (**F**) cells.

Figure 4. Dependence of chromatin accessibility of cell proliferation. **A-D**. HT1080 cells were incubated in the

presence of EdU for 1 hour. After that cells were fixed and stained for EdU and DAPI. **A**. Violin plots with quartiles

of nuclear fluorescence of of EdU positive and negative cells. Bleu squares correspond to the positions of quartiles

(0.25, 0.5 and 0.75) in EdU negative cells. **B**. Histogram of distribution of total nuclear fluorescence (Integral) in EdU

negative and positive cells. Blue squares show values of nuclear fluorescence used to select cells with DNA content

close to G1 or G2/M cells. **C, D**. Violin plots with quartiles showing mean nuclear fluorescence of EdU positive and

negative cells with DNA content close to G1 (**C**) or G2 /M (**D**). ns – p-value > 0.05. **E, F**. Comparison of mean nuclear

fluorescence of NDF (**E**) or HT1080 (**F**) cells growing in normal conditions or arrested with dense plating and medium

with 1% FBS for 48 hours. Before staining cells were treated with the indicated concentrations of CBL0137 for 60

minutes. Numbers are p-values of t-test comparing growing and arrested cells at each concentration of CBL0137.

Figure 5. Comparison of chromatin accessibility between normal, transformed and tumor cells. **A, B**. Split violin

plots of mean nuclear fluorescence of NDF and HT1080 cells treated with different concentrations of CBL0137 for

60 minutes and stained with DAPI (**A**) or PI (**B**). Numbers are p-value of t-test comparing NDF and HT1080 cells. **C,**

D. Violin plots with quartiles showing mean nuclear fluorescence (DAPI) of two biological replicates of transformed

NDF (**C**) or MCF10A (**D**) cells (GR1, GR2) and control non transformed cells (EV). Numbers are p-values of t-test

comparing each of transformed variants (GR) with control non-transformed EV cells. **E, F**. Comparison of nuclear

fluorescence (PI) of mouse normal, transformed and tumor cells. **E**. Violin plots with quartiles of mean nuclear

fluorescence. Numbers above the lines show p-values of ANOVA test comparing all groups. **F**. Distribution of total

nuclear fluorescence. Numbers show X-values of corresponding peaks on histograms. Dotted lines correspond to the

positions of G1 peaks of normal MEF (red), transformed (blue) and tumor (green) cells.

698