Widespread occurrence of benzimidazole resistance single nucleotide

polymorphisms in the canine hookworm, *Ancylostoma caninum,* in Australia

Swaid Abdullah[a,1,*], Thomas Stocker[b,1], Hyungsuk Kang[a], Ian Scott[c], Douglas

Hayward[d], Susan Jaensch[d], Michael P. Ward[b], Malcolm K. Jones[a], Andrew C.

Kotze[a], Jan Šlapeta[b,e,*]

[a] *The University of Queensland, School of Veterinary Science, Gatton 4343, QLD,*

*Australia*

[b] *Sydney School of Veterinary Science, Faculty of Science, University of Sydney,*

*Sydney, New South Wales, Australia*

[c] *School of Veterinary Science, Massey University, Palmerston North, New Zealand*

[d] *Vetnostics NSW - North Ryde Laboratory, Macquarie Park, New South Wales,*

*Australia*

[e] *Sydney Institute for Infectious Diseases, The University of Sydney, New South*

*Wales, Australia*

*Corresponding authors.

E-mail addresses: jan.slapeta@sydney.edu.au (Jan Šlapeta);

swaid.abdullah@uq.edu.au (Swaid Abdullah)

[1] These authors contributed equally.

**Abstract**

Canine hookworm (*Ancylostoma caninum*), a gastrointestinal nematode of domestic dogs, principally infects the small intestine of dogs and has the potential to cause zoonotic disease. In greyhounds and pet dogs in the USA, *A. caninum* has been shown to be resistant to multiple anthelmintics. We conducted a molecular survey of benzimidazole resistance in *A. caninum* from dogs at veterinary diagnostic centers in Australia and New Zealand. First, we implemented an internal transcribed spacer (ITS)-2 rDNA deep amplicon metabarcoding sequencing approach to ascertain the species of hookworms infecting dogs in the region. Then, we evaluated the frequency of the canonical F167Y and Q134H isotype-1 β-tubulin mutations, which confer benzimidazole resistance, using the same sequencing approach. The most detected hookworm species in diagnostic samples was *A. caninum* (90%; 83/92); the related Northern hookworm (*Uncinaria stenocephala*) was identified in 11% (10/92) of the diagnostic samples. There was a single sample with coinfection by *A. caninum* and *U. stenocephala*. Both isotype-1 β-tubulin mutations were present in *A. caninum*, 49% and 67% for Q134H and F167Y, respectively. Mutation F167Y in the isotype-1 β-tubulin mutation was recorded in *U. stenocephala* for the first known time. Canonical benzimidazole resistance codons 198 and 200 mutations were absent. Egg hatch assays performed on a subset of the *A. caninum* samples showed significant correlation between 50% inhibitory concentration ($IC_{50}$) to thiabendazole and F167Y, with an increased $IC_{50}$ for samples with >75% F167Y mutation. We detected 14% of dogs with >75% F167Y mutation in *A. caninum*. Given that these samples were collected from dogs across various regions of Australia, the present study suggests that benzimidazole resistance in *A. caninum* is widespread. Therefore, to mitigate the risk of resistance selection and further spread, adoption of

49    a risk assessment-based approach to limit unnecessary anthelmintic use should be a

50    key consideration for future parasite control.

52    *Keywords*:  *Ancylostoma caninum,* Multiple anthelmintic drug resistant (MADR),

53    Greyhounds, Anthelmintics, Canine health, Deep sequencing

## 1. Introduction

For many years the prevailing dogma regarding the use of anthelmintics in small animals such as dogs and cats, as well as in humans, had been that the development of anthelmintic resistance would be slow to occur, or may never occur, or be limited in spread (Schulz et al., 2018; Marsh and Lakritz, 2023). The theory behind this dogma was that, at the time of deworming, the refugia - the population of susceptible nematodes infecting the host - remains substantial (von Samson-Himmelstjerna et al., 2021). By contrast, in ruminants, routine and indiscriminate mass treatment with anthelminthics has led to widespread drug resistance because the refugia has not been retained (Vercruysse et al., 2012; von Samson-Himmelstjerna et al., 2021). However, despite these differences between ruminants and small animals, a number of reports over recent years have indicated that resistance in canine hookworms is indeed a widespread problem in North America (Jimenez Castro et al., 2019; Kitchen et al., 2019; Venkatesan et al., 2023).

Currently, the effectiveness of the benzimidazole anthelmintic class in ruminants varies, with many cases showing reduced efficacy (Ramunke et al., 2016; Jaeger and Carvalho-Costa, 2017; Ploeger and Everts, 2018). This is due to well-documented single nucleotide polymorphisms (SNPs) that result in nematodes carrying alleles resistant to these anthelmintics (Kwa et al., 1993; von Samson-Himmelstjerna et al., 2007). However, this may not apply universally across all countries, production systems, species of nematodes and species of livestock where benzimidazole remains efficacious (Mohammedsalih et al., 2021). In humans, hookworms are commonly treated using benzimidazoles, specifically albendazole and mebendazole (WHO, 2017). Between 1995 and 2015, the efficacy of both albendazole and mebendazole for the treatment of human hookworm infection has

decreased by as much as 15%, but resistance as such has not been documented

(Schulz et al., 2018).

In dogs, anthelminthic resistance was first documented 20 years ago in

hookworms to the pyrimidine derivative anthelmintic pyrantel in Queensland,

Australia (Kopp et al., 2007), and currently this anthelmintic is considered ineffective

for treating hookworms in dogs in Australia (Dale et al., 2024). However, resistance

to benzimidazoles and other anthelmintic classes for dogs in Australia has not been

reported. The first report of resistance to benzimidazoles in canine hookworms was

from the USA in 2019 (Kitchen et al., 2019). Molecular, phenotypic, and experimental

infection evidence demonstrating benzimidazole resistance in *A. caninum* was

published independently by two groups in 2019 (Jimenez Castro et al., 2019; Kitchen

et al., 2019). These reports first emerged in North America and were followed by an

extensive surveillance for the canonical F167Y (T$\underline{T}$C>T$\underline{A}$C) isotype-1 β-tubulin

mutation conferring benzimidazole resistance (Leutenegger et al., 2023, 2024;

Venkatesan et al., 2023). A new mutation in isotype-1 β-tubulin at Q134H

(CA$\underline{A}$>CA$\underline{T}$) was experimentally established as another mutation responsible for

resistance to benzimidazoles in *A. caninum* and shown to be widespread across

North America (Venkatesan et al., 2023).

It is speculated that benzimidazole resistance in *A. caninum* emerged in

racing greyhound dogs due to indiscriminate and frequent deworming in racing

kennels (Jimenez Castro et al., 2019; Kitchen et al., 2019; Leutenegger et al., 2024).

Evidence of multiple anthelmintic drug resistance (MADR) in *A. caninum* led the

American Association of Veterinary Parasitologists to establish a hookworm task

force to address this issue in 2021 (von Samson-Himmelstjerna et al., 2021; Marsh

and Lakritz, 2023). The screening of thousands of dogs positive for hookworms in

the USA and Canada demonstrated that the canonical F167Y (T<u>T</u>C>T<u>A</u>C) isotype-1

β-tubulin mutation was present in 11.3% (715/6,329) of faecal *A. caninum*-positive

samples (Leutenegger et al., 2024). In a smaller study of samples from the USA,

49.7% (156/314) of the individual samples tested positive for the F167Y (T<u>T</u>C>T<u>A</u>C)

mutation (Venkatesan et al., 2023). Reanalysis of next-generation sequence data

from Venkatesan et al. (2023) for the presence of non-*A. caninum* sequences

revealed the presence of other hookworms (*Ancylostoma braziliense*, *Ancylostoma*

*tubaeforme* and *Uncinaria stenocephala*), but only the benzimidazole-susceptible

residues of isotype-1β-tubulin were found in these species (Stocker et al., 2024).

The aim of this study was to investigate the extent of benzimidazole

resistance SNPs in canine hookworm (*A. caninum*) outside North America,

specifically in Australia and New Zealand. We utilised opportunistic sampling from

four different sources in Australia and New Zealand. All hookworm-positive samples

were screened using internal transcribed spacer (ITS)-2 rDNA PCR and deep

amplicon sequencing to determine the hookworm species present. To investigate

resistance to benzimidazoles, we used deep amplicon sequencing to evaluate the

nucleotide sequence of the isotype-1 β-tubulin mutations that are known to be

associated with resistance to this drug class, specifically the regions that encode

amino acids: glutamine 134 (Q134), phenylalanine 167 (F167), glutamic acid 198

(E198) and phenylalanine 200 (F200). We then undertook egg hatch assays (EHAs)

on a subset of the samples to correlate the molecular signatures, specifically

frequency of F167, with the benzimidazole-resistant phenotype of these field

isolates.

**2. Materials and methods**

*2.1. Ethics statement*

The use of residual clinical samples was in accordance with the University of Sydney (Australia) Animal Ethics Committee Protocol 2024/2508. Dog faecal samples for larval cultures were obtained in accordance with the University of Queensland (Australia) Animal Ethics Committee Protocol 2021/AE001169 and the faecal egg count reduction test on one dog was conducted in accordance with the University of Queensland Animal Ethics Committee Protocol SVS/111/20.

*2.2. Faecal samples with hookworm infection*

Faecal samples ($n$=64) were provided by registered veterinary practitioners in accordance with the Veterinary Practice and Animal Welfare Acts, Australia for the purpose of diagnostics used in decision making by the veterinary practitioner. Samples were submitted for parasitological diagnostics ($n$=50) to Vetnostics (Laverty Pathology, North Ryde Laboratory, Sydney, New South Wales, Australia) and ($n$=14) Veterinary Pathology Diagnostics Services (VPDS, The University of Sydney, New South Wales, Australia) for centrifugal qualitative flotation using saturated salt (NaCl, specific gravity = 1.21); refrigerated residual faecal samples that previously returned a positive result for hookworms at the clinical diagnostic laboratory were received for molecular species characterisation. These samples came from 63 dogs and one cat. The group of dogs included 45 greyhounds, four foxhounds, one boxer, one Labrador retriever, one Afghan, one German Shepherd, one King Charles Cavalier, one Irish wolfhound cross, and eight dogs of unknown breed. Material used for

hookworm species identification and benzimidazole susceptibility, and subsequent

results, were de-identified.

Material (*n*=22) from Queensland, Australia was obtained from dogs from

adoption and shelter facilities that receive rescued or surrendered animals from

greyhound rescue organisations and pounds across south-eastern Queensland.

These sample were from 16 greyhounds, one Siberian husky, three cross-breed

dogs and two dogs of an unknown breed. Dogs within these facilities are kept in

individual pens, which are cleaned and disinfected daily. Fresh faecal samples were

collected from each dog after natural defecation and screened for hookworm

infection using faecal flotation as described in section 2.3.

Samples (*n*=20) from Massey University, Palmerston North, New Zealand that

contained hookworm eggs were processed to isolate the eggs, which were stored in

80% ethanol for transfer to VPDS.

Details of samples used in this study, with available metadata, are available at

LabArchives (https://dx.doi.org/10.25833/hmvy-jn79).

*2.3. Faecal egg counts (FECs), hookworm egg recovery and DNA isolation*

Faecal egg counts (FECs) were performed with saturated salt (NaCl, specific

gravity = 1.21) solution using a Whitlock Universal McMaster four chamber worm egg

counting slide in two different laboratories.

At the University of Sydney, 3 g of faecal samples were homogenised with 60

ml of saturated salt solution and filtered via a coarse strainer. We counted two

compartments each of 0.5 ml volume, resulting in minimum detection of 20 egg per

gram of faeces (EPG) (absence of eggs indicates <20 EPG in the sample). Only strongyle-like (=hookworm) eggs were counted. The remainder of the filtered homogenate was poured into a 50 ml Falcon tube and centrifuged at 800 $g$ for 10 min for centrifugal floatation. The uppermost supernatant (~10 ml) was collected with a plastic Pasteur pipette into a new clean 50 ml Falcon tube, topped with tap water and centrifuged at 800 $g$ for 10 min. The supernatant was decanted and the pellet that contained the concentrated eggs was retained. Then ~1 ml of tap water was added with a clean plastic Pasteur pipette to disturb the pellet by repeated aspiration until a homogeneous solution formed. The homogenate was transferred to a clean 1.5ml Eppendorf tube and spun at 8,000 $g$ for 10 min. The supernatant was completely removed and the Eppendorf tube with the egg pellet was stored at -20 °C until further use.

At the University of Queensland, 3 g of faecal samples were homogenised with 42 ml of saturated salt solution, filtered through a tea strainer and the filtrate was immediately pipetted into the counting chamber (four compartments each of 0.5 ml volume). Counting all compartments (2 ml) delivered minimum detection of seven EPG (absence of eggs indicates <7 EPG in the sample). Eggs were purified for use in the EHA (see section 2.6). The remaining eggs were suspended in distilled water in 1.5 ml microtubes, incubated for 72 h at 28°C and allowed to hatch. The hatched larvae were stored at -20°C until further use.

All total genomic DNA was isolated at the University of Sydney. For partially cleaned hookworm eggs, an Isolate II Fecal DNA Kit (Meridian Bioscience, Australia) was used according to the manufacturer's instructions with the following homogenisation step. For purified eggs, 750 µL of lysis buffer were added to the Eppendorf tube with the pellet and homogenised by repeated aspiration with a

pipette, then all the contents were transferred into a bead-beating tube, and the eggs

disrupted and homogenised using a high-speed benchtop homogeniser, FastPrep-24

(MP Biomedicals, Australia), for 40 s at 6.0 m/s. Every 11[th] sample was a blank

sample that only contained the lysis buffer. Egg samples suspended in ethanol were

first dried before processing as described above. For hatched larvae, a Monarch

Genomic DNA Purification Kit (New England Biolab, Australia) was used according

to the manufacturer's instructions. Purified genomic DNA samples and blanks were

isolated into 100 μL of elution buffer (10 mM Tris-Cl, pH = 8.5) and stored at –20 °C

until further use.

*2.4. Amplicon metabarcoding and Illumina deep sequencing*

All isolated DNA samples were first screened for the presence of nematode

DNA using nematode ITS-2 rDNA quantitative PCR and samples with a cycle

threshold (Ct)<35 were considered suitable DNA material for follow-up processes.

Total DNA was subjected to three independent real-time PCR amplifications

at the University of Sydney as previously described (Stocker et al., 2023). ITS-2

rDNA (ITS-2) was amplified by the NC-1 and NC-2 primers designed by Gasser et al.

(1993), and two isotype-1 β-tubulin regions were amplified using assays BZ167 and

BZ200 developed by Jimenez Castro et al. (2019), which include key diagnostic

SNPs conferring benzimidazole resistance. All primers were adapted for Illumina

metabarcoding using Next Generation Sequencing (NGS). The ITS-2 target was

amplified using 32-36 amplification cycles, while 40 amplification cycles were used

for isotype-1 β-tubulin (BZ167, BZ200). All reactions utilized SYBR-chemistry using

SensiFAST™SYBR® No-ROX mix (Meridian Bioscience, Australia). Each real-time

PCR mix (25 μL) included 2 μL of template DNA (neat) and primers at final

concentrations of 400 nM. All PCRs were prepared using a Myra robotic liquid

handling system (Bio Molecular Systems, Australia). Reactions were run on a CFX96

Real-Time PCR Detection System with corresponding CFX Manager software

(BioRad, Australia). The protocol involved an initial denaturation at 95 °C for 3 min,

followed by 32 or 40 cycles of 95 °C for 5 s, 60 °C for 15 s, 72 °C for 15 s, and a final

melt curve analysis. Each run included a no-template control (ddH$_2$O) to monitor for

potential contamination. Amplicons in 96 well plate format were submitted for

indexing, purification and amplicon sequencing using Illumina NGS at the Ramaciotti

Centre for Genomics, University of New South Wales, Sydney, Australia and

sequenced on MiSeq using MiSeq Reagent Kits v2 (250PE, Illumina). De-

multiplexed FastQ files were generated via BaseSpace (Illumina).

*2.5. Analysis of the Illumina sequenced amplicons*

All FastQ files were processed through R package 'dada2' v1.26.0 (Callahan

et al., 2016) in R v4.2.2 (R Core Team, 2022), to obtain amplicon sequence variants

(ASV) and their abundance per sequenced amplicon. The dada2 pipeline at the

University of Sydney includes filtering, error estimation and denoising, before paired-

end reads were merged to reconstruct the full ASV before removal of chimeras. Prior

to processing through the dada2 pipeline, primers were removed from all forward

and reverse FastQ reads using 'cutadapt' version 4.0 (Martin, 2011).

The assignment of species based on ITS-2 ASVs was carried out in CLC

Main Workbench v22.0 (Qiagen Australia) as previously described (Stocker et al.,

2024). All ASVs were imported into CLC Main Workbench and top blastn hits against

the 'nr' database at NCBI [https://blast.ncbi.nlm.nih.gov/Blast.cgi] were identified.

ITS-2 sequences were discarded for any ASVs where the top blastn hit was not

hookworm (Ancylostomatidae). For species assignment we constructed alignment of

AVSs with an in-house curated reference ITS-2 hookworm sequence library

consisting of publicly available sequences (GenBank: **KP844736**; **MT345056**;

**JQ812692**; **LC036567**; **JX220891**; **EU344797**; **AB793527**). The resulting multiple

sequence alignment was subject to phylogenetic reconstruction, where species

identity was based on manual assignment of ASV to a species based on its position

on the phylogenetic tree relative to the reference ITS-2 sequences.

For the β-tubulin sequence data (BZ167, BZ200 assay), the assignment of

species was performed with the aid of CLC Main Workbench v22.0 (Qiagen

Australia) as previously described in Stocker et al. (2024). Briefly, all ASVs were

imported into CLC Main Workbench and top blastn hits against the 'nr' database at

NCBI were identified. All ASVs where the top blastn hit was a hookworm

(Ancylostomatidae) β-tubulin were retained for further analysis. The reference

hookworm alignments of BZ167 and BZ200 amplicons according to Stocker et al.

(2024) were used to unambiguously identify ASV identity and only those ASVs that

were confirmed based on phylogenetic analysis were included in further analyses.

The sequence alignment included β-tubulin reference sequences from *A. caninum*

(**DQ459314**) (Schwenkenbecher et al., 2007), *Ancylostoma duodenale* (**EF392850**),

*U. stenocephala* (Stocker et al., 2023), and *Necator americanus* (**EU182348**), which

enabled us to identify the exon/intron boundary and residues that code for

susceptibility to benzimidazoles, amino acids amino acids Q134, F167, E198 and

F200 (Venkatesan et al., 2023). Our reference alignment included sequence data for

isotype-1 β-tubulin from *A. caninum*, *Ancylostoma ceylanicum*, *A. duodenale* and *N.*

*americanus* at 'WormBase ParaSite' (Howe et al., 2017; Bolt et al., 2018; Harris et
al., 2020).

Following the dada2 pipeline, species assignment raw read counts were
converted into relative abundance (percentages). Any species read counts with <25
reads were disregarded as spurious. Proportions of SNPs for amino acids Q134,
F167, E198 and F200 were calculated per species for each of the amplicons of
isotype-1 β-tubulin (BZ167, BZ200).

The number of samples, and number of *A. caninum* samples with Q134H
(CA<u>A</u>>CA<u>T</u>) and F167Y (T<u>T</u>C>T<u>A</u>C) separately, were summed per postcode in
New South Wales. Proportional symbol maps were created by joining this data to
postcodes of Australia shapefile (Geocentric Datum of Australia, GDA 1994) in
ArcGIS® Pro. Spearman rank correlations ($r_{SP}$) were calculated between EPG per
sample and both frequency of Q134H (CA<u>A</u>>CA<u>T</u>) and F167Y (T<u>T</u>C>T<u>A</u>C). In
addition, the EPG between resistant versus not resistant allele status for both
mutations was assessed using a Kruskal-Wallis non-parametric analysis of
variance. Spearman rank correlations were also calculated between EPG,
frequency of Q134H (CA<u>A</u>>CA<u>T</u>), frequency of F167Y (T<u>T</u>C>T<u>A</u>C) and age
(years). Finally, the association between breed (greyhound versus other) and
EPG, resistance allele frequency for Q134H (CA<u>A</u>>CA<u>T</u>) and F167Y (T<u>T</u>C>T<u>A</u>C),
was assessed using a Kruskal-Wallis non-parametric ANOVA; DFn: degrees of
freedom for the numerator of the F ratio, DFd: degrees of freedom for the
denominator of the F ratio. The type I error for all statistical tests was set at 0.05
(IBM SPSS Statistics v 28).

*2.6. In vitro EHA*

Eggs were harvested from fresh faecal samples from hookworm-positive dogs

for use in in vitro EHAs. Saturated sugar solution (specific gravity = 1.30), three

times the weight of individual faecal samples, was added to the faecal samples in a

shaker bottle. The bottle was manually agitated to form a slurry, which was then

filtered through a 200 μm sieve into a beaker to remove large debris. All of the

filtered solution was then transferred into 50 ml Falcon tubes, filling each tube up to

40 ml, and then 10 ml of distilled water were slowly and carefully poured along the

edge of the tube to avoid mixing with the faecal solution. Each tube was then screw

capped and centrifuged at 400 *g* for 10 min. Following centrifugation, the 'fuzzy' layer

(water-sucrose interface) of each tube was removed with a pipette and pooled into a

200 mL beaker. The solution was passed through a 75 μm sieve, retaining the

filtrate, which was further passed through a 20 μm sieve, which retained the

hookworm eggs. This sieve was then retrograde washed with 20 mL of distilled water

into a glass Petri dish. The recovered eggs were carefully poured into a two 15 mL

centrifuge tubes, leaving the debris in the Petri dish. The eggs were concentrated

into 2 mL of water by centrifugation for 5 min at 400 *g* and the supernatant

discarded. The egg concentration was estimated by thoroughly mixing the 2 mL of

egg solution and counting the eggs in three 20 μl aliquots. The egg numbers were

adjusted to approximately two eggs per microlitre either by dilution with distilled

water or by concentration using centrifugation as described above.

The EHA was performed using 96-well plates containing thiabendazole

embedded in agar, as described previously for assays with human hookworms

(Kotze et al., 2009). A stock solution of thiabendazole (5 mg/mL, Sigma-Aldrich

Australia) was prepared in dimethyl sulfoxide (DMSO) and serially diluted two-fold in

the same solvent. Aliquots (2 μL) from a series of dilutions were added to 96-well

microtiter plates, such that each row of the plate comprised a double dilution, ranging

from 0.003 μg/mL up to 1.359 μg/mL or 0.011 μg/mL to 5.437 μg/mL. The first two

wells of each row were used as control wells (received 2 μL of DMSO only). Molten

agar (200 μL at 2% w/v) (Davis Gelatine, Australia; powdered agar Grade J) was

dispensed into each well of the plate and allowed to set. Plates were placed into

plastic press-seal bags stored at 4°C.

Prior to establishing the assays, the plates were allowed to warm in an

incubator set to 28°C. Thirty microlitres of concentrated hookworm egg solution (~2

eggs/μL) were pipetted into each well. Plates were incubated for 48 h at 28°C,

whereupon the assay was terminated by addition of Lugol's iodine solution. The

numbers of unhatched eggs and larvae in each well were counted with the aid of a

stereo microscope (Stemi 508, Carl Zeiss, Australia). All assays were performed in

triplicate for the sample isolated from each dog sample.

The number of larvae in each well was expressed as a percentage of the total

numbers of larvae and unhatched eggs to give the hatch rate in each well, and then

this was expressed relative to the mean hatch rate in the control wells. The data

were then analysed by non-linear regression using GraphPad Prism 10.3.0 software

(GraphPad Software Inc. USA). This analysis yielded 50% inhibitory concentration

($IC_{50}$) values representing the concentration of thiabendazole that resulted in an egg

hatch of 50% compared with control assays. The $IC_{50}$ values for the dogs examined

in this study were compared with those reported previously for drug-susceptible *A.*

*caninum* isolates: 0.046 and 0.048 μg/mL (Jimenez Castro et al., 2019) and 0.045

μg/mL (Diawara et al., 2013). An Australian fenbendazole-susceptible isolate of *A.*

*caninum* ('Dougie', from February 2021) *s*erved as a local positive control and

demonstrated faecal egg reduction of 100% 10 days post administration from an
initial 525 EPG.

The left-over eggs were suspended in distilled water in 1.5 ml microtubes,
incubated for 72 h at 28°C and allowed to hatch. The hatched larvae were stored at -
20°C and subjected to DNA isolation and follow up processes described earlier
(section 2.3).

*2.7. In vivo efficacy measurements*

One dog (sample D6) was treated by the shelter facilities and the FEC
monitored throughout treatment. The dog was treated according to its weight and the
product label on day 0 with Drontal Allwormer (Vetoquinol, Australia; actives:
febantel, praziquantel and pyrantel embonate) and on day 21 with Credelio Plus
(Elanco, Australia: milbemycin oxime and lotilaner). In animals, febantel is a prodrug
that is actively hydrolyzed to the active metabolite fenbendazole. Egg counts were
conducted within the period 10-21 days after Drontal Allwormer treatment to
calculate the faecal egg reduction percentage (>95% reduction is considered an
effective product).  Only a single sample was obtained at day 8 after Credelio Plus
because the dog was then rehomed.

*2.8. Data availability*

The EHA raw data, FEC reduction data and metadata for all samples are
available at LabArchives (https://dx.doi.org/10.25833/hmvy-jn79).  Raw   FastQ
sequence data was deposited at SRA NCBI BioProject: **PRJNA1149426**.

## 3. Results

*3.1. Dominance of A. caninum across south-eastern Australia using metabarcoded ITS-2 rDNA deep sequencing in dogs*

We used 106 faecal samples that were considered hookworm-positive. There were 20 dog samples from New Zealand. For 61 of the 106 samples the worm FEC ranged from 20 to 4,920 (average 526) EPG. The majority of the Australian samples came from a commercial diagnostic laboratory (*n*=50) and were collected over a period of 13 months between March 2023 and March 2024.

There were 86 samples from south-eastern Australia, 85 samples from dogs and one from a cat (TS42). Amplifiable DNA was obtained from 98% (84/86), yielding a Ct<35 (11.8 to 31.6) at ITS-2 in an rDNA assay targeting nematode DNA. All 84 samples yielded metabarcoded and deep sequenced ITS2 rDNA sequence data for nematode species identification (*n*=84; average 21,323 reads, range 794 to 32,899 reads per sample) (Fig. 1A). Two samples yielded a low (<2,000) number of ITS-2 rDNA sequence reads, 794 (TS100) and 1,537 (TS95); removing those the remaining samples yielded >2,000 reads per sample (*n*=82; average 21,815, range 2,736 to 32,899). The cat hookworm sample contained sequences that matched *Ancylostoma tubaeforme* (TS42). The remaining dog samples (*n*=81) contained sequences of *A. caninum* (*n*=77, 95%) and *U. stenocephala* (*n*=4, 5%). One sample (TS35) had both *A. caninum* (96%) and *U. stenocephala* (4%) ITS-2 rDNA sequences. The additional two samples with <2,000 ITS-2 rDNA sequence reads (TS95, TS100) included only *A. caninum* sequences.

*3.2. Widespread and high frequency of A. caninum canonical F167Y (TTC>TAC)*

*isotype-1 β-tubulin benzimidazole resistance mutation in Australia*

For 81 Australian samples, the isotype-1 β-tubulin BZ167 amplicons included

*A. caninum* isotype-1 β-tubulin with an average sequencing read count between 81

to 31,265 (Fig. 1B). Excluding six samples (TS6, TS7, TS39, TS55, TS63, TS90;

sequence reads 81 to 1,699) with <2,000 sequences, the average was 11,092

sequences (*n*=75; 2,213 to 31,265). There were 72 samples with >2,000 isotype-1 β-

tubulin BZ167 amplicon sequences belonging to *A. caninum* (average sequencing

reads 11,271) (Fig. 1B).

Among the samples, 49% (35/72) contained the Q134H (CAA>CAT) resistant

allele, with an average frequency of 10% (range 1% to 45%; 95% confidence interval

(95% CI): 7-13%). Similarly, 67% (48/72) of the samples contained the F167Y

(TTC>TAC) resistant allele, with an average frequency of 41% (range 1% to 100%;

95% CI: 33-51%) (Fig. 1C). There were 28% (20/72), 25% (18/72) and 14% (10/72)

of samples with >40%, >50% and >75% F167Y (TT̲C>TA̲C) frequency, respectively

(Fig. 1C). Moreover, 36% (26/72) of samples had both Q134H (CAA̲>CAT̲) and

F167Y (TT̲C>TA̲C) benzimidazole-resistant alleles. The six samples with <2,000

sequences for *A. caninum* had only the susceptible (coding for Q134, F167) alleles,

except TS90 with 100% of F167Y (TT̲C>TA̲C) mutation. Sequence analysis of

isotype-1 β-tubulin sequence (amino acids E198 and F200; BZ200 amplicons)

revealed no evidence of benzimidazole resistance signatures.

Besides *A. caninum* isotype-1 β-tubulin sequences, there were four *U.*

*stenocephala* and one (1/4) of the samples TS24 included F167Y (TT̲C>TA̲C) in the

isotype-1 β-tubulin mutation in 51% of the sequence reads. The remaining isotype-1

β-tubulin of *U. stenocephala* matched benzimidazole susceptible sequence residues,

including across BZ200 amplicon (E198 and F200). The cat TS42 sample with *A.*

*tubaeformae* and its isotype-1 β-tubulin included only the benzimidazole-susceptible

sequences (amino acids Q134, F167, E198 and F200).

For the New Zealand samples only 50% (10/20) of DNA yielded $C_t<35$ (20.2

to 33.6); nine samples yielded ITS-2 rDNA sequence data to consider species

identification (TS76-TS84; average 18,488; 1,118 to 27,624 sequencing reads) (Fig.

1A). Four of the nine samples contained sequence data that belonged to non-

hookworm sequence including ITS-2 rDNA sequencing matched to parasites of

sheep and rabbits; three DNA samples contained only ASVs that matched spurious

parasites (='pseudoparasite') DNA sequences. Subtracting the 'pseudoparasite'

sequences to retain hookworm sequence data revealed the presence of uniquely *U.*

*stenocephala* sequences, five samples with 8,027 to 27,624 sequencing reads

belonging to hookworms and one sample (TS76) with only 118 sequences.

Sequence analysis of the isotype-1 β-tubulin (amino acids Q134, F167, E198 and

F200) revealed no evidence of benzimidazole resistance signatures in New Zealand

samples.

*3.3. High frequency of Q134H (CAA>CAT) and F167Y (TTC>TAC) mutations in A.*

*caninum in greyhounds*

We then focused on 55 dog samples from two diagnostic laboratories

(Vetnostics, VPDS) for which we had *A. caninum* Q134H (CA<u>A</u>>CA<u>T</u>) and F167Y

(T<u>T</u>C>T<u>A</u>C) mutation frequency in the isotype-1 β-tubulin. There were 53 dogs with a

defined location to a postcode from New South Wales, Australia (one dog from New

South Wales did not have a postcode and the other dog was from Queensland).

Dogs came from 12 distinct New South Wales postcodes (Fig. 2). The *A. caninum*

faecal egg count across all 55 dogs was 534 EPG (minimum 20, maximum 4,920)

Greyhounds represented 80% (44/55) of the samples (42 from New South

Wales, one from Queensland, one from an unknown state). The F167Y (T<u>T</u>C>T<u>A</u>C)

allele was present in 30 (68%, 30/44) of them with 37% average frequency (mininum

2%, maximum 100%). The Q134H (CA<u>A</u>>CA<u>T</u>) allele was present in 28 (64%,28/44)

of them with 10% average frequency (minimum 1%, maximum 45%).

There was no significant correlation between Q134H (CA<u>A</u>>CA<u>T</u>) or F167Y

(T<u>T</u>C>T<u>A</u>C) and EPG (correlation coefficient=-0.169, *P*=0.217; correlation

coefficient=0.03, *P*=0.82). Considering the association of presence or absence of

either Q134H (CA<u>A</u>>CA<u>T</u>) or F167Y (T<u>T</u>C>T<u>A</u>C) and EPG, there was no significant

association for either SNP (Kruskal-Walis Test: H=0.126, degrees of freedom (df)=1,

*P*=0.72; H=0.56, df=1, P=0.45). We then considered the least stringent threshold of

>40% for F167Y (T<u>T</u>C>T<u>A</u>C) mutation and recalculated the association, but the test

remained not significant (Kruskal-Walis Test: H=0.02, df=1, *P*=0.89). There was only

a single sample with >40% of the Q134H (CA<u>A</u>>CA<u>T</u>) mutation, therefore preventing

calculation of these test statistics. Considering the ages of the dogs (for those

available, *n*=41) there was no significant correlation between age and EPG, Q134H

(CA<u>A</u>>CA<u>T</u>) or F167Y (T<u>T</u>C>T<u>A</u>C) (Spearman correlation, *P*>0.05).

The median resistance allele frequency in *A. caninum* samples from

greyhounds for both the Q134H (CA<u>A</u>>CA<u>T</u>) and F167Y (T<u>T</u>C>T<u>A</u>C) mutation (0.035

and 0.090, respectively) was significantly higher than that for other dog breeds

(median 0.000; Kruskal Wallis Test, $P \leq 0.01$) (Fig 2). There was no significant

difference between EPGs for the two breed categories (Kruskal-Wallis Test, $P$=0.22).

*3.4. Phenotypic evidence of benzimidazole resistance in A. caninum field sample*

*with >75% F167Y (T<u>T</u>C>T<u>A</u>C) mutation*

We tested 13 *A. caninum* samples (D1, D2, D3, D5, D6, D8, D9, D10, D12,

D15, D16, D17, D18/20) with EHAs that had their frequency of Q134H (CA<u>A</u>>CA<u>T</u>)

and F167Y (T<u>T</u>C>T<u>A</u>C) *A. caninum* SNP established (Fig. 1, Table 1). Dose

response curves and $IC_{50}$ formed two groups, those that had high $IC_{50}$>0.15 µg/mL

and those that had low $IC_{50}$<0.15 µg/mL of thiabendazole (Fig. 3A). Three

greyhounds (D1/D2: TS87/88, D3: TS89, D17: TS103) and the crossbreed dog (D6:

TS92, D18/20: TS104/106) were in the high $IC_{50}$ group and they all had >75% of

F167Y (T<u>T</u>C>T<u>A</u>C) *A. caninum* mutation in the isotype-1 β-tubulin (Fig. 1, Table 1).

The $IC_{50}$ of the remaining seven greyhound samples ranged from 0.02 to 0.06 µg/mL

of thiabendazole and the frequency of *A. caninum* F167Y (T<u>T</u>C>T<u>A</u>C) was <40%

(Fig. 1, Table 1). For one dog (represented by two samples, D6 and D18/20, in Fig.

1), we established an in vitro EHA on eggs purified from the dog's faeces on day 1

and day 9, showing an $IC_{50}$ of 0.24 (0.19 to 0.30) µg/mL and 0.17 (95%CI 0.12 to

0.24) µg/mL of thiabendazole, respectively. There was no significant difference

between the response curves for these two samples, demonstrated by a single

response curve fitting both sample responses to thiabendazole ($P$=0.19; F=1.735,

DFn=2, DFd=45). The benzimidazole susceptible control $IC_{50}$ was 0.11 (95%CI 0.10

to 0.12) µg/mL (Fig. 3), and the *A. caninum* DNA was devoid of canonical F167Y

(T<u>T</u>C>T<u>A</u>C) and Q134H (CA<u>A</u>>CA<u>T</u>) isotype-1 β-tubulin mutations which confer

benzimidazole resistance; the BZ167 and BZ200 amplicons coded for amino acids

Q134, F167, E198 and F200. There was significant correlation of the $IC_{50}$ for

thiabendazole in EHAs and F167Y (T$\underline{T}$C>T$\underline{A}$C) mutation in the isotype-1 β-tubulin

(Spearman's r=0.83 (95% CI=0.52 to 0.95, *P* < 0.001) (Fig. 3B).

In vivo evidence based on repeated FECs for one dog (D6 and D18/20 in Fig.

1) suggested decreased (<95%) ability of febantel/pyrantel to reduce egg shedding

post-treatment. This crossbreed dog (D6, D18/20) initially shed 540 EPG of *A.*

*caninum* hookworms on day 1 (day of treatment). The hookworm isolate carried 83%

of F167Y (T$\underline{T}$C>T$\underline{A}$C) mutation in the isotype-1 β-tubulin (D6: TS092; Fig. 1), and

was treated with 'Drontal Allwormer', but the faecal egg reduction only reached

88.8% (60 EPG) and 72.2% (195 EPG) on day 10 and day 14 post-treatment,

respectively. The same dog was re-treated with macrocyclic lactone ('Credelio Plus')

on day 20 (90 EPG) post original treatment, which resulted in 100% *A. caninum* egg

reduction in faeces on day 8 post second treatment (0 EPG).

**4. Discussion**

This survey of dogs from south-eastern Australia infected with hookworms

demonstrated a high frequency of *A. caninum* with canonical mutations F167Y and

Q134H in the isotype-1 β-tubulin that are known to be associated with benzimidazole

resistance. Both F167Y and Q134H mutations have been conclusively confirmed to

lead to benzimidazole resistance using *Caenorhabditis elegans* as a surrogate

(Kitchen et al., 2019; Venkatesan et al., 2023). The extent of both mutations, 67% for

F167Y and 49% for Q134H, in *A. caninum* tested in this study exceeds even the

recent USA report by Venkatesan et al. (2023) that utilised the same methodology

based on NGS amplicon metabarcoding. Our samples from New South Wales

correspond to hookworm-positive samples tested by a commercial diagnostic

laboratory (*n*=49) over a 13-month period. Our sampling effort for New South Wales

represents approximately 1,700 dog samples, assuming a conservative *A. caninum*

prevalence of 2.9% in temperate and subtropical regions of Australia (Massetti et al.,

2022).

Using our samples from New South Wales, we show that the isotype-1 β-tubulin

F167Y mutation is present in canine hookworms across the state and predominantly

in greyhounds that provided most of the samples. In North America, the isotype-1 β-

tubulin F167Y mutation was present in all geographic areas with detected

hookworms, including Canada (Leutenegger et al., 2024). Initially the F167Y

polymorphism was recognised in *A. caninum* infecting greyhounds; currently the

highest isotype-1 β-tubulin F167Y mutation prevalence was shown to be in the

hookworms infecting poodles (29%), followed by Bernese mountain dogs (25%),

cocker spaniels (23%) and greyhounds (22%) in North America (Jimenez Castro et

al., 2019; Kitchen et al., 2019; Leutenegger et al., 2024). Using faecal samples from

17,671,724 individual dogs in North America, greyhounds were found to have a

significantly higher risk of testing positive for hookworms on faecal flotation (odds

ratio = 15.3) (Burton et al., 2024). Additionally, the interval to test negative was 71–

72 days for presumably anthelminthic-treated greyhounds, compared with just 1–2

days for other breeds (Burton et al., 2024). In this study from Australia, in addition to

greyhounds, one crossbreed dog from Queensland was infected with *A. caninum*

and had 83% of the isotype-1 β-tubulin F167Y mutation, together with a reduced in

vitro $IC_{50}$ for thiabendazole.

The EHA tests for in vitro susceptibility to benzimidazoles, thus phenotypes including

*A. caninum* resistance to benzimidazole (Kotze et al., 2005, 2009; Diawara et al.,

2013). Importantly, we demonstrated correlation of the EHA and the isotype-1 β-

tubulin F167Y mutation, and those samples with elevated EHAs ($IC_{50}$) all showed

frequencies of F167Y mutation >75%. Known in vitro resistant *A. cainum* with an $IC_{50}$

of 8.76 μM (=1.76 μg/mL for KGR isolate) and 40.08 μM (=8.07 μg/mL for BCR

isolate) for thiabendazole had 100% and 76% of the F167Y mutation (McKean et al.,

2024). Our resistant *A. caninum* had an $IC_{50}$ from 0.17 to 0.43 μg/mL of

thiabendazole and are closer to the range (0.55 to 0.67 μg/mL) detected for resistant

*A. caninum* by Jimenez Castro et al. (2019). The in vitro resistance was further

confirmed by FEC reduction in one of the Australian dogs in vivo. In a study by

Jimenez Castro et al. (2021), no significant correlation was found between the

F167Y (TTC>TAC) mutation frequency and the $IC_{50}$ for thiabendazole in a small

study involving 15 *A. caninum* samples. However, three samples with an in vitro

resistance phenotype exhibited very low (<6%) F167Y (TTC>TAC) mutation

frequency (Jimenez Castro et al., 2021). This discrepancy was later explained by the

discovery of a new mutation in isotype-1 β-tubulin, Q134H (CAA>CAT), which was

present at a high frequency (≥90%) in all three in vitro resistant *A. caninum* samples

studied by Venkatesan et al. (2023).

The detection of the isotype-1 β-tubulin F167Y mutation in *U. stenocephala*,

the Northern hookworm that commonly infects dogs in cooler climates, is

unexpected, as drug resistance has not previously been reported for *U.*

*stenocephala* (Leutenegger et al., 2023; Venkatesan et al., 2023; Stocker et al.,

2024). In our samples from Australia, we detected *U. stenocephala* F167Y mutation

in one out of four dog samples. More frequently in Australia, *U. stenocephala* is a

parasite of red foxes which likely serves as the reservoir (Ryan, 1976). In a study of

930 foxes from New South Wales, the most prevalent nematodes were *Toxocara*

*canis* (35.2%) and *U. stenocephala* (30.6%), while *A. caninum* was detected in only

70 (7.5%) foxes (Ryan, 1976). Although *U. stenocephala* is often considered

relatively non-pathogenic, it has been acknowledged as a cause of subacute enteritis

and hypoproteinaemia in dogs (Gibbs, 1958; Miller, 1968). The significance of the

F167Y mutation in *U. stenocephala* in conferring benzimidazole resistance in the

presence of a large *refugium* in red foxes will require further confirmation from in

vitro and in vivo drug response studies.

The absence of resistant SNPs that would code for E198 and F200 in the

isotype-1 β-tubulin of *A. caninum* in Australia agrees with reports from the USA

(Venkatesan et al., 2023). The F200Y SNP has been documented once in *A.*

*caninum* dog samples from Brazil, but the in vitro or in vivo phenotype was not

measured (Furtado et al., 2014). The F200Y was detected in human *N. americanus*

from Haiti with frequencies of 20-90%, and in an in vitro EHA assay these hookworm

isolates were less sensitive to thiabendazole ($IC_{50}$ = 0.12 to 0.26 µg/µL) (Diawara et

al., 2013). Molecular dynamic simulations using isotype-1 β-tubulin showed that *A.*

*caninum* and *A. ceylanicum* both shared the same drug binding profile, while *A.*

*duodenale* and *N. americanus* both had different binding profiles with no overlap

except for the E198 interaction that is shared with all nematodes (Jones et al., 2022).

With growing evidence of anthelmintic resistance in *A. caninum*, the question

arises whether the fitness and ability of resistant strains to spread across the dog

population differ from non-resistant strains. Recently, a strain of triple-resistant *A.*

*caninum* (BCR) was shown to exhibit reduced 'larval activation', an initial step in the

transformation of infective non-feeding larvae into migrating and feeding larvae in a

permissive host (McKean et al., 2024). Larval activation in vitro is initiated when

infective L3s (iL3s) of *A. caninum* are incubated with a host-mimicking signal such as

dog serum to resume feeding over a period of 24 h (Hawdon and Schad, 1990;

Hotez et al., 1993). The consequences of the 40-50% reduced initial activation of the

*A. caninum* triple-resistant strain (BCR) reported by McKean et al. (2024) are not yet

known but suggest a possible reduced ability to establish infection. However, we did

not find any association between the SNPs and FEC in the dogs we studied. Our

study does not permit the examination of relationships between FECs and other

variables (e.g., worm fitness) because the infection histories (such as exposure to

larvae and drug treatment history) of the dogs are completely unknown. Our samples

were neither tested for larval activation nor for the presence of pyrantel and

macrocyclic lactone resistance, so reduced 'larval activation' may either be restricted

to the triple-resistant strains, or it may not lead to lower infection rates and changes

in egg output.

The molecular tools employed in the present study enable agile moderate to

high throughput tracking of anthelmintic resistance mutations. This offers advantages

in terms of speed and cost compared with phenotypic in vivo and in vitro

measurements of sensitivity to anthelmintic drugs. We identified the hookworm

species and simultaneously discovered widely distributed isotype-1 β-tubulin SNPs

across eastern Australia. Hookworms, whipworm and roundworms in dogs and cats

are traditionally managed by sustained routine administration of anthelminthics. The

recent discovery of MADR in canine hookworm (*A. caninum*) has raised concerns

within the veterinary profession (von Samson-Himmelstjerna et al., 2021; Marsh and

Lakritz, 2023). Adopting a risk assessment-based approach to limit unnecessary use

of parasiticides has the potential to mitigate the risk of resistance selection and its

spread, and should be an important consideration in responsible parasite control

(Selzer and Epe, 2021; Bagster and Elsheikha, 2022).

**Acknowledgements**

Funding was provided by the Margo Roslyn Flood Bequest (Sydney School of

Veterinary Science, The University of Sydney, Australia), and by The Canine

Research Foundation and Dogs Victoria (School of Veterinary Science, University of

Queensland, Australia).

**References**

Bagster, A., Elsheikha, H., 2022. Perception of UK companion animal veterinarians on risk assessment based parasite control. Vet Parasitol Reg Stud Reports 34, 100774. doi: 10.1016/j.vprsr.2022.100774

Bolt, B.J., Rodgers, F.H., Shafie, M., Kersey, P.J., Berriman, M., Howe, K.L., 2018. Using WormBase ParaSite: An integrated platform for exploring helminth genomic data. Methods Mol Biol 1757, 471-491. doi: 10.1007/978-1-4939-7737-6_15

Burton, K.W., Hegarty, E., Couto, C.G., 2024. Retrospective analysis of canine fecal flotation and coproantigen immunoassay hookworm positive results in Greyhounds and other dog breeds. Vet Parasitol Reg Stud Reports 51, 101026. doi: 10.1016/j.vprsr.2024.101026

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods 13, 581-583. doi: 10.1038/nmeth.3869

Dale, A., Xu, G., Kopp, S.R., Jones, M.K., Kotze, A.C., Abdullah, S., 2024. Pyrantel resistance in canine hookworms in Queensland, Australia. Vet Parasitol Reg Stud Reports 48, 100985. doi: 10.1016/j.vprsr.2024.100985

Diawara, A., Schwenkenbecher, J.M., Kaplan, R.M., Prichard, R.K., 2013. Molecular and biological diagnostic tests for monitoring benzimidazole resistance in human soil-transmitted helminths. Am J Trop Med Hyg 88, 1052-1061. doi: 10.4269/ajtmh.12-0484

Furtado, L.F., Bello, A.C., dos Santos, H.A., Carvalho, M.R., Rabelo, E.M., 2014. First identification of the F200Y SNP in the beta-tubulin gene linked to

benzimidazole resistance in *Ancylostoma caninum*. Vet Parasitol 206, 313-316.

doi: 10.1016/j.vetpar.2014.10.021

Gasser, R.B., Chilton, N.B., Hoste, H., Beveridge, I., 1993. Rapid sequencing of

rDNA from single worms and eggs of parasitic helminths. Nucleic Acids Res 21,

2525-2526. doi: 10.1093/nar/21.10.2525

Gibbs, H.C., 1958. On the gross and microscopic lesions produced by the adults and

larvae of *Dochmoides stenocephala* (Raillet, 1884) in the dog. Can J Comp

Med Vet Sci 22, 382-385.

Harris, T.W., Arnaboldi, V., Cain, S., Chan, J., Chen, W.J., Cho, J., Davis, P., Gao,

S., Grove, C.A., Kishore, R., Lee, R.Y.N., Muller, H.M., Nakamura, C., Nuin, P.,

Paulini, M., Raciti, D., Rodgers, F.H., Russell, M., Schindelman, G., Auken,

K.V., Wang, Q., Williams, G., Wright, A.J., Yook, K., Howe, K.L., Schedl, T.,

Stein, L., Sternberg, P.W., 2020. WormBase: a modern Model Organism

Information Resource. Nucleic Acids Res 48, D762-D767. doi:

10.1093/nar/gkz920

Hawdon, J.M., Schad, G.A., 1990. Serum-stimulated feeding in vitro by third-stage

infective larvae of the canine hookworm *Ancylostoma caninum*. J Parasitol 76,

394-398. doi: 10.2307/3282673

Hotez, P., Hawdon, J., Schad, G.A., 1993. Hookworm larval infectivity, arrest and

amphiparatenesis: the *Caenorhabditis elegans* Daf-c paradigm. Parasitol Today

9, 23-26. doi: 10.1016/0169-4758(93)90159-d

Howe, K.L., Bolt, B.J., Shafie, M., Kersey, P., Berriman, M., 2017. WormBase

ParaSite - a comprehensive resource for helminth genomics. Mol Biochem

Parasitol 215, 2-10. doi: 10.1016/j.molbiopara.2016.11.005

Jaeger, L.H., Carvalho-Costa, F.A., 2017. Status of benzimidazole resistance in

intestinal nematode populations of livestock in Brazil: a systematic review. BMC

Vet Res 13, 358. doi: 10.1186/s12917-017-1282-2

Jimenez Castro, P.D., Howell, S.B., Schaefer, J.J., Avramenko, R.W., Gilleard, J.S.,

Kaplan, R.M., 2019. Multiple drug resistance in the canine hookworm

*Ancylostoma caninum*: an emerging threat? Parasit Vectors 12, 576. doi:

10.1186/s13071-019-3828-6

Jimenez Castro, P.D., Venkatesan, A., Redman, E., Chen, R., Malatesta, A., Huff,

H., Zuluaga Salazar, D.A., Avramenko, R., Gilleard, J.S., Kaplan, R.M., 2021.

Multiple drug resistance in hookworms infecting greyhound dogs in the USA. Int

J Parasitol Drugs Drug Resist 17, 107-117. doi: 10.1016/j.ijpddr.2021.08.005

Jones, B.P., van Vliet, A.H.M., LaCourse, E.J., Betson, M., 2022. *In silico* docking of

nematode β-tubulins with benzimidazoles points to gene expression and

orthologue variation as factors in anthelmintic resistance. Front Trop Dis 3,

898814. doi: 10.3389/fitd.2022.898814

Kitchen, S., Ratnappan, R., Han, S., Leasure, C., Grill, E., Iqbal, Z., Granger, O.,

O'Halloran, D.M., Hawdon, J.M., 2019. Isolation and characterization of a

naturally occurring multidrug-resistant strain of the canine hookworm,

*Ancylostoma caninum*. Int J Parasitol 49, 397-406. doi:

10.1016/j.ijpara.2018.12.004

Kopp, S.R., Kotze, A.C., McCarthy, J.S., Coleman, G.T., 2007. High-level pyrantel

resistance in the hookworm *Ancylostoma caninum*. Vet Parasitol 143, 299-304.

doi: 10.1016/j.vetpar.2006.08.036

Kotze, A.C., Coleman, G.T., Mai, A., McCarthy, J.S., 2005. Field evaluation of

anthelmintic drug sensitivity using in vitro egg hatch and larval motility assays

with *Necator americanus* recovered from human clinical isolates. Int J Parasitol

35, 445-453. doi: 10.1016/j.ijpara.2004.12.007

Kotze, A.C., Lowe, A., O'Grady, J., Kopp, S.R., Behnke, J.M., 2009. Dose-response

assay templates for in vitro assessment of resistance to benzimidazole and

nicotinic acetylcholine receptor agonist drugs in human hookworms. Am J Trop

Med Hyg 81, 163-170. doi: 10.4269/ajtmh.2009.81.163

Kwa, M.S., Kooyman, F.N., Boersema, J.H., Roos, M.H., 1993. Effect of selection for

benzimidazole resistance in *Haemonchus contortus* on beta-tubulin isotype 1

and isotype 2 genes. Biochem Biophys Res Commun 191, 413-419. doi:

10.1006/bbrc.1993.1233

Leutenegger, C.M., Evason, M.D., Willcox, J.L., Rochani, H., Richmond, H.L.,

Meeks, C., Lozoya, C.E., Tereski, J., Loo, S., Mitchell, K., Andrews, J., Savard,

C., 2024. Benzimidazole F167Y polymorphism in the canine hookworm,

*Ancylostoma caninum*: Widespread geographic, seasonal, age, and breed

distribution in United States and Canada dogs. Int J Parasitol Drugs Drug

Resist 24, 100520. doi: 10.1016/j.ijpddr.2024.100520

Leutenegger, C.M., Lozoya, C.E., Tereski, J., Savard, C., Ogeer, J., Lallier, R., 2023.

Emergence of *Ancylostoma caninum* parasites with the benzimidazole

resistance F167Y polymorphism in the US dog population. Int J Parasitol Drugs

Drug Resist 21, 131-140. doi: 10.1016/j.ijpddr.2023.01.001

Marsh, A.E., Lakritz, J., 2023. Reflecting on the past and fast forwarding to present

day anthelmintic resistant *Ancylostoma caninum*-A critical issue we neglected

to forecast. Int J Parasitol Drugs Drug Resist 22, 36-43. doi:

10.1016/j.ijpddr.2023.04.003

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput

sequencing reads. EMBnet.journal 17, 10-12. doi: 10.14806/ej.17.1.200

Massetti, L., Wiethoelter, A., McDonagh, P., Rae, L., Marwedel, L., Beugnet, F.,

Colella, V., Traub, R.J., 2022. Faecal prevalence, distribution and risk factors

associated with canine soil-transmitted helminths contaminating urban parks

across Australia. Int J Parasitol 52, 637-646. doi: 10.1016/j.ijpara.2022.08.001

McKean, E.L., Grill, E., Choi, Y.J., Mitreva, M., O'Halloran, D.M., Hawdon, J.M.,

2024. Altered larval activation response associated with multidrug resistance in

the canine hookworm *Ancylostoma caninum*. Parasitology 151, 271-281. doi:

10.1017/S0031182023001385

Miller, T.A., 1968. Pathogenesis and immunity in hookworm infection. Trans R Soc

Trop Med Hyg 62, 473-489. doi: 10.1016/0035-9203(68)90130-2

Mohammedsalih, K.M., Krücken, J., Bashar, A., Juma, F.R., Abdalmalaik, A.A.H.,

Khalafalla, A., Abakar, A., Coles, G., von Samson-Himmelstjerna, G., 2021.

Susceptible trichostrongyloid species mask presence of benzimidazole-

resistant *Haemonchus contortus* in cattle. Parasit Vectors 14, 101. doi:

10.1186/s13071-021-04593-w

Ploeger, H.W., Everts, R.R., 2018. Alarming levels of anthelmintic resistance against

gastrointestinal nematodes in sheep in the Netherlands. Vet Parasitol 262, 11-

15. doi: 10.1016/j.vetpar.2018.09.007

R Core Team, 2022. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-

project.org/.

Ramunke, S., Melville, L., Rinaldi, L., Hertzberg, H., de Waal, T., von Samson-

Himmelstjerna, G., Cringoli, G., Mavrot, F., Skuce, P., Krucken, J., Demeler, J.,

2016. Benzimidazole resistance survey for *Haemonchus*, *Teladorsagia* and

*Trichostrongylus* in three European countries using pyrosequencing including

the development of new assays for *Trichostrongylus*. Int J Parasitol Drugs Drug

Resist 6, 230-240. doi: 10.1016/j.ijpddr.2016.10.002

Ryan, G.E., 1976. Helminth parasites of the fox (*Vulpes vulpes*) in New South

Wales. Aust Vet J 52, 126-131. doi: 10.1111/j.1751-0813.1976.tb05445.x

Schulz, J.D., Moser, W., Hurlimann, E., Keiser, J., 2018. Preventive Chemotherapy

in the Fight against Soil-Transmitted Helminthiasis: Achievements and

Limitations. Trends Parasitol 34, 590-602. doi: 10.1016/j.pt.2018.04.008

Schwenkenbecher, J.M., Albonico, M., Bickle, Q., Kaplan, R.M., 2007.

Characterization of beta-tubulin genes in hookworms and investigation of

resistance-associated mutations using real-time PCR. Mol Biochem Parasitol

156, 167-174. doi: 10.1016/j.molbiopara.2007.07.019

Selzer, P.M., Epe, C., 2021. Antiparasitics in Animal Health: Quo Vadis? Trends

Parasitol 37, 77-89. doi: 10.1016/j.pt.2020.09.004

Stocker, T., Scott, I., Šlapeta, J., 2023. Unambiguous identification of *Ancylostoma*

*caninum* and *Uncinaria stenocephala* in Australian and New Zealand dogs from

faecal samples. Aust Vet J 101, 373-376. doi: 10.1111/avj.13272

Stocker, T., Ward, M.P., Šlapeta, J., 2024. Nationwide USA re-analysis of amplicon

metabarcoding targeting β-tubulin isoform-1 reveals absence of benzimidazole

resistant SNPs in *Ancylostoma braziliense*, *Ancylostoma tubaeforme* and

*Uncinaria stenocephala*. Vet Parasitol 328, 110118. doi:

10.1016/j.vetpar.2024.110118

Venkatesan, A., Jimenez Castro, P.D., Morosetti, A., Horvath, H., Chen, R.,

Redman, E., Dunn, K., Collins, J.B., Fraser, J.S., Andersen, E.C., Kaplan, R.M.,

Gilleard, J.S., 2023. Molecular evidence of widespread benzimidazole drug

resistance in *Ancylostoma caninum* from domestic dogs throughout the USA

and discovery of a novel beta-tubulin benzimidazole resistance mutation. PLoS

Pathog 19, e1011146. doi: 10.1371/journal.ppat.1011146

Vercruysse, J., Levecke, B., Prichard, R., 2012. Human soil-transmitted helminths:

implications of mass drug administration. Curr Opin Infect Dis 25, 703-708. doi:

10.1097/QCO.0b013e328358993a

von Samson-Himmelstjerna, G., Blackhall, W.J., McCarthy, J.S., Skuce, P.J., 2007.

Single nucleotide polymorphism (SNP) markers for benzimidazole resistance in

veterinary      nematodes.      Parasitology      134,      1077-1086.      doi:

10.1017/S0031182007000054

von Samson-Himmelstjerna, G., Thompson, R.A., Krucken, J., Grant, W., Bowman,

D.D., Schnyder, M., Deplazes, P., 2021. Spread of anthelmintic resistance in

intestinal helminths of dogs and cats is currently less pronounced than in

ruminants and horses - Yet it is of major concern. Int J Parasitol Drugs Drug

Resist 17, 36-45. doi: 10.1016/j.ijpddr.2021.07.003

WHO, 2017. Guideline: Preventive chemotherapy to control soil-transmitted helminth

infections in at-risk population groups. World Health Organization, Geneva.

 **Figure legends**

**Fig. 1.** Molecular profiling of hookworm species using internal transcribed spacer

(ITS)-2 rDNA and benzimidazole susceptibility using isotype-1 β-tubulin. (A)

Percentage of hookworm species within each sample (depicted as percentage) and

number of sequencing reads per sample in the bar chart directly below. Sampled

with hookworm-like stages that were profiled using metabarcoded deep Illumina

amplicon sequencing at ITS-2 rDNA. All but one sample came from dogs. The

single, cat sample (TS042) is on the very right. The broad scale of the samples is

indicated above the bar chart. Amplicon sequence variants (ASVs) were pooled

according to species of nematodes to which they belonged using a reference

sequence alignment. Proportion of each species (*Ancylostoma caninum*,

*Ancylostoma tubaeforme, Uncinaria stenocephala*) is colour coded in the stacked bar

chart; ASVs that belonged to unrelated nematodes are depicted as 'pseudoparasites'

and represented spurious parasite sequences, specifically trichostrongylid species

from sheep and rabbits. The bar chart underneath represents the number of

sequencing reads for each sample, and samples with <2,000 sequence reads are

depicted with a star. Samples are vertically aligned across the bar charts. (B) Dog

samples profiled for hookworm species identification using the BZ167 amplicon of

the isotype-1 β-tubulin, specifically the region covering amino acids Q134 and F167

conferring benzimidazole resistance. Only ASVs of hookworm as a percentage of the

total are used and the sequencing read depth is depicted underneath. Samples that

did not amplify hookworms are indicated by squares and samples with <2,000

sequence reads are depicted with stars. (C) Bar chart for Q134 and F167 isotype-1

β-tubulin of *A. caninum* where the proportion of an alternative residue coding for an

amino acid that confers benzimidazole resistance are shown. Samples with no bar

chart are those that did not amplify *A. caninum*, thus are no *A. caninum* ASVs.

**Fig. 2.** Distribution of *Ancylostoma caninum* and benzimidazole-resistant isotype-1 β-

tubulin single nucleotide polymorphism. Proportional symbol maps of (A) the number

of *A. caninum*-positive samples, and (B) number of Q134H (CA<u>A</u>>CA<u>T</u>) and (C)

number of F167Y (T<u>T</u>C>T<u>A</u>C) isotype-1 β-tubulin of *A. caninum*-positive samples.

Samples were summed per postcode in New South Wales, Australia. Samples are

joined to postcodes of Australia shapefile (Geocentric Datum of Australia, GDA

1994) in ArcGIS® Pro.

**Fig. 3.** *Ancylostoma caninum* 50% inhibitory concentration ($IC_{50}$) for thiabendazole in

egg hatch assays. (A) Correlation $IC_{50}$ values for *A. caninum* samples for

thiabendazole agar-based egg hatch assay (EHA) from Queensland, Australia with

the percentage of F167Y (T<u>T</u>C>T<u>A</u>C) isotype-1 β-tubulin of *A. caninum* present in

the sample. Simple linear regression is depicted as a solid line with 95% confidence

interval (95% CI) depicted as a dotted line. (B) Histogram of $IC_{50}$ values for *A.*

*caninum* samples from Queensland, Australia. A susceptible *A. caninum* sample,

that was confirmed to have 100% faecal egg count (FEC) reduction, is colour coded

(black). The samples with >0.15 μg/mL thiabendazole $IC_{50}$ values have their

identifiers above the bars.

**Table 1.** Summary of *Ancylostoma caninum* egg hatch assay with frequency of Q134H and F167Y polymorphisms in isotype-1 β-

tubulin.

| Sample ID | Breed | EPG | $IC_{50}$ (µg/mL)[a] | $IC_{50}$ 95% CI (µg/mL)[a] | Q134H[c] | F167Y[c] |
|---|---|---|---|---|---|---|
| High $IC_{50}$>0.15 µg/mL of thiabendazole | | | | | | |
| D3:TS89 | Greyhound | 1,275 | 0.429 | 0.344 to 0.534 | 0% | 89.91% |
| D1:TS87[d] | Greyhound | 1,050 | 0.298 | 0.216 to 0.412 | 0% | 85.47% |
| D2:TS88[d] | Greyhound | 645 | 0.262 | 0.237 to 0.289 | 0% | 88.80% |
| D6:TS92[e] | Crossbreed | 540 | 0.239 | 0.201 to 0.285 | 0% | 83.11% |
| D18/D20:TS104/106[e] | Crossbreed | 225 | 0.170 | 0.121 to 0.239 | 0%/0% | 100%/77.54% |
| D17:TS103 | Greyhound | 5,535 | 0.194 | 0.150 to 0.250 | 2.55% | 81.23% |
| Low $IC_{50}$<0.15 µg/mL of thiabendazole | | | | | | |
| D5:TS91 | Greyhound | 14,910 | 0.056 | 0.048 to 0.066[b] | 0% | 10.69% |

| | | | | | | |
|---|---|---|---|---|---|---|
| D9:TS95 | Greyhound | 930 | 0.050 | 0.043 to 0.059[b] | 18.27% | 33.51% |
| D10:TS96 | Greyhound | 4,080 | 0.043 | 0.040 to 0.046[b] | 0% | 22.67% |
| D8:TS94 | Greyhound | 390 | 0.039 | 0.031 to 0.048[b] | 0% | 29.26% |
| D12:TS98 | Greyhound | 3,930 | 0.037 | 0.035 to 0.039[b] | 0% | 0% |
| D15:TS101 | Greyhound | 720 | 0.036 | 0.031 to 0.041[b] | 0% | 6.56% |
| D16:TS102 | Greyhound | 885 | 0.021 | 0.013 to 0.033[b] | 7.43% | 22.53% |

841    [a] Thiabendazole concentration in µg/mL.

842    [b] Drug-susceptible *A. caninum* - 0.046 and 0.048 µg/mL (Jimenez Castro et al., 2019) and 0.045 µg/mL (Diawara et al., 2013).

843    [c] Percentage of canonical benzimidazole resistance codons on the isotype-1 β-tubulin.

844    [d,e] Same dog.

845    EPG, *A. caninum* eggs per gram of faeces; $IC_{50}$, 50% inhibition concentration; 95% CI, 95% confidence interval.

A

*Ancylostoma caninum* samples

New South Wales, Australia

B

F167Y resistant *Ancylostoma caninum*

C

Q134H resistant *Ancylostoma caninum*