1

Widespread occurrence of benzimidazole resistance SNPs in the canine hookworm,

Ancylostoma caninum, in Australia

Swaid Abdullah[1*,&], Thomas Stocker[2,&], Hyungsuk Kang[1], Ian Scott[3], Douglas Hayward[4],

Susan Jaensch[4], Michael P. Ward[2], Malcolm K. Jones[1], Andrew C. Kotze[1], Jan Šlapeta[2,4*]

[1] The University of Queensland, School of Veterinary Science, Gatton 4343, QLD,

Australia

[2] Sydney School of Veterinary Science, Faculty of Science, University of Sydney, Sydney,

New South Wales, Australia

[3] School of Veterinary Science, Massey University, Palmerston North, New Zealand

[4] Vetnostics NSW - North Ryde Laboratory, Macquarie Park, New South Wales, Australia

[6] Sydney Institute for Infectious Diseases, The University of Sydney, New South Wales,

Australia.

*Corresponding authors: Jan Šlapeta, jan.slapeta@sydney.edu.au; Swaid Abdullah

swaid.abdullah@uq.edu.au

[&] Shared first authors

Abstract

Canine hookworm (*Ancylostoma caninum*), a gastrointestinal nematode of domestic dogs, principally infects the small intestine of dogs and has the potential to cause zoonotic disease. In greyhounds and pet dogs in the USA, *A. caninum* has been shown to be resistant to multiple anthelmintics. We conducted a molecular survey of benzimidazole resistance in *A. caninum* from dogs at veterinary diagnostic centers in Australia and New Zealand. First, we implemented an ITS-2 rDNA deep amplicon metabarcoding sequencing approach to ascertain the species of hookworms infecting dogs in the region. Then, we evaluated the frequency of the canonical F167Y and Q134H isoform-1 β-tubulin mutations, which confer benzimidazole resistance, using the same sequencing approach. The most detected hookworm species in diagnostic samples was *A. caninum* (90%; 83/92); the related Northern hookworm (*Uncinaria stenocephala*) was identified in 11% (10/92) of the diagnostic samples. There was a single sample with coinfection by *A. caninum* and *U. stenocephala*. Both isoform-1 β-tubulin mutations were present in *A. caninum*, 49% and 67% for Q134H and F167Y, respectively. Mutation F167Y in the isoform-1 β-tubulin mutation was recorded in *U. stenocephala* for the first time. Canonical benzimidazole resistance codons 198 and 200 mutations were absent. Egg hatch assays performed on a subset of the *A. caninum* samples showed significant correlation between IC_{50} to thiabendazole and F167Y, with an increased IC_{50} for samples with >75% F167Y mutation. We detected 14% of dogs with >75% F167Y mutation in *A. caninum*. Given that these samples were collected from dogs across various regions of Australia, the present study suggests that benzimidazole resistance in *A. caninum* is widespread. Therefore, to mitigate the risk of resistance selection and further spread,

adoption of a risk assessment-based approach to limit unnecessary anthelmintic use

should be a key consideration for future parasite control.

Keywords:

Ancylostoma caninum, Multiple-drug resistance (MDR), Multiple anthelmintic drug

resistant (MADR), Greyhounds, Anthelmintics, Canine health

Highlights:

• High prevalence of benzimidazole resistance mutations in Australian canine

hookworms.

• *Ancylostoma caninum* detected in 90% of samples and *Uncinaria stenocephala*

in 11% of diagnostic samples.

• F167Y and Q134H mutations found in *A. caninum* and F167Y in *U. stenocephala*.

• Egg hatch assay confirms correlation between F167Y mutation and

benzimidazole resistance.

Introduction

For many years the prevailing dogma regarding the use of anthelmintics in small animals, such as dogs and cats, as well as in humans, had been that the development of anthelmintic resistance would be slow to occur, or may never occur or be limited in spread (Schulz et al., 2018; Marsh and Lakritz, 2023). The theory behind this dogma was that, at the time of deworming, the *refugia* - the population of susceptible nematodes infecting the host - remains substantial (von Samson-Himmelstjerna et al., 2021). By contrast, in ruminants, routine and indiscriminate mass treatment with anthelminthics has led to widespread drug resistance, because the *refugia* has not been retained (Vercruysse et al., 2012; von Samson-Himmelstjerna et al., 2021). However, despite these differences between ruminants and small animals, a number of reports over recent years have indicated that resistance in canine hookworms is indeed a widespread problem in North America (Jimenez Castro et al., 2019; Kitchen et al., 2019; Venkatesan et al., 2023).

Nowadays, the anthelminthics class of benzimidazoles is largely ineffective in ruminants, the result of well documented single nucleotide polymorphisms (SNPs) that lead to nematodes carrying alleles resistant to these anthelminthics (Kwa et al., 1993; von Samson-Himmelstjerna et al., 2007). In humans, hookworms are commonly treated using benzimidazoles, specifically albendazole and mebendazole (WHO, 2017). Between 1995 and 2015, the efficacy of both albendazole and mebendazole for the treatment of human hookworm infection has decreased by as much as 15%, but resistance as such has not been documented (Schulz et al., 2018).

In dogs, anthelminthics resistance was first documented 20 years ago in hookworms to

the pyrimidine-derivative anthelmintic pyrantel in Queensland, Australia (Kopp et al.,

2007), and currently this anthelmintic is considered ineffective for treating hookworms

in dogs in Australia (Dale et al., 2024). However, resistance to benzimidazoles and other

anthelmintic classes for dogs in Australia has not been reported. The first report of

resistance to benzimidazoles in canine hookworms was from the USA in 2019 (Kitchen

et al., 2019). Molecular, phenotypic, and experimental infections evidence

demonstrated benzimidazole resistance in *A. caninum* was published independently by

two groups in 2019 (Jimenez Castro et al., 2019; Kitchen et al., 2019). These reports first

emerged in North America and were followed by an extensive surveillance for the

canonical F167Y (T<u>T</u>C>T<u>A</u>C) isoform-1 β-tubulin mutation conferring benzimidazole

resistance (Leutenegger et al., 2023; Venkatesan et al., 2023; Leutenegger et al., 2024).

A new mutation in isoform-1 β-tubulin at Q134H (CA<u>A</u>>CA<u>T</u>) was experimentally

established as another mutation responsible for resistance to benzimidazoles in *A.*

caninum (Venkatesan et al., 2023).

It is speculated that benzimidazole resistance in *A. caninum* emerged in racing

greyhound dogs because of indiscriminate and frequent deworming in racing kennels

(Jimenez Castro et al., 2019; Kitchen et al., 2019; Leutenegger et al., 2024). Evidence of

multiple anthelmintic drug resistance (MADR) in *A. caninum* led the American

Association of Veterinary Parasitologists to establish a hookworm task force to address

this issue in 2021 (von Samson-Himmelstjerna et al., 2021; Marsh and Lakritz, 2023). The

screening of thousands of dogs positive to hookworms in the USA and Canada

demonstrated that the canonical F167Y (T<u>T</u>C>T<u>A</u>C) isoform-1 β-tubulin mutation was

present in 11.3% (715/6,329) of faecal *A. caninum* positive samples (Leutenegger et al.,

2024). In a smaller study of samples from the USA, 49.7% (156/314) of the individual

samples tested positive for the F167Y (T$\underline{T}$C>T$\underline{A}$C) mutation (Venkatesan et al., 2023).

Reanalysis of next-generation sequence data from Venkatesan et al. (2023) for the

presence of non-*A. caninum* sequences revealed the presence of other hookworms

(*Ancylostoma braziliense*, *Ancylostoma tubaeforme* and *Uncinaria stenocephala*), but

only the benzimidazole susceptible residues of isoform-1β-tubulin were found in these

species (Stocker et al., 2024).

The aim of this study was to investigate the extent of benzimidazole resistance SNPs in

canine hookworm (*A. caninum*) outside North America, specifically in Australia and New

Zealand. We utilised opportunistic sampling from four different sources in Australia and

New Zealand. All hookworm-positive samples were screened using ITS-2 rDNA PCR and

deep amplicon sequencing to determine hookworms' species present. To investigate

resistance to benzimidazoles, we used deep amplicon sequencing to evaluate the

nucleotide sequence of the isoform-1 β-tubulin mutations that are known to be

associated with resistance to this drug class, specifically the regions that encode amino

acids Q_{134}, F_{167}, E_{198} and F_{200}. We then undertook egg hatch assay (EHA) on a subset of the

samples to correlate the molecular signatures, specifically frequency of F_{167}, with

benzimidazole-resistance phenotype of these field isolates.

Material and methods

Ethics statement

The use of residual clinical samples was in accordance with the University of Sydney

Animal Ethics Committee Protocol (2024/2508). Dog faecal samples for larval cultures

were obtained in accordance with the University of Queensland Animal Ethics

Committee Protocol (2021/AE001169) and faecal egg count reduction test on one dog

was conducted in accordance with University of Queensland Animal Ethics Committee

Protocol (SVS/111/20).

Faecal samples with hookworm infection

Faecal material from New South Wales, Australia and New Zealand was provided for the

purpose of diagnostics used in decision-making by the veterinary practitioner. Results

and material used for hookworm species and benzimidazole susceptibility were de-

identified. Faecal samples were provided by registered veterinary practitioners in

accordance with the Veterinary Practice and Animal Welfare Acts. Samples were

submitted for parasitological diagnostics to Vetnostics (Laverty Pathology—North Ryde

Laboratory, Sydney, New South Wales, Australia) and Veterinary Pathology Diagnostics

Services (VPDS, The University of Sydney, New South Wales, Australia); refrigerated

residual faecal samples that previously returned a positive result for hookworms at the

clinical diagnostic laboratory were received for molecular species characterisation. In

addition, samples from Massey University, Palmerston North, New Zealand that

contained hookworm eggs were processed to isolate the eggs, which were stored in 80%

ethanol for transfer to VPDS.

Material from Queensland, Australia was obtained from dogs from adoption and shelter

facilities that receive rescued or surrendered animals from greyhound rescue

organizations and pounds across southeast Queensland. Dogs within these facilities are

kept in individual pens, which are cleaned and disinfected daily. Fresh faecal samples

were collected from each dog after natural defecation and screened for hookworm

infection using faecal flotation as described below.

Faecal egg counts, hookworm egg recovery and DNA isolation

Faecal egg counts were performed with saturated (NaCl) solution using Whitlock

Universal McMaster 4 chamber worm egg counting slide in two different laboratories. At

the University of Queensland, 3 g of faecal samples was homogenised with 42 ml of

saturated salt solution (specific gravity = 1.21) and filtered through a tea strainer and the

filtrate was immediately pipetted into the counting chamber (4 compartments each of

0.5 ml volume). Counting all compartments (2 ml) delivered minimum detection of 7 egg

per gram (absence of eggs indicates <7 eggs per gram in the sample). At the University of

Sydney, 3 g of faecal samples was homogenised with 60 ml of saturated salt solution

(specific gravity = 1.21) and filtered via coarse strainer. We counted 2 compartments

each of 0.5 ml volume resulting in minimum detection of 20 egg per gram (absence of

eggs indicates <20 eggs per gram in the sample). Only strongyle-like (=hookworm) eggs

were counted.

At the University of Sydney, the remainder of the filtered homogenate was poured into 50

ml Falcon tube and centrifuged at 800×g for 10 minutes for centrifugal floatation. The

upper most supernatant (~10 ml) was collected with a plastic Pasteur pipette into a new

clean 50 ml Falcon tube and topped with tap water and centrifuged at 800×g for 10

minutes. The supernatant was decanted and the pellet that contained the concentrated

eggs was retained. Then ~1 ml of tap water was added with a clean plastic Pasteur pipette

to disturb the pellet by repeated aspiration until a homogeneous solution formed. The

homogenate was transferred to a clean 1.5ml Eppendorf tube and spun at 8,000×g for 10

minutes. The supernatant was completely removed and the Eppendorf tube with the egg

pellet was stored at -20 °C until further use.

At the University of Queensland, eggs were purified for use in the egg hatch assay (see

later). The remaining eggs were suspended in distilled water in 1.5 ml microtubes,

incubated for 72 hours at 28˚C and allowed to hatch. The hatched larvae were stored at

-20˚C until further use.

The total genomic DNA was isolated from hookworm eggs using the Isolate II Fecal DNA

Kit (Meridian Bioscience, Australia) according to the manufacturer's instructions with

following homogenisation step. For purified eggs,750 µL of lysis buffer was added to the

Eppendorf tube with the pellet and homogenised by repeated aspiration with a pipette,

then all the contents were transferred into bead-beating tube and the eggs disrupted and

homogenised using a high-speed benchtop homogeniser, FastPrep-24 (MP Biomedicals,

Australia), for 40 s at 6.0 m/s. Every 11[th] sample was a blank sample that only contained

the lysis buffer. For hatched larvae, Monarch Genomic DNA Purification Kit (New England

Biolab, Australia) was used according to the manufacturer's instructions. Purified

genomic DNA samples and blanks were isolated into 100 µL of elution buffer (10 mM Tris-

Cl, pH = 8.5) and stored at −20 °C until further use.

Amplicon metabarcoding and Illumina deep sequencing

All isolated DNA samples were first screened for the presence of nematode DNA using

nematode ITS2 rDNA quantitative PCR and samples with C_t<35 were considered as a

suitable DNA material for follow up processes.

Total DNA was subjected to three independent real-time PCR amplifications at the

University of Sydney as previously described (Stocker et al., 2023). ITS-2 rDNA (ITS-2) was

amplified by the NC-1 and NC-2 primers designed by Gasser et al. (1993), and two

isoform-1 β-tubulin regions were amplified using assays BZ167 and BZ200 developed by

Jimenez Castro et al. (2019), which   include key diagnostic SNPs conferring

benzimidazole resistance. All primers were adapted for Illumina metabarcoding using

Next Generation Sequencing (NGS). The ITS-2 target was amplified using 32-36

amplification cycles, while 40 amplification cycles were used for isoform-1 β-tubulin

(BZ167, BZ200). All reactions utilized SYBR-chemistry using SensiFAST™SYBR® No-ROX

mix (Meridian Bioscience, Australia). Each real-time PCR mix (25 μL) included 2 μL of

template DNA (neat) and primers at final concentrations of 400 nM. All PCR reactions

were prepared using a Myra robotic liquid handling system (Bio Molecular Systems,

Australia). Reactions were run on a CFX96 Real-Time PCR Detection System with

corresponding CFX Manager software (BioRad, Australia). The protocol involved an initial

denaturation at 95 °C for 3 min, followed by 32 or 40 cycles of 95 °C for 5 s, 60 °C for 15

s, 72 °C for 15 s, and a final melt curve analysis. Each run included a no-template control

(ddH$_2$O) to monitor for potential contamination. Amplicons were submitted for amplicon

sequencing using Illumina NGS at the Ramaciotti Centre for Genomics, University of New

South Wales, Sydney, Australia and sequenced on MiSeq using MiSeq Reagent Kits v2

(250PE, Illumina). De-multiplexed FastQ files were generated via BaseSpace (Illumina).

Analysis of the Illumina sequenced amplicons

All FastQ files were processed through R package 'dada2' v1.26.0 (Callahan et al., 2016)

in R v4.2.2 (R Core Team, 2022), to obtain amplicon sequence variants (ASV) and their

abundance per sequenced amplicon. The dada2 pipeline at the University of Sydney

includes filtering, error estimation and denoising, before paired-end reads were merged

to reconstruct the full ASV before removal of chimeras. Prior to processing through the

dada2 pipeline, primers were removed from all forward and reverse FastQ reads using

'cutadapt' version 4.0 (Martin, 2011).

The assignment of species based on ITS-2 ASVs was carried out in CLC Main Workbench

v22.0 (Qiagen Australia) as previously described (Stocker et al., 2024). All ASVs were

imported into CLC Main Workbench and top blastn hits against 'nr' database at NCBI

was identified. ITS-2 sequences were discarded for any ASVs where the top blastn hit

was not hookworm (Ancylostomatidae). For species assignment we constructed

alignment of AVSs with an in-house curated reference ITS-2 hookworm sequence library

consisting of publicly available sequences (KP844736; MT345056; JQ812692; LC036567;

JX220891; EU344797; AB793527). The resulting multiple sequence alignment was

subject to phylogenetic reconstruction, where species identity was based on manual

assignment of ASV to a species based on position on the phylogenetic tree relative to the

reference ITS-2 sequences.

For the β-tubulin sequence data (BZ167, BZ200 assay), the assignment of species was

carried with the aid of CLC Main Workbench v22.0 (Qiagen Australia) as previously

described in Stocker et al. (2024). Briefly, all ASVs were imported into CLC Main

Workbench and top blastn hits against 'nr' database at NCBI was identified. All ASVs

where the top blastn hit was a hookworm (Ancylostomatidae) β-tubulin were retained for

further analysis. The reference hookworm alignments of BZ167 and BZ200 amplicons

according to Stocker et al. (2024) was used to unambiguously identify ASVs identity and

only those ASVs that were confirmed based on phylogenetic analysis were included in

further analyses. The sequence alignment included β-tubulin reference sequences from

A. caninum (DQ459314) (Schwenkenbecher et al., 2007), *Ancylostoma duodenale*

(EF392850), *U. stenocephala* (Stocker et al., 2023), and *N. americanus* (EU182348),

which enabled us to identify the exon/intron boundary and residues that code for

susceptibility to benzimidazoles, amino acids amino acids Q_{134}, F_{167}, E_{198} and F_{200}

(Venkatesan et al., 2023). Our reference alignment included sequence data for isoform-

1 β-tubulin from *A. caninum*, *A. ceylanicum*, *A. duodenale* and *Necator americanus* at

'WormBase ParaSite' (Howe et al., 2017; Bolt et al., 2018; Harris et al., 2020).

Following the dada2 pipeline, and species assignment raw read counts were converted

into relative abundance (percentages) through. Any species read counts with less <25

reads were disregarded as spurious. Proportions of SNP for amino acids amino acids

Q_{134}, F_{167}, E_{198} and F_{200} were calculated per species for each of the amplicons of isoform-

1 β-tubulin (BZ167, BZ200).

The number of samples, and number of *A. caninum* samples with Q134H (CA<u>A</u>>CA<u>T</u>) and

F167Y (T<u>T</u>C>T<u>A</u>C) mutations separately, were summed per postcode in New South

Wales. Proportional symbol maps were created by joining this data to postcodes of

Australia shapefile (Geocentric Datum of Australia, GDA 1994) in ArcGIS® Pro. Spearman

rank correlations (r_{SP}) were calculated between EPG per sample and both percentages of

Q134H (CA<u>A</u>>CA<u>T</u>) and F167Y (T<u>T</u>C>T<u>A</u>C) mutation. In addition, the EPG between

resistant versus not resistant status for both mutations was assessed using a Kruskal-

Wallis non-parametric analysis of variance. Spearman rank correlations were also

calculated between EPG, percentage of Q134H (CA<u>A</u>>CA<u>T</u>) mutation, percentage of

F167Y (T$\underline{T}$C>T$\underline{A}$C) mutation and age. Finally, the association between breed (greyhound

versus other) and EPG, percentage of Q134H (CA$\underline{A}$>CA$\underline{T}$) mutation and percentage of

F167Y (T$\underline{T}$C>T$\underline{A}$C) mutation was assessed using a Kruskal-Wallis non-parametric

analysis of variance. Type I error for all statistical tests was set at 0.05 (IBM SPSS

Statistics v 28).

***In vitro* egg hatch assay (EHA)**

Eggs were extracted from fresh faecal samples from hookworm-positive dogs for use in

in vitro egg hatch assays (EHAs). Saturated sugar solution (specific gravity = 1.30), three

times the weight of individual faecal samples, was added to the faecal samples in a

shaker bottle. The bottle was manually agitated to form a slurry, which was then filtered

through a 200 µm sieve into a beaker to remove large debris. All of the filtered solution

was then transferred into 50 ml Falcon tubes, filling each tube up to the 40 ml, and then

10 ml of distilled water was slowly and carefully poured along the edge of the tube to

avoid mixing with the faecal solution. Each tube was then screw capped and centrifuged

at 2000 rpm (400×g) for 10 minutes. Following centrifugation, the 'fuzzy' layer (water-

sucrose interface) of each tube was removed with a pipette and pooled into a 200 mL

beaker. The solution was passed through a 75 µm sieve, retaining the filtrate, which was

further passed through a 20 µm sieve, which retained the hookworm eggs. This sieve was

then retrograde washed with 20 mL of distilled water into a glass Petri dish. The recovered

eggs were carefully poured into a two 15 mL centrifuge tubes, leaving the debris in the

Petri dish. The eggs were concentrated into 2 mL of water by centrifugation for 5 minutes

at 2000 rpm (400×g) and discarding the supernatant. The egg concentration was

estimated by thoroughly mixing the 2 mL egg solution and counting the eggs in three 20µl

aliquots. The egg numbers were adjusted to approximately 2 eggs per microlitre either

by dilution with distilled water or by concentration using centrifugation as described

above.

The egg hatch assay was performed using 96-well plates containing thiabendazole

embedded in agar, as described previously for assays with human hookworms (Kotze et

al., 2009). A stock solution of thiabendazole (5 mg/mL, Sigma-Aldrich Australia) was

prepared in dimethyl sulfoxide (DMSO) and serially diluted 2-fold in the same solvent.

Aliquots (2 µL) from a series of dilutions were added to 96-well microtiter plates, such

that each row of the plate comprised a double dilution, ranging from 0.003 µg/mL up to

1.359 µg/mL or 0.011 µg/mL to 5.437 µg/mL. The first two wells of each row were used as

control wells (received 2 µL of DMSO only). Molten agar (200 µL at 2% w/v) (Davis

Gelatine, Australia; powdered agar Grade J) was dispensed into each well of the plate

and allowed to set. Plates were placed into plastic press-seal bags stored at 4°C.

Prior to establishing the assays, the plates were allowed to warm in an incubator set to

28˚C. 30µL of concentrated hookworm egg solution (~2 eggs/µL) was pipetted into each

well. Plates were incubated for 48 hours at 28˚C, whereupon the assay was terminated

by addition of Lugol's iodine solution. The numbers of unhatched eggs and larvae in each

well were counted with the aid of a stereo microscope (Stemi 508, Carl Zeiss, Australia).

All assays were performed in triplicate for the sample isolated from each dog sample.

The number of larvae in each well was expressed as a percentage of the total numbers of

larvae and unhatched eggs in order to give the hatch rate in each well, and then this was

expressed relative to the mean hatch rate in the control wells. The data were then

analysed by non-linear regression using GraphPad Prism software (GraphPad Software

Inc. USA). This analysis yielded IC_{50} values representing the concentration of

thiabendazole that resulted in an egg hatch of 50% compared to control assays. The IC_{50}

values for the dogs examined in this study were compared to those reported previously

for drug-susceptible *A. caninum* isolates: 0.046 and 0.048 µg/mL (Jimenez Castro et al.,

2019) and 0.045 µg/mL (Diawara et al., 2013). An Australian fenbendazole susceptible

isolate of *A. caninum* ('Dougie', from February 2021) *s*erved as local positive control;

demonstrated faecal egg reduction of 100% 10 days post administration from initial 525

EPG.

The left-over eggs were suspended in distilled water in 1.5 ml microtubes, incubated for

72 hours at 28˚C and allowed to hatch. The hatched larvae were stored at -20˚C and

subjected to DNA isolation and follow up processes described earlier.

In vivo efficacy measurements

One dog (sample D6) was treated by the shelter facilities and faecal egg count monitored

throughout the treatment. The dog was treated according to its weight and product label

on day 0 with Drontal Allwormer (Vetoquinol, Australia; actives: febantel, praziquantel

and pyrantel embonate) and on day 21 with Credelio Plus (Elanco, Australia: milbemycin

oxime and lotilaner). In animals, febantel is a prodrug that is actively hydrolyzed to the

active metabolite fenbendazole. Egg counts were conducted within the period 10-21

days after Drontal Allwormer treatment to calculate faecal egg reduction percentage

(>95% reduction is considered effective product). Only a single sample was obtained at

day 8 after Credelio Plus because the dog was then rehomed.

Data availability

The EHA raw data, FEC reduction data and metadata for all samples are available at

LabArchives (https://dx.doi.org/10.25833/hmvy-jn79). Raw FastQ sequence data was

deposited at SRA NCBI BioProject: PRJNA1149426.

Results

Dominance of *A. caninum* across south-eastern Australia using metabarcoded ITS-

2 rDNA deep sequencing in dogs

We used 106 faecal samples that were considered hookworm positive. There were 20

samples from New Zealand from dogs. For 61 of the 106 samples the worm faecal egg

count (FEC) ranged from 20 to 4,920 (avg. 526) eggs per gram (EPG). The majority of the

Australian samples came from a commercial diagnostic laboratory (n=50) that were

collected over period of 13 months between March, 2023 and March, 2024.

There were 86 samples from south-eastern Australia, 85 samples from dogs and one

from a cat (TS42). Good quality DNA was obtained from 98% (84/86) yielding C_t<35 (11.8

to 31.6) at ITS-2 rDNA assay targeting nematode DNA. All 84 samples yielded

metabarcoded and deep sequenced ITS2 rDNA sequence data for nematode species

identification with the sequencing depth (n=84; avg. 21,323, 794 to 32,899) (Fig. 1A). Two

samples yielded low ITS-2 rDNA sequence depth, 794 (TS100) and 1,537 (TS95)

sequencing reads; removing those the remaining yielded >2,000 reads per sample (n=82;

avg. 21,815, range 2,736 to 32,899). The cat hookworm sample contained sequences

that matched *Ancylostoma tubaeforme* (TS42). The remaining dog samples (n=81)

contained sequences of *A. caninum* (n=77, 95%) and *U. stenocephala* (n=4, 5%). One

sample (TS35) had both *A. caninum* (96%) and *U. stenocephala* (4%) ITS-2 rDNA

sequences. The additional two samples with <2,000 ITS-2 rDNA sequences (TS95, TS100)

included only *A. caninum* sequences.

358 Widespread and high frequency of *A. caninum* canonical F167Y (TTC>TAC) isoform-

359 1 β-tubulin benzimidazole resistance mutation in Australia

For 81 Australian samples, the isoform-1 β-tubulin BZ167 amplicons were obtained with

an average sequencing depth for confirmed hookworm sequences between 81 to 31,265

(Fig. 1B). Excluding six samples (TS6, TS7, TS39, TS55, TS63, TS90; sequence depth 81 to

1,699 depth) with <2,000 sequences, the average sequencing depth was 11,092

sequences (n=75; 2,213 to 31,265). There were 72 samples with >2,000 isoform-1 β-

tubulin BZ167 amplicon sequences belonging to *A. caninum* (avg. depth 11,271) (Fig. 1B).

There were 35 (49%, 35/72) samples that harboured on average 10% (1% to 45%; 95%

confidence interval 7-13%) of the Q134H (CAA>CAT) resistant allele and 48 (67%, 48/72)

samples that harboured on average 41% (1% to 100%; 95% confidence interval 33-51%)

of the F167Y (T$\underline{T}$C>T$\underline{A}$C) benzimidazole resistant allele (Fig. 1C). There were 18 (25%,

18/72) and 14% (10/72) samples with >50% and >75% of the F167Y (T$\underline{T}$C>T$\underline{A}$C) (Fig. 1C).

There were 26 (36%, 26/72) samples that included both Q134H (CA$\underline{A}$>CA$\underline{T}$) and F167Y

(T$\underline{T}$C>T$\underline{A}$C) benzimidazole resistant alleles. The six samples with <2,000 sequences for

A. caninum had only the susceptible (coding for Q_{134}, F_{167}) alleles, except TS90 with 100%

of F167Y (T$\underline{T}$C>T$\underline{A}$C) mutation across its BZ167 sequences. Sequence analysis of

isoform-1 β-tubulin sequence (amino acids E_{198} and F_{200}) revealed no evidence of

benzimidazole resistance signatures.

There were four *U. stenocephala* and one (1/4) of the samples TS24 included F167Y

(T$\underline{T}$C>T$\underline{A}$C) in the isoform-1 β-tubulin mutation in 51% of the sequence reads. The

remaining isoform-1 β-tubulin of *U. stenocephala* matched benzimidazole susceptible

sequence residues. The cat TS42 sample with *A. tubaeformae* and its isoform-1 β-tubulin

included only the benzimidazole susceptible sequences (amino acids Q_{134}, F_{167}, E_{198} and

F_{200}).

For the New Zealand samples only 50% (10/20) of DNA yielded C_t<35 (20.2 to 33.6); nine

samples yielded ITS-2 rDNA sequence data to consider species identification (TS76-

TS84) with the sequencing depth (avg. 18,488; 1,118 to 27,624) (Fig. 1A). Four of the nine

samples contained sequence data that belonged to non-hookworm sequence including

ITS-2 rDNA sequencing matched to parasites of sheep and rabbits; three DNA samples

contained only ASVs that matched pseudoparasite DNA sequences. Subtracting the

pseudoparasite sequences to retain hookworm sequence data revealed presence of

uniquely *U. stenocephala* sequences, five samples with sequence depth for hookworms

from 8,027 to 27,624 and one sample (TS76) with only 118 sequences. Sequence

analysis of isoform-1 β-tubulin (amino acids Q_{134}, F_{167}, E_{198} and F_{200}) revealed no evidence

of benzimidazole resistance signatures in New Zealand samples.

High frequency of Q134H (CAA>CAT) or F167Y (TTC>TAC) mutation in *A. caninum* in

greyhounds

We then focused on 55 dog samples from two diagnostic labs (Vetnostics, VPDS) for

which we had *A. caninum* Q134H (CA$\underline{A}$>CA$\underline{T}$) and F167Y (T$\underline{T}$C>T$\underline{A}$C) mutation frequency.

There were 53 dogs with defined location to a postcode from New South Wales, Australia

(one dog from New South Wales did not have a postcode and the other dog was from

Queensland). Dogs came from 12 distinct New South Wales postcodes (Fig. 2).

There was no significant correlation between Q134H (CA<u>A</u>>CA<u>T</u>) or F167Y (T<u>T</u>C>T<u>A</u>C) and

EPG (correlation coefficient=-0.169, P=0.217; correlation coefficient=0.03, P=0.82).

Considering the association of presence or absence of either Q134H (CA<u>A</u>>CA<u>T</u>) or

F167Y (T<u>T</u>C>T<u>A</u>C) and EPG, there was no significant association for either SNP (Kruskal-

Walis Test: H=0.126, degrees of freedom=1, P=0.72; H=0.56, degrees of freedom=1,

P=0.45). We then considered the threshold of >40% for F167Y (T<u>T</u>C>T<u>A</u>C) mutation and

recalculated the association, but the rest remained not significant (Kruskal-Walis Test:

H=0.02, degrees of freedom=1, P=0.89). There was only a single sample with >40% of the

Q134H (CA<u>A</u>>CA<u>T</u>) mutation, therefore preventing calculation of these test statistics.

Considering the age of the dogs for which the age was available (n=41) there was no

significant correlation between age and EPG, Q134H (CA<u>A</u>>CA<u>T</u>) or F167Y (T<u>T</u>C>T<u>A</u>C)

(Spearman correlation, P>0.05).

The median resistance in Greyhounds for both the Q134H (CA<u>A</u>>CA<u>T</u>) and F167Y

(T<u>T</u>C>T<u>A</u>C) *A. caninum* mutation (0.035 and 0.090), respectively were significantly higher

than that for other breeds (median 0.000; Kruskal Wallis Test, P≤0.01) (Fig 2). There was

no significant difference between EPGs for the two breed categories (Kruskal-Wallis Test,

P=0.22).

Phenotypic evidence of benzimidazole resistance in *A. caninum* field sample with

>75% F167Y (T<u>T</u>C>T<u>A</u>C) mutation

We tested 13 *A. caninum* samples (D1, D2, D3, D5, D6, D8, D9, D10, D12, D15, D16, D17,

D18/20) with EHA that had their frequency of Q134H (CA<u>A</u>>CA<u>T</u>) and F167Y (T<u>T</u>C>T<u>A</u>C) *A.*

caninum SNP established (Fig. 1, Table 1). Dose response curves and IC_{50} formed two

groups, those that had high IC_{50}>0.15 µg/mL and those that had low IC_{50}<0.15 µg/mL of

thiabendazole (Fig. 3A). Three greyhounds (D1/D2: TS87/88, D3: TS89, D17: TS103) and

the crossbreed dog (D6: TS92, D18/20: TS104/106) were in the high IC_{50} group and they

all had >75% of F167Y (T$\underline{T}$C>T$\underline{A}$C) *A. caninum* mutation in the isoform-1 β-tubulin (Fig. 1,

Table 1). The IC_{50} of the remaining seven greyhound samples ranged from 0.02 to 0.06

µg/mL of thiabendazole and frequency of *A. caninum* F167Y (T$\underline{T}$C>T$\underline{A}$C) was <40% (Fig.1,

Table 1). For one dog (represented by two samples D6 and D18/20 in Fig. 1), we

established an *in vitro* EHA on eggs purified from dog's faeces on day 1 and day 9,

showing IC_{50} of 0.24 (0.19 to 0.30) µg/mL and 0.17 (95%CI 0.12 to 0.24) µg/mL of

thiabendazole, respectively. There was no significant difference between the response

curves for these two samples, demonstrated by single response curve fitting both

sample responses to thiabendazole (P=0.19; F=1.735, DFn=2, DFd=45). The

benzimidazole susceptible control IC_{50} was 0.11 (95%CI 0.10 to 0.12) µg/mL (Fig. 3), and

the *A. caninum* DNA was devoid of canonical F167Y (T$\underline{T}$C>T$\underline{A}$C) and Q134H (CA$\underline{A}$>CA$\underline{T}$)

isoform-1 β-tubulin mutations which confer benzimidazole resistance; the BZ167 and

BZ200 amplicons coded for amino acids Q_{134}, F_{167}, E_{198} and F_{200}. There was significant

correlation of IC_{50} for thiabendazole in EHA and F167Y (T$\underline{T}$C>T$\underline{A}$C) mutation in the

isoform-1 β-tubulin (Spearman's r=0.83 (95%CI=0.52 to 0.95, P-value < 0.001) (Fig. 3B).

In vivo evidence based on repeated FEC for one dog (D6 and D18/20 in Fig. 1) suggested

decreased (<95%) ability of febantel/pyrantel to reduce egg shedding post treatment.

This crossbreed dog (D6, D18/20) initially shed 540 EPG of *A. caninum* hookworms on

day 1 (day of treatment). The hookworm isolate carried 83% of F167Y (T$\underline{T}$C>T$\underline{A}$C)

mutation in the isoform-1 β-tubulin (D6: TS092; Fig. 1), and was treated with 'Drontal

Allwormer', but the faecal egg reduction only reached 88.8% (60 EPG) and 72.2% (195

EPG) on day 10 and day 14 post treatment, respectively. The same dog was re-treated

with macrocyclic lactone ('Credelio Plus') on day 20 (90 EPG) post original treatment that

resulted in 100% *A. caninum* egg reduction in faeces on day 8 post second treatment (0

EPG).

Discussion

This survey of dogs from south-east Australia infected with hookworms demonstrated a

high frequency of *A. caninum* with canonical mutations F167Y and Q134H in the isoform-

1 β-tubulin that are known to be associated with benzimidazole resistance. Both F167Y

and Q134H mutations have been conclusively confirmed to lead to benzimidazole

resistance using *Caenorhabditis elegans* as a surrogate (Kitchen et al., 2019; Venkatesan

et al., 2023). The extent of both mutations, 67% for F167Y and 49% for Q134H in *A.

caninum* tested in this study, exceeds even the recent USA report by Venkatesan et al.

(2023) that utilised the same methodology based on NGS amplicon metabarcoding. Our

samples from New South Wales, Australia correspond to hookworm-positive samples

tested by a commercial diagnostic laboratory (n=49) over a 13-month period. Our

sampling effort for New South Wales represents approximately 1,700 dog samples,

assuming a conservative *A. caninum* prevalence of 2.9% in temperate and subtropical

regions of Australia (Massetti et al., 2022).

Using our samples from New South Wales, Australia, we show that the isoform-1 β-

tubulin F167Y mutation is present in canine hookworms across the state and

predominantly in greyhounds that formed most of the samples. In North America, the

isoform-1 β-tubulin F167Y mutation was present in all geographic areas with detected

hookworms, including Canada (Leutenegger et al., 2024). Initially the F167Y

polymorphism was recognised in greyhounds; currently the highest isoform-1 β-tubulin

F167Y mutation prevalence was shown to be in poodles (29%), followed by Bernese

mountain dogs (25%), cocker spaniels (23%) and greyhounds (22%) that are infected with

A. caninum in North America (Jimenez Castro et al., 2019; Kitchen et al., 2019;

Leutenegger et al., 2024). Using faecal samples from 17,671,724 individual dogs in North

America, greyhounds were found to have a significantly higher risk of testing positive for

hookworms on faecal flotation (odds ratio = 15.3) (Burton et al., 2024). Additionally, the

interval to test negative was 71–72 days for presumably anthelminthic-treated

greyhounds, compared to just 1–2 days for other breeds (Burton et al., 2024). In this study

from Australia, in addition to greyhounds, one crossbreed dog from Queensland was

infected with *A. caninum* and had 83% of the isoform-1 β-tubulin F167Y mutation, along

with a reduced in vitro IC_{50} for thiabendazole.

The EHA tests for *in vitro* susceptibility to benzimidazoles thus phenotype including *A.*

caninum resistance to benzimidazole (Kotze et al., 2005; Kotze et al., 2009; Diawara et

al., 2013). Importantly, we demonstrated correlation of EHA and the isoform-1 β-tubulin

F167Y mutation, and those samples with elevated EHA (IC_{50}) all showed frequency of

F167Y mutation >75%. Known *in vitro* resistant *A. caninum* with IC_{50} of 8.76 μM (=1.76

μg/mL, KGR) and 40.08 μM (=8.07 μg/mL, BCR) for thiabendazole had 100% and 76% of

the F167Y mutation (McKean et al., 2024). Our resistant *A. caninum* had IC_{50} from 0.17 to

0.43 μg/mL of thiabendazole and are closer to the range (0.55 to 0.67 μg/mL) detected

for resistant *A. caninum* by Jimenez Castro et al. (2019). The *in vitro* resistance was

further confirmed by FEC reduction in one of the Australian dogs *in vivo*. There was no

significant correlation between F167Y (TTC>TAC) frequency and IC_{50} for thiabendazole in

a similar small study based on 15 *A. caninum* samples (Jimenez Castro et al., 2021).

The detection of the isoform-1 β-tubulin F167Y mutation in *U. stenocephala*, the

Northern hookworm that commonly infects dogs in cooler climates, is unexpected, as

drug resistance has not previously been reported for *U. stenocephala* (Leutenegger et al.,

2023; Venkatesan et al., 2023; Stocker et al., 2024). In our samples from Australia, we

detected *U. stenocephala* F167Y mutation in one out of four dog samples. More

frequently in Australia, *U. stenocephala* is a parasite of red foxes which likely serves as

the reservoir (Ryan, 1976). In a study of 930 foxes from New South Wales, Australia, the

most prevalent nematodes were *Toxocara canis* (35.2%) and *U. stenocephala* (30.6%),

while *A. caninum* was detected in only 70 (7.5%) foxes (Ryan, 1976). Although *U.*

stenocephala is often considered relatively non-pathogenic, it has been acknowledged

as a cause of subacute enteritis and hypoproteinaemia in dogs (Gibbs, 1958; Miller,

1968). The significance of the F167Y mutation in *U. stenocephala* in conferring

benzimidazole resistance in the presence of a large *refugium* in red foxes will require

further confirmation from *in vitro* and *in vivo* drug response studies.

The absence of resistant SNPs that would code for E_{198} and F_{200} in the isoform-1 β-tubulin

of *A. caninum* in Australia agrees with reports from the USA (Venkatesan et al., 2023). The

F200Y SNP has been documented once in *A. caninum* dog samples from Brazil, but the

in vitro or *in vivo* phenotype was not measured (Furtado et al., 2014). The F200Y was

detected in human *N. americanus* from Haiti with frequency of 20-90%, and in an *in vitro*

EHA assay these hookworm isolates were less sensitive to thiabendazole (IC_{50} = 0.12 to

0.26 μg/μL) (Diawara et al., 2013). Molecular dynamics simulations using isotype-1 β-

tubulin showed that *A. caninum* and *Ancylostoma ceylanicum* both shared the same

drug binding profile, while *A. duodenale* and *N. americanus* both had different binding

profiles with no overlap except for the E_{198} interaction that is shared with all nematodes

(Jones et al., 2022).

With growing evidence of anthelmintic resistance in *A. caninum*, the question arises

whether the fitness and ability of resistant strains to spread across the dog population

differ from non-resistant strains. Recently, a strain of triple-resistant *A. caninum* (BCR)

was shown to exhibit reduced 'larval activation', an initial step in the transformation of

infective non-feeding larvae into migrating and feeding larvae in a permissive host

(McKean et al., 2024). Larval activation *in vitro* is initiated when third-stage infective

larvae (iL3) of *A. caninum* are incubated with a host-mimicking signal, such as dog

serum, to resume feeding over a period of 24 hours (Hawdon and Schad, 1990; Hotez et

al., 1993). The consequences of the 40-50% reduced initial activation of the *A. caninum*

triple-resistant strain (BCR) reported by McKean et al. (2024) are not yet known but

suggest a possible reduced ability to establish infection. However, we did not find any

association between the SNPs and FEC in the dogs we studied. Our study does not

permit the examination of relationships between FECs and other variables (e.g., worm

fitness) because the infection histories (such as exposure to larvae and drug treatment

history) of the dogs are completely unknown. Our samples were neither tested for larval

activation nor for the presence of pyrantel and macrocyclic lactone resistance, so

reduced 'larval activation' may either be restricted to the triple-resistant strains, or it may

not lead to lower infection rate and changes in egg output.

Conclusion

The molecular tools employed in the present study enable agile moderate to high throughput tracking of anthelmintic resistance mutations. This offers advantages in terms of speed and cost compared to phenotypic *in vivo* and *in vitro* measurements of sensitivity to anthelmintic drugs. We identified the hookworm species and simultaneously discovered widely distributed isoform-1 β-tubulin SNPs across Eastern Australia. Hookworms, whipworm and roundworms in dogs and cats are traditionally managed by sustained routine administration of anthelminthics. The recent discovery of MADR in canine hookworm (*A. caninum*) has raised the concerns of the veterinary profession (von Samson-Himmelstjerna et al., 2021; Marsh and Lakritz, 2023). Adopting a risk assessment-based approach to limit unnecessary use of parasiticides has the potential to mitigate the risk of resistance selection and spread, and should be an important consideration in responsible parasite control (Selzer and Epe, 2021; Bagster and Elsheikha, 2022).

Acknowledgements

Funding was provided by the Margo Roslyn Flood Bequest (Sydney School of Veterinary Science, The University of Sydney). Funding for University of Queensland part of research was provided by The Canine Research Foundation and Dogs Victoria.

558 **Table 1:** Summary of *Ancylostoma caninum* egg hatch assay with frequency of Q134H and F167Y polymorphisms in isoform-1 β-tubulin.

Sample ID	Breed	EPG	IC50 µg/mL *	IC_{50} CI95% µg/mL *	Q134H[&]	F167Y[&]
High IC_{50}>0.15 µg/mL of thiabendazole						
D3:TS89	Greyhound	1,275	0.429	0.344 to 0.534	0%	89.91%
D1:TS87[^]	Greyhound	1,050	0.298	0.216 to 0.412	0%	85.47%
D2:TS88[^]	Greyhound	645	0.262	0.237 to 0.289	0%	88.80%
D6:TS92[^^]	Crossbreed	540	0.239	0.201 to 0.285	0%	83.11%
D18/D20:TS104/106[^^]	Crossbreed	225	0.170	0.121 to 0.239	0%/0%	100%/77.54%
D17:TS103	Greyhound	5,535	0.194	0.150 to 0.250	2.55%	81.23%
Low IC_{50}<0.15 µg/mL of thiabendazole						
D5:TS91	Greyhound	14,910	0.056	0.048 to 0.066[#]	0%	10.69%

D9:TS95	Greyhound	930	0.050	0.043 to 0.059[#]	18.27%	33.51%
D10:TS96	Greyhound	4,080	0.043	0.040 to 0.046[#]	0%	22.67%
D8:TS94	Greyhound	390	0.039	0.031 to 0.048[#]	0%	29.26%
D12:TS98	Greyhound	3,930	0.037	0.035 to 0.039[#]	0%	0%
D15:TS101	Greyhound	720	0.036	0.031 to 0.041[#]	0%	6.56%
D16:TS102	Greyhound	885	0.021	0.013 to 0.033[#]	7.43%	22.53%

[*] thiabendazole concentration in µg/mL

[#] drug-susceptible *A. caninum* - 0.046 and 0.048 µg/mL (Jimenez Castro et al., 2019) and 0.045 µg/mL (Diawara et al., 2013).

[&] percentage of canonical benzimidazole resistance codons on the isoform-1 β-tubulin

[^,^^] same dog

Notes: EPG – *A. caninum* eggs per gram; IC_{50} - 50% inhibition concentration; CI95% - 95% confidence interval.

Figure 1 | Molecular profiling of hookworm species using ITS-2 rDNA and benzimidazole susceptibility using isoform-1 β-tubulin.

(A) Sampled with hookworm-like stages that were profiled using metabarcoded deep Illumina amplicon sequencing at ITS-2 rDNA. All but one sample came from dogs. Th single, cat single sample (TS042) is on the very right. The broad scale of the samples is indicated above the bar chart. Amplicon sequence variants (ASVs) were pooled according to species of nematode they belonged using reference sequence alignment. Proportion of each species (*Ancylostoma caninum*, *A. tubaeforme, Uncinaria stenocephala*) is colour coded in the stacked bar chart; ASVs that belonged to unrelated nematodes are depicted as 'pseudoparasites' and represented trichostrongylid species from sheep and rabbits. The bar chart underneath represents sequence depth for each sample, and samples with <2,000 sequence reads are depicted with a star. Samples are vertically aligned across the bar charts. (B) Dog samples profiled using targeting BZ167 amplicon of the isoform-1 β-tubulin, specifically region covering amino acids Q_{134} and F_{167} conferring benzimidazole resistance. Only ASVs of hookworm are used and the sequencing depth is depicted underneath. Samples that did not amplify hookworms are indicated by square and samples with <2,000 sequence reads are depicted with a star. (C) Bar chart for Q_{134} and F_{167} isoform-1 β-tubulin of *A. caninum* where proportion of an alternative residue coding for amino acid that confers benzimidazole resistance are show. Samples with no bar chart are those that did not amplify *A. caninum* and so there are no *A. caninum* ASVs.

Figure 2 | Distribution of *Ancylostoma caninum* and benzimidazole resistant isoform-1 β-tubulin single nucleotide polymorphism.

Proportional symbol maps of (A) the number of *A. caninum* positive samples, and (B)

number of Q134H (CA$\underline{A}$>CA$\underline{T}$) and (C) number of F167Y (T$\underline{T}$C>T$\underline{A}$C) isoform-1 β-tubulin of

A. caninum positive samples. Samples were summed per postcode in New SouthWales,

Australia. Samples are joined to postcodes of Australia shapefile (Geocentric Datum of

Australia, GDA 1994) in ArcGIS® Pro.

Figure 3 | *Ancylostoma caninum* IC$_{50}$ for thiabendazole in egg hatch assays.

(A) Correlation IC$_{50}$ values for *A. caninum* samples for thiabendazole agar-based egg

hatch assay (EHA) from Queensland, Australia with percentage of F167Y (T$\underline{T}$C>T$\underline{A}$C)

isoform-1 β-tubulin of *A. caninum* present in the sample. Simple linear regression is

depicted as solid line with 95% confidence interval depicted as dotted line. (B) Histogram

of IC$_{50}$ values for *A. caninum* samples from Queensland, Australia.   A susceptible *A.*

caninum samples, that was confirmed to have 100% faecal egg count (FEC) reduction is

colour coded (black). The samples with >0.15 µg/mL thiabendazole IC$_{50}$ values have their

identifiers above the bars.

605   **References**
606
Bagster, A., Elsheikha, H., 2022. Perception of UK companion animal veterinarians on risk
assessment based parasite control. Vet Parasitol Reg Stud Reports 34, 100774. doi:
10.1016/j.vprsr.2022.100774
Bolt, B.J., Rodgers, F.H., Shafie, M., Kersey, P.J., Berriman, M., Howe, K.L., 2018. Using
WormBase ParaSite: An integrated platform for exploring helminth genomic data. Methods Mol
Biol 1757, 471-491. doi: 10.1007/978-1-4939-7737-6_15
Burton, K.W., Hegarty, E., Couto, C.G., 2024. Retrospective analysis of canine fecal flotation and
coproantigen immunoassay hookworm positive results in Greyhounds and other dog breeds. Vet
Parasitol Reg Stud Reports 51, 101026. doi: 10.1016/j.vprsr.2024.101026

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., Holmes, S.P., 2016. DADA2:
High-resolution sample inference from Illumina amplicon data. Nat Methods 13, 581-583. doi:
10.1038/nmeth.3869

Dale, A., Xu, G., Kopp, S.R., Jones, M.K., Kotze, A.C., Abdullah, S., 2024. Pyrantel resistance in
canine hookworms in Queensland, Australia. Vet Parasitol Reg Stud Reports 48, 100985. doi:
10.1016/j.vprsr.2024.100985

Diawara, A., Schwenkenbecher, J.M., Kaplan, R.M., Prichard, R.K., 2013. Molecular and
biological diagnostic tests for monitoring benzimidazole resistance in human soil-transmitted
helminths. Am J Trop Med Hyg 88, 1052-1061. doi: 10.4269/ajtmh.12-0484

Furtado, L.F., Bello, A.C., dos Santos, H.A., Carvalho, M.R., Rabelo, E.M., 2014. First
identification of the F200Y SNP in the beta-tubulin gene linked to benzimidazole resistance in
Ancylostoma caninum. Vet Parasitol 206, 313-316. doi: 10.1016/j.vetpar.2014.10.021

Gasser, R.B., Chilton, N.B., Hoste, H., Beveridge, I., 1993. Rapid sequencing of rDNA from single
worms and eggs of parasitic helminths. Nucleic Acids Res 21, 2525-2526. doi:
10.1093/nar/21.10.2525

Gibbs, H.C., 1958. On the gross and microscopic lesions produced by the adults and larvae of
Dochmoides stenocephala (Raillet, 1884) in the dog. Can J Comp Med Vet Sci 22, 382-385. doi:

Harris, T.W., Arnaboldi, V., Cain, S., Chan, J., Chen, W.J., Cho, J., Davis, P., Gao, S., Grove, C.A.,
Kishore, R., Lee, R.Y.N., Muller, H.M., Nakamura, C., Nuin, P., Paulini, M., Raciti, D., Rodgers,
F.H., Russell, M., Schindelman, G., Auken, K.V., Wang, Q., Williams, G., Wright, A.J., Yook, K.,
Howe, K.L., Schedl, T., Stein, L., Sternberg, P.W., 2020. WormBase: a modern Model Organism
Information Resource. Nucleic Acids Res 48, D762-D767. doi: 10.1093/nar/gkz920

Hawdon, J.M., Schad, G.A., 1990. Serum-stimulated feeding in vitro by third-stage infective
larvae of the canine hookworm *Ancylostoma caninum*. J Parasitol 76, 394-398. doi:
10.2307/3282673

Hotez, P., Hawdon, J., Schad, G.A., 1993. Hookworm larval infectivity, arrest and
amphiparatenesis: the *Caenorhabditis elegans* Daf-c paradigm. Parasitol Today 9, 23-26. doi:
10.1016/0169-4758(93)90159-d

Howe, K.L., Bolt, B.J., Shafie, M., Kersey, P., Berriman, M., 2017. WormBase ParaSite - a
comprehensive resource for helminth genomics. Mol Biochem Parasitol 215, 2-10. doi:
10.1016/j.molbiopara.2016.11.005

Jimenez Castro, P.D., Howell, S.B., Schaefer, J.J., Avramenko, R.W., Gilleard, J.S., Kaplan, R.M.,
2019. Multiple drug resistance in the canine hookworm *Ancylostoma caninum*: an emerging
threat? Parasit Vectors 12, 576. doi: 10.1186/s13071-019-3828-6

Jimenez Castro, P.D., Venkatesan, A., Redman, E., Chen, R., Malatesta, A., Huff, H., Zuluaga
Salazar, D.A., Avramenko, R., Gilleard, J.S., Kaplan, R.M., 2021. Multiple drug resistance in
hookworms infecting greyhound dogs in the USA. Int J Parasitol Drugs Drug Resist 17, 107-117.
doi: 10.1016/j.ijpddr.2021.08.005

Jones, B.P., van Vliet, A.H.M., LaCourse, E.J., Betson, M., 2022. *In silico* docking of nematode β-
tubulins with benzimidazoles points to gene expression and orthologue variation as factors in
anthelmintic resistance. Frontiers in Tropical Diseases 3, 898814. doi: 10.3389/fitd.2022.898814

Kitchen, S., Ratnappan, R., Han, S., Leasure, C., Grill, E., Iqbal, Z., Granger, O., O'Halloran, D.M.,
Hawdon, J.M., 2019. Isolation and characterization of a naturally occurring multidrug-resistant
strain of the canine hookworm, *Ancylostoma caninum*. Int J Parasitol 49, 397-406. doi:
10.1016/j.ijpara.2018.12.004

Kopp, S.R., Kotze, A.C., McCarthy, J.S., Coleman, G.T., 2007. High-level pyrantel resistance in
the hookworm *Ancylostoma caninum*. Vet Parasitol 143, 299-304. doi:
10.1016/j.vetpar.2006.08.036

Kotze, A.C., Coleman, G.T., Mai, A., McCarthy, J.S., 2005. Field evaluation of anthelmintic drug
sensitivity using in vitro egg hatch and larval motility assays with *Necator americanus* recovered
from human clinical isolates. Int J Parasitol 35, 445-453. doi: 10.1016/j.ijpara.2004.12.007

Kotze, A.C., Lowe, A., O'Grady, J., Kopp, S.R., Behnke, J.M., 2009. Dose-response assay
templates for in vitro assessment of resistance to benzimidazole and nicotinic acetylcholine
receptor agonist drugs in human hookworms. Am J Trop Med Hyg 81, 163-170. doi:
10.4269/ajtmh.2009.81.163

Kwa, M.S., Kooyman, F.N., Boersema, J.H., Roos, M.H., 1993. Effect of selection for
benzimidazole resistance in *Haemonchus contortus* on beta-tubulin isotype 1 and isotype 2
genes. Biochem Biophys Res Commun 191, 413-419. doi: 10.1006/bbrc.1993.1233

Leutenegger, C.M., Evason, M.D., Willcox, J.L., Rochani, H., Richmond, H.L., Meeks, C., Lozoya,
C.E., Tereski, J., Loo, S., Mitchell, K., Andrews, J., Savard, C., 2024. Benzimidazole F167Y
polymorphism in the canine hookworm, *Ancylostoma caninum*: Widespread geographic,
seasonal, age, and breed distribution in United States and Canada dogs. Int J Parasitol Drugs
Drug Resist 24, 100520. doi: 10.1016/j.ijpddr.2024.100520

Leutenegger, C.M., Lozoya, C.E., Tereski, J., Savard, C., Ogeer, J., Lallier, R., 2023. Emergence of
Ancylostoma caninum parasites with the benzimidazole resistance F167Y polymorphism in the
US dog population. Int J Parasitol Drugs Drug Resist 21, 131-140. doi:
10.1016/j.ijpddr.2023.01.001

Marsh, A.E., Lakritz, J., 2023. Reflecting on the past and fast forwarding to present day
anthelmintic resistant *Ancylostoma caninum*-A critical issue we neglected to forecast. Int J
Parasitol Drugs Drug Resist 22, 36-43. doi: 10.1016/j.ijpddr.2023.04.003

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet.journal 17, 10-12. doi: 10.14806/ej.17.1.200

Massetti, L., Wiethoelter, A., McDonagh, P., Rae, L., Marwedel, L., Beugnet, F., Colella, V., Traub,
R.J., 2022. Faecal prevalence, distribution and risk factors associated with canine soil-
transmitted helminths contaminating urban parks across Australia. Int J Parasitol 52, 637-646.
doi: 10.1016/j.ijpara.2022.08.001

McKean, E.L., Grill, E., Choi, Y.J., Mitreva, M., O'Halloran, D.M., Hawdon, J.M., 2024. Altered larval
activation response associated with multidrug resistance in the canine hookworm *Ancylostoma*
caninum. Parasitology 151, 271-281. doi: 10.1017/S0031182023001385

Miller, T.A., 1968. Pathogenesis and immunity in hookworm infection. Trans R Soc Trop Med Hyg
62, 473-489. doi: 10.1016/0035-9203(68)90130-2

R Core Team, 2022. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Ryan, G.E., 1976. Helminth parasites of the fox (*Vulpes vulpes*) in New South Wales. Aust Vet J
52, 126-131. doi: 10.1111/j.1751-0813.1976.tb05445.x

Schulz, J.D., Moser, W., Hurlimann, E., Keiser, J., 2018. Preventive Chemotherapy in the Fight
against Soil-Transmitted Helminthiasis: Achievements and Limitations. Trends Parasitol 34, 590-
602. doi: 10.1016/j.pt.2018.04.008

Schwenkenbecher, J.M., Albonico, M., Bickle, Q., Kaplan, R.M., 2007. Characterization of beta-
tubulin genes in hookworms and investigation of resistance-associated mutations using real-
time PCR. Mol Biochem Parasitol 156, 167-174. doi: 10.1016/j.molbiopara.2007.07.019

Selzer, P.M., Epe, C., 2021. Antiparasitics in Animal Health: Quo Vadis? Trends Parasitol 37, 77-
89. doi: 10.1016/j.pt.2020.09.004

Stocker, T., Scott, I., Šlapeta, J., 2023. Unambiguous identification of *Ancylostoma caninum* and
Uncinaria stenocephala in Australian and New Zealand dogs from faecal samples. Aust Vet J 101,
373-376. doi: 10.1111/avj.13272

Stocker, T., Ward, M.P., Šlapeta, J., 2024. Nationwide USA re-analysis of amplicon
metabarcoding targeting β-tubulin isoform-1 reveals absence of benzimidazole resistant SNPs
in *Ancylostoma braziliense*, *Ancylostoma tubaeforme* and *Uncinaria stenocephala*. Vet Parasitol
328, 110118. doi: 10.1016/j.vetpar.2024.110118

Venkatesan, A., Jimenez Castro, P.D., Morosetti, A., Horvath, H., Chen, R., Redman, E., Dunn, K.,
Collins, J.B., Fraser, J.S., Andersen, E.C., Kaplan, R.M., Gilleard, J.S., 2023. Molecular evidence

of widespread benzimidazole drug resistance in *Ancylostoma caninum* from domestic dogs
throughout the USA and discovery of a novel beta-tubulin benzimidazole resistance mutation.
PLoS Pathog 19, e1011146. doi: 10.1371/journal.ppat.1011146
Vercruysse, J., Levecke, B., Prichard, R., 2012. Human soil-transmitted helminths: implications
of mass drug administration. Curr Opin Infect Dis 25, 703-708. doi:
10.1097/QCO.0b013e328358993a
von Samson-Himmelstjerna, G., Blackhall, W.J., McCarthy, J.S., Skuce, P.J., 2007. Single
nucleotide polymorphism (SNP) markers for benzimidazole resistance in veterinary nematodes.
Parasitology 134, 1077-1086. doi: 10.1017/S0031182007000054
von Samson-Himmelstjerna, G., Thompson, R.A., Krucken, J., Grant, W., Bowman, D.D.,
Schnyder, M., Deplazes, P., 2021. Spread of anthelmintic resistance in intestinal helminths of
dogs and cats is currently less pronounced than in ruminants and horses - Yet it is of major
concern. Int J Parasitol Drugs Drug Resist 17, 36-45. doi: 10.1016/j.ijpddr.2021.07.003
WHO, 2017. Guideline: Preventive chemotherapy to control soil-transmitted helminth infections
in at-risk population groups Geneva: World Health Organization Geneva.