Deep Generative Optimization of mRNA Codon Sequences for Enhanced mRNA Translation and Therapeutic Efficacy

Yupeng Li[1,#], Fan Wang[1,#], Jiaqi Yang[1], Zirong Han[2,3], Linfeng Chen[1], Wenbing Jiang[1], Hao Zhou[1], Tong Li[1], Zehua Tang[1], Jianxiang Deng[1], Xin He[1], Gaofeng Zha[4], Jiekai Hu[5], Yong Hu[5], Linping Wu[6], Changyou Zhan[7], Caijun Sun[2,3,8,9], Yao He[1,*], Zhi Xie[1,*]

[1]State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-Sen University, Guangdong Provincial Key Laboratory of Ophthalmology and Visual Science, Guangzhou 510060, China.

[2]School of Public Health (Shenzhen), Sun Yat-Sen University, Shenzhen, 518107, China.

[3]Shenzhen Key Laboratory of Pathogenic Microbes and Biosafety, Shenzhen Campus of Sun Yat-Sen University, Shenzhen, 518107, China.

[4]Scientific Research Center, The Seventh Affiliated Hospital. Sun Yat-Sen University, Shenzhen, 518107, China.

[5]ENO Bio mRNA Innovation Institute, Rhegen Biotechnology Co., Ltd. Shenzhen, China

[6]Center for Chemical Biology and Drug Discovery, China-New Zealand Joint Laboratory of Biomedicine and Health, Guangdong Provincial Key Laboratory of Biocomputing, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, 510530, China.

[7]Department of Pharmacology, School of Basic Medical Sciences, Fudan University, Shanghai, 200032, China.

[8]Key Laboratory of Tropical Disease Control (Sun Yat-Sen University), Ministry of Education; Guangzhou, 514400, China.

[9]State Key Laboratory of Anti-Infective Drug Discovery and Development, School of Pharmaceutical Sciences, Sun Yat-Sen University, Guangzhou, 510006, China.

[#]Equally contributed.

[*]Corresponding author. Email: xiezhi@gmail.com (Z.X.), scheyao@hotmail.com (Y. He)

ABSTRACT

Messenger RNA (mRNA) therapeutics show immense promise, but their efficacy is limited by suboptimal protein expression. Here, we present RiboDecode, a deep learning framework that generates mRNA codon sequences for enhanced mRNA translation. RiboDecode introduces several advances, including direct learning from large-scale ribosome profiling data and generative exploration of a large sequence space. *In silico* analysis demonstrates RiboDecode's robust predictive accuracy for unseen genes and cellular environments. *In vitro* experiments showed substantial improvements in protein expression, significantly outperforming past methods. In addition, RiboDecode enables mRNA design with consideration of cellular context and demonstrates robust performance across different mRNA formats, including $m^1\Psi$-modified and circular mRNAs, an important feature for mRNA therapeutics. *In vivo* mouse studies showed that optimized influenza hemagglutinin mRNAs induce ten times stronger neutralizing antibody responses against influenza virus compared to the unoptimized sequence. In an optic nerve crush model, optimized nerve growth factor mRNAs achieve equivalent neuroprotection of retinal ganglion cells at one-fifth the dose of the unoptimized sequence. Collectively, RiboDecode represents a paradigm shift from rule-based to a data-driven, context-aware approach for mRNA therapeutic applications, enabling the development of more potent and dose-efficient treatments.

INTRODUCTION

Messenger RNA (mRNA) therapy has emerged as a promising approach for treating diseases. This innovative therapeutic strategy harnesses the cell's protein synthesis machinery to produce desired proteins encoded by the delivered mRNA[1–3], leading to the application of mRNA therapies in various fields, such as vaccine development and protein replacement therapy[4]. The successful development and deployment of mRNA vaccines during the COVID-19 pandemic have further highlighted the transformative potential of this technology[5].

Despite the remarkable progress in mRNA vaccines, achieving efficient and consistent protein translation from delivered mRNA molecules remains a key challenge, particularly critical for protein replacement therapy where sustained, precise, and often higher levels of protein expression are required in specific cellular contexts. However, the biological instability of mRNA and the complex regulatory mechanisms governing mRNA translation in cells can lead to suboptimal protein expression[6–8]. Therefore, improving the expression of mRNA is a key challenge for enhancing the therapeutic efficacy and reducing the required dose of mRNA-based treatments.

An amino acid can be encoded by multiple synonymous codons, ranging from one to six codons per amino acid. Codon optimization is a strategy to improve protein expression by changing the synonymous codon of an mRNA molecule while maintaining the encoded amino acid sequence. The choice of synonymous codons can largely impact the efficiency of mRNA translation and the stability of the mRNA molecule[6,7]. For example, it has been shown that optimal codon usage can enhance ribosome engagement and increase translation elongation rates, ultimately leading to higher protein production[8]. Additionally, codon choice can influence mRNA structure. Previous studies have demonstrated that mRNA structure critically influences its stability *in vivo*[9], in solution[10], and translation[11]. Therefore, codon optimization is a

critical step in the design of mRNA-based therapies to achieve maximal protein

production, leading to better therapeutic efficacy.

Computational tools have been developed for codon optimization, most of which

were designed for DNA, employing various strategies to select optimal codons. Past

methods rely on codon usage bias derived from highly expressed genes in a given

species, such as codon adaptation index (CAI)[12]. These methods aim to mimic the

codon usage patterns of efficiently translated endogenous mRNAs. More recently,

LinearDesign has been developed for mRNA optimization, aiming to jointly optimize

translation and mRNA stability by increasing CAI and reducing minimum free energy

(MFE)[6], which is a computational metric for evaluating mRNA secondary structure.

LinearDesign uses a linear programming approach to explore a wider space of

sequence variants compared to previous methods and showed superior performance

over the previous codon optimization methods. Additionally, other indices have been

used to guide sequence optimization. For instance, higher GC content (GC%) has

been associated with enhanced gene expression[13].

Despite the development of the previous methods, several limitations hinder their

effectiveness in consistently improving the protein expression of mRNA molecules.

Firstly, the existing methods primarily rely on predefined sequence features, such as

CAI, to guide codon selection. However, these metrics often fail to correlate with the

experimentally measured protein expression levels[14,15], indicating that they do not

accurately capture the complex factors governing mRNA translation. Secondly, the

existing methods do not adequately account for the activity of translational regulators

that influence mRNA translation, such as translation factors and RNA-binding

proteins[16,17]. This lack of context-aware optimization may reduce the effectiveness of

the optimized mRNA sequences in specific cellular environments. Furthermore, the

existing methods explore a limited space of codon sequences due to computational

constraints and the reliance on predefined rules. This restricted search space may

prevent the discovery of novel and highly optimized sequences that could potentially

yield significant improvements in protein expression.

Deep learning has achieved remarkable success in tasks such as image

recognition, natural language processing, and protein structure prediction, where it

has outperformed conventional algorithms by learning complex patterns and

relationships from vast amounts of data[18,19]. In the context of mRNA codon

optimization, a deep learning approach may enable the model to capture the complex

interplay between codon usage and cellular context, without relying on predefined

rules. Moreover, deep learning models can explore a vast sequence space and discover

novel patterns that may not be apparent to human experts or accessible through

traditional optimization methods[20]. This ability has been exemplified in the field of

protein engineering, where deep learning has been used to design novel protein

sequences with improved stability, binding affinity, and catalytic activity[21–23]. Recent

advances in codon optimization research have seen the emergence of deep learning-

based algorithms, particularly large language models (LLMs) trained on cross-species

nucleotide sequences. These models have been implemented for predictive modeling

of mRNA translation efficiency and degradation kinetics[24,25]. Nevertheless, there

persists an urgent demand for developing a rigorously validated optimization

framework to improve mRNA-encoded protein expression specifically tailored for

therapeutic applications.

Massive parallel reporter assays (MPRA) are commonly used to study the effects

of regulatory sequences on gene expression[26]. However, it is not suitable for

optimizing coding sequences due to the short sequence limitation, which is generally

less than 300 base pairs, for high-throughput DNA synthesis. Additionally, MPRA

experiments often rely on artificial reporter constructs and may not fully recapitulate

the complex regulatory landscape of endogenous mRNA molecules. Ribosome

profiling sequencing (Ribo-seq) is a powerful experimental technique that provides a

snapshot of actively translating ribosomes on mRNA molecules[27,28], where the

translation level of an mRNA can be derived from the reads per kilobase per million

(RPKM) of Ribo-seq. Recent studies have leveraged Ribo-seq to develop translation-

focused deep learning models[29–31]. For instance, RiboNN predicts mRNA translation

efficiency in mammalian cells by integrating mRNA sequences with ribosome

profiling data, revealing translation-stability regulatory mechanisms[32]. However, the

field critically requires a rational design framework that translates data-derived

translational signatures into codon optimization strategies, enabling high-throughput

exploration of sequence space and generation of optimized mRNA constructs for

therapeutic development.

In this study, we present RiboDecode, a deep learning model for mRNA codon

optimization that enhances mRNA translation by directly learning complex

relationship of mRNA codon sequences to their translation level from large-scale

Ribo-seq data. Our prediction model demonstrated robust performance, while analysis

of RiboDecode's optimization strategies revealed a complex interplay between

sequence characteristics and translation. *In vitro* experiments showed significantly

increased in mRNA translation and protein expression, outperforming past methods.

RiboDecode also considered cellular context, and maintained robust performance

across unmodified, $m^1\Psi$-modified, and circular mRNA formats. *In vivo*, optimized

influenza virus hemagglutinin (HA) mRNA induced approximately ten times stronger

neutralizing antibody responses in mice, while optimized nerve growth factor (NGF)

mRNA achieved equivalent neuroprotection of retinal ganglion cells at one-fifth the

dose in an optic nerve crush mouse model. This data-driven approach to codon

optimization advances our understanding of mRNA translation and facilitates the

development of more effective mRNA therapeutics.

RESULTS

RiboDecode is a Deep Learning Framework for mRNA Codon Optimization

RiboDecode is a deep learning-based framework for optimizing mRNA codon sequences. It integrates three components: a translation prediction model, an MFE prediction model, and a codon optimizer that explores and optimizes codon choices guided by the prediction models (Figure 1a).

The translation prediction model estimates the translation level of a given codon sequence by learning the translational expression of diverse mRNA sequences from Ribo-seq experiments (Figures 1b and S1, Methods). In contrast to previous tools that rely on optimizing predefined features such as CAI, our deep learning model automatically extracts relevant features by training on 320 paired Ribo-seq and RNA sequencing (RNA-seq) datasets from 24 different human tissues and cell lines, encompassing translation measurements of over 10,000 mRNAs per dataset (Supplementary file 1 and Methods)[33,34]. In addition, the model incorporates not only codon sequences but also mRNA abundances and cellular context that is presented by gene expression profiles from RNA-seq (Methods). This approach enables the prediction of mRNA translation by jointly considering these important factors influencing translation.

To address mRNA stability, we developed an MFE prediction model. Current MFE prediction tools, such as RNAfold[35] and Linearfold[36], use dynamic programming. However, these methods are non-differentiable and thus incompatible with our codon optimizer described below. Our MFE model employs a deep neural network architecture and undergoes an iterative optimization process, to simultaneously improve its predictive capability and optimize sequences for lower MFE values (Figure S2, Methods).

The codon optimizer of RiboDecode begins with the original codon sequence of a given protein (Figure 1c). The prediction models then predict a fitness score for this

sequence. Using a gradient ascent optimization approach based on activation

maximization (AM)[37], the optimizer adjusts the codon distribution to maximize the

fitness score (Figure S3). A synonymous codon regularizer ensures that only

synonymous codons encoding the same amino acids as the original sequence are

considered, preserving the protein's amino acid sequence. Through iterative cycles of

sequence generation, prediction, and optimization, the system produces codon

sequences with improved properties. RiboDecode can optimize mRNA translation,

stability or both, by interfacing with both the translation and MFE models. This uses a

parameter, w: $w=0$ optimizes translation only, $w=1$ optimizes MFE only, and a value

$0<w<1$ jointly optimizes both (Methods).

By combining data-driven predictions with high-throughput sequence generation,

RiboDecode overcomes limitations of conventional heuristic approaches. It enables

the exploration of a vast mRNA codon space, potentially uncovering optimized

sequences.

Evaluation of Translation Prediction Model

We first evaluated the RiboDecode's performance and generalizability using three

cross-validation datasets: "unseen genes", "unseen environments", and "unseen genes

and environments", which represented unseen genes and unseen cell types during

training (Figure S4, Methods). The model achieved a coefficient of determination (R^2)

of 0.81, 0.89, and 0.81 for the three datasets, respectively (Figures 2a), indicating its

robustness and ability to generalize.

To understand the relative importance of the three model inputs, we performed

ablation analysis, revealing that mRNA abundances were the most important

contributor to the prediction of translation (Figure 2b, Table S2), in agreement with an

early study of yeast translation that found that the most predictive variable for

translation was the mRNA expression of the gene[28]. The incorporation of codon

sequences lifted the R^2 by 0.15, and further inclusion of cellular environment

improved the R^2 by 0.06. The ablation analysis demonstrated that all the inputs

contributed to predicting mRNA translation.

We next investigated whether our model captured complex sequence features

beyond common translation-related metrics. While our model learned relevant

sequence features directly from the raw codon sequences, we tried to include common

translation-related sequence metrics including CAI, MFE, and codon frequencies as

additional model inputs and found these metrics did not improve prediction accuracy

(Table S1). This suggested that the model could capture the sequence patterns that

were predictive of translation, beyond these sequence metrics.

We explored alternative approaches to incorporating cellular context information.

We directly incorporated the meta information of Ribo-seq datasets into the model,

including cell types and experimental conditions and found it did not improve the

performance (Table S1). This indicated that the gene expression profiles used in the

model were an effective proxy to capture the relevant cellular environment

influencing mRNA translation.

Finally, we investigated the positional importance of coding sequences in

translation prediction. We analyzed the importance of each nucleotide position for the

model's prediction (Methods). The results showed that the coding sequences close to

the translation start site (TSS) were more important (Figure 2c). This is consistent

with a general knowledge that codons near TSS have a greater impact on protein

synthesis, by influencing translation initiation[8].

Overall, the data-driven approach of RiboDecode enabled robust predictive

capabilities with biological relevance by learning important sequence patterns directly

from the Ribo-seq data.

RiboDecode's Optimization Strategies for Enhanced mRNA Translation

Having established the efficacy of our translation prediction model, we next explored

how this model could be leveraged to generate sequences with enhanced translation

potential. We first generated codon sequences of Gaussia luciferase (Gluc) (Figure

S5). T-distributed stochastic neighbor embedding (t-SNE) indicated that the model established an association between the sequence space and translation levels (similar to Ribo-seq-derived RPKM values). The red area in the upper right showed that a wide space of high translation sequences was explored (Figure 3a). We next explored how RiboDecode optimized translation and stability independently or jointly. A widely used Gluc sequence was used as a reference for comparison, which had a predicted translation level of 5.9 and an MFE value of -216 (Figure 3b). By optimizing the sequence for translation (w=0), the predicted translation level increased to around 25. On the other hand, codon sequences optimized for MFE (w=1) reduced the MFE from -150 kcal/mol to around -350 kcal/mol, with a similar translation level to the reference. With joint optimization (0<w<1), RiboDecode explored a wider sequence space, achieving both enhanced translation and reduced MFE. Moreover, RiboDecode-generated sequences spanned a broader embedding space compared to those produced by Ribotree, CDSfold, and LinearDesign (Figure S6, see Methods), suggesting enhanced sequence diversity. Furthermore, designing Gluc codons for different cell lines showed that the generated codons had distinct sequence patterns for different cellular contexts, reflecting differences in cellular environment (Figure S7).

To understand RiboDecode's optimization strategy, we analyzed codon usage patterns between generated sequences with enhanced and reduced translation, as well as between high- and low-translated endogenous sequences. We found that codons preferentially used in highly translated endogenous sequences were also favored in RiboDecode-generated sequences with enhanced translation. Notably, the differences in codon usage between RiboDecode's enhanced and reduced translation sequences were more pronounced than the differences observed in endogenous sequences (Figures 3c, Methods). To assess the generalizability of these findings, we extended our analysis to multiple genes across various cell types. Consistently, we observed the same pattern of biased codon usage in all the cases (Figure S8). This suggests that

RiboDecode not only mimics but amplifies the codon usage patterns of efficiently

translated endogenous mRNAs, potentially leading to even greater improvements in

translation.

We next examined how RiboDecode utilized sequence features during generation

and optimization. Analysis of sequence features across different mRNAs revealed

complex and variable relationships with translation (Figures 3d and S9, Methods).

Notably, highly translated mRNAs generally showed an increase in uridine content

(U%), which may reduce secondary structure formation and facilitate smoother

ribosome movement during translation[7]. Additionally, these mRNAs mostly exhibited

a decrease in Effective Number of Codons (ENC), suggesting a selection against rare

or inefficient codon pairs to enhance translation[38]. Variations in CAI, Codon Pair Bias

(CPB), GC content (GC%) and MFE across different mRNAs suggested that while

these features could influence translation, their impact might be more mRNA- or

context-dependent, due to complex sequence or structure feature changes.

In short, these findings highlight RiboDecode's ability to capture complex

sequence-translation relationships, offering a sophisticated approach to mRNA

optimization that goes beyond traditional codon optimization metrics.

Experimental Validation Demonstrates RiboDecode's Versatility and Efficacy

While our *in silico* analyses demonstrated the potential of RiboDecode to optimize

codon sequences, we next sought to validate these findings experimentally. We first

validated RiboDecode's ability to optimize codon sequences for enhanced protein

expression. For Gluc, protein expression levels of the RiboDecode-optimized

sequences surpassed the reference and were more than twice as high as that of the

best-performing sequences designed by LinearDesign (p-value=0.019, one-sided

Mann-Whitney U test, Figure 4a and S17a, Table S3). The predicted translational

levels showed a positive correlation with the experimentally measured protein levels

(Correlation coefficient=0.71, p-value=0.077, Pearson's correlation, Figure S10).

However, the p value is marginal, potentially due to the small number of constructs

tested. Further validation with larger datasets is needed. In contrast, the CAI showed negative correlation with the experimental measurements (Correlation coefficient=-0.15), indicating that CAI is not a reliable predictor of protein expression levels and ineffective optimization strategy in this context. We noticed that the two RiboDecode-designed mRNAs (RD1 and RD2) had the best performance which also had the highest MFE values of around -200. In contrast, the LinearDesign sequences had MFE values of -350 and -300. To rule out the increased protein production was associated with higher MFE value, we tested additional LinearDesign sequences with higher MFEs (-266.8 and -246.70 kcal/mol, respectively). Compared to four LinearDesign sequences, RiboDecode-designed sequences outperformed (Figure S11).

We additionally optimized another commonly used reporter gene, firefly luciferase (Fluc). Seven sequences, including four RiboDecode-optimized (RD1-RD4), two LinearDesign-optimized (LD1, LD2) and a WT, were transfected into HEK293T cells, and activity was measured over 72 hours (Figure S12, Table S6). All optimized sequences significantly outperformed the WT. The LD1 (CAI of 0.766) and LD2 (CAI=0.952) yielded the increased expression with about 7-and 41-fold changes, respectively. RiboDecode sequences (CAI of ~0.71-0.73) also achieved substantial improvements with 6- to 16-fold over the WT. The superior performance of the LD2 sequence with a high CAI value here—contrasting with results for Gluc—underscores that the optimal codon strategy is gene-specific. This highlights the need for context-dependent approaches like RiboDecode that capture complex features beyond single metrics like CAI.

We next evaluated RiboDecode's ability to design for cellular context by optimizing Gluc mRNA for preferential expression in HEK293T cells over A549 and ARPE19 cells. The designed variants successfully exhibited the intended higher expression in HEK293T compared to both cell lines (Figure 4b, Table S4). Predicted expression ratios favoring HEK293T closely matched experimental results in the comparison against A549 (~1.7-fold predicted vs. ~1.55-fold experimental).

Preferential expression was also achieved against ARPE19 (Figure 4b, Table S5),

although the experimental fold-change (~2.8-fold) doubled the prediction (~1.4-fold).

These results confirm RiboDecode's capability for context-aware design, while the

discrepancy in the ARPE19 comparison indicates potential for refining the model to

better capture quantitative differences across diverse cellular environments.

Modified mRNAs, such as those with 1-methylpseudouridine (m$^1\Psi$)

modifications, and circular RNAs are used in mRNA therapy instead of unmodified

mRNAs due to their improved stability and reduced immunogenicity[3,7,39]. We

therefore assessed the effectiveness of RiboDecode in enhancing translation in these

alternative mRNA forms. Among the four codon variants, m$^1\Psi$-modified RD2 and

RD4 showed higher protein expression levels compared to the reference, with up to a

4.6-fold higher expression at 48 hours post-transfection (Figure 4c and S17b).

Moreover, all four RiboDecode-generated codon variants in the circular form

outperformed the reference (Figure 4d and S17c). These results demonstrate that

RiboDecode optimization enhances protein production in both m$^1\Psi$-modified and

circular mRNAs, illustrating its reliability and versatility.

These experimental validations demonstrate RiboDecode's ability to significantly

enhance protein expression, optimize in specific cell types, and improve translation

across various mRNA forms, highlighting its potential as a powerful tool for mRNA

therapeutic development.

RiboDecode Enhances Immunogenicity of mRNA-based Influenza Vaccines

Having established the robustness of our optimization approach, we next aimed to

demonstrate its practical application in the development of mRNA-based vaccines.

Influenza A viruses are responsible for causing respiratory infections, leading to

annual epidemics that result in millions of human infections worldwide[40]. HA, a

glycoprotein found on the surface of influenza A viruses, plays a crucial role in the

viral infection process and is the primary target for the development of influenza

vaccines. Although most of the vaccines were developed using inactivated influenza viruses, mRNA-based influenza vaccines are currently actively developed[41].

To enhance the expression of HA and potentially improve the efficacy of HA-based vaccines, we optimized the HA coding sequence. Three out of four RiboDecode-optimized HA sequences and two LinearDesign-optimized sequences showed higher *in vitro* protein expression compared to the WT (Figures 5a, Table S7). Particularly, RD3 showed approximately 6-fold increase compared to the WT and LinearDesign-optimized sequences. In addition, RD3 exhibited considerably higher expression levels compared to the WT sequence in both $m^1\Psi$-modified and circular mRNA forms (Figures 5b and 5c). These results again highlight the robustness and versatility of the RiboDecode-optimized sequence.

We further assessed the *in vivo* immunogenicity induced by the optimized sequence for both the prime and boost responses, where split virus influenza vaccine (SV) was served as the positive control (Figure 5d, Methods). The RD3 sequence induced significantly stronger neutralizing antibody responses, measured by the micro-neutralization (MN) titers, compared to the WT sequence and SV. For the prime response, RD3 elicited significantly higher MN titers compared to WT, with approximately 4.4-fold increase (Figure 5e, mean MN titers: RD3=2,560, WT=580; p-value=0.008, one-sided Wilcoxon test). The difference was more pronounced for the boost response, with RD3 inducing a 9.6-fold increase in MN titers compared to WT (Figure 5e, mean MN titers: RD3=83,200, WT=8,640; p-value=0.002, one-sided Wilcoxon test). These results demonstrated that the RiboDecode-optimized sequence significantly enhanced both the initial and boosted immune responses. This dramatic improvement in immunogenicity underscores RiboDecode's potential to enable more effective vaccines with lower doses.

 Enhanced Protein Expression and Therapeutic Efficacy with optimized NGF mRNA

Having demonstrated the efficacy of RiboDecode in optimizing mRNA for vaccine

development, we next explored its potential in protein replacement therapy. We

focused on NGF as a promising candidate for treating glaucoma, which is a leading

cause of irreversible blindness[42] and causes death of retinal ganglion cells (RGCs).

Our recent study demonstrated that mRNA-based NGF therapy provided robust

neuroprotection for RGCs in an optic nerve crush (ONC) mouse model[43].

To improve the neuroprotection efficacy, we optimized the codon sequences of

human NGF mRNA. The protein expression levels of three RiboDecode-designed

sequences were more than 2-fold higher compared to that of the WT whereas the

LinearDesign sequences did not show improvement (Figure 6a and S18a, Table S8).

We further assessed the best performing sequence (RD3) in both $m^1\Psi$-modified

mRNA and circular mRNA forms. Notably, with $m^1\Psi$-modification, RD3 achieved

8.4- and 9.8-fold higher protein levels compared to the WT at 24h and 48h,

respectively (Figure 6b and S18b). With mRNA circulation, RD3 also achieved a

more than 2-fold higher expression than the WT at both 24h and 48h (Figure 6c and

S18c). These results again demonstrated the robustness of RiboDecode-optimized

sequences across different mRNA forms.

Based on its superior performance in initial *in vitro* tests, we selected RD3 for

further *in vivo* studies. To evaluate the *in vivo* expression of optimized NGF mRNA,

we intravitreally administered both the RD3 and WT sequences. Each mRNA was

$m1\Psi$-modified and encapsulated within LNP and administered at two doses: 100

ng/μl and 500 ng/μl. The RD3 sequence demonstrated significantly higher NGF

protein expression compared to the WT sequence at both doses. Remarkably, RD3 at

100 ng/μl achieved even slightly higher expression than WT at 500 ng/μl (Figure 6d).

We then investigated the therapeutic potential of optimized NGF mRNA using an

ONC mouse model, which mimics RGC injury and resulted in significant RGC loss

(Figures 6e-g). Treatment with NGF mRNA showed clear neuroprotective effects,

preserving more RGCs after injury. Notably, mice treated with 100 ng/µl RD3 showed

significantly higher RGC counts than those treated with the same dose of WT mRNA

(Figures 6h and 6i, p-value=0.0002, one-sided Wilcoxon test). Moreover, these counts

were comparable to those in mice treated with 500 ng/µl WT mRNA.

To sum, the optimized sequence exhibited superior protein expression both *in*

vitro and *in vivo*, while maintaining therapeutic efficacy at one-fifth the dose of the

unoptimized sequence. These results demonstrated the effectiveness of RiboDecode in

optimizing NGF mRNA for the treatment of RGC injury.

DISCUSSION

In this study, we present RiboDecode, a novel deep learning-based framework for

mRNA codon optimization. The generative optimization framework, guided by the

deep learning prediction model, enables the efficient exploration of the immense

space of possible codon sequences. This allows RiboDecode to discover novel, highly

optimized sequences that may not be accessible to traditional optimization methods.

RiboDecode-optimized sequences demonstrate superior performance in various

mRNA formats, including unmodified, $m^1\Psi$-modified, and circular mRNAs,

highlighting its broad applicability in the rapidly evolving field of mRNA therapeutics.

In vitro and *in vivo* experiments using the optimized sequences of therapeutically

relevant proteins show substantial enhancements in protein expression compared to

the unoptimized sequences. These improvements further translate into increased

therapeutic efficacy, as demonstrated by significantly enhanced immune responses to

an optimized influenza vaccine and markedly improved RGC protection in mice with

optic nerve injury.

While mRNA abundance serves as both an input feature and intrinsically linked to

the Ribo-seq RPKM target variable, potentially explaining its high predictive weight

in ablation analysis, the model demonstrated significant improvements in prediction

accuracy by integrating codon sequences and cellular context. Importantly,

RiboDecode's *in vitro* and *in vivo* validation, where optimized sequences substantially enhanced protein expression and therapeutic efficacy, confirms its ability to optimize biologically meaningful translational signals rather than merely reflecting transcript abundance. The superior performance of RiboDecode may be attributed to several factors. Firstly, RiboDecode's deep learning model learns directly from diverse nature sequences with translation measurements, enabling it to capture complex patterns of codon sequences for mRNAs with high translation level. Second, the model considered the cellular contexts of mRNA translation. Third, RiboDecode's generative optimization framework allows it to explore a large sequence space and to discover novel, highly optimized sequences that may not be accessible to the traditional approaches.

These findings also align with the broader framework of gene expression regulation: while mRNA abundance is a prerequisite for protein synthesis (consistent with its predictive dominance), our approach directly targets translational control, a critical layer highlighted by Schwanhäusser *et al.*[44], by leveraging Ribo-seq-derived codon usage and cellular context to maximize translational efficiency. Although the model excludes post-translational events (as Ribo-seq captures ribosome activity prior to protein maturation), the achieved gains in protein output and *in vivo* efficacy robustly validate translational optimization as a key determinant of functional protein levels, complementing established regulatory hierarchies.

The findings of our study have important implications for the field of mRNA therapeutics. Firstly, RiboDecode can generate and evaluate a vast number of novel codon combinations. This capability allows RiboDecode to optimize mRNA sequences beyond the limitations of evolutionary constraints, potentially uncovering more efficient codon usage patterns not found in natural transcripts. Second, by substantially increasing protein production, the optimized sequences can improve the potency and reduce the required dose of mRNA-based treatments, potentially mitigating side effects and enhancing patient outcomes. This is particularly relevant

for applications such as protein replacement therapies, where achieving high levels of protein expression is crucial for therapeutic success. Third, RiboDecode's robustness and versatility across different mRNA formats, including modified and circular mRNAs, expand the range of therapeutic applications for which it can be employed. As the field of mRNA therapeutics continues to evolve and new mRNA formats are developed to enhance stability, reduce immunogenicity, and improve delivery[39,45], RiboDecode's ability to optimize sequences for these diverse formats will be invaluable.

While our study demonstrates the significant potential of RiboDecode in optimizing mRNA codon sequences for enhanced mRNA translation and therapeutic efficacy, there are several future directions to explore. Firstly, we focused exclusively on optimizing the codon sequences while not explicitly modeling the 5'UTR. Recognizing the critical role of the 5'UTR in regulating translation initiation, our future work aims to expand RiboDecode to jointly optimize both UTRs and the codon sequence. Second, our results indicate that while our primary goal was to enhance translation through codon optimization, incorporating MFE optimization synergistically modulating mRNA secondary structure stability and translation efficiency. However, we observed the best w value appears context-dependent, suggesting a one-size-fits-all approach may be suboptimal. Consequently, we recommend that experimental designs include the testing of multiple w values to identify the optimal balance for each mRNA target. Further research into the determinants of the optimal w value will be critical to refine this strategy and could lead to more systematic approaches for integrating MFE into mRNA design. Third, our MFE model cannot predict MFEs for unseen mRNAs. For an unseen mRNA, the model must first train the sequences with MFEs labelled by RNAfold. A general MFE model should be developed and used in future. Finally, the model was trained exclusively on Ribo-seq data from endogenous, unmodified mRNAs. Although our results showed significant expression enhancements for optimized sequences in both

m1Ψ-modified and circular forms compared to their respective controls, the relative

fold-improvement compared to the unmodified format varied between constructs.

Future iterations could potentially incorporate data from modified and circular

transcripts to further refine optimization rules.

In conclusion, RiboDecode represents a paradigm shift from rule-based to data-

driven mRNA optimization, potentially uncovering entirely new principles of efficient

translation that were previously inaccessible. RiboDecode will provide a versatile tool

for researchers to maximize the potential of mRNA-based therapeutics, paving the

way for more effective treatments in various medical applications.

MATERIALS AND METHODS

Data collection and processing

Data preprocessing and filtering

We downloaded translation counts of Ribo-seq datasets from the RPFdb

database[33,34,46]. The following steps were implemented for processing the Ribo-seq

data in accordance with RPFdb: First, to prevent adapter interference in downstream

analyses, the 3' adapter sequences were manually extracted for each dataset from the

original publications or the corresponding MultiQC[47] outputs. Adapter sequences, if

present at the ends of sequencing reads, were subsequently removed using Cutadapt

(version 1.16)[48]. Next, to minimize rRNA and tRNA contamination, sequences

corresponding to rRNA and tRNA were retrieved for each species from ENSEMBL[49]

and UCSC[50] databases and removed post-mapping using Bowtie2[51]. Finally, to ensure

the retention of high-quality ribosome-protected footprints, which exhibit a

characteristic read-length distribution reflecting the size of a translating ribosome on

the RNA, only footprints within the 25–34 nucleotide length range were retained after

contaminant removal and alignment. The count tables were transformed to reads per

kilobase per million (RPKM). Because some of the paired RNA-seq were not

available in RPFdb, we reprocessed the RNA-seq datasets. The raw FASTQ files of

RNA-seq were trimmed by sickle[52] for adapter removal and quality control. Then, to

filter out reads from tRNA or rRNA, we mapped the reads to human tRNA and rRNA

reference sequences (hg38) using bowtie2[53] (-L 20). The unmapped reads were then

mapped   to   the   human   genome   (GRCh38,   gencode.v28,

https://www.gencodegenes.org/) using STAR[54] (2.7.4a). Finally, read counts for each

gene were summarized by featureCounts[55] (v2.0.1, -t exon). The expression counts of

Ribo-seq and RNA-seq were ln(RPKM $\times$ 5+1) transformed. Genes with low

expression (median RPKM<1) were filtered out. Finally, 11,725 coding genes from

320 samples with 24 cell types were included in this study (Supplementary file 1).

Cross-validation dataset preparation

Out of 11,725 genes in the Ribo-seq data, we randomly selected 1,173 genes (1/10 of

the total genes) that were not included in the training datasets, as the "unseen genes"

dataset. Out of 320 Ribo-seq datasets, we randomly selected 120 datasets whose cell

types were not included in the training datasets, as the "unseen environments" dataset

(Supplementary file 1). The "unseen genes and environments" dataset was also

defined (Figure S4).

mRNA isoform selection and sequence encoding

To address the complexity of alternative splicing while managing computational

feasibility, we utilized the major isoform of each gene as a representative for mRNA

codon variants. This approach was necessitated by the inherent limitations of NGS

data in accurately quantifying the proportions of individual isoforms. We defined the

major isoform as the transcript with the highest expression level, estimated using

RSEM[56]. In total, our dataset contained 60,255 different mRNA sequences.

Translation Model Architecture

Translation is influenced by multiple factors, including codon sequences as the pivotal

signals modulating translation[57], trans-acting elements modulating the cellular

environment of translation[58,59], and mRNA abundance which provides more templates

for translation[60,61]. To capture these interacting variables, a translation model was

developed using a deep neural network architecture comprising 2 convolutional layers

and 5 fully connected (FC) layers[62,63]. The model inputs are a codon sequence in one-

hot encoding, a transcript abundance, and a vector of gene expression profiles from

RNA-seq presenting cellular environment (Figure S1). The codon sequences were

fixed to 4,500 base pairs (bp) starting from the translation start site, which covers over

98% of coding sequences, and those shorter than 4,500 bp were zero-padded on 3' end.

To process these inputs, 3 sequential FC layers extract features from the cellular

environment vector, which are then concatenated with the transcript abundance to

form an attention vector. In parallel, convolutional neural networks (CNNs) processes

the one-hot encoded codon sequence, generating embeddings with four distinct

feature sets. Subsequently, the averaged feature is encoded through the other

convolutional layers and flattened. Finally, 2 FC layers yield the predicted translation

level output. To enhance model robustness and prevent overfitting, batch

normalization[64] and dropout[65] techniques are employed after each layer, and max-

pooling[66] is utilized following the convolutional layers. For optimization, the AdamW

optimizer[67] is adopted in conjunction with the SmoothL1Loss loss function and ReLU

activation functions[68]. Additionally, gradient clipping and learning rate decay

strategies are implemented to mitigate gradient explosions and instability during

training. The model was parameterized by a total of 15 hyper-parameters, as detailed

in Table S9. The model was pretrained for 20 epochs before being used by the codon

optimizer of RiboDecode. Model training and validation were assessed using the R-

squared metric (R^2). The best-performing model was selected based on the highest R^2

value achieved on the validation set. The R^2 metric, defined as R^2 = 1 -

($SS_{Residual}/SS_{Total}$), quantifies the goodness of fit by comparing the sum of squared

residuals ($SS_{Residual}$) to the total sum of squares (SS_{Total}). The PyTorch 1.12.0

framework was leveraged for the implementation of the deep model.

The translation model's architecture, which uses a fixed 4,500 bp input, results in

a high nominal parameter count. This is an artifact of flattening the zero-padded input

sequences required for the majority of genes, which have a median length of only

1,200 nt. To mitigate the resulting risk of overfitting, we implemented a strong

dropout regularization strategy (rate of 0.9) before the final layers. This high dropout

rate forces the model to learn robust features rather than fitting to noise in the padded

regions. A comparative study (Figure S15) confirms this strategy is essential: models

with no or low dropout overfit immediately, whereas the 0.9 dropout rate eliminates

overfitting and ensures stable, improving performance on all validation sets.

MFE Model Architecture

The MFE prediction model was formed by a deep neural network architecture

comprising 2 convolutional layers, 9 residual blocks derived from ResNet[69], and 3 FC

layers (Figure S2a). The model input is the one-hot encoded codon sequence, which is

the same as the codon sequence used in the translation model. Each residual block

consists of 2 convolutional layers with a residual connection. Initially, shallow

features are extracted through 2 convolutional layers, followed by 5 residual blocks

with max-pooling for deep feature extraction. Subsequently, 4 residual blocks without

max-pooling are used to maintain spatial resolution. After the final residual block,

features are flattened and fed into 3 FC layers to predict the MFE value. Batch

normalization[64] is applied after the first 2 convolutional layers, and dropout[65] is

introduced after the subsequent 2 residual blocks. The Fast Gradient Method (FGM)[70]

is integrated during training to enhance generalization by applying perturbations to

input sequences. We utilized the AdamW optimizer[67], SmoothL1Loss function, and

LeakyReLU activations[71]. The final loss is defined as $Loss = Loss_{mfe} + Loss_{fgm}$, where

$Loss_{mfe}$ is the SmoothL1Loss between predictions and RNAfold[35] MFE values, and

$Loss_{fgm}$ is the SmoothL1Loss with added perturbations. The model is trained

alongside optimization by the codon optimizer of RiboDecode using generated

sequences as the training data, with performance evaluated using the R^2 metric.

Details of 15 hyperparameters are provided in Table S10. The PyTorch 1.12.0

framework and RNAfold 2.4.18 were used through Python interfaces. We evaluated

the MFE values of mRNAs between our model and RNAfold. We found that our MFE

value highly agreed with the one from RNAfold (Figure S13a), showing the reliability

of our MFE model. To determine the optimal training set size, we compared our

model trained on a compact set of 210,000 sequences against a model trained on an

extended dataset of 11.5 million sequences (achieving a data-to-parameter ratio > 1).

Both models achieved comparable optimization performance and yielded MFE

predictions highly correlated with values from RNAfold (Figure S13). Given the

similar performance, we selected the 210K training set for our framework, as it

reduces computational time by approximately 75% without sacrificing model

reliability or accuracy.

We evaluated our MFE optimization framework against existing MFE

optimization methods, including the general-purpose, differentiable model (JAX-

RNAfold) [72]. Our analysis revealed that the high computational complexity of JAX-

RNAfold severely limits its use to short sequences (under 600 nt on our V100 GPU

with 32 GB video RAM), making it unsuitable for our primary goal of optimizing

full-length human mRNAs. To perform a direct, head-to-head comparison, we used a

compatible sequence (Gaussia luciferase, 558 nt). Our model achieved a lower (more

favorable) MFE, completing the optimization faster and with substantially less GPU

memory than JAX-RNAfold (Table S12). While LinearDesign is another powerful

tool, it is not a differentiable model and is thus incompatible with our gradient-based

optimization framework. Given the critical need for scalability to therapeutically

relevant sequence lengths and the efficiency observed in direct comparison, we

proceeded with our sequence-specific MFE optimization approach.

Codon Optimizer Architecture

Fitness score

The fitness score combines mRNA translation level and MFE predicted by above
models.

For translation level, the prediction of a mRNA sequence is conducted using the
pre-trained translation model with the cellular environment and transcript abundance
holding constant during the optimization. The loss function is designed as follows:

$$L_t = \sqrt{(r - S_t)^2}/\alpha$$

625   Here, S_t is the predicted translation level, r represents the desired output value of
626   the translation model, and α is a constant based on S_t.

627   For MFE, the loss function is formulated based on model parameters of the
628   current epoch since the MFE model is trained alongside the optimization:

$$L_m = -\beta/S_m$$

where, S_m is the predicted value of the current epoch and β is a constant set based on
it.

The introduction of α and β is intended to balance the loss in the translation
optimization and MFE optimization processes. The output range of the translation
model ($S\square$) typically lies between 0 and ~100 (e.g., 0 to 25 for Gluc), while
RNAfold-derived MFE values generally range between -100 and -1000 (e.g., -350 to -
150 for Gluc). To ensure comparable magnitudes and numerical stability during
optimization, we introduced scaling factors α and β to normalize both loss
components. For most codon sequences exhibiting translation prediction values < 100
and MFE values > -1000 kcal/mol, $\alpha=\beta=100$ provide appropriate normalization (Table
S11). However, we recommend parameter adjustments under extreme value
conditions: α should be increased to 1000 when translation predictions exceed 100,
while β should be elevated to 1000 when MFE values fall below -1000 kcal/mol.

The final optimized loss function is defined as follows:

$$Loss = L_m * w + L_t * (1 - w),$$

Where w is a constant ranging from 0 to 1. When w is set to 0, only the

translation of mRNA is optimized. When w set to 1, only the MFE of mRNA is

optimized. When w set to a constant value between 0 and 1, both models are

optimized simultaneously, with the magnitude indicating the relative emphasis on

optimizing each model.

Optimization process of the codon optimizer

The codon optimizer uses a gradient ascent optimization approach based on

activation maximization (AM)[30] to generate synonymous codon sequences with

optimized fitness score. The optimization process involves the following steps:

1. Initial representation: the codon distribution is initially represented in a one-

hot manner, where each position is assigned to the specific codon of the original

codon sequence.

2. Optimization: the optimized codon distribution by AM becomes probabilistic,

assigning a likelihood to each possible synonymous codon at every position, with the

goal of maximizing the fitness score.

3. Regularization: during the optimization, a synonymous codon regularizer is

used to ensure that the optimization process only adjusts the selection probabilities

within the synonymous codons capable of encoding the same amino acids as in the

original sequence. The regularizer applies a constraint on the codon distribution, T,

given by a synonymous substitution mask matrix W:

$$W = \begin{bmatrix} w_{11} & \cdots & w_{i1} \\ \vdots & \ddots & \vdots \\ w_{1j} & \cdots & w_{ij} \end{bmatrix}$$

663   where w_{ij} signifies the selection of the j^{th} coding category at the i^{th} position, with

664   value 1 for a synonymous codon, and 0 otherwise. Here, $i = 1, 2, \ldots, L$ and $j = 1, 2, \ldots,$

665   C.

666   Subsequently, regularization is performed on T:

$$T = \frac{T \odot W}{\sum_{i=1}^{L}(T \odot W)_i}$$

667 The maximum index values are then converted to a one-hot representation to

668 obtain $T_{one\text{-}hot}$:

$$T_{one-hot_{ij}} = \begin{cases} 1, & if\ j = argmax(T_i) \\ 0, & otherwise \end{cases}$$

4. Sequence generation: after the gradient updates, a new codon sequence is

generated by selecting codons with the highest probabilities.

5. Iteration: This sequence then re-enters the codon optimizer for further rounds

of optimization.

We employed the Adam optimizer and utilized a learning rate decay strategy at

different training stages. The optimization process was regularized with a weight

decay of 1×10^{-4}. The learning rate was initialized at 5×10^{-4} and decayed by a factor of

10 when the generated data reached 1/2, 3/4, 7/8, and 16/17 of the total 40,000

sequences for one epoch. The optimization process was conducted over 20 epochs.

However, we observed that all genes converged to maximum translation levels before

the 7th epoch. Consequently, a total of 280,000 sequences were utilized to plot the

progression of the generation.

Integrative training of MFE model

To train the MFE model, we adopted an active learning strategy that integrates

model training with sequence optimization by the codon optimizer. This approach

uses the RNAfold tool[31] for labeling and involves simultaneous training, optimization,

and sequence generation. The process consists of four interconnected steps that run

concurrently with the sequence optimization:

Initial sampling: we begin by generating 20,000 sequences through random

synonymous substitutions (up to 10% of codons) of the original sequence.

Initial training: these 20,000 sequences are input into both the MFE model for

predictions and RNAfold for ground truth MFE labeling.

Sequence generation: utilizing the trained model of current epoch, new sequences

are generated by the codon optimizer of RiboDecode. This pool is sorted by predicted

MFE, with lower values being better. The top 480 sequences are selected as generated

sequences. The top sequence undergoes two operations: random replacement (up to

10% of codons) and distributed replacement according to its generating codon

distribution. A total of 10,000 sequences are generated through this process.

Model retraining: The generated sequences, along with those from distributed and

random replacements, are used as input for MFE prediction and RNAfold labeling.

The model is then retrained on this new data.

Model running time

The computational infrastructure employed for our model training and sequence

optimization comprised a single NVIDIA Tesla V100 SXM2 GPU with 32 GB of

memory and an Intel Xeon Gold 5218 CPU operating at 2.30 GHz.

The translation model required approximately 24 hours for training completion.

For the Minimum Free Energy (MFE) model, training and optimization were

conducted concurrently. The duration of this process varied based on input sequence

length, ranging from 1 hour to 3 hours. Upon completion of model training, the

optimization phase was executed, which took about 1 hour for any sequences.

Optimized codon sequences screening strategy for experimental validation

To select optimal codon sequences for experimental validation, we employed a multi-

step screening process for each gene:

1. Sequence Generation: We generated three rounds of candidate sequences using

different optimization strategies: a) Translation-only optimization (w=0) b)

Joint optimization with moderate MFE consideration (w=0.5) c) Joint

optimization with stronger MFE consideration (w=0.7).

2. Initial Filtering: For each round, we filtered out potentially over-optimized

sequences by retaining only those variants with a predicted translation level

below the 90th percentile of all generated variants. This step helps avoid

unreliable over-optimization that might not translate to real-world

performance.

3. Selection Criteria: From the filtered sequences in each round, we selected

candidate sequences based on two criteria: a) High predicted translation level

b) Low MFE value.

4. Final Selection: We recommend selecting one or more candidates from each of

the three optimization rounds (w=0, w=0.5, w=0.7) for experimental validation.

This ensures a diverse set of optimized sequences, balancing pure translation

optimization with different levels of MFE consideration.

Model evaluation and analysis

Translation model evaluation

To evaluate the importance of each input component, we independently trained

multiple translation models on the cellular environment vector, transcript abundance,

and codon sequence and their combinations[20]. During training, the hyperparameter

settings and dataset partitioning were consistent for each model. We replaced the input

to be ablated with a zero tensor of the same shape and independently observed the

impact of each input component on the final translation prediction.

When evaluating the performance of the model with CAI, MFE and cellular

information as additional inputs, these new features were concatenated into the

attention vector of cellular environment features and transcript abundance for mRNA

translation prediction.

Furthermore, potential associations between genes in the unseen dataset and

those in the seen dataset may lead to overestimation of model performance. To

evaluate this potential issue, we obtained gene family annotations from the HUGO

Gene Nomenclature Committee (HGNC; https://www.genenames.org/) and excluded

genes in the unseen dataset that belonged to the same gene families as those in the

seen dataset (resulting in the removal of 130 genes). The results demonstrated only a

marginal decrease in prediction accuracy on the "unseen gene test set" (with R^2

decreasing from 0.81329 to 0.81295). Therefore, although some of the genes in the

unseen dataset are related to those in the seen dataset, the performance of the model

was not overestimated.

Nucleotide position contribution for translation prediction

To evaluate the importance of each nucleotide at different positions in the codon

sequence, we explored the attribution of translation model predictions to their input

features. Here, we implemented the attribution method of Integrated Gradients[73],

obtaining an importance score for each nucleotide position. This method combines the

implementation invariance of gradients with the sensitivity of techniques such as

LRP[74] or DeepLift[75]. Firstly, we determined a vector with all features set to zero as

the baseline value. Then, we linearly interpolated the input features from the baseline

value to the actual input values, with these intermediate values representing different

strengths or combinations of features. Subsequently, for each interpolated input, we

calculated the gradient of the model output relative to that input, and multiplied the

gradient at each interpolation point with the difference between the input feature value

and the baseline value, obtaining the contribution of that feature at each interpolation

point. Finally, we weighted and summed the contribution values at all interpolation

points to obtain the Integrated Gradient for that feature. For the final analytical

outcomes and visualization, the importance score assigned to each nucleotide position

was calculated as the mean of absolute values derived from non-padding regions.

Cellular environment and mRNA abundance for translation model

The default mRNA abundance level was set to 4.5 (ln-transformed RPKM×5+1,

median transcript abundance; Figure S14). To evaluate robustness, we additionally

tested input values of 3.82 and 5.19, corresponding to half and double the original

RPKM expression level, respectively. Notably, Gluc codon sequence predictions

remained highly consistent across these input variations, demonstrating the stability of

our model.

When optimizing mRNA sequences using RiboDecode, we recommend using the

RNA expression profile of the target or the most similar tissue or cell type through

either experimental sequencing or publicly available datasets. To facilitate

implementation, we have incorporated a user-defined environment input parameter

into the software package, accompanied by comprehensive documentation and step-

by-step instructions.

For example, to predict translation in HEK293T, the input of cellular environment

was constructed as follows. First, the mRNA expression levels of genes representing

environmental factors were obtained from RNA-seq data of untreated HEK293T cells.

Then, the batch effect arising from different data sources was corrected with

ComBat[76]. Finally, the mean value of the mRNA expression was taken as the cellular

environment input.

Codon usage analysis

Codon sequence variants with enhanced and reduced predicted translation levels were

generated in different cellular contexts, including HEK293T, HeLa, and A549. Codon

sequences of endogenous genes with high (top 10%) and low (bottom 10%)

translation level were selected from Ribo-seq data. We further evaluated the impact of

varying expression thresholds (top/bottom 5%, 10%, and 20%, Figures S8 and S16),

with consistent results observed across all cutoff values, except Gluc in A549 cells at

the 5% threshold. Then, the codon usage of generated and endogenous sequences was

calculated by the proportion of each codon among synonym codons. To get the

codons that appeared more in high-translated sequences, we performed t-test (p-value

< 0.05, after FDR adjustment for multiple-testing) on high and low-translated

sequences for both endogenous and generated sets. Codons with the significant and

greatest differences on codon usage in endogenous sequences were chosen as the top-

10 codons.

Sequence feature analysis

To analyze the sequence features of RiboDecode-optimized sequences compared to

non-optimized sequences, we followed these steps:

1. Sequence Generation: a) We randomly selected 2,000 RiboDecode-optimized codon sequences. b) We generated 2,000 non-optimized sequences by performing random synonymous codon substitutions on the unoptimized input sequence.

2. Feature Calculation: We calculated several sequence features for both sets of sequences, including Codon Adaptation Index (CAI), Codon Pair Bias (CPB), Effective Number of Codons (ENC), GC content (GC%), Minimum Free Energy (MFE) and Uracil content (U%).

3. Fold Change Calculation: For each feature, we calculated the fold change by dividing the median value of the optimized sequences by the median value of the non-optimized sequences.

4. Data Transformation: We ln-transformed the fold change values for each feature to normalize the distribution.

5. Data Scaling: The ln-transformed fold change values were scaled to a range of -1 to 1 for visualization purposes.

6. Visualization: The scaled values were used to create a heatmap representation of the feature changes (Figure 3d). For detailed distributions of each feature, refer to Figure S9.

Parameters used for sequence optimization

For Gluc, HA, and NGF sequence optimization, four sequences were designed by RiboDecode (RD1, RD2, RD3, and RD4: w=0, 0, 0.7 and 0.5, respectively), and two sequences were designed by LinearDesign (LD1 and LD2: λ=0 and 4). The WT sequence was also used as a reference.

To preliminarily evaluate mRNA optimization *in vitro*, we first performed codon optimization for HA and NGF for the HEK293T cellular environment and measured their protein expression in HEK293T cells (Figures 5a-c and 6a-c).

Comparative analysis of sequence space across design methods

We generated 1,000 full-length Gluc codon sequences using Ribotree, LinearDesign,

CDS-fold, and RiboDecode, respectively. High-dimensional sequence embeddings

were extracted using CodonBERT[24], followed by t-SNE dimensionality reduction for

visualization.

mRNA preparation

Plasmid construction

The 5'homology sequence, IRES sequence, 3'homology sequence, E1/E2 sequence

and protein coding sequence, were chemically synthesized, and cloned into the vector

pUC57, which contains a T7 RNA polymerase promoter.

For linear mRNA, the plasmids contain the 5'UTR, protein coding region, 3'UTR

and 105nt poly-A elements. A 3xflag tag was added after the coding region, in order

to detect the protein expression by western blot. The UTR and coding sequences were

listed in Supplementary file 2.

Linear mRNA production and modification

The linear mRNAs were produced using the HiScribe T7 High Yield RNA Synthesis

Kit and capped with m7G(5')ppp(5')G RNA Cap Structure Analog (NEB, #S1404).

Then, the RNA was column-purified.

For mRNA $m^1\Psi$-modification, N1-Me-Pseudo UTP (Yeasen Biotechnology,

#10651ES) was used to replace the unmodified UTP.

Circular mRNA production and purification

The *in vitro* transcription (IVT) of circular mRNA was carried out from linearized

circular mRNA plasmid templates with the HiScribe T7 High Yield RNA Synthesis

Kit (New England Biolabs, E2040S) following the kit manual. After IVT, circular

mRNA was purified using the Monarch RNA Cleanup Kit (New England Biolabs,

#T2050L). Then, the RNA precursors were heated to 70°C for 3 minutes and

immediately placed on ice for 2 minutes. GTP was added to a final concentration of

2mM with a buffer (50mM Tris-HCl, 10mM MgCl2, 1mM DTT, pH 7.5) for 8 minutes at 55°C to catalyze the cyclization. Then RNA was column-purified.

For circular mRNA purification, we collected RNA fractions through UV absorbance at 260nm on an Agilent 1260 Series HPLC (Agilent) system with a 4.6×300mm column (Sepax Technologies, #215980P-4630) at a flow rate of 0.3mL/minute. The fractions were concentrated with a 4ml Ultracel-10 regenerated cellulose membrane (Millipore, #UFC8010) and purified by column chromatography. Then, the RNA was treated with RNase R (Beyotime, R7092L) for further enrichment. Finally, RNase R-digested RNA was column-purified.

In vitro experiments

mRNA transfection

HEK293T cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) (BasalMedia, #L110KJ) containing 10% fetal bovine serum (NATOCOR, #SFBE) and 1% penicillin-streptomycin (GIBCO, #15140122) at the condition of 37°C and 5% CO_2. mRNAs were transfected into HEK293T cells using Lipofectamine MessengerMax (Invitrogen, #LMRNA015) according to the manufacturer's instructions. At the appropriate time after transfection, the cell lysate or supernatant was collected for protein detection.

***In vitro* protein expression measurements**

For measurement of Gaussia luciferase activities, HEK293T cells were seeded in 96-well plate and transfected with 150ng mRNA per well. The cells were lysed at 6, 24, 48 and 72 hours after transfection using 1× cell lysis buffer from Dual Luciferase Reporter Assay Kit (Vazyme Biotech, DL101-01). Then, the luminescence signal was detected following the provided instructions.

For *in vitro* quantitative measurement of nerve growth factor (NGF), HEK293T cells were seeded in 24-well plate and 500ng of RNA was transfected into the cells per well. The cell culture supernatants were collected 6, 24, 48 and 72 hours after

transfection. Then, the protein expression level was detected using Enzyme-linked

Immunosorbent Assay Kit (Cloud-Clone Corp, #SEA105Mu), following the provided

instructions.

For measurement of HA protein level, HEK293T cells were seeded in 12-well

plate and transfected with 1.25ug mRNA per well. After 24 hours, cells were

harvested and collected in 300ul RIPA lysis buffer (HANGZHOU DUDE

BIOLOGICAL CO.LTD, #FD009) that contained 1% PMSF (HANGZHOU DUDE

BIOLOGICAL CO.LTD, #FD0100). Then, protein level was analyzed with western

blot using anti-flag primary antibody (Sigma-Aldrich, #F1804).

Each experiment was repeated three or four times from distinct samples.

Cell type specificity experiments

We used RiboDecode to design three Gluc mRNA variants optimized for preferential

expression in HEK293T cells. The optimization process considered the cellular

context of HEK293T cells while maintaining or reducing expression levels in A549

and ARPE19 cells. Gluc protein expression was measured 24 hours post-transfection.

Each experiment was performed in quadruplicate. Expression levels were normalized

to the reference (MF882921.1) Gluc mRNA for each cell type.

In vivo experiments

Mouse retina histology and microscopy

For retinal whole-mounts immunofluorescence, eyes were surgically removed from

perfused mice and fixed with 4% PFA at room temperature for 1 hour. Retinas were

detached and whole mount staining was performed. The retinas were blocked for 1

hour in PBS staining buffer containing 5% normal donkey serum (Solarbio, SL050)

and 0.1% Triton X-100 (Sigma, X100-100). The retinas were incubated with the

primary antibody (Novus, NBP2-20112) overnight at 4°C and washed 3 times with

PBS for 5 minutes each before incubation with the secondary antibody (CST, 4413S)

for 2 hours at room temperature. The retinas were washed again with PBS 3 times for

5 minutes each and then mounted.

Confocal images were obtained using a Zeiss LSM 980 microscope. To count

retinal ganglion cells (RGCs), we analyzed 320×320μm samples from the peripheral

retina. These samples were taken ~500μm from the center to the edge in all four

quadrants of the retina. We then processed the data using ImageJ and ZEN software.

mRNA intravitreal injection

Adult mice were anesthetized by intraperitoneal injection of 1% sodium pentobarbital

solution (25 mg/kg). Subsequently, a minor incision was made in the eyelid using a

30-gauge needle to facilitate eye exposure. For intravitreal injections, a micropipette

was carefully inserted through the serosal opening, and formulations such as LNP-

mRNA or other substances were administered into the vitreous body of the eye. To

prevent the backflow of the injected solution, the needle was maintained in position

for approximately 10 seconds after the injection before being gently withdrawn. To

protect the cornea post-procedure, tobramycin was applied.

In vivo **NGF protein expression**

The m1Ψ-modified NGF mRNAs were injected into mouse retina and protein level

were measured after 48 hours. The detachment and processing of the mouse retina

were performed in the same way as mentioned in the section above. Then, the retinas

were collected in 300ul RIPA lysis buffer (Beyotime, P0013B) that contained 1%

PMSF (Sigma, 10837091001). Protein level was analyzed with western blot using

anti-NGF Antibody- BSA and Azide free (Abcam, #ab6199).

Optic nerve crush mouse model

Mice were anesthetized by intraperitoneal injection of 1% sodium pentobarbital

solution (25 mg/kg). Then, the eye surface was dilated with tropicamide drops and

surface anesthesia was provided with proparacaine hydrochloride. The mice were

fixed on the animal operating table. The optic nerve was completely exposed by

cutting open the bulbar fascia under the surgical microscope and using microforceps

to separate the surrounding tissues and hold the optic nerve for 5 s with a 0.07 mm

wide reverse forceps at 1 mm posterior to the globe in the vertical direction of the

longitudinal axis of the optic nerve. Tobramycin was applied daily to the superior

orbital rim incision for 3 days postoperatively. Five to nine biological replicates were

performed for each experiment.

***In vivo* immunogenicity**

First, $m^1\Psi$-modified mRNA with WT and RD1 codon sequences were encapsulated

within lipid nanoparticles (LNP). BALB/c mice were received two intramuscular

doses of 10μg mRNA each, with the dose determined by previous studies [77,78],

administered on day 0 (prime) and day 14 (boost). (Figure 5d). We collected mouse

serum and performed micro-neutralization (MN) assays to quantify neutralizing

antibodies at two time points: day 14 (for prime response) and day 28 (for boost

response) (Figure 5e, Methods). PBS buffer and the inactivated Split-virus

(A/Victoria/2570/2019 (H1N1)) Influenza Vaccine was used as negative and positive

control, respectively.

Micro-neutralization (MN) assay

To measure the titer of anti-influenza virus neutralizing antibodies, we treated mouse

serum with receptor-destroying enzyme □ (RDE □) (Denka-Seiken) at 37°C for 16

hours, followed by heat-inactivation at 56°C for 30 minutes. The experimental

procedure of MN was the same as previously reported[79]. MN titration was defined as

the reciprocal of the maximum serum dilution that neutralizes the 50% influenza

H1N1/PR8 virus infections in MDCK cells. The minimum MN titer was set to 10.

Eight biological replicates were performed for each experiment.

Ethics statement

For NGF mRNA *in vivo* assay, C57BL/6 mice were supplied by GemPharmatech. All

experimental procedures involving these mice were conducted in strict accordance

with the animal protocols that received approval from the Institutional Animal Care

and Use Committee (IACUC) at the Zhongshan Ophthalmic Center, Sun Yat-Sen

University, under the animal ethics approval number Z2021067. The *in vivo*

experiments used male C57BL/6 mice aged 8 weeks. For HA mRNA *in vivo* assay, all

the experimental procedures involving BALB/c mice were conducted in strict

accordance with the animal protocols that received approval from the Animal Care

and Use Committee Institutional at the Sun Yat-Sen University, under the animal

ethics approval number 2022002794. The *in vivo* experiments used male BALB/c

mice aged 6-8 weeks.

DATA AVAILABILITY

The datasets used for modeling and analysis are listed in Supplementary file 1. The

translation model and optimization framework of RiboDecode are available on

GitHub (https://github.com/wangfanfff/RiboDecode) and figshare

(https://figshare.com/articles/software/RiboDecode_zip/28916288?file=54138881).

The codon sequences optimized by RiboDecode and UTR sequences in this study are

listed in Supplementary file 2. Source Data are provided with this paper.

ACKNOWLEDGMENTS

We gratefully acknowledge the researchers who made their Ribo-seq and RNA-seq

data publicly available. We also extend our sincere appreciation to the reviewers for

their insightful comments and suggestions, which have significantly improved this

manuscript.

AUTHORS' CONTRIBUTIONS

Z.X. conceived, designed and supervised the project. YP.L. collected and

preprocessed the datasets, evaluated the model performance and analyzed the data. Y.

He, F.W., and ZH.T. developed the translation deep model. H.Z. developed the MFE

deep model. F.W. and H.Z. developed the AM framework. JX.D. tested the model.

JQ.Y. and LF.C. prepared mRNAs and conducted *in vitro* experiments. WB.J.

conducted *in vivo* NGF experiments. ZR.H. and CJ.S. conducted the *in vivo* HA

experiments. GF.Z encapsulated mRNA with LNP. X.H., H.J., Y. Hu, LP.W., and CY.Z.

contributed to project discussion. YP.L., Z.X., Y. He and F.W. wrote the manuscript.

All authors read and approved the manuscript.

FUNDING

This project was supported by the National Key R&D Program of China (Grant No.

2022YFF1203100, Y.H.); National Natural Science Foundation of China (Grant No.

32470705, Z.X.); Science and Technology Program of Guangzhou, China (Grant No.

2025A03J3990, Z.X.).

COMPETING INTERESTS

The authors declare that they have no competing interests.

REFERENCES

1.   Baden, L. R. *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2

Vaccine. *N Engl J Med* **384**, 403–416 (2021).

2.   Gebre, M. S. *et al.* Optimization of non-coding regions for a non-modified

mRNA COVID-19 vaccine. *Nature* **601**, 410–414 (2022).

3.   Pardi, N., Hogan, M. J., Porter, F. W. & Weissman, D. mRNA vaccines —

a new era in vaccinology. *Nat Rev Drug Discov* **17**, 261–279 (2018).

4.   Qin, S. *et al.* mRNA-based therapeutics: powerful and versatile tools to

combat diseases. *Signal Transduct Target Ther* **7**, 166 (2022).

5.   Fang, E. *et al.* Advances in COVID-19 mRNA vaccine development. *Sig

Transduct Target Ther* **7**, 94 (2022).

6.   Zhang, H. *et al.* Algorithm for optimized mRNA design improves stability

and immunogenicity. *Nature* **621**, 396–403 (2023).

7.   Leppek, K. *et al.* Combinatorial optimization of mRNA structure, stability,

and translation for RNA-based therapeutics. *Nat Commun* **13**, 1536 (2022).

8.   Hanson, G. & Coller, J. Codon optimality, bias and usage in translation

and mRNA decay. *Nat Rev Mol Cell Biol* **19**, 20–30 (2018).

9.   Mauger, D. M. *et al.* mRNA structure regulates protein expression through

changes in functional half-life. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 24075–24083

(2019).

10. Wayment-Steele, H. K. *et al.* Theoretical basis for stabilizing messenger
RNA through secondary structure design. *Nucleic Acids Res* **49**, 10604–
10617 (2021).

11. Hedaya, O. M. *et al.* Secondary structures that regulate mRNA translation
provide insights for ASO-mediated modulation of cardiac hypertrophy. *Nat*
Commun **14**, 6166 (2023).

12. Sharp, P. M. & Li, W. H. The codon Adaptation Index--a measure of
directional synonymous codon usage bias, and its potential applications.
Nucleic Acids Res **15**, 1281–1295 (1987).

13. Kudla, G., Lipinski, L., Caffin, F., Helwak, A. & Zylicz, M. High guanine and
cytosine content increases mRNA levels in mammalian cells. *PLoS Biol* **4**,
e180 (2006).

14. Lu, P., Vogel, C., Wang, R., Yao, X. & Marcotte, E. M. Absolute protein
expression profiling estimates the relative contributions of transcriptional and
translational regulation. *Nat Biotechnol* **25**, 117–124 (2007).

15. Li, G.-W., Burkhardt, D., Gross, C. & Weissman, J. S. Quantifying
absolute protein synthesis rates reveals principles underlying allocation of
cellular resources. *Cell* **157**, 624–635 (2014).

16. Waldman, Y. Y., Tuller, T., Shlomi, T., Sharan, R. & Ruppin, E. Translation
efficiency in humans: tissue specificity, global optimization and differences
between developmental stages. *Nucleic Acids Research* **38**, 2964–2974
(2010).

17. Fabbri, L., Chakraborty, A., Robert, C. & Vagner, S. The plasticity of
mRNA translation during cancer progression and therapy resistance. *Nat Rev*
Cancer **21**, 558–577 (2021).

18. Jumper, J. *et al.* Highly accurate protein structure prediction with
AlphaFold. **596**, 583–589 (2021).

19. Cao, C. *et al.* Deep Learning and Its Applications in Biomedicine.
Genomics, Proteomics & Bioinformatics **16**, 17–32 (2018).

20. Zrimec, J. *et al.* Deep learning suggests that gene expression is encoded
in all parts of a co-evolving interacting gene regulatory structure. *Nat Commun*
11, 6141 (2020).

21. Dauparas, J. *et al.* Robust deep learning–based protein sequence design
using ProteinMPNN. *Science* **378**, 49–56 (2022).

22. Sumida, K. H. *et al.* Improving Protein Expression, Stability, and Function
with ProteinMPNN. *J. Am. Chem. Soc.* **146**, 2054–2061 (2024).

23. Bennett, N. R. *et al.* Improving de novo protein binder design with deep
learning. *Nat Commun* **14**, 2625 (2023).

24. Li, S. *et al.* CodonBERT: Large Language Models for mRNA design and
optimization.

25. Richard, G. *et al.* ChatNT: A Multimodal Conversational Agent for DNA,
RNA and Protein Tasks. Preprint at https://doi.org/10.1101/2024.04.30.591835
(2024).

26. Melnikov, A. *et al.* Systematic dissection and optimization of inducible
enhancers in human cells using a massively parallel reporter assay. *Nat*
Biotechnol **30**, 271–277 (2012).

27. Ingolia, N. T., Ghaemmaghami, S., Newman, J. R. S. & Weissman, J. S.
Genome-wide analysis in vivo of translation with nucleotide resolution using
ribosome profiling. *Science* **324**, 218–223 (2009).

28. Weinberg, D. E. *et al.* Improved Ribosome-Footprint and mRNA
Measurements Provide Insights into Dynamics and Regulation of Yeast
Translation. *Cell Rep* **14**, 1787–1799 (2016).

29. Liu, T.-Y. & Song, Y. S. Prediction of ribosome footprint profile shapes from
transcript sequences.

30. Shao, B. *et al.* Riboformer: a deep learning framework for predicting
context-dependent translation dynamics. *Nat Commun* **15**, 2011 (2024).

31. Clauwaert, J. *et al.* Deep learning to decode sites of RNA translation in
normal and cancerous tissues. *bioRxiv* 2024.03.21.586110 (2024)
doi:10.1101/2024.03.21.586110.

32. Linder, J., Srivastava, D., Yuan, H., Agarwal, V. & Kelley, D. R. Predicting
RNA-seq coverage from DNA sequence as a unifying model of gene
regulation. *Nat Genet* (2025) doi:10.1038/s41588-024-02053-6.

33. Xie, S.-Q. *et al.* RPFdb: a database for genome wide information of
translated mRNA generated from ribosome profiling. *Nucleic Acids Res* **44**,
D254-258 (2016).

34. Wang, H. *et al.* RPFdb v2.0: an updated database for genome-wide
information of translated mRNA generated from ribosome profiling. *Nucleic*
Acids Research **47**, D230–D234 (2019).

35. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**, 26
(2011).

36. Huang, L. *et al.* LinearFold: linear-time approximate RNA folding by 5'-to-3'
dynamic programming and beam search. *Bioinformatics* **35**, i295–i304 (2019).

37. Linder, J. & Seelig, G. Fast activation maximization for molecular
sequence design. *BMC Bioinformatics* **22**, 510 (2021).

38. Wright, F. The 'effective number of codons' used in a gene. *Gene* **87**, 23–
29 (1990).

39. Chen, R. *et al.* Engineering circular RNA for enhanced protein production.
Nat Biotechnol **41**, 262–272 (2023).

40. Molinari, N.-A. M. *et al.* The annual impact of seasonal influenza in the US:
measuring disease burden and costs. *Vaccine* **25**, 5086–5096 (2007).

41. Myers, M. L. *et al.* Commercial influenza vaccines vary in HA-complex
structure and in induction of cross-reactive HA antibodies. *Nat Commun* **14**,
1763 (2023).

42. Lambiase, A. *et al.* Experimental and clinical evidence of neuroprotection
by nerve growth factor eye drops: Implications for glaucoma. *Proc Natl Acad*
Sci U S A **106**, 13469–13474 (2009).

43. Jiang, W. *et al.* Circular RNA-based therapy provides sustained and
robust neuroprotection for retinal ganglion cells. *Molecular Therapy - Nucleic*
Acids **35**, 102258 (2024).

44. Schwanhäusser, B. *et al.* Global quantification of mammalian gene
expression control. *Nature* **473**, 337–342 (2011).

45. Chaudhary, N., Weissman, D. & Whitehead, K. A. mRNA vaccines for
infectious diseases: principles, delivery and clinical translation. *Nat Rev Drug*
Discov **20**, 817–838 (2021).

46. Wang, Y., Tang, Y., Xie, Z. & Wang, H. RPFdb v3.0: an enhanced
repository for ribosome profiling data and related content. *Nucleic Acids Res*
53, D293–D298 (2025).

47. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize
analysis results for multiple tools and samples in a single report.
Bioinformatics **32**, 3047–3048 (2016).

48. Martin, M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet j.* **17**, 10 (2011).

49. Zerbino, D. R. *et al.* Ensembl 2018. *Nucleic Acids Res* **46**, D754–D761
(2018).

50. Casper, J. *et al.* The UCSC Genome Browser database: 2018 update.
Nucleic Acids Res **46**, D762–D769 (2018).

51. Johnstone, T. G., Bazzini, A. A. & Giraldez, A. J. Upstream ORFs are
prevalent translational repressors in vertebrates. *EMBO J* **35**, 706–723 (2016).

52. Joshi NA & Fass JN. Sickle: A sliding-window, adaptive, quality-based
trimming tool for FastQ files. (2011).

53. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie
2. *Nat Methods* **9**, 357–359 (2012).

54. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29, 15–21 (2013).

55. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general
purpose program for assigning sequence reads to genomic features.
Bioinformatics **30**, 923–930 (2014).

56. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-
Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323
(2011).

57. Jia, L. & Qian, S.-B. Therapeutic mRNA Engineering from Head to Tail.
Acc Chem Res **54**, 4272–4282 (2021).

58. Ho, J. J. D. *et al.* A network of RNA-binding proteins controls translation
efficiency to activate anaerobic metabolism. *Nat Commun* **11**, 2677 (2020).

59. Luan, Y. *et al.* Deficiency of ribosomal proteins reshapes the
transcriptional and translational landscape in human cells. *Nucleic Acids*
Research **50**, 6601–6617 (2022).

60. Liu, Y., Beyer, A. & Aebersold, R. On the Dependency of Cellular Protein
Levels on mRNA Abundance. *Cell* **165**, 535–550 (2016).

61. Schwanhäusser, B. *et al.* Global quantification of mammalian gene
expression control. *Nature* **473**, 337–342 (2011).

62. Zrimec, J., Buric, F., Kokina, M., Garcia, V. & Zelezniak, A. Learning the
Regulatory Code of Gene Expression. *Front. Mol. Biosci.* **8**, 673363 (2021).

63. Kelley, D. R., Snoek, J. & Rinn, J. L. Basset: learning the regulatory code
of the accessible genome with deep convolutional neural networks. *Genome*
Res. **26**, 990–999 (2016).

64. Ioffe, S. & Szegedy, C. Batch Normalization: Accelerating Deep Network
Training by Reducing Internal Covariate Shift. in *International conference on*
machine learning (2015).

65. Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I. & Salakhutdinov, R.
Dropout: A Simple Way to Prevent Neural Networks from Overfitting. *The*
journal of machine learning research **15**, 1929–1958 (2014).

66. Krizhevsky, A., Sutskever, I. & Hinton, G. E. ImageNet classification with
deep convolutional neural networks. *Commun. ACM* **60**, 84–90 (2017).

67. Loshchilov, I. & Hutter, F. Decoupled Weight Decay Regularization.
Preprint at http://arxiv.org/abs/1711.05101 (2019).

68. Nair, V. & Hinton, G. E. Rectified Linear Units Improve Restricted
Boltzmann Machines. in *Proceedings of the 27th international conference on*
machine learning (ICML-10) (2010).

69. He, K., Zhang, X., Ren, S. & Sun, J. Deep Residual Learning for Image
Recognition. in *2016 IEEE Conference on Computer Vision and Pattern*
Recognition (CVPR) 770–778 (2016).

70. Miyato, T., Dai, A. M. & Goodfellow, I. Adversarial Training Methods for
Semi-Supervised Text Classification. Preprint at
http://arxiv.org/abs/1605.07725 (2021).

71. Maas, A. L., Hannun, A. Y. & Ng, A. Y. Rectifier nonlinearities improve
neural network acoustic models. in *Proc. icml* (2013).

72. Krueger, R. & Ward, M. Scalable Differentiable Folding for mRNA Design.
Preprint at https://doi.org/10.1101/2024.05.29.594436 (2024).

73. Sundararajan, M., Taly, A. & Yan, Q. Axiomatic Attribution for Deep
Networks. in *International conference on machine learning* 3319–3328 (2017).

74. Bach, S. *et al.* On Pixel-Wise Explanations for Non-Linear Classifier
Decisions by Layer-Wise Relevance Propagation. *PLoS ONE* **10**, e0130140
(2015).

75. Shrikumar, A., Greenside, P. & Kundaje, A. Learning Important Features
Through Propagating Activation Differences. in *International conference on*
machine learning 3145–3153 (2017).

76. Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in
microarray expression data using empirical Bayes methods. *Biostatistics* **8**,
118–127 (2007).

77. Pardi, N. *et al.* Nucleoside-modified mRNA immunization elicits influenza
virus hemagglutinin stalk-specific antibodies. *Nat Commun* **9**, 3361 (2018).

78. Zhuang, X. *et al.* mRNA Vaccines Encoding the HA Protein of Influenza A
H1N1 Virus Delivered by Cationic Lipid Nanoparticles Induce Protective
Immune Responses in Mice. *Vaccines (Basel)* **8**, 123 (2020).

79. Wang, Y. *et al.* L226Q Mutation on Influenza H7N9 Virus Hemagglutinin
Increases Receptor-Binding Avidity and Leads to Biased Antigenicity
Evaluation. *J Virol* **94**, e00667-20 (2020).

1199
1200

1201

Figure 1. Predictive and Generative Optimization of RiboDecode.

a. RiboDecode contains three main components, a codon optimizer, a translation prediction model
and an MFE prediction model.

b. The framework of the prediction model for translation. The input includes codon sequences in
one-hot encoding, corresponding mRNA abundances and, the cellular environment which is
presented by vectors of gene expression profiles from RNA-seq. The model integrates these inputs
using CNNs. From the fused representations, a convolutional neural network extracts features and
outputs the predicted translation level of mRNA (see Methods).

c. Iterative optimization of codon sequences. RiboDecode predicts fitness of an original sequence
(T=0), then uses activation maximization to generate optimized synonymous variants (T+=1). A
synonymous regularizer maintains amino acid sequence. This process iterates until peak fitness is
achieved.

1214

Figure 2. Evaluation of the Translation Prediction Model.

a. Experimentally measured translation levels by Ribo-seq versus predicted translation levels in

the three validation datasets. Red lines denote the linear fit. The translation levels from Ribo-seq

were ln-transformed (see Methods).

b. Ablation analysis shows the contributions of the three inputs to the prediction model. The table

below shows the ablation status of the inputs, with dots and crosses representing the presence and

absence of corresponding elements, respectively. The error bars denote standard deviation.

c. The importance of each nucleotide position for the translation prediction. The x-axis represents

the nucleotide position from the TSS (translation starting site). Integrated Gradients attribution

method was used to obtain the importance score for each nucleotide position (black dots). The red

line denotes the local polynomial regression fit.

1228

1229

1230

1231

Figure 3. Strategies of Enhanced Translation in Generated Sequences.

a. Generation of Gluc codon sequences with low translation level (the upper left area) and high

translation level (the upper right area) (w=0, Figure S5). T-SNE of codon sequences is shown.

Each dot represents one sequence, and the color represents the predicted translation level.

b. Generation and optimization of Gluc codon sequences using different w of 0, 0.5, 0.7, and 1.

Each dot represents one sequence, positioned in its predicted translation level (y-axis) and MFE

(x-axis). The position of the reference sequence (MF882921.1) is shown.

c. Codons that appeared more frequently in highly translated endogenous sequences were also

used more often in highly translated Gluc sequences generated by RiboDecode. "RD":

RiboDecode-generated sequences, "Endo.": endogenous genes. (***: $p<0.001$, t-test).

d. Changes of sequence features of optimized sequences compared to the unoptimized, for

different mRNAs. For each column (feature), cells in red represent that the values were higher in

optimized sequences than unoptimized ones while cells in blue represent the opposite. The

difference of ENC for INS shows no significance (the cell in gray). Abbreviations: GFP: green

fluorescent protein, Fluc: firefly luciferase, INS: insulin, VZV: varicella zoster virus glycoprotein

E, and HA: influenza A hemagglutinin.

1248

1249

Figure 4. Robustness of Optimization across Unmodified, Modified, and Circular mRNAs.

a. Protein expression of Gluc was measured by fluorescence intensity. RD sequences were

designed by RiboDecode, with *w* parameter indicated in parentheses. LD sequences were designed

by LinearDesign, with *λ* parameter indicated in parentheses. "RLU": relative light units. The error

bars denote standard deviation.

b. Relative expression of experimentally measured protein expression values of mRNA variants

designed by RiboDecode at 24 hours in different cells. The error bars denote standard deviation.

One-sided Wilcoxon test was used to calculate p-values shown in the figure.

c, d. Protein expression of generated Gluc codon variants in (c) the linear mRNA form with m^{1}Ψ

modification and (d) the circular mRNA form. The error bars denote standard deviation.

1260

1261

1262

1263

Figure 5. More Effective mRNA-based Influenza Vaccines Through Codon Optimization

a. Western blot analysis shows protein expression of the HA variants in HEK293T cells 24 hours

after transfection. RD sequences were designed by RiboDecode (*w* parameter were set to 0, 0, 0.7,

and 0.5 for RD1-4). LD sequences were designed by LinearDesign (λ parameter were set to 0 and

4 for LD1 and LD2). The expression values were quantified using GelAnalyzer.

b, c. Protein expression RD3 in (b) the linear mRNA form with $m^1\Psi$ modification and (c) the

circular mRNA form.

d. HA mRNA immunization and analysis: BALB/c mice were intramuscularly inoculated with two

doses (10μg mRNA for each dose) with an interval of two weeks. The mouse serum was collected

at 14 days and 28 days for MN assay.

e. Levels of neutralizing antibodies against influenza viruses after prime and boost vaccination.

IC50, half-maximal inhibitory concentration. PBS and split virus influenza vaccine (SV) were

used as the negative and positive controls, respectively. One-sided Wilcox test was used to

calculate p values shown in the figure. The error bars denote standard deviation.

1278

1279

1282 **Figure 6. Enhanced Protein Expression and Therapeutic Efficacy with Optimized NGF**
1283 **mRNA.**

1284 **a.** NGF protein expression in HEK293T cells, measured by ELISA. RD sequences were designed
1285 by RiboDecode, with *w* parameter indicated in parentheses. LD sequences were designed by
1286 LinearDesign. The error bars denote standard deviation.

1287 **b, c.** Protein expression levels of RD3 in $m^1\Psi$-modified (b) and circular (c) mRNA formats.

1288 **d.** *In vivo* protein expression of RD3. The $m^1\Psi$-modified mRNAs were injected into mouse retinas,
1289 and protein levels were measured by western blot 48 hours post-injection. The expression values
1290 were quantified using GelAnalyzer.

e. Timeline of NGF mRNA therapy in ONC model: At Day 0 (D0), the m¹Ψ-modified NGF
mRNAs were injected into mouse retinas. At Day 4 (D4), the optic nerve was subjected to a
physical crush injury. At Day 18 (D18), RGC numbers were quantified using immunofluorescence
staining.

f, g. RGC numbers measured by immunofluorescence staining in the ONC mouse were
significantly reduced compared to that of the control (one-sided Wilcox test). Scale bar, 50μm.

h. RGC numbers in the ONC mouse retina after injection of NGF $m^1\Psi$-modified mRNA with 100
and 500 ng/μl dosages. Scale bar, 50μm.

i. The RGC number in mice treated with 100 ng/μl RD3 was similar to those treated with 500
ng/μl WT and significantly higher than those treated with 100 ng/μl WT. One-sided Wilcoxon test
was used to calculate p-values shown in the figure; ns: $p>0.05$.
1302
1303