1 **An RNA binding regulatory cascade controls the switch from proliferation to**

2 **differentiation in the *Drosophila* male germ cell lineage**

Devon E. Harris[1], Jongmin J. Kim[2,3], Sarah R. Stern[1], Hannah M. Vicars[1], Neuza R. Matias[1],

Lorenzo Gallicchio[1], Catherine C. Baker[1], and Margaret T. Fuller[1,4,#]

Author Affiliations:

[1]Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA

94305, USA

[2]Department of Chemical and Systems Biology, Stanford University School of Medicine,

Stanford, CA 94305, USA

[3]Current address: Department of Biomedical Sciences, Cornell University, Ithaca NY, 14853,

USA

[4]Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305, USA

[#]Corresponding Author (mtfuller@stanford.edu)

Short title: Bam represses *how* to set the switch from mitosis to male meiosis

Key words: differentiation, proliferation, spermatogenesis, RNA-binding proteins, adult stem

cell lineage

Abstract

The switch from precursor cell proliferation to onset of differentiation in adult stem cell

lineages must be carefully regulated to produce sufficient progeny to maintain and repair tissues,

yet prevent overproliferation that may enable oncogenesis. In the *Drosophila* male germ cell

lineage, spermatogonia produced by germ line stem cells undergo a limited number of transit

amplifying mitotic divisions before switching to the spermatocyte program that sets up meiosis

and eventual spermatid differentiation. The number of transit amplifying divisions is set by

accumulation of the *bag-of-marbles* (Bam) protein to a critical threshold. In *bam* mutants,

spermatogonia proliferate through several extra rounds of mitosis then die without becoming

spermatocytes. Here we show that a key role of Bam for the mitosis to differentiation switch is

repressing expression of Held Out Wings (*how*), homolog of mammalian Quaking. Knockdown

of *how* in germ cells was sufficient to allow spermatogonia mutant for *bam* or its partner *benign*

gonial cell neoplasm (*bgcn*) to differentiate, while forced expression of nuclear-targeted How

protein in spermatogonia wild-type for *bam* resulted in continued proliferation at the expense of

differentiation. Our findings suggest that Bam targets *how* RNA for degradation by acting as an

adapter to recruit the CCR4-NOT deadenylation complex via binding its subunit, Caf40. As How

is itself an RNA binding protein with roles in RNA processing, our findings reveal that the

switch from proliferation to meiosis and differentiation in the *Drosophila* male germ line adult

stem cell lineage is regulated by a cascade of RNA-binding proteins.

45
46

Introduction

In a common feature of the adult stem cell lineages that build and repair tissues

throughout the body, relatively less differentiated precursor cells undergo a limited series of

transit amplifying divisions, which serve to increase the number of differentiated progeny

produced from a single adult stem cell division. Blood, skin, intestinal epithelia, and male germ

cell lineages all employ transit amplifying (TA) divisions to produce large numbers of

differentiated cells (1). In this context, the switch from proliferation to differentiation must be

carefully regulated. Premature switching can lead to inadequate differentiated cell replacement,

with defects in tissue maintenance or repair. Conversely, failure to switch resulting in

overproliferation of precursor cells may increase danger of accumulating oncogenic mutations

leading to cancer (2).

Genetic analysis of the *Drosophila* male germ line adult stem cell lineage has revealed

key players in the switch from the mitotic program of transit amplifying spermatogonia to onset

of meiosis and the spermatocyte gene expression program that sets up differentiation. Germ line

stem cells at the testis apical tip divide with an oriented spindle (2) so that one daughter remains

next to the apical hub and maintains stem cell identity, while the daughter displaced away from

the hub becomes a gonialblast, founding a clone of transit amplifying spermatogonia (Figure 1A)

(3). Because daughters of a single founding gonialblast divide in synchrony with incomplete

cytokinesis and are enclosed by a pair of somatic cyst cells, counting the number of germ cells in

each cyst can reveal the number of rounds of mitotic divisions that took place in the clone prior

to the switch to the spermatocyte differentiation program. The number of transit amplifying

mitotic divisions differs among *Drosophilid* species, so is under genetic control (4), with the

number in wild-type *Drosophila melanogaster* almost always exactly four (Figure 1A).

Mutational analysis has identified three interacting proteins - Bag of marbles (Bam),

Benign gonial cell neoplasm (Bgcn), and Tumorous testes (Tut), that are required for

spermatogonia to stop proliferating and become spermatocytes (5-8). While Bgcn and Tut have

RNA recognition motifs (8, 9), Bam does not contain any known RNA-binding domains. Bam

appears to form a bridge in a ternary complex, with sequences in the Bam N-terminal third

binding Tut and sequences in the Bam C-terminal third binding Bgcn (8). Bam protein

expression detected by immunofluorescence staining showed the protein present in 4-cell cysts,

higher in 8-cell cysts, and remaining high during premeiotic S phase, after which the protein

quickly disappeared (4). Gene dosage experiments indicated that the number of transit

amplifying divisions is set by the length of time it takes Bam protein to reach a critical threshold

(4). In flies with one mutant and one wild type allele of *bam*, spermatogonial cysts sometimes

undergo one or occasionally more rounds of mitotic divisions before switching to spermatocyte

differentiation. Conversely, in flies carrying a frameshift allele that deleted the Bam C-terminal

PEST sequence, Bam protein accumulated more rapidly in transit amplifying spermatogonia and

the cells often switched to spermatocyte state a round earlier, resulting in spermatocyte cysts

with 8 germ cells rather than the normal 16 (4).

Bam protein has been shown to directly bind the Caf40 subunit of the CCR4-NOT

deadenylation complex (CNOT9 in humans), suggesting that Bam may target RNAs for

degradation or translational repression by recruiting the CCR4-NOT complex (10). While some

targets of Bam in the male germ line are known, none of the genes identified so far appear to be

key regulators in the switch from mitosis to meiosis downstream Bam. Previous work showed

that Bgcn, Tut, and possibly Bam bind the 3'UTR of *mei-P26* and reduce expression of Mei-P26

protein. However, overexpression of *mei-P26* did not prevent the switch to spermatocytes, and

decreasing *mei-P26* did not allow the switch to occur, suggesting that Mei-P26 is not the key

target of Bam for the switch from mitosis to meiosis in males (8, 11).

Here we show that a key role of Bam and Bgcn in the switch from mitosis to meiosis in

the *Drosophila* male germ line is to repress expression of the RNA-binding protein Held Out

Wings (How), a homolog of mammalian Quaking, with roles in alternative RNA processing,

export, and translation. How protein is normally downregulated in germ cells soon after Bam

protein becomes expressed, but remained high in the TA spermatogonia that overproliferate in

bam mutant males. Strikingly, knocking down *how* expression in *bam* mutant spermatogonia by

RNAi allowed the germ cells to switch to spermatocyte state and differentiate. Reciprocally,

forced expression of nuclear targeted How blocked differentiation of otherwise wild-type

spermatogonia, even in the presence of Bam protein. Consistent with the model that Bam down

regulates *how* RNA by recruiting Caf40 and the CCR4-NOT RNA degradation complex,

knockdown of *Caf40* in early spermatogonia resulted in persistence of *how* RNA and

overproliferation of TA cells, even in the presence of Bam protein. How is an RNA binding

protein that regulates alternative RNA processing, export, and translation. Our findings reveal

that an irreversible cell fate transition from mitosis to terminal differentiation in an adult stem

cell lineage is controlled by a regulatory cascade of RNA-binding proteins.

Results

Downregulation of *how* is an early consequence of Bam action

Function of *bag-of-marbles* (*bam*) is required for proliferating spermatogonia to turn

down expression of Held Out Wings (How). In testes wild-type for *bam*, immunofluorescence

staining showed that How protein, present in the nucleus of early germ cells, is downregulated in

mid-stage transit amplifying spermatogonia, soon after the Bam protein was first detected by

immunofluorescence staining (Figure 1B), as previously documented by Monk *et al.* (2010) (12).

Some mid-stage spermatogonia up regulating Bam protein still showed staining for How protein

in their nuclei (Figure 1B, yellow dotted outline), while adjacent Bam positive cysts lacked

detectable How protein (Figure 1B, white dotted outline). How protein continued to be detected

in the nuclei of the somatic cyst cells that enclose spermatocyte cysts (Figure 1B, C,

arrowheads), as well as in the testis sheath, but was below the level of detection in germ cells by

early spermatocyte stages. In contrast, How protein persisted at high levels in the nuclei of the

spermatogonia that continued to overproliferate in *bam* mutant males (Figure 1D).

When *bam* mutant spermatogonia were induced to differentiate in response to a burst of

Bam expression under heat shock control, down regulation of *how* RNA was one of the earliest

responses detected both by RNA-Seq and independent microarray analysis of whole testes

Figures 1E; S1). In the heat shock Bam time course strategy developed by Kim *et al.*, (2017)

(13), when *bam; hs-Bam* males were shifted as late pupae to 37^{o}C for 30 minutes to induce

expression of Bam then returned to 25^{o}C, a wave of spermatogonia initiate differentiation,

complete mitosis and a final S phase by 24 hours post heat shock (PHS), and differentiate into

spermatocytes, with onset of expression of early spermatocyte-specific transcripts beginning by

24h PHS (Figure S1).

Analysis of PolyA+ RNAs expressed in whole testes at early time points in the heat

shock-Bam differentiation time course showed that downregulation of *how* RNA was among the

earliest changes detected. The level of *how* transcripts detected fell by > 2-fold by 8h PHS, long

before the germ cells began to express spermatocyte-specific markers. *how* was one of six genes

showing greater than 2-fold decrease in transcript level by 8h PHS by both RNA-sequencing and

microarray analysis (Figure 1E and Figure S1A). In the gene expression comparisons in Figures

1E and S1, only those genes that met the following criteria were colored blue (for

downregulated) or red (for upregulated): both RNA-Seq and independent microarray analysis of

testes from *hs-Bam;bam* flies subjected to heat shock showed transcript levels changed >2 fold

compared to the same genotype not subjected to heat shock, and testes from control *bam* mutant

flies lacking the *hs-bam* did not show >2 fold change expression by microarray analysis for the

same gene at the indicated times PHS compared to testes from *bam* flies not subjected to heat

shock (Figure S1). Transcripts from additional genes became downregulated > 2-fold by later

time points, with the number of genes with lowered transcripts growing from six at 8h PHS to 28

(16h), 83 (24h) and 114 by 32h PHS, by which time transcripts from a number of genes

expressed specifically in spermatocytes began to be detected (Figure S1D,E).

Reclustering of snRNA-seq data from early germ cells generated by Li *et al.,* (14)

showed that most early germ cell nuclei expressed either *how* RNA or *bam* RNA, while a small

subset of the nuclei were positive for both (Figure 1F, white dots, merge). Fluorescence *in situ*

hybridization (FISH) confirmed that levels of *how* RNA became down regulated as germ cells

differentiate into spermatocytes (Figure S2B-E) and abruptly decreased in early germ cells soon

after onset of Bam protein expression in wild-type testes (Figures 1G; S2C-E). In testes carrying

a *Bam-GFP* transgene to allow visualization of Bam protein expression in mid to late transit

amplifying spermatogonia, Hybridization Chain Reaction (HCR) FISH with probes recognizing

the How protein coding sequence (Table 1) detected *how* transcripts in the nucleus and

cytoplasm of cells near the testis tip, apical to the region where germ cells expressing Bam-GFP

were detected (Figures 1G; S2C-E). *how in situ* signal was apparent in the cytoplasm of germ

cells in some early spermatogonial cysts expressing Bam-GFP (yellow outline), while other Bam

positive cysts showed little or no signal for *how* RNA (white and cyan outlines, respectively)

(Figures 1G; S2C-E), suggesting that the level of *how* RNA in spermatogonia decreases rapidly

once Bam protein becomes expressed. HCR FISH with a different probe set, directed against the

3'UTR associated with *how* mRNAs that encode nuclear targeted How(L) protein isoforms

(Table 2) also detected *how* RNA signal at the testis apical tip that decreased with distance from

the testis hub, like signal from the *how* protein coding region probe correlating with progression

of transit amplifying spermatogonia to more mature stages (Figure S2B).

***how* is a key target of Bam for the proliferation to differentiation switch**

Knocking down expression of *how* in *bam* mutant TA spermatogonia by RNAi under

control of *bam-Gal4* allowed *bam* mutant spermatogonia to differentiate into spermatocytes.

Whereas squashed preparations of control testes wild-type for *bam* viewed by phase contrast

microscopy showed abundant large spermatocytes and elongating spermatid bundles (Figure

2A), testes from males mutant for *bam* were considerably smaller, lacked spermatocytes and

spermatids and contained large numbers of small germ cells that proliferate in cysts then

eventually die (Figure 2B). However, if the *bam* mutant flies also carried a *UAS-how-RNAi*

construct forcibly expressed in transit amplifying spermatogonia under control of *bamGal4*, the

bam$^{-/-}$ germ cells successfully differentiated into spermatocytes and elongating spermatids

(Figures 2C; S3), indicating that the main requirement for Bam for the switch from mitosis to

meiosis in males is reducing function of How. Similar rescue allowing differentiation of *bam*

mutant germ cells to spermatocytes and elongating spermatids resulted from knockdown of

expression of *how* by RNAi using a non overlapping shRNA construct from the TRiP collection

(HMC03820) (Figure S3A-F). The *bam* mutant males in which *how* had been knocked down by

bamGal4 induced RNAi were fertile and produced viable offspring, similar to control males,

although in both cases the number of progeny was quite low, possibly due to culturing the males

at 29˚C (Figure S3J).

Testes from males mutant for *bgcn*, a binding partner of *bam*, had the same phenotype as

bam mutants, overproliferation of transit amplifying spermatogonia that maintained expression

of nuclear How protein (Figure S4A), and eventual germ cell death, with no cells at later stages

of development (Figure S4B). Knockdown of *how* function in mid-to-late transit amplifying

spermatogonia by RNAi under control of *bamGal4* also restored ability of *bgcn* mutant

spermatogonia to differentiate into spermatocytes and elongating spermatids (Figure S4C).

Brief incubation of testes in EdU to label cells in S phase and immunofluorescence

staining for the spermatocyte marker Kmg confirmed that knockdown of *how* by RNAi in mid-

to-late spermatogonia under control of *bamGal4* restored the ability of *bam* mutant

spermatogonia to switch to the spermatocyte state. Testes from control or RNAi knockdown flies

were incubated in EdU for 5 minutes to label nuclei undergoing DNA replication, then

immediately fixed and processed for immunofluorescence staining and imaging. Control testes

from flies carrying the *bamGal4* transgene and raised under the same temperature regimen used

for RNAi knockdowns showed a few small clusters of EdU-positive nuclei located near the testis

apical tip, marking cysts of spermatogonia undergoing S phase in synchrony. As expected, no

EdU positive nuclei were identified further from the tip, where germ cells showed expression of

the spermatocyte specific marker Kmg (Figure 2D). In contrast, testes from flies mutant for *bam*

raised under the RNAi knockdown temperature shift regime showed many more EdU-positive

cysts per testis, several with more than 16 EdU-positive nuclei, indicating spermatogonial

overproliferation, and no spermatocytes expressing Kmg (Figure 2E). Testes from *bam* mutant

males in which expression of *how* was knocked down in late spermatogonia and early

spermatocytes by RNAi under control of *bamGal4* showed many fewer EdU-positive cysts per

testis, again confined to near the testis apical tip, and abundant Kmg-positive spermatocytes

(Figure 2F). Similar results were observed when expression of How was knocked down by a

different RNAi construct expressed under control of *bamGal4* (Figure S3D-F).

Notably, testes from *bam*$^{-/-}$; *bamGal4;UAS-How RNAi* males raised under knockdown

conditions were unusually wide, with many early germ cells (Figures 2C, F; S3I). The increased

width of *bam*$^{-/-}$; *bamGal4;UAS-How RNAi* testes was likely because downregulation of *how* due

to RNAi under control of *bamGal4* may occur later than in 4-8 cell cysts as in wild type, because

of the time it takes for the *bamGal4* to be expressed, activate transcription of the RNAi construct,

then for the RNAi to act upon *how* transcripts. Such delay would allow the *bam* mutant

spermatogonia to go through additional rounds of mitotic transit amplifying divisions before

downregulation of How. Consistent with this, spilling out individual cysts revealed that control

testes raised under the knockdown temperature regimen almost always had 16 spermatocytes per

cyst. In contrast, *bam*$^{-/-}$; *bamGal4;UAS-How RNAi* testes normally had 32, 64, and sometimes

more spermatocytes per cyst, indicating five, six, or more rounds of transit amplifying divisions

prior to the switch to spermatocyte state, rather than the normal four (Figure 2G-I).

Conversely, forced expression of nuclear-targeted How but not cytoplasmic How in mid-

stage spermatogonia was sufficient to largely block differentiation of otherwise wild-type

spermatogonia into spermatocytes. The *how* locus encodes several transcript and protein

isoforms (Figure S2A) (15). Some transcripts encode long forms of How protein that have a C-

terminal nuclear localization signal (How(L)). Others encode shorter forms of the protein that

lack the nuclear localization signal (How(S)) and are cytoplasmic.

Flies bearing a transgene with UAS sequences controlling inducible expression of

How(L) protein (isoform A in Figure S2A) cloned in frame with a C-terminal HA epitope tag

followed by terminator sequences from SV40 (*UAS-How(L)HA-SV40*) were crossed to flies

bearing *bamGal4* to drive expression in mid-to-late transit amplifying spermatogonia. The mated

flies were cultured at 25°C for three days then adults were removed and the progeny shifted to

and maintained at 29°C to boost expression from the *UAS-How(L)HA-SV40* transgene. Under

these conditions, testes from the newly eclosed *bamGal4; UAS-How(L)-HA-SV4* progeny had

extensive apical regions filled with small germ cells (Figure 3B red bracket, compare to A),

followed by areas with dying cells, as in *bam* mutant testes. Some testes also had a few cysts

containing spermatocytes or post-meiotic spermatids, usually located far down the testis away

from the apical tip, distal to the region with cell death.

Immunofluorescence staining after a short pulse of EdU revealed that most (~80%) of the

bamGal4; UAS-How(L)HA SV40 testes had a much larger than normal number of cysts with

germ cells undergoing synchronous DNA replication per testis than controls (Figure 3E'). The

germ cell cysts subjected to forced expression of How(L) often had more than 16 nuclei

undergoing S phase, indicating overproliferation. In addition, the nuclei remained small and

stained brightly with DAPI, unlike in spermatocytes, and lacked expression of the spermatocyte

marker Kmg (Figure 3E'). Immunofluorescence staining with anti-HA confirmed that How(L)-

HA was localized to germ cell nuclei and that the *bamGal4* driver did not force expression of

How(L)HA in germ line stem cells and early spermatogonia (Figure 3H). Notably,

immunofluorescence staining with anti-Bam showed that Bam protein was expressed in the

spermatogonia that overproliferated when *How(L)HA-SV40* was forcibly expressed, indicating

that the failure of spermatogonia to differentiate was not due to repression of Bam by nuclear

targeted How(L) (Figure 3K).

In contrast, testes from males expressing *UAS-How(S)HA-SV40* (isoform B in Figure

S2A, which lacks the NLS) under control of *bamGal4* raised under the same conditions did not

show massive overproliferation of spermatogonia, but instead had a modest population of small

germ cells at the testis apical tip followed by plentiful differentiating spermatocytes, visible as

large cells with large nuclei in unfixed squashed preparations viewed by phase contrast

microscopy (Figure 3C). Immunofluorescence staining confirmed that the switch to

spermatocyte state occurred after a limited number of mitotic divisions when *How(S)HA* was

forcibly expressed in mid-to-late spermatogonia. Testes from *bamGal4; UAS-How(S)HA-SV40*

males subjected to a brief incubation in EdU showed only a small number of EDU positive cysts,

all of which were close to the apical tip. The testes also had many spermatocytes, marked by

large nuclei positive for Kmg protein, starting from within a few cell diameters of the testis

apical tip (Figure 3F'). Immunofluorescence staining with anti-HA confirmed that How(S)HA

was cytoplasmic (Figure 3I). As expected for the *bamGal4* driver, anti-HA was not detected in

very early germ cells at the testis tip. Together these data indicate that it is nuclear rather than

cytoplasmic forms of How protein that block ability of spermatogonia to differentiate into

spermatocytes.

Addition of 3'UTR sequences from *how(L)* mRNA reduced the effect of forced

expression of *How(L)HA* on ability of spermatogonia to stop proliferating and differentiate into

spermatocytes. *How(L)* mRNA isoforms have overlapping 1566nt - 2424nt 3'UTRs, which do

not share sequences with the much shorter 3'UTRs of mRNA isoforms that encode How(S)

proteins (15) (see Figure S2A). The 2424nt How(L) 3'UTR from *how* mRNA isoform A (Figure

S2A) was cloned into the *UAS-How(L)HA-SV40* overexpression construct between the HA tag

and the SV40 terminator to generate *UAS-How(L)HA-How(L)3'UTR-SV40* (Figure 4, top

diagram) and introduced into flies as a transgene (Materials and Methods).

Flies bearing either the newly constructed *UAS-How(L)HA-How(L)3'UTR-SV40*

transgene (hereafter termed *UAS-How(L)-3'UTR*) or a parallel *UAS-How(L)-SV40* transgene

lacking the *How(L) 3'UTR* (hereafter termed *UAS-How(L)-SV40*) inserted at the same genomic

site were then crossed to flies bearing *bamGal4* to drive expression starting in mid-to-late

spermatogonia and the progeny grown continuously at 18˚C. Under these conditions, forced

expression of *UAS-How(L)-SV40* had a range of phenotypes, visualized and scored by EdU

labeling and anti-Kmg staining. Some testes (37%) had no spermatocytes detected at all (Figure

4A), while the remaining 63% of testes contained at least some individual Kmg-positive

spermatocytes (Figure 4B), with 53% of the testes scored containing entire cysts of Kmg-positive

spermatocytes (n = 30 testes) (Figure 4C,G). However, in flies in which the *UAS-How(L)-3'UTR*

construct was forcibly expressed under control of *bamGal4* at 18˚C, 100% of testes had at least

some spermatocyte cysts (n = 29 testes) (Figure 4D-F,G). Briefly incubating testes in EdU to

label nuclei in S phase revealed that flies overexpressing How(L)-HA with and without the

How(L)3'UTR both contained spermatogonial cysts that had undergone additional rounds of

proliferation beyond the normal 4, visualized as EdU positive cysts with many more than 16

labeled nuclei (Figure 4A-F,H). However, *UAS-How(L)-3'UTR* testes had overall less

spermatogonial overproliferation, based on fewer cysts undergoing S phase per testis (Figure 4H)

and fewer cysts with more than 16 EdU labeled nuclei per cyst (Figure 4I), compared to testes

from flies overexpressing *UAS-How(L)-SV40* (without the How(L) 3'UTR) grown in parallel.

The milder effect of forced overexpression of *How(L)-3'UTR* compared to *How(L)-SV40* under

control of *bamGal4* raised the possibility that Bam may downregulate How expression via the

How(L) 3'UTR. □

Bam may downregulate *how* by recruiting the CCR4-NOT complex

The reduction of *how* transcript levels in transit amplifying spermatogonia soon after the

appearance of Bam protein in the cytoplasm (Figures 1G; S2C-E) suggested that Bam may

downregulate *how* at least in part due to effects on the *how* RNA. Protein structure studies by

Sgromo *et al.* (2018) showed that a 23 amino acid sequence near the N-terminus of the Bam

protein binds in a groove of Caf40, a subunit of the CCR4-NOT complex (10). Further, Sgromo

et al. showed that when Bam or an N-terminal fragment of Bam containing the Caf40-binding

motif (CBM) was tethered to a luciferase reporter RNA, it was able to recruit CCR4-NOT for

transcript degradation, decreasing levels of luciferase RNA and protein (10). These findings

raise the possibility that Bam and Bgcn may downregulate *how* by recruiting Caf40. Consistent

with this, knockdown of *Caf40* in early spermatogonia by RNAi under control of *nosGal4*

resulted in massive overproliferation of small germ cells, similar to the phenotype observed in

bam mutant males (Figures 5C; S5B-E). Digesting the testis sheath to spill out intact cysts

confirmed that the overproliferating small cells were organized in cysts (Figure 5D), as in *bam*

mutants. Brief incubation in EdU confirmed that large clusters of germ cells were undergoing

DNA synthesis in synchrony in testes in which expression of *Caf40* had been knocked down

under control of *nosGal4*, while immunofluorescence staining confirmed failure to turn on

expression of the spermatocyte marker Kmg (Figure 5F), as in *bam* mutant males. Strikingly,

immunofluorescence staining with anti-Bam revealed that the hundreds of small germ cells that

overproliferated when *Caf40* was knocked down by RNAi under control of *nosGal4* expressed

Bam protein (Figure 5H), indicating that they were germ cells and that the failure to differentiate

was not due to failure to express Bam.

Analysis by fluorescence HCR *in situ* hybridization (FISH) confirmed that *how*

transcripts remained in the germ cells that overproliferated after function of the CCR4-NOT

component *Caf40* was knocked down by RNAi. In testes from control males carrying the

nosGal4 driver but not the *UAS-Caf40 RNAi* construct (Figure 5I) raised under RNAi

temperature shift conditions, *how* mRNA was detected in early germ cells near the testis apical

tip but was not detected in the spermatocyte region further down the testes (as in Figure 1).

However, despite the expression of Bam protein, *how* transcripts remained high in the

overproliferating germ cells that accumulated in testes in which *Caf40* had been knocked down

in early germ cells under control of *nosGal4*, as observed in *bam$^{-/-}$* males (Figure 5J,K). The

persistence of *how* transcripts despite the presence of Bam protein suggested that Caf40 does not

act upstream of Bam but instead might be part of the mechanism through which Bam protein

downregulates *how* RNA.

Discussion

A key function of Bam in spermatogonia is repression of *how*

Our results indicate that the major role of Bam and Bgcn in the switch from mitotic proliferation to onset of the meiotic program in the male germ line is to down regulate expression of Held Out Wings, the *Drosophila* homolog of mammalian Quaking (Figure 6A). Most telling, male germ cells lacking *bam* function can differentiate if expression of *how* is knocked down in mid-to-late stage transit amplifying spermatogonia by RNAi. In addition, forced expression of a nuclear targeted isoform of How, How(L), was sufficient to cause spermatogonia to overproliferate and largely fail to become spermatocytes. Strikingly, spermatogonia subjected to forced expression of How(L) expressed abundant Bam protein, indicating that the failure to differentiate was not due to How(L) repressing expression of Bam.

Our finding that knockdown of *Caf40* by RNAi driven by *nosGal4* led to overproliferation of spermatogonial cysts and persistence of *how* RNA and protein, even though Bam protein was present, suggested that *Caf40* is required for Bam to downregulate *how*. Sgromo *et al.* showed that Bam protein has a 23-amino acid N-terminal domain that binds in a groove of *Caf40* (homolog of human NOT9), a subunit of the CCR4-NOT complex (10). Through this Caf40-binding motif (CBM) domain, Bam tethered to a reporter RNA was able to recruit the CCR4-NOT complex to degrade the reporter RNA (10). Bam protein appears to participate in a ternary complex, bridging between the RNA binding protein Tut bound to sequences in the Bam N-terminal third and the RNA binding protein Bgcn bound to sequences in the Bam C-terminal third (8). Consistent with models suggested by others, we propose that Bam, recruited to target RNAs by its structural partners Bgcn and Tut, may act as an adaptor to recruit

the CCR4-NOT deadenylation complex to destabilize the bound RNA and/or lead to its

translational repression (8, 10, 16) (Figure 6B). In support of the model that Bam targets *how*

RNA for degradation and/or translational repression by recruiting CCR4-NOT, knockdown of

several CCR4-NOT subunits by RNAi under control of the *bamGal4* driver showed a mild

overproliferation phenotype in spermatogonial cysts (16).

Bam and Bgcn have been shown to downregulate expression of E-Cadherin via

sequences in the E-Cadherin 3'UTR (17). Assays conducted in S2 cells carrying reporter

constructs with the Firefly *luciferase* coding sequence attached to the 3'UTR from E-Cadherin

showed that expression of luciferase was down regulated compared to a renilla control with a

heterologous 3'UTR if Bam and Bgcn were co-expressed in the cells. The inhibition of luciferase

expression was not observed when either Bam or Bgcn was expressed without its partner, but

was detected if Bam was tethered to the E-Cadherin 3'UTR. Bgcn, Bam and Tut have also been

shown to repress expression of Mei-P26 via sequences in the *mei-P26* 3'UTR (8, 11).

Bam serves as a differentiation factor in both the male and female germ lines (6, 7). The

regulatory logic is similar: in both cases, action of *bam* downregulates a factor required for

maintenance of the precursor state. In *Drosophila* females, the germ line stem cell state is

maintained by *nanos* and *pumilio*, which are thought to repress translation of transcripts that

drive differentiation (7, 18-20). Loss of function of *nanos* or *pumilio* in females resulted in loss

of germ line stem cells to differentiation (19). In female early germ cells, expression of *nanos*

protein was abruptly downregulated when Bam protein became expressed in late cystoblasts.

Nanos protein expression persisted in the oogonia that overproliferate in *bam* mutant ovaries, and

the down regulation of *nanos* in response to Bam depended on sequences in the *nanos* 3'UTR (7,

21). Similarly, in males, How protein is downregulated when Bam becomes expressed in mid

stage transit amplifying spermatogonia, How protein persisted in spermatogonia mutant for *bam*,

and addition of the How(L) 3'UTR somewhat reduced the effects of forced expression of

How(L).

The timing of Bam expression and action is different in male than in female germ cells.

In females, Bam protein becomes upregulated enough to be detected by immunofluorescence

staining in the late cystoblast, the immediate daughter of a stem cell division, which will found a

clone of female germ cells that will eventually generate one oocyte and fifteen nurse cells. In

males, Bam protein is upregulated later, during the spermatogonial transit amplifying divisions,

so that it was initially detected by immunofluorescence staining in four cell cysts, appeared to

increase in level by the 8 cell stage, and remained present through pre-meiotic S phase before

being abruptly degraded (4). In both sexes, expression of How protein in germ cells appears

reciprocal to expression of Bam. In females, How protein was present in the nucleus of female

germ line stem cells but was abruptly down regulated by the two cell stage (22). Male germ line

stem cells also showed nuclear How, which persisted in gonialblasts and two cell transit

amplifying stages but dropped precipitously once Bam protein was expressed and began to

accumulate (12, this study).

In males, knockdown of *how* in germ line stem cells and early transit amplifying cells

under control of *nosGal4* led to loss of germ line stem cells, as did induction of germ line clones

homozygous for a strong loss of function *how* allele (12). Germ cells homozygous mutant for

how appeared stalled in G2 at the two cell cyst stage, likely due to defects in expression of

Cyclin B, and eventually died (12). The effect of loss of How function in female germ line stem

cells was different: induction of germ line clones mutant for *how* led to gradual loss of female

germ line stem cells, possibly to differentiation, but not to cell cycle arrest and apoptosis (22).

These results suggest that although the regulatory relationship between How and Bam may be

similar, the roles of How (and Bam) in early germ cells in the two sexes likely differs.

408   Monk *et al.*, 2010 (12) observed that using *nosGal4* to drive overexpression of *How(L)-*

SV40 in early germ cells (approximately the time when How is normally expressed) caused

shortening of cell cycle time in TA spermatogonia, likely due to increased expression of Cyclin

B, and frequent 32 cell spermatocyte cysts, indicating switching from spermatogonia to

spermatocytes after 5 rather than the usual 4 rounds of transit amplifying mitotic divisions.

Monk *et al.*, 2010 proposed that How(L) represses expression of Bam protein in early

spermatogonia (12). In contrast, we found that using *bamGal4* to drive forced expression of

How(L)-SV40 later, a time when it is normally downregulated, resulted in massive

overproliferation of spermatogonia. Strikingly, Bam protein was expressed in the spermatogonia

overproliferating under these conditions, arguing that How(L) does not repress expression of

Bam in later stage spermatogonia.

419   Our data suggest that Bam down regulates expression of How in mid-to-late transit

amplifying spermatogonia, possibly through the How(L) 3'UTR, which was not included in the

How(L)-SV40 construct. Such a mutual repression relationship may provide a mechanism to

convert a gradual rise in Bam expression (perhaps driven at the level of transcription) into a

sharp switch in cell state. In early transit amplifying male germ cells, abundant How(L) protein

may repress premature accumulation of Bam protein and maintain a stem cell competent, early

TA state. As expression *of bam* transcript and protein rise, however, Bam may reach levels

sufficient to bind via a Tut-Bam-Bgcn complex to all or most of the How(L) mRNA molecules,

targeting them for translational repression and/or degradation and so throwing the switch to onset

of differentiation. Alternatively, the 32 spermatocyte cell cysts detected by Monk *et al.,* (2010)

could be due to the observed shortening of the cell cycle in spermatogonia after forced

expression of How(L) under control of *nosGal4* (12) rather than to How(L) repression of *bam* in

early spermatogonia. As shown by Insco *et al.* (2009), shortening the cell cycle can allow more

cysts to complete 5 rounds of cell division before Bam protein reaches the critical level required

for the switch to spermatocyte state (4).

The fact that testes from *bamGal4>UAS-How(L)-3'UTR* males still showed some

aberrant overproliferation of spermatogonia may be because getting the proportions of *how* RNA

and Bam protein just right so that all the *how* RNA can be bound and repressed/degraded is not

easy. The amount of How(L)-3'UTR mRNA produced in the *bamGal4>UAS-How(L)-3'UTR*

males may titrate out the endogenous Bam protein and so overwhelm the capacity of Bam to

repress either the *UAS-How(L)* or endogenous *how* RNA.

A central lesson from genetic analysis of the *Drosophila* male and female germ line adult

stem cell lineages is that the switch from precursor cell state to onset of differentiation is

controlled by a cascade of RNA-binding proteins. In each case, Bam protein with its RNA-

binding partner Bgcn acts to trigger differentiation by downregulation of another RNA-binding

protein. In females, the key target is the translational inhibitor *nanos* (7). Here we show that in

males the key target is the RNA-binding protein How. It will be interesting in future work to test

whether mRNAs bound and regulated by How(L) in early spermatogonia maintain the germ cells

in a proliferative state or block their differentiation. Our results suggest that it will also be

important to investigate whether similar regulation by RNA-binding proteins controls the switch

from proliferation to differentiation in adult stem cell lineages in other organisms.

Methods

451
452   **Fly Strains and husbandry**

453
Flies were raised on molasses food. Overexpression and knockdown crosses were raised

at 25˚C for 3 days, then shifted to 29˚C after discarding adults, unless otherwise noted. RNAi

lines were obtained from The Vienna Drosophila Resource Center (VDRC): UAS *how* RNAi

(100775), UAS *caf40* RNAi (101462) and the Bloomington Drosophila Stock Center: UAS *how*

RNAi (Bloomington stock #55665, containing TRiP line HMC03820) and UAS *caf40* RNAi

(67987, containing TRiP line HMS05850). Results using the How RNAi line VDRC100775 are

shown throughout except for Figures 2I, S3, and S6, which also show results from knocking

down How using the TRiP RNA construct HMC03820 carried in Bloomington stock #55665.

For Figure 4, flies were raised at 18˚C throughout. For the heat shock time course, flies were

raised at 25˚C until there were mid- and late-stage pupa. Then bottles were placed in a 37˚C

water bath for 30 minutes to induce heat shock and returned to 25˚C, as spelled out in Kim *et al*.,

(2017) (13). For the RNA-Seq analysis shown in Figure S5, the *bam* mutant testes were from

CyO/+; bam^1^/bam^{Δ86} males that were grown at 22˚C until the late pupal stage, then the bottles

were incubated in a 37˚C water bath for 30min, then cultured at 25˚C for 48hrs, at which time the

testes plus seminal vesicle were obtained by dissection and prepped for RNA extraction. In all

other cases, flies were raised at 25˚C. *UAS-How(Long)-3xHA* and *UAS-How(Short)-3xHA* fly

stocks with an SV40 heterologous 3'UTR were a gift from T. Volk (23). Two Gal4 driver lines

were used: *nanosGal4-VP16* for germ line stem cells and early spermatogonia and *bamGal4* for

the mid- to late spermatogonia (24, 25). Gal4 driver strains for knockdowns also contained *UAS-*

Dicer2. The *bam* alleles were *bamP-bam-GFP* (25), *bam^1^* (5), and *bam^{Δ86}* (26). Fertility tests

were performed with males from crosses that had been shifted to 29˚C for Gal4 driver

expression. One male was placed in a vial with three virgin females and kept at 25°C. Adults

were removed after six days and vials were scored for the presence of pupa and adult offspring

after at least 10 days.

478
479   **Phase Contrast Microscopy**
480
For testis squashes, testes from 0-1 day old males were dissected in PBS. Whole testes or

cysts were flattened into a monolayer under a coverslip by wicking away PBS and observed by

phase contrast microscopy using a Zeiss Axioskop. Images were taken with a Spot Imaging

camera and software. To count the number of spermatocytes per cyst, dissected testes were

treated with 0.5 mg/mL collagenase (Sigma C7657) + 0.5 mg/mL dispase (Worthington,

LSO2109) in PBS on a slide for 1.5-3 minutes (larger testes burst open sooner) (as described in

Lu *et al*., 2020) (27). The reaction mix was replaced with PBS and cysts were gently flattened

under a coverslip by wicking away liquid before counting and imaging.

Transgenics

Plasmids containing *how* coding sequences with HA tags expressed under control of UAS

were a gift from Talila Volk (23). The *how*(L) coding sequence (FBgn0264491, isoform A on

FlyBase) was amplified and inserted into a pUASTattB vector at NotI and XhoI restriction sites

(15). To add the *how*(L) 3'UTR (FBtr0084177) was amplified in two parts from *bam* testis

cDNA (to exclude the intron) and inserted upstream of the SV40 polyadenylation site, using

XhoI and XbaI restriction sites. Differences in the *how* coding sequence from published fly

genome (Dm6) were corrected to match *bam^1*/*bam$^{\Delta 86}$* mutant flies, using a Q5 Site-Directed

Mutagenesis Kit (New England Biosciences E0554S). Plasmids were injected into PhiC31

integrase transgenic flies and the constructs were integrated at attP40 on chromosome 2L, then

selected for transformants by BestGene Inc. (Chino Hills, CA) (28).

Immunofluorescence

For whole mount preparations, whole testes were dissected in PBS, fixed in 4%

paraformaldehyde (PFA) for 20 minutes and washed twice in PBS. The tissue was permeabilized

in a solution of 1X PBS, 0.6% sodium deoxycholate, and 0.6% Triton for 1 hour at room

temperature, then washed twice before blocking overnight in PBST(Triton)-3% BSA (bovine

serum albumin) at 4°C. Primary antibodies were added at the following concentrations and testes

were incubated rotating for two days: rabbit anti-HOW (1:50), mouse anti-Bam from DSHB

(1:10), rabbit anti-Kmg (1:200) (Kim et al., 2017, 13), mouse anti-HA (1:200), and goat anti-

Vasa (1:100, Santa Cruz Biotechnology). After two PBS washes at room temperature, donkey

secondary antibodies (Alexa Fluor, ThermoFischer) were added at 1:500 and the samples were

incubated for 2 hours at room temperature. Testes were then mounted on a slide in DAPI mount

(Vectashield). To decrease background/non-specific binding, the rabbit anti-HOW antibody

(gifted by T. Volk) was preabsorbed with ~10 new wild-type testes overnight 3 times (fixed as

for whole mount preparations.

For squashed preparations (samples stained with anti-HOW and anti-Bam), testes were

placed on a SuperFrost Plus slide in a square drawn with a hydrophobic marker. Then the tissues

were flattened live under a coverslip before being flash frozen in liquid nitrogen. After removing

the coverslip, slides were incubated in cold 95% ethanol for 10 minutes before being fixed in 4%

PFA for 7 minutes. Testes were washed in PBST (Triton-X) before being transferred to a wet

chamber for blocking overnight in BSA at 4˚C, after which antibody staining continued as for

whole mounts.

EdU Labeling

Cells in S phase were labeled with the Click-iT EdU Cell Proliferation Kit for Imaging -

Alexa Fluor 555 dye (Invitrogen - C10338). Testes were dissected in Schneider's (S2) medium

and transferred into a drop of the same medium on a slide. The medium was then removed and

replaced with 100 μM 5-ethynyl-2′-deoxyuridine (EdU) in S2 cell medium and incubated for 5

minutes at room temperature. Testes were then washed twice in S2 medium by removing the

liquid with a pipette and replacing it with S2 medium. After the washes, testes were transferred

with tweezers from the drop of S2 medium into a 1.7 mL Eppendorf tube with ~200 uL 1xPBS,

then immediately the PBS was drawn off and replaced with 200 μL 4% paraformaldehyde in

PBS for 20 minutes (rotating at room temperature) to fix, followed by two washes in PBST. The

testes sheaths were then permeabilized in PBS with 0.6% Triton and 0.6% sodium deoxycholate

for one hour. The permeabilization mix was removed and the testes were rinsed once with PBST

before adding the EdU detection reaction mix per the manufacturer's instructions. Testes were

then incubated in the dark with the reaction mix for 30 minutes, the reaction mix was removed,

and the testes washed twice in 500 μL of PBST at room temperature. Blocking and antibody

protocols continued in the same way as for squashes and whole mounts.

Hybridization chain reaction *in situs*

Probes to label the *how* coding sequence and the *how(L) 3'UTR* were designed following

the method of Bedbrook et al., 2023 (29), which generated 44 probes (22 pairs) then ordered

from Integrated DNA Technologies at 50 pmol/oligo. All other reagents were from Molecular

Instruments (Los Angeles, CA), including H1 and H2 hairpins conjugated with 488. Testes were

dissected in 1xPBS, fixed in 4% paraformaldehyde for 20 minutes, then permeabilized in 0.6%

sodium deoxycholate for 30 minutes. Samples were then washed twice in PBS at RT.

Hybridization chain reaction *in situ* hybridization followed the "sample in solution"

HCR™ RNA-FISH protocol from Molecular Instruments with the following specifications:

Probe solution was made with 1 μM of probes. Hybridization buffer and probe wash buffer were

reduced to 200 μL from 500 and pre-amplification was done in 250 μL of amplification buffer.

After completing the HCR protocol, DAPI mounting media was added, testes were mounted on a

slide and imaged by confocal microscopy.

Imaging and quantification

Immunofluorescence stained or HCR FISH testes were imaged on a Leica SP8 Confocal

microscope. Image brightness was adjusted in FIJI. To quantify overproliferation, testes were

scored for the overall number of EdU positive cysts per testis and the number of EdU positive

cysts with >16 cells each. Testes were also stained for the spermatocyte marker Kmg and scored

for the presence of individual Kmg positive spermatocytes or spermatocyte cysts. For the

quantification in Figure 4, EdU and Kmg scoring was performed with the scorer blind to the

genotypes.

Analysis of transcripts expressed in testes

Microarray experiments were performed as described in Kim *et al.,* 2017 (13). To

control for possible persistent changes in gene expression resulting from the heat shock

treatment, control *bam¹/bam^{Δ86}* flies lacking the *hsBam* transgene were subjected to 30min of

heat shock then cultured at 25ºC for 8, 16, 24, or 32hr in parallel with the experimental *hsBam;*

bam¹/bam^{Δ86} flies and the genes expressed in testes from the two genotypes were assessed by

microarray in parallel. Briefly, RNA was extracted from about 30 pairs of dissected testes with

seminal vesicles but without accessory glands using Trizol (Invitrogen). Reverse transcription

was performed with oligo(dT)24 primer with a T7 promoter using ~200 ng of total RNA per

Affymetrix protocol. The second strand was synthesized from the cDNA, and cRNA was

produced by *in vitro* transcription. Fragmented cRNA was hybridized to the *Drosophila* genome

2.0 arrays (Affymetrix, Cat# 900532). All microarray experiments were performed by the

Stanford Protein and Nucleic Acid (PAN) facility. *how* transcript isoforms were distinguished by

probes binding at the 3' end of transcripts, identified by Affymetrix Probeset ID (1637943_at).

For analysis, all the raw CEL files were background adjusted and quantile normalized together

by using R/BioConductor (v3.0.2) package GCRMA (30). Gene annotation was based on the

Affymetrix file: "Drosophila_2.na32.annot.csv".

For analysis of transcript expression by RNA-Sequencing, 10~20 μg of total RNA was

extracted from ~100 pairs of testes plus seminal vesicle from *hsBam; bam¹/bam^{Δ86}* flies for each

time point using Trizol (Invitrogen) followed by RNeasy cleanup (Qiagen) according to kit

instructions. PolyA-tailed RNA was purified using the Oligotex mRNA kit (Qiagen,

Cat#70022). Purified PolyA RNA was fragmented in the presence of random hexamer primers

in first strand synthesis buffer (Invitrogen, Cat# 18080093) at 85˚C for 8 minutes. Fragmented

RNAs were reverse transcribed using Superscript III reverse transcriptase (Invitrogen, Cat#

18080093) in the presence of RNaseOUT (Invitrogen Cat# 10777019) at 50˚C for one hour.

From this step, we followed the directions in the NEBNext mRNA Library Prep Master Mix Set

for Illumina (E6110s) to make libraries. DNA was purified using a QIAquick PCR purification

kit (Qiagen Cat# 28104) after second strand synthesis, end repair, dA tailing, and adapter

ligation. After adapter ligation, 300~500 bp fragments were size-selected by gel extraction (1.5%

low-melt NUSIEVE gel in TBE). Pooled libraries were sequenced with Illumina HiSeq: 100bp,

each paired-end with single indices.

For analysis of time course RNA-Seq data, raw fastq reads were trimmed using trim

galore (version 0.4.3) to remove low-quality (Phred score 20) and adapter-containing (stringency

1) reads (31). Trimmed reads were mapped to the *Drosophila melanogaster* genome

(BDGP6.46) with default parameters using STAR (version 2.5.3) (32). Mapped reads with

quality scores smaller than 10 (-q 10) or not properly paired reads (-f 2) were discarded using

SAMtools (version 1.4.1) (33). Counts per gene were obtained using the featureCounts function

of the subread package (version 1.5.0) (34). Expression levels at different time points were

TMM-normalized (Trimmed Mean of M-values), assuming the majority of housekeeping genes

have the same expression levels in different time points using the R (version 4.1.0) package

edgeR (version 3.34.0) (35). Axes for scatter plots were log2 transformed normalized CPM

(Counts Per Million) + 1. Scatter plots were generated by Matlab (R2021a). Analyses scripts are

available in: https://github.com/jongminkmg/HeldOutWings2024.

To ensure knockdown of *caf40,* we analyzed gene expression levels using RNA-seq

for testes plus seminal vesicle from *bam* mutant (48hrs after heat shock) and *nosGal4*

driven *caf40* knockdown flies (Figure S5). Testes were dissected from 0–2 day old males in

1xPBS, as batches of 50 flies, in a cyclops dissecting dish for <30min at room temperature. Each

batch was transferred to a 1.7 ml Eppendorf tube containing 1ml of 1xPBS, the PBS was then

immediately removed, and the testes were snap-frozen in liquid nitrogen and stored at −80°C.

RNA was extracted from 150 pairs of testes using the RNeasy Plus Mini Kit from QIAGEN

(74104). Frozen tissue was dissociated using a 1 ml syringe with a 27-gauge needle aspirating up

and down ∼10 times in 300µl of lysis buffer (from RNeasy kit) supplemented with 1:100 β-

mercaptoethanol. Library preparation was carried out using the NEBNext® Ultra™ II

Directional RNA Library Prep Kit for Illumina (#E7760S), with the NEBNext® Poly(A) mRNA

Magnetic Isolation Module (#E7490S). Sequencing was performed by Novogene on a

NOVAseq, PE 150 Illumina platform. Adapters and low-quality bases were trimmed with

trimGalore and then aligned to the *Drosophila* dm6 genome using STAR. Differential expression

analysis was performed using DEseq2.

Acknowledgements

We thank members of the Fuller Lab for their insights on the project and helpful

suggestions and T. Templin for help with statistical analysis. DH thanks Anne Villeneuve,

Lauren Goins, and Roel Nusse for their guidance on experiments, interpreting data, and data

visualization. Fly food and solutions were made by The Developmental Biology Department Fly

Food Facility and Media Facility. Talila Volk generously provided UAS-HOW fly stocks and

How antibodies. The Stanford Protein and Nucleic Acid (PAN) facility assisted with the

microarray experiments. Kitty Lee and Jon Mulholland of Stanford's Cell Sciences Imaging

Facility (CSIF) provided invaluable training and assistance on the Leica SP8 Confocal

Microscope.

This work was funded by NIGMS MIRA #1R35GM136433, the Katherine Dexter

McCormick and Stanley McCormick Memorial Professorship, and the Reed-Hodgson

Professorship in Human Biology to MTF, the Stanford University Genetics and Developmental

Biology Training Grant 5T32GM007790 (DH), and The Italian American Cancer Research

Fellowship (LG). RNA-sequencing from the heat shock time course was partially funded by

1U54HD068158 (PI: Renee Reijo Pera). The project described was also supported, in part, by

Award Number 1S10OD010580-01A1 from the National Center for Research Resources

(NCRR). Its contents are solely the responsibility of the authors and do not necessarily represent

the official views of the NCRR or the National Institutes of Health.

References

1.	E. Rangel-Huerta, E. Maldonado, Transit-Amplifying Cells in the Fast Lane from Stem Cells towards Differentiation. *Stem Cells Int.* **2017**, 7602951 (2017).

2.	M. F. Clarke, M. Fuller, Stem Cells and Cancer: Two Faces of Eve. *Cell* **124**, 1111–1115 (2006).

3.	M. T. Fuller, "Spermatogenesis" in *The Development of Drosophila Melanogaster*, Cold Spring Harbor Laboratory Series., (Cold Spring Harbor Laboratory Press, 1993), pp. 71-147.

4.	M. L. Insco, A. Leon, C. H. Tam, D. M. McKearin, M. T. Fuller, Accumulation of a differentiation regulator specifies transit amplifying division number in an adult stem cell lineage. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 22311–22316 (2009).

5.	D. M. McKearin, A. C. Spradling, bag-of-marbles: a Drosophila gene required to initiate both male and female gametogenesis. *Genes Dev.* **4**, 2242–2251 (1990).

6.	P. Gönczy, E. Matunis, S. DiNardo, bag-of-marbles and benign gonial cell neoplasm act in the germline to restrict proliferation during Drosophila spermatogenesis. *Dev. Camb. Engl.* **124**, 4361–4371 (1997).

7.	Y. Li, N. T. Minor, J. K. Park, D. M. McKearin, J. Z. Maines, Bam and Bgcn antagonize Nanos-dependent germ-line stem cell maintenance. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 9304–9309 (2009).

8.	D. Chen, *et al.*, Three RNA binding proteins form a complex to promote differentiation of germline stem cell lineage in Drosophila. *PLoS Genet.* **10**, e1004797 (2014).

9.   B. Ohlstein, C. A. Lavoie, O. Vef, E. Gateff, D. M. McKearin, The Drosophila

cystoblast differentiation factor, benign gonial cell neoplasm, is related to DExH-box

proteins and interacts genetically with bag-of-marbles. *Genetics*, **155**, 1809-1819 (2000).

10.   A. Sgromo, *et al.*, Drosophila Bag-of-marbles directly interacts with the CAF40

subunit of the CCR4-NOT complex to elicit repression of mRNA targets. *RNA N. Y. N* **24**,

381–395 (2018).

11.   M. L. Insco, *et al.*, A self-limiting switch based on translational control regulates

the transition from proliferation to differentiation in an adult stem cell lineage. *Cell Stem*

Cell **11**, 689–700 (2012).

12.   A. C. Monk, *et al.*, HOW is required for stem cell maintenance in the Drosophila

testis and for the onset of transit-amplifying divisions. *Cell Stem Cell* **6**, 348–360 (2010).

13.   J. Kim, *et al.*, Blocking promiscuous activation at cryptic promoters directs cell

type-specific gene expression. *Science* **356**, 717–721 (2017).

14.   H. Li, *et al.*, Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit

fly. *Science* **375**, eabk2432 (2022).

15.   V. K. Jenkins, A. Larkin, J. Thurmond, FlyBase Consortium, Using FlyBase: A

Database of Drosophila Genes and Genetics. *Methods Mol. Biol.* **2540**, 1-34 (2022).

16.   L. Shan, *et al.*, Regulators of alternative polyadenylation operate at the transition

from mitosis to meiosis. *J. Genet. Genomics Yi Chuan Xue Bao* **44**, 95–106 (2017).

17.   Z. Jin, *et al.*, Differentiation-defective stem cells outcompete normal stem cells

for niche occupancy in the Drosophila ovary. *Cell Stem Cell* **2**, 39-49 (2008).

18.   L. Gilboa, R. Lehmann, Repression of primordial germ cell differentiation

parallels germ line stem cell maintenance. *Curr. Biol. CB* **14**, 981–986 (2004).

19.   A. Forbes, R. Lehmann, Nanos and Pumilio have critical roles in the development

and function of Drosophila germline stem cells. *Dev. Camb. Engl.* **125**, 679–690 (1998).

20.   Y. Li, *et al.*, Mei-P26 Cooperates with Bam, Bgcn and Sxl to Promote Early

Germline Development in the Drosophila Ovary. *PLoS One* **8**, e58301 (2013).

21.   S. Malik, W. Jang, J. Y. Kim, C. Kim, Mechanisms ensuring robust repression of

the Drosophila female germline stem cell maintenance factor Nanos via posttranscriptional

regulation. *FASEB J.* **34**, 11421–11430 (2020).

22.   A. C. Monk, N. A. Siddall, B. Fraser, E. A. McLaughlin, G. R. Him, Differential

roles of HOW in male and female Drosophila germline differentiation. *PLoS One* **6**, e28508

(2011).

23.   F. Rodrigues, L. Thuma, C. Klämbt, The regulation of glial-specific splicing of

Neurexin IV requires HOW and Cdk12 activity. *Dev. Camb. Engl.* **139**, 1765–1776 (2012).

24.   M. Van Doren, A. L. Williamson, R. Lehmann, Regulation of zygotic gene

expression in *Drosophila* primordial germ cells. *Curr. Biol.* **8**, 243–246 (1998).

25.   D. Chen, D. M. McKearin, A discrete transcriptional silencer in the bam gene

determines asymmetric division of the Drosophila germline stem cell. *Dev. Camb. Engl.*

130, 1159–1170 (2003).

26.   D. McKearin, B. Ohlstein, A role for the Drosophila Bag-of-marbles protein in

the differentiation of cystoblasts from germline stem cells. *Development* **121**, 2937–2947

(1995).

27.   D. Lu, H. S. Sin, C. Lu, M. T. Fuller, Developmental regulation of cell type-

specific transcription by novel promoter-proximal sequence elements. *Genes Dev.* **34**, 663–

677 (2020).

28.   J. Bischof, R. K. Maeda, M. Hediger, F. Karch, K. Basler, An optimized

transgenesis system for Drosophila using germ-line-specific φC31 integrases. *PNAS* **104**,

3312-3317 (2007).

29.   C. N. Bedbrook, R. D. Nath, R. Nagvekar, K. Deisseroth, A. Brunet, Rapid and

precise genome engineering in a naturally short-lived vertebrate. *eLife* **12**:e80639 (2023).

30.   B. M. Bolstad, R. A. Irizarry, M. Astrand, T. P. Speed, A comparison of

normalization methods for high density oligonucleotide array data based on variance and

bias. *Bioinforma. Oxf. Engl.* **19**, 185–193 (2003).

31.   M. Martin, Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).

32.   A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinforma. Oxf.*

Engl. **29**, 15–21 (2013).

33.   H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinforma.*

Oxf. Engl. **25**, 2078–2079 (2009).

34.   Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinforma. Oxf. Engl.* **30**, 923–

930 (2014).

35.   M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. *Bioinforma. Oxf. Engl.*

26, 139–140 (2010).

Figure 1. Nuclear How persists in *bam* mutant spermatogonia.

(A) Diagram of *Drosophila* early male germ line development.

(B-B''') Immunofluorescence images of apical tip of a wild type testis stained with anti-How,

anti-Vasa, and anti-Bam. (B) merge of (green) How and (magenta) Bam. (B'-B''') Black and

white single channels showing (B') How, (B'') Bam, and (B''') Vasa to mark germ cells.

(C-D) Immunofluorescence images of (C) wild type and (D) $bam^{1/\triangle86}$ testis apical tips stained

with anti-How and anti-Vasa. Dashed cyan line in (C) marks border between spermatogonia (left

of line) and the spermatocyte region (right). Arrowheads in B and C: somatic cyst cell nuclei.

(E) Scatter plot of RNA-sequencing data from *hs-Bam*; $bam^{1/\triangle86}$ comparing $\log_2$ transcript

expression levels per gene for testes from flies not treated with heat shock (no HS) to testes from

flies 8 hours post heat shock (PHS). (Blue dots) *how* is one of the 6 genes for which transcript

levels decreased by over 2-fold by 8 hours post induction of Bam expression by heat shock in

both the RNA-seq and separate microarray analysis (details in Figure S1). (Red dots)

Transcripts from two genes increased in expression level in both the RNA-seq and microarray

analysis. Diagonal lines mark 2-fold change.

(F) UMAP visualization of single nuclear RNA sequencing data from the Fly Cell Atlas (14),

after the nuclei in the two earliest male germ line clusters (Leiden resolution 6.0) were

reclustered. (Green) nuclei scoring positive for *how*. (Magenta) nuclei scoring positive for *bam*.

(White) nuclei positive for both *how* and *bam* transcripts.

(G-G'') Fluorescence images of testis apical tip showing (magenta) Bam-GFP protein and

(green) *how* mRNA visualized by hybridization chain reaction (HCR) using probes to protein

coding sequences of *how*. Dashed outlines: spermatogonial cysts positive for Bam protein either

(yellow) with how RNA, (white) with fewer loci of how RNA signal, or (cyan) no how RNA

signal detected. Scale bars: 25 μm in B-D and 12.5 μm in G.

Figure 2. Knocking down *how* allowed spermatogonia lacking *bam* to differentiate into

spermatocytes.

(A-C) Phase contrast images of apical regions of testis in live squash preparations. (A) *bamGal4*

expression driver only. (B) *bam*$^{1/\Delta 86}$. (C) *bam*$^{1/\Delta 86}$; *bamGal*4 > *how* RNAi (VDRC100775). All

flies were raised under the same temperature shift regimen used for RNAi. Arrowheads:

spermatocytes. Arrows: spermatid bundles. SG: spermatogonia. Scale Bars: 100 μm.

(D-F) Immunofluorescence images of testis apical tips stained for (red) EdU to mark nuclei in S

phase and (blue) Kmg to mark spermatocyte nuclei. (D) *bamGal4* driver only control. (E)

bam$^{1/\Delta 86}$. (F) *bam*$^{1/\Delta 86}$; *bamGal4* > *how* RNA. Scale Bars: 25 μm.

(G, H) Phase contrast images of intact spermatocyte cysts marked with the number of

spermatocytes in the cyst. (G) *bamGal4* driver only. (H) *bam*$^{1/\Delta 86}$; *bamGal4* > *how* RNAi. Scale

Bars: 50 μm.

(I) Number of spermatocytes per cyst from: (Left side) *bamGal4* sibling controls (N=24 cysts) or

bam$^{1/\Delta 86}$; *bamGal4* > *how RNAi* VDRC100775 (N=43 cysts). (***) p-value = 1.716e-10, based

on a Wilcoxon rank-sum test with continuity correction. (Right side) *bamGal4* control (N=52

cysts) or *bam*$^{1/\Delta 86}$; *bamGal4* > *how RNAi* TRiP HMC03820 from Bloomington Stock #55665

(N=70 cysts). (****) p-value = 1.033e-14 based on a Wilcoxon rank-sum test with continuity

correction.

Figure 3. Overexpression of How(L) but not How(S) blocked the transition from

spermatogonia to spermatocytes.

(A-C) Phase contrast images of testis apical tips from (A) *bamGal4* driver only. (B) *bamGal4 >*

UAS-How(L)HA-SV40. (C) *bamGal4 > UAS-How(S)HA-SV40*. SG: spermatogonia.

Arrowheads: spermatocytes. Arrows: elongating spermatids. Red line in B marks region with

overproliferating spermatogonia. Scale bars: 50 μm.

(D-F) Immunofluorescence images of apical tips of testes stained with (top row) DAPI to mark

nuclei and (bottom row) EdU to mark nuclei in S phase and anti-Kmg to mark spermatocyte

nuclei. (D) *bamGal4* driver only. (E) *bamGal4 > UAS-How(L)HA-SV40*. (F) *bamGal4 > UAS-*

How(S)HA-SV40. Scale Bars: 25 μm.

(G-I) Immunofluorescence images of testis apical tips stained with anti-HA to detect the forcibly

expressed How(L) and How(S). (G) *bamGal4* driver alone control. (H) *bamGal4*, *UAS-*

How(L)HA-SV40 showing nuclear localization of How(L). Note persistence of How(L)HA

expression in the overproliferating spermatogonia. (I) *bamGal4 > UAS-How(S)HA-SV40*

showing cytoplasmic localization of How(S). Note shut off of How(S)HA expression in

differentiating spermatocytes.

(J-L) Immunofluorescence images of testis apical tips stained with anti-Bam. (J) *bamGal4*

control. (K) *bamGal4 > UAS-How(L)HA-SV40*. (L) *bamGal4 > UAS-How(S)HA-SV40*.

Scale bars: 25 μm.

Figure 4. Adding the How(L) 3'UTR reduced the effect of forced expression.

(A-F) Top: cartoons depicting design of the overexpression constructs. (Left) *UAS-How(L)HA*

with the SV40 terminator, (right) *UAS-How(L)HA* with the 2424nt 3'UTR from How(L) (Figure

S2A) added before the SV40 terminator. Immunofluorescence images of testes stained for

(white) Kmg to mark spermatocyte nuclei and (red) EdU to label cysts in S phase in the

phenotypic classes indicated in G. (A-C) *bamGal4 > UAS-How(L)HA-SV40*. (D-F) *bamGal4 >*

UAS-How(L)HA-3'UTR-SV40. Flies were grown at 18°C for their entire life. Arrowheads: EdU

positive cysts. Blue arrow: single Kmg positive spermatocyte. Yellow arrows: spermatocytes in

cysts. Scale Bars in A-F: 25 μm. (G) Graph of the percentage of testes from each genotype with

either (gray) no spermatocytes detected, (blue) single spermatocytes or (orange) cysts of multiple

spermatocytes, scored based on the spermatocyte marker Kmg. *SV40* n = 30 testes.

How(L)3'UTR n = 29 testes. Percentages were compared using a two-sample t-test of

proportions. Percentage of testes containing spermatocyte cysts (p = 2.5 x 10-5) and percentage

of testes containing any spermatocytes (p = 3.0 x 10-4) differed significantly between the two

genotypes. (H) Distribution of the number of EdU positive cysts in *bamGal4 > UAS-How(L)-*

SV40 testes and *bamGal4 > UAS-How(L)-3'UTR-SV40* testes. Statistical tests performed via 1-

way between-groups ANOVA. (I) Number of overproliferating (>16 nuclei) EdU positive cysts

from testes overexpressing *How(L)-SV40* (n = 30 testes) compared to *How(L)-3'UTR-SV40* (n =

29 testes). P-value calculated as for (H). The median in *How(L)-3'UTR-SV40* was zero

overproliferating cysts/testis. Medians marked by red arrowheads in all conditions.

Figure 5. The CCR4-NOT component Caf40 is required for *how* repression.

(A, C) Phase contrast images of testis apical tips from males carrying (A) *nosGal4* driver only

control and (C) *nosGal4 > Caf40 RNAi* (TRiP HMS05850 Bloomington stock #67987). White

arrowhead: spermatocytes. White arrow: spermatid bundles. SG = spermatogonia. (B, D) Spilled

out cysts from (B) *nosGal4* control testes including many spermatocyte cysts as well as more

uncommon cysts of smaller spermatogonia. Spilled out cysts from (D) *nosGal4 > Caf40* RNAi

males contained many small cells and no spermatocytes. (E, F) Immunofluorescence images of

testes apical tips from (E) *nosGal4* driver only control and (F) *nosGal4 > Caf40* RNAi labeled

with (red) EdU to mark nuclei in S phase and (cyan) anti-Kmg to mark spermatocyte nuclei.

Yellow arrowheads: cysts of EdU positive spermatogonia. (G, H) Immunofluorescence images

of testis apical tips stained with anti-Bam from (G) *nosGal4* and (H) *nosGal4 > Caf40* RNAi

flies. Yellow line: boundary between early spermatogonia and Bam positive spermatogonia. (I-

K) Apical tips of testes stained for *how* RNA by HCR with probes complementary to the How

protein coding sequence. (I) *nosGal4* control. Green bracket: region with *how* transcripts in

spermatogonia. (J) *nosGal4 > Caf40* RNAi. (K) *bam*$^{1/\triangle 86}$. Scale bars: 50 μm A-D; 25 μm E-K.

All control flies were raised under the same temperature regimen used for the RNAi (3 days at

25˚C, then cultured at 29˚C).

Figure 6. Model for mechanism of Bam action on *how* RNA.

(A) Cartoon of *How* (green) and *Bam* (purple) protein expression in the germ line stem cell and

spermatogonia. Levels of *how* decrease as Bam expression appears at the four cell cyst stage.

Bam levels increase in mid-to-late spermatogonia until sufficient to repress expression of *How*,

allowing the subsequent switch from proliferation to differentiation.

(B) Model for regulation by Bam, Bgcn and Tut: Bam protein binds the CCR4-NOT subunit

Caf40, acting as an adaptor between the CCR4-NOT complex and its target transcripts.

 SOM Figure Legends

Figure S1. *how* is among the earliest transcripts to decrease after Bam is turned on.

(A, B) Scatterplots of transcript levels from microarray analysis comparing no heat shock and 8

hours PHS in (A) $bam^{1/\Delta86}$; *hs-Bam* or (B) $bam^{1/\Delta86}$ males lacking the *hs-Bam* construct, but

subjected to the same heat shock regimen in parallel, to control for the effects of heat shock on

gene expression. Genes colored blue (downregulated) or red (upregulated) were 1) not changed

>2 fold in the microarray analysis of testes from control $bam^{1/\Delta86}$ males lacking the *hs-Bam*

construct but heat shocked then incubated for the indicated time and 2) also detected by

independent RNA-seq as up or down >2 fold in $bam^{1/\Delta86}$; *hs-Bam* males by RNA-sequencing at

the time points indicated compared to testes from flies of the same genotype not subjected to heat

shock. (C) Level of *how(L)* transcripts detected by microarray throughout the time course,

showing decrease by 8h PHS in flies carrying the *hs-Bam* construct, but not in testes from

control $bam^{1/\Delta86}$ flies lacking the hs-Bam construct but subjected to the 30-minute pulse of

incubation at high temperature, then shifted back to 25˚C as for the experimental genotype.

(D, E) Microarray data from later time points after heat shock, showing the transcripts that

increase in expression (red) and decrease in expression (blue) by the cutoff criteria (detected as

up or down regulated >2 fold) in both this microarray comparison and in independent analysis by

RNA-seq. (D) $bam^{1/\Delta86}$; *hs-Bam* testes (red oval in *hs-Bam* 32h: genes expressed specifically in

spermatocytes. (E) testes from control $bam^{1/\Delta86}$ males that did not have the *hs-Bam* transgene, but

were subjected to heat shock then incubated at 25^{o}C for the indicated times. Black oval in (E)

16h PHS marks genes expressed in accessory glands that contaminated this sample.

Figure S2. How transcripts were detected in spermatogonia but not in spermatocytes

(A) Diagram of the *how* locus showing mRNA isoforms from FlyBase, designated A-F as in

Flybase. The How(L) cDNA construct utilized in Figures 3 and 4 is RA and the How(S) cDNA

construct utilized in Figure 3 is RB. Grey: UTRs. Green: protein coding sequence. Lines denote

introns. Red arrowheads: nuclear localization signal. Orange arrowhead: additional coding

sequence in isoform RF, distinguishing it from RA. Black bars: Sites of RNAi constructs

utilized. (B) Apical tip of wild type testis showing distribution of *how* transcripts detected by

Hybrid Chain Reaction (HCR FISH) using two different probe sets. (B) Merge. Magenta:

Coding sequence (CDS) probe set; Green: 3'UTR probe set. Zoom in marked by red box.

Yellow rectangles mark white loci of overlapping probes sets. (B') How(L) 3'UTR probe set.

(B") Coding sequence (CDS) probe set. Scale Bar: 25 µm. (C-E) High magnification

immunofluorescence images of apical tips of *Bam-GFP* testes with How protein coding sequence

RNA labeled by HCR (additional examples for Figure 1G). Left: merge with (blue) DAPI,

(green) *how* RNA, and (magenta) Bam-GFP. Middle: *how* RNA only. Right: Bam protein only.

Yellow dashed outlines: early Bam positive cysts with *how* RNA present. White dashed

outlines: later stage Bam positive cysts with fewer h*ow* RNA foci. Cyan dashed outlines: Bam

positive cysts with no or very low *how* RNA signal detected. Scale bar: 12.5 µm.

Figure S3. Knockdown of *how* in mid- to late spermatogonia in *bam* mutant males resulted

in production of spermatocytes for two different RNAi lines.

(A-C) Phase contrast images of testis apical tips from (A) *bamGal4* driver only; (B) *bam*$^{1/\triangle86}$;

and (C) *bam*$^{1/\triangle86}$; *bamGal4 > how* RNAi (TRiP HMC03820 from Bloomington Stock #55665).

Arrowheads: spermatocytes. Arrows: spermatid bundles. SG: spermatogonia. Scale bars: 100

µm, insets Scale Bars: 25 µm. (D-F) Immunofluorescence images of apical testis tips stained for

EdU to mark S-phase nuclei (red) and Kmg to mark spermatocyte nuclei (blue). (D) Control:

*bam*Gal4 driver only. (E) *bam$^{1/\triangle 86}$*. (F) *bam$^{1/\triangle 86}$*; *bam*Gal4 > *how* RNAi (TRiP HMC03820 from

Bloomington Stock #55665). (G-I) Whole testis from (G) *bamGal4* driver only; (H) *bam$^{1/\triangle 86}$* ;

and (I) *bam$^{1/\triangle 86}$*; *bamGal4* > *how* RNAi (VDRC100775) for size comparison. Arrowhead:

spermatocytes. Arrow: spermatid bundles. Scale bars: 100 μm. (J) Fertility tests of *bamGal4*

driver only, *bam$^{1/\triangle 86}$*; and *bam$^{1/\triangle 86}$*; *bamGal4* > *how* RNAi (VDRC 100775) males. Males from

all genotypes were progeny from crosses where the mated parents were allowed to lay eggs for 3

days at 25˚C then the parents were removed and the progeny shifted to 29˚C. Once the progeny

males were collected, they were subjected to fertility tests at RT, ~ 25˚C.

Figure S4: Knockdown of *how* in mid- to late spermatogonia in *bgcn* mutant males resulted

in production of spermatocytes.

Immunofluorescence images of apical tip of *bgcn$^{1/63-44}$* mutant testis stained with (A) anti-How

and (A') anti-Vasa. Scale bars: 25 μm. (B,C) Phase contrast images of testis apical tips from (B)

bgcn$^{1/63-44}$ versus (C) *bgcn$^{1/63-44}$*; *bam-Gal4* > *how RNAi* (VDRC 100775). Arrowhead:

spermatocytes. Arrow: spermatid bundles. Scale bars: 50 μm.

Figure S5. Knockdown of *caf40* in early germ cells by RNAi hairpin resulted in early germ

cell overproliferation similar to *bam* mutants.

(A) Diagram of *Caf40* locus based on Flybase, showing sites of RNAi constructs tested (black

lines). (B,C) Phase contrast images of testis apical regions showing two additional examples of

nosGal4 VP16 driving knockdown of *Caf40* using a different *RNAi* line (VDRC 101462).

Although the knockdown phenotype was not as strong as for the RNAi construct carried in

Bloomington Stock #67987 (Figure 5), there was still overproliferation of small early germ cells

followed by cell death. SG: spermatogonia. Arrowhead: spermatocytes. Scale bars: 50 μm.

(D,E) Comparison of gene expression levels in testes from *nosGal4;UASCaf40 RNAi*

(Bloomington 67987) *versus bam*$^{1/\Delta 86}$ males 48hrs after heat shock. D) RNA-Seq data showing

expression levels per gene plotted as log10(normalized counts + 1). E) Volcano plot comparing

RNA-Seq data from (D). (Blue dot) expression of *Caf40*: RNA-Seq confirmed that the

knockdown lowered levels of *Caf40* RNA by approximately 14-fold. (Magenta dot) expression

of *bam*: The level of *bam* mRNA detected was higher in *Caf40* knockdown than in *bam* mutant

testes, likely in part because *bam*$^{\Delta 86}$ deletes most of the *bam* coding sequence (26). (Green dot)

expression of *how*: RNA-Seq confirmed that levels of *how* RNA in *nosGal4;UASCaf40 RNAi*

testes were similar to the levels in *bam* mutant testes.

Figure S6. Knockdown of *how* in wild-type mid to late spermatogonia did not affect the

switch to differentiation.

(A,B) Phase contrast images of testis apical regions from flies expressing different *how* RNAi

hairpins driven by *bamGal4*: (A) Vienna Drosophila Resource Center (VDRC)100775. (B) TRiP

HMC03820 from Bloomington Drosophila Stock #55665. Arrowheads: spermatocytes. Arrows:

elongating spermatids. Scale bars: 50 μm.

Figure 1

A hub, GSC, Gb, TA spermatogonia, mitosis, terminal differentiation, premeiotic S phase, spermatocytes, meiotic divisions, spermatid elongation

B WT — How (green), Bam (magenta)
B' How
B'' Bam
B''' Vasa

C WT — How
C' Vasa
D *bam⁻/⁻* — How
D' Vasa

E No HS vs 8 hours PHS — How

F how, bam, merge — umap_2 vs umap_1

G DAPI, How RNA, Bam Protein
G' How RNA
G'' Bam Protein

Figure 2

Figure 3

bamGal4 control	*bamGal4 > How(L)HA*	*bamGal4 > How(S)HA*

A — **B** SG — **C**

D DAPI — **E** DAPI — **F** DAPI

D' EdU Kmg — **E'** EdU Kmg — **F'** EdU Kmg

G HA — **H** HA — **I** HA

J Bam — **K** Bam — **L** Bam

Figure 4

Figure 5

A — *nosGal4* control

B

C — SG only — *nosGal4 > Caf40* RNAi

D

E — EdU Kmg — *nosGal4* control

F — EdU Kmg — *nosGal4 > Caf40* RNAi

G — Bam — *nosGal4* control

H — Bam — *nosGal4 > Caf40* RNAi

I — *how* RNA — *nosGal4* control

J — *how* RNA — *nosGal4 > Caf40* RNAi

K — *how* RNA — *bam⁻/⁻*

Figure 6

A

B