New evidence for the presence and function of phosphoinositides (PPIs) in the chloroplast

Mastoureh Sedaghatmehr[1,2], Frieda Rößler[1,3], Alexander P. Hertle*[1,2,4]

[1]Max-Planck Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam, Germany

[2]Heinrich-Heine-Universität Düsseldorf, 40225 Düsseldorf, Germany

[3]Leibnitz Institute of Plant Biochemistry, Weinberg 3, 06120 Halle, Germany

[4]Cluster of Excellence on Plant Sciences, Heinrich Heine University, Düsseldorf, Germany

***Correspondence:** Alexander P. Hertle, email: Alexander.Hertle@hhu.de

Abstract

Essential proteins involved in thylakoid formation, including chloroplast-localized Sec14-like protein (CPSFL1) and the vesicle-inducing protein in plastids (VIPP1), bind PPIs *in vitro* with high affinity. PPIs are a class of phospholipids characterized by a phosphorylated inositol head group. Although PPIs constitute a relatively small proportion of the total phospholipids, they play essential roles in various regulatory processes. The exact subcellular localization of most PPI species in plants is incomplete due to their rapid turnover and low abundance. Currently there is no documented evidence for the presence and function of phosphoinositides (PPIs) in chloroplasts. In our study, we developed genetically encoded biosensors targeted into plastids, enabling the detection of various PPI isoforms (PI3P, PI4P, PI5P, PI(4,5)P$_2$ and PI(3,5)P2) within the chloroplasts. We effectively demonstrated the specificity of our PPI biosensors in detection of PPIs, as evidenced by the alterations in PPI biosensor distribution patterns upon co-expression of PPI modifying enzymes (*cTP-SAC7*, *cTP-PTEN*, and *cTP-dOCRL*). Additionally, our research confirmed the capability of the generated PPI biosensors to detect PPIs within the chloroplasts of both tobacco and *Arabidopsis* in a concentration-dependent manner. Furthermore, we unveiled the association and potential interaction of PI3P with VIPP1. We could show that the increased PPI flux within the cell during heat stress affects PPI levels in the chloroplasts, resulting in different distribution patterns of the PPI biosensors. Finally, plants expressing PPI modifiers *cTP-SAC7*, *cTP-PTEN*, and *cTP-dOCRL,* in the chloroplast, showed increased sensitivity to drought stress, likely due to impaired PPI signaling.

Keywords: Phosphatidylinositol phosphates, chloroplasts, signaling, PIP biosensors impaired PPI signaling.

Keywords: Phosphatidylinositol phosphates, chloroplasts, signaling, PPI biosensors

Introduction

Phosphatidylinositols (PtdIns or PIs) are low-abundance lipids with significant regulatory functions, governing a wide range of physiological processes in eukaryotes. PtdIns are produced in the endoplasmic reticulum (ER) and subsequently transported to various subcellular sites through a combination of lipid transfer proteins and vesicle trafficking mechanisms (Phillips et al., 2006; Heilmann, 2009). Phosphorylated derivatives of PtdIns are generated by a group of enzymes, including phosphatidylinositol kinases (PIKs) and phosphatidylinositol phosphatases (Mueller-Roeber and Pical, 2002). PIK enzymes phosphorylate three out of the five available hydroxyl groups on the inositol ring, specifically at positions 3′, 4′, and 5′. This process generates seven phosphatidylinositol phosphates (PtdInsPs or PPIs) consisting of three monophosphates (PI3P, PI4P, and PI5P), three bisphosphates ($PI(3,4)P_2$, $PI(4,5)P_2$, and $PI(3,5)P_2$) and one triphosphate ($PI(3,4,5)P_3$). Among them, the presence of $PI(3,4)P_2$ and $PI(3,4,5)P_3$ has not yet been confirmed in plants (Mueller-Roeber and Pical, 2002; Zarreen et al., 2023). To date, no homologs of the heterodimeric inositol lipid 3-kinases have been identified in plant cells to phosphorylate PI4P to $PI(3,4)P_2$ and $PI(4,5)P_2$ to $PI(3,4,5)P_3$. PI5P is found in plants, yet an enzyme responsible for its production remains unidentified (Mueller-Roeber and Pical, 2002; Heilmann, 2009; Zarreen et al., 2023).

PPIs are being recognized as new secondary messengers within plant cells. Besides their crucial role under favorable conditions, they are quickly generated when plants are exposed to a wide range of environmental stresses, indicating their importance under stress conditions (Munnik and Meijer, 2001; Meijer and Munnik, 2003; Wang et al., 2008; Xue et al., 2009; Munnik and Vermeer, 2010; Pribat et al., 2011; Zhai et al., 2013; Zarreen et al., 2023). PI3P, PI5P, $PI(3,5)P_2$, and $PI(4,5)P_2$ exhibited increased levels in response to salt and osmotic stress, likely due to the induction of relevant PI kinases or the action of lipid phosphatases, resulting from the degradation products of other PPIs. (Meijer et al., 1999; Pical et al., 1999; DeWald et al., 2001; Meijer et al., 2001; Meijer and Munnik, 2003). Individual studies have reported alteration in PPI kinase and phosphatase activity under stress conditions. Salt and osmotic stress have been shown to boost both PI 3-kinase and PI3P 5-kinase activities in plants. (Meijer et al., 1999; Meijer et al., 2001; Munnik and Meijer, 2001; Leshem et al., 2007). PTEN (phosphatase and tensin homolog deleted on chromosome ten) phosphatases in vascular plants play a crucial role in regulating 3-phosphorylated PI signaling events. The transcript levels of *AtPTEN2* exhibited changes in response to various stresses, including conditions of salt and osmotic stress (Pribat et al., 2011). Similarly, environmental stresses such as cold, salt, and drought stimulate the expression and activity of phosphoinositide-specific phospholipase C (PLC) (Hirayama et al., 1995; Munnik and Meijer, 2001). PLC induces the hydrolysis of $PI(4,5)P_2$ to generate the two signaling molecules, inositol-$_{1,4,5}$-trisphosphate ($InsP_3$) and diacylglycerol (DAG) (Hirayama et al., 1995; Testerink and Munnik, 2011; Zarreen et al., 2023). PLC has the ability to enhance salt tolerance in rice as well as drought tolerance in maize and tobacco (Wang et al., 2008; Zhai et al., 2013; Li et al., 2017). The expression of the well-characterized PI4K kinase, *AtPI4Kγ3*, is induced upon exposure to salt stress. *Arabidopsis* plants overexpressing PI4Kγ3 exhibit increased stress tolerance and inducible expression of stress-responsive genes (Akhter et al., 2016). All these investigations suggest that alterations in PPI concentration are tightly regulated and play a critical role in facilitating precise PPI signaling.

The unique characteristics of PPIs, including their low abundance, rapid turnover, and diversity, make them dynamic signaling molecules. PPIs have the ability to bind to specific proteins, which can result in changes in the proteins' localization, conformation, and activity. Additionally, PPIs can influence membrane curvature. These capabilities are crucial aspects of PPI function in facilitating downstream signaling in all eukaryotes (Meijer and Munnik, 2003; Xue et al., 2009; Munnik and Vermeer, 2010).

While the functions of PtdIns in membrane trafficking, endomembrane organization, and cytoskeletal
rearrangement are well-established, their pivotal roles in signaling and their subcellular localization
are only emerging in plants. Due to their rapid turnover and limited abundance, the precise subcellular
localization of most PPI species in plants remains incomplete. The existence and function of PPI in
the nucleus of animals, yeast, and plant cells has been documented (Horton et al., 2007; Dieck et al.,
2012; Gerth et al., 2017). It is widely recognized that nuclear PtdIns play a significant role in
regulating various processes, including DNA replication, chromatin remodeling, transcription, stress
responses, and hormone signaling in plant, animal, and yeast cells (Dieck et al., 2012; Zarreen et al.,
2023). The unique role of PPIs in the nucleus, distinct from their function in the plasma membrane,
highlights the existence of distinct PPI signaling pathways within various cellular organelles.

Only few biochemical studies employing radiolabeling of isolated spinach chloroplasts indicate the
presence of PPIs in plastids. (Siegenthaler et al., 1997; Müller et al., 2000; Bovet et al., 2001). A PPI
kinase activity could be assigned to the envelope membranes (Siegenthaler et al., 1997). However,
sub-organelle fractionation of chloroplast membranes failed to demonstrate PPI distribution and
abundances (Müller et al., 2000). A more recent study revealed a role of PI4P in chloroplast division
by inhibiting PI4Ks through chemical inhibitor treatment in *Arabidopsis* seedlings a It has also been
reported that PI4P specifically binds to PLASTID DIVISION1 (PDV1) and PDV2, exerting a
negative regulatory effect on chloroplast division. Of note, PDV1 and PDV2 proteins are located at
the outer envelope membrane of the chloroplast, and the PI4P lipid-signaling pathway, involved in
chloroplast division via PDVs takes place at the outer envelope membrane (Okazaki et al., 2015).
Thus, presence and function of PPIs inside the chloroplasts remain unexplored thus far (Okazaki et
al., 2015). Recently we identified a chloroplast localized Sec14-like protein (CPSFL1) homologues to
yeast Sec14 (Hertle et al., 2020). Yeast Sec14 proteins regulate cytosolic vesicle transport between the
trans-Golgi network and the plasma membrane by modulating PIP levels (Bankaitis et al., 1990).
CPSFL1 was shown to act *in vitro* as a phosphatidylinositol transfer protein (PITP) which directly
transports PPIs between membranes. Expression of CPSFL1 rescues yeast *sec14* mutants. *Arabidopsis*
cpsfl1 mutants are seedling lethal, show reduced thylakoids and a defect in chloroplast vesicle
transport and terpenoid biosynthesis (Hertle et al., 2020; Garcia et al., 2020; Kim et al., 2022). Also,
the vesicle-inducing protein in plastids (VIPP1), a member of the ancient ESCRTIII family of
membrane remodelling proteins, has been shown to engulf PPI-containing membranes *in vitro*,
highlighting its essential role in thylakoid formation (Theis et al., 2019; Gupta et al., 2021). All
together this indicates a function of PPIs in plastids but does not prove their presence.

In this study, our results present evidence for the presence of PPIs within the chloroplasts of tobacco,
as indicated by various PPI biosensors and immunodetection. We observed rapid relocation of PPI
biosensors within the chloroplasts of *Arabidopsis* under abiotic stress conditions. Additionally, plants
expressing chloroplast-targeted PPI modifiers exhibited increased sensitivity to drought and heat
stress. Overall, our investigation sheds light on the existence of PPIs in the chloroplast and their
potential involvement in the PPI signaling pathway during stress conditions.

Results

Generation and sub-organelle localization of a set of PPI biosensors in chloroplasts

Monitoring PPI poses a consistent challenge due to the rapid turnover, tight regulation, and low
quantities of PPIs present within all living cells (Meijer and Munnik, 2003; Jensen et al., 2022).
Indirect evidence indicating the presence of PPIs in the chloroplast is mainly limited to *in vitro* studies
(Gou et al., 2015; Okazaki et al., 2015; Theis et al., 2019; Hertle et al., 2020). Nevertheless, a current
research gap exists concerning the *in vivo* presence of PPIs within chloroplasts.

Genetically encoded biosensors have been widely employed to study the distribution of PPIs in
functional living cells (Varnai and Balla, 2006; Balla, 2013; Simon et al., 2014). Biosensors consist of
lipid-binding domains that interact specifically with known PPIs and can be easily   tracked   by
fusion with a fluorescent protein. To investigate the presence of various PPIs in the lipid composition
of chloroplasts, we generated a collection of biosensors that contain a chloroplast transit peptide
(cTP), yellow fluorescence protein or mCherry respectively, and a lipid-binding domain.   To detect
PI3P, PI4P, PI5P, PI(4,5)P_2 and PI(3,5)P_2 individually and specifically (Figure 1A)   the coding
sequence of pleckstrin homology (PH) domain of human ARHGEF9 for PI5P binding, the PH domain
of human rho guanidine nucleotide exchange factor 7 (ARHGEF7) for PI(3,5)P_2 detection (Singh et
al., 2021), the four phosphate adaptor protein 1 and 2 (PH) domain of rat FAPP1 for PI4P detection,
the PH domain of mouse 1-phosphatidylinositol-4,5-bisphosphate diesterase delta (Plcd1) for
PI(4,5)P_2 detection and putative phosphatidylinositol 3-phosphate binding site encoded by the FYVE
domain from human hepatocyte growth factor related tyrosine kinase substrate (HGS11) (Simon et al.,
2014) was codon optimized for expression in *Arabidopsis* and fused to the C-terminal region of the
chloroplast targeted fluorescent reporter (Figure 1A). Whereas some PH domains bind to more than
one PPIs isoform, the employed domains for PI3P, PI4P and PI4,5P detection have been shown in
numerous *in vivo* studies in plants to be specific for their substrate (Varnai and Balla, 2006; Balla,
2013; Simon et al., 2014). In contrast, specific recognition of PI5P and PI(3,5)P_2 by the lipid binding
domains of ARHGEF7 and 9 respectively has only been shown *in vitro* (Singh et al.2021). Thus,
recognition of more than one PPI isoform might occur *in vivo*.

To visualize their distribution pattern in plastids, we performed a transient expression of each of
the designed biosensor constructs in tobacco leaves using infiltration with transgenic *Agrobacteria* to
initiate transient expression (shown in Figure 1A). 48 hours after Agrobacterium infiltration, YFP
fluorescence was observed in the chloroplast of all transformed cells due to the presence of
chloroplast transit peptide. We further analyzed the sub-organelle distribution of PPI biosensors in the
chloroplasts. We observed a diffuse localization of YFP-PI3P and YFP-PI5P biosensors within the
chloroplast. In contrast, YFP-PI4P, YFP-PI(4,5)P_2, and YFP-PI(3,5)P_2 biosensors to PPI-containing
membrane leads to a different localization reflected in distinct punctate structures within or
surrounding the chloroplasts (Figure 1B and C). Counting puncta in several chloroplasts gave a
constant and characteristic number of puncta for each individual sensor. Whereas both, PI(3,5)P2 and
PI(4,5)P2 biosensors, showed a similar and low number of puncta colocalized with the thylakoid
mediated chlorophyll fluorescence, PI4P sensors localized with a much higher number of smaller
puncta also in a rather peripheral region of the chloroplast as compared to the chlorophyll signal
(Supplementary Figure 1 A, B).

The presence of PI4P in the outer envelope of the chloroplast has been previously demonstrated
(Okazaki et al., 2015). To confirm that the YFP-PI4P biosensors detect their substrate from within the
chloroplast and not at the outer envelope of the chloroplast, we treated isolated chloroplasts from
plants expressing YFP-PI4P biosensor with the protease thermolysin. Thermolysin effectively breaks
down proteins located on the surface of chloroplasts, but it does not penetrate the intermembrane
space (Zufferey et al., 2017). Chloroplasts subjected to thermolysin treatment were subsequently
separated using SDS-PAGE and subjected to immunoblot analysis. The presence of the TOC159
protein band, a known thermolysin-sensitive outer membrane protein, was notably reduced following
thermolysin treatment. In contrast, the YFP-PI4P biosensor protein remained largely unaffected by the
treatment (Supplementary Figure 1C). This observation indicates that the YFP-PI4P biosensor is
mainly associated with membranes within the chloroplasts and is not readily accessible for
thermolysin-mediated degradation at the outer surface membrane. To further support the presence of
PI4P within the inner membranes of chloroplasts, we performed fat blot assays and immunologically
detected PI4P using PI4P specific antibodies (Supplementary Figure 1D). The same antibodies also

recognized lipid extracts from chloroplasts and thylakoids isolated from *Arabidopsis* and spinach
(Supplementary Fig. 1 D).
182

Chloroplast localized-PPI biosensors change their distribution upon PPI phosphatase expression

To further support the specificity of our PPI biosensors, we verified the sub-organelle distribution
pattern of PPI biosensors in the chloroplasts of tobacco plants in the presence of PPI modifiers. For
this purpose, we generated three distinct PPI phosphatases that feature a chloroplast transit peptide.
These phosphatases include SUPPRESSOR OF ACTIN 7 (SAC7) phosphatase and tensin homolog
deleted on chromosome ten (PTEN), and the phosphatase domain of *Drosophila melanogaster*
Oculocerebrorenal Syndrome of Lowe protein (dOCRL) (Goulden et al., 2019; Thole et al., 2008;
Doumane et al., 2021) . SAC7, PTEN, and dOCRL can reversibly dephosphorylate PI4P, PI3P, and
PI5P respectively, by their action as lipid phosphatases (Figure 2A). Co-expression of the PI(4,5)P$_2$
biosensor with *cTP-SAC7* and *cTP-dOCRL*, modifiers specific to PI4P and PI5P, respectively,
resulted in a significant and substantial alteration in the distribution pattern of the PI(4,5)P$_2$ biosensor.
The distinctive punctate structures of the PI(4,5)P$_2$ biosensor were notably reduced and transformed
into diffuse fluorophore signals in the presence of the SAC7 and dOCRL phosphatases (Figure 2 B
and C). In contrast, the punctate-like structures of the PI(4,5)P$_2$ biosensor remained unchanged when
co-expressed with the PI3P modifier, *cTP-PTEN* (Figure 2C). The punctate signals of the PI4P
biosensor within the chloroplasts was also predominantly converted into diffuse fluorophore signals in
the presence of the PI4P modifier SAC7. Since PI3P biosensors were equally distributed when
targeted to plastids, co-expression of PTEN or SAC7 did not show any alternation (Figure 2 D).
Observed changes were quantified and analyzed for statistical relevance (Supplementary Figure 2D).
A significant reduction of fluorescent puncta was observed for PI4,5P2 sensors upon dOCRL
expression. In contrast, co-expression of SAC7 and PI4P sensors resulted in a significant reduction of
the fluorescent puncta but an increase in a diffuse fluorescent signal.
We also examined the distribution of the same PI(4,5)P$_2$ biosensor tagged with different fluorophores
following co-expression. As expected, biosensors with identical substrate, YFP-PI(4,5)P$_2$ and
mCherry-PI(4,5)P$_2$, co-localized with the same fluorescence pattern in the chloroplasts
(Supplementary Figure 2A). Next, we co-expressed biosensors sensitive for PI(3,5)P$_2$ tagged with
YFP and a biosensor specific for PI4P and tagged with mCherry transiently in tobacco leaves. The
sensors differed in their sub-organelle localization and in respect to the chlorophyll autofluorescence.
This supports, that different sensors recognize different substrates in the chloroplast (Supplementary
Figure 2 B).
214

VIPP1 and PI3P sensor proteins colocalize

To date, our understanding of the interaction between chloroplast proteins and PPIs has primarily
relied on *in vitro* studies employing protein-lipid blot overlay assays (Gou et al., 2015; Okazaki et al.,
2015; Theis et al., 2019; Hertle et al., 2020). We further assayed the colocalization of the chloroplast
protein, VESICLE-INDUCING PROTEIN IN PLASTIDS 1 (VIPP1) with individual PPI biosensors.
In vitro studies have already demonstrated the robust binding of *Chlamydomonas reinhardtii* VIPP1
to liposomal membranes containing PI4P, PI(4,5)P$_2$, and PI(3,4,5)P$_3$ (Theis et al., 2019).
To assess the *in vivo* association of VIPP1 with PPIs, we conducted colocalization between VIPP1-
CFP and PPI biosensors within the chloroplasts. Interestingly, the simultaneous expression of *VIPP1-*
CFP and *cTP-mEOS4b-PI3P* led to a significant alteration in the distribution pattern of mEOS4b-
PI3P, resulting in the formation of distinct dot-like structures (Figure 3A). Significantly, these dot-like

structures of mEOS4b-PI3P exhibited co-localization with VIPP1-CFP (Figure 3A). Interestingly, the overrepresentation of VIPP1 did not alter the sub-organelle distribution pattern of the PI(4,5)P$_2$ biosensor (Figure 3A) .

Similarly, co-expression of the *VIPP1-CFP* with *cTP-mCherry-PTEN*, modifier specific to PI3P, resulted in a substantial alteration in the typical distribution pattern of VIPP1 within chloroplasts (Figure 3B). Subsequently, we examined whether the stability of VIPP1 and the configuration of its homooligomeric complexes were affected in the presence of PTEN. As shown in Supplementary Figure 3A, the stability of VIPP1 remained unaltered in plants expressing *cTP-PTEN*. To verify the complex state of VIPP1 in *cTP-PTEN*-expressing plants, thylakoid proteins from both WT and *cTP-PTEN* plants were subjected to treatment with Bis (Sulfosuccinimidyl) suberate (BS^3). BS^3 serves as a membrane-impermeable crosslinking agent to stabilize the complex state. Nevertheless, no significant differences   were observed in terms of the multiple complex states of VIPP1 in *cTP-PTEN*-expressing plants (Supplementary Figure 3B). Our observation suggests that overexpression of VIPP1 may concentrate PI3P lipids in a specific location within the chloroplast, making them detectable by the PI3P biosensor, while in the absence of VIPP1 overexpression, the condensation of PI3P appears to be below the detection threshold of the PI3P biosensor (Figure 1C). Thus, we wanted to proof the sensitivity and specificity of the PI3P sensor by an independent approach. For this we purified chloroplasts and thylakoids purified form *Arabidopsis* and spinach respectively. Following lipid extraction, we performed fat blot assays together with a dilution series of PI3P lipid standards and probed them with PI3P specific FYVE-mEOS4b sensor proteins. Recombinant FYVE sensors recognized PI3P lipid standards as well as chloroplast and thylakoid lipid extracts from both species. Still, with a low concentration (<6.3 pmol) and obscured by a bright spot in the center of the lipid dot resembling the distribution of coextracted chlorophyll. Thus, we fractionated spinach chloroplasts and probed chlorophyll free inner envelope membranes, thlykoids and an array of chloroplast lipids to support specific recognition of PI3P sensor proteins. PI3P and inner chloroplast envelope membranes were recognized with high specificity (Supplementary Figure 3D).

As a last attempt, we investigated the PI3P dependent membrane recognition of VIPP1 within plant lipid-based giant unilamellar vesicles (GUVs). While low binding of VIPP1 was observed in GUVs composed of MGDG, DGDG, PG, and SQDG, the inclusion of 0.1 mol% PI3P into the GUV lipid mixtures led to a pronounced recruitment of recombinant VIPP1 to the GUV surfaces in punctate pattern (Figure 3 C). Next, we confirmed the specific binding of FVYE sensor proteins to GUVs with but not without PI3P in a chloroplast typical lipid composition (Supplementary Figure 3E). However, the punctate and uneven distribution pattern of VIPP1could not be observed for FYVE sensor proteins which suggests a non-homogeneous distribution or local concentration of phosphatidylinositols (PPIs) could be induced by VIPP1 (Figure 3A, C).

Taken together, these results suggest the association and potential function of VIPP1 with PI3P *in vivo*.

The localization of PPI sensors changes upon heat in chloroplasts of *Arabidopsis*

To investigate the presence and function of various PPIs in the chloroplasts, we generated transgenic *Arabidopsis* lines stably expressing our chloroplast localized biosensors (*cTP-YFP-PI4P*, *cTP-YFP-PI5P*, *cTP-YFP-PI(4,5)P$_2$*, and *cTP-YFP-PI(3,5)P$_2$*). Remarkably, none of the transgenic lines showed any visible developmental phenotypes. Next, we assessed the sub-chloroplasts distribution patterns of YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P$_2$, YFP-PI(3,5)P$_2$ biosensors, and cTP-GFP within the chloroplasts of 10-day-old *Arabidopsis* seedlings under standard control conditions using confocal microscopy. We noted that all the PPI sensors displayed a diffuse localization within the chloroplast (Figure 4A). This observation suggests that the biosensors may not have encountered their intended

targets for binding, or alternatively, the concentration of PPIs in the chloroplast of *Arabidopsis*
seedlings might be below the detection limits of our designed PPI biosensors.

Furthermore, we decided to elevate the PPI flux within the cell, potentially including the chloroplast,
and subsequently assess the sub-chloroplast distribution patterns of the PPI biosensors. Studies have
demonstrated that the metabolism of PI and PPIs is significantly activated in response to heat stress in
various organisms (Morales-Johansson et al., 2004; Lu et al., 2020) and diverse plant species
(Mishkind et al., 2009; Annum et al., 2022; Zhu et al., 2022; Annum et al., 2023). To elevate the PPI
flux, we subjected 10-day-old Arabidopsis seedlings to a 40°C temperature treatment for 30 minutes,
after which we conducted confocal microscopy analysis. Surprisingly, we observed that all PPI
biosensors re-localized within the chloroplasts in response to short heat stress treatment. As shown in
Figure 4B, there was no change in the sub-chloroplast localization of GFP in plants expressing GFP
targeted into chloroplasts. In contrast, we observed a clear translocation of sensors to PPI-containing
membranes resulting in punctate-like structures in the plant expressing PPI biosensors (Figure 4B).
Quantification of fluorescence in plastids containing PI4P, PI5P and PI(3,5)P$_2$ sensor proteins showed
similar numbers of fluorescent puncta per chloroplast. Consistent with transient expression PI(4,5)P2
sensor containing plastids showed significantly fewer puncta as compared to other sensor lines
(Supplementary Figure 4 A, B). Although, expression of PPI biosensors is under the control of the
35S promoter, we additionally confirmed, through western blot analysis, that the heat stress treatment
did not result in any noticeable changes in their abundance when compared to the control condition
(Figure 4C). Moreover, we noticed that the distribution patterns of all PPI biosensors remained
unchanged in heat-stressed root cells that lacked fully mature chloroplasts, in comparison to untreated
root cells (Supplementary Figure 4C). These results indicates that increased PPI flux within the cell
upon heat stress may affect PPI levels in the chloroplasts, resulting in different disruption pattern of
PPI biosensors.

Potential function of PPIs as a lipid signaling molecule in chloroplasts under abiotic stresses

Multiple studies have demonstrated stress-induced alterations in PI and PPI signaling, implying the
pivotal roles of PPI signaling in plant adaptation to stress conditions (Hirayama et al., 1995; Meijer et
al., 1999; Pical et al., 1999; DeWald et al., 2001; Meijer et al., 2001; Heilmann, 2009; Munnik and
Vermeer, 2010; Zhu et al., 2022; Annum et al., 2023). Next, to further explore the possible
involvement of various PPIs within the chloroplasts in response to abiotic stress, we generated
transgenic Arabidopsis lines that constitutively express PPI modifiers (phosphatases) containing a
chloroplast transit peptide, namely *cTP-SAC7.OX, cTP-PTEN.OX*, and *cTP-dOCRL.OX*. As
previously mentioned, SAC7, PTEN, and dOCRL are capable of reversibly dephosphorylating PI4P,
PI3P, and PI5P, respectively (Figure 2A). Remarkably, none of the transgenic lines expressing *cTP-*
SAC7, cTP-PTEN, and *cTP-dOCRL* showed any visible developmental phenotypes, nor did they
exhibit any abnormalities when examined with electron microscopy (Figure 5A and B). This
observation aligns with the absence of binding of PPI biosensors to any PPIs in the chloroplast of
plants growing under optimal conditions (Figure 4A).

Subsequently, plant overexpressing plastidial phosphatase were subjected to drought stress by
withholding water. As shown in Figure 5B, plants overexpressing *cTP-SAC7*, *cTP-PTEN*, and *cTP-*
dOCRL began to exhibit leaf wilting symptoms approximately 8 days into the drought period. These
symptoms became progressively more pronounced in the subsequent days of drought compared to the
control plants. Furthermore, the measurements of the maximum quantum efficiency of photosystem
(PS) II (Fv/Fm) in drought-treated plants revealed a photosynthetic impairment in plants expressing

cTP-SAC7, *cTP-PTEN*, and *cTP-dOCRL* when compared to WT (Figure 5A). In addition, the effect of heat stress on PTEN expressing plants was investigated. Exposing seedlings overexpressing *cTP-PTEN* to 30°C for 9 days resulted in pale cotyledons as compared to WT (Supplementary Fig. 5 A). Plants expressing SAC7 did not show a significant difference. Next, we performed an ultrastructural analysis of plants overexpressing PPI modifying enzymes targeted to the plastids. No significant differences were observed when grown under normal growth conditions (Fig. 5B, supplementary Fig. 5B). In contrast,   ultrastructural analysis of drought treated WT and *cTP-PTEN* overexpression lines showed a strong swelling of the stromal fraction and thylakoid membranes indicating altered osmotic pressure within plastids (Figure 5B). Interestingly, similar phenotypes have been described for VIPP1 knockdown lines (Zhang et al., 2012). In order to show that PPI signaling molecules could play a direct role in the phenotype observed when phosphatases are overexpressed, we also analyzed the PPI sensor patterns in response to drought stress using transgenic *Arabidopsis* PPI sensor lines (Figure 5C). As described for heat stress treatments, PI4P, PI(3,5)P_2 and PI(4,5)P_2 expressing lines formed punctate structures upon drought stress treatment (Fig. 5C). Quantification of fluorescent puncta per chloroplast showed a significant difference between PI4P and PI(4,5)P2 and PI(3,5)P2 lines. In contrast, no punctate structures were observed for chloroplasts of PI3P and PI5P lines (Supplementary Fig. 5C).

Taken together this data suggest that PPIs may serve as lipid signaling molecules within chloroplasts, fine-tuning the response to abiotic stresses.

Discussion

Chloroplasts contain the most important membrane system in plant cells, thylakoid membranes, where the light-dependent reaction of photosynthesis takes place. Notably, chloroplast membranes show a distinctive lipid composition, mainly composed out of galactolipids and sulfolipids and very few phospholipids. Whereas recent studies have shown the bulk lipid composition of chloroplasts (LaBrant et al., 2018), presence and distribution of minor or signaling lipids remains elusive. Hence, a better understanding of the lipid composition of thylakoids and more broadly chloroplasts is currently crucial to understand their biogenesis maintenance driving photosynthesis under steady state and stress conditions. To date, presence of PPIs in the lipid composition of chloroplast membranes remains vague. This is most probably due to their low abundance and high turnover rates, which make them difficult to detect. Here, we indeed show for the first time the presence of PPIs within the chloroplasts of tobacco (Figure 1 and 2) and heat stressed *Arabidopsis* plants (Figure 4), using our PPI biosensors.
Synthesis and turnover of PPI pools are vital for regulating various membrane trafficking processes such as thylakoid membrane biogenesis and maintenance, along with signaling transduction pathways (Meijer and Munnik, 2003; Di Paolo and De Camilli, 2006; Heilmann, 2009). Therefore, it is likely that arranging PPI flux within the chloroplasts serves a similar role. However, it remains unclear if PPIs can be produced within the chloroplasts. If so, it suggests the potential presence of unknown PI kinases and PPI phosphatases within this organelle. ARC3, a chloroplast division factor, represents a chimera of prokaryotic FtsZ and part of eukaryotic phosphatidyl-4-phosphate 5-kinase and localizes to the stroma (Shimada et al. 2004, Maple et al., 2007). Whether it functions as PI Kinase remains so far unexplored. Alternatively, PPI might be transferred from ER to chloroplast envelopes through membrane contact sites (LaBrant et al., 2018) or the action of flipases and flopases (Rajasekharan and Gummadi, 2011). Once inside the inner leaflet of the inner envelope membrane PPIs could subsequently be transported within the chloroplast via the action of lipid binding proteins or envelope vesicles (LaBrant et al., 2018, Hertle et al., 2021, Schroda et al., 2021). So far, interactions between

PPIs and chloroplast proteins, PDV1 and PDV2, Chloroplast Sec14-like protein 1 (CPSFL1), WHEAT KINASE START1 (WKS1), and VIPP1 have only been shown *in vitro*, using protein-lipid blot overlay assays (Gou et al., 2015; Okazaki et al., 2015; Theis et al., 2019; Hertle et al., 2020). PDV1 and PDV2 are integral components of the chloroplast division machinery, and they are responsible for regulating the rate of chloroplast division (Okazaki et al., 2009). CPSFL1 plays a crucial role in ensuring proper chloroplast development, promoting plant growth, and supporting optimal photosynthetic function in Arabidopsis and *Chlamydomonas* (García-Cerdán et al., 2020; Hertle et al., 2020; Kim et al., 2022). WKS1 provides resistance to *Puccinia striiformis f. sp tritici* in wheat (*Triticum aestivum).* VIPP1 plays essential roles in thylakoid membrane biogenesis, maintenance and repair under both normal and stress conditions in *Arabidopsis* and *Chlamydomonas* (Nordhues et al., 2012; Zhang et al., 2016; Theis et al., 2020). Here we demonstrated the co-localization of VIPP1 and the PI3P biosensor within the chloroplasts (Figure 3A). Additionally, we observed alterations in the sub-chloroplast distribution pattern of VIPP1 in the presence of the PI3P modifier, PTEN (Figure 3B). Our findings showed that plants expressing *cTP-PTEN* were sensitive to heat and drought stress and exhibited reduced photosynthetic efficiency (Figure 5A). Interestingly, ultrastructural analysis of *ctp-PTEN* overexpressing plants indicated osmolarity changes by swollen plastids and thylakoids (Figure 5B). Given the critical role of VIPP1 in thylakoid maintenance and repair under stress conditions (Zhang et al., 2016; Theis et al., 2020), it is possible that PI_3P may influence the function of VIPP1 under stress conditions. However, we did not detect any changes in VIPP1 stability or its association with multiple complexes in *cTP-PTEN*-expressing plants (Supplementary Figure 4). Nevertheless, the *in vivo* association of PI3P with VIPP1 (Figure 3) might fine-tune the function of VIPP1 under stress conditions.

In fact, phosphoinositide signaling pathways are a pivotal component mediating plant responses to various forms of stresses (Sui et al., 2008; Xue et al., 2009; Munnik and Vermeer, 2010; Zhai et al., 2013; Morales et al., 2019). Extensive research has confirmed that elevated temperatures trigger PPI flux in plant cells. This phenomenon is not limited to the model plant *Arabidopsis* (Mishkind et al., 2009; Annum et al., 2023) but extends across the plant kingdom, encompassing species such as rice (Mishkind et al., 2009), wheat (Annum et al., 2022)*, Cycas multipinnata*, and *C. panzhihuaensis* (Zhu et al., 2022). Hence, the inducibility of PPI flux within the cells in response to heat stress appears to be evolutionarily conserved, indicating the crucial role of these phospholipids in the response to high-temperature stress. We verified, that high temperature and subsequently induction of PPI flux within the cell leads to relocation of PPI biosensors, indicating an increase in PPI concentration inside the chloroplasts (Figure 4). Thus, changing cytoplasmic PI or PPI pools most probably also affects PI and PPI flux in the plastid.

Drought induces water deficiency and hinders plant growth. Numerous studies have illustrated the crucial role of the phosphoinositide signaling pathway in drought stress across different plants (Mikami et al., 1998; Perera et al., 2008; Wang et al., 2008; Zhai et al., 2013; Carpaneto et al., 2017). The instant induction of *AtPIP5K1*, Phosphatidylinositol-4-phosphate 5-kinase, transcript in Arabidopsis plants subjected to drought or abscisic acid treatment suggests the participation of AtPIP5K1 in altering PPI pools and, consequently, its involvement in water-stress signal transduction (Mikami et al., 1998). Furthermore, dehydration induces the levels of PI5P through the activity of a myotubularin homolog (*AtMTM1*). MTM1 is itself induced during dehydration and can produce PI5P from $PI(3,5)P_2$ (Ndamukong et al., 2010). Moreover, it has been reported that drought stress induces synthesis of $PI(3,5)P_2$ in plants (Meijer et al., 1999) in which $PI(3,5)P_2$ controls the decrease in vacuolar volume to counteract water deficiency and facilitates the closure of stomata (Carpaneto et al., 2017). Analysis of expression patterns showed that six out of nine Arabidopsis *PLC* genes are induced by dehydration (Tasma et al., 2008). Similarly, the expression of maize *ZmPLC1* is induced under dehydration, and this increased expression contributes to the improved drought tolerance of

maize (Wang et al., 2008). In addition, it has been reported that *Arabidopsis* transgenic plants, overexpressing mammalian inositol polyphosphate 5-phosphatase display modified responses to abscisic acid and significantly improved drought tolerance (Perera et al., 2008). These studies provide evidence for a close association between phosphoinositide signaling pathways and water-stress responses in plants. We observed that perturbation of PPI balance in chloroplasts upon dehydration in plant overexpressing different PPI phosphatases resulted in drought sensitivity (Figure 5). Our results suggest the generation and turnover of PPI pools within the chloroplasts may play crucial roles in regulating signaling transduction pathways.

The alterations in PPI levels or localization triggered by dehydration can be detected by downstream protein targets possessing PPI-binding domains, consequently altering their activity or localization (Munnik and Vermeer, 2010). For instance, the activity of the *Arabidopsis* trithorax-like factor ATX1 was modified upon dehydration stress due to the binding of PI5P (Ndamukong et al., 2010). Hence, further research is required to comprehensively characterize chloroplast proteins possessing PIP binding domains.

While organelles have distinct specialized functions as individual compartments, there is evident that inter-organelle communication plays a vital role in preserving cellular function. This collaborative interaction occurs through various membrane-mediated processes and signaling molecules. It has been reported that phospholipids are a notable class of molecules that promote these interactions (Meijer and Munnik, 2003; Ravi et al., 2021). Hence, it is reasonable to hypothesize that PPIs may play a role in mediating cross-communication between the chloroplast and other cellular organelles, such as the nucleus. Comprehensive research efforts are required to elucidate the involvement of PPIs in orchestrating signaling pathways and regulating crosstalk between chloroplasts and other organelles.

Material and methods

Generation of transgenic plants

For fluorescent reporters the cTP of RbcS (sequence in supplement) was amplified using oligos (AH1 and AH2) containing recombination sites for InFusion cloning into pDK366 linearized by SpeI. Resulting plasmids were linearized by SpeI and synthetic DNA of YFP, mCherry, CFP or mEOS4b respectively (and codon optimized for expression in A. thaliana) was inserted in frame to the n-terminal cTP by InFusion cloning. The resulting plamids were used to create chloroplast targeted fluorescent lipid sensor and lipid modifying proteins. In brief, the coding sequence of the lipid binding domains of the pleckstrin homology (PH) domain of human ARHGEF9 for PI5P binding, switch-associated protein 70 PH domain of human differentially expressed in FDCP6 homolog guanidine exchange factor DEF6 for PI3,4P detection, the PH domain of human rho guanidine nucleotide exchange factor 7 (ARHGEF7) for PI3,5P detection (Singh et al., 2021), the four phosphate adaptor protein 1 and 2 (PH) domain of rat FAPP1 for PI4P detection, the PH domain of mouse 1-phosphatidylinositol-4,5-bisphosphate diesterase delta (Plcd1) for PI4,5P detection and putative phosphatidylinositol 3-phosphate binding site encoded by the FYVE domain from human hepatocyte growth factor related tyrosine kinase substrate (HGS11) (Simon et al., 2014) was codon optimized for expression in Arabidopsis were codon optimized and synthesized   with recombination sites (sequences in supplement) to be integrated into pDK366 containing cTP_YFP, cTP_mCherry or cTP_mEOS4b fusion proteins under the control of the 35S promotor by InFusion clonng.   , mCherry-FKBP-PTEN was a gift from Gerry Hammond (Addgene plasmid # 116866 ; http://n2t.net/addgene:116866 ; RRID:Addgene_116866)   , SAC7/RHD4 (At3g51460) was amplified from Arabidopsis thaliana cDNA. dOCRL was used as codon optimized synthetic DNA of

the sequence publishes in Doumane et al., 2021. VIPP1 was amplified from Arabidopsis cDNA using
oligos (AH15/AH16). Subsequent VIPP1 lacking the stop codon was fused in frame to codon
optimized CFP (sequence in supplement) by fusion PCR using oligos coding for recombination sites
for infusion cloning into pDK488 linearuzed by SpeI   . For stable expression, we introduced the
constructs into *Arabidopsis* (Col-0) through a floral dip method utilizing Agrobacterium tumefaciens
(strain GV3101)(Clough and Bent, 1998). Transient expression in *N. benthamiana* was conducted
following the established procedure (Li, 2011).

Plant materials, growth conditions, and the procedures for heat and drought treatments

Arabidopsis thaliana Columbia (Col-0) was used as the WT. The seeds were germinated on 0.5×
Murashige and Skoog agar medium enriched with 1% (w/v) sucrose. The seedlings were cultivated
following a 16-hour light cycle (120 μmol $m^{-2} s^{-1}$) at 22 °C, and 8 h dark at 22 °C. To induce PIP flux
with the cells, 10-day-old Arabidopsis seedlings were subjected to a heat regime of 40 °C for 30 min.
To assess heat stress tolerance, 7-day-old Arabidopsis seedlings were exposed to a heat regime of 45.5
°C for 45 min. To assess drought tolerance, the seedlings were cultivated for 4 weeks in soil under a
diurnal cycle comprising 16 hours of light (120 μmol $m^{-2} s^{-1}$) and 8 hours of darkness at 22 °C.
Subsequently, to induce drought stress, watering was discontinued for a period of 10-14 days.

Chloroplast or thylakoid Isolation

To isolate chloroplasts, leaf material from the various genotypes was initially ground in chloroplast
isolation buffer. The buffer composition included 0.45 M sorbitol, 20 mM Tricine (pH 8.4), 10 mM
EDTA, 10 mM NaHCO3, and 0.1% BSA. Chloroplasts were collected by centrifugation for 5 min at
3000g, 4°C from the collected material obtained by filtering it through two layers of Miracloth. The
released chloroplasts were then carefully layered onto a Percoll gradient made with 5X resuspension
buffer (0.3 M sorbitol, 20 mM Tricine (pH 8.4), 2.5 mM EDTA, and 5 mM MgCl2). Centrifugation
was performed in a swing-out rotor at 4°C for 10 minutes at 3300g. Intact chloroplasts were collected
from the lower green band and subsequently subjected to a wash using 1X resuspension buffer.
Thylakoid membrane preparation from leaves of 3-week-old WT and *PTEN.OX* plants was conducted
in accordance with the method described in (Bassi et al., 1985). Leaf material was homogenized in an
ice-cold buffer comprising 0.4 M sorbitol, 0.1 M Tricine/KOH (pH 7.8), and 1 mM PMSF. The
resulting homogenate was filtered through a nylon mesh and then centrifuged at 4°C, 3000g for 10
minutes. Subsequently, chloroplasts were disrupted in a buffer consisting of 20 mM HEPES/KOH
(pH 7.8) and 10 mM EDTA (pH 8.0). Thylakoids were collected by centrifugation at 12,000g for 10
minutes and resuspended in a buffer containing 50% glycerol, 10 mM HEPES/KOH (pH 7.5), and 1
mM EDTA (pH 8.0). Fractionation of spinach chloroplasts was performed as described in (Hertle et.
al 2020). Envelope fractions were separated into inner and outer envelopes using sucrose gradients as
discribed in (Block et al., 2018).

Thermolysin treatment

495 Intact chloroplasts were treated with thermolysin based on Fitzpatrick et al., 2001. The chloroplast
496 pellet, equivalent to 30 μg of chlorophyll was resuspended in import buffer (330-mM sorbitol and 50-

497 mM HEPES–KOH, pH 8.0) and incubated with thermolysin (36   units mg^{-1}; 0.05 units μg^{-1}
498 chlorophyll) for 30 minutes on ice. Intact chloroplasts were re-isolated over a 40% v/v Percoll.
499 chloroplasts were dissolved in 100ul of 4xSDS buffer.

Bis(sulfosuccinimidyl)suberate (BS3) treatment

Isolated thylakoids from WT and *PTEN.OX* were washed 5 times with 20 mM HEPES/ KOH pH 7.5 to remove the EDTA. Crosslinking was performed by adding 2mM BS3 to a thylakoid suspension containing 150 µg/ml of chlorophyll, followed by incubation on ice for 20 minutes. The reaction was stopped with 150 mM TRIS HCL pH:7.5. The plates were dissolved in 5X SDS sample buffer for further analysis.

Protein extraction and immunoblotting

Soluble proteins were extracted from Arabidopsis seedlings expressing PIP biosensors following the method outlined in (Sedaghatmehr et al., 2016). The protein concentration was determined using the Bradford protein assay. Subsequently, the proteins were separated by SDS-PAGE, and the PageRule Plus Prestained Protein Ladder (10–250 kDa; Thermo Scientific) served as the molecular weight marker.

For immunoblot analysis, the proteins were transferred onto either a Protan nitrocellulose or Polyvinylidene Difluoride membrane. Primary antibodies used included a rabbit anti-VIPP1 polyclonal antibody from   Agrisera, rabbit anti-TOC159 from Agrisera, and mouse anti-green fluorescent protein (GFP) antibody from Clon   tech. Secondary antibodies, IRDye 800CW-conjugated goat anti-rabbit or anti-mouse IgG (H+L), were employed at a 1:10,000 dilution (LI-COR Biosciences). Band detection was performed using the Odyssey Infrared Imaging System (LI-COR Biosciences).

GUV formation

To form GUVs, PVA assisted swelling was performed in PBS buffer according to Weinberger et al., 2013. PVA-coated substrate was prepared by spreading and drying of 100–300 μL of PVA (5%) solution on a glass slide. Afterwards 10–20 μL of plant lipid mixtures (2 mM final) of MGDG/DGDG/SQDG/PG/PI (55/27/7/10/1) according to Block et al,.1983, with or without 0.1% PI3P and with 0.01mol% DiI. dissolved in chloroform were spead on the dried PVA film and placed under vacuum for 30 min for solvent evaporation. A chamber was formed using vitrex and a coverglass filled with sucrose (200 mM) containing PBS buffer. Following 15 min. incubation, GUVs were transferred into an Eppendorf tube using a pipette until further use. For protein binding studies proteins were added in PBS containing glucose in equal osmolarity in a final concentration of 0.5 µM.

Expression of recombinant his tagged VIPP1-mEOS4b and FYVE-mEOS4b

VIPP1 lacking the cTP and the stop codon was amplified and fused to mEOS4b by fusion PCR using the primers AH13 and AH14. mEOS-FYVE was amplified from plant plasmids described above using primers overlapping to pET28b. The PCR product was integrated into pET28b linearized by restriction using BamHI and EcoRI by in Fusion cloning. Resulting plasmid was used for inducible expression of 6xHis tagged VIPP1-mEOS. VIPP1 was expressed by adding 0,5 mM IPTG to a culture of O.D.600=0.5 grown at 37°C. The temperature was lowered to 30°C and the culture was grown for 3 more hours. The bacteria were harvested by centrifugation and the pellet was snap frozen in liquid nitrogen and stored at -70°C until further use. Recombinant protein purification was performed using Ni-NTA(Qiagen) according to the manufacturers protocol for native proteins. The lysis buffer was supplemented with 2% Triton-X-100 (Serva). To remove access detergent and imidazole, buffer exchange of the obtained purified protein solution was achieved by 10x centrifugation of the solution

543 through a 10.000 MW cut off centricon (Millipore) using PBS. The protein was stored at -70°C until
544 used.

Fat blot assays

546 Fat blot assays were performed as discribed in (Munnik and Wierzchowiecka, 2013). FYVE-mEOS
547 was detected using a gel doc fluorescence blot reader (Biorad) at 488 nm excitation.

TEM microscopy

Leaf samples were fixed in 2.5% (w/v) glutaraldehyde, 2% (w/v) paraformaldehyde in 50 mM sodium
cacodylate (pH 7.4) for 1 hr under vacuum at RT. Subsequent fixation was continued overnight at
4°C. After washing in PBS Post-fixation with was performed in 2% OsO4 and 0.8% $K_3Fe(CN)_6$ in
50 mM cacodylate buffer (pH 7.4) for 2 hr at RT. After rinsing, leaf samples were post-stained with
2% (w/v) uranyl acetate for 2 h at RT. Subsequent dehydration was perfomed using acetone and
Spurr`s resin (Science Services) was infiltered. Polymerisation was carried out at 60°C for 48hrs.
Ultrathin sections (50-70 nm) were cut on a UC-6 ultramicrotome (Leica) using a diamond knife. For
transmission electron microscopy, sections were deposited on formvar and carbon coated nickel grids
and contrasted with 2% uranyl acetate and Reynold´s lead citrate. Images were acquired on a JEOL
JEM F-200 (JEOL).

Confocal microscopy

560 Plants were imaged using a Leica TCS SP8 or SP5 confocal microscope. The excitation/emission
561 settings used were 405/415-475nm for CFP, 488/505-535 nm for GFP, 512/520-540 nm for YFP and
562 mEOS4b, 561/570-600 nm for RFP and DiI, and 633/650-720 nm for chlorophyll fluorescence. All
563 images were acquired using sequential scanning settings. The images analysis and quantification of
564 fluorescence intensities was performed using the ImageJ software.

Acknowledgments

We thank the Deutsche Forschungsgemeinschaft (DFG) and the Max Planck Society (MPG) for
funding. We thank Prof. Dr. Ralph Bock and his team from the Max Planck Institute of Molecular
Plant Physiology (MPIMP, Potsdam, Germany) for supporting and hosting the project and engaging
in insightful discussions.

We thank Dr. Nadja Tarakina and Bolortuya Badamdorj from the Max Planck Institute of colloids
and interfaces for providing access and excellent technical assistance to the transmission electron
microscope.

We thank Dr. Rumiana Dimova and Dr. Shreya Pramanik at Max Planck Institute of colloids and
interfaces (MPIKGF, Potsdam, Germany) for help and discussions with plant GUV experiments

We thank Prof. Petra Bauer at the Institute of Botany, at the Heinrich Heine University (Düsseldorf,
Germany) for hosting the project and discussions.

Author contributions

580 AH conceived the study. MS and AH designed and performed the experiments; FR generated PIP
581 biosensor constructs. MS and AH wrote   the manuscript   . All authors agreed to the submission of
582 the manuscript.

Funding

The project was funded by a research grant of the Deutsche Forschungsgemeinschaft (DFG), (HE8905/2-1) projektnummer: 452589609 and the Max Planck Society (MPG). The work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy – EXC-2048/1 – project ID 390686111. The work was supported by the Deutsche Froschungsgemeinschaft (DFG, German Research Foundation) Project no. 267205415-SFB 1208.

Conflict of interest

The authors declare no conflict of interest.

Supplemental data

The following materials are available in the online version of this article.

Supplementary Figure 1. Intensity plots of PPI biosensors in chloroplasts following transient expression in tobacco.

Supplementary Figure 2. Colocalization of PPI biosensors with two different fluorescent proteins in the chloroplasts.

Supplementary Figure 3. VIPP1 stability and/or its multiple complexes in *PTEN-expressing* plants and PI3P detection in chloroplasts.

Supplementary Figure 4. Distribution pattern of YFP-PI4P, YFP-PI5P, YFP-PI4,5P, and YFP-PI3,5P biosensors in   the chloroplasts of Arabidopsis under heat stress conditions.

Supplementary Figure 5. Distribution pattern of YFP-PI4P, YFP-PI5P, YFP-PI4,5P, and YFP-PI3,5P biosensors in chloroplasts of Arabidopsis under drought stress conditions.

Supplementary Table 1. Oligos used in the study

Supplementary Table 2. Sequences of synthetic DNA or fluorescent reporters used in the study

References:

Akhter S, Uddin MN, Jeong IS, Kim DW, Liu XM, Bahk JD (2016) Role of Arabidopsis AtPI4Kγ3, a type II phosphoinositide 4-kinase, in abiotic stress responses and floral transition. Plant Biotechnol J **14:** 215-230

Annum N, Ahmed M, Imtiaz K, Mansoor S, Tester M, Saeed NA (2022) (32)P(i) Labeled Transgenic Wheat Shows the Accumulation of Phosphatidylinositol 4,5-bisphosphate and Phosphatidic Acid Under Heat and Osmotic Stress. Front Plant Sci **13:** 881188

Annum N, Ahmed M, Tester M, Mukhtar Z, Saeed NA (2023) Physiological responses induced by phospholipase C isoform 5 upon heat stress in Arabidopsis thaliana. Front Plant Sci **14:** 1076331

Balla T (2013) Phosphoinositides: tiny lipids with giant impact on cell regulation. Physiol Rev **93:**
1019-1137
Bankaitis VA, Aitken JR, Cleves AE, Dowhan W (1990) An essential role for a phospholipid transfer
protein in yeast Golgi function. Nature. 347:561-2.
Bassi R, dal Belin Peruffo A, Barbato R, Ghisi R (1985) Differences in chlorophyll-protein complexes
and composition of polypeptides between thylakoids from bundle sheaths and mesophyll
cells in maize. Eur J Biochem **146:** 589-595
Block MA. (2018) Isolation of Inner and Outer Membranes of the Chloroplast Envelope from Spinach
and Pea. Methods Mol Biol 1829:137-144
Block MA, Dorne AJ, Joyard J, Douce R. (1983) Preparation and characterization of membrane
fractions enriched in outer and inner envelope membranes from spinach chloroplasts. II.
Biochemical characterization. J Biol Chem. **258**:13281-6.
Bovet L, Müller MO, Siegenthaler PA. (2001) Three distinct lipid kinase activities are present in
spinach chloroplast envelope membranes: phosphatidylinositol phosphorylation is sensitive
to wortmannin and not dependent on chloroplast ATP. Biochem Biophys Res Commun. 289:
269-75
Carpaneto A, Boccaccio A, Lagostena L, Di Zanni E, Scholz-Starke J (2017) The signaling lipid
phosphatidylinositol-3,5-bisphosphate targets plant CLC-a anion/H(+) exchange activity.
EMBO Rep **18:** 1100-1107
Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J **16:** 735-743
DeWald DB, Torabinejad J, Jones CA, Shope JC, Cangelosi AR, Thompson JE, Prestwich GD, Hama H
(2001) Rapid Accumulation of Phosphatidylinositol 4,5-Bisphosphate and Inositol 1,4,5-
Trisphosphate Correlates with Calcium Mobilization in Salt-Stressed Arabidopsis. Plant
Physiology **126:** 759-769
Di Paolo G, De Camilli P (2006) Phosphoinositides in cell regulation and membrane dynamics.
Nature **443:** 651-657
Dieck CB, Boss WF, Perera IY (2012) A role for phosphoinositides in regulating plant nuclear
functions. Front Plant Sci **3:** 50
Doumane M, Lebecq A, Colin L, Fangain A, Stevens FD, Bareille J, Hamant O, Belkhadir Y, Munnik T,
Jaillais Y, Caillaud MC (2021). Inducible depletion of PI(4,5)P_2 by the synthetic iDePP system
in *Arabidopsis*. *Nat. Plants* **7**: 587–597.

García-Cerdán JG, Schmid EM, Takeuchi T, McRae I, McDonald KL, Yordduangjun N, Hassan AM,
Grob P, Xu CS, Hess HF, Fletcher DA, Nogales E, Niyogi KK (2020) Chloroplast Sec14-like 1
(CPSFL1) is essential for normal chloroplast development and affects carotenoid
accumulation in <i>Chlamydomonas</i>. Proceedings of the National Academy of Sciences
117: 12452-12463
Gerth K, Lin F, Daamen F, Menzel W, Heinrich F, Heilmann M (2017) Arabidopsis
phosphatidylinositol 4-phosphate 5-kinase 2 contains a functional nuclear localization
sequence and interacts with alpha-importins. Plant J **92:** 862-878
Gou JY, Li K, Wu K, Wang X, Lin H, Cantu D, Uauy C, Dobon-Alonso A, Midorikawa T, Inoue K,
Sánchez J, Fu D, Blechl A, Wallington E, Fahima T, Meeta M, Epstein L, Dubcovsky J (2015)
Wheat Stripe Rust Resistance Protein WKS1 Reduces the Ability of the Thylakoid-Associated
Ascorbate Peroxidase to Detoxify Reactive Oxygen Species. Plant Cell **27:** 1755-1770
Goulden BD, Pacheco J, Dull A, Zewe JP, Deiters A, Hammond GRV (2019). A high-avidity biosensor
reveals plasma membrane PI(3,4)P_2 is predominantly a class I PI3K signaling product. J Cell
Biol. **218**:1066-1079.
Gupta TK, Klumpe S, Gries K, Heinz S, Wietrzynski W, Ohnishi N, Niemeyer J, Spaniol B, Schaffer M,
Rast A, Ostermeier M, Strauss M, Plitzko JM, Baumeister W, Rudack T, Sakamoto W,
Nickelsen J, Schuller JM, Schroda M, Engel BD (2021). Structural basis for VIPP1
oligomerization and maintenance of thylakoid membrane integrity. Cell. **184**:3643-3659

Heilmann I (2009) Using genetic tools to understand plant phosphoinositide signalling. Trends in
Plant Science **14:** 171-179
Hertle AP, García-Cerdán JG, Armbruster U, Shih R, Lee JJ, Wong W, Niyogi KK (2020) A Sec14
domain protein is required for photoautotrophic growth and chloroplast vesicle formation in
<i>Arabidopsis thaliana</i>. Proceedings of the National Academy of Sciences **117:** 9101-
9111
Hirayama T, Ohto C, Mizoguchi T, Shinozaki K (1995) A gene encoding a phosphatidylinositol-
specific phospholipase C is induced by dehydration and salt stress in Arabidopsis thaliana.
Proc Natl Acad Sci U S A **92:** 3903-3907
Horton P, Park KJ, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai K (2007) WoLF PSORT:
protein localization predictor. Nucleic Acids Res **35:** W585-587
Jensen JB, Falkenburger BH, Dickson EJ, de la Cruz L, Dai G, Myeong J, Jung S-R, Kruse M, Vivas O,
Suh B-C, Hille B (2022) Biophysical physiology of phosphoinositide rapid dynamics and
regulation in living cells. Journal of General Physiology **154**
Kim E-H, Poudyal RS, Lee K-R, Yu H, Gi E, Kim HU (2022) Chloroplast-localized PITP7 is essential for
plant growth and photosynthetic function in Arabidopsis. Physiologia Plantarum **174:**
e13760
LaBrant E, Barnes AC, Roston RL (2018) Lipid transport required to make lipids of photosynthetic
membranes. Photosynthesis Research **138:** 345-360
Leshem Y, Seri L, Levine A (2007) Induction of phosphatidylinositol 3-kinase-mediated endocytosis
by salt stress leads to intracellular production of reactive oxygen species and salt tolerance.
The Plant Journal **51:** 185-197
Li L, Wang F, Yan P, Jing W, Zhang C, Kudla J, Zhang W (2017) A phosphoinositide-specific
phospholipase C pathway elicits stress-induced Ca(2+) signals and confers salt tolerance to
rice. New Phytol **214:** 1172-1187
Li X (2011) Infiltration of Nicotiana benthamiana protocol for transient expression via
Agrobacterium. Bio-protocol**:** e95-e95
Lu K-Y, Pasaje CFA, Srivastava T, Loiselle DR, Niles JC, Derbyshire E (2020) Phosphatidylinositol 3-
phosphate and Hsp70 protect Plasmodium falciparum from heat-induced cell death. eLife **9:**
e56773
Maple J, Vojta L, Soll J, Møller SG. (2007) ARC3 is a stromal Z-ring accessory protein essential for
plastid division. EMBO Rep 8(3):293-9
Matsuoka K, Bassham DC, Raikhel NV, Nakamura K (1995) Different sensitivity to wortmannin of
two vacuolar sorting signals indicates the presence of distinct sorting machineries in tobacco
cells. J Cell Biol **130:** 1307-1318
Meijer HJ, Berrie CP, Iurisci C, Divecha N, Musgrave A, Munnik T (2001) Identification of a new
polyphosphoinositide in plants, phosphatidylinositol 5-monophosphate (PtdIns5P), and its
accumulation upon osmotic stress. Biochem J **360:** 491-498
Meijer HJG, Divecha N, van den Ende H, Musgrave A, Munnik T (1999) Hyperosmotic stress induces
rapid synthesis of phosphatidyl-D-inositol 3,5-bisphosphate in plant cells. Planta **208:** 294-
298
Meijer HJG, Munnik T (2003) Phospholipid-Based Signaling in Plants. Annual Review of Plant Biology
54: 265-306
Mikami K, Katagiri T, Iuchi S, Yamaguchi-Shinozaki K, Shinozaki K (1998) A gene encoding
phosphatidylinositol-4-phosphate 5-kinase is induced by water stress and abscisic acid in
Arabidopsis thaliana. Plant J **15:** 563-568
Mishkind M, Vermeer JEM, Darwish E, Munnik T (2009) Heat stress activates phospholipase D and
triggers PIP2 accumulation at the plasma membrane and nucleus. The Plant Journal **60:** 10-
21

Morales-Johansson H, Jenoe P, Cooke FT, Hall MN (2004) Negative regulation of
phosphatidylinositol 4,5-bisphosphate levels by the INP51-associated proteins TAX4 and
IRS4. J Biol Chem **279:** 39604-39610

Morales JA, Gonzalez-Kantun WA, Rodriguez-Zapata LC, Ramón-Ugalde J, Castano E (2019) The
effect of plant stress on phosphoinositides. Cell Biochemistry and Function **37:** 553-559

Müller MO, Meylan-Bettex M, Eckstein F, Martinoia E, Siegenthaler PA, Bovet L. (2000) Lipid
phosphorylation in chloroplast envelopes. Evidence for galactolipid CTP-dependent kinase
activities. J Biol Chem 275: 19475-81

Mueller-Roeber B, Pical C (2002) Inositol Phospholipid Metabolism in Arabidopsis. Characterized
and Putative Isoforms of Inositol Phospholipid Kinase and Phosphoinositide-Specific
Phospholipase C. Plant Physiology **130:** 22-46

Munnik T, Meijer HJ (2001) Osmotic stress activates distinct lipid and MAPK signalling pathways in
plants. FEBS Lett **498:** 172-178

Munnik T, Wierzchowiecka M. (2013) Lipid-binding analysis using a fat blot assay. Methods Mol Biol
1009:253-9

Munnik T, Vermeer JE (2010) Osmotic stress-induced phosphoinositide and inositol phosphate
signalling in plants. Plant Cell Environ **33:** 655-669

Ndamukong I, Jones DR, Lapko H, Divecha N, Avramova Z (2010) Phosphatidylinositol 5-phosphate
links dehydration stress to the activity of ARABIDOPSIS TRITHORAX-LIKE factor ATX1. PLoS
One **5:** e13396

Nordhues A, Schöttler MA, Unger AK, Geimer S, Schönfelder S, Schmollinger S, Rütgers M, Finazzi
G, Soppa B, Sommer F, Mühlhaus T, Roach T, Krieger-Liszkay A, Lokstein H, Crespo JL,
Schroda M (2012) Evidence for a role of VIPP1 in the structural organization of the
photosynthetic apparatus in Chlamydomonas. Plant Cell **24:** 637-659

Okazaki K, Kabeya Y, Suzuki K, Mori T, Ichikawa T, Matsui M, Nakanishi H, Miyagishima SY (2009)
The PLASTID DIVISION1 and 2 components of the chloroplast division machinery determine
the rate of chloroplast division in land plant cell differentiation. Plant Cell **21:** 1769-1780

Okazaki K, Miyagishima SY, Wada H (2015) Phosphatidylinositol 4-phosphate negatively regulates
chloroplast division in Arabidopsis. Plant Cell **27:** 663-674

Perera IY, Hung C-Y, Moore CD, Stevenson-Paulik J, Boss WF (2008) Transgenic Arabidopsis Plants
Expressing the Type 1 Inositol 5-Phosphatase Exhibit Increased Drought Tolerance and
Altered Abscisic Acid Signaling. The Plant Cell **20:** 2876-2893

Phillips SE, Vincent P, Rizzieri KE, Schaaf G, Bankaitis VA, Gaucher EA (2006) The diverse biological
functions of phosphatidylinositol transfer proteins in eukaryotes. Crit Rev Biochem Mol Biol
41: 21-49

Pical C, Westergren T, Dove SK, Larsson C, Sommarin M (1999) Salinity and Hyperosmotic Stress
Induce Rapid Increases in Phosphatidylinositol 4,5-Bisphosphate, Diacylglycerol
Pyrophosphate, and Phosphatidylcholine in Arabidopsis thaliana Cells*. Journal of Biological
Chemistry **274:** 38232-38240

Pribat A, Sormani R, Rousseau-Gueutin M, Julkowska Magdalena M, Testerink C, Joubès J,
Castroviejo M, Laguerre M, Meyer C, Germain V, Rothan C (2011) A novel class of PTEN
protein in Arabidopsis displays unusual phosphoinositide phosphatase activity and efficiently
binds phosphatidic acid. Biochemical Journal **441:** 161-171

Rajasekharan A, Gummadi SN. (2011) Flip-flop of phospholipids in proteoliposomes reconstituted
from detergent extract of chloroplast membranes: kinetics and phospholipid specificity. PLoS
One6(12):e28401

Ravi A, Palamiuc L, Emerling BM (2021) Crucial Players for Inter-Organelle Communication: PI5P4Ks
and Their Lipid Product PI-4,5-P(2) Come to the Surface. Front Cell Dev Biol **9:** 791758

Sedaghatmehr M, Mueller-Roeber B, Balazadeh S (2016) The plastid metalloprotease FtsH6 and
small heat shock protein HSP21 jointly regulate thermomemory in Arabidopsis. Nature
Communications **7:** 12439

Shimada H, Koizumi M, Kuroki K, Mochizuki M, Fujimoto H, Ohta H, Masuda T, Takamiya K. (2004) ARC3, a chloroplast division factor, is a chimera of prokaryotic FtsZ and part of eukaryotic phosphatidylinositol-4-phosphate 5-kinase. Plant Cell Physiol 45(8):960-7

Siegenthaler PA, Müller MO, Bovet L. Evidence for lipid kinase activities in spinach chloroplast envelope membranes (1997). FEBS Lett. 416: 57-60

Simon ML, Platre MP, Assil S, van Wijk R, Chen WY, Chory J, Dreux M, Munnik T, Jaillais Y (2014) A multi-colour/multi-affinity marker set to visualize phosphoinositide dynamics in Arabidopsis. Plant J **77:** 322-337

Singh N, Reyes-Ordoñez A, Compagnone MA, Moreno JF, Leslie BJ, Ha T, Chen J. (2021) Redefining the specificity of phosphoinositide-binding by human PH domain-containing proteins. Nat Commun:12:4339.

Sui Z, Niu L, Yue G, Yang A, Zhang J (2008) Cloning and expression analysis of some genes involved in the phosphoinositide and phospholipid signaling pathways from maize (Zea mays L.). Gene **426:** 47-56

Tasma IM, Brendel V, Whitham SA, Bhattacharyya MK (2008) Expression and evolution of the phosphoinositide-specific phospholipase C gene family in Arabidopsis thaliana. Plant Physiology and Biochemistry **46:** 627-637

Testerink C, Munnik T (2011) Molecular, cellular, and physiological responses to phosphatidic acid formation in plants. J Exp Bot **62:** 2349-2361

Theis J, Gupta TK, Klingler J, Wan W, Albert S, Keller S, Engel BD, Schroda M (2019) VIPP1 rods engulf membranes containing phosphatidylinositol phosphates. Sci Rep **9:** 8725

Theis J, Niemeyer J, Schmollinger S, Ries F, Rütgers M, Gupta TK, Sommer F, Muranaka LS, Venn B, Schulz-Raffelt M, Willmund F, Engel BD, Schroda M (2020) VIPP2 interacts with VIPP1 and HSP22E/F at chloroplast membranes and modulates a retrograde signal for HSP22E/F gene expression. Plant, Cell & Environment **43:** 1212-1229

Thole JM, Vermeer JE, Zhang Y, Gadella TW Jr, Nielsen E. (2008) Root hair defective4 encodes a phosphatidylinositol-4-phosphate phosphatase required for proper root hair development in Arabidopsis thaliana. Plant Cell;**20**:381-95.

Varnai P, Balla T (2006) Live cell imaging of phosphoinositide dynamics with fluorescent protein domains. Biochim Biophys Acta **1761:** 957-967

Walker EH, Pacold ME, Perisic O, Stephens L, Hawkins PT, Wymann MP, Williams RL (2000) Structural Determinants of Phosphoinositide 3-Kinase Inhibition by Wortmannin, LY294002, Quercetin, Myricetin, and Staurosporine. Molecular Cell **6:** 909-919

Wang CR, Yang AF, Yue GD, Gao Q, Yin HY, Zhang JR (2008) Enhanced expression of phospholipase C 1 (ZmPLC1) improves drought tolerance in transgenic maize. Planta **227:** 1127-1140

Weinberger A, Tsai FC, Koenderink GH, Schmidt TF, Itri R, Meier W, Schmatko T, Schröder A, Marques C. (2013) Gel-assisted formation of giant unilamellar vesicles. Biophys J.;**105**:154-64.

Xue HW, Chen X, Mei Y (2009) Function and regulation of phospholipid signalling in plants. Biochem J **421:** 145-156

Zarreen F, Kumar K, Chakraborty S (2023) Phosphoinositides in plant-pathogen interaction: trends and perspectives. Stress Biology **3:** 4

Zhai S, Gao Q, Liu X, Sui Z, Zhang J (2013) Overexpression of a Zea mays phospholipase C1 gene enhances drought tolerance in tobacco in part by maintaining stability in the membrane lipid composition. Plant Cell, Tissue and Organ Culture (PCTOC) **115:** 253-262

Zhang L, Kato Y, Otters S, Vothknecht UC, Sakamoto W. (2012) Essential role of VIPP1 in chloroplast envelope maintenance in Arabidopsis. Plant Cell. **24**:3695-707.

Zhang L, Kondo H, Kamikubo H, Kataoka M, Sakamoto W (2016) VIPP1 Has a Disordered C-Terminal Tail Necessary for Protecting Photosynthetic Membranes against Stress. Plant Physiol **171:** 1983-1995

821   **Zhu H, Wu Y, Zheng Y** (2022) Effects of heat shock on photosynthesis-related characteristics and
822   lipid profile of Cycas multipinnata and C. panzhihuaensis. BMC Plant Biology **22:** 442

823   **Zufferey M, Montandon C, Douet V, Demarsy E, Agne B, Baginsky S, Kessler F** (2017) The novel
824   chloroplast outer membrane kinase KOC1 is a required component of the plastid protein
825   import machinery. J Biol Chem **292:** 6952-6964

Fig. 1 Expression of YFP-PI3P, YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P$_2$ and YFP-PI(3,5)P$_2$ biosensors in chloroplasts of tobacco and identification of their sub-organelle distribution. A. The collection of biosensors is constructed to detect the different PPIs individually. Each specific biosensor consists of a chloroplast transit peptide and a fusion between lipid-binding domains that interacts specifically with each PPI and a yellow fluorescent protein (YFP), mCherry or mEOS4b respectively. B. scheme representing the distribution pattern of PPI biosensors in chloroplast with or without binding to specific PPIs. The upper panel represents a lack of binding of the PPI biosensor with diffuse localization of PPI biosensors in the chloroplast. Lower panel represents the binding of PPI biosensors to PPI-containing membranes with distinct localization of PPI biosensors. C. Confocal images of tobacco leaves transiently expressing PPI biosensors in the chloroplasts. Lacking the binding of the PI3P and PI5P biosensors results in diffuse localization of mEOS4b-PI3P and YFP-PI5P biosensors in the chloroplast. The binding of YFP-PI4P, YFP-PI(4,5)P2 and YFP-PI(3,5)P2 biosensors results in distinct punctate localization of PI biosensors indicating substrate recognition.

Fig. 2 Sub-organelle distribution pattern of mEOS4b-PI3P, mEOS4b-PI4P and YFP-PI(4,5)P$_2$ biosensors in the presence of the different PPI phosphatases in the chloroplasts. A. The illustration shows a simplified view of the phosphatase activity of PTEN, SAC7, and dOCRL in chloroplasts. B. Distribution pattern of mEOS4b-PI4P biosensors in the presence of phosphatase, SAC7, in the chloroplasts of transiently expressing tobacco. C. Distribution pattern YFP-PI(4,5)P$_2$ biosensors without and in the presence of presence of PI5P and PI3P phosphatases, dOCRL-mCherry and PTEN-mCherry in the chloroplasts of tobacco. D. Distribution pattern of PI3P biosensors in the presence of PI3P and PI4P phosphatases, PTEN-mCherry and dOCRL-mCherry in the chloroplasts of tobacco. Fluorescence signals were visualized in tobacco leaves co-expressing PPI biosensors and phosphatases (as indicated) by fluorescence confocal microscopy.

Fig. 3 Co-localization of VIPP1 with cTP-mEOS4b-PI3P (FYVE) biosensor and its differential sub-organelle localization in the presence of the PtdIns3P phosphatase, PTEN in the chloroplasts. A. Confocal images of tobacco leaves transiently expressing mEOS4b-PI3P (FYVE) biosensor in the presence and absence of VIPP1-CFP (middle and upper panel). Middle panels show colocalization of mEOS4b-PI3P biosensor with VIPP1-CFP in the chloroplasts. Lower panel shows different localization of mEOS4b-PI(4,5)P$_2$ biosensor and VIPP1-CFP in the chloroplasts. B. Sub-organelle distribution pattern of VIPP1-CFP in the presence of PI3P phosphatases, PTEN in the chloroplasts. C. Plant lipid based GUVs (magenta) with and without PI3P in the presence or absence of recombinant mEOS4b-VIPP1 (green).

Fig. 4 Sub-organelle distribution pattern of YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P$_2$ and YFP-PI(3,5)P$_2$ biosensors as well as cTP-GFP in the chloroplast of *Arabidopsis* seedlings under control and heat stress (HS) conditions. A and B. Confocal images of 10-days-old *Arabidopsis* seedlings stably expressing *cTP-YFP-PI4P*, *cTP-YFP-PI5P*, *cTP-YFP-PI(4,5)P$_2$* and c*TP-YFP-PI(3,5)P$_2$* biosensors. (A) control and (B) HS (at 40°C for 30 minutes). The diagrams at the right represent PIP flux upon control and HS. C. Immunoblot analysis of cTP-GFP, YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P and YFP-PI(3,5)P$_2$ proteins under control and HS conditions using GFP antibody.

Fig. 5 Perturbation of PIP abundance leads to drought sensitivity in *Arabidopsis* plants expressing PI4P phosphatase (SAC7), PI3P phosphatase (PTEN), and PI5P phosphatase (dOCRL). A. Sub-organelle distribution pattern of mEOS4b-PI3P, YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P2 and YFP-PI(3,5)P2 biosensors in the chloroplast of *Arabidopsis* seedlings under control and drought stress

conditions. Confocal images of 10-days-old *Arabidopsis* seedlings stably expressing *cTP-mEOS4b-PI3P*, *cTP-YFP-PI4P*, *cTP-YFP-PI5P*, *cTP-YFP-PI(4,5)P₂* and c*TP-YFP-PI(3,5)P₂* biosensors. For drought stress leaf samples from plants growing on petri dishes were detached and kept for 30 minutes at RT prior to imaging. B. Phenotypes of WT, *cTP-PTEN.OX* (left), *cTP-SAC7.OX* (middle) and *cTP-DOCRL.OX* (right) plants under control condition (well watered; upper panel) and after drought stress (middle panel). Seedlings were grown for 4 weeks in soil under a diurnal cycle of 16 h light ($120\,\mu E\,m^{-2}\,s^{-1}$) and 8 h dark at 22 °C. Drought stress was applied by withholding water for 10-14 days. The lower panel indicates *Fv/Fm* measurements of drought-stressed *cTP-PTEN.OX* (left) and *cTP-SAC7.OX* (right) plants. B. TEM images of *Arabidopsis* WT and *cTP-PTEN.OX* (upper images). TEM images of Arabidopsis WT and plants expressing *cTP-PTEN.OX* following drought treatments (lower images).

Parsed Citations

Akhter S, Uddin MN, Jeong IS, Kim DW, Liu XM, Bahk JD (2016) Role of Arabidopsis AtPI4Kγ3, a type II phosphoinositide 4-kinase, in abiotic stress responses and floral transition. Plant Biotechnol J 14: 215-230
Google Scholar: Author Only Author and Title

Annum N, Ahmed M, Imtiaz K, Mansoor S, Tester M, Saeed NA (2022) (32)P(i) Labeled Transgenic Wheat Shows the Accumulation of Phosphatidylinositol 4,5-bisphosphate and Phosphatidic Acid Under Heat and Osmotic Stress. Front Plant Sci 13: 881188
Google Scholar: Author Only Author and Title

Annum N, Ahmed M, Tester M, Mukhtar Z, Saeed NA (2023) Physiological responses induced by phospholipase C isoform 5 upon heat stress in Arabidopsis thaliana. Front Plant Sci 14: 1076331
Google Scholar: Author Only Title Only Author and Title

Balla T (2013) Phosphoinositides: tiny lipids with giant impact on cell regulation. Physiol Rev 93: 1019-1137
Google Scholar: Author Only Title Only Author and Title

Bankaitis VA, Aitken JR, Cleves AE, Dowhan W (1990) An essential role for a phospholipid transfer protein in yeast Golgi function. Nature. 347:561-2.
Google Scholar: Author Only Title Only Author and Title

Bassi R, dal Belin Peruffo A, Barbato R, Ghisi R (1985) Differences in chlorophyll-protein complexes and composition of polypeptides between thylakoids from bundle sheaths and mesophyll cells in maize. Eur J Biochem 146: 589-595
Google Scholar: Author Only Title Only Author and Title

Block MA (2018) Isolation of Inner and Outer Membranes of the Chloroplast Envelope from Spinach and Pea. Methods Mol Biol 1829:137-144
Google Scholar: Author Only Title Only Author and Title

Block MA, Dorne AJ, Joyard J, Douce R. (1983) Preparation and characterization of membrane fractions enriched in outer and inner envelope membranes from spinach chloroplasts. II. Biochemical characterization. J Biol Chem. 258:13281-6.
Google Scholar: Author Only Title Only Author and Title

Bovet L, Müller MO, Siegenthaler PA. (2001) Three distinct lipid kinase activities are present in spinach chloroplast envelope membranes: phosphatidylinositol phosphorylation is sensitive to wortmannin and not dependent on chloroplast ATP. Biochem Biophys Res Commun. 289: 269-75
Google Scholar: Author Only Title Only Author and Title

Carpaneto A, Boccaccio A, Lagostena L, Di Zanni E, Scholz-Starke J (2017) The signaling lipid phosphatidylinositol-3,5-bisphosphate targets plant CLC-a anion/H(+) exchange activity. EMBO Rep 18: 1100-1107
Google Scholar: Author Only Author and Title

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J 16: 735-743
Google Scholar: Author Only Title Only Author and Title

DeWald DB, Torabinejad J, Jones CA, Shope JC, Cangelosi AR, Thompson JE, Prestwich GD, Hama H (2001) Rapid Accumulation of Phosphatidylinositol 4,5-Bisphosphate and Inositol 1,4,5-Trisphosphate Correlates with Calcium Mobilization in Salt-Stressed Arabidopsis. Plant Physiology 126: 759-769
Google Scholar: Author Only Author and Title

Di Paolo G, De Camilli P (2006) Phosphoinositides in cell regulation and membrane dynamics. Nature 443: 651-657
Google Scholar: Author Only Title Only Author and Title

Dieck CB, Boss WF, Perera IY (2012) A role for phosphoinositides in regulating plant nuclear functions. Front Plant Sci 3: 50
Google Scholar: Author Only Title Only Author and Title

Doumane M, Lebecq A, Colin L, Fangain A, Stevens FD, Bareille J, Hamant O, Belkhadir Y, Munnik T, Jaillais Y, Caillaud MC (2021). Inducible depletion of PI(4,5)P2 by the synthetic iDePP system in Arabidopsis. Nat. Plants 7: 587–597.
Google Scholar: Author Only Author and Title

García-Cerdán JG, Schmid EM, Takeuchi T, McRae I, McDonald KL, Yordduangjun N, Hassan AM, Grob P, Xu CS, Hess HF, Fletcher DA, Nogales E, Niyogi KK (2020) Chloroplast Sec14-like 1 (CPSFL1) is essential for normal chloroplast development and affects carotenoid accumulation in *Chlamydomonas*. Proceedings of the National Academy of Sciences 117: 12452-12463
Google Scholar: Author Only Title Only Author and Title

Gerth K, Lin F, Daamen F, Menzel W, Heinrich F, Heilmann M (2017) Arabidopsis phosphatidylinositol 4-phosphate 5-kinase 2 contains a functional nuclear localization sequence and interacts with alpha-importins. Plant J 92: 862-878
Google Scholar: Author Only Title Only Author and Title

Gou JY, Li K, Wu K, Wang X, Lin H, Cantu D, Uauy C, Dobon-Alonso A, Midorikawa T, Inoue K, Sánchez J, Fu D, Blechl A, Wallington E, Fahima T, Meeta M, Epstein L, Dubcovsky J (2015) Wheat Stripe Rust Resistance Protein WKS1 Reduces the Ability of the Thylakoid-Associated Ascorbate Peroxidase to Detoxify Reactive Oxygen Species. Plant Cell 27: 1755-1770
  Google Scholar: Author Only Title Only Author and Title

Goulden BD, Pacheco J, Dull A, Zewe JP, Deiters A, Hammond GRV (2019). A high-avidity biosensor reveals plasma membrane PI(3,4)P2 is predominantly a class I PI3K signaling product. J Cell Biol. 218:1066-1079.
  Google Scholar: Author Only Author and Title

Gupta TK, Klumpe S, Gries K, Heinz S, Wietrzynski W, Ohnishi N, Niemeyer J, Spaniol B, Schaffer M, Rast A, Ostermeier M, Strauss M, Plitzko JM, Baumeister W, Rudack T, Sakamoto W, Nickelsen J, Schuller JM, Schroda M, Engel BD (2021). Structural basis for VIPP1 oligomerization and maintenance of thylakoid membrane integrity. Cell. 184:3643-3659
  Google Scholar: Author Only Title Only Author and Title

Heilmann I (2009) Using genetic tools to understand plant phosphoinositide signalling. Trends in Plant Science 14: 171-179
  Google Scholar: Author Only Title Only Author and Title

Hertle AP, García-Cerdán JG, Armbruster U, Shih R, Lee JJ, Wong W, Niyogi KK (2020) A Sec14 domain protein is required for photoautotrophic growth and chloroplast vesicle formation in *Arabidopsis thaliana*. Proceedings of the National Academy of Sciences 117: 9101-9111
  Google Scholar: Author Only Title Only Author and Title

Hirayama T, Ohto C, Mizoguchi T, Shinozaki K (1995) A gene encoding a phosphatidylinositol-specific phospholipase C is induced by dehydration and salt stress in Arabidopsis thaliana. Proc Natl Acad Sci U S A 92: 3903-3907
  Google Scholar: Author Only Title Only Author and Title

Horton P, Park KJ, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai K (2007) WoLF PSORT: protein localization predictor. Nucleic Acids Res 35: W585-587
  Google Scholar: Author Only Title Only Author and Title

Jensen JB, Falkenburger BH, Dickson EJ, de la Cruz L, Dai G, Myeong J, Jung S-R, Kruse M, Vivas O, Suh B-C, Hille B (2022) Biophysical physiology of phosphoinositide rapid dynamics and regulation in living cells. Journal of General Physiology 154
  Google Scholar: Author Only Title Only Author and Title

Kim E-H, Poudyal RS, Lee K-R, Yu H, Gi E, Kim HU (2022) Chloroplast-localized PITP7 is essential for plant growth and photosynthetic function in Arabidopsis. Physiologia Plantarum 174: e13760
  Google Scholar: Author Only Title Only Author and Title

LaBrant E, Barnes AC, Roston RL (2018) Lipid transport required to make lipids of photosynthetic membranes. Photosynthesis Research 138: 345-360
  Google Scholar: Author Only Title Only Author and Title

Leshem Y, Seri L, Levine A (2007) Induction of phosphatidylinositol 3-kinase-mediated endocytosis by salt stress leads to intracellular production of reactive oxygen species and salt tolerance. The Plant Journal 51: 185-197
  Google Scholar: Author Only Title Only Author and Title

Li L, Wang F, Yan P, Jing W, Zhang C, Kudla J, Zhang W (2017) A phosphoinositide-specific phospholipase C pathway elicits stress-induced Ca(2+) signals and confers salt tolerance to rice. New Phytol 214: 1172-1187
  Google Scholar: Author Only Title Only Author and Title

Li X (2011) Infiltration of Nicotiana benthamiana protocol for transient expression via Agrobacterium. Bio-protocol: e95-e95
  Google Scholar: Author Only Title Only Author and Title

Lu K-Y, Pasaje CFA, Srivastava T, Loiselle DR, Niles JC, Derbyshire E (2020) Phosphatidylinositol 3-phosphate and Hsp70 protect Plasmodium falciparum from heat-induced cell death. eLife 9: e56773
  Google Scholar: Author Only Title Only Author and Title

Maple J, Vojta L, Soll J, Møller SG. (2007) ARC3 is a stromal Z-ring accessory protein essential for plastid division. EMBO Rep 8(3):293-9
  Google Scholar: Author Only Title Only Author and Title

Matsuoka K, Bassham DC, Raikhel NV, Nakamura K (1995) Different sensitivity to wortmannin of two vacuolar sorting signals indicates the presence of distinct sorting machineries in tobacco cells. J Cell Biol 130: 1307-1318
  Google Scholar: Author Only Title Only Author and Title

Meijer HJ, Berrie CP, Iurisci C, Divecha N, Musgrave A, Munnik T (2001) Identification of a new polyphosphoinositide in plants, phosphatidylinositol 5-monophosphate (PtdIns5P), and its accumulation upon osmotic stress. Biochem J 360: 491-498
  Google Scholar: Author Only Title Only Author and Title

Meijer HJG, Divecha N, van den Ende H, Musgrave A, Munnik T (1999) Hyperosmotic stress induces rapid synthesis of

phosphatidyl-D-inositol 3,5-bisphosphate in plant cells. Planta 208: 294-298
  Google Scholar: Author Only Author and Title

Meijer HJG, Munnik T (2003) Phospholipid-Based Signaling in Plants. Annual Review of Plant Biology 54: 265-306
  Google Scholar: Author Only Title Only Author and Title

Mikami K, Katagiri T, Iuchi S, Yamaguchi-Shinozaki K, Shinozaki K (1998) A gene encoding phosphatidylinositol-4-phosphate 5-kinase is induced by water stress and abscisic acid in Arabidopsis thaliana. Plant J 15: 563-568
  Google Scholar: Author Only Title Only Author and Title

Mishkind M, Vermeer JEM, Darwish E, Munnik T (2009) Heat stress activates phospholipase D and triggers PIP2 accumulation at the plasma membrane and nucleus. The Plant Journal 60: 10-21
  Google Scholar: Author Only Title Only Author and Title

Morales-Johansson H, Jenoe P, Cooke FT, Hall MN (2004) Negative regulation of phosphatidylinositol 4,5-bisphosphate levels by the INP51-associated proteins TAX4 and IRS4. J Biol Chem 279: 39604-39610
  Google Scholar: Author Only Author and Title

Morales JA, Gonzalez-Kantun WA, Rodriguez-Zapata LC, Ramón-Ugalde J, Castano E (2019) The effect of plant stress on phosphoinositides. Cell Biochemistry and Function 37: 553-559
  Google Scholar: Author Only Title Only Author and Title

Müller MO, Meylan-Bettex M, Eckstein F, Martinoia E, Siegenthaler PA, Bovet L. (2000) Lipid phosphorylation in chloroplast envelopes. Evidence for galactolipid CTP-dependent kinase activities. J Biol Chem 275: 19475-81
  Google Scholar: Author Only Title Only Author and Title

Mueller-Roeber B, Pical C (2002) Inositol Phospholipid Metabolism in Arabidopsis. Characterized and Putative Isoforms of Inositol Phospholipid Kinase and Phosphoinositide-Specific Phospholipase C. Plant Physiology 130: 22-46
  Google Scholar: Author Only Title Only Author and Title

Munnik T, Meijer HJ (2001) Osmotic stress activates distinct lipid and MAPK signalling pathways in plants. FEBS Lett 498: 172-178
  Google Scholar: Author Only Title Only Author and Title

Munnik T, Wierzchowiecka M. (2013) Lipid-binding analysis using a fat blot assay. Methods Mol Biol 1009:253-9
  Google Scholar: Author Only Title Only Author and Title

Munnik T, Vermeer JE (2010) Osmotic stress-induced phosphoinositide and inositol phosphate signalling in plants. Plant Cell Environ 33: 655-669
  Google Scholar: Author Only Title Only Author and Title

Ndamukong I, Jones DR, Lapko H, Divecha N, Avramova Z (2010) Phosphatidylinositol 5-phosphate links dehydration stress to the activity of ARABIDOPSIS TRITHORAX-LIKE factor ATX1. PLoS One 5: e13396
  Google Scholar: Author Only Title Only Author and Title

Nordhues A, Schöttler MA, Unger AK, Geimer S, Schönfelder S, Schmollinger S, Rütgers M, Finazzi G, Soppa B, Sommer F, Mühlhaus T, Roach T, Krieger-Liszkay A, Lokstein H, Crespo JL, Schroda M (2012) Evidence for a role of VIPP1 in the structural organization of the photosynthetic apparatus in Chlamydomonas. Plant Cell 24: 637-659
  Google Scholar: Author Only Title Only Author and Title

Okazaki K, Kabeya Y, Suzuki K, Mori T, Ichikawa T, Matsui M, Nakanishi H, Miyagishima SY (2009) The PLASTID DIVISION1 and 2 components of the chloroplast division machinery determine the rate of chloroplast division in land plant cell differentiation. Plant Cell 21: 1769-1780
  Google Scholar: Author Only Title Only Author and Title

Okazaki K, Miyagishima SY, Wada H (2015) Phosphatidylinositol 4-phosphate negatively regulates chloroplast division in Arabidopsis. Plant Cell 27: 663-674
  Google Scholar: Author Only Title Only Author and Title

Perera IY, Hung C-Y, Moore CD, Stevenson-Paulik J, Boss WF (2008) Transgenic Arabidopsis Plants Expressing the Type 1 Inositol 5-Phosphatase Exhibit Increased Drought Tolerance and Altered Abscisic Acid Signaling. The Plant Cell 20: 2876-2893
  Google Scholar: Author Only Title Only Author and Title

Phillips SE, Vincent P, Rizzieri KE, Schaaf G, Bankaitis VA, Gaucher EA (2006) The diverse biological functions of phosphatidylinositol transfer proteins in eukaryotes. Crit Rev Biochem Mol Biol 41: 21-49
  Google Scholar: Author Only Title Only Author and Title

Pical C, Westergren T, Dove SK, Larsson C, Sommarin M (1999) Salinity and Hyperosmotic Stress Induce Rapid Increases in Phosphatidylinositol 4,5-Bisphosphate, Diacylglycerol Pyrophosphate, and Phosphatidylcholine in Arabidopsis thaliana Cells*. Journal of Biological Chemistry 274: 38232-38240
  Google Scholar: Author Only Author and Title

Pribat A, Sormani R, Rousseau-Gueutin M, Julkowska Magdalena M, Testerink C, Joubès J, Castroviejo M, Laguerre M, Meyer C, Germain V, Rothan C (2011) A novel class of PTEN protein in Arabidopsis displays unusual phosphoinositide phosphatase activity and efficiently binds phosphatidic acid. Biochemical Journal 441: 161-171
Google Scholar: Author Only Title Only Author and Title

Rajasekharan A, Gummadi SN. (2011) Flip-flop of phospholipids in proteoliposomes reconstituted from detergent extract of chloroplast membranes: kinetics and phospholipid specificity. PLoS One6(12):e28401
Google Scholar: Author Only Title Only Author and Title

Ravi A, Palamiuc L, Emerling BM (2021) Crucial Players for Inter-Organelle Communication: PI5P4Ks and Their Lipid Product PI-4,5-P(2) Come to the Surface. Front Cell Dev Biol 9: 791758
Google Scholar: Author Only Author and Title

Sedaghatmehr M, Mueller-Roeber B, Balazadeh S (2016) The plastid metalloprotease FtsH6 and small heat shock protein HSP21 jointly regulate thermomemory in Arabidopsis. Nature Communications 7: 12439
Google Scholar: Author Only Title Only Author and Title

Shimada H, Koizumi M, Kuroki K, Mochizuki M, Fujimoto H, Ohta H, Masuda T, Takamiya K. (2004) ARC3, a chloroplast division factor, is a chimera of prokaryotic FtsZ and part of eukaryotic phosphatidylinositol-4-phosphate 5-kinase. Plant Cell Physiol 45(8):960-7
Google Scholar: Author Only Author and Title

Siegenthaler PA, Müller MO, Bovet L. Evidence for lipid kinase activities in spinach chloroplast envelope membranes (1997). FEBS Lett. 416: 57-60
Google Scholar: Author Only Title Only Author and Title

Simon ML, Platre MP, Assil S, van Wijk R, Chen WY, Chory J, Dreux M, Munnik T, Jaillais Y (2014) A multi-colour/multi-affinity marker set to visualize phosphoinositide dynamics in Arabidopsis. Plant J 77: 322-337
Google Scholar: Author Only Title Only Author and Title

Singh N, Reyes-Ordoñez A, Compagnone MA, Moreno JF, Leslie BJ, Ha T, Chen J. (2021) Redefining the specificity of phosphoinositide-binding by human PH domain-containing proteins. Nat Commun:12:4339.
Google Scholar: Author Only Title Only Author and Title

Sui Z, Niu L, Yue G, Yang A, Zhang J (2008) Cloning and expression analysis of some genes involved in the phosphoinositide and phospholipid signaling pathways from maize (Zea mays L.). Gene 426: 47-56
Google Scholar: Author Only Title Only Author and Title

Tasma IM, Brendel V, Whitham SA, Bhattacharyya MK (2008) Expression and evolution of the phosphoinositide-specific phospholipase C gene family in Arabidopsis thaliana. Plant Physiology and Biochemistry 46: 627-637
Google Scholar: Author Only Title Only Author and Title

Testerink C, Munnik T (2011) Molecular, cellular, and physiological responses to phosphatidic acid formation in plants. J Exp Bot 62: 2349-2361
Google Scholar: Author Only Title Only Author and Title

Theis J, Gupta TK, Klingler J, Wan W, Albert S, Keller S, Engel BD, Schroda M (2019) VIPP1 rods engulf membranes containing phosphatidylinositol phosphates. Sci Rep 9: 8725
Google Scholar: Author Only Title Only Author and Title

Theis J, Niemeyer J, Schmollinger S, Ries F, Rütgers M, Gupta TK, Sommer F, Muranaka LS, Venn B, Schulz-Raffelt M, Willmund F, Engel BD, Schroda M (2020) VIPP2 interacts with VIPP1 and HSP22E/F at chloroplast membranes and modulates a retrograde signal for HSP22E/F gene expression. Plant, Cell & Environment 43: 1212-1229
Google Scholar: Author Only Title Only Author and Title

Thole JM, Vermeer JE, Zhang Y, Gadella TW Jr, Nielsen E. (2008) Root hair defective4 encodes a phosphatidylinositol-4-phosphate phosphatase required for proper root hair development in Arabidopsis thaliana. Plant Cell;20:381-95.
Google Scholar: Author Only Title Only Author and Title

Varnai P, Balla T (2006) Live cell imaging of phosphoinositide dynamics with fluorescent protein domains. Biochim Biophys Acta 1761: 957-967
Google Scholar: Author Only Title Only Author and Title

Walker EH, Pacold ME, Perisic O, Stephens L, Hawkins PT, Wymann MP, Williams RL (2000) Structural Determinants of Phosphoinositide 3-Kinase Inhibition by Wortmannin, LY294002, Quercetin, Myricetin, and Staurosporine. Molecular Cell 6: 909-919
Google Scholar: Author Only Author and Title

Wang CR, Yang AF, Yue GD, Gao Q, Yin HY, Zhang JR (2008) Enhanced expression of phospholipase C 1 (ZmPLC1) improves drought tolerance in transgenic maize. Planta 227: 1127-1140

Google Scholar: Author Only Title Only Author and Title

Weinberger A, Tsai FC, Koenderink GH, Schmidt TF, Itri R, Meier W, Schmatko T, Schröder A, Marques C. (2013) Gel-assisted formation of giant unilamellar vesicles. Biophys J.;105:154-64.
  Google Scholar: Author Only Title Only Author and Title

Xue HW, Chen X, Mei Y (2009) Function and regulation of phospholipid signalling in plants. Biochem J 421: 145-156
  Google Scholar: Author Only Title Only Author and Title

Zarreen F, Kumar K, Chakraborty S (2023) Phosphoinositides in plant-pathogen interaction: trends and perspectives. Stress Biology 3: 4
  Google Scholar: Author Only Title Only Author and Title

Zhai S, Gao Q, Liu X, Sui Z, Zhang J (2013) Overexpression of a Zea mays phospholipase C1 gene enhances drought tolerance in tobacco in part by maintaining stability in the membrane lipid composition. Plant Cell, Tissue and Organ Culture (PCTOC) 115: 253-262
  Google Scholar: Author Only Title Only Author and Title

Zhang L, Kato Y, Otters S, Vothknecht UC, Sakamoto W. (2012) Essential role of VIPP1 in chloroplast envelope maintenance in Arabidopsis. Plant Cell. 24:3695-707.
  Google Scholar: Author Only Title Only Author and Title

Zhang L, Kondo H, Kamikubo H, Kataoka M, Sakamoto W (2016) VIPP1 Has a Disordered C-Terminal Tail Necessary for Protecting Photosynthetic Membranes against Stress. Plant Physiol 171: 1983-1995
  Google Scholar: Author Only Title Only Author and Title

Zhu H, Wu Y, Zheng Y (2022) Effects of heat shock on photosynthesis-related characteristics and lipid profile of Cycas multipinnata and C. panzhihuaensis. BMC Plant Biology 22: 442
  Google Scholar: Author Only Title Only Author and Title

Zufferey M, Montandon C, Douet V, Demarsy E, Agne B, Baginsky S, Kessler F (2017) The novel chloroplast outer membrane kinase KOC1 is a required component of the plastid protein import machinery. J Biol Chem 292: 6952-6964
  Google Scholar: Author Only Title Only Author and Title