New evidence for the presence and function of phosphoinositides (PPIs) in the chloroplast

Mastoureh Sedaghatmehr[1,2], Frieda Rößler[1,3], Alexander P. Hertle*[1,2,4]

[1]Max-Planck Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam,
Germany
[2]Heinrich-Heine-Universität Düsseldorf, 40225 Düsseldorf, Germany
[3]Leibnitz Institute of Plant Biochemistry, Weinberg 3, 06120 Halle, Germany
[4]Cluster of Excellence on Plant Sciences, Heinrich Heine University, Düsseldorf, Germany

***Correspondence:** Alexander P. Hertle, email: Alexander.Hertle@hhu.de

Abstract

Essential proteins such as the chloroplast-localized Sec14-like protein (CPSFL1) and the
vesicle-inducing protein in plastids (VIPP1) have a high affinity for binding phosphoinositides
(PPIs) in vitro. PPIs are a specific class of phospholipids characterized by a phosphorylated
inositol head group, and while they make up a small fraction of total phospholipids, they are
crucial for various regulatory functions. However, the precise subcellular localization of most
PPI species in plants remains unclear due to their rapid turnover and low abundance Currently
there is no documented evidence for the presence and function of phosphoinositides (PPIs) in
chloroplasts. In our study, we developed genetically encoded biosensors targeted to plastids,
allowing for the detection of several PPI isoforms, including PI3P, PI4P, PI5P, PI(4,5)P_2, and
PI(3,5)P_2 within chloroplasts. We demonstrated the specificity of these biosensors through
immunological methods and observed changes in their distribution patterns when co-expressed
with PPI-modifying enzymes (cTP-SAC7, cTP-PTEN, and cTP-dOCRL). Our findings also
revealed the association and potential interaction between PI3P and VIPP1. Importantly, we
found that elevated PPI levels during stress conditions led to altered biosensor localization, and
plants expressing PPI modifiers showed increased sensitivity to drought stress, highlighting the
role of PPIs in plant stress responses.

Keywords: Phosphatidylinositol phosphates, chloroplasts, signaling, PPI biosensors, abiotic
stress, lipids

Introduction

Phosphatidylinositols (PtdIns or PIs) are low-abundance lipids with significant regulatory functions, governing a wide range of physiological processes in eukaryotes. PtdIns are produced in the endoplasmic reticulum (ER) and subsequently transported to various subcellular sites through a combination of lipid transfer proteins and vesicle trafficking mechanisms (Phillips et al., 2006; Heilmann, 2009). Phosphorylated derivatives of PtdIns are generated by a group of enzymes, including phosphatidylinositol kinases (PIKs) and phosphatidylinositol phosphatases (Mueller-Roeber and Pical, 2002). PIK enzymes phosphorylate three out of the five available hydroxyl groups on the inositol ring, specifically at positions 3′, 4′, and 5′. This process generates seven phosphatidylinositol phosphates (PtdInsPs or PPIs) consisting of three monophosphates (PI3P, PI4P, and PI5P), three bisphosphates PI(3,4)P$_2$, PI(4,5)P$_2$, and PI(3,5)P$_2$ and one triphosphate PI(3,4,5)P$_3$. Among them, the presence of PI(3,4)P$_2$ and PI(3,4,5)P$_3$ has not yet been confirmed in plants (Mueller-Roeber and Pical, 2002; Zarreen et al., 2023). To date, no homologs of the heterodimeric inositol lipid 3-kinases have been identified in plant cells to phosphorylate PI4P to PI(3,4)P$_2$ and PI(4,5)P$_2$ to PI(3,4,5)P$_3$. PI5P is found in plants, yet an enzyme responsible for its production remains unidentified (Mueller-Roeber and Pical, 2002; Heilmann, 2009; Zarreen et al., 2023).

PPIs are being recognized as new secondary messengers within plant cells. Besides their crucial role under favorable conditions, they are quickly generated when plants are exposed to a wide range of environmental stresses, indicating their importance under stress conditions (Munnik and Meijer, 2001; Meijer and Munnik, 2003; Wang et al., 2008; Xue et al., 2009; Munnik and Vermeer, 2010; Pribat et al., 2011; Zhai et al., 2013; Zarreen et al., 2023). PI3P, PI5P, PI(3,5)P$_2$, and PI(4,5)P$_2$ exhibited increased levels in response to salt and osmotic stress, likely due to the induction of relevant PI kinases or the action of lipid phosphatases, resulting from the degradation products of other PPIs. (Meijer et al., 1999; Pical et al., 1999; DeWald et al., 2001; Meijer et al., 2001; Meijer and Munnik, 2003, König et al., 2008). Individual studies have reported alteration in PPI kinase and phosphatase activity under stress conditions. Salt and osmotic stress have been shown to boost both PI 3-kinase and PI3P 5-kinase activities in plants. (Meijer et al., 1999; Meijer et al., 2001; Munnik and Meijer, 2001; Leshem et al., 2007, König et al., 2008). PTEN (phosphatase and tensin homolog deleted on chromosome ten) proteins in vascular plants play a crucial role in regulating 3-phosphorylated PI signaling events. The transcript levels of *AtPTEN2* exhibited changes in response to various stresses, including conditions of salt and osmotic stress (Pribat et al., 2011). Similarly, environmental

stresses such as cold, salt, and drought stimulate the expression and activity of

phosphoinositide-specific phospholipase C (PLC) (Hirayama et al., 1995; Munnik and Meijer,

2001). PLC induces the hydrolysis of PI(4,5)P$_2$ to generate the two signaling molecules,

inositol-$_{1,4,5}$-trisphosphate (InsP$_3$) and diacylglycerol (DAG) (Hirayama et al., 1995; Testerink

and Munnik, 2011; Zarreen et al., 2023). PLC has the ability to enhance salt tolerance in rice

as well as drought tolerance in maize and tobacco (Wang et al., 2008; Zhai et al., 2013; Li et

al., 2017). The expression of the well-characterized PI4K kinase, *AtPI4Kγ3*, is induced upon

exposure to salt stress. *Arabidopsis* plants overexpressing PI4Kγ3 exhibit increased stress

tolerance and inducible expression of stress-responsive genes (Akhter et al., 2016). All these

investigations suggest that alterations in PPI concentration are tightly regulated and play a

critical role in facilitating precise PPI signaling.

The unique characteristics of PPIs, including their low abundance, rapid turnover, and

diversity, make them dynamic signaling molecules. PPIs have the ability to bind to specific

proteins, which can result in changes in the proteins' localization, conformation, and activity.

Additionally, PPIs can influence membrane curvature. These capabilities are crucial aspects of

PPI function in facilitating downstream signaling in all eukaryotes (Meijer and Munnik, 2003;

Xue et al., 2009; Munnik and Vermeer, 2010).

For instance, PPI affinity chromatography and mass spectrometry analysis confirmed that

PI(3,5)P$_2$ and PI3P bind to ROF1 (ROTAMASE FKBP1) in Arabidopsis (Karali et al., 2012).

ROF1 is associated with responses to heat, salt, and osmotic stress (Meiri and Breiman, 2009;

Karali et al., 2012). The authors suggested that the interaction of ROF1 with PPI causes a

change in its localization under salt stress (Karali et al., 2012). Similarly, PI(4,5)P$_2$ plays a role

in regulating ion channels and proteins associated with cytoskeletal modification and

membrane trafficking (McLaughlin et al., 2002; Suh et al., 2006; Munnik and Vermeer, 2010).

While the functions of PtdIns in membrane trafficking, endomembrane organization, and

cytoskeletal rearrangement are well-established, their pivotal roles in signaling and their

subcellular localization are only emerging in plants. Due to their rapid turnover and limited

abundance, the precise subcellular localization of most PPI species in plants remains

incomplete. The existence and function of PPI in the nucleus of animals, yeast, and plant cells

has been documented (Horton et al., 2007; Dieck et al., 2012; Gerth et al., 2017). It is widely

recognized that nuclear PtdIns play a significant role in regulating various processes, including

DNA replication, chromatin remodeling, transcription, stress responses, and hormone signaling

in plant, animal, and yeast cells (Dieck et al., 2012; Zarreen et al., 2023). The unique role of

PPIs in the nucleus, distinct from their function in the plasma membrane, highlights the
existence of distinct PPI signaling pathways within various cellular organelles.

A recent biochemical study has highlighted the critical role of PI4P in the functioning of
another cellular organelle, chloroplasts. They revealed that inhibiting PI4Ks through chemical
inhibitor treatment in *Arabidopsis* seedlings accelerates chloroplast division. It has also been
reported that PI4P specifically binds to PLASTID DIVISION1 (PDV1) and PDV2, exerting a
negative regulatory effect on chloroplast division. Of note, PDV1 and PDV2 proteins are
located at the outer envelope membrane of the chloroplast, and the PI4P lipid-signaling
pathway, involved in chloroplast division via PDVs takes place at the outer envelope
membrane. Therefore, the presence and function of PPIs inside the chloroplasts remain
unexplored thus far (Okazaki et al., 2015).

Lipid transport across the envelope membranes is highly controlled and restricted (Karki et al.,
2019). Furthermore, the enzymes responsible for PPI biosynthesis in plastids have so far not
been identified. This makes the presence of these "eukaryotic" lipids in the inner chloroplast
membranes (e.g. inner envelope and thylakoids) rather unlikely. Nevertheless, experiments
using radioactive labeling with ATP, as well as mass spectrometric analysis of plastidial
endomembranes, have detected low amounts of individual PPI species (König et al., 2008;
Siegenthaler et al., 1997). Contamination through membrane contacts with the ER or
thylakoid/envelope membranes would be one explanation. Yet, the presence and function of
PPIs in plastids cannot be denied either. Additionally, studies on membrane remodeling protein
VIPP1 and chloroplast vesicle traffic regulator CPSFL1 support the latter by to PPIs with high
specificity *in vitro* (Theis et al.2019, Hertle et al., 2020)

In this study, our results present evidence for the presence of PPIs within the chloroplasts of
tobacco, as indicated by various PPI biosensors and immunodetection. We observed rapid
relocation of PPI biosensors within the chloroplasts of *Arabidopsis* under abiotic stress
conditions. Additionally, plants expressing chloroplast-targeted PPI modifiers exhibited
increased sensitivity to drought and heat stress. Overall, our investigation sheds light on the
existence of PPIs in the chloroplast and their potential involvement in the PPI signaling
pathway during stress conditions.

Results

Generation and sub-organelle localization of a set of PPI biosensors in chloroplasts

Monitoring PPI poses a consistent challenge due to the rapid turnover, tight regulation, and low
quantities of PPIs present within all living cells (Meijer and Munnik, 2003; Jensen et al., 2022).

Indirect evidence indicating the presence of PPIs in the chloroplast is mainly limited to *in vitro*

studies (Gou et al., 2015; Okazaki et al., 2015; Theis et al., 2019; Hertle et al., 2020).

Nevertheless, a current research gap exists concerning the *in vivo* study of PPIs within

chloroplasts.

Genetically encoded biosensors have been widely employed to study the distribution of PPIs

in functional living cells (Varnai and Balla, 2006; Balla, 2013; Simon et al., 2014). Biosensors

consist of lipid-binding domains that interact specifically with known PPIs and can be easily

tracked by fusion with a fluorescent protein. To investigate the presence of various PPIs in the

lipid composition of chloroplasts, we generated a collection of biosensors that contain a

chloroplast transit peptide (cTP), yellow fluorescence protein or mCherry , and a lipid-binding

domain. To individually and specifically detect PI3P, PI4P, PI5P, PI(4,5)P_2, and PI(3,5)P_2, the

coding sequences of the following domains were used. The pleckstrin homology (PH) domain

of human ARHGEF9 was used for PI5P binding. The PH domain of human Rho guanine

nucleotide exchange factor 7 (ARHGEF7) was used for PI(3,5)P_2 detection (Singh et al., 2021).

The PH domain of rat four-phosphate adaptor protein 1 (FAPP1) was used for PI4P detection.

The PH domain of mouse 1-phosphatidylinositol-4,5-bisphosphate diesterase delta (Plcd1) was

used for PI(4,5)P_2 detection. The FYVE domain of human hepatocyte growth factor-regulated

tyrosine kinase substrate (HGS11) was used for PI3P detection (Simon et al., 2014). These

domains were codon-optimized for expression in *Arabidopsis* and fused to the C-terminal

region of a chloroplast-targeted fluorescent reporter (Fig. 1a). Whereas some PH domains bind

to more than one PPIs isoform, the employed domains for PI3P, PI4P and PI(4,5)P_2 detection

have been shown in numerous *in vivo* studies in plants to be specific for their substrate (Varnai

and Balla, 2006; Balla, 2013; Simon et al., 2014). Moreover, specific recognition of PI5P and

PI(3,5)P_2 by the lipid binding domains of ARHGEF7 and 9 has been shown *in vitro* (Singh et

al.2021).

To visualize their distribution pattern in plastids, we performed a transient expression of

each of the designed biosensor constructs in tobacco leaves using infiltration with transgenic

Agrobacteria to initiate transient expression (shown in Fig. **1a**). YFP fluorescence was

observed 48 hours after Agrobacterium infiltration in the chloroplast of all transformed cells

due to the presence of chloroplast transit peptide. We further analyzed the sub-organelle

distribution of PPI biosensors in the chloroplasts. We observed a diffuse localization of YFP-

PI3P and YFP-PI5P biosensors within the chloroplast. In contrast, binding of YFP-PI4P, YFP-

PI(4,5)P_2, and YFP-PI(3,5)P_2 biosensors to PPI-containing membrane leads to a different

localization reflected in distinct punctate structures within or surrounding the thylakoids

represented by chlorophyll autofluorescence of chloroplasts (Fig. **1b, c**). Counting puncta in

several chloroplasts gave a constant and characteristic number of puncta for respective sensors.

Furthermore, $PI(3,5)P_2$ and $PI(4,5)P_2$ biosensors, with a lower number of puncta colocalized

with the chlorophyll fluorescence, whereas PI4P sensors with a much higher number of smaller

puncta localized differently in a peripheral region of the chloroplast surrounding the thylakoids

as indicated by chlorophyll fluorescence (Supporting Information Fig. S1 a, b).

The presence of PI4P in the outer envelope of the chloroplast has been previously demonstrated

(Okazaki et al., 2015). To confirm that the YFP-PI4P biosensors detect their substrate within

the chloroplast and not at the outer envelope of the chloroplast, we treated isolated chloroplasts

from plants expressing YFP-PI4P biosensor with the protease thermolysin. Thermolysin

effectively breaks down proteins located on the surface of chloroplasts, but it does not penetrate

the intermembrane space (Zufferey et al., 2017). Chloroplasts subjected to thermolysin

treatment were subsequently analyzed by SDS-PAGE and subjected to immunoblot analysis.

TOC159, a resident of the outer chloroplast envelope membrane and known to be thermolysin-

sensitive was notably reduced following thermolysin treatment. In contrast, the YFP-PI4P

biosensor protein remained largely unaffected by the treatment (Supporting Information Fig.

S1c). This observation indicates that the YFP-PI4P biosensor is mainly associated with

membranes within the chloroplasts and is not readily accessible for thermolysin-mediated

degradation at the outer surface membrane. To further support the presence of PI4P within the

inner membranes of chloroplasts as well as its specificity, we performed fat blot assays using

lipid standards together with membrane extracts and detected PI4P immunologically using

PI4P specific antibodies (Weber et al., 2006) (Supporting Information Fig 1d). The same

antibodies also recognized lipid extracts from chloroplasts and thylakoids isolated from

Arabidopsis (Supporting Information Fig S1d).

Chloroplast localized-PPI biosensors change their distribution upon PPI phosphatase

expression

To further support the specificity of our PPI biosensors, we verified the sub-organelle

distribution pattern of PPI biosensors in the chloroplasts of tobacco plants in the presence of

PPI modifiers. For this purpose, we generated three distinct PPI phosphatases that feature a

chloroplast transit peptide. These phosphatases include SUPPRESSOR OF ACTIN 7 (SAC7)

phosphatase and tensin homolog deleted on chromosome ten (PTEN), and the phosphatase

domain of *Drosophila melanogaster* Oculocerebrorenal Syndrome of Lowe protein (dOCRL)

(Goulden et al., 2019; Thole et al., 2008; Doumane et al., 2021) . SAC7, PTEN, and dOCRL

can reversibly dephosphorylate PI4P, PI3P, and PI5P respectively, by their action as lipid

phosphatases (Fig. **2a**). Co-expression of the PI(4,5)P$_2$ biosensor with *cTP-SAC7* and *cTP-*

dOCRL, modifiers specific to PI4P and PI5P, respectively, resulted in a significant and

substantial alteration in the distribution pattern of the PI(4,5)P$_2$ biosensor. The distinctive

punctate structures of the PI(4,5)P$_2$ biosensor were notably reduced and transformed into

diffuse fluorophore signals in the presence of the SAC7 and dOCRL phosphatases (Fig. **2b, c**).

In contrast, the punctate-like structures of the PI(4,5)P$_2$ biosensor remained unchanged when

co-expressed with the PI3P modifier, *cTP-PTEN* (Fig. **2c**). The punctate signals of the PI(4,5)P$_2$

biosensor within the chloroplasts was also predominantly converted into diffuse fluorophore

signals in the presence of the PI4P modifier SAC7. Since PI3P biosensors were equally

distributed when targeted to plastids, co-expression of PTEN or SAC7 did not show any

alternation (Fig. **2d**). Observed changes were quantified and analyzed for statistical relevance

(Supporting Information Fig. S2a). A significant reduction of fluorescent puncta was observed

for PI4,5P$_2$ sensors upon dOCRL expression. Moreover, co-expression of SAC7 and PI4P

sensors resulted in a significant reduction of the fluorescent puncta but an increase in a diffuse

fluorescent signal.

We also examined the distribution of the same PI(4,5)P$_2$ biosensor tagged with different

fluorophores following co-expression. As expected, biosensors with identical substrate, YFP-

PI(4,5)P$_2$ and mCherry-PI(4,5)P$_2$, co-localized with the same fluorescence pattern in the

chloroplasts (Supporting Information Fig. S2b). Next, we co-expressed biosensors sensitive

for PI(3,5)P$_2$ tagged with YFP and a biosensor specific for PI4P and tagged with mCherry

transiently in tobacco leaves. The sensors differed in their sub-organelle localization and in

respect to the chlorophyll autofluorescence. In summary, these observations collectively

demonstrate the specificity of our PPI biosensors in selectively detecting individual PPIs within

the chloroplasts (Supporting Information Fig. S2c, d).

VIPP1 may colocalize with PI3P rich membranes

To date, our understanding of the interaction between chloroplast proteins and PPIs has

primarily relied on *in vitro* studies employing protein-lipid blot overlay assays (Gou et al.,

2015; Okazaki et al., 2015; Theis et al., 2019; Hertle et al., 2020). We further assayed the

colocalization of the chloroplast protein, VESICLE-INDUCING PROTEIN IN PLASTIDS 1

(VIPP1) with individual PPI biosensors. *In vitro* studies have already demonstrated the robust

binding of *Chlamydomonas reinhardtii* VIPP1 to liposomal membranes containing PI4P,

PI(4,5)P$_2$, and PI(3,4,5)P$_3$ (Theis et al., 2019).

To assess the *in vivo* association of VIPP1 with PPIs, we conducted colocalization between
VIPP1-CFP and PPI biosensors within the chloroplasts. Interestingly, the simultaneous
expression of *VIPP1-CFP* and *cTP-mEOS4b-PI3P* led to a significant alteration in the
distribution pattern of mEOS4b-PI3P, resulting in the formation of distinct dot-like structures
(Figure 3A). Significantly, these dot-like structures of mEOS4b-PI3P exhibited co-localization
with VIPP1-CFP (Fig. **3a**). Interestingly, the overrepresentation of VIPP1 did not alter the sub-
organelle distribution pattern of the PI(4,5)P$_2$ biosensor (Fig. **3a**)   .
Similarly, co-expression of the *VIPP1-CFP* with *cTP-mCherry-PTEN*, modifier specific to
PI3P, resulted in a substantial alteration in the typical distribution pattern of VIPP1 within
chloroplasts (Fig. **3b**). Subsequently, we examined whether the stability of VIPP1 and the
configuration of its homooligomeric complexes were affected in the presence of PTEN. The
amounts of VIPP1 remained unaltered in plants expressing *cTP-PTEN* (Supplementing
Information Fig. S3a). To verify the complex state of VIPP1 in *cTP-PTEN*-expressing plants,
thylakoid proteins from both WT and *cTP-PTEN* plants were subjected to treatment with Bis
(Sulfosuccinimidyl) suberate (BS^3). BS^3 serves as a membrane-impermeable crosslinking
agent to stabilize the complex state. Nevertheless, no significant differences   were observed
in terms of the multiple complex states of VIPP1 in *cTP-PTEN*-expressing plants (Supporting
Information Fig. S3b). Our observation suggests that overexpression of VIPP1 may concentrate
PI3P lipids in a specific location within the chloroplast, making them detectable by the PI3P
biosensor, while in the absence of VIPP1 overexpression, the condensation of PI3P appears to
be below the detection threshold of the PI3P biosensor (Fig. **1c**). Thus, we wanted to proof the
sensitivity and specificity of the PI3P sensor by an independent approach. For this we purified
chloroplasts and thylakoids purified form *Arabidopsis* and spinach respectively. Following
lipid extraction, we performed fat blot assays together with a dilution series of PI3P lipid
standards and probed them with PI3P specific FYVE-mEOS4b sensor proteins. Recombinant
FYVE sensors recognized PI3P lipid standards as well as chloroplast and thylakoid lipid
extracts from both species. Still, with a low concentration (<6.3 pmol) and obscured by a bright
spot in the center of the lipid dot resembling the distribution of coextracted chlorophyll
(Supporting Information Fig. S3c). Thus, we fractionated spinach chloroplasts and probed
chlorophyll free inner envelope membranes, thylakoids and an array of chloroplast lipids to
support specific recognition of PI3P sensor proteins. PI3P and inner chloroplast envelope
membranes were recognized with high specificity (Supporting Information Fig. S3d).
The specificity of FYVE-mEOS4b sensors in the presence of thylakoid lipids and PI3P was
further investigated by using recombinant sensor proteins and using giant unilammelar vesicles

(GUVs) composed of MGDG, DGDG, SQDG, PG and PI in a stöchiometry of 52:26:6.5:9.5:1

(v/v) (Hölzl & Dörmann, 2019; LaBrant et al., 2018). Only PI3P containing GUVs were

recognized by PI3P biosensor proteins (Supporting Information Fig. S3e). Next, we tested the

binding of VIPP1 to plant lipid-based giant unilamellar vesicles (GUVs). While low binding

of VIPP1 was observed in GUVs composed of MGDG, DGDG, PG, and SQDG, the inclusion

of 0.1 mol% PI3P into the GUV lipid mixtures led to a pronounced recruitment of recombinant

VIPP1 to the GUV surfaces in punctate pattern (Fig. **3c**). This punctate pattern of VIPP1could

not be observed for FYVE sensor proteins which suggests a non-homogeneous distribution or

local concentration of phosphatidylinositols (PPIs) induced by VIPP1 (Fig. **3a, c**).

Taken together, these results suggest the association and potential function of VIPP1 with PI3P

in vivo.

The localization of PPI sensors changes upon heat in chloroplasts of *Arabidopsis*

To investigate the presence and function of various PPIs in the chloroplasts, we generated

transgenic *Arabidopsis* lines stably expressing our chloroplast localized biosensors (*cTP-YFP-*

PI4P, *cTP-YFP-PI5P*, *cTP-YFP-PI(4,5)P$_2$*, and *cTP-YFP-PI(3,5)P$_2$*. Of note, none of the

transgenic lines showed any visible developmental phenotypes. Next, we assessed the sub-

chloroplasts distribution patterns of YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P$_2$, YFP-PI(3,5)P$_2$

biosensors, and cTP-GFP within the chloroplasts of 10-day-old *Arabidopsis* seedlings under

standard control conditions using confocal microscopy. We noted that all the PPI sensors

displayed a diffuse localization within the chloroplast (Fig. **4a**). This observation suggests that

the biosensors may not have encountered their intended targets for binding, or alternatively,

the concentration of PPIs in the chloroplast of *Arabidopsis* seedlings might be below the

detection limits of our designed PPI biosensors.

Furthermore, we decided to elevate the PPI flux within the cell, potentially including the

chloroplast, and subsequently assess the sub-chloroplast distribution patterns of the PPI

biosensors. Studies have demonstrated that the metabolism of PI and PPIs is significantly

activated in response to heat stress in various organisms (Morales-Johansson et al., 2004; Lu

et al., 2020) and diverse plant species (Mishkind et al., 2009; Annum et al., 2022; Zhu et al.,

2022; Annum et al., 2023). To elevate the PPI flux, we subjected 10-day-old Arabidopsis

seedlings to a 40°C temperature treatment for 30 minutes, after which we conducted confocal

microscopy analysis. Surprisingly, we observed that all PPI biosensors re-localized within the

chloroplasts in response to short heat stress treatment. As shown in Fig. **4b**, there was no change

in the sub-chloroplast localization of GFP in plants expressing GFP targeted into chloroplasts.

In contrast, we observed a clear translocation of sensors to PPI-containing membranes resulting

in punctate-like structures in the plant expressing PPI biosensors (Fig. **4b**). Quantification of

fluorescent loci in plastids containing PI4P, PI5P and PI(3,5)P$_2$ sensor proteins showed similar

numbers per chloroplast. Consistent with transient expression PI(4,5)P$_2$ sensor containing

plastids showed significantly fewer puncta as compared to other sensor lines (Supporting

Information Fig. S4a, b). Although, expression of PPI biosensors is under the control of the

35S promoter, we additionally confirmed, through western blot analysis, that the heat stress

treatment did not result in any noticeable changes in their abundance when compared to the

control condition (Fig. **4c**). Moreover, we noticed that the distribution patterns of PI(4,5)P$_2$-

YFP biosensors remained unchanged in plastids of heat-stressed root cells, in comparison to

untreated root cells (Supporting Information Fig. S4c). These results indicates that the

relocation PI(4,5)P$_2$ sensors observed in leaves is dependent on chloroplast development or

thylakoid differentiation.

Potential function of PPIs as a lipid signaling molecule in chloroplasts under abiotic

stresses

Multiple studies have demonstrated stress-induced alterations in PI and PPI signaling, implying

the pivotal roles of PPI signaling in plant adaptation to stress conditions (Hirayama et al., 1995;

Meijer et al., 1999; Pical et al., 1999; DeWald et al., 2001; Meijer et al., 2001; Heilmann, 2009;

Munnik and Vermeer, 2010; Zhu et al., 2022; Annum et al., 2023). We observed that almost

all PPI biosensors re-localized within the chloroplasts in response to drought stress treatment,

(Fig. **5a** and Supporting Information Fig. S5a) most probably due to increased PPI flux within

the chloroplasts following drought stress. Quantification of fluorescent puncta per chloroplast

showed a significant difference between PI4P and PI(4,5)P$_2$ and PI(3,5)P$_2$ lines. In contrast, no

punctate structures were observed for chloroplasts of PI3P and PI5P lines (Supporting

Information Fig. S5a). Next, to further explore the possible involvement of various PPIs within

the chloroplasts in response to abiotic stress, we generated transgenic Arabidopsis lines that

constitutively express PPI modifiers (phosphatases) containing a chloroplast transit peptide,

namely *cTP-SAC7.OX, cTP-PTEN.OX*, and *cTP-dOCRL.OX*. As previously mentioned, SAC7,

PTEN, and dOCRL are capable of reversibly dephosphorylating PI4P, PI3P, and PI5P,

respectively (Fig. **2a**). Of note, none of the transgenic lines expressing *cTP-SAC7, cTP-PTEN,*

and *cTP-dOCRL* showed any visible developmental phenotypes, nor did they exhibit any

abnormalities (Fig. **5b**). This observation aligns with the absence of binding of PPI biosensors

to any PPIs in the chloroplast of plants growing under optimal conditions (Fig. **4a**).

Subsequently, plant overexpressing plastidial phosphatase were subjected to drought stress by

withholding water. As shown in Fig. **5b**, plants overexpressing *cTP-SAC7*, *cTP-PTEN*, and

cTP-dOCRL began to exhibit leaf wilting symptoms approximately 8 days into the drought

period. These symptoms became progressively more pronounced in the subsequent days of

drought compared to the control plants. Furthermore, the measurements of the maximum

quantum efficiency of photosystem (PS) II (Fv/Fm) in drought-treated plants revealed a

photosynthetic impairment in plants expressing *cTP-SAC7*, *cTP-PTEN*, and *cTP-dOCRL* when

compared to WT (Fig. **5b**). In addition, exposing seedlings overexpressing *cTP-PTEN* to 30°C

for 9 days resulted in pale cotyledons as compared to WT (Supporting Information Fig. S5b).

Next, we performed an ultrastructural analysis of plants overexpressing PPI modifying

enzymes targeted to the plastids. No significant differences were observed compared to WT

plants when grown under normal growth conditions (Fig. **5b, c** and Supporting Information

Fig. S5c). In contrast, ultrastructural analysis of drought treated *cTP-PTEN* overexpression

lines showed a strong swelling of the stromal fraction and thylakoid membranes compared to

drought treated WT, indicating altered osmotic pressure within plastids of *PTEN.OX* plants

(Fig. **5c**). Interestingly, similar phenotypes have been described for *VIPP1* knockdown lines

(Zhang et al., 2012). Of note, we showed that VIPP1 and PI3P sensor protein colocalized (Fig.

3).

Taken together this data suggest that PPIs may serve as lipid signaling molecules within

chloroplasts, fine-tuning the response to abiotic stresses.

Discussion

Chloroplasts contain the most important membrane system in plant cells, thylakoid

membranes, where the light-dependent reaction of photosynthesis takes place. Notably,

chloroplast membranes show a distinctive lipid composition, mainly composed out of

galactolipids and sulfolipids and very few phospholipids. Whereas recent studies have shown

the bulk lipid composition of chloroplasts (LaBrant et al., 2018), presence and distribution of

minor or signaling lipids remains elusive. Hence, a better understanding of the lipid

composition of thylakoids and more broadly chloroplasts is currently crucial to understand their

biogenesis maintenance driving photosynthesis under steady state and stress conditions. To

date, presence of PPIs in the lipid composition of chloroplast membranes remains vague. This

is most probably due to their low abundance and high turnover rates, which make them difficult

to detect. Here, we show the detection of PPIs within the chloroplasts of tobacco (Fig. **1**

and **2**) and heat stressed *Arabidopsis* plants (Fig. **4**), using our PPI biosensors *in vivo*.

Synthesis and turnover of PPI pools are vital for regulating various membrane trafficking

processes membrane maintenance along with signaling transduction pathways (Meijer and

Munnik, 2003; Di Paolo and De Camilli, 2006; Heilmann, 2009). Therefore, it is likely that

PPI could play a similar role in chloroplasts. However, it remains unclear if PPIs can be

produced within the chloroplasts. If so, it suggests the potential presence of unknown PI kinases

and PPI phosphatases within this organelle. ARC3, a chloroplast division factor, represents a

chimera of prokaryotic FtsZ and part of eukaryotic phosphatidyl-4-phosphate 5-kinase and

localizes to the stroma (Shimada et al. 2004, Maple et al., 2007). Whether it functions as PI

Kinase remains so far unexplored. Alternatively, PPI might be transferred from ER to

chloroplast envelopes through membrane contact sites (LaBrant et al., 2018). However, the

exchange of lipids between ER and plastid outer envelope membranes is highly restricted

(Karkie et., 2019). Lipid exchange mediated by the action of flipases and flopases had been

speculated (Rajasekharan and Gummadi, 2011). Once inside the inner leaflet of the inner

envelope membrane PPIs could subsequently be transported within the chloroplast via the

action of lipid binding proteins or envelope vesicles (LaBrant et al., 2018, Hertle et al., 2021,

Schroda et al., 2021). So far, interactions between PPIs and chloroplast proteins, PDV1 and

PDV2, Chloroplast Sec14-like protein 1 (CPSFL1), WHEAT KINASE START1 (WKS1), and

VIPP1 have only been shown *in vitro*, using protein-lipid blot overlay assays (Gou et al., 2015;

Okazaki et al., 2015; Theis et al., 2019; Hertle et al., 2020). PDV1 and PDV2 are integral

components of the chloroplast division machinery, and they are responsible for regulating the

rate of chloroplast division (Okazaki et al., 2009). CPSFL1 plays a crucial role in ensuring

proper chloroplast development, promoting plant growth, and supporting optimal

photosynthetic function in Arabidopsis and *Chlamydomonas* (García-Cerdán et al., 2020;

Hertle et al., 2020; Kim et al., 2022). WKS1 provides resistance to *Puccinia striiformis f.*

sp tritici in wheat (*Triticum aestivum).* VIPP1 plays essential roles in thylakoid membrane

biogenesis, maintenance and repair under both normal and stress conditions in *Arabidopsis* and

Chlamydomonas (Nordhues et al., 2012; Zhang et al., 2016; Theis et al., 2020). Here we

demonstrated the co-localization of VIPP1 and the PI3P biosensor within the chloroplasts (Fig.

3a). Additionally, we observed alterations in the sub-chloroplast distribution pattern of VIPP1

in the presence of the PI3P modifier, PTEN (Fig. **3b**). Our findings showed that plants

expressing *cTP-PTEN* were sensitive to heat and drought stress and exhibited reduced

photosynthetic efficiency (Fig. **5b**). Interestingly, ultrastructural analysis of *ctp-PTEN*
overexpressing plants indicated osmolarity changes by swollen plastids and thylakoids (Fig.
5c). Given the critical role of VIPP1 in thylakoid maintenance and repair under stress
conditions (Zhang et al., 2016; Theis et al., 2020), it is possible that PI3P may influence the
function of VIPP1 under stress conditions. However, we did not detect any changes in VIPP1
stability or its association with multiple complexes in *cTP-PTEN*-expressing plants
(Supporting Information Fig. S3a, b) . Nevertheless, the *in vivo* association of PI3P with VIPP1
(Fig. **3**) might fine-tune the function of VIPP1 under stress conditions.

In fact, phosphoinositide signaling pathways are a pivotal component mediating plant
responses to various forms of stresses (Sui et al., 2008; Xue et al., 2009; Munnik and Vermeer,
2010; Zhai et al., 2013; Morales et al., 2019). Extensive research has confirmed that elevated
temperatures trigger PPI flux in plant cells. This phenomenon is not limited to the model plant
Arabidopsis (Mishkind et al., 2009; Annum et al., 2023) but extends across the plant kingdom,
encompassing species such as rice (Mishkind et al., 2009), wheat (Annum et al., 2022)*, Cycas
multipinnata*, and *C. panzhihuaensis* (Zhu et al., 2022). Hence, the inducibility of PPI flux
within the cells in response to heat stress appears to be evolutionarily conserved, indicating the
crucial role of these phospholipids in the response to high-temperature stress. We verified, that
high temperature and subsequently induction of PPI flux within the cell leads to relocation of
PPI biosensors, indicating an increase in PPI concentration inside the chloroplasts (Fig. **4**).
Thus, changing cytoplasmic PI or PPI pools most probably also affects PI and PPI flux in the
plastid.

Drought induces water deficiency and hinders plant growth. Numerous studies have illustrated
the crucial role of the phosphoinositide signaling pathway in drought stress across different
plants (Mikami et al., 1998; Perera et al., 2008; Wang et al., 2008; Zhai et al., 2013; Carpaneto
et al., 2017). The instant induction of *AtPIP5K1*, Phosphatidylinositol-4-phosphate 5-kinase,
transcript in Arabidopsis plants subjected to drought or abscisic acid treatment suggests the
participation of AtPIP5K1 in altering PPI pools and, consequently, its involvement in water-
stress signal transduction (Mikami et al., 1998). Furthermore, dehydration induces the levels
of PI5P through the activity of a myotubularin homolog (*AtMTM1*). MTM1 is itself induced
during dehydration and can produce PI5P from $PI(3,5)P_2$ (Ndamukong et al., 2010). Moreover,
it has been reported that drought stress induces synthesis of $PI(3,5)P_2$ in plants (Meijer et al.,
1999) in which $PI(3,5)P_2$ controls the decrease in vacuolar volume to counteract water
deficiency and facilitates the closure of stomata (Carpaneto et al., 2017). Analysis of expression
patterns showed that six out of nine Arabidopsis *PLC* genes are induced by dehydration (Tasma

et al., 2008). Similarly, the expression of maize *ZmPLC1* is induced under dehydration, and
this increased expression contributes to the improved drought tolerance of maize (Wang et al.,
2008). In addition, it has been reported that *Arabidopsis* transgenic plants, overexpressing
mammalian inositol polyphosphate 5-phosphatase display modified responses to abscisic acid
and significantly improved drought tolerance (Perera et al., 2008). These studies provide
evidence for a close association between phosphoinositide signaling pathways and water-stress
responses in plants. We observed that perturbation of PPI balance in chloroplasts upon
dehydration in plant overexpressing different PPI phosphatases resulted in drought sensitivity
(Fig. **5**). Our results suggest the generation and turnover of PPI pools within the chloroplasts
may play crucial roles in regulating signaling transduction pathways.

The alterations in PPI levels or localization triggered by dehydration can be detected by
downstream protein targets possessing PPI-binding domains, consequently altering their
activity or localization (Munnik and Vermeer, 2010). For instance, the activity of the
Arabidopsis trithorax-like factor ATX1 was modified upon dehydration stress due to the
binding of PI5P (Ndamukong et al., 2010). Hence, further research is required to
comprehensively characterize chloroplast proteins possessing PPI binding domains.

While organelles have distinct specialized functions as individual compartments, there is
evident that inter-organelle communication plays a vital role in preserving cellular function.
This collaborative interaction occurs through various membrane-mediated processes and
signaling molecules. It has been reported that phospholipids are a notable class of molecules
that promote these interactions (Meijer and Munnik, 2003; Ravi et al., 2021). Hence, it is
reasonable to hypothesize that PPIs may play a role in mediating cross-communication between
the chloroplast and other cellular organelles, such as the nucleus. Comprehensive research
efforts are required to elucidate the involvement of PPIs in orchestrating signaling pathways
and regulating crosstalk between chloroplasts and other organelles.

Material and methods

Generation of transgenic plants

For fluorescent reporters the cTP of RbcS (sequence in supplement) was amplified using
oligos (AH1 and AH2) containing recombination sites for InFusion® cloning into pDK366
linearized by SpeI. Resulting plasmids were linearized by SpeI and synthetic DNA of YFP,
mCherry, CFP or mEOS4b respectively (and codon optimized for expression in A. thaliana)
was inserted in frame to the n-terminal cTP by InFusion cloning. The resulting plasmids were
used to create chloroplast targeted fluorescent lipid sensor and lipid modifying proteins. In

brief, the coding sequence of the lipid binding domains of the pleckstrin homology (PH)
domain of human ARHGEF9 for PI5P binding, the PH domain of human rho guanidine
nucleotide exchange factor 7 (ARHGEF7) for PI(3,5)P$_2$ detection (Singh et al., 2021), the four
phosphate adaptor protein 1 and 2 (PH) domain of rat FAPP1 for PI4P detection, the PH domain
of mouse 1-phosphatidylinositol-4,5-bisphosphate diesterase delta (Plcd1) for PI(4,5)P$_2$
detection and putative phosphatidylinositol 3-phosphate binding site encoded by the FYVE
domain from human hepatocyte growth factor related tyrosine kinase substrate (HGS11)
(Simon et al., 2014) was codon optimized for expression in Arabidopsis were codon optimized
and synthesized with recombination sites (sequences in supplement) to be integrated into
pDK366 containing cTP_YFP, cTP_mCherry or cTP_mEOS4b fusion proteins under the
control of the 35S promotor by InFusion® cloning. mCherry-FKBP-PTEN was a gift from
Gerry Hammond (Addgene plasmid # 116866; http://n2t.net/addgene:116866;
RRID:Addgene_116866), SAC7/RHD4 (At3g51460) was amplified from Arabidopsis thaliana
cDNA. dOCRL was used as codon optimized synthetic DNA of the sequence publishes in
Doumane et al., 2021. VIPP1 was amplified from Arabidopsis cDNA using oligos
(AH15/AH16). Subsequent VIPP1 lacking the stop codon was fused in frame to codon
optimized CFP (sequence in supplement) by fusion PCR using oligos coding for recombination
sites for infusion cloning into pDK488 linearized by SpeI. For stable expression, we introduced
the constructs into *Arabidopsis* (Col-0) through a floral dip method utilizing Agrobacterium
tumefaciens (strain GV3101) (Clough and Bent, 1998). Transient expression in *N.*
benthamiana was conducted following the established procedure (Li et al., 2011).

Plant materials, growth conditions, and the procedures for heat and drought treatments
Arabidopsis thaliana Columbia (Col-0) was used as the WT. The seeds were germinated on
0.5× Murashige and Skoog agar medium enriched with 1% (w/v) sucrose. The seedlings were
cultivated following a 16-hour light cycle (120 μmol m^{-2} s^{-1}) at 22 °C, and 8 h dark at 22 °C.
To induce elevated PPI levels in the cells, 10-day-old Arabidopsis seedlings were subjected to
a heat regime of 40 °C for 30 min. To assess heat stress tolerance, 7-day-old Arabidopsis
seedlings were exposed to a heat regime of 45.5 °C for 45 min. To assess drought tolerance,
the seedlings were cultivated for 4 weeks in soil under a diurnal cycle comprising 16 hours of
light (120 μmol m^{-2} s^{-1}) and 8 hours of darkness at 22 °C. Subsequently, to induce drought
stress, watering was discontinued for a period of 10-14 days.

Chloroplast or thylakoid Isolation

To isolate chloroplasts, leaf material from the various genotypes was initially ground in chloroplast isolation buffer. The buffer composition included 0.45 M sorbitol, 20 mM Tricine (pH 8.4), 10 mM EDTA, 10 mM NaHCO3, and 0.1% BSA. Chloroplasts were collected by centrifugation for 5 min at 3000g, 4°C from the collected material obtained by filtering it through two layers of Miracloth. The released chloroplasts were then carefully layered onto a Percoll gradient made with 5X resuspension buffer (0.3 M sorbitol, 20 mM Tricine (pH 8.4), 2.5 mM EDTA, and 5 mM MgCl2). Centrifugation was performed in a swing-out rotor at 4°C for 10 minutes at 3300g. Intact chloroplasts were collected from the lower green band and subsequently subjected to a wash using 1X resuspension buffer.

Thylakoid membrane preparation from leaves of 3-week-old WT and *PTEN.OX* plants was conducted in accordance with the method described in (Bassi et al., 1985). Leaf material was homogenized in an ice-cold buffer comprising 0.4 M sorbitol, 0.1 M Tricine/KOH (pH 7.8), and 1 mM PMSF. The resulting homogenate was filtered through a nylon mesh and then centrifuged at 4°C, 3000g for 10 minutes. Subsequently, chloroplasts were disrupted in a buffer consisting of 20 mM HEPES/KOH (pH 7.8) and 10 mM EDTA (pH 8.0). Thylakoids were collected by centrifugation at 12,000g for 10 minutes and resuspended in a buffer containing 50% glycerol, 10 mM HEPES/KOH (pH 7.5), and 1 mM EDTA (pH 8.0). Fractionation of spinach chloroplasts was performed as described in (Hertle et. al 2020). Envelope fractions were separated into inner and outer envelopes using sucrose gradients as discribed in (Block et al., 2018).

Thermolysin treatment

Intact chloroplasts were treated with thermolysin based on Fitzpatrick et al., 2001. The chloroplast pellet, equivalent to 30 μg of chlorophyll was resuspended in import buffer (330-mM sorbitol and 50 mM HEPES–KOH, pH 8.0) and incubated with thermolysin (36 units mg^{-1}; 0.05 units $μg^{-1}$ chlorophyll) for 30 minutes on ice. Intact chloroplasts were re-isolated over a 40% v/v Percoll. chloroplasts were dissolved in 100ul of 4xSDS buffer.

Bis(sulfosuccinimidyl)suberate (BS3) treatment

Isolated thylakoids from WT and *PTEN.OX* were washed 5 times with 20 mM HEPES/ KOH pH 7.5 to remove the EDTA. Crosslinking was performed by adding 2mM BS3 to a thylakoid suspension containing 150 μg/ml of chlorophyll, followed by incubation on ice for 20 minutes.

The reaction was stopped with 150 mM TRIS HCL pH:7.5. The plates were dissolved in 5X

SDS sample buffer for further analysis.

Protein extraction and immunoblotting

Soluble proteins were extracted from Arabidopsis seedlings expressing PPI biosensors

following the method outlined in (Sedaghatmehr et al., 2016). The protein concentration was

determined using the Bradford protein assay. Subsequently, the proteins were separated by

SDS-PAGE, and the PageRule Plus Prestained Protein Ladder (10–250 kDa; Thermo

Scientific) served as the molecular weight marker.

For immunoblot analysis, the proteins were transferred onto either a Protan nitrocellulose or

Polyvinylidene Difluoride membrane. Primary antibodies used included a rabbit anti-VIPP1

polyclonal antibody from Agrisera, rabbit anti-TOC159 from Agrisera, and mouse anti-green

fluorescent protein (GFP) antibody from Clontech. Secondary antibodies, IRDye 800CW-

conjugated goat anti-rabbit or anti-mouse IgG (H+L), were employed at a 1:10,000 dilution

(LI-COR Biosciences). Band detection was performed using the Odyssey Infrared Imaging

System (LI-COR Biosciences).

GUV formation

To form GUVs, PVA assisted swelling was performed in PBS buffer according to Weinberger

et al., 2013. PVA-coated substrate was prepared by spreading and drying of 100–300 μL of

PVA (5%) solution on a glass slide. Afterwards 10–20 μL of plant lipid mixtures (2 mM final)

of MGDG/DGDG/SQDG/PG/PI (52:26:6.5:9.5:1) (v/v) according to Hölzl et al., 2019 Labrant

et al., 2018; Block et al,.1983), with or without 0.1% PI3P and with 0.01mol% DiI. dissolved

in chloroform were spead on the dried PVA film and placed under vacuum for 30 min for

solvent evaporation. A chamber was formed using vitrex and a coverglass filled with sucrose

(200 mM) containing PBS buffer. Following 15 min. incubation, GUVs were transferred into

an Eppendorf tube using a pipette until further use. For protein binding studies proteins were

added in PBS containing glucose in equal osmolarity in a final concentration of 0.5 μM.

Expression of recombinant his tagged VIPP1-mEOS4b and FYVE-mEOS4b

VIPP1 lacking the cTP and the stop codon was amplified and fused to mEOS4b by fusion PCR

using the primers AH13 and AH14. mEOS-FYVE was amplified from plant plasmids described

above using primers overlapping to pET28b. The PCR product was integrated into pET28b

linearized by restriction using BamHI and EcoRI by in Fusion cloning. Resulting plasmid was

used for inducible expression of 6xHis tagged VIPP1-mEOS. VIPP1 was expressed by adding

0,5 mM IPTG to a culture of O.D.600=0.5 grown at 37°C. The temperature was lowered to

30°C and the culture was grown for 3 more hours. The bacteria were harvested by

centrifugation and the pellet was snap frozen in liquid nitrogen and stored at -70°C until further

use. Recombinant protein purification was performed using Ni-NTA(Qiagen) according to the

manufacturers protocol for native proteins. The lysis buffer was supplemented with 2% Triton-

X-100 (Serva). To remove access detergent and imidazole, buffer exchange of the obtained

purified protein solution was achieved by 10x centrifugation of the solution through a 10.000

MW cut off centricon (Millipore) using PBS. The protein was stored at -70°C until used.

Fat blot assays

Fat blot assays were performed as described in (Munnik and Wierzchowiecka, 2013). FYVE-

mEOS was detected using a gel doc fluorescence blot reader (Biorad) at 488 nm excitation.

TEM microscopy

Leaf samples were fixed in 2.5% (w/v) glutaraldehyde, 2% (w/v) paraformaldehyde in 50 mM

sodium cacodylate (pH 7.4) for 1 hr under vacuum at RT. Subsequent fixation was continued

overnight at 4°C. After washing in PBS Post-fixation with was performed in 2% OsO4 and

0.8% $K_3Fe(CN)_6$ in 50 mM cacodylate buffer (pH 7.4) for 2 hr at RT. After rinsing, leaf

samples were post-stained with 2% (w/v) uranyl acetate for 2 h at RT. Subsequent dehydration

was performed using acetone and Spurr`s resin (Science Services) was infiltered.

Polymerisation was carried out at 60°C for 48hrs. Ultrathin sections (50-70 nm) were cut on a

UC-6 ultramicrotome (Leica) using a diamond knife. For transmission electron microscopy,

sections were deposited on formvar and carbon coated nickel grids and contrasted with

2% uranyl acetate and Reynold´s lead citrate. Images were acquired on a JEOL JEM F-200

(JEOL).

Confocal microscopy

Plants were imaged using a Leica TCS SP8 or SP5 confocal microscope. The

excitation/emission settings used were 405/415-475nm for CFP, 488/505-535 nm for GFP,

512/520-540 nm for YFP and mEOS4b, 561/570-600 nm for RFP and DiI, and 633/650-720 nm

for chlorophyll fluorescence. All images were acquired using sequential scanning settings. The

images analysis and quantification of fluorescence intensities was performed using the ImageJ software.

Acknowledgments

We thank the Deutsche Forschungsgemeinschaft (DFG) and the Max Planck Society (MPG) for funding. We thank Prof. Dr. Ralph Bock and his team from the Max Planck Institute of Molecular Plant Physiology (MPIMP, Potsdam, Germany) for supporting and hosting the project and engaging in insightful discussions.

We thank Dr. Nadja Tarakina and Bolortuya Badamdorj from the Max Planck Institute of colloids and interfaces for providing access and excellent technical assistance to the transmission electron microscope.

We thank Dr. Rumiana Dimova and Dr. Shreya Pramanik at Max Planck Institute of colloids and interfaces (MPIKGF, Potsdam, Germany) for help and discussions with plant GUV experiments

We thank Prof. Petra Bauer at the Institute of Botany, at the Heinrich Heine University (Düsseldorf, Germany) for hosting the project and discussions.

Author contributions

AH conceived the study. MS and AH designed and performed the experiments; FR generated PIP biosensor constructs. MS and AH wrote the manuscript. All authors agreed to the submission of the manuscript.

Funding

The project was funded by a research grant of the Deutsche Forschungsgemeinschaft (DFG), (HE8905/2-1) projektnummer: 452589609 and the Max Planck Society (MPG). The work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy – EXC-2048/1 – project ID 390686111. The work was supported by the Deutsche Froschungsgemeinschaft (DFG, German Research Foundation) Project no. 267205415-SFB 1208.

Conflict of interest

The authors declare no conflict of interest.

References:

Akhter S, Uddin MN, Jeong IS, Kim DW, Liu XM, Bahk JD (2016) Role of Arabidopsis AtPI4Kγ3, a type II phosphoinositide 4-kinase, in abiotic stress responses and floral transition. Plant Biotechnol J **14:** 215-230

Annum N, Ahmed M, Imtiaz K, Mansoor S, Tester M, Saeed NA (2022) (32)P(i) Labeled Transgenic Wheat Shows the Accumulation of Phosphatidylinositol 4,5-bisphosphate and Phosphatidic Acid Under Heat and Osmotic Stress. Front Plant Sci **13:** 881188

Annum N, Ahmed M, Tester M, Mukhtar Z, Saeed NA (2023) Physiological responses induced by phospholipase C isoform 5 upon heat stress in Arabidopsis thaliana. Front Plant Sci **14:** 1076331

Balla T (2013) Phosphoinositides: tiny lipids with giant impact on cell regulation. Physiol Rev **93:** 1019-1137

Bankaitis VA, Aitken JR, Cleves AE, Dowhan W (1990) An essential role for a phospholipid transfer protein in yeast Golgi function. Nature. 347:561-2.

Bassi R, dal Belin Peruffo A, Barbato R, Ghisi R (1985) Differences in chlorophyll-protein complexes and composition of polypeptides between thylakoids from bundle sheaths and mesophyll cells in maize. Eur J Biochem **146:** 589-595

Block MA. (2018) Isolation of Inner and Outer Membranes of the Chloroplast Envelope from Spinach and Pea. Methods Mol Biol 1829:137-144

Block MA, Dorne AJ, Joyard J, Douce R. (1983) Preparation and characterization of membrane fractions enriched in outer and inner envelope membranes from spinach chloroplasts. II. Biochemical characterization. J Biol Chem. **258**:13281-6.

Bovet L, Müller MO, Siegenthaler PA. (2001) Three distinct lipid kinase activities are present in spinach chloroplast envelope membranes: phosphatidylinositol phosphorylation is sensitive to wortmannin and not dependent on chloroplast ATP. Biochem Biophys Res Commun. 289: 269-75

Carpaneto A, Boccaccio A, Lagostena L, Di Zanni E, Scholz-Starke J (2017) The signaling lipid phosphatidylinositol-3,5-bisphosphate targets plant CLC-a anion/H(+) exchange activity. EMBO Rep **18:** 1100-1107

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J **16:** 735-743

DeWald DB, Torabinejad J, Jones CA, Shope JC, Cangelosi AR, Thompson JE, Prestwich GD, Hama H (2001) Rapid Accumulation of Phosphatidylinositol 4,5-Bisphosphate and Inositol 1,4,5-Trisphosphate Correlates with Calcium Mobilization in Salt-Stressed Arabidopsis. Plant Physiology **126:** 759-769

Di Paolo G, De Camilli P (2006) Phosphoinositides in cell regulation and membrane dynamics. Nature **443:** 651-657

Dieck CB, Boss WF, Perera IY (2012) A role for phosphoinositides in regulating plant nuclear functions. Front Plant Sci **3:** 50

Doumane M, Lebecq A, Colin L, Fangain A, Stevens FD, Bareille J, Hamant O, Belkhadir Y, Munnik T, Jaillais Y et al. (2021). Inducible depletion of PI(4,5)P$_2$ by the synthetic iDePP system in *Arabidopsis*. *Nat. Plants* **7**: 587–597.

García-Cerdán JG, Schmid EM, Takeuchi T, McRae I, McDonald KL, Yordduangjun N, Hassan AM, Grob P, Xu CS, Hess HF et al. (2020) Chloroplast Sec14-like 1 (CPSFL1) is essential for normal chloroplast development and affects carotenoid accumulation in Chlamydomonas. Proceedings of the National Academy of Sciences **117:** 12452-12463

Gerth K, Lin F, Daamen F, Menzel W, Heinrich F, Heilmann M (2017) Arabidopsis phosphatidylinositol 4-phosphate 5-kinase 2 contains a functional nuclear localization sequence and interacts with alpha-importins. Plant J **92:** 862-878

Gou JY, Li K, Wu K, Wang X, Lin H, Cantu D, Uauy C, Dobon-Alonso A, Midorikawa T, Inoue K et al. (2015) Wheat Stripe Rust Resistance Protein WKS1 Reduces the Ability of the Thylakoid-Associated Ascorbate Peroxidase to Detoxify Reactive Oxygen Species. Plant Cell **27:** 1755-1770

Goulden BD, Pacheco J, Dull A, Zewe JP, Deiters A, Hammond GRV (2019). A high-avidity biosensor reveals plasma membrane PI(3,4)P$_2$ is predominantly a class I PI3K signaling product. J Cell Biol. **218**:1066-1079.

Gupta TK, Klumpe S, Gries K, Heinz S, Wietrzynski W, Ohnishi N, Niemeyer J, Spaniol B, Schaffer M, Rast A et al. (2021). Structural basis for VIPP1 oligomerization and maintenance of thylakoid membrane integrity. Cell. **184**:3643-3659

Heilmann I (2009) Using genetic tools to understand plant phosphoinositide signalling. Trends in Plant Science **14:** 171-179

Hertle AP, García-Cerdán JG, Armbruster U, Shih R, Lee JJ, Wong W, Niyogi KK (2020) A Sec14 domain protein is required for photoautotrophic growth and chloroplast

vesicle formation in Arabidopsis thaliana. Proceedings of the National Academy of
Sciences **117**: 9101-9111

Hirayama T, Ohto C, Mizoguchi T, Shinozaki K (1995) A gene encoding a
phosphatidylinositol-specific phospholipase C is induced by dehydration and salt stress
in Arabidopsis thaliana. Proc Natl Acad Sci U S A **92**: 3903-3907

Hölzl, G., & Dörmann, P. (2019). Chloroplast Lipids and Their Biosynthesis. *Annual Review*
of Plant Biology, **70**, 51–81.

Horton P, Park KJ, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai K (2007)
WoLF PSORT: protein localization predictor. Nucleic Acids Res **35**: W585-587

Jensen JB, Falkenburger BH, Dickson EJ, de la Cruz L, Dai G, Myeong J, Jung S-R,
Kruse M, Vivas O et al. (2022) Biophysical physiology of phosphoinositide rapid
dynamics and regulation in living cells. Journal of General Physiology **154**

Karki, N., Johnson, B. S., & Bates, P. D. (2019). Metabolically Distinct Pools of
Phosphatidylcholine
Are Involved in Trafficking of Fatty Acids out of and into the Chloroplast for
Membrane Production. *The Plant Cell*, *31*(11), 2768–2788.

Kim E-H, Poudyal RS, Lee K-R, Yu H, Gi E, Kim HU (2022) Chloroplast-localized PITP7
is essential for plant growth and photosynthetic function in Arabidopsis. Physiologia
Plantarum **174:** e13760

König, S., Ischebeck, T., Lerche, J., Stenzel, I., & Heilmann, I. (2008). Salt-stress-induced
association   of phosphatidylinositol 4,5-bisphosphate with clathrin-coated vesicles
in   plants. *Biochemical   Journal*, **415**(3), 387–399

LaBrant E, Barnes AC, Roston RL (2018) Lipid transport required to make lipids of
photosynthetic membranes. Photosynthesis Research **138:** 345-360

Leshem Y, Seri L, Levine A (2007) Induction of phosphatidylinositol 3-kinase-mediated
endocytosis by salt stress leads to intracellular production of reactive oxygen species
and salt tolerance. The Plant Journal **51:** 185-197

Li L, Wang F, Yan P, Jing W, Zhang C, Kudla J, Zhang W (2017) A phosphoinositide-
specific phospholipase C pathway elicits stress-induced Ca(2+) signals and confers salt
tolerance to rice. New Phytol **214:** 1172-1187

Li X (2011) Infiltration of Nicotiana benthamiana protocol for transient expression via
Agrobacterium. Bio-protocol**:** e95-e95

Lu K-Y, Pasaje CFA, Srivastava T, Loiselle DR, Niles JC, Derbyshire E (2020)
Phosphatidylinositol 3-phosphate and Hsp70 protect Plasmodium falciparum from
heat-induced cell death. eLife **9:** e56773

Maple J, Vojta L, Soll J, Møller SG. (2007) ARC3 is a stromal Z-ring accessory protein
essential for plastid division. EMBO Rep 8(3):293-9

Matsuoka K, Bassham DC, Raikhel NV, Nakamura K (1995) Different sensitivity to
wortmannin of two vacuolar sorting signals indicates the presence of distinct sorting
machineries in tobacco cells. J Cell Biol **130:** 1307-1318

Meijer HJ, Berrie CP, Iurisci C, Divecha N, Musgrave A, Munnik T (2001) Identification
of a new polyphosphoinositide in plants, phosphatidylinositol 5-monophosphate
(PtdIns5P), and its accumulation upon osmotic stress. Biochem J **360:** 491-498

Meijer HJG, Divecha N, van den Ende H, Musgrave A, Munnik T (1999) Hyperosmotic
stress induces rapid synthesis of phosphatidyl-D-inositol 3,5-bisphosphate in plant
cells. Planta **208:** 294-298

Meijer HJG, Munnik T (2003) Phospholipid-Based Signaling in Plants. Annual Review of
Plant Biology **54:** 265-306

Mikami K, Katagiri T, Iuchi S, Yamaguchi-Shinozaki K, Shinozaki K (1998) A gene
encoding phosphatidylinositol-4-phosphate 5-kinase is induced by water stress and
abscisic acid in Arabidopsis thaliana. Plant J **15:** 563-568

Mishkind M, Vermeer JEM, Darwish E, Munnik T (2009) Heat stress activates
phospholipase D and triggers PIP2 accumulation at the plasma membrane and nucleus.
The Plant Journal **60:** 10-21

Morales-Johansson H, Jenoe P, Cooke FT, Hall MN (2004) Negative regulation of
phosphatidylinositol 4,5-bisphosphate levels by the INP51-associated proteins TAX4
and IRS4. J Biol Chem **279:** 39604-39610

Morales JA, Gonzalez-Kantun WA, Rodriguez-Zapata LC, Ramón-Ugalde J, Castano E
(2019) The effect of plant stress on phosphoinositides. Cell Biochemistry and Function
37: 553-559

Müller MO, Meylan-Bettex M, Eckstein F, Martinoia E, Siegenthaler PA, Bovet L. (2000)
Lipid phosphorylation in chloroplast envelopes. Evidence for galactolipid CTP-
dependent kinase activities. J Biol Chem 275: 19475-81

Mueller-Roeber B, Pical C (2002) Inositol Phospholipid Metabolism in Arabidopsis.
Characterized and Putative Isoforms of Inositol Phospholipid Kinase and
Phosphoinositide-Specific Phospholipase C. Plant Physiology **130:** 22-46

Munnik T, Meijer HJ (2001) Osmotic stress activates distinct lipid and MAPK signalling pathways in plants. FEBS Lett **498:** 172-178

Munnik T, Wierzchowiecka M. (2013) Lipid-binding analysis using a fat blot assay. Methods Mol Biol 1009:253-9

Munnik T, Vermeer JE (2010) Osmotic stress-induced phosphoinositide and inositol phosphate signalling in plants. Plant Cell Environ **33:** 655-669

Ndamukong I, Jones DR, Lapko H, Divecha N, Avramova Z (2010) Phosphatidylinositol 5-phosphate links dehydration stress to the activity of ARABIDOPSIS TRITHORAX-LIKE factor ATX1. PLoS One **5:** e13396

Nordhues A, Schöttler MA, Unger AK, Geimer S, Schönfelder S, Schmollinger S, Rütgers M, Finazzi G, Soppa B, Sommer F et al. (2012) Evidence for a role of VIPP1 in the structural organization of the photosynthetic apparatus in Chlamydomonas. Plant Cell **24:** 637-659

Okazaki K, Kabeya Y, Suzuki K, Mori T, Ichikawa T, Matsui M, Nakanishi H, Miyagishima SY (2009) The PLASTID DIVISION1 and 2 components of the chloroplast division machinery determine the rate of chloroplast division in land plant cell differentiation. Plant Cell **21:** 1769-1780

Okazaki K, Miyagishima SY, Wada H (2015) Phosphatidylinositol 4-phosphate negatively regulates chloroplast division in Arabidopsis. Plant Cell **27:** 663-674

Perera IY, Hung C-Y, Moore CD, Stevenson-Paulik J, Boss WF (2008) Transgenic Arabidopsis Plants Expressing the Type 1 Inositol 5-Phosphatase Exhibit Increased Drought Tolerance and Altered Abscisic Acid Signaling. The Plant Cell **20:** 2876-2893

Phillips SE, Vincent P, Rizzieri KE, Schaaf G, Bankaitis VA, Gaucher EA (2006) The diverse biological functions of phosphatidylinositol transfer proteins in eukaryotes. Crit Rev Biochem Mol Biol **41:** 21-49

Pical C, Westergren T, Dove SK, Larsson C, Sommarin M (1999) Salinity and Hyperosmotic Stress Induce Rapid Increases in Phosphatidylinositol 4,5-Bisphosphate, Diacylglycerol Pyrophosphate, and Phosphatidylcholine in Arabidopsis thaliana Cells*. Journal of Biological Chemistry **274:** 38232-38240

Pribat A, Sormani R, Rousseau-Gueutin M, Julkowska Magdalena M, Testerink C, Joubès J, Castroviejo M, Laguerre M, Meyer C et al. (2011) A novel class of PTEN protein in Arabidopsis displays unusual phosphoinositide phosphatase activity and efficiently binds phosphatidic acid. Biochemical Journal **441:** 161-171

Rajasekharan A, Gummadi SN. (2011) Flip-flop of phospholipids in proteoliposomes reconstituted from detergent extract of chloroplast membranes: kinetics and phospholipid specificity. PLoS One6(12):e28401

Ravi A, Palamiuc L, Emerling BM (2021) Crucial Players for Inter-Organelle Communication: PI5P4Ks and Their Lipid Product PI-4,5-P(2) Come to the Surface. Front Cell Dev Biol 9: 791758

Sedaghatmehr M, Mueller-Roeber B, Balazadeh S (2016) The plastid metalloprotease FtsH6 and small heat shock protein HSP21 jointly regulate thermomemory in Arabidopsis. Nature Communications 7: 12439

Shimada H, Koizumi M, Kuroki K, Mochizuki M, Fujimoto H, Ohta H, Masuda T, Takamiya K. (2004) ARC3, a chloroplast division factor, is a chimera of prokaryotic FtsZ and part of eukaryotic phosphatidylinositol-4-phosphate 5-kinase. Plant Cell Physiol 45(8):960-7

Siegenthaler PA, Müller MO, Bovet L. Evidence for lipid kinase activities in spinach chloroplast envelope membranes (1997). FEBS Lett. 416: 57-60

Simon ML, Platre MP, Assil S, van Wijk R, Chen WY, Chory J, Dreux M, Munnik T, Jaillais Y (2014) A multi-colour/multi-affinity marker set to visualize phosphoinositide dynamics in Arabidopsis. Plant J 77: 322-337

Singh N, Reyes-Ordoñez A, Compagnone MA, Moreno JF, Leslie BJ, Ha T, Chen J. (2021) Redefining the specificity of phosphoinositide-binding by human PH domain-containing proteins. Nat Commun: 12:4339.

Sui Z, Niu L, Yue G, Yang A, Zhang J (2008) Cloning and expression analysis of some genes involved in the phosphoinositide and phospholipid signaling pathways from maize (Zea mays L.). Gene 426: 47-56

Tasma IM, Brendel V, Whitham SA, Bhattacharyya MK (2008) Expression and evolution of the phosphoinositide-specific phospholipase C gene family in Arabidopsis thaliana. Plant Physiology and Biochemistry 46: 627-637

Testerink C, Munnik T (2011) Molecular, cellular, and physiological responses to phosphatidic acid formation in plants. J Exp Bot 62: 2349-2361

Theis J, Gupta TK, Klingler J, Wan W, Albert S, Keller S, Engel BD, Schroda M (2019) VIPP1 rods engulf membranes containing phosphatidylinositol phosphates. Sci Rep 9: 8725

Theis J, Niemeyer J, Schmollinger S, Ries F, Rütgers M, Gupta TK, Sommer F, Muranaka LS, Venn B, Schulz-Raffelt M et al. (2020) VIPP2 interacts with VIPP1

and HSP22E/F at chloroplast membranes and modulates a retrograde signal for

HSP22E/F gene expression. Plant, Cell & Environment **43:** 1212-1229

Thole JM, Vermeer JE, Zhang Y, Gadella TW Jr, Nielsen E. (2008) Root hair defective4

encodes a phosphatidylinositol-4-phosphate phosphatase required for proper root hair

development in Arabidopsis thaliana. Plant Cell;**20**:381-95.

Varnai P, Balla T (2006) Live cell imaging of phosphoinositide dynamics with fluorescent

protein domains. Biochim Biophys Acta **1761:** 957-967

Walker EH, Pacold ME, Perisic O, Stephens L, Hawkins PT, Wymann MP, Williams RL

(2000) Structural Determinants of Phosphoinositide 3-Kinase Inhibition by

Wortmannin, LY294002, Quercetin, Myricetin, and Staurosporine. Molecular Cell **6:**

909-919

Wang CR, Yang AF, Yue GD, Gao Q, Yin HY, Zhang JR (2008) Enhanced expression of

phospholipase C 1 (ZmPLC1) improves drought tolerance in transgenic maize. Planta

227: 1127-1140

Weinberger A, Tsai FC, Koenderink GH, Schmidt TF, Itri R, Meier W, Schmatko T,

Schröder   A, Marques C. (2013) Gel-assisted formation of giant unilamellar

vesicles. Biophys J.;**105**:154-64.

Xue HW, Chen X, Mei Y (2009) Function and regulation of phospholipid signalling in plants.

Biochem J **421:** 145-156

Zarreen F, Kumar K, Chakraborty S (2023) Phosphoinositides in plant-pathogen

interaction: trends and perspectives. Stress Biology **3:** 4

Zhai S, Gao Q, Liu X, Sui Z, Zhang J (2013) Overexpression of a Zea mays phospholipase

C1 gene enhances drought tolerance in tobacco in part by maintaining stability in the

membrane lipid composition. Plant Cell, Tissue and Organ Culture (PCTOC) **115:** 253-

262

Zhang L, Kato Y, Otters S, Vothknecht UC, Sakamoto W. (2012) Essential role of VIPP1

in   chloroplast envelope maintenance in Arabidopsis. Plant Cell. **24**:3695-707.

Zhang L, Kondo H, Kamikubo H, Kataoka M, Sakamoto W (2016) VIPP1 Has a

Disordered C-Terminal Tail Necessary for Protecting Photosynthetic Membranes

against Stress. Plant Physiol **171:** 1983-1995

Zhu H, Wu Y, Zheng Y (2022) Effects of heat shock on photosynthesis-related characteristics

and lipid profile of Cycas multipinnata and C. panzhihuaensis. BMC Plant Biology **22:**

442

Zufferey M, Montandon C, Douet V, Demarsy E, Agne B, Baginsky S, Kessler F (2017)
The novel chloroplast outer membrane kinase KOC1 is a required component of the
plastid protein import machinery. J Biol Chem **292:** 6952-6964

Figure 1: Expression of YFP-PI3P, YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P$_2$ and YFP-PI(3,5)P$_2$
biosensors in chloroplasts of tobacco and identification of their sub-organelle distribution. A.
The collection of biosensors is constructed to detect the different PPIs individually. Each
specific biosensor consists of a chloroplast transit peptide and a fusion between lipid-binding
domains that interacts specifically with each PPI and a yellow fluorescent protein (YFP),
mCherry or mEOS4b. B. scheme representing the distribution pattern of PPI biosensors in
chloroplast with or without binding to specific PPIs. The upper panel represents a lack of
binding of the PPI biosensor with diffuse localization of PPI biosensors in the chloroplast.
Lower panel represents the binding of PPI biosensors to PPI-containing membranes with

distinct localization of PPI biosensors. C. Confocal images of tobacco leaves transiently expressing PPI biosensors in the chloroplasts. Lacking the binding of the PI3P and PI5P biosensors results in diffuse localization of mEOS4b-PI3P and YFP-PI5P biosensors in the chloroplast. The binding of YFP-PI4P, YFP-PI(4,5)P$_2$ and YFP-PI(3,5)P$_2$ biosensors results in distinct punctate localization of PI biosensors indicating substrate recognition.

Figure 2: Sub-organelle distribution pattern of mEOS4b-PI3P, mEOS4b-PI4P and YFP-PI(4,5)P$_2$ biosensors in the presence of the different PPI phosphatases in the chloroplasts. A. The illustration shows a simplified view of the phosphatase activity of PTEN, SAC7, and dOCRL in chloroplasts. B. Distribution pattern of mEOS4b-PI4P biosensors in the presence of phosphatase, SAC7, in the chloroplasts of transiently expressing tobacco. C. Distribution pattern YFP-PI(4,5)P$_2$ biosensors without and in the presence of presence of PI5P and PI3P phosphatases, dOCRL-mCherry and PTEN-mCherry in the chloroplasts of tobacco. D. Distribution pattern of PI3P biosensors in the presence of PI3P and PI4P phosphatases, PTEN-mCherry and dOCRL-mCherry in the chloroplasts of tobacco. Fluorescence signals were visualized in tobacco leaves co-expressing PPI biosensors and phosphatases (as indicated) by fluorescence confocal microscopy.

Figure 3: Co-localization of VIPP1 with cTP-mEOS4b-PI3P (FYVE) biosensor and its differential sub-organelle localization in the presence of the PtdIns3P phosphatase, PTEN in the chloroplasts. A. Confocal images of tobacco leaves transiently expressing mEOS4b-PI3P (FYVE) biosensor in the presence and absence of VIPP1-CFP (middle and upper panel). Middle panels show colocalization of mEOS4b-PI3P biosensor with VIPP1-CFP in the chloroplasts. Lower panel shows different localization of mEOS4b-PI(4,5)P$_2$ biosensor and VIPP1-CFP in the chloroplasts. B. Sub-organelle distribution pattern of VIPP1-CFP in the presence of PI3P phosphatases, PTEN in the chloroplasts. C. Plant lipid based GUVs (magenta) with and without PI3P in the presence or absence of recombinant mEOS4b-VIPP1 (green).

Figure 4: Sub-organelle distribution pattern of YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P_2 and YFP-PI(3,5)P_2 biosensors as well as cTP-GFP in the chloroplast of *Arabidopsis* seedlings under control and heat stress (HS) conditions. A and B. Confocal images of 10-days-old *Arabidopsis* seedlings stably expressing *cTP-YFP-PI4P*, *cTP-YFP-PI5P*, *cTP-YFP-PI(4,5)P_2* and c*TP-YFP-PI(3,5)P_2* biosensors. (A) control and (B) HS (at 40°C for 30 minutes). The diagrams at

Figure 5: Perturbation of PIP abundance leads to drought sensitivity in *Arabidopsis* plants expressing PI4P phosphatase (SAC7), PI3P phosphatase (PTEN), and PI5P phosphatase (dOCRL). A. Sub-organelle distribution pattern of mEOS4b-PI3P, YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P$_2$ and YFP-PI(3,5)P$_2$ biosensors in the chloroplast of *Arabidopsis* seedlings under control and drought stress conditions. Confocal images of 10-days-old *Arabidopsis* seedlings stably expressing *cTP-mEOS4b-PI3P*, *cTP-YFP-PI4P*, *cTP-YFP-PI5P*, *cTP-YFP-PI(4,5)P$_2$* and c*TP-YFP-PI(3,5)P$_2$* biosensors. For drought stress leaf samples from plants growing on petri dishes were detached and kept for 30 minutes at RT prior to imaging. B. Phenotypes of WT, *cTP-PTEN.OX* (left), *cTP-SAC7.OX* (middle) and *cTP-DOCRL.OX* (right) plants under control condition (well watered; upper panel) and after drought stress (middle panel). Seedlings were grown for 4 weeks in soil under a diurnal cycle of 16 h light (120 µE m^{-2} s^{-1}) and 8 h dark at 22 °C. Drought stress was applied by withholding water for 10-14 days. The lower panel indicates *Fv/Fm* measurements of drought-stressed *cTP-PTEN.OX* (left) and *cTP-SAC7.OX* (right) plants. C. TEM images of *Arabidopsis* WT and *cTP-PTEN.OX* (left two rows). TEM images of Arabidopsis WT and plants expressing *cTP-PTEN.OX* following drought treatments (right two rows).

Supporting Information

Figure S1. Intensity plots of PPI biosensors in chloroplasts following transient expression in tobacco.

Figure S2. Colocalization of PPI biosensors with two different fluorescent proteins in the chloroplasts.

Figure S3. VIPP1 stability and/or its multiple complexes in *PTEN-expressing* plants and PI3P detection in chloroplasts.

Figure S4. Distribution pattern of YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P$_2$, and YFP-PI(3,5)P$_2$ biosensors in   the chloroplasts of Arabidopsis under heat stress conditions.

Figure S5. Distribution pattern of YFP-PI4P, YFP-PI5P, YFP-PI(4,5)P$_2$, and YFP-PI(3,5)P$_2$ biosensors in chloroplasts of Arabidopsis under drought stress conditions.

Supplementary Table 1. Oligos used in the study

Supplementary Table 2. Sequences of synthetic DNA or fluorescent reporters used in the study

969