Title: Genome composition predicts physiological responses to temperature in polyploid
salamanders

Keywords: metabolism, mitonuclear mismatch, total resistance to water loss, unisexual,
respiration efficiency ratio

Word count: 6558

Elements of manuscript: abstract, introduction, methods, results, discussion, references, table (1),
and figures (5)

Article type: main article

Short title: Subgenomes predict unisexual physiology

Isabella J. Burger[1], Michael Itgen[2], Lynn Tan[2], Parker Woodward[2], Linet Rivas-Moreno[2], Tamyra
Hunt[2], Hailey R. Ready[2], Xochitl G. Martin Geronimo[2], Robert D. Denton[2], Eric A. Riddell[1]

[1]Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599

[2]Department of Biology, Marian University, Indianapolis, IN 46222

Abstract

Multi-trait analyses can be used to measure the differential performance of phenotypic traits in

species complexes. Hybridization within these complexes can result in a mismatch between

mitochondrial and nuclear DNA that may lead to reduced performance and acclimation capacity

in hybrids. To test the effect of this mismatch on physiology, we compared physiological

performance and acclimation capacity of metabolic rate ($V\text{CO}_2$) and total resistance to water loss

(r_T) between two sexual *Ambystoma* species and a closely related unisexual lineage. We also

separated unisexuals by their unique biotypes to determine how physiology varies with

subgenomic composition. We found that unisexual biotypes exhibited phenotypes more like their

related sexual species than to other unisexuals. We also found a trade-off between r_T and $V\text{CO}_2$,

with increasing r_T resulting in a decrease in $V\text{CO}_2$. Although we did not find evidence for

mitonuclear mismatch, our results indicate that genomic composition of hybrids may be a

suitable predictor of hybrid trait performance. Multi-trait analyses are imperative for

understanding variation in phenotypic diversity, providing insight into how this diversity affects

species responses to environmental change.

Introduction

The formation of hybrids plays a key role in the process of speciation and extinction (Arnold, 1992). Hybrids can positively or negatively impact parental species through gene flow, competition, or the reinforcement of geographic ranges (Arnold, 1992; Barton, 2001; Coyne and Orr, 2004; Edelman et al., 2019). The outcome of these interactions can depend on differential fitness driven by phenotypic variation between hybrids and parental species (Barton, 2001; Thompson et al., 2021; Moran et al., 2021). Hybrids can outcompete one or both parental species by exhibiting beneficial phenotypes (i.e. heterosis; Schluter, 2000; Birchler et al., 2006; Hessenauer et al., 2020). Alternatively, hybrids can experience disadvantages due to Dobzhansky-Muller genetic incompatibilities (DMIs), in which the interactions between divergent alleles can result in reduced performance and, in some cases, hybrid lethality (Dobzhansky, 1982; Arnold and Martin, 2010; Thompson et al., 2023; Moran et al., 2024). Thus, the success of a hybrid often depends upon their performance relative to the parental lineages (Barton, 2001; Thompson et al., 2021). Understanding the effects of hybridization on performance can provide insight into the mechanisms that facilitate or impede hybrid success in species complexes.

The process of allopolyploidization—hybridization events that result in offspring with more than two sets of chromosomes—can reduce performance through the introduction of genetic incompatibilities and fundamental effects on cell size (Gregory and Mable, 2005; Balao et al., 2011; Mable, 2013; Wertheim et al., 2013; Moran et al., 2024). For instance, several key proteins involved in oxidative phosphorylation are comprised of interacting subunits originating from the nuclear and mitochondrial genomes. Thus, divergent nuclear and mitochondrial genomes may result in intergenomic incompatibilities between nuclear- and mitochondrial-

encoded protein subunits ('mitonuclear mismatch'), resulting in a reduction in mitochondrial

efficiency and capacity (Harrison and Burton 2006, Toews and Brelsford, 2012; Wolff et al.

2014), which in turn can lead to decreased performance (Sackton et al. 2003; Comai, 2005; Lee-

Yaw et al. 2014; Denton et al. 2017; Rank et al. 2020; Klabacka et al., 2022; Moran et al., 2024)

and fitness in hybrid offspring (Yee et al. 2013; Wolff et al. 2014). Therefore, mitonuclear

mismatch influences hybrid success by shaping physiological performance.

A secondary consequence of allopolyploidization is its effects on DNA content and,

consequently, cell size (Gregory, 2001; Balao et al., 2011). The total amount of DNA within a

cell (i.e., genome size) is strongly correlated with the size of the cell; therefore, polyploidization

causes a universal increase in cell size throughout the organism (Hanken and Wake, 1993;

Gregory, 2001; Itgen et al., 2022). Large genome and cell size have been hypothesized to drive

the evolution of physiological traits through the scaling effects of lower surface-area-to-volume

ratios in larger cells (Szarski, 1983; Vinogradov 1995; Kozłowski et al., 2003). Although broad-

scale taxonomic studies have found conflicting evidence for this relationship (Licht and

Lowcock, 1991; Uyeda et al., 2017; Gardner et al., 2020), there is empirical support for a cell

size-mediated role in determining metabolic rate. For example, larger cell size via polyploidy has

been shown to lower the metabolic cost of cells and, by extension, larvae in polyploid *Xenopus*

frogs by reducing the energetic costs of maintaining ion gradients (Cadart et al., 2023). In

addition, organisms with larger cell sizes may have thicker skin, which reduces desiccation risk

by lowering water loss rates (Feder and Burggren, 1985; Johnson et al., 2021). Therefore,

allopolyploidization might introduce two key consequences (i.e., cytonuclear incompatibility and

cell size) that affect physiological performance. Investigating the effects of allopolyploidization

on physiological traits is particularly important because these traits are highly integrated and

shape variation in suites of behavioral and life history traits that influence performance and

fitness (Ricklefs and Wikelski 2002; Wikelski et al. 2003; Martin et al. 2006).

Metabolic rate and water loss are two important physiological traits that covary in

terrestrial animals (Woods and Smith, 2010). Specifically, these traits are linked by the properties

of respiratory surfaces (e.g., lungs and skin) that facilitate the diffusion of gases, like oxygen and

carbon dioxide. Respiratory surfaces are thin and highly vascularized, which enables oxygen to

diffuse rapidly across the tissue (Czopek, 1962; Feder and Burggren, 1985). However, the

properties that facilitate oxygen uptake also result in high rates of water loss (Feder and

Burggren, 1985). This relationship between gas exchange and water loss results in a trade-off:

physiological changes to reduce water loss can also limit an organism's ability to breathe (Addo-

Bediako et al. 2001; Woods and Smith, 2010; Riddell et al. 2018a). By lowering water loss rates

(e.g., as an acclimation response to desiccation), organisms often experience a concurrent

reduction in metabolic rate due to lower rates of gas exchange (Anderson, 1970; Riddell et al.

2018a). This reduction in metabolic rate might lower an organism's ability to fuel energetically

demanding activities, such as foraging, reproduction, and territory-defense (Burggren and Vitalis,

2005; Biro and Stamps, 2010; Sadowska et al., 2015; Sokolova, 2021). However, the strength of

the trade-off can vary across species and hybrids (Burger et al., 2024), which may allow these

two linked traits to evolve more independently in different lineages. The relationship between

these two traits can also be evaluated through the water-gas exchange ratio (WGER) – the ratio

of oxygen uptake to water lost – that provides an index of an organism's respiration efficiency

(Burger et al., 2024). Understanding these linkages may help to predict organismal and

evolutionary responses to environmental change.

In this study, we explore the physiological consequences of allopolyploidization in a

lineage of allopolyploid salamanders (genus *Ambystoma*). We measured the thermal sensitivity

of metabolic rate, water loss rate, respiration efficiency, and the trade-off between water loss and

oxygen uptake in the unisexual *Ambystoma* complex. We also investigated the acclimation

capacity (defined as the capacity of an organism to reversibly adjust a physiological trait in

response to a new environment) of these traits because acclimation provides insight into a

species' ability to respond to environmental change (Riddell et al. 2018b). Specifically, we

compared traits between two sexual, diploid *Ambystoma* species (Jefferson salamander [*A.*

jeffersonianum] and blue-spotted salamander [*A. laterale*]) and unisexual, polyploid salamander

hybrids (Figure 1). We hypothesize that the unisexual allopolyploid salamanders will differ in

physiological performance compared to the sexual species due to the consequences of

allopolyploidization (i.e., cytonuclear incompatibility and large cell size) (see *Study species* in

Methods). Due to the potential for cytonuclear incompatibilities, we predicted that unisexual

salamanders would have lower metabolic rates, respiration efficiency, and acclimation capacity

compared to *A. jeffersonianum* and *A. laterale*. Due to larger cell sizes (as a result of increased

ploidy), we predicted that unisexuals will exhibit lower metabolic rates as well as water loss rates

(Johnson et al., 2021; Cadart et al., 2023). Analyzing multiple facets of an organism's

ecophysiology will allow us to develop a more comprehensive understanding of the effects of

hybridization and environmental variation on whole-organism performance.

Materials and methods

Study species

Unisexual *Ambystoma* are an all-female salamander lineage that is the oldest recognized

unisexual vertebrate group, persisting for ~5 million years (Bi and Bogart, 2010). Unisexuals

reproduce via kleptogenesis, in which they must "steal" sperm from a congeneric male to

stimulate egg development (Bogart et al. 2007; Gibbs and Denton, 2016). The genetic material

from males can then either be discarded (offspring are clones of mother) or incorporated into the

chromosome array of the embryos. This process has resulted in a range of polyploidy biotypes

(2N-5N) that combine genomes from five different species, but the majority of populations are

triploid with genomic contributions from *A. laterale* and *A. jeffersonianum* (Robertson et al.

2006; Bi and Bogart, 2010, Bogart et al. 2007). While frequent genome introgression from

sexual species into unisexual populations has produced highly variable combinations of nuclear

genomes (Bogart, 2019; Gibbs and Denton, 2016), the mitochondrial genome is strictly inherited

matrilineally and monophyletic (Hedges et al., 1992; Spolsky et al., 1992). The lack of co-

evolutionary history between the mitochondrial and nuclear genomes may result in mitonuclear

mismatch and likely has energetic consequences for unisexual salamanders by reducing

endurance relative to sympatric sexual species (Denton et al. 2017). Additionally, triploid

unisexuals have larger cells than diploid individuals (Licht and Bogart, 1987), and cell size has

been linked to variation in metabolic rate and water loss rates in amphibians (Johnson et al.

2021; Cadart et al., 2023; but see Licht and Bogart, 1990). However, studies have yet to compare

unisexual biotypes with related sexual species across their range (Figure 1) to make

comprehensive comparisons and understand if unisexual biotypes exhibit phenotypes more like

the sexual species to which they are most genetically similar. In this study, we compare a suite of

physiological traits between *A. jeffersonianum, A. laterale* and unisexuals from the same

populations with different genomic composition to determine if mitonuclear mismatch affected

unisexual physiological performance. Each of these 'biotypes' is described by using the first

letter of the parental species epithet, e.g. a LLJ individual is triploid with two sets of

chromosomes from *A. laterale* and one set of chromosomes from *A. jeffersonianum.*

Animal collections and housing

We collected animals between February and April 2020, 2022, and 2023 during breeding

migrations. We collected *Ambystoma jeffersonianum* from Litchfield, CT; Campbell, KY; and

Delaware, OH; *A. laterale* from Calumet, WI; Marquette, MI; Livingston, MI; Windham, CT;

and Addison, VT; and unisexuals from each location (Figure 1B, see figure for sample sizes).

These sites were chosen to maximize representation across the range of the unisexual polyploids

and target previously surveyed sites that have biotypes most representative of the lineage

(triploids, either LJJ or LLJ). We captured sexually mature individuals using a combination of

nighttime surveys, terrestrial drift fences, bucket traps, and/or cover boards, depending on the

approved collection methods for each state. We acquired scientific collection permits where

required (KY:SC2311177, MI:FSCP12302022102023, OH:18-164/SRP23-01, VT:SR-2023-03,

WI:SRLN-23-46) and worked with local field personnel when possible to help state agencies as

part of larger survey efforts.

We housed salamanders in individual clear boxes (19 x 33 x 10 cm) within refrigeration

units set at 12-14˚C immediately following collection. Each container contained moist paper

towels that were changed weekly and a PVC hide. We fed animals 1-2 appropriately sized

European Nightcrawlers (*Eisenia hortensis*) per week. We covered the front glass of the

refrigeration units except for narrow slits that would let in ambient light from the room during

normal building hours (~9am-5pm). We weighed each animal once per month and checked

animals each day to monitor for any signs of stress. The animal care protocol identified signs of

stress as general lethargy, unresponsive to stimulus, weight loss, or unusual skin coloration. None

of the animals in this study demonstrated those behaviors, and therefore animals were not

removed from our final dataset at this stage (see below for removing salamanders that were

active during respirometry trials). Prior to the initiation of this study, we acquired approval from

IACUC (protocol #RD-2021-1R). In May 2023, we transported salamanders to [university

redacted for double-blind review] for physiological experiments. For transport, we stored

individuals in smaller plastic containers lined with dampened paper towels. Total transport lasted

approximately 7.5 hours, at which time we immediately removed salamanders from the vehicle

and transferred them to a walk-in environmental chamber (2.5 x 3 m) set to 14°C for the duration

of the experiment (except for the acclimation trials – see *Physiological experiments*). We kept

individuals in the same plastic containers described above. We ensured that lights were kept off

in the housing unit, but salamanders were exposed to passive light through a small access

window during the day. We made all efforts to keep the housing at the experimental laboratory as

similar as possible to that of the location where animals were housed after capture.

Genetic analysis

We clipped a small tissue sample from the tail of each captured animal to determine their genetic

identity. First, we sequenced a highly variable portion of the mitochondrial genome and

compared it to known reference sequences for multiple *Ambystoma* species to determine species

identity (primers F-THR and R-651; McKnight and Shaffer, 1997; Bogart et al. 2007). For

individuals that were identified as unisexuals, we then used fragment length data from three

microsatellite loci to determine their specific ploidy and subgenome composition (*following*

Teltser and Greenwald, 2015). These subgenomes make up the separate unisexual biotypes (LLJ,

LJJ, etc.). One microsatellite locus amplifies only in *A. jeffersonianum*, whereas the other two

amplify in both species at different fragment size ranges. As in other studies (Bogart et al. 2007;

Bogart et al. 2009; Julian et al. 2003; Ramsden et al. 2005), this approach identifies subgenome

composition and can be validated across multiple individuals from the same site due to most

polyploid individuals in the same populations sharing the same genomic composition (Bogart

and Klemens 2008). Finally, the sites used for sample collection have undergone previous

surveys, establishing *a priori* expectations for the biotypes located there.

Study design

We conducted physiological experiments from May to July 2023. Prior to experimentation, we

separated salamanders into five batches of 28 individuals. We further divided these batches into

groups of seven, as only seven salamanders could be placed in the flow-through system at a time.

We randomized each group by site and species to ensure equal representation in each batch.

Salamanders were given one week at 14°C to adjust to laboratory conditions prior to taking

physiological measurements in the flow-through system (description below).

We exposed each salamander to three acute temperature treatments in the flow-through

system that represent a typical range of spring temperatures they would experience during active

periods in the wild: 6°C, 14°C, and 22°C (Lowcock et al., 1991; Hoffman et al., 2018; Mitchel,

2020). To ensure that salamanders reached thermal equilibrium prior to the run, we placed a

group in the incubator (Percival, Inc.; Model #I-36VL) at the experimental temperature for one

hour prior to physiological measurements. We ran two groups (14 salamanders) a day, and

individuals were given a rest day after each run to allow time for rehydration. We determined if a

salamander was rehydrated by monitoring their mass prior to and following the respirometry

trials. Therefore, it took two days to run a full batch at one temperature.

After the initial physiological measurements at each temperature, we moved the

salamanders to a separate incubator (Percival, Inc.; Model #I-36VL) set to 20°C for 20 days to

evaluate their capacity to acclimate by adjusting physiological traits. During this time, we

increased the rate of feeding to twice per week. We conducted the same physiological

measurements following this 20-day period to determine acclimation capacity of each individual.

For example, we ran a batch of salamanders at 6□, placed them in the 20□-acclimation chamber

for 20 days, then measured their physiology again at 6□. We compared physiological

performance between the two 6□ measurements (before and after exposure to the warm

acclimation chamber) to determine acclimation capacity. We conducted all physiological

measurements between 7:00 and 17:00 to measure physiological traits when salamanders would

be inactive in the field (McNab, 1997). Additionally, we stopped feeding salamanders one week

prior to experimental treatments to ensure a post-absorptive state. We ensured each individual

was at rest during the trials using a Reolink Network Video Recorder (Model RLN8-410) to

monitor and record salamander activity. We checked that salamanders were inactive through

visual inspection of the data output and verification using the video recordings. Activity is

characterized by erratic peaks in metabolic rate. Using this information, we removed all trials

where activity was clear and verified inactivity in the other trials using the recorded videos. We

removed all trials where the salamanders were active from the final dataset.

Flow-through system

In preparation for the physiological trials, we placed salamanders on a hardwire mesh platform in

a glass chamber (20 x 3.6 cm; volume *c*. 207 mL). Our primary objective was to understand

physiological responses to temperature; thus, the mesh restricted the salamander's ability to

behaviorally reduce water loss by curling onto themselves. We placed chambers in a

temperature- and humidity-controlled incubator (Percival, Inc.; Model #I-36VL) and connected

to a flow-through system (Sable Systems International (SSI), Las Vegas, NV). We delayed

physiological measurements by 30 minutes once the chambers were connected to allow the

salamanders to come to rest in the new environment.

For the 14°C and 22°C treatments, we used a bench pump (PP-2; SSI) to pull air from the

incubator. We also adjusted the relative humidity at each temperature to ensure a similar vapor

pressure deficit (VPD) across the treatments. The VPD is the difference between the saturated

vapor pressure (the amount of water the air can hold) and actual vapor pressure (the amount of

water in the air). As temperature increases, saturation vapor pressure increases, potentially

exposing organisms to higher VPDs (and thus greater evaporative demand) (Riddell et al. 2017,

Weaver et al. 2022). By maintaining the same VPD across temperature treatments, we ensured

that variation in water loss rates could be attributed to organismal responses to temperature and

not the effects of humidity. We adjusted humidity across these temperatures to maintain a VPD

of $0.75 \pm 5\%$ kPa. We were unable to use the incubator to control humidity at 6°C, so we split the

airstream being pulled by the pump into two columns: one pulling air straight from the incubator

and one pulling air from the incubator and through a column of Drierite^{TM}. We adjusted the ratio

of these two columns until we reached the target VPD (0.75 kPa).

After air was pumped out of the incubator, it passed through a flow-through manifold

(SSI) and separated into different streams with a continuous flow rate of 100 mL·min^{-1}. These

streams passed through the chambers containing the salamanders, as well as through an

additional, empty chamber that we used to collect baseline data. A multiplexer (MUX8; SSI)

cycled through the eight chambers (seven experimental and one baseline), and excurrent air

samples from each passed through a water-vapor analyzer (RH300; SSI) and a carbon dioxide

analyzer (CA-10A; SSI) to measure water vapor and CO_2, respectively, in volts. We calibrated

the analyzers once a month throughout the duration of the experiment to ensure accurate data

collection. We calibrated the CA-10 by passing N_2 gas and a known concentration of CO_2 gas

through the analyzer to zero and span the system, respectively. We calibrated the RH-300 by

passing N_2 gas through the system to set zero and using our humidity-controlled incubator to set

the span. The empty chamber was sampled between each salamander measurement to correct for

any drift throughout the duration of the experiment. We measured individuals three times at five-

minute intervals over a three-and-a-half-hour period.

We connected each instrument to a universal interface (UI-3; SSI) that uploaded the

voltage data to Expedata. We then linearly transformed the data in Expedata using equations

provided by the manufacturer (SSI). We used lag- and drift-corrections to correct for the time

delay in gas exchange/water loss and for drift that occurred during the experiment, respectively.

We also corrected the flow rate and carbon dioxide measurements for barometric pressure and

water vapor pressure of the excurrent air (Lighton 2008). We then transformed CO_2 values from

percentages into partial pressure by dividing the corrected data by 100. We measured the

asymptote of water loss values and used z-transformations to calculate stable water loss

measurements. We used equations 3-5 in Riddell et al. (2017) and 10.4 in Lighton (2008) to

calculate evaporative water loss (EWL), total resistance to water loss (r_T), and metabolic rate

(VCO_2) using the collected data. We measured total resistance to water loss as the inverse of

water loss, while accounting for the evaporative demand of the air (Riddell et al. 2017). We

averaged VCO_2, r_T, and EWL values across the measurements when salamanders were at rest.

We evaluated the relationship between gas exchange and water loss using two metrics:

the water-gas exchange ratio (WGER; VCO_2[mol]:EWL [mol]) and the trade-off between oxygen

uptake and resistance to water loss. The WGER is calculated as the rate of oxygen consumption

per unit water loss, providing an index of an organism's respiration efficiency from a water loss

perspective (Burger et al., 2024). The trade-off between oxygen consumption and resistance to

water loss more effectively captures physiological changes to minimize water loss (e.g.,

vasoconstriction of the skin, breathing rate; Riddell et al., 2019) and how these changes affect

oxygen uptake. However, resistance controls for the evaporative demand of the air and therefore

does not reflect the total amount of water loss; thus, we propose that analyzing both WGER and

the trade-off between oxygen uptake and resistance provides the most comprehensive

understanding of the relationship between gas exchange and water loss (Burger et al. 2024).

Statistical analysis

We ran three groups of analyses on the collected data: analysis of the traits, change in the traits

(acclimation), and the trade-off between resistance to water loss and gas exchange. For the trait

analyses, we ran three separate linear mixed effects models (LMER) to determine how VCO_2, r_T,

and respiration efficiency (WGER) varied throughout the experiment (*A. laterale*: n = 34; *A.*

jeffersonianum: n = 18; unisexual: n = 80). We included mass as a covariate and species,

temperature treatment, and time period (before and after acclimation) as factors in each separate

analysis. We also included interactions between species, temperature treatment, and time period

to assess differences between species. We included time period as an effect in these analyses to

account for the variation introduced by this treatment; however, we restricted our discussion of

the effects of acclimation to the change in trait analyses (see below). VCO_2 and WGER values

were log-transformed to meet assumptions of normality. For the change in the traits, we

conducted additional analyses that used the change in VCO_2, r_T, and respiration efficiency

between acclimation periods as the response variables (ΔVCO_2 [*A. laterale*: n = 32; *A.*

jeffersonianum: n = 18; unisexual: n = 76], Δr_T [*A. laterale*: n = 32; *A. jeffersonianum*: n = 18;

unisexual: n = 75]*,* and ΔWGER [*A. laterale*: n = 32; *A. jeffersonianum*: n = 18; unisexual: n =

75], respectively). Acclimation was calculated as the change in trait value after exposure to the

20□-acclimation treatment minus the trait value before exposure to the acclimation treatment for

each individual at each temperature treatment. For these analyses, we only included data from

individuals that had a value for both pre- and post-acclimation. We also removed time period as a

factor since we calculated the change in traits across the time periods. We conducted these two

groups of analyses (trait and change in the trait) to explicitly test our hypotheses on trait

performance and acclimation capacity. Additionally, we used a separate linear regression to

evaluate the trade-off between the change in VCO_2 and the change in r_T [*A. laterale*: n = 32; *A.*

jeffersonianum: n = 18; unisexual: n = 75]. All models included mass as a covariate and

salamander ID nested inside batch number and year collected as random effects. Year collected

was removed from models when it did not account for any additional variation in the data. We

were unable to include collection site in the above models due to its confounding relationship

with biotype (specific biotypes are only found in specific sites, Fig. 1); thus, we ran additional

analyses for each individual biotype to evaluate the effect of collection site on trait performance

(Appendix 1) and acknowledge that collection site may influence physiological rates in our

global models.

We were also interested in whether VCO_2, r_T, and respiration efficiency differed between

the unisexual biotypes. Therefore, we ran additional analyses on the traits above but included

biotype as a fixed effect as opposed to species (analysis of traits: LLJ: n = 39; LJJ: n = 36; trade-

offs: LLJ: n = 37; LJJ: n = 34). In these models, we treated biotype (*A. jeffersonianum* [JJ]*, A.*

laterale [LL]*,* LJJ, and LLJ), temperature, and time period as factors. For the change in the trait

analyses (ΔVCO_2 = LLJ: n = 37; LJJ: n = 35; Δr_T = LLJ: n = 37; LJJ: n = 34; ΔWGER = LLJ: n =

37; LJJ: n = 34), we removed time period as a factor (similar to the above analysis). We also

included mass as a covariate and the same random effects as the models described above. Sample

size of *A. jeffersonianum* and *A. laterale* were the same as above for all biotype analyses. If any

analysis was significant, we conducted pairwise analyses using the emmeans package to

investigate the differences in traits between specific biotypes. We then compared the species-

based models with the biotype-based models using AIC to determine which set of models were a

better predictor for physiological performance.

Additionally, we calculated ω^2 to measure the effect size of each variable in each analysis

using the following equation:

335
$$\omega^2 = \frac{SS_{treatment} - df_{treatment} \times MS_{error}}{SS_{total} + MS_{error}}$$

where $SS_{treatment}$ is the sum of squares for the parameter, $df_{treatment}$ is the degrees of freedom for the

parameter, MS_{error} is the mean square error, and SS_{total} is the total sum of squares (Olejnik and

Algina, 2003). We also calculated marginal and conditional R^2 to describe the variation

accounted for by the fixed effects and the variation accounted for by all effects in the model

(fixed and random), respectively. We show all data as mean ± SEM using the *effects* package in

R, which returns an adjusted mean incorporating the effects of all variables in the model. We

conducted all analyses in R (version 4.3.1). Table 1 reports the outputs from the above linear

mixed effects models, and Table S1 reports the outputs from the pairwise comparisons in our

biotype models.

Results

Metabolic rate (VCO_2)

Our results indicated that VCO_2 was positively related to mass, with larger individuals having

higher metabolic rates (see Table 1 for all statistics). We also found that VCO_2 increased with

temperature, and *A. jeffersonianum* had a higher VCO_2 than *A. laterale* and unisexuals. The

interaction between temperature and species also influenced VCO_2, with *A. jeffersonianum*

having the highest sensitivity to the changes in temperature (Figure S1A). Moreover, 78% of the

variance was explained by our treatments and mass as indicated by the marginal R^2. In the

ΔVCO_2 analysis, salamanders reduced metabolic rate by 19% following the 20□-acclimation

period (Figure S1B). Additionally, ΔVCO_2 was positively related to mass and was greatest at

22°C. There was also an effect of the interaction between temperature and species on ΔVCO_2,

with *Ambystoma jeffersonianum* having the lowest change in metabolic rate at 6°C but the

highest change at 22°C (Figure S1B). We found a marginal R^2 of 12%, with an additional 26% of

the variation being explained by the random effects (conditional $R^2 = 38\%$).

When we separated unisexuals by their triploid biotype (LJJ, LLJ), we found that VCO_2

increased with mass and temperature (Figure 2A). We also found that biotypes had different

metabolic rates across temperatures, with unisexual biotypes exhibiting metabolic rates most

similar to the sexual species with which they share the greater number of chromosome sets,

specifically at the extreme temperatures (e.g. LJJ similar to *A. jeffersonianum* (JJ) but not to LLJ;

Figure 2A; Table S1). For the ΔVCO_2 analysis, ΔVCO_2 was positively related to mass and was

greatest at 22°C. We also found a significant effect of the interaction between temperature and

biotype on ΔVCO_2, with ΔVCO_2 varying across biotypes and temperatures (Figure 2B). Follow-

up pairwise analyses revealed a difference between JJ and LLJ at 6□ for the acclimation analysis

($p = 0.03$), but there were no other pairwise differences between biotype pairs ($p > 0.05$; Table

S1). Using AIC model comparison, we also found that models with biotype were more supported

by the data than models that used species for the trait analysis (species: AIC = 281.18; biotype:

AIC = 225.78) and acclimation analysis (species: AIC = 2922.08; biotype: AIC = 2901.88).

Total resistance to water loss (r_T)

For total resistance to water loss (r_T), we found larger individuals and warmer temperatures had a

higher r_T (Figure S2A). Additionally, the interaction between temperature and species affected r_T,

with *A. laterale* having the lowest thermal sensitivity of resistance (Figure S2A). We also found

that our treatments and mass explained 97% of the variation in our data. When analyzing Δr_T, we

found that total resistance to water loss declined by 3% following exposure to the 20□-

acclimation chamber (Figure S2B). We also found that Δr_T was greater in *A. jeffersonianum* than

A. laterale or unisexuals (Figure S2B). Mass and treatments accounted for 7% of the variation, as

shown by the marginal R^2 value, with an additional 10% being explained by the random effects

(conditional $R^2 = 17\%$).

When we separated unisexuals by their biotype, we found that r_T was positively related to

body mass, with larger individuals having the highest resistance, and r_T increased with

temperature (Figure 3A). The interaction between biotype and temperature also influenced r_T,

with LJJ and *A. jeffersonianum* (JJ) having the highest thermal sensitivity of resistance (Figure

3A). Similar to the VCO_2 results, unisexual species exhibited a r_T more similar to the sexual

species they are more genetically like, specifically at the two extreme temperatures (Figure 3A;

Table S1). For the Δr_T analysis, there was no effect of the interaction between temperature and

biotype. However, there were differences among biotypes, with *Ambystoma jeffersonianum* (JJ)

having the highest Δr_T value (Figure 3B). Using AIC model comparison, we found that models

with biotype were more supported by the data than models that used species for the trait analysis

(species: AIC = 2320.77; biotype: AIC = 2208.51). However, both models had similar AIC

values for the change in trait analysis (species: AIC = 1185.87; biotype: AIC = 1184.54).

Respiration efficiency (WGER)

Respiration efficiency and mass were positively related, as were efficiency and temperature

(Figure S3A). Furthermore, *A. jeffersonianum* had the highest respiration efficiency. We also

found that the interaction between temperature and species influenced efficiency, with *A. laterale*

having the least variation in efficiency across temperatures (Figure S3A). Our treatments and

mass accounted for 83% of the variation, based on the marginal R^2. In the ΔWGER analysis, we

found that salamanders were 18% less efficient following the acclimation period, and they had

the largest decrease in efficiency at 22°C (Figure S3B). Additionally, the interaction between

temperature and species affected efficiency, with *Ambystoma jeffersonianum* having the greatest

fluctuation in ΔWGER across temperatures (Figure S3B). Lastly, we found a marginal R^2 of

15%, with an additional 25% of the variation being explained by the random effects (conditional

$R^2 = 40\%$).

When we separated the unisexual group by biotype, we found that larger individuals had

the highest respiration efficiency, and respiration efficiency increased as temperatures warmed

(Figure 4A). We also found that the interaction between temperature and biotype was significant.

Specifically, *A. jeffersonianum* and LJJ exhibited similar efficiencies at 6□ and 22□, as did *A.*

laterale and LLJ (Figure 4A; Table S1). For ΔWGER, we found that all values were negative,

indicating that individuals had lower respiratory efficiency following the acclimation treatment

(Figure 4B). Additionally, ΔWGER was positively related to both mass and temperature. We also

found an effect of the interaction between biotype and temperature on ΔWGER, with acclimation

capacity varying by biotype and temperature (Figure 4B). However, the pairwise comparison did

not identify any significant differences between biotypes in the ΔWGER analysis (Table S1).

Using AIC model comparison, we found that models with biotype were more supported by the

data than models that used species for the trait analysis (species: AIC = 217.35; biotype: AIC =

158.96); however, models with species were more supported by the data than models that used

biotype for the change in trait analysis (species: AIC = -2667.24; biotype: AIC = -2623.78).

Trade-offs between gas exchange and water loss

We also found a trade-off between the change in r_T and $V\text{CO}_2$, in which the change in resistance

was negatively related to the change in $V\text{CO}_2$ (Figure 5). We did not find any differences in the

strength of the trade-off between species (Figure 5A) or temperature (Figure 5B). When

separating unisexuals by biotype, we also found a significant trade-off between the change in r_T

and $V\text{CO}_2$. However, we did not find any variation in the strength of the trade-off by biotype

(Figure 5C). Using AIC model comparison, we found that models with biotype were more

supported by the data than models that used species for the tradeoff analysis (species: AIC =

2536.86; biotype: AIC = 2502.35).

Discussion

The unisexual salamander system combines two potential consequences of allopolyploidization

that may negatively impact metabolic processes: genetic incompatibilities from the interactions

between divergent genomes and larger cell size via polyploidy. Our initial predictions were that

unisexuals, regardless of biotype, would have reduced metabolic rate and respiration efficiency

due to mitonuclear mismatch and lower metabolic and water loss rates due to cell size. However,

we found that physiological performance of unisexual polyploids mirrored the most

representative species in their sub-genome composition (Figure 1A, 2A, 3A), and unisexuals

exhibited similar acclimation capacity compared to the sexual species (Figure 1B, 2B, 3B).

Therefore, we found no evidence of allopolyploidization influencing physiological performance

in unisexual salamanders, similar to other studies investigating metabolic and water loss rates in

this system (Licht and Bogart, 1990; Licht and Lowcock, 1991) but unlike studies that

investigated endurance (Denton et al., 2017). To the best of our knowledge, this study is the first

to include both sexual species that contribute genomes to the unisexual lineage (*A. laterale* and

A. jeffersonianum) and subsequently the first to show that subgenome composition may play a

larger role than ploidy in estimating physiological performance in this system. These results

highlight the importance of including the performance of the parental species when measuring

and interpreting the performance of hybrid offspring.

Accounting for the subgenome composition in hybrids may be necessary for

understanding variation in hybrid performance across traits. Unisexual *Ambystoma* have greater

performance in certain traits (i.e., tissue regeneration) compared to congenerics (Saccucci et al.

2016), but they have lower performance in traits related to endurance compared to sexual

species, possibly due to mitonuclear mismatch (Denton et al. 2017). Alternatively, the unisexual

Ambystoma salamanders in this study broadly display intermediate values between the sexual

species and mirror the values of their major subgenome when comparing unique biotypes (i.e.

LLJ is more like *A. laterale* than LJJ). The inconsistencies of unisexual performance compared

to sexual congenerics across these studies may be due to the differences in the genomic array of

the unisexuals in each study. While the unisexual lineage is very old (Bogart et al. 2007; Bi and

Bogart 2010), it readily adds genomic variation from nearby species into its genome (Bi et al.

2008; Gibbs and Denton 2016; Denton et al. 2018). Therefore, variation in unisexual

performance across different traits (e.g., metabolism and total resistance to water loss, this study;

tissue regeneration, Saccucci et al. 2016; and endurance/dispersal rates, Denton et al. 2017)

likely depends on the genomic composition of each individual and the relative performance of

the sexual species from which each genome was received. Additionally, our study measured

resting metabolic rate as opposed to active or maximal metabolic rate (as measured in Denton et

al. 2017). Metabolic rate during endurance trials relies on a higher rate of ATP production to fuel

the greater energetic demand (Clarke and Fraser, 2004); therefore, mitonuclear mismatch may

have more of an impact during activity since it reduces the production of ATP. Therefore,

accounting for the genomic array of the unisexuals and the physiological traits being measured is

necessary to understand hybrid performance across studies.

The similarities between unisexuals and sexual species in this study provide support for

the persistence of unisexuals across millennia. Unisexuals commonly underperform sexual

congenerics, having lower fecundity (Uzzell 1969), being avoided by sexual males in mate-

choice trials (Dawley and Dawley, 1986), and having lower endurance and dispersal rates

(Denton et al., 2017) compared to sexual species. Unisexuals also require the presence of

sympatric sexual species to reproduce, with unisexual populations declining when sexual species

are absent (Bogart et al., 2016). In fact, the persistence of unisexuals may even rely on cyclical

patterns of population fluctuations between competing unisexuals and sexual species (Greenwald

et al., 2016). Thus, these combined effects suggest that unisexuals should be outcompeted by

sexual species and not be able to persist. However, unisexuals exhibit faster growth rates (Licht

and Bogart, 1987), have a competitive advantage over certain congenerics as larvae (outcompete

A. laterale larvae but not *A. texanum*; Brodman and Krouse, 2007), can potentially drive sexual

populations to extinction (Clanton, 1934), and as shown here, do not exhibit any differences in

metabolic rate, water loss rate, or acclimation capacity. Understanding the coexistence between

sexual and unisexual species may require linking physiology to organismal performance across

ecological and environmental contexts, especially given that the species in this study are

considered threatened or protected in many parts of their range (Canada's Species at Risk Act,

2002; Illinois DNR, 2020; Ohio DNR, 2022;).

Species responses to environmental change may be partly explained by the environmental

conditions experienced throughout their geographic range (Navas, 1996; Wilson, 2001; Lancaster

et al., 2020). *Ambystoma laterale* is the most cold-tolerant salamander in North America (Storey

and Storey 1992) and is primarily found in higher latitudes and previously glaciated regions

(Figure 1B), so higher performance at cooler temperatures may be an adaptive advantage for this

species. We observed higher metabolic rates, resistance to water loss, and respiration efficiency

in *A. laterale* at cool temperatures, which might facilitate prolonged activity while minimizing

desiccation risk at temperatures they experience more regularly. *Ambystoma laterale* also have

the most widespread geographic range and therefore experience a broader thermal gradient

(Figure 1B). Their lower thermal sensitivity to acute changes in temperature (Figures 2-4) may

lessen any negative effects of higher thermal variation by minimizing energetic costs at different

temperatures. Alternatively, *A. jeffersonianum* is subject to warmer average temperatures, which

could explain the higher metabolic rate, resistance to water loss, and respiration efficiency at

higher temperatures. An increase in these traits at warmer temperatures may facilitate a greater

capacity for work while minimizing desiccation risk. *Ambystoma jeffersonianum* also have a

much smaller geographic range (Figure 1B), providing a potential explanation for their relatively

higher thermal sensitivity of physiological traits. Unisexuals experience a transitional climate,

overlapping both *A. laterale* and *A. jeffersonianum* ranges; therefore, their intermediate

phenotypes may be the result of evolving in this geographic region (Figure 1B) or simply the

additive effects of their sub-genomes (McElroy et al. 2017). Geographic locations may also

explain unisexual performance, since LLJ biotypes are often found in *A. laterale* populations,

and LJJ are found in *A. jeffersonianum* populations (Figure 1B; Appendix 1). These interspecific

differences and the environmental conditions that explain them could inform predictions on how

environmental change will affect species physiological responses, performance, and fitness

(Riddell et al., 2018).

Including phenotypic measurements of parental species is imperative for an accurate

interpretation of hybrid performance. In this study, we discovered that subgenome composition

was highly indicative of hybrid trait response, with biotypes performing more like the parental

species of their major subgenomes. Although we found no effect of allopolyploidization on the

traits studied here, measuring other metrics of performance, such as maximal metabolic rate,

could better elucidate the effects of allopolyploidization on hybrid performance. Therefore,

further investigation into phenotypic variation and the mechanistic drivers of this variation could

give insight into not only the persistence of certain lineages throughout millennia, but also the

persistence and vulnerability of various species towards current and future environmental

change.

References

Addo-Bediako, A., S. L. Chown, and K. J. Gaston. 2001. Revisiting water loss in insects: a large scale view. Journal of Insect Physiology 47.

Anderson, J. F. 1970. Metabolic rates of spiders. Pergamon Press 33:51-72.

Arnold, M. L. 1992. Natural hybridization as an evolutionary process. Annual Review of Ecology and Systematics 23:237-261.

Arnold, M. L., and N. H. Martin. 2010. Hybrid fitness across time and habitats. Trends in ecology & evolution 25:530–536.

Balao, F., J. Herrera, and S. Talavera. 2011. Phenotypic consequences of polyploidy and genome size at the microevolutionary scale: a multivariate morphological approach. The New phytologist 192:256–265.

Barton, N. H. 2001. The role of hybridization in evolution. Molecular ecology 10:551–568.

Bi, K., and J. P. Bogart. 2010. Time and time again: unisexual salamanders (genus Ambystoma) are the oldest unisexual vertebrates. BMC evolutionary biology 10:1-14.

Bi, K., J. P. Bogart, and J. Fu. 2008. The prevalence of genome replacement in unisexual salamanders of the genus Ambystoma (Amphibia, Caudata) revealed by nuclear gene genealogy. BMC evolutionary biology 8:158.

Birchler, J. A., H. Yao, and S. Chudalayandi. 2006. Unraveling the genetic basis of hybrid vigor. Proceedings of the National Academy of Sciences of the United States of America 103:12957-12958.

Biro, P. A. and J. A. Stamps. 2010. Do consistent individual differences in metabolic rate promote consistent individual differences in behavior? Trends in Ecology & Evolution 25:653–659.

Bogart J.P. 2019. A family study to examine clonal diversity in unisexual salamanders (genus *Ambystoma*). Genome 62:549-561.

Bogart, J. P. and M. W. Klemens. 2008. Additional distributional records of *Ambystoma laterale, A. jeffersonianum* (Amphibia: Caudata) and their unisexual kleptogens in northeastern North America. American Museum Novitates 3627:1-58.

Bogart, J. P., J. Bartoszek, D. W. A. Noble, and K. Bi. 2009. Sex in unisexual salamanders: discovery of a new sperm donor with ancient affinities. Heredity 103:483–493.

Bogart, J. P, J. E. Linton, and A. Sandilands. 2016. A population in limbo: unisexual salamanders (genus Ambystoma) decline without sperm-donating species. Herpetological Conservation and Biology. 12:41-55.

Bogart, J. P., K. Bi, J. Fu, D. W. Noble, and J. Niedzwiecki. 2007. Unisexual salamanders (genus Ambystoma) present a new reproductive mode for eukaryotes. Genome 50:119-136.

Brodman, R., and H.i D. Krouse. 2007. How blue-spotted and small-mouthed salamander larvae coexist with their unisexual counterparts. Herpetologica 63:135–43.

Burger, I. J., E. T. Carter, L. M. Magner, M. M. Muñoz, M. W. Sears, B. M. Fitzpatrick, and E. A. Riddell. 2024. Assessing hybrid vigour using the thermal sensitivity of physiological trade￿offs in tiger salamanders. Functional Ecology 38:143-152.

Burggren, W. W. and T. Z. Vitalis. 2005. The interplay of cutaneous water loss, gas exchange and blood flow in the toad, Bufo woodhousei: Adaptations in a terrestrially adapted amphibian. Journal of Experimental Biology 208:105–112.

Cadart, C., J. Bartz, G. Oaks, M. Z. Liu, and R. Heald. 2023. Polyploidy in Xenopus lowers metabolic rate by decreasing total cell surface area. Current Biology 33:1744-1752.e7.

Canada's Species at Risk Act. 2002. An Act respecting the protection of wildlife species at risk in Canada: List of Wildlife Species at Risk. Minister of Justice.

Clanton, W. 1934. An unusual situation in the salamander *Ambystoma jeffersonianum* (green). Occassioanl papers of the museum of zoology, University of Michigan 290.

Clarke, A. and K. P. P. Fraser. 2004. Why does metabolism scale with temperature? Functional Ecology 18:243–251.

Comai, L. 2005. The advantages and disadvantages of being polyploid. Nature Reviews Genetics 6:836–846.

Coyne, J. A., and H. A. Orr. 2004. Speciation: a catalogue and critique of species concepts. Philosophy of biology: an anthology 272-92.

Czopek, J. 1962. Vascularization of respiratory surfaces in some caudata. Copeia 1962:576-587.

Dawley, E. M., and R. M. Dawley. 1986. Species discrimination by chemical cues in a unisexual-bisexual complex of salamanders. Journal of Herpetology 20:114.

Denton, R. D., K. R. Greenwald, and H. L. Gibbs. 2017. Locomotor endurance predicts differences in realized dispersal between sympatric sexual and unisexual salamanders. Functional Ecology 31:915-926.

Denton, R. D., A. E. Morales, and H. L. Gibbs. 2018. Genome-specific histories of divergence and introgression between an allopolyploid unisexual salamander lineage and two ancestral sexual species. Evolution; international journal of organic evolution 72:1689-1700.

Dobzhansky T. 1982. Genetics and the Origin of Species. New York, NY: Columbia University Press.

Edelman, N. B., P. B. Frandsen, M. Miyagi, B. Clavijo, J. Davey, R. B. Dikow, G. García-Accinelli, et al. 2019. Genomic architecture and introgression shape a butterfly radiation. Science 366:594-599.

Feder, M. E. and W. W. Burggren. 1985. Skin breathing in vertebrates. Scientific American
253:126-143.

Gardner, J. D., M. Laurin, and C. L. Organ. 2020. The relationship between genome size and
metabolic rate in extant vertebrates. Philosophical transactions of the Royal Society of
London. Series B, Biological sciences 375:20190146.

Gibbs, H. L. and R. D. Denton. 2016. Cryptic sex? Estimates of genome exchange in unisexual
mole salamanders (Ambystoma sp.). Molecular Ecology 25:2805-2815.

Greenwald, K. R., R. D. Denton, and H. L. Gibbs. 2016. Niche partitioning among sexual and
unisexual *Ambystoma* salamanders. Ecosphere 7.

Gregory, T. R. 2001. The Bigger the C-Value, the Larger the Cell: Genome Size and Red Blood
Cell Size in Vertebrates. Blood Cells, Molecules, and Diseases 27: 830–843.

Gregory, T. R., and B. K. Mable. 2005. Polyploidy in animals. The evolution of the genome.
Academic Press 427-517.

Hanken, J., and D. B. Wake. 1993. Miniaturization of Body Size: Organismal Consequences and
Evolutionary Significance. Annual review of ecology and systematics 24:501–519.

Harrison, J.S. and R. S. Burton. 2006. Tracing hybrid incompatibilities to single amino acid
substitutions. Molecular Biology and Evolution 23:559–564.

Hedges, S. B., J. P. Bogart, and L. R. Maxson. 1992. Ancestry of unisexual salamanders. Nature
356:708-710.

Hessenauer, P., A. Fijarczyk, H. Martin, J. Prunier, G. Charron, J. Chapuis, L. Bernier, et al.
2020. Hybridization and introgression drive genome evolution of Dutch elm disease
pathogens. Nature ecology & evolution 4:626–638.

Hoffmann, K., M. Hunter, A. J. K. Calhoun, and J. Bogart. 2018. Post-breeding migration and
habitat of unisexual salamanders in Maine, USA. Journal of Herpetology 52:273–81.

Illinois Endangered Species Protection Board. 2020. Checklist of Illinois endangered and
threatened animals and plants. Illinois Department of Natural Resources,

Itgen, M. W., G. R. Natalie, D. S. Siegel, S. K. Sessions, and R. L. Mueller. 2022. Genome size
drives morphological evolution in organ-specific ways. Evolution; international journal
of organic evolution 76:1453–1468.

Johnson, B. B., J. B. Seale, and J. P. Sparks. 2021. Genome size influences adaptive plasticity of
water loss, but not metabolic rates, in lungless salamanders. The Journal of Experimental
Biology 224:242196.

Julian, S. E., T. L. King, and W. K. Savage. 2003. Novel Jefferson salamander, *Ambystoma*
jeffersonianum, microsatellite DNA markers detect population structure and hybrid
complexes. Molecular Ecology Notes. 3:95-97.

Klabacka, R. L., H. A. Parry, K. N. Yap, R. A. Cook, V. A. Herron, L. M. Horne, M. E. Wolak, J.
A. Maldonado, M. K. Fujita, A. N. Kavazis, J. R. Oaks, and T. S. Schwartz. 2022.
Reduced mitochondrial respiration in hybrid asexual lizards. American Naturalist.
199:719-728.

Kozłowski, J., M. Konarzewski, and A. T. Gawelczyk. 2003. Cell size as a link between
noncoding DNA and metabolic rate scaling. Proceedings of the National Academy of
Sciences of the United States of America 100:14080–14085.

Lancaster, L. T. and A. M. Humphreys. 2020. Global variation in the thermal tolerances of plants.
Proceedings of the National Academy of Sciences 117:13580–13587.

Lee☐Yaw, J. A., C. G. Jacobs, and D. E. Irwin. 2014. Individual performance in relation to
cytonuclear discordance in a northern contact zone between long☐toed salamander
(Ambystoma macrodactylum) lineages. Molecular Ecology 23:4590-4602.

Licht, L. E., and J. P. Bogart. 1987. Comparative Size of Epidermal Cell Nuclei from Shed Skin
of Diploid, Triploid and Tetraploid Salamanders (Genus Ambystoma). Copeia 1987:284–
290.

———. 1990. Comparative rates of oxygen consumption and water loss in diploid and polyploid
salamanders (genus ambystoma). Comparative biochemistry and physiology. Part A,
Physiology 97:569–572.

Licht, L. E., and L. A. Lowcock. 1991. Genome size and metabolic rate in salamanders.
Comparative Biochemistry and Physiology Part B: Comparative Biochemistry 100:83–
92.

Lighton, J. R. 2008. Measuring metabolic rates: a manual for scientists. Oxford University Press.

Lowcock, L. A., H. Griffith, and R. W. Murphy. 1991. The *Ambystoma laterale-jeffersonianum*
complex in central Ontario: ploidy structure, sex ratio, and breeding dynamics in a
bisexual-unisexual community. Copeia 1991:87.

Mable, B. K. 2013. Polyploids and hybrids in changing environments: winners or losers in the
struggle for adaptation? Heredity 110:95–96.

Martin I., B. Lynn, D. Hasselquist, and M. Wikelski. 2006. Investment in immune defense is
linked to pace of life in house sparrows. Oecologia 147:565-575.

McElroy, K. E., R. D. Denton, J. Sharbrough, L. Bankers, M. Neiman, and H. L. Gibbs. 2017.
Genome expression balance in a triploid trihybrid vertebrate. Genome Biology and
Evolution 9:968-980.

McKnight, M. L., and H. B. Shaffer. 1997. Large, rapidly evolving intergenic spacers in the
mitochondrial DNA of the salamander family Ambystomatidae (Amphibia:
Caudata). Molecular biology and evolution 14:1167-1176.

McNab, B. K. 1997. On the utility of uniformity in the definition of basal rate of metabolism.
Physiological Zoology 70:718–720.

Mitchel, K. B. 2020. Temporal changes in unisexual and sexual *Ambystoma* salamander
populations in Southesast Michigan. Master's thesis. Eastern Michigan University.

Moran, B. M., C. Payne, Q. Langdon, D. L. Powell, Y. Brandvain, and M. Schumer. 2021. The
genomic consequences of hybridization. eLife 10.

Moran, B. M., C. Y. Payne, D. L. Powell, E. N. K. Iverson, A. E. Donny, S. M. Banerjee, Q. K.
Langdon, et al. 2024. A lethal mitonuclear incompatibility in complex I of natural
hybrids. Nature 626:119–127.

Navas, C. A. 1996. Metabolic physiology, locomotor performance, and thermal niche breadth in
neotropical anurans. Physiological Zoology 69:1481–1501.

Ohio Division of Wildlife. 2022. Ohio's Listed Species: Wildlife that are considered to be
endangered, threatened, species of concern, special interest, extirpated, or extinct in Ohio
(Report No. 5356). Ohio Department of Natural Resources.

Olejnik, S., and J. Algina. 2003. Generalized eta and omega squared statistics: measures of effect
size for some common research designs. Psychological methods 8:434.

Ramsden, C. K. Beriault, and J. P. Bogart. 2005. A nonlethal method of identification of
Ambystoma laterale, A. jeffersonianum, and sympatric unisexuals. Molecular Ecology
Notes. 6:261-264.

Rank, N. E., P. Mardulyn, S. J. Heidl, K. T. Roberts, N. A. Zavala, J. T. Smiley, and E. P.
Dahlhoff. 2020. Mitonuclear mismatch alters performance and reproductive success in
naturally introgressed populations of a montane leaf beetle. Evolution 74:1724-1740.

Ricklefs, R. E., and M. Wikelski. 2002. The Physiology/Life-History Nexus. Trends in Ecology
& Evolution 17:462–68.

Riddell, E. A., E. K. Apanovitch, J. P. Odom, and M. W. Sears. 2017. Physical calculations of
total resistance to water loss improve predictions of species range models. Ecological
Monographs 87:21-33.

Riddell, E. A., E. Y. Roback, C. E. Wells, K. R. Zamudio, and M. W. Sears. 2019. Thermal cues
drive plasticity of desiccation resistance in montane salamanders with implications for
climate change. Nature Communications 10: 4091.

Riddell, E. A., J. McPhail, J. D. Damm, and M. W. Sears. 2018a. Trade□offs between water loss
and gas exchange influence habitat suitability of a woodland salamander. Functional
Ecology 32:916-925.

Riddell, E. A., J. P. Odom, J. D. Damm, and M. W. Sears. 2018b. Plasticity reveals hidden
resistance to extinction under climate change in the global hotspot of salamander
diversity. Science Advances 4:eaar5471.

Robertson, A. V., C. Ramsden, J. Niedzwiecki, J. Fu, and J. P. Bogart. 2006. An unexpected
recent ancestor of unisexual Ambystoma. Molecular Ecology 15:3339-3351.

Saccucci, M. J., R. D. Denton, M. L. Holding, and H. L. Gibbs. 2016. Polyploid unisexual
salamanders have higher tissue regeneration rates than diploid sexual relatives. Journal of
zoology 300:77–81.

Sackton T.B., R. A. Haney, and D. M. Rand. 2003. Cytonuclear coadaptation in Drosophila:
disruption of cytochrome c oxidase activity in backcross genotypes. Evolution 57:2315–
2325.

Sadowska, J., A. K. Gebczynski, K. Paszko, and M. Konarzewski. 2015. Milk Output and
Composition in Mice Divergently Selected for BMR. Journal of Experimental Biology
jeb111245.

Schluter, D. 2000. The Ecology of Adaptive Radiation. OUP Oxford.

Sokolova, I. 2021. Bioenergetics in environmental adaptation and stress tolerance of aquatic
ectotherms: linking physiology and ecology in a multi-stressor landscape. Journal of
Experimental Biology jeb236802.

Spolsky, C. M., C. A. Phillips, and T. Uzzell. 1992. Antiquity of clonal salamander lineages
revealed by mitochondrial DNA. Nature 356:706-708.

Storey, K. B., and J. M. Storey. 1992. Natural freeze tolerance in ectothermic vertebrates. Annual
review of physiology 54:619–637.

Szarski, H. 1983. Cell size and the concept of wasteful and frugal evolutionary strategies. Journal
of theoretical biology 105:201–209.

Teltser, C., and K. R. Greenwald. 2015. Survivorship of ploidy-variable unisexual Ambystoma
salamanders across developmental stages. Herpetologica 71:81-87.

Thompson, K. A., Y. Brandvain, J. M. Coughlan, K. E. Delmore, H. Justen, C. R. Linnen, D.
Ortiz-Barrientos, et al. 2023. The Ecology of Hybrid Incompatibilities. Cold Spring
Harbor perspectives in biology.

Thompson, K. A., M. Urquhart-Cronish, K. D. Whitney, L. H. Rieseberg, and D. Schluter. 2021.
Patterns, Predictors, and Consequences of Dominance in Hybrids. The American
naturalist 197:E72–E88.

Toews, D. P., and A. Brelsford. 2012. The biogeography of mitochondrial and nuclear
discordance in animals. Molecular ecology 21:3907-3930.

Uyeda, J. C., M. W. Pennell, E. T. Miller, R. Maia, and C. R. McClain. 2017. The Evolution of
Energetic Scaling across the Vertebrate Tree of Life. The American naturalist 190:185–
199.

Uzzell, T. M. 1964. Relations of the Diploid and Triploid Species of the Ambystoma
jeffersonianum Complex (Amphibia, Caudata). Copeia 1964:257-300.

Vinogradov, A. 1995. Nucleotypic effect in homeotherms: body-mass-corrected basal metabolic
rate of mammals is related to genome size. Evolution 49:1249-1259.

Weaver, S. J., H. Edwards, T. McIntyre, S. M. Temple, Q. Alexander, M. C. Behrens, ... and E. N.
Taylor. 2022. Cutaneous evaporative water loss in lizards is variable across body regions
and plastic in response to humidity. Herpetologica 78:169-183.

Wertheim, B., L. W. Beukeboom, and L. van de Zande. 2013. Polyploidy in animals: effects of
gene expression on sex determination, evolution and ecology. Cytogenetic and Genome
Research 140:256-269.

Wikelski, M., L. Spinney, W. Schelsky, A. Scheuerlein, and E. Gwinner. 2003. Slow pace of life
in tropical sedentary birds: a common-garden experiment on four stonechat populations
from different latitudes. Proceedings of the Royal Society of London. Series B:
Biological Sciences 270:2383-2388.

Wilson, R. S. 2001. Geographic variation in the thermal sensitivity of jumpint performance in the
frog *Limnodynastes peronii*. Journal of Experimental Biology 204:4227-4236.

Wolff, J.N., E. D. Ladoukakis, J. A. Enriquez, and D. K. Dowling. 2014. Mitonuclear
interactions: evolutionary consequences over multiple biological scales. Philosophical
Transactions of the Royal Society B: Biological Sciences 369:20130443.

Woods, H. A., and J. N. Smith. 2010. Universal model for water costs of gas exchange by
animals and plants. Proceedings of the National Academy of Sciences of the United
States of America 107:8469–8474.

Yee W. K. W., K. L. Sutton, and D. K. Dowling. 2013. In vivo male fertility is affected by
naturally occurring mitochondrial haplotypes. Current Biology 23:R55–56.

Table 1: Results for linear models from the trait value analyses, change of trait analyses, and trade-off analysis*.

Trait	Fixed Effects	Trait value analysis			Change in trait analysis		
		F statistic (df)	P value	Effect size (ω^2)	F statistic (df)	P value	Effect size (ω^2)
Metabolic rate – by species	Mass	335.38 (1,134)	< 0.001	0.71	3.50 (1,116)	0.06	0.02
	Temperature	726.77 (2, 538)	< 0.001	0.68	13.41 (2, 193)	< 0.001	0.12
	Species	3.88 (2, 116)	0.02	0.04	0.32 (2,107)	0.72	< 0.001
	Acclimation	120.41 (1, 549)	< 0.001	0.14	-	-	-
	Species × Temperature	15.46 (4, 535)	< 0.001	0.09	3.08 (4, 190)	0.02	0.04
	Species × Acclimation	3.53 (2, 539)	0.03	0.01	-	-	-
	Temperature × Acclimation	5.33 (2, 538)	0.005	< 0.001	-	-	-
	Species × Temperature × Acclimation	1.85 (4, 536)	0.12	< 0.001	-	-	-
Metabolic rate – by biotype	Mass	221.84 (1,142)	< 0.001	0.60	3.48 (1,114)	0.06	0.02
	Temperature	843.13 (2,531)	< 0.001	0.75	13.36 (2,191)	< 0.001	0.12
	Biotype	2.73 (3,115)	0.05	0.03	0.27 (3, 107)	0.85	< 0.001
	Acclimation	134.64 (1, 549)	< 0.001	0.18	-	-	-
	Biotype × Temperature	26.64 (6,530)	< 0.001	0.22	3.21 (6,190)	0.005	0.06
	Biotype × Acclimation	3.73 (3, 533)	0.01	0.02	-	-	-
	Temperature × Acclimation	5.36 (2, 531)	0.005	0.01	-	-	-
	Biotype × Temperature × Acclimation	1.83 (6, 530)	0.09	< 0.001	-	-	-
Total resistance to water loss – by species	Mass	7386.57 (1,122)	< 0.001	0.98	1.57 (1,121)	0.21	< 0.001
	Temperature	2555.63 (2,498)	< 0.001	0.88	1.03 (2,192)	0.36	< 0.001
	Species	1.37 (2,107)	0.26	< 0.001	6.83 (2,112)	0.002	0.09
	Acclimation	2.54 (1, 504)	0.11	< 0.001	-	-	-
	Species × Temperature	40.64 (4,493)	< 0.001	0.24	1.13 (4,190)	0.34	< 0.001
	Species × Acclimation	0.84 (2, 495)	0.43	< 0.001	-	-	-
	Temperature × Acclimation	1.46 (2, 495)	0.23	< 0.001	-	-	-
	Species × Temperature × Acclimation	0.62 (4, 493)	0.65	< 0.001	-	-	-

Total resistance to water loss – by biotype	Mass	3716.60 (1,139)	< 0.001	0.96	1.18 (1,111)	0.28	< 0.001
	Temperature	3271.40 (2,486)	< 0.001	0.93	1.00 (2,188)	0.37	< 0.001
	Biotype	1.16 (3,111)	0.33	<0.001	4.56 (3,109)	0.005	0.09
	Acclimation	4.64 (1, 506)	0.03	< 0.001	-	-	-
	Biotype × Temperature	58.07 (6,484)	< 0.001	0.41	1.43 (6,188)	0.21	0.01
	Biotype × Acclimation	1.14 (3, 485)	0.33	< 0.001	-	-	-
	Temperature × Acclimation	2.00 (2,483)	0.14	< 0.001	-	-	-
	Biotype × Temperature × Acclimation	0.92 (6, 482)	0.48	< 0.001	-	-	-
Respiration efficiency – by species	Mass	229.52 (1,136)	< 0.001	0.62	2.38 (1,85)	0.13	0.02
	Temperature	1332.91 (2, 503)	< 0.001	0.80	23.06 (2,164)	< 0.001	0.21
	Species	3.62 (2,116)	0.03	0.04	0.27 (2,85)	0.76	< 0.001
	Acclimation	111.84 (1, 512)	< 0.001	0.14	-	-	-
	Species × Temperature	17.72 (4,499)	< 0.001	0.11	2.53 (4,161)	0.04	0.04
	Species × Acclimation	3.23 (2,501)	0.04	0.01	-	-	-
	Temperature × Acclimation	3.31 (2, 501)	0.04	< 0.001	-	-	-
	Species × Temperature × Acclimation	1.40 (4, 500)	0.23	< 0.001	-	-	-
Respiration efficiency – by biotype	Mass	148.54 (1,146)	< 0.001	0.50	2.79 (1,101)	0.09	0.02
	Temperature	1574.74 (2,494)	< 0.001	0.86	22.43 (2,163)	< 0.001	0.21
	Biotype	2.54 (3,115)	0.06	0.03	0.26 (3,87)	0.85	< 0.001
	Acclimation	131.37 (1, 512)	< 0.001	0.19	-	-	-
	Biotype × Temperature	29.86 (6,493)	< 0.001	0.25	2.45 (6,163)	0.03	0.05
	Biotype × Acclimation	3.30 (3, 494)	0.02	0.02	-	-	-
	Temperature × Acclimation	3.71 (2, 492)	0.02	< 0.001	-	-	-
	Biotype × Temperature × Acclimation	1.26 (6, 492)	0.27	< 0.001	-	-	-
Tradeoff – by species (VCO_2)	Total resistance to water loss	-	-	-	17.79 (1,241)	< 0.001	0.02
	Temperature × Total resistance to water loss	-	-	-	0.16 (2,209)	0.85	< 0.001
	Species × Total resistance to water loss	-	-	-	1.84 (2,240)	0.16	< 0.001
Tradeoff – by	Total resistance to water loss	-	-	-	17.24 (1,237)	< 0.001	0.05

biotype (VCO_2)	Temperature × Total resistance to water loss	-	-	-	0.07 (2,210)	0.93	< 0.001
	Biotype × Total resistance to water loss	-	-	-	1.40 (3,227)	0.24	< 0.001

*Results for linear models from the trait value analyses and change of trait analyses for the effects of mass, temperature, species/biotype, and the interaction of species/biotype and temperature on metabolic rate, total resistance to water loss, respiration efficiency, and the trade-off between metabolic rate and total resistance to water loss.

Figure 1 Inset of United States and Canada with the area of interest outlined (A). Area of interest showing distribution, sample size (N), and range of different biotypes across collection sites. M-MI: Marquette, Michigan; L-MI: Livingston, Michigan; WI: Calumet, Wisconsin; KY: Campbell, Kentucky; OH: Delaware, Ohio; VT: Addison, Vermont; W-CT: Windham, Connecticut; L-CT: Litchfield, Connecticut; JJ: *Ambystoma jeffersonianum;* LJJ and LLJ: different *Ambystoma* unisexual biotypes; LL: *Ambystoma laterale.* Pie charts indicate relative numbers of individuals collected at each site (B). Pictures of *Ambystoma jeffersonianum* (C), unisexual polyploid (D), and *Ambystoma laterale* (E). Illustrations provided by Katie Garrett.

Figure 2 Variation in metabolic rate (VCO$_2$) across temperature and biotype. (A) The effects of temperature on log-scaled VCO$_2$ separated by biotype. (B) The effects of temperature on the change in VCO$_2$ separated by biotype. Negative values in acclimation plot (B) show that salamanders acclimated by lowering their VCO$_2$ following the acclimation treatment. Data for all panels are shown as the adjusted mean ± SEM, incorporating the effects of all variables in the model.

Figure 3 Variation in total resistance to water loss (r_T) across temperature and biotype. (A) The effects of temperature on r_T separated by biotype. (B) The effects of temperature on the change in r_T separated by biotype. Negative values in the acclimation plot (B) show that salamanders acclimated by lowering their r_T following the acclimation treatment. Data for all panels are shown as the adjusted mean ± SEM, incorporating the effects of all variables in the model.

Figure 4 Variation in respiration efficiency (WGER) across temperature and biotype. (A) The effects of temperature on log-scaled WGER separated by biotype. (B) The effects of temperature on the change in WGER separated by biotype. Negative values in acclimation plot (B) show that salamanders acclimated by lowering their WGER following the acclimation treatment. Data for all panels are shown as the adjusted mean ± SEM, incorporating the effects of all variables in the model.

Figure 5 Trade-off between the acclimation of metabolic rate (*V*CO$_2$) and change in total resistance to water loss (*r*$_T$) across species (A), temperature (B), and biotype (C). Data for all panels are shown as adjusted mean ± SEM, incorporating the effects of all variables in the model.