Target protein identification in live cells and organisms with a non-diffusive proximity tagging system

Yingjie Sun[1†], Changheng Li[1†], Xiaofei Deng[2], Wenjie Li[1], Xiaoyi Deng[1], Weiqi Ge[1], Miaoyuan Shi[1], Ying Guo[1], Yanxun V Yu[1,3]*, Hai-Bing Zhou[2,4]*, Youngnam N Jin[1,3]*

[1]Department of Neurology, Medical Research Institute, Zhongnan Hospital of Wuhan University, Wuhan University, Wuhan, China; [2]Department of Hematology, Zhongnan Hospital of Wuhan University, School of Pharmaceutical Sciences, Wuhan University, Wuhan, China; [3]Frontier Science Center for Immunology and Metabolism, Wuhan University, Wuhan, China; [4]State Key Laboratory of Virology, Hubei Province Engineering and Technology Research Center for Fluorinated Pharmaceuticals, Key Laboratory of Combinatorial Biosynthesis and Drug Discovery (Wuhan University), Ministry of Education, Frontier Science Center for Immunology and Metabolism, Wuhan University, Wuhan, China

[†]These authors contributed equally to this work.

*For correspondence: yanxunyu@whu.edu.cn (YVY); zhouhb@whu.edu.cn (H-BZ); youngnam_jin@whu.edu.cn (YNJ)

Abstract

Identifying target proteins for bioactive molecules is essential for understanding their mechanisms, developing improved derivatives, and minimizing off-target effects. Despite advances in target identification (target-ID) technologies, significant challenges remain, impeding drug development. Most target-ID methods use cell lysates, but maintaining an intact cellular context is vital for capturing specific drug-protein interactions, such as those with transient protein complexes and membrane-associated proteins. To address these limitations, we developed POST-IT (Pup-On-target for Small molecule Target Identification Technology), a non-diffusive proximity tagging system for live cells, orthogonal to the eukaryotic system. POST-IT utilizes an engineered fusion of proteasomal accessory factor A (PafA) and HaloTag to transfer Pup to proximal proteins upon directly binding to the small molecule. After significant optimization to eliminate self-pupylation and polypupylation, minimize depupylation, and optimize chemical linkers, POST-IT successfully identified known targets and discovered a new binder, SEPHS2, for dasatinib, and VPS37C as a new target for hydroxychloroquine, enhancing our understanding these drugs' mechanisms of action. Furthermore, we demonstrated the application of POST-IT in live zebrafish embryos, highlighting its potential for broad biological research and drug development.

Introduction

The landscape of modern pharmaceutical medicine has significantly evolved with the advent of enhanced drugs, targeted therapies, and combination treatments, yet variability in patient outcomes and toxic responses remain prevalent. This unpredictability stems from a complex interplay of factors, including incomplete target identification, unanticipated off-target effects, intricate polypharmacology, and the influence of individual genetics and environmental factors, all of which collectively contribute to the high failure rate of new drug candidates. This highlights the critical need for accurate identification of protein targets (target-ID). Phenotype-based drug screening offers a promising strategy for discovering novel or superior bioactive molecules, yet precise target-ID within this approach stands as a major barrier to advancing drug development further (Hughes et al., 2021; Vincent et al., 2022). Target-ID is fundamental in unraveling the mechanistic pathways through which drugs exert their therapeutic effects, serving as a critical step in the refinement and development of targeted therapies. By enabling the precise tailoring of drugs to specific biological targets, target-ID significantly enhances drug specificity, efficacy, and safety, while also facilitating the repurposing of existing drugs and advancing our understanding of polypharmacology. Despite its critical importance, accurately identifying biological targets within the complex cellular environment of proteins and other molecules remains a challenge, often hindering progress from drug discovery to therapeutic application.

Significant advancements have been made in target-ID technologies, with each approach offering its own advantages and limitations (Ha et al., 2021). Affinity-based pull-down methods, traditionally employed for target-ID, are hindered by their dependence on strong binding interactions, potentially overlooking proteins that interact transiently or are present at low levels but are crucial for the drug's action. Additionally, affinity-based methods do not adequately preserve the intricate intracellular environment where these interactions occur. As an alternative, photo-affinity labeling (PAL) has emerged, utilizing analogs of bioactive molecules with photo-reactive groups to detect interactions in the natural cellular state. Despite its advantages, PAL has

limitations such as nonspecific labeling, low photoreaction yields, orientation dependency, and in
vivo incompatibility. In response, newer, modification-free approaches like drug affinity
responsive target stability (DARTS) (Lomenick et al., 2009), cellular thermal shift assay
(CETSA) (Molina et al., 2013), and thermal proteome profiling (TPP) (Savitski et al., 2014) have
been developed. These techniques rely on protein stabilization upon binding, offering benefits
like avoiding the need for cumbersome synthesis of derivatives and preserving the cellular
context (Molina et al., 2013; Savitski et al., 2014). However, these methods may have difficulty
detecting subtle or little changes in protein stability, underscoring the ongoing need for continued
development in target identification methodologies.
Proximity labeling (PL) systems such as APEX (Martell et al., 2012)/APEX2 (Lam et al.,
2015), BioID (Roux et al., 2012), and TurboID (Branon et al., 2018) are rapidly emerging as
effective tools for dissecting protein-protein or RNA-protein molecular interactions. While these
methods offer significant advantages in mapping interactomes and networks, their diffusive
labeling presents a limitation for accurate target-ID. Although certain developments such as the
proximity-dependent target-ID system (PROCID) (Kwak et al., 2022) and biotin targeting
chimera (BioTAC) (Tao et al., 2023) using TurboID and miniTurbo, respectively, have shown
promise, the inherent diffusive nature still remains an obstacle. Wells and colleagues developed a
non-diffusive PL system for protein-protein interactions by fusing the E2-conjugating enzyme
Ubc12 for NEDD8, a ubiquitin homolog, with a substrate binding domain/protein (Zhuang et al.,
2013). This system was further adapted for target-ID by combining Ubc12 with a SNAP-tag for
linkage to a specific small molecule, allowing Ubc12 to covalently attach a biotinylated NEDD8
derivative to adjacent target proteins (Hill et al., 2016a). While effective for target-ID, it has
limitations. Exogenous expression of NEDD8 can lead to unintended labeling by the
ubiquitination machinery (Enchev et al., 2015), thus restricting its application in live cells.
Consequently, this system is generally utilized in vitro with cell lysates. While capable of
tagging weakly interacting proteins, the system does not utilize the intact intracellular context,
which is crucial for certain interactions. Additionally, the possibility of NEDD8 polyneddylation
may introduce bias in labeling efficiency and affinity pulldown. Furthermore, the typical
neddylation conditions, occurring at 37 °C for 18 h, may cause protein instability and
precipitation, although adjustments in incubation time or the addition of mild detergents may
alleviate these concerns. Hence, there is a pressing need to develop innovative target-ID methods
that enhance existing techniques.
To overcome existing limitations and establish a streamlined approach for target-ID in live
cells and animal models such as zebrafish, we envisioned a non-diffusive PL system that
leverages a prokaryotic ubiquitin-like protein, orthogonal to the eukaryotic ubiquitin proteasome
system, thereby ensuring specificity and minimizing cross-reactivity. We chose the proteasomal
accessory factor A (PafA) as a ligase and the prokaryotic ubiquitin-like protein (Pup) as its
cognate substrate. Liu et al. had previously demonstrated its effectiveness as a PL system for
studying protein-protein interactions in live cells (Liu et al., 2018). Here, we introduce a target-
ID system, named Pup-On-target for Small molecule Target Identification Technology (POST-
IT). This system integrates PafA with a HaloTag to attach a specific small molecule. After
extensive optimization, we showed that POST-IT successfully identified many known targets
and a novel binder, SEPHS2, for dasatinib in live cells. We then applied POST-IT for
hydroxychloroquine (HCQ) and discovered VPS37C as a novel target, providing mechanistic
insight into HCQ's action. Moreover, we have successfully demonstrated the application of

POST-IT in live zebrafish embryos, highlighting its potential for broad usage in biological
research.
116
117
118 **Results**
119
Design and optimization of POST-IT
To create a non-diffusive PL system suitable for universal and robust target-ID in live cells and
animal models, we integrated PafA from *Corynebacterium glutamicum* (*Cglu*) as a non-diffusive
PL effector with HaloTag as an anchor for small molecules of interest (***Figure 1***). PafA
covalently transfers Pup to proximal prey proteins via direct contact on the ε-amino group of
lysine residues (Özcelik et al., 2012; Watrous et al., 2010), a process known as pupylation. This
approach offers an advantage due to PafA's inherent characteristic of promiscuous, low
selectivity for the amino acid sequence (Liu et al., 2018; Watrous et al., 2010). We selected
HaloTag over alternatives like SNAP-tag due to its superior live-cell labeling efficiency
(Erdmann et al., 2019), readily accessible ligand surface (Kang et al., 2017; Los et al., 2008), and
relatively simple chemical synthesis with a chloroalkane moiety (Los et al., 2008). To simplify
our system for in vivo applications without requiring exogenous biotin, we fused Pup with either
a streptavidin binding peptide (SBP) or a Strep-tag II (STII) as a handle to capture the targets,
rather than using the commonly used biotin carboxyl carrier protein (BCCP). When a target
protein binds to the compound linked to HaloTag, PafA is positioned in close proximity,
allowing for the covalent attachment of Pup to the target's lysine residue. Pupylated targets can
then be captured and identified by mass spectrometry.
To initiate the optimization of POST-IT, we reasoned that using smaller Pup substrates
could minimize interference with the cellular functions of pupylated proteins. It has been shown
that the N terminus of Pup can be removed without compromising its function, generating a short
truncated protein with 28 amino acids from the C terminus (Liu et al., 2018). This truncated Pup
protein will be referred to hereafter as short Pup (sPup). We fused sPup with either an SBP or a
twin-STII (TS) tag at its N terminus (***Figure 1—figure supplement 1A***). HaloTag was
introduced at the N terminus of PafA. In vitro pupylation tests with recombinant proteins showed
that the fusion of HaloTag did not impair PafA activity based on self-pupylation (***Figure 1—***
figure supplement 1B). Notably, both types of sPup substrates exhibited similar pupylation
levels to the full-length Pup. Additionally, PafA retained its robust activity even at 20 °C, a
valuable feature for using this system in other animal models such as zebrafish and *C. elegans*,
which naturally inhabit environments around this temperature. Moreover, we observed robust
self-pupylation within 10 min at 37 °C. However, extensive incubation resulted in the formation
of high molecular weight Halo-PafA, indicating polypupylation of Pup substrates (***Figure 1—***
figure supplement 1C), similar to that observed with ubiquitin.
Polypupylation can lead to unwanted biases in pupylation or pull-down efficiency. To
counter this, we mutated lysine residues to arginine in TS-sPup and SBP-sPup (***Figure 1—figure***
supplement 2A). Since the TS-tag contains two lysines, mutating sPup alone did not affect
polypupylation (***Figure 1—figure supplement 2B and C***). However, further mutations within
the TS-tag completely abolished polypupylation in TS-sPup (***Figure 1—figure supplement 2D***).
Interestingly, Halo-PafA with polypupylation branches showed a distinct migration pattern
compared to multipupylation by TS^{K8R}-sPup^{K61R} at 0.5 h incubation. Similarly, modifying a
lysine residue in the SBP-tag, either through mutation to SBP^{K4R} or by deleting the N-terminal

four amino acids to create sSBP, also completely prevented polypupylation (***Figure 1—figure***
supplement 2E and F). Notably, we observed enhanced levels of multipupylation with lysine-
removed sPup substrates over time (***Figure 1—figure supplement 2D and F***). Together, these
results demonstrate that lysine to arginine (KR) mutation in sPup and modifications to the tags
effectively eliminated polypupylation, while maintaining or enhancing the pupylation efficiency
in a dose- and time-dependent manner.
In addition to polypupylation, we reasoned that the robust self-pupylation activity of Halo-
PafA could diminish the pupylation of target proteins. To address this, we replaced all eight
lysines with arginines in HaloTag, creating Halo^{8KR}-PafA (***Figure 1—figure supplement 3A***).
This mutant exhibited a marked decrease in self-pupylation, similar to PafA alone, indicating
inherent self-pupylation in PafA (***Figure 1—figure supplement 3B***). Previous research reported
that PafA from *M. tuberculosis* (*Mtb*) possesses depupylase activity (Jiang et al., 2018), which
may negatively impact pupylation levels of target proteins. Notably, an *Mtb* PafA mutant with a
S119A mutation showed significantly reduced depupylase activity with minimal changes in
pupylase activity (Jiang et al., 2018). Based on this, we hypothesized that inhibiting depupylation
activity might enhance pupylation. Surprisingly, a PafA mutant with a S126A substitution
(analogous to S119A in *Mtb* PafA) exhibited substantially increased pupylation activity with
reduced polypupylation (***Figure 1—figure supplement 3A and B***).
178
Validation of POST-IT with dasatinib in vitro
To assess the efficiency of our optimized Halo-PafA and Pup in pupylating target proteins for a
small molecule, we synthesized DH1, a HaloTag ligand (HTL) derivative of dasatinib (***Figure***
2A), a well-characterized tyrosine kinase inhibitor used for leukemia treatment (Rix et al., 2007;
Shi et al., 2012). Initially, we evaluated the functionality of Halo^{8KR} in Halo^{8KR}-PafA.
Fluorescence polarization (FP) assay demonstrates the rapid binding of the Halo-AF488 ligand to
Halo^{8KR}-PafA, reaching a plateau in 15-20 min, consistent with a previous report (Los et al.,
2008) (***Figure 2—figure supplement 1A***). A Halo-biotin competition assay confirmed further
that DH1 could be efficiently linked to Halo^{8KR}-PafA within 15 min (***Figure 2—figure***
supplement 1B and C).
After demonstrating the functionalities of Halo^{8KR} and DH1, we tested whether SRC, a
known target of dasatinib, could be labeled by Halo-PafA in the presence of DH1. Surprisingly,
robust pupylation of purified recombinant short SRC occurred exclusively with PafA harboring
Halo^{8KR} at its N terminus in the presence of DH1, whereas PafA with Halo^{8KR} at the C terminus
showed only modest labeling. This result highlights the importance of eliminating self-
pupylation and the N-terminal positioning of HaloTag for effectively labeling the target protein
(***Figure 2B***). We next examined the influence of KR mutations in sPup substrates. Both SBP^{K4R}-
sPup^{K61R} and TS^{K8R}-sPup^{K61R} showed substantial levels of pupylation, especially at lower
pupylation bands, suggesting increased multipupylation (***Figure 2C***). The pupylation of SRC by
Halo^{8KR}-PafA with DH1 diminished with increasing concentrations of dasatinib, affirming the
specificity of this system (***Figure 2D***). Notably, even with high concentration of SRC and
Halo^{8KR}-PafA in the in vitro reaction, no pupylation was detected in the absence of DH1,
indicating that the labeling by our PL system is non-diffusive, specific, and proximity-dependent.
Like SBP^{K4R}-sPup^{K61R}, we observed a significant increase in pupylation by PafA with the N-
terminal, but not the C-terminal, Halo^{8KR} when using TS^{K8R}-sPup^{K61R} (***Figure 2—figure***
supplement 2A) to a degree similar to that of SBP^{K4R}-sPup^{K61R} (***Figure 2—figure supplement***
2B). Although TS^{K8R}-sPup^{K61R} serves as an excellent substrate for PafA without causing
polypupylation, the lysine mutations in the TS-tag led to a loss of binding to Strep-Tactin beads.

Conversely, SBP^{K4R} and sSBP retained their binding to streptavidin, as the core binding sequence
of SBP does not require the 1-10 amino acids at the N-terminus (Barrette-Ng et al., 2013).
Therefore, we decided to use the SBP-tag for further studies.
210
Validation of POST-IT with dasatinib in cellulo
Having successfully verified target protein pupylation by Halo-PafA in vitro, we evaluated our
PL system under cellular conditions to minimize artifacts from non-physiological environments
such as cell lysates. To mimic drug-target interaction-induced pupylation in live cells, we
incorporated the rapamycin-induced interaction between FRB and FKBP into our PL system
(*Figure 3—figure supplement 1A*). Rapamycin treatment led to robust pupylation of FKBP-
EGFP by PafA fused with FRB in a Pup-dose dependent manner (*Figure 3—figure supplement
1B and C*), along with significant self-pupylation independent of rapamycin. Next, we
investigated how KR mutations in SBP-sPup substrates affect pupylation efficiency in live cells.
Both SBP-sPup and SBP^{K4R}-sPup^{K61R} exhibited efficient pupylation (*Figure 3—figure
supplement 1D*), while sSBP-sPup^{K61R} failed to induce self-pupylation as well as pupylation of
FKBP-EGFP, presumably due to its low expression or instability. Consequently, we chose
SBP^{K4R}-sPup^{K61R} for subsequent studies to avoid polypupylation, although SBP-sPup did not
cause any detectable polypupylation in this case. Interestingly, the presence of rapamycin greatly
reduced self-pupylation, suggesting that PafA constantly pupylates either itself or proximal target
proteins. This underscores the importance of minimizing self-pupylation as a critical factor in
developing for a better POST-IT system.
Given these findings, we sought to identify additional self-pupylation sites in PafA through
mass analysis, uncovering six lysine residues susceptible to self-pupylation (*Table 1*). Focusing
on the most frequently pupylated residue, we introduced an additional KR mutation at K172
(*Figure 3—figure supplement 2*), generating Halo^{8KR}-PafA^{S126A,K172R}.
To further refine our PL system, we tested the effect of linker length between Halo^{8KR} and
PafA on SRC labeling in the presence of DH1. We observed enhanced labeling with a linker
longer than 10 amino acids (*Figure 3A*), prompting us to select an 18-amino acid linker for
further studies. We then confirmed that the S126A and K172R mutations additively enhanced
pupylation activity under cellular conditions (*Figure 3B*). This optimized version of Halo-PafA,
Halo^{8KR}-PafA^{S126A,K172R}, in combination with a polypupation-free version of sPup will be referred
to as POST-IT. Unlike wild type Halo-PafA, which displayed negligible labeling under cellular
conditions, POST-IT exhibited significant labeling activity on both exogenous and endogenous
SRC in the presence of DH1, generating POST-IT^{DH1} (*Figure 3C-E*). Labeling intensity of short
SRC increased with higher DH1 concentrations and was competitively inhibited by increasing
concentrations of dasatinib (*Figure 3F and G*), demonstrating the specificity of POST-IT^{DH1} in
target labeling.
In our quest for a superior PafA variant, we investigated the potential of *Mtb* PafA as a PL
system. However, we detected no labeling of short SRC by *Mtb* PafA, irrespective of the
positioning of Halo^{8KR} at the N- or C-terminus (*Figure 3—figure supplement 3*). In contrast,
Halo^{8KR}-PafA (*Cglu*) exhibited significant pupylation with sPup^{K61R} substrates from both *Cglu*
and *Mtb*, suggesting a lack of orthogonality among different prokaryotic PafA systems.
249
Optimization of the linker of dasatinib-HTL (DH) derivatives
Encouraged by the successful labeling of the target protein by POST-IT^{DH1} in live cells, we next
explored its ability to identify known targets and potentially reveal novel binding proteins of

dasatinib. To this end, we investigated whether the linker length or structure between dasatinib
and the Halo chloroalkane moiety could influence labeling efficiency, as the linker's
characteristics in biomolecules, including PROTAC (Békés et al., 2022), significantly impact
their functionalities. Based on a previous report highlighting the enhanced performance of HTLs
containing carbamate linkers (Friedman Ohana et al., 2015), we synthesized four additional DH
derivatives (DH2-DH5) with varying linker lengths longer than that in DH1 (***Figure 4—figure***
supplement 1). These new DH derivatives with carbamate linkers displayed significantly
enhanced covalent binding to recombinant Halo^{8KR}-PafA in vitro and to transiently expressed
Halo^{8KR}-PafA in intact cells, as measured by FP assays, compared to DH1 (***Figure 4A and B***).
Importantly, the new DH derivatives also showed more robust pupylation of short SRC in vitro
and endogenous SRC under cellular conditions (***Figure 4C and D***). DH5, containing the longest
linker, was selected for further experiments due to its excellent efficiency. We then verified that
POST-IT could efficiently pull down endogenous SRC in the presence of DH5, an effect
attenuated by an excess of dasatinib (***Figure 4E***). Furthermore, we measured the inhibitory effect
of DH5 on SRC kinase activity and confirmed that DH5 retains comparable activity to dasatinib
(***Figure 4—figure supplement 2***).
269
Identification of target proteins for dasatinib by POST-IT
With our optimized POST-IT system and DH5, POST-IT^{DH5}, in hand, we conducted labeling
experiments in live cells with or without an excess of dasatinib. After pulling down cell lysates
with streptavidin beads, we analyzed biotin-eluted proteins via mass spectrometry. Remarkably,
we identified 8 known target proteins (FYN, RIPK2, MAPK14, ABL2, ABL1, CSK, LYN, and
SRC) with significant enrichment and 2 known target proteins (GAK and SIK2) with lower
confidence (***Figure 5A***). Among these ten known targets, five are tyrosine kinases (FYN,
ABL1/2, CSK, LYN, and SRC) and four are serine/threonine kinases (RIPK2, MAPK14, GAK,
and SIK2). These results validate the effectiveness of our POST-IT system as a target-ID
methodology and underscore its potential superiority to other PL systems for target-ID (Hill et
al., 2016b; Kwak et al., 2022).
Additionally, we identified selenophosphate synthetase 2 (SEPHS2) as a novel protein target
that binds to dasatinib. SEPHS2, a crucial enzyme for selenoprotein synthesis, exhibits ATP-
binding and kinase activities, supporting its potential interaction with dasatinib. To confirm
SEPHS2 as a genuine target of dasatinib, we purified the recombinant SEPHS2 protein and
measured its binding affinity to DH5 using microscale thermophoresis (MST) analysis (***Figure***
5B and C). This analysis yielded a dissociation constant (K_d) of 11.6 ± 5.7 μM for DH5 and 9.1
± 0.8 μM for dasatinib. To elucidate the binding mode of dasatinib to SEPHS2, we performed
molecular docking analysis to model their interaction. A previous study using molecular
dynamics simulations identified Sec60, Lys63, and Gly319 as active sites and 15 amino acids as
key residues or HUB nodes (Nunziata et al., 2019). Remarkably, the docking analysis revealed a
favorable binding mode for dasatinib within the active sites of SEPHS2, with a Vina score of -
9.9 (***Figure 5D***). In the binding pocket of SEPHS2, dasatinib engages in multiple interactions,
including pi-cation or cation-π interaction with Lys63 and Van der Waals interaction with
Gly319 in the active sites, along with hydrophobic interactions with Phe130 and Phe140, Van
der Waals interaction with Phe320, and a hydrogen bond with His325 among the HUB nodes
(***Figure 5E***). Collectively, these results suggest that SEPHS2 is a novel protein target for
dasatinib, indicating that dasatinib may impact SEPHS2 function.
298

Identification of target proteins for chloroquine by POST-IT

To further demonstrate the effectiveness of the POST-IT system in target-ID, we applied it to identify target proteins for hydrochloroquine (HCQ), using it as an additional model ligand. HCQ has a long history of clinical use for treating malaria, rheumatoid arthritis, and lupus. Moreover, it has been widely used in research as an autophagy inhibitor, a property gaining interest in the development of cancer therapy (Jain et al., 2023). Despite its broad application, the precise mechanisms through which HCQ inhibits autophagy and the pathways through which HCQ or chloroquine (CQ) elicit adverse effects such as psychiatric symptoms, retinal and ototoxicity, cardiac toxicity, and even death (Muller, 2021; Nirk et al., 2020) remain to be fully understood.

First, we synthesized a dimer form, DC661-H1, of a CQ HTL derivative with the same linker used for DH5 (*Figure 6—figure supplement 1A*), assuming that a dimer would enhance binding affinity as previously described (Rebecca et al., 2019). We confirmed that DC661-H1 and DC660, an analog of DC661-H1 without HTL, induced robust LC3-II bands in two mammalian cell lines to a similar extent as HCQ, indicating that they function as potent autophagy inhibitors (*Figure 6—figure supplement 1B*). To test the versatility of POST-IT as a tool for mass analysis, POSTT-IT was coupled with stable isotope labeling with amino acid in cell culture (SILAC) for target-ID. Heavy-Lys/Arg-labeled HEK293T cells expressing POST-IT were treated with DC661-H1 and DMSO, while light-Lys/Arg-labeled HEK293T cells expressing POST-IT were treated with DC661-H1 and an excess of competitive DC660, as described in *Figure 6—figure supplement 2*. Intriguingly, POST-IT identified several target proteins directly linked to autophagy regulation (*Figure 6A*). Hence, we aimed to verify whether DC660 and HCQ bind to these autophagy-related proteins. Among them, we purified five recombinant proteins, including TOM1, TOM1L2, TRAPPC3, VPS29, and VPS37C, and measured their binding properties using MST analysis. TOM1, TOM1L2, and TRAPPC3 appeared to bind well to DC660 but failed to acquire K_d values due to abnormal thermophoretic movement, suggesting non-homogeneous aggregation (*Figure 6—figure supplement 3A-C*). VPS29 showed dose-dependent binding to DC660, yielding a K_d of 24.95 ± 11.71 μM (*Figure 6—figure supplement 3D*). However, binding to HCQ was negligible in these proteins (*Figure 6—figure supplement 3E-H*). These results support the validity of our POST-IT as a PL system for live cell target-ID, as all tested proteins showed binding ability to DC660.

Next, we investigated the binding affinity of VPS37C to DC660 and HCQ using purified recombinant protein. MST analysis revealed a very high binding affinity of VPS37C to DC660 with a K_d value of 21.5 ± 9.8 nM (*Figure 6B*). Notably, VPS37C also exhibited moderate binding affinity to HCQ (*Figure 6C*). This result prompted us to further explore the binding ability of VPS37C under cellular conditions. First, we performed POST-IT labeling in HEK293T cells expressing VPS37C with a V5-tag, followed by pulldown using streptavidin beads. VPS37C was substantially enriched in the presence of DC661-H1, but its binding was almost completely abolished by competitive DC660 or HCQ (*Figure 6D and E*). Furthermore, through a CETSA experiment, we observed that VPS37C became more stable in the presence of DC660 or HCQ than in the DMSO condition, upon exposure to increasing temperatures from cell lysates (*Figure 6F and G*). Collectively, these results provide strong evidence that VPS37C binds to DC661 (or DC660) as well as HCQ in vitro and in live cells, suggesting a potential role of VPS37C in the action mechanism of CQ or HCQ in inhibiting autophagy.

In vivo application of POST-IT as a target-ID system

Phenotype-based screening is gaining increased recognition as a powerful strategy for
discovering novel compounds, superior drugs, or uncovering unknown biological mechanisms,
such as the non-canonical translation discovered in our previous study using this approach (Jin et
al., 2018). Zebrafish have emerged as an excellent in vivo whole-animal model system for small
molecule screening and drug discovery. They are vertebrates, highly prolific, and amenable to
high-throughput chemical screens, often leading to the serendipitous discovery of novel
compounds (Patton et al., 2021). However, target-ID remains a major hurdle in understanding
the mechanisms of action of novel compounds, thereby delaying the advancements in drug
development. Therefore, we set out to test the applicability of POST-IT as an efficient method
for target-ID in zebrafish.
To this end, we first generated a robust, ubiquitous expression plasmid for POST-IT in
zebrafish using an optimized QF-binary system (Burgess et al., 2020). This system combines a
DNA binding domain from the QF transactivator fused with an activation domain from the Gal4
transactivator, creating QFGal4, which binds to five repeats of QUAS, leading to gene
activation. POST-IT, tagged with a 3×FLAG, is inserted following a 2A ribosome-skipping
sequence, allowing for the simultaneous expression of QFGal4 and POST-IT under the control of
a ubiquitous promoter, *ubb* (***Figure 7A***). We chose the *ubb* promoter over tissue-specific
promoters to more accurately evaluate the levels and potential toxicity of transgene expression.
The expression of SBP^{K4R}-sPup^{K61R} is driven by a 5×QUAS promoter, and EGFP is incorporated
upstream of a 2A sequence as an expression marker. All these components are assembled in a
single plasmid to simplify transgenesis. Embryos injected with this plasmid displayed robust
EGFP expression throughout their entire bodies without observable toxicity, indicating the
suitability of the POST-IT system in zebrafish (***Figure 7B***). Note that the expression pattern of
EGFP is non-homogeneous due to the mosaic expression of the injected plasmid. After treating
live embryos with DH5 or DMSO, we observed marked enrichment of endogenous SRC in
embryos treated with DH5 but not in those treated with DMSO, although expression levels of
POST-IT were slightly lower in the DH5 group due to injection variability (***Figure 7C***).
Together, these results demonstrated that POST-IT can be effectively utilized as a transgene in
zebrafish without toxicity and applied for in vivo target-ID.
374
375
376 **Discussion**
377
In this study, we present POST-IT (Pup-On-target for Small molecule Target Identification
Technology), a new non-diffusive PL system for target-ID in live cells and in vivo animal
models. We systematically employed mutagenesis to eliminate self-pupylation, polypupylation,
and depupylase activity. Moreover, we optimized the linker length between HaloTag and PafA
and developed PEG linkers for HTL derivatives. POST-IT provides a straightforward, yet highly
robust and reliable method for target-ID within intact cellular context, a feature that significantly
reduces artifacts from cell lysates and may be crucial for certain ligand/drug-protein interactions.
We further demonstrated the applicability of POST-IT in an animal model, zebrafish. POST-IT
does not cause noticeable toxicity and allows for tissue-specific and temporally controlled
expression using a variety of transgenic zebrafish systems. We believe that POST-IT can be
readily applied to other organisms, from *C. elegans* to mice, while retaining its effectiveness.
This is evidenced by the robust activity of Halo-PafA, which remains active even at temperatures
as low as 20 °C (***Figure 1—figure supplement 1B***).

While TurboID is a widely used tool for studying protein networks, it suffers from persistent high background labeling, even without biotin. This necessitates meticulous experimental adjustments, such as employing dialyzed FBS and fine-tuning biotin labeling levels (May et al., 2020). Alternative strategies, such as light-regulation (Lee et al., 2023) and split-TurboID (Cho et al., 2020), have been introduced to mitigate high background labeling, yet their suitability for target-ID and in vivo applications remains to be explored. Additionally, previous studies have reported cellular toxicity associated with TurboID (Branon et al., 2018; May et al., 2020), although recent work shows no obvious toxicity in zebrafish (Xiong et al., 2021). Furthermore, the labeling radius of TurboID can exceed 35 nm (May et al., 2020), complicating the precise identification of target proteins. While these PL systems can provide valuable insights into a drug's protein network (Kwak et al., 2022; Tao et al., 2023), direct target-ID is essential for understanding the drug's mechanism of action.

A recent study introduced BmTyr, a PL system based on *Bacillus megaterium* tyrosinase (Zhu et al., 2024), which oxidizes phenol or catechol derivatives into *o*-quinones for labeling adjacent proteins. This method provides fast and lower-background labeling compared to TurboID. Unlike APEX, BmTyr does not require hydrogen peroxide, enhancing its biocompatibility for in vivo use. However, the need for copper might introduce toxicity and limitations. Oxygen dependency and glutathione reactivity may also affect its performance. While in vivo application was demonstrated by injecting purified BmTyr into mouse brains, the feasibility of its transgenic expression and broader application remains untested. Importantly, the diffusive nature of BmTyr's labeling might challenge its use for precise target-ID, despite being an innovative alternative PL toolkit.

Using our POST-IT system, we identified SEPHS2 and VPS37C as novel target proteins for dasatinib and HCQ/CQ, respectively. SEPHS2 detoxifies selenide, a toxic intermediate of selenocysteine biosynthesis, by converting it with ATP into selenophosphate, an essential component for selenoprotein production. This function is particularly important in cancer cells that require enhanced defense against oxidative stress. These cancer cells frequently exhibit elevated SLC7A11/xCT expression, leading to increased selenide import and a heightened demand for selenide detoxification. Similarly, abnormal upregulation of selenium uptake and selenocysteine biosynthesis occurs in many cancer cells, including breast cancers (Carlisle et al., 2020) and acute myeloid leukemia (Eagle et al., 2022), making SEPHS2 crucial for their survival but not for normal cells. Therefore, SEPHS2 is a promising target for cancer therapy. Given that SEPHS2 has been identified as a target for dasatinib, potentially binding to the active site, investigating whether SEPHS2 inhibition by dasatinib contributes to its anticancer effects is warranted. Binding information of dasatinib to SEPHS2 could facilitate the development of new SEPHS2 inhibitors.

Despite decades of research using HCQ/CQ as an autophagy inhibitor, its exact mechanism of action remains elusive. Although it has been widely assumed that HCQ/CQ increases lysosomal pH, Mauthe et al. demonstrated that its primary effect is to inhibit the fusion between autophagosomes and lysosomes (Mauthe et al., 2018). Additionally, HCQ/CQ induces disorganization in the Golgi and endosomal systems independently of autophagy. These findings highlight the significance of pH-independent mechanisms in HCQ/CQ's action. Furthermore, components of the Endosomal Sorting Complex Required for Transport (ESCRT) complexes have been shown to regulate the final stages of autophagosome formation (Javed et al., 2023; Takahashi et al., 2019, 2018; Zhen et al., 2020). ESCRT-I consists of three core components (TSG101/VPS23, VPS28, and one of VPS37A/B/C/D) and a single auxiliary protein among

UBAP1, MVB12A, and MVB12B. While VPS37A has been identified as a key player in phagophore closure (Javed et al., 2023; Takahashi et al., 2019), the roles of other VPS37 proteins remain unexplored. Considering the structural similarities among the VPS37 proteins (***Figure 6—figure supplement 4***), it is plausible that HCQ/CQ could interact with multiple members of this family to some extent, potentially inhibiting their role in autophagosome maturation. This suggests two potential mechanisms for HCQ/CQ-mediated autophagy inhibition. One involves interfering with the function of VPS37 proteins in autophagosome closure. The other suggests that HCQ/CQ is actively transported into lysosomes by ESCRT-I, leading to lysosomal alkalization. Further research is crucial to elucidate these mechanisms. Understanding these pathways will not only illuminate therapeutic strategies for HCQ/CQ but also guide the development of superior derivatives with minimized side effects.

There remain some limitations in utilizing POST-IT system. One limitation is the requirement to synthesize an HTL derivative for the compound of interest, a challenge common to all target-ID methods except for those that are modification-free. To facilitate derivatization, we provide guidelines for linkers incorporating carbamate and PEG, as shown in ***Figure 4*** and ***Figure 4—figure supplement 1***. PEG linkers are advantageous for their increased solubility and cell permeability. Provided the derivatives maintain comparable binding affinity to the target protein or preserve the compound's activity, we believe that POST-IT can be effectively applied from in vitro conditions to live cells and organisms. Secondly, POST-IT may require significantly longer incubation times in live cells compared to in vitro reactions, which could be attributed to several factors, such as insufficient expression of POST-IT or Pup, or the instability or metabolism of an HTL derivative of the drug. Further improvement of POST-IT through directed evolution, rational protein engineering, and structure-aided design is warranted. Given that many drugs function in specific cellular locations, such as the cell surface and mitochondria, targeting POST-IT to precise subcellular locations could enhance the likelihood of identifying true biological target proteins rather than off-target binding partners. Therefore, exploring the functionalities of POST-IT in specific subcellular contexts is an essential direction for future research. Additionally, in this study, we utilized HEK293T cells for proof-of-concept. However, investigating protein targets in diverse cell types, particularly cancer cells, may reveal a broader spectrum of targets. To gain a more comprehensive understanding of a drug's target profile, we recommend that users employ POST-IT in various cell lines or in vivo models, such as zebrafish.

In conclusion, POST-IT represents a significant advancement in the field of target identification, offering an innovative, non-diffusive proximity tagging system designed for precise target-ID within live cells and in vivo models. By preserving the cellular context, POST-IT overcomes the limitations associated with traditional cell lysate methods and offers superior specificity compared to existing PL systems like TurboID. We successfully identified SEPHS2 as a target for dasatinib and VPS37C as a target for HCQ, exemplifying its effectiveness. Its application in both live cells and zebrafish embryos underscores its versatility and potential to provide valuable insights into the molecular mechanisms of drug action, thereby enhancing the scope of biomedical research and therapeutic development.

Materials and methods

Materials

General reagents and solvents including NaCl, KCl, EDTA·Na$_2$, KH$_2$PO$_4$, Na$_2$HPO$_4$·12H$_2$O,
NaF, NH$_4$OH, NaOH, urea, chloroform, glycerol, methanol, ethanol, isopropanol, and others,
were obtained from Sinopharm Chemical Reagent, China. Regents related to protein expression,
such as Isopropyl β-D-1-thiogalactopyranoside (IPTG), Tricine, sodium dodecyl sulfate (SDS),
acrylamide, N,N′-methylene-bis-acrylamide, tetramethylethylenediamine (TEMED), ammonium
persulfate, phenylmethylsulfonyl fluoride (PMSF), dithiothreitol (DTT), imidazole, Tris-HCl,
glycine, ATP·Na$_2$, and biotin, were obtained from aladdin or Biofroxx, China. Triton X-100 and
Tween-20 were from Sangon Biotech, China, and NP-40 was from Biosharp, China. Ni
Sepharose 6 Fast Flow (17531801) was from Cytiva, USA, glutathione resin (SA008010) was
from Smart-Lifesciences, China, and Sephadex-G100 (BS210) was from Biosharp, China.
Amicon Ultra Centrifugal Filters (UFC9050, UFC9030, UFC9010) were purchased from
Millipore, USA. For molecular cloning, 2×MultiF Seamless Assembly Mix (RK21020) was from
ABclonal, China, and Phanta Max Super-Fidelity DNA Polymerase (P505-d1) and HiScript III
1st strand cDNA Synthesis Kit (R312) were from Vazyme, China. All restriction enzymes were
from New England Biolabs, USA. HaloTag Alexa Fluor 488 ligand (G1001) and HaloTag Biotin
ligand (G8281) were from Promega, USA. Dasatinib (S45672-25MG) was purchased from
Shanghai yuanye Bio-Technology, China. Hydrochloroquine (HCQ) (T9287) was obtained from
TargetMol Chemicals, USA. BCA assay (P0011) was from Beyotime, China. Clarity Western
ECL (1705060) was purchased from Bio-Rad, USA. Antibodies against V5-tag (AE017), Flag-
tag (AE092), HA-tag (AE008), Myc-tag (AE070), and β-actin (AC026), as well as secondary
antibodies, were purchased from ABclonal, China. Anti-SRC (36D10) (2109) and Streptavidin-
HRP (3999) were from Cell Signaling Technology, USA. Streptavidin magnetic beads (L-1012)
was purchased from BioLinkedin, China. Information on all other materials is provided in the
methods section as they appear.
506
Construction of plasmids
Plasmids were typically constructed by the HiFi DNA assembly method using 2×MultiF
Seamless Assembly Mix. PCR amplification was carried out with a high-fidelity DNA
polymerase, Phanta Max Super-Fidelity DNA Polymerase. In general, the pTol2-EGFP-2ApA-
CK2 or pTol2-EGFPpA vector, developed in our lab (Jin et al., 2018), was used for mammalian
expression, and the pET28a vector was utilized for recombinant protein expression in *E. coli*,
unless specified otherwise. All DNA constructs were validated by Sanger sequencing.
To construct Halo-PafA plasmids, HaloTag and PafA were PCR amplified using HaloTag7
(a gift from Yang Yu, Chinese Academy of Sciences) and pEF6a-kozak-CD28-PafA-myc (a gift
from Min Zhuang, ShanghaiTech University), respectively, and then subcloned into the NotI and
EcoRI sites of pTol2-EGFP-2ApA-CK2, or into the EcoRI site of pET28a. HaloTag^{8KR}, TS-
sPup, and SBP-sPup DNA fragments were synthesized by Tsingke Biotechnology (Wuhan).
Halo^{8KR}-PafA was subcloned into the NotI and EcoRI sites of pTol2-EGFP-pA or into the
BamHI and NotI sites of pET28a. TS-sPup and SBP-sPup were subcloned into the KpnI and
NotI sites of pEF6a-HB-PUP (a gift from Min Zhuang) for mammalian expression and inserted
into the EcoRI site of pET28a for *E. coli* expression. Point mutations, including SBP^{K4R}, TS^{K8R},
sPup^{K61R}, PafA^{S126A}, and PafA^{K172R}, were introduced using a modified site-directed mutagenesis
(Liu and Naismith, 2008). For recombinant protein expression of PafA in *E. coli*, PCR-amplified
PafA was fused to a GST-tag into the BamHI and EcoRI sites of pGEX-6P-2, generating pGEX-
6P-2-PafA.

Target proteins, such as SRC, SEPHS2, VPS37C, TOM1, TOM1L2, TRAPPC3, and VPS29, were PCR amplified using a cDNA library synthesized with the HiScript III 1st strand cDNA Synthesis Kit from total RNA extracted from HEK293T cells with RNAiso Plus (9109, Takara Bio, Japan). A catalytic active, truncated version of SRC spanning amino acids 247-536, SRC(247-536), was inserted into the NotI and EcoRI sites of pTol2-EGFP-2ApA-CK2, and into the BamHI and NotI sites of pET28a. SEPHS2-2×V5 was subcloned into the BamHI and EcoRI sites of pE28a-N-SBPII, engineered by inserting a SBPII-tag into pET28a vector in our lab for *E. coli* expression. The selenocysteine (Sec60) in SEPHS2 was mutated to cysteine for expression in *E. coli*. VPS37C-2×V5 was subcloned into the NotI and MluI sites of pHAGE-IRES-puro (a gift from Lingling Chen, ShanghaiTech University) for mammalian expression, and the codon-optimized VPS37C, coVPS37C, for *E. coli* was inserted into the EcoRI and SalI sites of pE28a-N-SBPII for *E. coli* expression. TOM1, TOM1L2, TRAPPC3, and VPS29 were subcloned into the EcoRI site of pET28a.

FKBP and FRB were synthesized by Tsingke Biotechnology (Wuhan). FKBP, tagged with a 3×HA, was inserted into the NotI and EcoRI sites of pTol2-EGFPpA, generating pTol2-3×HA-FKBP-EGFPpA. FRB, tagged a 3×V5, and mKate2 were assembled into the NotI and XmaI sites of pTol2-Myc-Halo-PafA-2ApA-CK2, resulting in pTol2-3×V5-FRB-mKate2-PafA-pA. To make a stable cell line, CMV-HA-Halo^{8KR}-PafA^{S126A,K172R}, SBP^{K4R}-sPup^{K61R}, TRE3G (a Tet-On promoter), Tet-On 3G transactivator, IRES-EGFP, and Puro were PCR amplified and assembled into pPB-mU6pro (a gift from Zhou Yan, Wuhan University), generating PB-puro-CMV-POST-IT-Tet-SBP^{K4R}-sPup^{K61R}-iresGFP. The expression of SBP^{K4R}-sPup^{K61R} was induced by the application of doxycycline via the Tet-On promoter. The P2A, a 2A ribosome-skipping sequence, was placed between HA-Halo^{8KR}-PafA^{S126A,K172R}, the Tet-On 3G transactivator, and Puro^{R} to allow for simultaneous expression of the three genes. For the expression of POST-IT in zebrafish, QFGal4, 3×Flag-Halo^{8KR}-PafA^{S126A,K172R}, 5×*QUAS*, EGFP, and SBP^{K4R}-sPup^{K61R}, were PCR amplified and assembled into the ClaI site of pUbi-QF-CK, which was developed in our lab for the Tol2 transposon system, resulting in pUbi-QFGal4-2A-3×Flag-POST-IT. P2A was inserted between QFGal4 and 3×Flag-Halo^{8KR}-PafA^{S126A,K172R}, and between EGFP and SBP^{K4R}-sPup^{K61R}.

Recombinant protein expression and purification
E. coli BL21(DE3) cells were transformed with a plasmid for expression of His-tagged or GST-tagged protein, and cultured in LB medium containing 100 µg/ml of either ampicillin or kanamycin overnight at 37 °C until the OD600 reached 0.6-0.7. Protein expression was induced by adding 0.1-0.5 mM IPTG and incubating at 16 °C for 16-20 h. The bacterial culture was then centrifuged at 6,000g for 15 min at 4 °C. The pellet was resuspended in *E. coli*-lysis buffer (50 mM Tris-HCl pH 8.0, 300 mM NaCl, 10 mM imidazole, 5% glycerol, 1 mM PMSF), lysed by either sonication or a high-pressure homogenizer (AH-1500, ATS Engineering, China), and subsequently centrifuged at 24,000g at 4 °C for 30 min. PMSF stock solution (200 mM in isopropanol) was freshly added to the appropriate buffer to achieve the final concentration.

For Ni column purification via a 6×His-tag, the cleared supernatant was loaded onto Ni Sepharose 6 Fast Flow using a gravity flow. After extensive washing with more than 20 column volume (CV) of Ni-wash buffer (50 mM Tris-HCl pH 8.0, 500 mM NaCl, 10% glycerol, 0.1% Triton X-100, 10 mM imidazole, 1 mM PMSF), the recombinant protein was eluted with 2 CV of elution buffer 1-4 (50 mM Tris-HCl pH 8.0, 200 mM NaCl, varying concentrations of imidazole (250, 300, 350, 400 mM for elution buffer 1-4), 1 mM PMSF). Each fraction's purity

was assessed by SDS-PAGE, followed by staining with R-250 Coomassie Brilliant Blue (CBB).
The cleaner fractions were collected, concentrated, and stored in storage buffer (50 mM Tris-HCl
pH 8.0, 150 mM NaCl, 1 mM DTT, 10% glycerol) through buffer exchange using Amicon Ultra
Centrifugal Filter. All protein purification was performed via Ni column purification except for
the following cases.
For GST purification via a GST-tag, the cleared supernatant was loaded onto glutathione
resin and allowed to bind at 4 °C for 1 h. The resin was washed with more than 20 CV of Ni-
wash buffer, with 10 CV of PreScission Protease reaction buffer (50 mM Tris-HCl pH 8.0, 150
mM NaCl, 2 mM EDTA, 1 mM DTT). 50 U of PreScission Protease (SLP00501, Smart-
Lifesciences, China) was added and incubated at 4 °C overnight. The cleaved protein was
collected, concentrated, and stored in storage buffer through buffer exchange using Amicon Ultra
Centrifugal Filter.
For coVPS37C purification, a protocol using a mutant streptavidin, SAVSBPM18, was
applied with minor modifications as previously shown (Wu et al., 2019). The synthesized
SAVSBPM18 and DBD (dextran binding domain) were subcloned into the NcoI and XhoI sites
of pET28a, creating pET-28a-SAVSBPM18-XTEN-DBD, which was then transformed into *E.*
coli BL21(DE3) cells. The coVPS37C plasmid was transformed into the *E. coli* ArcticExpress
(DE3) strain. The *E. coli* cells were cultured at 37 °C until the OD600 reached 0.6, at which
point protein expression was induced by adding 0.2 mM IPTG. The *E. coli* cells were
subsequently cultured at 12 °C overnight and processed to get cleared cell lysates as described
above. *E. coli* cell lysates expressing SAVSBPM18-XTEN-DBD were prepared and mixed with
coVPS37C lysates at a 1:4 ratio. After 1 h of incubation at 4 °C, the mixture was loaded onto Ni
Sepharose 6 Fast Flow resin, washed with Ni-wash buffer, and eluted with elution buffer 2. The
eluate was then mixed with Sephadex-G100 for 1 h at 4 °C, washed with Ni-wash buffer, and
eluted with SBP elution buffer (10 mM biotin, 50 mM Tris-HCl pH 8.0, 300 mM NaCl). The
final eluate was collected, concentrated, and stored in storage buffer through buffer exchange
using Amicon Ultra Centrifugal Filter. All purified proteins were stored at -80 °C until use.
600
Cell culture and transfection
HEK293T cells (CRL-3216, ATCC, USA) were confirmed to be Mycoplasma-negative and
cultured in Dulbecco's modified Eagle's medium (DMEM) (SH30243.01, Cytiva, USA),
containing 4.5 g/L glucose, 4 mM glutamine, and 1 mM sodium pyruvate, supplemented with
10% FBS (SA211.02, Cellmax, China), 100 U/ml penicillin, and 100 µg/ml streptomycin at 37
°C with 5% CO_2. Transient transfections were performed using LipoJet (SL100468, SignaGen,
USA), following the manufacturer's instruction.
608
Fluorescence polarization assay
The fluorescence polarization (FP) assay was conducted using a SpectraMax i3x microplate
reader (Molecular Devices, USA) equipped with a fluorescence polarization cartridge (Ex. 485
nm, Em. 535 nm) and a grating (G) factor set to 1.5. Black opaque 96-well micro plates
(Beyotime, China) were used for all experiments. For the in vitro time course measurement, 30
nM of Halo^{8KR}-PafA protein and 2 nM of HaloTag Alexa Fluor 488, Halo-AF488, were prepared
in PBS buffer, and the parallel (Iv) and perpendicular (Ih) emission intensities were measured in
kinetic mode at 20-s intervals for 30 min. The millipolarization units (mP) were calculated using
the formula mP = $1000 \times [(Iv - G \times Ih) / (Iv + G \times Ih)]$. In the competition assay with DH
derivates, 15 nM of Myc-Halo-PafA and 1 nM of Halo-AF488 were incubated with various DH

derivatives in serial dilution at 37 °C for 30 min before FP measurement, with Iv and Ih
measured at the endpoint. For the cellular binding assay, HEK293T cells were co-transfected
with 4 µg of pTol2-myc-Halo-PafA-2ApA-CK2 per well in a 6-well plate and incubated for 48
h. Different DH derivatives at 1 µM or DMSO were added to each well and incubated for 3 h.
Cell lysates were prepared in PBST (PBS with 0.5% Triton X-100), supplemented with a
protease inhibitor cocktail (MB2678, MeilunBio, China), through brief sonication followed by
centrifugation. Protein concentration was determined by the BCA assay. Cleared cell lysates (5
µg/ul) were incubated with 2 nM Halo-AF488 in a 60 µl volume at 37 °C for 30 min. Iv and Ih
were measured, and mP was calculated.
628
Western blot analysis
Cells transfected and/or treated with different compounds were harvested at the end of treatment.
Cells were washed with ice-cold PBS and lysed with cell-lysis buffer that included RIPA buffer
(50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% Triton X-100, 1 mM EDTA, 5% glycerol),
supplemented with 20 mM NaF, 2 mM Na_3VO_4, 0.4% SDS, 0.2% sodium deoxycholate (DOC),
and a protease inhibitor cocktail. Following brief sonication, cell lysates were centrifuged at
maximum speed for 10 min at 4 °C. The supernatant was collected, and the protein concentration
was determined using the BCA assay. Samples were diluted to the same concentration, mixed
with SDS sample buffer for SDS-PAGE, and boiled for 5-10 min. Proteins were separated by 6%
or 8% SDS–PAGE and transferred to polyvinylidene fluoride (PVDF) membrane. The
membrane was blocked with 5% skim milk in Tris-buffered saline containing 0.05% Tween-20
(TBST) and incubated overnight at 4 °C with the specific antibodies diluted in TBST containing
2% BSA: 1:5,000 anti-V5, 1:2,500 anti-HA, 1:5,000 anti-Myc, 1:5,000 anti-SRC, 1:10,000 anti-
β-Actin, 1:5,000 Streptavidin-HRP. After three to four TBST washes, the membrane was
incubated with a 1:10,000 HRP-conjugated secondary antibody against rabbit or mouse in TBST
containing 5% skim milk for 1 h at room temperature. Following additional washes, the
membranes were developed using chemiluminescence with Clarity Western ECL.
646
In vitro pupylation assay
For the self-pupylation assay, the reaction consisted of 1 µM Halo-PafA (or a mutant protein)
and 10 µM of a Pup substrate in 20 µl of pupylation buffer (PBS supplemented with 15 mM
$MgCl_2$ and 10 mM ATP), incubated at 37 °C for durations ranging from 10 to 180 min as
specified in each figure, and stopped by the addition of 2× SDS sample buffer. Proteins were
separated in three layers (4, 6, and 10%) Tricine-SDS-PAGE and stained with CBB. To test the
effect of different DH derivatives on the pupylation of the target, 1 µM of a Halo-PafA
derivative, 0.5 µM SRC(247-536)-2×V5, and 500 nM of a DH derivative were mixed in the
pupylation buffer and incubated at room temperature for 30 min. Ten µM of a Pup substrate was
then added and the mixture was incubated at 37 °C for another 30 min. The reaction was stopped
by adding 2× SDS sample buffer, and the western blot assay was performed using an anti-V5
antibody. In the competition assay, a serial dilution of dasatinib from 0.2 µM to 10 µM was
added along with 500 nM DH1 to the reaction before the addition of the Pup substrate. For the
identification of pupylated lysine residues in PafA, 2.5 µM PafA and 20 µM HA-Pup were
mixed in 50 µl of the pupylation buffer at 37 °C for 2 h. The reaction was stopped by adding 2×
SDS sample buffer and separated by SDS-PAGE. CBB-stained bands were excised and analyzed
by LC-MS/MS.
664

In cellulo pupylation assay

For the western blot assay, HEK293T cells were plated on 6-well plates and co-transfected as described below. For rapamycin-induced pupylation, HEK293T cells were co-transfected with 1 µg of pTol2-3×HA-FKBP-EGFP-pA, 1 µg of pTol2-3×V5-FRB-mKate2-PafA-2ApA-CK2, and 2 µg of pEF6a-HB-Pup (or one of pEF6a-SBP-sPup derivatives). After 24 h, 100 nM rapamycin was added to induce binding between HA-FKBP-EGFP and V5-FRB-mKate2-PafA. After 18 h, cells were harvested and lysed with cell-lysis buffer. Samples were then analyzed by western blot assay using antibodies against V5 and HA. To evaluate the levels of target labeling by different Halo-PafA variants or the effect of different drug-HTL derivatives, plates or dishes were coated with 10 µg/ml poly-D-lysine, and HEK293T cells were co-transfected with 1 µg of one of Halo-PafA variants, 2.5 µg of pEF6a-SBP^{K4R}-sPup^{K61R}, and 0.5 µg of pTol2-SRC(247-536)-2×V5. Twenty-four hours later, 250 (or 500 nM) of a DH derivative or 200 nM DC661-H1, unless stated otherwise, was added for an additional 24 h. Samples were then prepared for western blot analysis. For the competition assay with dasatinib, 100 nM DH1 and a serially increasing concentration of dasatinib, ranging from 0.025 to 25.6 µM with a four-fold increment, were cotreated for 24 h. For the pulldown assay, cells were plated on 60 mm dishes, co-transfected with 2 µg of one of Halo-PafA variants and 5 µg of pEF6a-SBP^{K4R}-sPup^{K61R}, and lysed using RIPA buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% Triton X-100, 1 mM EDTA, 5% glycerol) supplemented with a protease inhibitor cocktail. To exogenously expression a target protein, 1 µg of a target plasmid, such as SRC(247-536)-2×V5 and VPS37C-2×V5, was included in the transfection. Cleared cell lysates were incubated with 50 µl of streptavidin magnetic beads for 2 h at 4 °C. The beads were washed five times with wash buffer A (50 mM Tris-HCl pH 7.4, 400 mM NaCl, 1% Triton X-100, 1 mM EDTA, 5% glycerol). Proteins were eluted with 2× SDS sample buffer and processed for western blot analysis.

Preparation of proteomics samples for target-ID of DH5

HEK293T cells were plated on two 100 mm dishes for each condition and co-transfected next day with 8 µg of pTol2-HA-Halo^{8KR}-PafA^{S126A,K172R} and 16 µg of pEF6a-SBP^{K4R}-sPup^{K61R} per dish. After 24 h, treatments with DMSO, DH5 (250 nM), and a competition containing 250 nM DH5 and 2.5 µM dasatinib were applied and incubated for an additional 24 h. Cells were harvested in 1 ml of RIPA buffer, supplemented with a protease inhibitor cocktail for each dish. Following brief sonication, cell debris was removed by centrifugation at maximum speed for 10 min at 4 °C. The supernatants were collected and mixed with 200 µl of streptavidin magnetic beads that had been prewashed with RIPA buffer. The beads were washed five times with wash buffer A, then three times with wash buffer B (wash buffer A supplemented with 1 M urea), and eluted twice with 100 µl of 5 mM biotin and 2.5 µM dasatinib in wash buffer B for 1 h. The eluates were combined and precipitated using trichloroacetic acid (TCA)/DOC. The resultant protein pellets were neutralized with 1 M Tris-HCl pH 8.8, and separated by Tricine-SDS-PAGE, and the CBB-stained bands were excised, with those smaller than 10 kD being removed, and processed for mass analysis.

Preparation of proteomics samples for target-ID of DC661-H1 using SILAC

A stable HEK293T cells expressing the POST-IT system was generated using the PiggyBac transposon system. Briefly, HEK293T cells on a 60 mm dish were co-transfected with 8 µg of PB-puro-CMV-POST-IT-Tet-SBP^{K4R}-sPup^{K61R}-iresGFP and 2 µg of the Super PiggyBac Transposase expression vector (PB200PA-1, System Biosciences, USA). Four days later, cells

were treated with 5 µg/ml puromycin for about two weeks. Media containing puromycin were changed every 3–4 days.

The stable cell line was cultured in SILAC DMEM (280001200, Silantes, Germany) without arginine, lysine, and glutamine, supplemented with 10% dialyzed FBS (C3820-0100, VivaCell Biosciences, China) and 2 mM glutamine. The SILAC medium included 600 mg/L proline (P120032, Aladdin, China) to prevent the conversion of arginine to proline (Bendall et al., 2008). The heavy medium, K8R10, contained 152.1 mg/L heavy labeled lysine K8 ($^{13}C_6{}^{15}N_2$) (211603902, Silantes, Germany) and 44.1 mg/L heavy labeled arginine R10 ($^{13}C_6{}^{15}N_4$) (201603902, Silantes, Germany). The light medium, K0R0, contained 146 mg/L normal lysine K0 (L113006, aladdin, China) and 42 mg/L normal arginine R0 (1119-34, aladdin, China). For SILAC experiments, cells were subcultured for at least 5 cell passages in either heavy or light DMEM medium.

The expression of POST-IT was induced by the addition of 100 ng/ml doxycycline. Two days later, 500 nM DC661-H1 was added to the heavy medium, or 500 nM DC661-H1 and 5 µM DC660 to the light medium. After 24 h of incubation, cells were rinsed with ice-cold PBS and collected with 1 ml RIPA buffer, supplemented with a protease inhibitor cocktail, per dish. Cell debris was removed by centrifugation, and protein concentration was determined by the BCA assay. The protein concentration was adjusted to 2 mg/ml, and 200 µl of prewashed streptavidin magnetic beads was added. After about 3 h of incubation at 4 °C, the beads were washed 5 times with wash buffer A, three times with wash buffer B, and eluted twice with 100 µl of 5 mM biotin and 10 µM DC660 in wash buffer B for 1 h. The eluates were combined and precipitated using TCA/DOC. The resultant protein pellets were neutralized with 1 M Tris-HCl pH 8.8, and separated by Tricine-SDS-PAGE, and the CBB-stained bands were excised, with those smaller than 10 kD being removed, and processed for mass analysis.

LC-MS/MS analysis

Excised gels were sliced into small pieces of ~1 mm^3, rinsed three times with Milli-Q water, decolorized with 50% acetonitrile and 100 mM NH_4HCO_3, and dehydrated with 100% acetonitrile. The dehydrated gel pieces were then reduced with 10 mM DTT in 50 mM NH_4HCO_3 at 56 °C for 1 h, carboxymethylated with 55 mM iodoacetamide in the dark at room temperature for 30 min, and digested with trypsin at a 1:50 enzyme to protein mass ratio at 37 °C overnight. The peptides were collected, desalted with ZipTipC18 (ZTC18S096, Millipore, USA), dried under vacuum, and stored at -20 °C until MS analysis. The resulting tryptic peptides were analyzed by LC-MS/MS at the Institute of Hydrobiology (Chinese Academy of Sciences, Wuhan, China). LC-MS/MS data acquisition was performed using a Q Exactive HF-X mass spectrometer coupled with an Easy-nLC 1200 system (Thermo Fisher Scientific, USA). The peptide mixtures were initially loaded onto a C18 trap column and subsequently separated on a C18 reverse-phase analytical HPLC column, Acclaim PepMap C18 column (75 µm ID × 250 mm, 2 µm particle size, 100 Å pore size), Thermo Fisher Scientific, USA), employing a 100-min gradient program with mobile phase A (0.1% formic acid) and mobile phase B (80% acetonitrile, 0.1% formic acid). The gradient program was as follows: 0-65 min, 5-23% B; 65-85 min, 23-45% B; 85-86 min, 45-90% B; 87-90 min, 90% B; 90-90.1min, 90–5% B; 90.1-100 min, 5% B, maintained at a constant flow rate of 300 nL/min. For data-dependent acquisition (DDA) mode analysis, each scan cycle consisted of one full-scan mass spectrum (R = 60 K, AGC = 3e6, max IT = 20 ms, scan range = 350-1800 m/z) followed by 20 MS/MS events (R = 15 K, AGC = 2e5, max IT = 50 ms). The higher-energy collisional dissociation (HCD) collision energy was set to

757 28 for ion fragmentation. The isolation window for precursor selection was set to 1.6 Da. The
758 former target ion exclusion was set for 25 s.
759

Mass spectrometry and data analysis
For the identification of DH5 targets, raw MS data were analyzed using MaxQuant (version
2.1.0.0, Max Planck Institute of Biochemistry) (Cox and Mann, 2008) with the Andromeda
search engine. The MS data were aligned to the UniProt human protein database (Proteome ID:
UP000005640). Additional sequences, including Halo^{8KR}-PafA^{S126A,K172R} and SBP^{K4R}-sPup^{K61R},
were integrated to identify potential proteomics contaminants. Variable modifications included
oxidation (M), acetyl (Protein N-term), deamidation (NQ), and GGE (K). The false discovery
rate (FDR) for both peptide and protein identification was set at 1%. The "Match Between run"
option were enabled, with other settings at their default values. Missing values from identified
proteins were filled through random imputation and further analyzed statistically to derive *P*
values through unpaired SAM analysis using PANDA-view (Chang et al., 2018). Proteins
commonly present under DMSO condition were considered nonspecific background interactions
and excluded from further analysis.
For the SILAC experiments to identify DC661 targets, raw MS data were analyzed with
FragPipe software v20.0 employing the MSFragger search engine (Kong et al., 2017). Alignment
was performed against the UniProt human protein database (Proteome ID: UP000005640).
Halo^{8KR}-PafA^{S126A,K172R} and SBP^{K4R}-sPup^{K61R} sequences were incorporated as proteomics
contaminants. The SILAC3 workflow was utilized with the default settings for light and heavy
SILAC labels. Proteins detected fewer than 2 times across three biological replicates or
exhibiting a high frequency of occurrence (> 0.15) in CRAPOME (Mellacheruvu et al., 2013)
were filtered out from the analysis.
781

Microscale thermophoresis (MST) assay
SEPHS2-2×V5, VPS37C-2×V5, VPS29, TRAPPC3, TOM1, and TOM1L2 were labeled using
Cy5-NHS ester (A100932, Sangon Biotech, China). Briefly, each target protein was diluted to 10
μM in PBS and underwent buffer exchange with PBS using a Zeba spin desalting column
(89882, Thermo Scientific, USA) to remove Tris. The labeling reaction consisted of 80 μl of 10
μM target protein, 10 μl of 1 M NaHCO$_3$ pH 8.3, and 10 μl of 1 mM Cy5-NHS ester in DMSO,
and was incubated at 4 °C overnight. The next day, unlabeled dyes were removed using Zeba
spin desalting columns, and labeling efficiency was assessed by measuring the concentration of
Cy5 with a UV-Vis spectrophotometer (SMA5000, Merinton, China). Dasatinib, DH5, HCQ, or
DC660 was serially diluted in ligand buffer (PBS with 0.1% Tween-20). Cy5-labeled SEPHS2-
2×V5 (5 nM), VPS37C-2×V5 (5 nM), VPS29 (10 nM), TRAPPC3 (10 nM), TOM1 (10 nM), or
TOM1L2 (10 nM) was mixed with the corresponding diluted compound in a total volume of 20
μl, and then loaded into Monolith premium capillaries (MO-K025, NanoTemper, Germany).
Binding affinity analysis was performed using a NanoTemper Monolith NT.115 instrument
(NanoTemper, Germany). K$_d$ values were obtained using MO.Affinity Analysis 2.3.0 software
(NanoTemper, Germany).
798

Molecular docking simulation for protein-ligand interaction
The docking simulation was performed using CB-Dock2 (https://cadd.labshare.cn/cb-
dock2/index.php), a web server that provides protein-ligand blind docking utilizing Autodock
Vina (version 1.1.2). The blind docking simulation was conducted via protein-surface curvature-

based cavity detection approach following the website procedures. The SEPHS2 structure
information, ma-y60vo.cif, was obtained from the ModelArchive database
(https://doi.org/10.5452/ma-y6ovo) (Nunziata et al., 2019). The PDB files of VPS37 family
proteins (Q8NEZ2, VPS37A; Q9H9H4, VPS37B; A5D8V6, VPS37C; Q86XT2, VPS37D) were
downloaded from AlphaFold database (https://alphafold.ebi.ac.uk/), and the structure of the yeast
ESCRT-I heterotetramer core (2P22) was retrieved from the RCSB protein data bank
(https://www.rcsb.org/). Ligand files were downloaded from PubChem as sdf files. The
superposition of VPS37C with other VPS37A/B/D proteins was performed using the US-align
(Universal Structural alignment) website (https://zhanggroup.org/US-align/) (Zhang et al., 2022).
UCSF ChimeraX 1.6.1 was employed for analyzing, editing, and visualizing the resulting PDB
files.
814
Cellular thermal shift assay (CETSA)
HEK273T cells were cultured on a 100 mm dish and transfected with 10 μg of VPS37C-2×V5
plasmid. Forty-eight hours later, the cells were washed twice with ice-cold PBS, harvested in 1
ml of PBS buffer supplemented with 0.4% NP-40 and 0.5× protease inhibitor cocktail, incubated
for 15 min with gently rotation at room temperature, and centrifuged at maximum speed for 15
min at 4 °C. The supernatant was transferred to a new 1.5 ml microtube, and protein centration
was determined by a BCA assay. Cell lysates were diluted to a concentration of 2 mg/ml and
divided into two groups. The drug group was treated with 20 μM of DC660 or HCQ, while the
control group received the same amount of DMSO. After a 20 min incubation at room
temperature, cell lysates were gently mixed and aliquoted into PCR tubes with 40 μl each, then
subjected to heat treatment for 4 min, ranging from 47 °C to 59 °C in 2 °C increments, and
cooled to room temperature on a T100 thermal cycler (BioRad, USA). Subsequently, cell lysates
were transferred to 1.5 ml microtubes and centrifuged at maximum speed for 20 min at 4 °C to
precipitate unstable insoluble proteins. The supernatant was transferred to a new 1.5 ml
microtube, and 10 μl of 5× SDS sample buffer was added to each tube. Samples were boiled at
95 °C for 5 min and processed for western blot analysis.
831
SRC kinase activity assay
Purified recombinant short SRC(247-536) (0.8 μM), 10 μM ATP, and 40 μM of RR-SRC
peptide, RRLIEDAEYAARG, (T510264-0001, Sangon Biotech, China) were prepared in SRC
Kinase Buffer containing 40 mM Tris-HCl pH 7.5, 20 mM $MgCl_2$, 0.1 mg/ml BSA, 2 mM
$MnCl_2$, and 50 μM DTT. After a 1-h incubation at room temperature, ATP levels were
determined using the Kinase-Lumi luminescent kinase assay (S0150S, Beyotime, China)
according to the manufacturer's instruction. To measure the IC_{50} values for dasatinib and DH5,
dasatinib or DH5 in serial dilution was added to the SRC kinase reaction, and ATP levels were
measured as outlined above. Relative SRC activities were calculated as compared to the DMSO
condition, set as 100%.
842
843 **Zebrafish husbandry**
844 Wild-type AB zebrafish were maintained under standard conditions at 28.5 °C, with a 14-h light
845 and 10-h dark cycle. Embryos were obtained from several matings, typically involving one male
846 and one to two females, and were incubated in zebrafish E3 medium (5 mM NaCl, 0.17 mM
847 KCl, 0.33 mM $CaCl_2$, and 0.33 mM $MgSO_4$) at 28.5 °C with the same light-dark cycle. The

Animal Care and Use Committee of Wuhan University (No. AF078) provided approval for all
procedures involving animals.

In vivo pupylation assay using zebrafish embryos
To express POST-IT in early zebrafish embryos, 20 pg of pUbi-QFGal4-2A-3×Flag-POST-IT
plasmid and 25 pg of transposase mRNA were injected into an embryo at the one-cell stage. The
next day, GFP-positive and live embryos were selected, dechorionated with Pronase E (HY-
114158, MedChemExpress, USA), and divided into two groups of approximately 1,000 embryos
each. One group was treated with DMSO and the other with 10 µM DH5 for 24 h. The embryos
were then collected, rinsed with PBS, and lysed in cell-lysis buffer using a motorized plastic
pestle. The resulting lysates were cleared by centrifugation, and protein concentrations were
measured using the BCA assay. Samples were diluted to 2 mg/ml in 1 ml, mixed with 200 µl of
prewashed streptavidin magnetic beads, and incubate at 4 °C for 3 h. The beads were washed
five times with wash buffer A. Proteins were eluted with 2× SDS sample buffer and
subsequently processed for western blot analysis.

Statistical analysis
Data plotting and statistical analysis were conducted using GraphPad Prism 9.5 software
(GraphPad Software, Inc.). Data are presented as mean ± s.e.m. or mean ± s.d., as specified in
each figure and its legend. *P* values were determined using Student's unpaired two-sided *t*-test,
unless otherwise mentioned. *P* < 0.05 was considered as a statistical difference in this study.

Materials availability statement
Plasmids and cell lines generated in this study are available upon a reasonable request from the
corresponding author (youngnam_jin@whu.edu.cn).

Data availability
All data generated or analyzed during this study are included in the paper and/or the supporting
files. This study includes no data deposited in external repositories.

References

Barrette-Ng IH, Wu S-C, Tjia W-M, Wong S-L, Ng KKS. 2013. The structure of the SBP-Tag–streptavidin complex
    reveals a novel helical scaffold bridging binding pockets on separate subunits. *Acta Crystallogr D Biol
    Crystallogr* **69**:879–887. doi:10.1107/S0907444913002576
Békés M, Langley DR, Crews CM. 2022. PROTAC targeted protein degraders: the past is prologue. *Nat Rev Drug
    Discov* **21**:181–200. doi:10.1038/s41573-021-00371-6
Bendall SC, Hughes C, Stewart MH, Doble B, Bhatia M, Lajoie GA. 2008. Prevention of Amino Acid Conversion
    in SILAC Experiments with Embryonic Stem Cells. *Molecular & Cellular Proteomics* **7**:1587–1597.
    doi:10.1074/mcp.M800113-MCP200
Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL, Perrimon N, Ting AY. 2018.
    Efficient proximity labeling in living cells and organisms with TurboID. *Nat Biotechnol* **36**:880–887.
    doi:10.1038/nbt.4201
Burgess J, Burrows JT, Sadhak R, Chiang S, Weiss A, D'Amata C, Molinaro AM, Zhu S, Long M, Hu C, Krause
    HM, Pearson BJ. 2020. An optimized QF-binary expression system for use in zebrafish. *Developmental
    Biology* **465**:144–156. doi:10.1016/j.ydbio.2020.07.007
Carlisle AE, Lee N, Matthew-Onabanjo AN, Spears ME, Park SJ, Youkana D, Doshi MB, Peppers A, Li R, Joseph
    AB, Smith M, Simin K, Zhu LJ, Greer PL, Shaw LM, Kim D. 2020. Selenium detoxification is required for
    cancer-cell survival. *Nat Metab* **2**:603–611. doi:10.1038/s42255-020-0224-7

Chang C, Xu K, Guo C, Wang J, Yan Q, Zhang J, He F, Zhu Y. 2018. PANDA-view: an easy-to-use tool for
statistical analysis and visualization of quantitative proteomics data. *Bioinformatics* **34**:3594–3596.
doi:10.1093/bioinformatics/bty408
Cho KF, Branon TC, Rajeev S, Svinkina T, Udeshi ND, Thoudam T, Kwak C, Rhee H-W, Lee I-K, Carr SA, Ting
AY. 2020. Split-TurboID enables contact-dependent proximity labeling in cells. *PNAS* **117**:12143–12154.
doi:10.1073/pnas.1919528117
Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass
accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**:1367–1372. doi:10.1038/nbt.1511
Eagle K, Jiang Y, Shi X, Li M, Obholzer ND, Hu T, Perez MW, Koren JV, Kitano A, Yi JS, Lin CY, Nakada D.
2022. An oncogenic enhancer encodes selective selenium dependency in AML. *Cell Stem Cell* **29**:386-
399.e7. doi:10.1016/j.stem.2022.01.003
Enchev RI, Schulman BA, Peter M. 2015. Protein neddylation: beyond cullin–RING ligases. *Nat Rev Mol Cell Biol*
16:30–44. doi:10.1038/nrm3919
Erdmann RS, Baguley SW, Richens JH, Wissner RF, Xi Z, Allgeyer ES, Zhong S, Thompson AD, Lowe N, Butler
R, Bewersdorf J, Rothman JE, St Johnston D, Schepartz A, Toomre D. 2019. Labeling Strategies Matter for
Super-Resolution Microscopy: A Comparison between HaloTags and SNAP-tags. *Cell Chemical Biology*
26:584-592.e6. doi:10.1016/j.chembiol.2019.01.003
Friedman Ohana R, Kirkland TA, Woodroofe CC, Levin S, Uyeda HT, Otto P, Hurst R, Robers MB, Zimmerman K,
Encell LP, Wood KV. 2015. Deciphering the Cellular Targets of Bioactive Compounds Using a
Chloroalkane Capture Tag. *ACS Chem Biol* **10**:2316–2324. doi:10.1021/acschembio.5b00351
Ha J, Park H, Park J, Park SB. 2021. Recent advances in identifying protein targets in drug discovery. *Cell Chemical
Biology* **28**:394–423. doi:10.1016/j.chembiol.2020.12.001
Hill ZB, Pollock SB, Zhuang M, Wells JA. 2016a. Direct Proximity Tagging of Small Molecule Protein Targets
Using an Engineered NEDD8 Ligase. *J Am Chem Soc* **138**:13123–13126. doi:10.1021/jacs.6b06828
Hill ZB, Pollock SB, Zhuang M, Wells JA. 2016b. Direct Proximity Tagging of Small Molecule Protein Targets
Using an Engineered NEDD8 Ligase. *J Am Chem Soc* **138**:13123–13126. doi:10.1021/jacs.6b06828
Hughes RE, Elliott RJR, Dawson JC, Carragher NO. 2021. High-content phenotypic and pathway profiling to
advance drug discovery in diseases of unmet need. *Cell Chemical Biology* **28**:338–355.
doi:10.1016/j.chembiol.2021.02.015
Jain V, Singh MP, Amaravadi RK. 2023. Recent advances in targeting autophagy in cancer. *Trends in
Pharmacological Sciences* **44**:290–302. doi:10.1016/j.tips.2023.02.003
Javed R, Jain A, Duque T, Hendrix E, Paddar MA, Khan S, Claude-Taupin A, Jia J, Allers L, Wang F, Mudd M,
Timmins G, Lidke K, Rusten TE, Akepati PR, He Y, Reggiori F, Eskelinen E, Deretic V. 2023.
Mammalian ATG8 proteins maintain autophagosomal membrane integrity through ESCRTs. *The EMBO
Journal* **42**:e112845. doi:10.15252/embj.2022112845
Jiang H-W, Czajkowsky DM, Wang T, Wang X-D, Wang J, Zhang H-N, Liu C-X, Wu F-L, He X, Xu Z-W, Chen
H, Guo S-J, Li Y, Bi L-J, Deng J-Y, Xie J, Pei J-F, Zhang X-E, Tao S-C. 2018. Identification of Serine 119
as an Effective Inhibitor Binding Site of M. tuberculosis Ubiquitin-like Protein Ligase PafA Using Purified
Proteins and M. smegmatis. *EBioMedicine* **30**:225–236. doi:10.1016/j.ebiom.2018.03.025
Jin YN, Schlueter PJ, Jurisch-Yaksi N, Lam P-Y, Jin S, Hwang WY, Yeh J-RJ, Yoshigi M, Ong S-E, Schenone M,
Hartigan CR, Carr SA, Peterson RT. 2018. Noncanonical translation via deadenylated 3′ UTRs maintains
primordial germ cells. *Nat Chem Biol* **14**:844–852. doi:10.1038/s41589-018-0098-0
Kang M-G, Lee H, Kim BH, Dunbayev Y, Seo JK, Lee C, Rhee H-W. 2017. Structure-guided synthesis of a protein-
based fluorescent sensor for alkyl halides. *Chem Commun* **53**:9226–9229. doi:10.1039/C7CC03714G
Kong AT, Leprevost FV, Avtonomov DM, Mellacheruvu D, Nesvizhskii AI. 2017. MSFragger: ultrafast and
comprehensive peptide identification in mass spectrometry–based proteomics. *Nat Methods* **14**:513–520.
doi:10.1038/nmeth.4256
Kwak C, Park C, Ko M, Im CY, Moon H, Park Y-H, Kim SY, Lee S, Kang M-G, Kwon HJ, Hong E, Seo JK, Rhee
H-W. 2022. Identification of proteomic landscape of drug-binding proteins in live cells by proximity-
dependent target ID. *Cell Chemical Biology* **29**:1739-1753.e6. doi:10.1016/j.chembiol.2022.10.001
Lam SS, Martell JD, Kamer KJ, Deerinck TJ, Ellisman MH, Mootha VK, Ting AY. 2015. Directed evolution of
APEX2 for electron microscopy and proximity labeling. *Nat Methods* **12**:51–54. doi:10.1038/nmeth.3179
Lee S-Y, Cheah JS, Zhao B, Xu C, Roh H, Kim CK, Cho KF, Udeshi ND, Carr SA, Ting AY. 2023. Engineered
allostery in light-regulated LOV-Turbo enables precise spatiotemporal control of proximity labeling in
living cells. *Nat Methods* **20**:908–917. doi:10.1038/s41592-023-01880-5

Liu H, Naismith JH. 2008. An efficient one-step site-directed deletion, insertion, single and multiple-site plasmid
953   mutagenesis protocol. *BMC Biotechnology* **8**:91. doi:10.1186/1472-6750-8-91
954   Liu Q, Zheng J, Sun W, Huo Y, Zhang L, Hao P, Wang H, Zhuang M. 2018. A proximity-tagging system to identify
955   membrane protein–protein interactions. *Nature Methods* **15**:715. doi:10.1038/s41592-018-0100-5
956   Lomenick B, Hao R, Jonai N, Chin RM, Aghajan M, Warburton S, Wang J, Wu RP, Gomez F, Loo JA,
957   Wohlschlegel JA, Vondriska TM, Pelletier J, Herschman HR, Clardy J, Clarke CF, Huang J. 2009. Target
958   identification using drug affinity responsive target stability (DARTS). *Proceedings of the National
959   Academy of Sciences* **106**:21984–21989. doi:10.1073/pnas.0910040106
960   Los GV, Encell LP, McDougall MG, Hartzell DD, Karassina N, Zimprich C, Wood MG, Learish R, Ohana RF, Urh
961   M, Simpson D, Mendez J, Zimmerman K, Otto P, Vidugiris G, Zhu J, Darzins A, Klaubert DH, Bulleit RF,
962   Wood KV. 2008. HaloTag: A Novel Protein Labeling Technology for Cell Imaging and Protein Analysis.
963   *ACS Chem Biol* **3**:373–382. doi:10.1021/cb800025k
964   Martell JD, Deerinck TJ, Sancak Y, Poulos TL, Mootha VK, Sosinsky GE, Ellisman MH, Ting AY. 2012.
965   Engineered ascorbate peroxidase as a genetically encoded reporter for electron microscopy. *Nat Biotechnol*
966   **30**:1143–1148. doi:10.1038/nbt.2375
967   Mauthe M, Orhon I, Rocchi C, Zhou X, Luhr M, Hijlkema K-J, Coppes RP, Engedal N, Mari M, Reggiori F. 2018.
968   Chloroquine inhibits autophagic flux by decreasing autophagosome-lysosome fusion. *Autophagy* **14**:1435–
969   1455. doi:10.1080/15548627.2018.1474314
970   May DG, Scott KL, Campos AR, Roux KJ. 2020. Comparative Application of BioID and TurboID for Protein-
971   Proximity Biotinylation. *Cells* **9**:1070. doi:10.3390/cells9051070
972   Mellacheruvu D, Wright Z, Couzens AL, Lambert J-P, St-Denis NA, Li T, Miteva YV, Hauri S, Sardiu ME, Low
973   TY, Halim VA, Bagshaw RD, Hubner NC, al-Hakim A, Bouchard A, Faubert D, Fermin D, Dunham WH,
974   Goudreault M, Lin Z-Y, Badillo BG, Pawson T, Durocher D, Coulombe B, Aebersold R, Superti-Furga G,
975   Colinge J, Heck AJR, Choi H, Gstaiger M, Mohammed S, Cristea IM, Bennett KL, Washburn MP, Raught
976   B, Ewing RM, Gingras A-C, Nesvizhskii AI. 2013. The CRAPome: a contaminant repository for affinity
977   purification–mass spectrometry data. *Nat Methods* **10**:730–736. doi:10.1038/nmeth.2557
978   Molina DM, Jafari R, Ignatushchenko M, Seki T, Larsson EA, Dan C, Sreekumar L, Cao Y, Nordlund P. 2013.
979   Monitoring Drug Target Engagement in Cells and Tissues Using the Cellular Thermal Shift Assay. *Science*
980   **341**:84–87. doi:10.1126/science.1233606
981   Muller R. 2021. Systemic toxicity of chloroquine and hydroxychloroquine: prevalence, mechanisms, risk factors,
982   prognostic and screening possibilities. *Rheumatol Int* **41**:1189–1202. doi:10.1007/s00296-021-04868-6
983   Nirk EL, Reggiori F, Mauthe M. 2020. Hydroxychloroquine in rheumatic autoimmune disorders and beyond. *EMBO
984   Molecular Medicine* **12**:e12476. doi:10.15252/emmm.202012476
985   Nunziata C, Polo A, Sorice A, Capone F, Accardo M, Guerriero E, Marino FZ, Orditura M, Budillon A, Costantini
986   S. 2019. Structural analysis of human SEPHS2 protein, a selenocysteine machinery component, over-
987   expressed in triple negative breast cancer. *Sci Rep* **9**:16131. doi:10.1038/s41598-019-52718-0
988   Özcelik D, Barandun J, Schmitz N, Sutter M, Guth E, Damberger FF, Allain FH-T, Ban N, Weber-Ban E. 2012.
989   Structures of Pup ligase PafA and depupylase Dop from the prokaryotic ubiquitin-like modification
990   pathway. *Nat Commun* **3**:1014. doi:10.1038/ncomms2009
991   Patton EE, Zon LI, Langenau DM. 2021. Zebrafish disease models in drug discovery: from preclinical modelling to
992   clinical trials. *Nat Rev Drug Discov* **20**:611–628. doi:10.1038/s41573-021-00210-8
993   Rebecca VW, Nicastri MC, Fennelly C, Chude CI, Barber-Rotenberg JS, Ronghe A, McAfee Q, McLaughlin NP,
994   Zhang G, Goldman AR, Ojha R, Piao S, Noguera-Ortega E, Martorella A, Alicea GM, Lee JJ, Schuchter
995   LM, Xu X, Herlyn M, Marmorstein R, Gimotty PA, Speicher DW, Winkler JD, Amaravadi RK. 2019.
996   PPT1 Promotes Tumor Growth and Is the Molecular Target of Chloroquine Derivatives in Cancer. *Cancer
997   Discovery* **9**:220–229. doi:10.1158/2159-8290.CD-18-0706
998   Rix U, Hantschel O, Dürnberger G, Remsing Rix LL, Planyavsky M, Fernbach NV, Kaupe I, Bennett KL, Valent P,
999   Colinge J, Köcher T, Superti-Furga G. 2007. Chemical proteomic profiles of the BCR-ABL inhibitors
imatinib, nilotinib, and dasatinib reveal novel kinase and nonkinase targets. *Blood* **110**:4055–4063.
doi:10.1182/blood-2007-07-102061
Roux KJ, Kim DI, Raida M, Burke B. 2012. A promiscuous biotin ligase fusion protein identifies proximal and
interacting proteins in mammalian cells. *Journal of Cell Biology* **196**:801–810. doi:10.1083/jcb.201112098
Savitski MM, Reinhard FBM, Franken H, Werner T, Savitski MF, Eberhard D, Molina DM, Jafari R, Dovega RB,
Klaeger S, Kuster B, Nordlund P, Bantscheff M, Drewes G. 2014. Tracking cancer drugs in living cells by
thermal profiling of the proteome. *Science* **346**:1255784. doi:10.1126/science.1255784

Shi H, Zhang C-J, Chen GYJ, Yao SQ. 2012. Cell-Based Proteome Profiling of Potential Dasatinib Targets by Use
of Affinity-Based Probes. *J Am Chem Soc* **134**:3001–3014. doi:10.1021/ja208518u
Takahashi Y, He H, Tang Z, Hattori T, Liu Y, Young MM, Serfass JM, Chen L, Gebru M, Chen C, Wills CA,
Atkinson JM, Chen H, Abraham T, Wang H-G. 2018. An autophagy assay reveals the ESCRT-III
component CHMP2A as a regulator of phagophore closure. *Nat Commun* **9**:2855. doi:10.1038/s41467-018-
05254-w
Takahashi Y, Liang X, Hattori T, Tang Z, He H, Chen H, Liu X, Abraham T, Imamura-Kawasawa Y, Buchkovich
NJ, Young MM, Wang H-G. 2019. VPS37A directs ESCRT recruitment for phagophore closure. *Journal of
Cell Biology* **218**:3336–3354. doi:10.1083/jcb.201902170
Tao AJ, Jiang J, Gadbois GE, Goyal P, Boyle BT, Mumby EJ, Myers SA, English JG, Ferguson FM. 2023. A biotin
targeting chimera (BioTAC) system to map small molecule interactomes in situ. *Nat Commun* **14**:8016.
doi:10.1038/s41467-023-43507-5
Vincent F, Nueda A, Lee J, Schenone M, Prunotto M, Mercola M. 2022. Phenotypic drug discovery: recent
successes, lessons learned and new directions. *Nat Rev Drug Discov* **21**:899–914. doi:10.1038/s41573-022-
00472-w
Watrous J, Burns K, Liu W-T, Patel A, Hook V, Bafna V, Barry 3rd CE, Bark S, Dorrestein PC. 2010. Expansion of
the mycobacterial "PUPylome." *Mol BioSyst* **6**:376–385. doi:10.1039/B916104J
Wu S-C, Wang C, Chin J, Wong S-L. 2019. A bio-coupling approach using a dextran-binding domain to immobilize
an engineered streptavidin to Sephadex for easy preparation of affinity matrix. *Sci Rep* **9**:3359.
doi:10.1038/s41598-019-40044-4
Xiong Z, Lo HP, McMahon K-A, Martel N, Jones A, Hill MM, Parton RG, Hall TE. 2021. In vivo proteomic
mapping through GFP-directed proximity-dependent biotin labelling in zebrafish. *eLife* **10**:e64631.
doi:10.7554/eLife.64631
Zhang C, Shine M, Pyle AM, Zhang Y. 2022. US-align: universal structure alignments of proteins, nucleic acids,
and macromolecular complexes. *Nat Methods* **19**:1109–1115. doi:10.1038/s41592-022-01585-1
Zhen Y, Spangenberg H, Munson MJ, Brech A, Schink KO, Tan K-W, Sørensen V, Wenzel EM, Radulovic M,
Engedal N, Simonsen A, Raiborg C, Stenmark H. 2020. ESCRT-mediated phagophore sealing during
mitophagy. *Autophagy* **16**:826–841. doi:10.1080/15548627.2019.1639301
Zhu H, Oh JH, Matsuda Y, Mino T, Ishikawa M, Nakamura H, Tsujikawa M, Nonaka H, Hamachi I. 2024.
Tyrosinase-Based Proximity Labeling in Living Cells and In Vivo. *J Am Chem Soc*.
doi:10.1021/jacs.3c13183
Zhuang M, Guan S, Wang H, Burlingame AL, Wells JA. 2013. Substrates of IAP Ubiquitin Ligases Identified with
a Designed Orthogonal E3 Ligase, the NEDDylator. *Molecular Cell* **49**:273–282.
doi:10.1016/j.molcel.2012.10.022
1041

Acknowledgments

We thank the staff in the core facility of the Medical Research Institute at Wuhan University for their technical support with equipment, and the Institute of Hydrobiology for their support in proteomics. We are also grateful to Dr. Min Zhuang and Dr. Yang Yu for providing the PafA and HaloTag7 plasmids, respectively. This work was supported by the Fundamental Research Funds for the Central Universities (2042022dx0003 to Y.N.J.) and the National Natural Science Foundation of China (32070832 and 32150610476 to Y.N.J.; 82273774 to H.-B.Z.).

Author contributions

Yingjie Sun, Formal analysis, Investigation, Methodology, Writing—original draft; Changheng Li, Formal analysis, Investigation, Methodology, Writing—original draft; Xiaofei Deng, Investigation, Methodology, Writing—original draft; Wenjie Li, Investigation; Xiaoyi Deng, Visualization, Investigation; Weiqi Ge, Investigation; Miaoyuan Shi, Investigation; Ying Guo, Investigation; Yanxun V Yu, Supervision, Project administration, Funding acquisition, Writing—original draft, Writing—review and editing; Hai-Bing Zhou, Supervision, Methodology, Project administration, Funding acquisition, Writing—original draft, Writing—

review and editing; Youngnam N Jin, Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Project administration, Supervision, Funding acquisition, Writing—original draft, Writing—review and editing.

Competing interest
The authors declare that no competing interests exist.

Figures legends

Figure 1. Schematic illustration of the POST-IT system for target identification. Upon the introduction of a drug-specific HaloTag ligand derivative, POST-IT tags target proteins with sPup, a tagging process known as pupylation, in live cells or organisms. Subsequently, these proteins are enriched and identified via mass spectrometry (MS) analysis.
The online version of this article includes the following source data and figure supplement(s) for figure 1:
Figure supplement 1. PafA fused with HaloTag exhibits high activity across a wide temperature range.
Figure supplement 2. Lysine-mutated Pup substrates bypass polypupylation while retaining activities comparable to wild-type Pup.
Figure supplement 3. Optimization of Halo-PafA.

Figure 2. Optimized Halo-PafA and Pup substrates efficiently promote pupylation in vitro in a proximity-dependent manner. (**A**) The chemical structure of DH1, an HTL derivative of dasatinib. (**B**) Comparison of proximity-tagging by different Halo-PafA derivatives on in vitro pupylation. (**C**) Comparison of different Pup substrates on in vitro pupylation with 500 nM DH1. (**D**) Pupylation levels by DH1 decrease with an increasing amount of dasatinib as a competitor, ranging from 0.2 μM to 10 μM. (**B-D**) All reactions were conducted with 1 μM of a Halo-PafA derivative, 10 μM of one of the different Pup substrates, and 0.5 μM of a purified short SRC with a 2×V5 tag, SRC(247-536)-2×V5, at 37 °C for 30 min. Pupylation levels of SRC were assessed by immunoblot (IB) using an anti-V5 antibody.
The online version of this article includes the following source data and figure supplement(s) for figure 2:
Figure 2—source data 1. Original western blots for **Figure 2**.
Figure supplement 1. Halo^{8KR}-PafA demonstrates efficient labeling with HaloTag ligands.
Figure supplement 2. Optimized Halo-PafA and Pup substrates efficiently promote pupylation in vitro in a proximity-dependent manner.
Figure supplement 1—source data 1. Original western blots for **Figure 2—figure supplement 1**.
Figure supplement 2—source data 1. Original western blots for **Figure 2—figure supplement 2**.

Figure 3. POST-IT can label target proteins in live cells.
(**A**) Effect of linker length between Halo^{8KR} and PafA on target-labeling. HEK293T cells were co-transfected with HA-Halo^{8KR}-PafA containing the specified length of linker, SBP^{K4R}-sPup^{K61R}, and SRC(247-536)-2×V5. (**B**) Introducing mutations S126A and K172R into PafA significantly enhances its proximity-tagging efficiency on target. (**C-E**) Comparison of labeling activity of Halo-PafA and Halo^{8KR}-PafA^{S126A,K172R}, a form used for POST-IT. (**F**, **G**) POST-IT^{DH1} mediates proximity-tagging in a DH1-dose dependent manner (**F**), an effect completely inhibited by competitive dasatinib, ranging from 0.025 to 25.6 μM (**G**). Pupylation levels of exogenously expressed short SRC (**C**, **F**, **G**), endogenous SRC (**D**), or pull-downed endogenous SRC (**E**) were assessed by immunoblot (IB) analysis using antibodies against V5-tag or SRC for exogenous or endogenous SRC, respectively. In all experiments, SBP^{K4R}-sPup^{K61R} was used for co-transfection, and cells were treated with 500 nM DH1, except in (**F**, **G**).

The online version of this article includes the following source data and figure supplement(s) for figure 3:

Figure 3—source data 1. Original western blots for **Figure 3**.

Figure supplement 1. Test of small molecule-induced proximity-tagging by PafA in HEK293T cells.

Figure supplement 2. Identification of self-pupylated lysine residues in PafA.

Figure supplement 3. PafA from *Cglu* outperforms PafA from *Mtb* in the POST-IT system in live cells.

Figure supplement 1—source data 1. Original western blots for **Figure 3—figure supplement 1**.

Figure supplement 3—source data 1. Original western blots for **Figure 3—figure supplement 3**.

Figure 4. Dasatinib-HTL (DH) derivatives with modified and longer linkers enhances POST-IT performance.

(**A**) In vitro binding assay for DH derivatives via fluorescence polarization (FP) assay. Purified Myc-Halo-PafA (15 nM) and 1 nM Halo-AF488 were incubated with various DH derivatives in a serial dilution at 37 °C for 30 min prior to FP measurement. Data are shown as mean ± s.d. (**B**) Cellular binding assay for DH derivatives by FP measurement. HEK293T cells transfected with Myc-Halo-PafA were later incubated with 1 µM of each DH derivative or DMSO for 3 h, followed by cell lysate incubation with 2 nM of Halo-AF488 at 37 °C for 30 min. $n = 4$. Data are shown as mean ± s.e.m. *P* values were calculated by an unpaired two-sided *t*-test. Immunoblot presents a representative image of the input levels for each condition, demonstrating that treatment with DH derivatives did not alter the expression levels of Myc-Halo-PafA. (**C**) In vitro labeling activity comparison among DH derivatives. Purified recombinant Halo^{8KR}-PafA (0.5 µM), SBP^{K4R}-sPup^{K61R} (10 µM), and SRC(247-536)-2×V5 (0.5 µM) were incubated with 0.5 µM of each DH derivative at 37 °C for 30 min. (**D, E**) In cellular labeling activity comparison among DH derivatives. HA-Halo^{8KR}-PafA and SBP^{K4R}-sPup^{K61R} were used for co-transfection, and cells were treated with 250 nM of various DH derivatives. Pupylation levels of purified short SRC (**C**), endogenous SRC (**D**), or pull-downed endogenous SRC (**E**) were assessed by immunoblot (IB) using anti-V5 or anti-SRC antibodies for exogenous or endogenous SRC, respectively.

The online version of this article includes the following source data and figure supplement(s) for figure 4:

Figure 4—source data 1. Original western blots for **Figure 4**.

Figure supplement 1. Chemical structures of dasatinib and its HTL derivatives with varied linkers.

Figure supplement 2. DH5 maintained robust kinase inhibitory activity.

Figure 5. Target-ID by POST-IT^{DH5}.

(**A**) Volcano plot displaying the relative fold change (FC) of DH5-binding proteins compared to those with dasatinib competition. Known target proteins for dasatinib were highlighted in red, and a newly identified target protein, SEPHS2, was marked in blue. (**B, C**) MST analysis demonstrates a direct interaction between DH5 and Cy5-labeled SEPHS2 (**B**) or between dasatinib and Cy5-labeled SEPHS2 (**C**), yielding K_d values of 11.6 ± 5.7 µM and 9.1 ± 0.8 µM, respectively. Data are shown as mean ± s.d. (**D**) Molecular docking binding pose between dasatinib and SEPHS2. The yellow box indicates the binding pocket, expanded for a closer view.

The binding pocket of dasatinib precisely fits into the active sites of SEPHS2. (**E**) Diagram of 2D molecular docking interaction between SEPHS2 and dasatinib. Red boxes highlight the active sites, and blue boxes indicate residues among HUB nodes.

Figure 6. Target-ID for HCQ by POST-IT$^{DC661-H1}$.
(**A**) Rank plot analysis of SILAC ratio values of proteins in three biological replicates. Autophage-related proteins ranked highly are marked in red. Heavy-labeled cells were treated with 200 nM DC661-H1, whereas light-labeled cells underwent incubation with both DC661-H1 (200 nM) and competitive DC660 (2 µM). (**B**, **C**) MST analyses demonstrate direct interactions of DC660 (**B**) and HCQ (**C**) with VPS37C, yielding K_d values of 21.5 ± 9.8 nM and 16.9 ± 8.7 µM, respectively. Data are shown as mean $\pm$ s.d. (**D**, **E**) Immunoblot results indicate that the VPS37C-V5 protein from cells treated with DC661-H1 was significantly enriched after streptavidin pulldown. The competition between DC661-H1 and either DC660 (**D**) or HCQ (**E**) nearly completely abolished VPS37C binding. HEK293T cells were co-transfected with HA-Halo^{8KR}-PafA^{S126A,K172R}, SBP^{K4R}-sPup^{K61R}, and VPS37C-V5. After 24 h, cells were incubated with 200 nM of DC661-H1, with or without 2 µM of DC660 (**D**) or HCQ (**E**). Twenty-four hours later, cells were collected for further analysis. (**F**, **G**) CETSA results demonstrate that VPS37C becomes thermostable when exposed to DC660 (**F**) or HCQ (**G**). Selective stabilization of VPS37C is evident at temperatures of 53 °C or higher, a phenomenon not observed in β-actin.
The online version of this article includes the following source data and figure supplement(s) for figure 6:
Figure 6—source data 1. Original western blots for **Figure 6**.
Figure supplement 1. A DC661-HTL derivative, DC661-H1, is highly active as an autophagy inhibitor.
Figure supplement 2. Schematic of the experimental flow for applying POST-IT in SILAC.
Figure supplement 3. MST binding analysis of candidate proteins identified with DC661-H1.
Figure supplement 4. Molecular docking binding poses of DC661 and HCQ with the VPS37 family.
Figure supplement 1—source data 1. Original western blots for **Figure 6—figure supplement 1**.

Figure. 7. POST-IT enables in vivo target-ID.
(**A**) Schematic of experimental design for applying POST-IT in zebrafish. A plasmid construct containing the POST-IT system was injected into embryos at the one-cell stage. The injected embryos were dechorionated 24 h later, and then treated with DH5 or DMSO for an additional 24 h. Embryos lysates were collected and analyzed by immunoblot. (**B**) Representative images of injected embryos. Bright field (BF) images reveal no overt toxicity from POST-IT expression. EGFP images display robust expression of POST-IT. Scale bar, 500 µm. (**C**) Immunoblot analysis demonstrates that POST-IT exhibits significant enrichment of SRC in the presence of DH5 but not DMSO.
Figure 7—source data 1. Original western blots for **Figure 7**.

Table legend

Table 1. Identification of lysine residues for pupylation in PafA by mass spectrometry.

Sun Y *et al*., Fig. 1

1216

Sun Y *et al*., Fig. 2

1226

1233 **Sun Y *et al*., Fig. 3**
1234
1235

1236

Sun Y *et al*., Fig. 4

Sun Y *et al*., Fig. 5

1256

1263   **Sun Y *et al*., Fig. 6**
1264
1265

Sun Y *et al*., Fig. 7

1276

Table 1. Identification of lysine residues for pupylation in PafA by mass spectrometry.

1278

1279

1280

1281

1282

Protein	Score	Coverage (%)	# PSMs	Detected pupylated sequences	MH+ [Da]	Xcorr	Lysine	Rank of pupylation
PafA	607.055	89	1197					
			1	RIMGIETEYGLTFVDGDSKK	2502.21808	5.40		
			2	RIMGIETEYGLTFVDGDSKK	2518.21299	5.35	K29	3
			2	IMGIETEYGLTFVDGDSKK	2362.11188	3.71		
			1	IMGIETEYGLTFVDGDSKK	2346.11697	3.46	K30	5
			1	MFRPIVEKYSSSNIFIPNGSR	2702.32428	2.93	K47	5
			1	MAVDAEESLAK	1406.64679	3.39		
			2	MAVDAEESLAKEDIAGQVYLFK	2670.29672	3.86	K106	2
			4	MAVDAEESLAKEDIAGQVYLFK	2686.29164	3.86		
			5	IHHPNPLDKGESFPLGYCISQR	2808.35222	5.10	K172	1
			3	IHHPNPLDKGESFPLGYCISQR	2809.33624	2.88		
			1	ILKAADTLGVPVTVDWMR	2228.17436	3.61	K427	4
			1	ILKAADTLGVPVTVDWMR	2244.16928	3.49		