A KCa2.2/2.3 opener reverses ET-1 induced NLRP3 activation in hypertensive mice

Fais, RS[1,2,3]; Comerma-Steffensen, SG[2,4]; Pinilla, E[2]; Matchkov, VV[2]; Tostes RC[1]; Carneiro, FS[1]; Simonsen, U[2].

[1] Department of Pharmacology, Ribeirao Preto Medical School, University of Sao Paulo, Ribeirao Preto-SP, Brazil

[2] Department of Biomedicine, Pulmonary and Cardiovascular Pharmacology, Aarhus University, Aarhus, Denmark

[3] Division of Pulmonary, Critical Care, and Sleep Medicine, National Jewish Health, Denver-CO, USA

[4] Department of Biomedical Sciences/Animal Physiology, Faculty of Veterinary, Central University of Venezuela, Venezuela.

Corresponding Authors: Ulf Simonsen, MD, Ph.D., Department of Biomedicine, Pulmonary and Cardiovascular Pharmacology, Aarhus University, Ole Worms Allé 4, building 1160, 219, 8000 Aarhus C, Denmark, us@biomed.au.dk, Phone: +4560202613, Mobile: +4560202613

Rafael Sobrano Fais, Ph.D., Division of Pulmonary, Critical Care, and Sleep Medicine, National Jewish Health, 1400 Jackson St., Goodman Hall, room 1007, Denver, CO 80206, USA. Phone +1(317)4760228; rsfais@gmail.com

Clinical Perspective

What Is New?

● Our data indicate that the immune system, through the NLRP3 inflammasome, has a central role in endothelin-1- and hypertension-associated erectile dysfunction.

● Endothelin-1 mediates the inflammasome activation through $K_{Ca}2.3$ inhibition in endothelial cells of erectile tissue

● An opener of $K_{Ca}2.3$ channels, NS13001 counteracts endothelin-1-induced suppression of Ca^{2+}-activated K^+ current in the endothelial cell membrane and development of erectile dysfunction in DOCA mice

What Are the Clinical Implications?

● Endothelin-1 contributes to hypertension and erectile dysfunction by provoking an endothelial dysfunction through vascular inflammation,

● NLRP3 inflammasome activation in endothelial cells is a key pathway behind endothelin-1-induced erectile dysfunction and is potentially involved in cardiovascular complications in patients with hypertension.

● The $K_{Ca}2.3$ channel is a putative novel therapeutic target in erectile dysfunction, especially when the endothelin-1-NLRP3 inflammasome is involved.

A KCa2.2/2.3 opener reverses ET-1 induced NLRP3 activation in hypertensive mice

Fais, RS; Comerma-Steffensen, SG; Pinilla, E; Matchkov, V; Tostes RC; Carneiro, FS; Simonsen, U.

Membrane depolarization is implicated in the activation of the NLRP3 inflammasome. Downregulation of endothelial Ca2+-activated K+ channels type 2.3 (K_{Ca} 2.3) and endothelin-1 (ET-1) upregulation in the corpus cavernosum (CC) are associated with erectile dysfunction. We hypothesized that opening K_{Ca} 2.2/2.3 channels could reverse erectile dysfunction and NLRP3 activation in hypertensive DOCA/salt mice. Methods: Hypertension was induced in mice using a DOCA/salt model, with unilaterally nephrectomized mice as controls; blood pressure was measured by tail-cuff. Intracavernous pressure (ICP) and CC reactivity were assessed. Western blot analysis for K_{Ca} 2.3, caspase-1, and interleukin-1β (IL-1β) was performed. Endothelial cells from erectile tissue were isolated and stimulated with ET-1, and K_{Ca} 2.2/2.3-dependent currents were evaluated via voltage-clamp electrophysiology. Results: DOCA/salt mice exhibited impaired erectile function, increased pro-caspase-1 and caspase-1 expression, and reduced relaxations induced by acetylcholine (ACh), sodium nitroprusside (SNP), and electrical field stimulation (EFS). Treatment with either the endothelin receptor antagonist bosentan or the K_{Ca} 2.2/2.3 channel opener NS13001 reversed these dysfunctions and reduced ET-1-induced NLRP3 activation in DOCA/salt mice. NS13001 also restored the decreased current observed in primary endothelial cells exposed to ET-1. Apamin, a K_{Ca} 2.2/2.3 channel blocker, inhibited erectile responses in unilaterally nephrectomized mice and restored erectile responses in DOCA mice. Apamin did not affect EFS-, ACh-, or SNP-induced relaxation in CC from hypertensive DOCA/salt mice. Conclusion: NS13001 reversed ET-1-induced NLRP3 activation and erectile dysfunction in DOCA/salt mice, suggesting that K_{Ca} 2.2/2.3 channel modulation may restore erectile function in hypertension-related diseases.

Introduction

Endothelial dysfunction is commonly associated with arterial hypertension and erectile dysfunction, which may be triggered by altered function of ion channels such as Ca^{2+}-activated K^+ channels (K_{Ca}), mainly the small (SK_{Ca}) conductance[1-3]. The K_{Ca} form the largest and most diverse group of ion channels and are divided into small SK_{Ca} ($K_{Ca}2.1$, $K_{Ca}2.2$, and, $K_{Ca}2.3$), intermediate (IK_{Ca} or ($K_{Ca}3.1$), and large conductance Ca^{2+}-activated K^+ channels (BK_{Ca} or $K_{Ca}1.1$)[4]

The $K_{Ca}2.3$ channel is essential for nitric oxide (NO) signaling through endothelial NO synthase (eNOS) activation. NO produced by eNOS in the vascular bed mediates not only vasodilation, but it also has anti-inflammatory functions[5, 6]. Downregulation of $K_{Ca}2.3$ leads to endothelial and erectile dysfunction, and the expression and/or function of $K_{Ca}2.3$ channels is reduced in arterial hypertension[1, 7].

The activation of $K_{Ca}2.3$ reduces blood pressure[8] and improves erectile function in rats. Furthermore, the $K_{Ca}2.3$ has an anti-inflammatory effect, mainly, because it inhibits the release of proinflammatory cytokines such as interleukin (IL)-6, tumor necrosis factor-alpha (TNF-α), and the migration of immune system cells. Thus, $K_{Ca}2.3$ channel plays an important role in the modulation of the immune system[5, 6].

ET-1 is closely linked with endothelial dysfunction, mainly by the inhibition of various populations of K^+ channels, such as inward-rectifying (K_{ir}), ATP-sensitive (K_{ATP}), voltage-gated (K_v), and K_{Ca} channels[1, 9]. ET-1 has been shown to activate mononuclear phagocytes and to promote the release of TNF-α, IL-1β, and IL-6, which increases the interaction of endothelial cells and monocytes. Moreover, a recent study showed that ET-1 activates NLRP3 via Ca^{2+}-induced ROS generation and impairs erectile function[10-12].

Membrane depolarization is thought to be involved in the activation of NLRP3[13]. Therefore, the present study hypothesized that opening $K_{Ca}2.3$ channels prevents the activation of NLRP3 and reverses erectile dysfunction. To address the hypothesis, studies were conducted on the effect of ET-1 and an opener of KCa 2.2/2.3 channels, NS13001, on the endothelial cell K^+ currents, and the effect of NS13001 on erectile function was examined in DOCA/salt hypertensive mouse model.

Materials and Methods

For a detailed description of the methods and composition of solutions, see Data Supplement. The data that support the findings of this study are available from the corresponding author upon reasonable request.

Animals

10-12 weeks-old C57BL/6 (~25 g) male mice were housed in a room with controlled temperature (20 - 22°C) and on light/dark cycles of 12 hrs with free access to standard chow and filtered water. All experimental protocols adhered to the ARRIVE guidelines[14], followed Danish legislation of animal use for scientific procedures as described in the 'Animal Testing Act' (Consolidation Act No. 726 of 9 September 1993 as amended by Act No. 1081 of 20 December 1995) and were approved by the Danish Animal Experiments Inspectorate (permission 2019-15-0201-00009).

DOCA-Salt Hypertension and Arterial Blood Pressure Measurements

DOCA-salt hypertension was induced as previously described[15].

Culture of primary endothelial cells of *Corpus cavernosum*

Endothelial cells from CC of intact C57BL/6 mice were isolated as previously described.[8]

Patch-Clamp Electrophysiology

Endothelial cells were detached from a Petri dish with trypsin-EDTA, washed, and patch-clamped using a Port-a-Patch automated system (Nanion Technologies, Munich Germany), as previously described[8].

Statistical analysis

The declared group size is the number of mice. Statistical analysis was undertaken only for experiments where each group size was at least n=5. The Western blot, CC reactivity, and immunohistochemistry results were analyzed by one- or two-way ANOVA and followed by Tukey post-test according to each analysis. The *in vivo* results were analyzed by repeated-measures two-way ANOVA and followed by Tukey post-test. The models' assumptions were investigated by inspecting Q-Q plots and data

were logarithmically transformed when necessary to generate a Gaussian-distributed data set. Relaxation responses were expressed as the percentage change from pre-contraction induced by phenylephrine. Concentration-effect curves were submitted to nonlinear regression analysis using the GraphPad Prism program (GraphPad Prism 9.0; GraphPad Software Inc., San Diego, CA, USA). Relaxant potency and maximal response were expressed as pEC_{50} (negative logarithm of molar concentration producing 50% of the maximal response) and E_{max} (maximal effect produced by the agonist), respectively. Statistical analysis of the E_{max} and pEC_{50} values was performed using nonlinear regression followed by Student's t-test or one-way ANOVA. Values of p less than 0.05 were considered statistically significant.

Results

Effect of endothelin-1 receptor inhibition on erectile function and NLRP3 inflammasome activity in DOCA/salt model of hypertension

In vivo stimulation of the cavernosal nerve in mice induced frequency-dependent increase of ICP (figure **2a** and **2b**). When mice were treated with vehicle (for bosentan), ICP/MAP was reduced in DOCA/salt mice at the frequencies from 4 to 20 Hz when compared with UNI mice (Figure **2b**). Additionally, DOCA/salt mice had gradual increment of the blood pressure from day 6 to 9, but then, it was kept sustainedly increased until day 21 (Figure **2c**). Bosentan, a non-selective endothelin-1 (ET-1) receptor antagonist, treatment restored the ICP/MAP (Figure **2b**) and systolic blood pressure (Figure **2c**) in DOCA/salt rats.

DOCA/salt model of arterial hypertension increased caspase-1, and $K_{Ca}2.3$ expression in mice CC (Figure **2d** & **2f**). No alteration to IL-1β expression was observed (Figure **2e**). Additionally, bosentan prevented caspase-1 increment in CC from DOCA/salt mice (Figure **2d**). Bosentan prevented also the $K_{Ca}2.3$ expression increment in CC of DOCA/salt mice (Figure **2f**). Furthermore, bosentan prevented pro-caspase-1 and pro-IL-1β increment in CC of DOCA/salt mice (Figure **S1a & S1b**).

DOCA/salt intervention did not modify the contractility to phenylephrine or the relaxation in response to electrical field stimulation of CC (Figure **3a** & **3b**). Bosentan did not change phenylephrine-induced contraction or the electrical field stimulation-induced relaxation of CC (**3c** & **3d**, Table **1**). However, DOCA/salt reduced endothelium-dependent and endothelium-independent relaxations to ACh and SNP,

respectively, shifting the concentration-response curves to the right (Figure **3c** & **3d**,

Table **1**). Bosentan treatment of DOCA/salt treated mice prevented these impairments

(**3c** & **3d**, Table **1**).

ET-1 inhibited the $K_{Ca}2.2/2.3$ channels in endothelial cells, impaired the erectile

function and activated the NLRP3.

The activity of $K_{Ca}2.2/2.3$ channels using whole cell voltage clamp was

measured. K^+ currents were measured in response to a voltage-step protocol. The total

membrane conductance was significantly blocked in the presence of ET-1 (Figure **4b**).

The activity of NLRP3 in endothelial cells was evaluated by Western blot of caspase-1

expression (**Figure 4c**), or IL-1β expression (**Figure 4d**). These Western blot results

suggest that ET-1 administration increased the NLRP3 activity.

The incubation of CC endothelial cells with NS13001, a $K_{Ca}2.2/2.3$ opener,

significantly reversed the inhibition of the background membrane conductance (Figure

4b) and reduced the NLRP3 activity (Figure **4c** & **4d**) promoted by ET-1. Additionally,

the incubation with apamin, a $K_{Ca}2.2/2.3$ blocker, did not alter blockage to the total

membrane conductance (Figure **4b**) and the NLRP3 activity (Figure **4c** & **4d**) promoted

by ET-1 in endothelial cells from murine CC.

Intracavernosal administration of NS13001 rescued ET-1-induced reduction in

the ICP/MAP responses to submaximal stimuli at 45, 55 and 65 min. Additionally,

apamin did not affect ET-1 impairment of ICP/MAP after its administration (Figure

S2b). ET-1 did not change ICP/MAP responses evoked by maximal stimuli (Figure

S**2a**). Additionally, NS13001 rescued ET-1-induced reduction in the ICP/MAP

responses to maximal stimuli (Figure S**2c**).

Effect of $K_{Ca}2.2/2.3$ opener on ET1-induced erectile dysfunction and NLRP3

activity in DOCA/salt model of hypertension

A selective $K_{Ca}2.2/2.3$ opener NS13001 restored erectile function (Figure **5b**)

and normalized systolic blood pressure (Figure **5c**) in DOCA/salt treated mice.

NS13001 also prevented caspase-1 increment in CC from DOCA/salt mice (Figure **5d**).

Pro-caspase-1 and pro-IL-1β expression was reduced by bosentan treatment in CC from

DOCA/salt mice (Figure **S1c** & **S2d**). Furthermore, NS13001 prevented an increase in

the $K_{Ca}2.3$ expression in CC of DOCA/salt mice (Figure **5f**). NS13001 impaired

electric-field-stimulation-induced relaxation at the frequencies 2 and 4 Hz (Figure **6b**).
NS13001 prevented the impaired CC relaxation to ACh and SNP promoted by
DOCA/salt (Figure **6c** & **6d**, Table **1**).

The $K_{Ca}2.2/2.3$ blocker apamin did not affect the ET1-induced erectile dysfunction
and NLRP3 activity in DOCA/salt model of hypertension

A selective $K_{Ca}2.2/2.3$ blocker, apamin, impaired erectile function (Figure **7b**)
and elevated the systolic blood pressure in UNI mice (Figure **7c**). Curiously, apamin
treatment restored erectile function (Figure **7b**) and normalized systolic blood pressure
(Figure **6c**) in DOCA/salt rats. Apamin also prevented caspase-1 and IL-1β increment in
CC from DOCA/salt mice (Figure **7d and e**). Pro-caspase-1 expression was reduced by
apamin treatment in CC of DOCA/salt mice (Figure **S1e** & **S1f**). Furthermore, apamin
prevented $K_{Ca}2.3$ expression increase in CC of DOCA/salt mice (Figure **7f**). Apamin
impaired electric-field-stimulation-induced relaxation at the frequencies 8, 16 and 20 Hz
(Figure **8b**, Table **1**). Apamin also prevented the impaired relaxation to acetylcholine
and sodium nitroprusside promoted by DOCA/salt treatment (Figure **8c** & **8d**, Table **1**).

Discussion

In this study, we found KCa 2.3 channels play a pivotal role in mediating ET-1-
induced NLRP3 activation and subsequent impairment in vascular and erectile function
in DOCA/salt hypertensive mice. This study shows that NS13001, $K_{Ca}2.3/K_{Ca}2.2$
channel opener, has therapeutic potential by inhibiting the ET-1 and NLRP3
inflammatory pathways and restoring erectile function. It has been shown that NS13001
has a higher affinity for $K_{Ca}2.3$, and there is no strong evidence for KCa2.2 modulation
on endothelial and erectile dysfunction[16, 17]. The increase in $K_{Ca}2.3$ expression in
hypertensive mice highlights its relevance in hypertension. Furthermore, as the ET-1
receptor antagonist, bosentan, normalized blood pressure and erectile function in
DOCA/salt mice, it underscores the central role of ET-1 in these dysfunctions, and we
suggest that it is modulated through the potassium channels. These findings offer
valuable insights into the intricate interplay among ET-1, $K_{Ca}2.2/2.3$ channels, and
NLRP3 inflammasome, suggesting potential targets for novel therapeutic interventions
for inflammatory-related diseases and the restoration of erectile function.

In accordance with our study, several previous studies reported that DOCA/salt

treatment leads to hypertension in rodents [2, 3, 18]. We found that the DOCA/salt mouse

model presented a gradual increase in blood pressure, impairment in erectile function,

and reduced relaxation to acetylcholine and sodium nitroprusside, i.e., endothelium-

dependent and endothelium-independent relaxations, respectively[15, 19]. The detrimental

effect of DOCA/salt hypertension in rodents on erectile dysfunction was reported

previously [15, 20] but the mechanism behind it was not elucidated. We found in this study

that it is associated with ET-1 signaling, leading to expression changes of the $K_{Ca}2.3$

and NLRP3 inflammasome.

ET-1 treatment leads to endothelial dysfunction, mainly by the inhibition of

various populations of potassium channels, such as inward rectifier (K_{ir})[21], sensitive

ATP (K_{ATP})[22, 23], voltage-gated (K_v)[24], and calcium-activated (K_{Ca})[9] potassium

channels. Moreover, it was already shown that ET-1 exposure releases pro-

inflammatory cytokines, like TNF-α [25] and e IL-1β, in monocytes [26] and smooth muscle

cells [27]. Accordingly, in this study, treatment with bosentan, an ET-1 type A (ET$_A$) and

B (ET$_B$) receptor antagonist[28], attenuated the increment in the $K_{Ca}2.3$, reduced the pro-

caspase-1 and caspase-1 expression in DOCA/salt mice. Furthermore, the treatment

with bosentan restored blood pressure, erectile function, and ACh and SNP-induced

relaxation, suggesting the contribution of ET-1 signaling to these abnormalities in

DOCA/salt model of hypertension. This is in line with previous reports that

administration of ET-1 antagonists reduces the elevation of blood pressure, reverts

cardiac hypertrophy and arterial remodeling [10], left ventriculus fibrosis [11, 12],

inflammation, and renal hypertrophy [12].

Accordingly, ET-1 reduces the total membrane conductance in endothelial cells

and our manipulations with the $K_{Ca}2.2/2.3$ activator and inhibitor suggest that this effect

is primarily because of the $K_{Ca}2.2/2.3$ inhibition. Importantly, the KCa2.3 is specifically

expressed in endothelial but not smooth muscle cells of corpus cavernosum (Comerma-

Steffensen et al., 2017). The KCa2.2 is also expressed in mouse corpus cavernosum, but

the role and location of the expression is less well known (Comerma-Steffensen et al., J

sex med, 2022). It has previously been shown that ET-1 leads to endothelial dysfunction

in the pulmonary artery, but mainly, this effect was mediated via inhibition of the

intermediate-conductance calcium-activated potassium channels, i.e., $K_{Ca}3.1$[9]. ET-1 was

also shown to inhibit the $K_{Ca}1.1$ in cultured smooth muscle cells from porcine coronary

artery [29]. Therefore, this is the first time the effect of ET-1 on $K_{Ca}2.3/2.2$ activity has
been reported.

Administration of the $K_{Ca}2.2/2.3$ modulator, NS13001, restored the $K_{Ca}2.2/2.3$
current in endothelial cells incubated with ET-1. Interestingly, it also reduced the
NLRP3 activity and restored the ET-1 impairment, both endothelium-dependent and
endothelium-independent relaxations. $K_{ca}2.3$ channels were suggested to be involved in
endothelium-dependent vasodilatation in human and rat intracavernous penile arteries [5,
6]. That is, the $K_{Ca}2.3$ activation regulates l-arginine uptake through the y+ transporter
system[30]. We propose that vasculature administration of l-arginine requires additional
mechanisms, such as $K_{Ca}2.3$ channels, to restore endothelium-dependent vasodilatation
[30]. According to the previous reports[8, 16], we propose that re-establishing the
endothelium-independent relaxation by NS13001 is probably a result of restoration of
both the endothelial function and the KCa1.1 channels activity.

The treatment with NS13001 also reduced the increase of pro-caspase-1,
caspase-1, and pro-IL-β expression. It can also be mediated via an anti-inflammatory
effect of the $K_{Ca}2.2/2.3$ channels, mainly, because it known to inhibit the release of
proinflammatory cytokines such as IL-6, TNF-α, and the migration of immune system
cells[31]. Interestingly, the treatment with apamin increased $K_{Ca}2.3$ expression in CC of
unilaterally nephrectomized mice with no effect in DOCA/salt hypertensive mice,
possibly because it is already upregulated. Following our suggestion about the anti-
inflammatory effect of the $K_{Ca}2.3$, apamin administration reduced pro-caspase-1,
caspase-1, pro-IL-β and IL-β expression. Here, it is possible to affirm that the increment
in the $K_{Ca}2.3$ expression correlates with the impairment of its function once the apamin
treatment increases the $K_{Ca}2.3$ expression. Additionally, apamin has an anti-
inflammatory and antioxidant role [32].

Importantly, the MCC950, a specific NLRP3 inhibitor, restores a mix of $K_{Ca}2.3$
and other ionic currents. In fact, several Ca^{2+}-channels [transient related canonic (TRC)
or potential (TRP) channels] or H^{+}-channels could also lead to K^{+} efflux from the cell,
and this mechanism could be responsible for NLRP3 activation [13]. Then, the MCC950
mechanism of NLRP3 inhibition could be related to the inhibition of these channels.
Additionally, a previous administration of MCC950 prevented the erectile dysfunction
induced by ET-1. Previous data from our lab had already shown that the genetic

deletion or pharmacological inhibition of NLRP3 prevents the impairment of

endothelium-dependent relaxation evoked by ET-1 in mice CC[33]. Indeed, ET receptors

increase intracellular $[Ca^{2+}]$ by activating Ca^{2+} influx or release in vascular smooth

muscle cells [34], liver cells and cardiomyocytes[35, 36].

We found that the inhibition of K_{Ca} 2.2/2.3 by the ET-1 possibly contributes to

the activation of NLRP3. Interestingly, although apamin did not further reduce the

K_{Ca}2.2/2.3 current in endothelial cells in the presence of ET-1 and the ACh- and SNP-

induced relaxations, it reduced the NLRP3 activity. It has previously been shown that

the endothelium-dependent vasodilator, ACh, relaxed *CC* by a mechanism sensitive to

inhibition of NO synthase, high extracellular potassium, and apamin, suggesting the

involvement of K_{Ca} 2.2/2.3 channels in CC relaxations in rats [8]. We suggest the same

mechanism is present in mouse tissue. Importantly, nitric oxide (NO) acts as a potent

inhibitor of cytokine, like TNF-α and IL-1β, production by stimulated human alveolar

macrophages [37]. This mechanism involves the nuclear factor- κB (NF-κB) blockage [38].

Accordingly, chronic treatment with apamin reduces the viral load and IL-1β, IL -2, IL -

6, and IL -10 after 9 month of therapy in type B and C hepatitis [39]. Moreover, apamin

suppresses TNF-α regulation of the NF-kB and AP-1 in chondrocytes [40]. Altogether,

these data support the involvement of K_{Ca}2.3 in ET-1-induced NLRP3 activation.

In conclusion, our study elucidates the pivotal role of KCa 2.3 channels in ET-1-

induced NLRP3 activation and subsequent impairment in vascular and erectile function

in DOCA/salt hypertensive mice. The novel findings highlight the therapeutic potential

of the KCa2.2/2.3 channel modulator, NS13001, in inhibiting the ET-1 and NLRP3

inflammatory pathways, ultimately leading to the restoration of erectile function. The

observed compensatory increase in the KCa2.3 expression in hypertensive mice

underscores its significance in hypertension. Furthermore, the efficacy of the ET-1

receptor antagonist, bosentan, in attenuating these effects emphasizes the central role of

ET-1 in inducing vascular and erectile dysfunction through the modulation of potassium

channels. Our results provide valuable insights into the complex interplay between ET-

1, KCa2.2/2.3 channels, and NLRP3 inflammasome, paving the way for potential

therapeutic interventions targeting these pathways to ameliorate inflammatory-related

diseases and restore erectile function.

Acknowledgements:

We want to express profound gratitude for the support from the funding grants

PHAB Level 3, AHA Postdoctoral fellowship (24POST1243617), the Sao Paulo

Research Foundation (FAPESP, 2016/11988-5; 2016/07641-0; 2018/05638-7;

2019/19749-8); Novo Nordisk Foundation (NFF130C0007739; NFF60C0023284) and

Wilhelm Pedersen and Wife's memorial grant.

References:

1.   Kloza M, Baranowska-Kuczko M, Malinowska B, Karpinska O, Harasim-Symbor E,
Kasacka I and Kozlowska H. The influence of DOCA-salt hypertension and chronic
administration of the FAAH inhibitor URB597 on K(Ca)2.3/K(Ca)3.1-EDH-type relaxation in rat
small mesenteric arteries. *Vascul Pharmacol*. 2017;99:65-73.
2.   Fhaner MJ, Galligan JJ and Swain GM. Increased catecholamine secretion from single
adrenal chromaffin cells in DOCA-salt hypertension is associated with potassium channel
dysfunction. *ACS Chem Neurosci*. 2013;4:1404-13.
3.   Matsumoto T, Szasz T, Tostes RC and Webb RC. Impaired beta-adrenoceptor-induced
relaxation in small mesenteric arteries from DOCA-salt hypertensive rats is due to reduced
K(Ca) channel activity. *Pharmacol Res*. 2012;65:537-45.
4.   Alexander SP, Mathie A and Peters JA. Guide to Receptors and Channels (GRAC), 5th
edition. *British journal of pharmacology*. 2011;164 Suppl 1:S1-324.
5.   Kun A, Kiraly I, Pataricza J, Marton Z, Krassoi I, Varro A, Simonsen U, Papp JG and Pajor
L. C-type natriuretic peptide hyperpolarizes and relaxes human penile resistance arteries. *J Sex
Med*. 2008;5:1114-1125.
6.   Schjorring O, Kun A, Flyvbjerg A, Kirkeby HJ, Jensen JB and Simonsen U. Flow-evoked
vasodilation is blunted in penile arteries from Zucker diabetic fatty rats. *J Sex Med*.
2012;9:1789-800.
7.   Behringer EJ and Hakim MA. Functional Interaction among K(Ca) and TRP Channels for
Cardiovascular Physiology: Modern Perspectives on Aging and Chronic Disease. *Int J Mol Sci*.
2019;20.
8.   Comerma-Steffensen SG, Carvacho I, Hedegaard ER and Simonsen U. Small and
Intermediate Calcium-Activated Potassium Channel Openers Improve Rat Endothelial and
Erectile Function. *Front Pharmacol*. 2017;8:660.
9.   Peng W, Michael JR, Hoidal JR, Karwande SV and Farrukh IS. ET-1 modulates KCa-
channel activity and arterial tension in normoxic and hypoxic human pulmonary vasculature.
Am J Physiol. 1998;275:L729-39.
10.   Li JS, Lariviere R and Schiffrin EL. Effect of a nonselective endothelin antagonist on
vascular remodeling in deoxycorticosterone acetate-salt hypertensive rats. Evidence for a role
of endothelin in vascular hypertrophy. *Hypertension*. 1994;24:183-8.
11.   Ammarguellat F, Larouche I and Schiffrin EL. Myocardial fibrosis in DOCA-salt
hypertensive rats: effect of endothelin ET(A) receptor antagonism. *Circulation*. 2001;103:319-
24.
12.   Ammarguellat FZ, Gannon PO, Amiri F and Schiffrin EL. Fibrosis, matrix
metalloproteinases, and inflammation in the heart of DOCA-salt hypertensive rats: role of
ET(A) receptors. *Hypertension*. 2002;39:679-84.
13.   Hafner-Bratkovic I and Pelegrin P. Ion homeostasis and ion channels in NLRP3
inflammasome activation and regulation. *Curr Opin Immunol*. 2018;52:8-17.
14.   McGrath JC and Lilley E. Implementing guidelines on reporting research using animals
(ARRIVE etc.): new requirements for publication in BJP. *Br J Pharmacol*. 2015;172:3189-93.

15.   Carneiro FS, Nunes KP, Giachini FR, Lima VV, Carneiro ZN, Nogueira EF, Leite R, Ergul A,
Rainey WE, Clinton Webb R and Tostes RC. Activation of the ET-1/ETA pathway contributes to
erectile dysfunction associated with mineralocorticoid hypertension. *J Sex Med*. 2008;5:2793-
807.
16.   Comerma-Steffensen S, Kun A, Hedegaard ER, Mogensen S, Aalkjaer C, Kohler R,
Monster Christensen B and Simonsen U. Down-regulation of K(Ca)2.3 channels causes erectile
dysfunction in mice. *Sci Rep*. 2017;7:3839.
17.   Kasumu AW, Hougaard C, Rode F, Jacobsen TA, Sabatier JM, Eriksen BL, Strobaek D,
Liang X, Egorova P, Vorontsova D, Christophersen P, Ronn LC and Bezprozvanny I. Selective
positive modulator of calcium-activated potassium channels exerts beneficial effects in a
mouse model of spinocerebellar ataxia type 2. *Chem Biol*. 2012;19:1340-53.
18.   Kloza M, Baranowska-Kuczko M, Malinowska B, Karpinska O, Harasim-Symbor E,
Kasacka I and Kozlowska H. The influence of DOCA-salt hypertension and chronic
administration of the FAAH inhibitor URB597 on KCa2.3/KCa3.1-EDH-type relaxation in rat
small mesenteric arteries. *Vascular pharmacology*. 2017;99:65-73.
19.   Carneiro FS, Giachini FR, Lima VV, Carneiro ZN, Nunes KP, Ergul A, Leite R, Tostes RC
and Webb RC. DOCA-salt treatment enhances responses to endothelin-1 in murine corpus
cavernosum. *Can J Physiol Pharmacol*. 2008;86:320-8.
20.   Jin L, Lagoda G, Leite R, Webb RC and Burnett AL. NADPH oxidase activation: a
mechanism of hypertension-associated erectile dysfunction. *The journal of sexual medicine*.
2008;5:544-51.
21.   Park WS, Han J, Kim N, Youm JB, Joo H, Kim HK, Ko JH and Earm YE. Endothelin-1
inhibits inward rectifier K+ channels in rabbit coronary arterial smooth muscle cells through
protein kinase C. *Journal of cardiovascular pharmacology*. 2005;46:681-9.
22.   Miyoshi Y, Nakaya Y, Wakatsuki T, Nakaya S, Fujino K, Saito K and Inoue I. Endothelin
blocks ATP-sensitive K+ channels and depolarizes smooth muscle cells of porcine coronary
artery. *Circulation research*. 1992;70:612-6.
23.   Watanuki M, Horie M, Tsuchiya K, Obayashi K and Sasayama S. Endothelin-1 inhibition
of cardiac ATP-sensitive K+ channels via pertussis-toxin-sensitive G-proteins. *Cardiovascular
research*. 1997;33:123-30.
24.   Rainbow RD, Norman RI, Everitt DE, Brignell JL, Davies NW and Standen NB.
Endothelin-I and angiotensin II inhibit arterial voltage-gated K+ channels through different
protein kinase C isoenzymes. *Cardiovascular research*. 2009;83:493-500.
25.   Browatzki M, Schmidt J, Kubler W and Kranzhofer R. Endothelin-1 induces interleukin-6
release via activation of the transcription factor NF-kappaB in human vascular smooth muscle
cells. *Basic research in cardiology*. 2000;95:98-105.
26.   Helset E, Sildnes T, Seljelid R and Konopski ZS. Endothelin-1 stimulates human
monocytes in vitro to release TNF-alpha , IL-1beta and IL-6. *Mediators of inflammation*.
1993;2:417-22.
27.   Kowalczyk A, Kleniewska P, Kolodziejczyk M, Skibska B and Goraca A. The role of
endothelin-1 and endothelin receptor antagonists in inflammatory response and sepsis. *Arch
Immunol Ther Exp (Warsz)*. 2015;63:41-52.
28.   Opitz CF, Ewert R, Kirch W and Pittrow D. Inhibition of endothelin receptors in the
treatment of pulmonary arterial hypertension: does selectivity matter? *Eur Heart J*.
2008;29:1936-48.
29.   Minami K, Hirata Y, Tokumura A, Nakaya Y and Fukuzawa K. Protein kinase C-
independent inhibition of the Ca(2+)-activated K+ channel by angiotensin II and endothelin-1.
Biochemical pharmacology. 1995;49:1051-6.
30.   Simonsen U, Winther AK, Olivan-Viguera A, Comerma-Steffensen S, Kohler R and Bek T.
Extracellular l-arginine Enhances Relaxations Induced by Opening of Calcium-Activated SKCa
Channels in Porcine Retinal Arteriole. *Int J Mol Sci*. 2019;20.

31.   Dolga AM, Letsche T, Gold M, Doti N, Bacher M, Chiamvimonvat N, Dodel R and
Culmsee C. Activation of KCNN3/SK3/K(Ca)2.3 channels attenuates enhanced calcium influx
and inflammatory cytokine production in activated microglia. *Glia*. 2012;60:2050-64.
32.   Sobral F, Sampaio A, Falcao S, Queiroz MJ, Calhelha RC, Vilas-Boas M and Ferreira IC.
Chemical characterization, antioxidant, anti-inflammatory and cytotoxic properties of bee
venom collected in Northeast Portugal. *Food and chemical toxicology : an international journal*
published for the British Industrial Biological Research Association. 2016;94:172-7.
33.   Sobrano Fais R, Menezes da Costa R, Carvalho Mendes A, Mestriner F, Comerma-
Steffensen SG, Tostes RC, Simonsen U and Silva Carneiro F. NLRP3 activation contributes to
endothelin-1-induced erectile dysfunction. *J Cell Mol Med*. 2023;27:1-14.
34.   Kawanabe Y and Nauli SM. Involvement of extracellular Ca2+ influx through voltage-
independent Ca2+ channels in endothelin-1 function. *Cellular signalling*. 2005;17:911-6.
35.   Jouneaux C, Mallat A, Serradeil-Le Gal C, Goldsmith P, Hanoune J and Lotersztajn S.
Coupling of endothelin B receptors to the calcium pump and phospholipase C via Gs and Gq in
rat liver. *The Journal of biological chemistry*. 1994;269:1845-51.
36.   Zeng Q, Li X, Zhong G, Zhang W and Sun C. Endothelin-1 induces intracellular [Ca2+]
increase via Ca 2+ influx through the L-type Ca2+ channel, Ca 2+ -induced Ca2+ release and a
pathway involving ET A receptors, PKC, PKA and AT1 receptors in cardiomyocytes. *Science in*
China Series C, Life sciences. 2009;52:360-70.
37.   Thomassen MJ, Buhrow LT, Connors MJ, Kaneko FT, Erzurum SC and Kavuru MS. Nitric
oxide inhibits inflammatory cytokine production by human alveolar macrophages. *American*
journal of respiratory cell and molecular biology. 1997;17:279-83.
38.   Raychaudhuri B, Dweik R, Connors MJ, Buhrow L, Malur A, Drazba J, Arroliga AC,
Erzurum SC, Kavuru MS and Thomassen MJ. Nitric oxide blocks nuclear factor-kappaB
activation in alveolar macrophages. *American journal of respiratory cell and molecular biology*.
1999;21:311-6.
39.   Elfiky HAGSRMHAEBMA-ZHASSEAA. Cytokine assessment and Immunomodulatory
Effect of Bee Venom in HBV&HCV Infected Patients. *Bulletin of Egyptian Society for*
Physiological Sciences. 2017;37:199-207.
40.   Jeong YJ, Shin JM, Bae YS, Cho HJ, Park KK, Choe JY, Han SM, Moon SK, Kim WJ, Choi
YH, Kim CH, Chang HW and Chang YC. Melittin has a chondroprotective effect by inhibiting
MMP-1 and MMP-8 expressions via blocking NF-kappaB and AP-1 signaling pathway in
chondrocytes. *International immunopharmacology*. 2015;25:400-5.

Figure Legends

Graphical Abstract 1. Overview of the $K_{Ca}2.2/2.3$ regulation on the ET-1-induced
NLRP3 inflammasome activation in ECs. NLRP3 inflammasome activation in ECs
depends on endothelin receptor B. On activation, NLRP3 recruits and forms a complex
with ASC as well as procaspase 1. In the final step, the assembled inflammasome
platform cleaves pro-caspase-1, and caspase-1 cleaves pro–IL-1β to activate IL-1β.
NS13001 activates KCa2.2/2.3, which inhibits ET-1-induced NLRP3 activation.
Apamin inhibits KCa2.2/2.3 opening. Bosentan directly inhibits ETB receptors in ECs,
preventing the NLRP3 inflammasome activation.

Figure 1. The endothelin receptor antagonist bosentan restores erectile function

and inhibits inflammation in DOCA/salt mice. (**a**) Original tracings showing the

intracavernosal pressure (ICP, lower traces) and mean arterial pressure (MAP, upper

traces) in response to cavernosal nerve stimulation assessed over a range of frequencies

(0.5-20 Hz) in a unilateral nephrectomized (UNI) and a DOCA/salt (DOCA)

hypertensive mice treated with vehicle or bosentan. (**b**) Average responses in ICP/MAP

and (**c**) systolic blood pressure of UNI and DOCA hypertensive mice. Immunoblot

analysis of (**d**) caspase-1, (**e**) IL-1β, and (**f**) $K_{Ca}2.3$ expression in CC strips of UNI and

DOCA hypertensive mice treated with bosentan or vehicle. The arrow indicates when

the treatment started (c). Data represents the mean ± SEM values of the groups. *

$p<0.05$. n=5-11. The comparison of each value was performed by two-way ANOVA

followed by the Holm-Šídák post-test. ICP=intracavernosal pressure; MAP=mean

arterial pressure. The entire membranes for pro-caspase-1, caspase-1, pro-IL-1β, IL-1β,

and $K_{Ca}2.3$ are shown in figure S4.

Figure 2. Bosentan restores the impaired endothelium-dependent and -

independent relaxations in corpus cavernosum of DOCA/salt mice. (**a**)

Concentration-response curves for phenylephrine, (**b**) frequency-response curves for

electrical-field-stimulation-induced relaxation, and concentration-response curves for

(c) acetylcholine and (d) sodium nitroprusside in CC strips of DOCA/salt (DOCA)

hypertensive and unilaterally nephrectomized (UNI) mice treated with bosentan or

vehicle. The values of the graphics represent mean ± SEM. * $p <0.05$. n = 6-8. The

comparison of pEC_{50} and Emax (table 1) and each value was performed by two-way

ANOVA followed by the Holm-Šídák post-test.

Figure 3. An opener of KCa2.2/2.3 channels restores the background membrane conductance and inhibits inflammation in primary endothelial cells of corpus cavernosum (CC) after endothelin-1 treatment. (**a**) Representative traces from the (**b**) Current evoked by the voltage-step protocol from −100 to +100 mV in endothelial cells in presence of vehicle or ET-1 (100 nM) and the effect of NS13001 (100 nM) or apamim (100 nM) administration. The representative traces (upper panel) and the average current-voltage relations (lower panel). Immunoblotting (upper panel) and expression (lower panel) for (**c**) caspase-1 and (**d**) IL-1β in murine CC after the administration of vehicle or ET-1 alone or combined with NS13001 and apamin. The data presented as mean ± SEM. *p ≤ 0.05. n = 3-6. The comparison of each value was performed by two-way ANOVA followed by the Holm-Šídák post-test. The entire membranes for pro-caspase-1, caspase-1, pro-IL-1β, and IL-1β are shown in figure S5.

Figure 4. **An opener of KCa2 channels, NS13001, restores erectile function and inhibits inflammation in DOCA/salt mice.** (**a**) Original tracings showing the intracavernosal pressure (ICP, lower traces) and mean arterial pressure (MAP, upper traces) in response to cavernosal nerve stimulation assessed over a range of frequencies (0.5-20 Hz) in unilateral nephrectomized (UNI) and DOCA/salt (DOCA) hypertensive mice treated with vehicle or NS13001. (**b**) Average responses in ICP/MAP and (**c**) systolic blood pressure of UNI and DOCA mice. Analysis of the (**d**) caspase-1, (**e**) IL-1β, **(f)** and $K_{Ca}2.3$ expression in CC strips from DOCA or UNI mice treated with NS13001 or vehicle. Data represent as mean ± SEM values of the groups. * p<0.05. n=5-6. The comparison of each value was performed by two-way ANOVA followed by

the Holm-Šídák post-test. ICP=intracavernosal pressure; MAP=mean arterial pressure.

The entire membranes for pro-caspase-1, caspase-1, pro-IL-1β, IL-1β, and KCa2.3 are

shown in figure S6. Figure 5 (**a to c**) ICP/MAP and Systolic Blood Pressure share the

same UNI and DOCA vehicles as in Figure 2 (**a** to **c**). Only UNI NS13001 and DOCA

NS13001 curves were statistically compared.

Figure 5. NS13001 restores the impaired endothelium-dependent and -independent

relaxations in corpus cavernosum of DOCA/salt mice. Concentration-response

curves for phenylephrine, (b) frequency-response curves for electrical-field-stimulation-

induced relaxation (b), and concentration-response curves for (c) acetylcholine and (d)

and sodium nitroprusside in CC strips of DOCA/salt (DOCA) hypertensive and

unilaterally nephrectomized (UNI) mice treated with NS13001 or vehicle. The values of

the graphics represent as mean ± SEM. * $p < 0.05$. n = 6-8. The comparison of pEC_{50}

and Emax (table 1) and each value was performed by two-way ANOVA followed by the

Holm-Šídák post-test. Figure 5 (**a to d**) curves share the same UNI and DOCA

vehicles as in Figure 3 (**a** to **d**). Only UNI NS13001 and DOCA NS13001 curves were

statistically compared.

Figure 6. A blocker of KCa2.2/2.3 channels, apamin, inhibits erectile responses in

unilaterally nephrectomized control mice and restore responses in DOCA/salt

treated mice. Original tracings showing the intracavernosal pressure (ICP, lower traces)

and mean arterial pressure (MAP, upper traces) in response to cavernosal nerve

stimulation assessed over a range of frequencies (0.5-20 Hz) in unilaterally

nephrectomized (UNI) and DOCA/salt (DOCA) hypertensive mice treated with vehicle

or apamin. (a) Average responses in ICP/MAP and (b) systolic blood pressure of UNI

and DOCA hypertensive mice. Analysis of the (c) caspase-1, (d) IL-1β, and (e) $K_{Ca}2.3$

expression in CC strips of DOCA hypertensive or UNI rats treated with apamin or

vehicle. Data represent as mean ± SEM values of the groups. * $p<0.05$. n=5-6. The

comparison of each value was performed by two-way ANOVA followed by the Holm-

Šídák post-test. ICP=intracavernosal pressure; MAP=mean arterial pressure. The entire

membranes for pro-caspase-1, caspase-1, pro-IL-1β, IL-1β, and KCa2.3 are shown in

figure S7. Figure 7 (**a** to **c**) ICP/MAP and Systolic Blood Pressure share the same

UNI and DOCA vehicles as in Figure 2 (**a** to **c**). Only UNI apamin and DOCA apamin

curves were statistically compared.

Figure 7. **Apamin inhibits relaxation in corpus cavernosum (CC) of unilaterally**

nephrectomized control mice and has no effect on responses in DOCA/salt mice.

Concentration-response curves for (**a**) phenylephrine, (**b**) frequency-response curves for

electrical-field-induced relaxation, and concentration-response curves for (**c**)

acetylcholine and (**d**) and sodium nitroprusside in CC strips of DOCA/salt (DOCA)

hypertensive and unilaterally nephrectomized (UNI) mice treated with vehicle or

apamin. The values of the graphics represent as mean ± SEM. * $p <0.05$. n = 5-6. The

comparison of pEC_{50} and Emax (table 1) and each value was performed by two-way

ANOVA followed by the Holm-Šídák post-test. Figure 8 (**a** to **d**) curves share the

same UNI and DOCA vehicles as Figure 2 (**a** to **d**). Only UNI apamin and DOCA

apamin curves were statistically compared.

Table 1. Values of Emax and pEC_{50} for the phenylephrine, acetylcholine, and sodium nitroprusside concentration-effect curves in CC of mice. Curves were performed in CC of unilaterally nephrectomized (UNI) or DOCA/salt-treated mice combined with the treatment with vehicle, bosentan, NS13001, or apamin.

Pharmacological inhibition of ET-1 on DOCA/salt-induced impairment of the CC reactivity				
Drug	**UNI**	**DOCA/salt**	**UNI bosentan**	**DOCA/salt bosentan**
phenylephrine				
pEC_{50}	5.890 ± 0.26	5.848 ± 0.30	6.134 ± 0.18	6.039 ± 0.12
Emax	633.9 ± 67.65	689.9 ± 87.56	406.1 ± 29.16	487.4 ± 22.95
acetylcholine				
pEC_{50}	7.054 ± 0.07	$6.545 \pm 0.11*$	7.143 ± 0.07	7.221 ± 0.06
Emax(%)	99.35 ± 2.95	87.68 ± 4.60	97.30 ± 2.80	95.04 ± 2.12
sodium nitroprusside				
pEC_{50}	7.180 ± 0.05	$6.583 \pm 0.05*$	7.588 ± 0.08	7.314 ± 0.06
Emax(%)	100 ± 5.21	100 ± 3.70	100 ± 2.22	100 ± 3.42

KCa2.3 activity on DOCA/salt-induced impairment in the CC reactivity				
Drug	**UNI NS13001**	**DOCA/salt NS13001**	**UNI apamin**	**DOCA/salt apamin**
phenylephrine				
pEC_{50}	6.004 ± 0.14	5.977 ± 0.16	5.877 ± 0.12	6.169 ± 0.17
Emax	473.6 ± 28.05	434.0 ± 28.99	541.7 ± 27.23	513.6 ± 33.95
acetylcholine				
pEC_{50}	7.157 ± 0.13	7.423 ± 0.08	6.828 ± 0.09	6.795 ± 0.11
Emax(%)	99.96 ± 4.86	97.33 ± 3.00	90.90 ± 2.36	89.13 ± 2.76
sodium nitroprusside				
pEC_{50}	$7.000 \pm$	$7.017 \pm$	$6.658 \pm 0.10*$	$6.569 \pm 0.10*$
Emax(%)	100 ± 3.25	100 ± 2.70	100 ± 2.21	100 ± 2.40

Values are mean $\pm$ SEM (n=5-11 in each group). * p<0.05 significantly different. The comparison of pEC_{50} and Emax parameters was performed by two-way ANOVA followed by the Holm-Šídák post-test.