A K$_{Ca}$ 2.2/2.3 Opener Reverses ET-1-induced NLRP3 Activation in Hypertensive Mice Corpora Cavernosa

Fais, RS[1,2,3]*; Comerma-Steffensen, SG[2,4]; Pinilla, E[2]; Matchkov, VV[2]; Tostes RC[1]; Carneiro, FS[1]; Simonsen, U[2]*

[1]   Department of Pharmacology, Ribeirao Preto Medical School, University of Sao Paulo, Ribeirao Preto-SP, Brazil; rsfais@gmail.com, rtostes@usp.br, fsilvac@usp.br

[2]   Department of Biomedicine, Pulmonary and Cardiovascular Pharmacology, Aarhus University, Aarhus, Denmark; us@biomed.au.dk, vvm@biomed.au.dk

[3]   Division of Pulmonary, Critical Care, and Sleep Medicine, National Jewish Health, Denver-CO, USA;

[4]   Department of Biomedical Sciences/Animal Physiology, Faculty of Veterinary, Central University of Venezuela, Venezuela. simoncomerma@biomed.au.dk.

*   Correspondence: US - us@biomed.au.dk, RSF - rsfais@gmail.com.

Abstract

Hypertension-induced erectile dysfunction is associated with endothelial dysfunction in the corpus cavernosum. Membrane depolarization activates the NLRP3 inflammasome, with downregulation of endothelial Ca^{2+}-activated K^+ channels type 2.3 (K_{Ca} 2.3) and upregulation of endothelin-1 (ET-1) linked to erectile dysfunction. However, underlying mechanisms remain incompletely understood. We hypothesized that activating K_{Ca} 2.2/2.3 channels reverses erectile dysfunction and ET-1-induced NLRP3 activation in hypertensive DOCA/salt mice. Hypertension was induced in mice using a DOCA/salt model, with unilaterally nephrectomized mice as controls. We measured blood pressure, intracavernous pressure (ICP), and corpus cavernosum (CC) contractility, and performed immunoblots for K_{Ca} 2.3, caspase-1, and interleukin-1β (IL-1β). DOCA/salt mice showed impaired erectile function and increased IL-1β activity and K_{Ca} 2.3 expression. Treatment with the endothelin receptor antagonist bosentan or the K_{Ca} 2.2/2.3 channel opener NS13001 reversed these dysfunctions and reduced ET-1-induced NLRP3 activation. NS13001 also restored decreased currents in endothelial cells exposed to ET-1. These findings establish that hypertension-induced erectile dysfunction involves an ET-1/membrane depolarization/NLRP3 inflammasome axis in corpus cavernosum endothelial cells, and that targeting endothelial K_{Ca}2.2/2.3 channels represents a promising therapeutic strategy to counteract erectile dysfunction.

Keywords: Erectile dysfunction; NLRP3 inflammasome; K_{Ca} 2.2/2.3, arterial hypertension.

1. Introduction

Erectile dysfunction (ED) commonly accompanies arterial hypertension and reflects systemic endothelial dysfunction. Penile erection is mediated by endothelium-dependent vasodilation within the corpus cavernosum (CC), which enhances cavernosal blood inflow and increases sinusoidal compliance [1-5]. A key endothelial mechanism underlying this response is membrane hyperpolarization mediated by small-conductance Ca^{2+}-activated K^+ channels ($K_{Ca}2.x$), which amplifies Ca^{2+} entry, sustains endothelial nitric oxide synthase (eNOS) activity, and promotes the release of vasodilators that relax cavernosal smooth muscle [6-8]. The K_{Ca} is a diverse group of potassium channels and are divided into small SK_{Ca} ($K_{Ca}2.1$, $K_{Ca}2.2$, and, $K_{Ca}2.3$), intermediate (IK_{Ca} or ($K_{Ca}3.1$), and large conductance Ca^{2+}-activated K^+ channels (BK_{Ca} or $K_{Ca}1.1$) [9].

ED is a predictor of cardiovascular disease, stroke, and perhaps all-cause mortality – through an augmentation in cardiovascular mortality and independently of traditional risk factors[10,11].The $K_{Ca}2.3$ channel is essential for nitric oxide (NO) signaling through endothelial NO synthase (eNOS) activation. NO produced by eNOS in the vascular bed not only mediates vasodilatation but also has anti-inflammatory functions [12,13]. The downregulation of $K_{Ca}2.3$ leads to endothelial dysfunction and ED, and expression and/or function of $K_{Ca}2.3$ channels are reduced in arterial hypertension. [6,14]

The activation of $K_{Ca}2.3$ reduces blood pressure[3] and improves erectile function in rats. Furthermore, KCa2.3 has an anti-inflammatory effect, mainly by inhibiting the release of proinflammatory cytokines such as interleukin (IL)-6 and tumor necrosis factor-alpha (TNF-α) and by inhibiting the migration of immune cells. Thus, $K_{Ca}2.3$ channel plays an essential role in the modulation of the immune system [12,13].

ET-1 is closely linked with endothelial dysfunction, mainly by the modulation of various

populations of K^+ channels, such as inward-rectifying (K_{ir}), ATP-sensitive (K_{ATP}), voltage-gated

(K_v), and K_{Ca} channels[6,15]. ET-1 has been shown to activate mononuclear phagocytes and to

promote the release of TNF-α, IL-1β, and IL-6, thereby increasing the interaction between

endothelial cells and monocytes. Moreover, a recent study showed that ET-1 activates NLRP3 via

Ca^{2+}-induced ROS generation and impairs erectile function[16-18].

Membrane depolarization is thought to be involved in the activation of NLRP3[19].

Therefore, the present study hypothesized that opening $K_{Ca}2.3$ channels prevents the activation of

NLRP3 and reverses ED. To address the hypothesis, studies were conducted on the effect of ET-

1 and an opener of KCa 2.2/2.3 channels, NS13001, on the endothelial cell K^+ currents, and the

effect of NS13001 on erectile function was examined in a DOCA/salt hypertensive mouse model.

Based on these results, this study provides a novel insight into the role of $K_{Ca}2.3$ channels in

mediating the relationship between hypertension and ED, conditions that often co-exist. By

revealing that ET-1-induced hypertension (DOCA/salt) impairs $K_{Ca}2.3$ function and activates the

NLRP3 inflammasome, leading to vascular and ED, it uncovers a previously underexplored critical

mechanism. The therapeutic potential of the $K_{Ca}2.3/K_{Ca}2.2$ channel opener NS13001 in reversing

these effects is a key finding, offering a targeted approach to treat ED and hypertension. This

timely research highlights potassium channels as promising therapeutic targets for inflammatory-

related vascular diseases and ED.

2. Materials and Methods

2.1. Animals

10-12 weeks-old C57BL/6 (~25 g) male mice were housed in a room with controlled temperature (20 - 22°C) and on light/dark cycles of 12 hrs with free access to standard chow with 0.2% sodium (Altromin 1324, Brogaarden, Denmark) and filtered water. All experimental protocols adhered to the ARRIVE guidelines [20], followed Danish legislation of animal use for scientific procedures as described in the 'Animal Testing Act' (Consolidation Act No. 726 of 9 September 1993 as amended by Act No. 1081 of 20 December 1995) and were approved by the Danish Animal Experiments Inspectorate (permission 2019-15-0201-00009).

2.2. DOCA-Salt Hypertension and Arterial Blood Pressure Measurements

Deoxycorticosterone acetate (DOCA) -salt hypertension was induced as previously described [21]. Mice were anesthetized using isoflurane (5% for induction and 2% for maintenance) in 100% oxygen (2 L/min). Briefly, mice were unilaterally nephrectomized and deoxycorticosterone-acetate (DOCA; 1 g/Kg) silastic pellets were implanted subcutaneously in the scapular region. After the DOCA surgery, the mice received buprenorphine (0.05mg/kg every 8 hours for 24 hours). DOCA mice received water containing 1% NaCl and 0.2% KCl, for 3 weeks. Unilaterally nephrectomized control mice (UNI) received silastic pellets without DOCA and tap water. Animals received either the non-selective ET-1 antagonist bosentan [30 mg/day/kg (diluted in saline) of body weight, per gavage], NS13001 [1 mg/day/kg diluted in polyethylene glycol) of body weight, per gavage], apamin [1 mg/day/kg (diluted in saline) of body weight, subcutaneous],

or its respective vehicle for 11 days starting 10 days after the surgery. Systolic blood pressure

(SBP) was measured in mice with a tail-cuff method using a RTBP1001 blood pressure system

(Kent Scientific Corporation, CT).

2.3. Isolation of corpus cavernosum (CC) strips and CC primary endothelial cells

For isolation of primary endothelial cells and corpus cavernosum (CC) strips, the mouse was

gently placed in a transparent chamber (19 cm wide x 29 cm deep x 13 cm high) and exposed to

isoflurane 3% in 100% oxygen (2L/min). Once the pain reflexes disappeared, the animals were

decapitated. After this procedure, the CC was isolated and split into 2 strips, one was used for CC

contractility measurements, and the other for immunoblotting.

2.4. Drugs and solutions

Physiological Krebs-Henseleit buffer of the following composition was used: NaCl 130

mM, KCl 4.7 mM, KH_2PO_4 1.18 mM, $MgSO_4 \cdot 7H_2O$ 1.17 mM, $NaHCO_3$ 14.9 mM, EDTA 0.026

mM, $CaCl_2 \cdot 2H_2O$ 1.6 mM and D-glucose 5.55 mM. A high concentration of KCl buffer of the

following composition was used: NaCl 14 mM, KCl 120 mM, KH_2PO_4 1.18 mM, $MgSO_4 \cdot 7H_2O$

1.17 mM, $NaHCO_3$ 14.9 mM, EDTA 0.026 mM, $CaCl_2 \cdot 2H_2O$ 1.6 mM and D-glucose 5.55 mM.

pH in all buffers was adjusted to 7.35 before its use. The incubations were performed with ET-1

(100 nM, Tocris 1160; diluted in 5% BSA + 95% deionized water), MCC950 (1 μM, Cayman

Chemical 17510; diluted in 5% DMSO and 95% deionized water), NS13001 (100 nM,

ChemShuttle, CA, USA; diluted in polyethylene glycol), apamin (100 nM, Latoxan L8407; diluted

in deionized water). The in vivo experiments were performed with bosentan (30 mg/day/kg, Sigma

Aldrich 157212-55-0; diluted in saline), NS13001 (1 mg/day/kg, ChemShuttle, CA, USA; diluted

in polyethylene glycol), apamin (1 mg/day/kg; Latoxan L8407; diluted in saline).

2.5. Culture of primary endothelial cells of Corpus cavernosum

Endothelial cells from CC of intact C57BL/6 mice were isolated as previously described[3].

Briefly, a freshly isolated CC strip was placed in sterile PBS and cut into small pieces together

with collagenase II (400 mg ml−1, Worthington Biochemical Corporation, Lakewood, NJ, USA)

for 60 min at 37°C. The sampled tissue was centrifuged at 900 RPM with isolation of the pellet

and transferred to a coated gelatin and collagen Petri dish at 37°C. The culture media was

Dulbecco's Modified Eagle Medium with 1% Fungizone, 1% penicillin streptomycin, newborn

fetal calf serum and HEPES. When reached confluence after 1–4 days, the cells were incubated

with the endothelium-specific CD102 [intracellular adhesion molecule 2 (ICAM2)] rat anti-mouse

(diluted 1:500, BD Biosciences 553326. RRID: AB_394784) for 30 min followed by incubation

with Dynabeads with sheep anti-rat IgG (Life Technologies, Oslo, Norway) for 30 min at 4°C.

Trypsin was added to separate the cells from Petri dish and transferred to dynamag 5TM (Life

Technologies, Oslo, Norway) for 2 min cleaning. The pellet was dissolved and left in a coated

Petri dish for further cultivation in endothelial cell growth serum (Provitro AG, Berlin, Germany).

The endothelial cells were treated with ET-1 (4 h) in the presence of NS13001 (30 min), apamin

(30 min), or vehicle. After the treatment, the cells were separated for patch clamp or Western blot

experiments.

2.6. Patch-Clamp Electrophysiology

Endothelial cells were detached from Petri dish with trypsin-EDTA, washed, and patch clamped using a Port-a-Patch automated system (Nanion Technologies, Munich Germany), as previously described [3]. Briefly, 3–5 MΩ resistance microchips were used. Currents were recorded using an extracellular solution containing (in mM): 140 NaCl, 4 KCl, 10 Glucose, 10 HEPES, 2 CaCl$_2$ and 1 MgCl$_2$, pH 7.3–7.4. The intracellular solution was prepared as described previously (Stankevicius et al., 2011) and contained (in mM): 30 KCl, 100 K-Aspartate, 1 mM MgCl$_2$, 10 mM EGTA, 3 Na$_2$ATP, 5 mM HEPES, and 8,5 CaCl$_2$, pH 7.2. The osmolarity of all solutions was 290–320 mOsm. All voltages were corrected for calculated junction potentials present between the internal and external solutions before seal formation. The K$^+$ currents were activated by voltage steps from −140 to 140 mV in 40 mV increments for 250 ms and using a holding membrane potential of −80 mV. Currents measured at a step of 100 mV were used to compare the effects of the K$_{Ca}$2.2/2.3 openers or blockers used in this study. All current amplitudes were expressed as current density (pA/pF) normalized to cell capacitance.

2.7. In vivo measurements of intracavernosal pressure and mean arterial pressure

Mice were anesthetized with 2% isoflurane for induction and maintained with 1.5% in 100% oxygen (2 L/min) [4]. Then, the left carotid artery and right CC of each mouse were cannulated for continuous monitoring of mean arterial pressure (MAP) and intracavernosal pressure (ICP), respectively.

In the DOCA and UNI mice treated with either vehicle, bosentan, NS13001, or apamin, the cavernosal nerve was electrically stimulated with silver electrodes at different frequencies (5 V, 1 ms pulses, and frequencies between 0.5 and 20 Hz for 60 sec stimulations) to induce changes in ICP. Each frequency was followed by a 2-minute resting period. Cavernosal nerve mediates the relaxation and contraction of cavernous smooth muscle, respectively, together regulating the erectile function of penile tissue(Dean and Lue, 2005).

In intact C57BL/6 mice, the cavernosal nerve was stimulated electrically with silver electrodes at 6 V, 1 ms pulse width, for 60 sec, and the frequency at 16 Hz to induce changes in ICP and obtain the maximum (max) ICP/MAP. Subsequently, the cavernosal nerve was stimulated at the same parameters and varying the frequency from 2 to 8 Hz to determine the 50% (submaximal) of ICP/MAP for each mouse(Sobrano Fais et al., 2023). Then, we administered intracavernosal NS13001 (0.1 µg/Kg), apamin (20.27 µg/Kg), or vehicle. A new submaximal stimulation was performed at 3, 10, 20, and 30 minutes, and the maximal stimulation at 25 minutes. A second intracavernosal administration with ET-1 (0.2492 µg/Kg) or vehicle was performed, and the submaximal stimulation was performed at 3, 10, 20, and 30 minutes, and the maximal stimulation at 25 minutes.

2.8. Cavernosal tissue contractility

Cavernosal strips from intact C57BL/6 mice were isolated and mounted in 10 mL-myograph chambers (Danish Myo Technology, Aarhus, Denmark) containing buffer continuously bubbled with a mixture of BioAir (5% CO_2 and 19.82% O_2 in nitrogen) and maintained at 37°C as previously described [2]. The tissues were stretched to a resting force of 2.5 mN and allowed to

equilibrate for 60 min. Changes in isometric force were recorded using the PowerLab/8SP data

acquisition system (Chart software, version 7; ADInstruments, Colorado Springs, CO). At the end

of the equilibration period, phenylephrine (1 μM) was added to the organ bath to verify the

preparations' contractile ability.

The contraction was evaluated by cumulative concentration-response curves to phenylephrine

(1 nM to 100 μM) in CC strips. Relaxation was evaluated by constructing cumulative

concentration-response curves for acetylcholine (ACh) (100 pM – 3 µM; endothelium-dependent

relaxation) and sodium nitroprusside (SNP) (10 pM – 100 µM; NO donor) in CC strips pre-

contracted with phenylephrine (10 µM). To investigate neurogenic relaxations in CC strips, the

preparations were treated with guanethidine (30 µM) and atropine (1 µM) as previously

described(Simonsen et al., 2001). Then, the preparations were contracted with phenylephrine, and

electrical field stimulation (0.5 to 32.0 Hz) was performed.

2.9. Immunoblotting

DOCA or UNI animals received either the non-selective ET-1 antagonist bosentan [30

mg/day/Kg (diluted in saline) of body weight, per gavage], NLRP3 inhibitor [10 mg/day/Kg

(diluted in saline) of body weight, subcutaneous], NS13001 [1 mg/day/Kg diluted in polyethylene

glycol) of body weight, per gavage], apamin [1 mg/day/kg diluted in saline) of body weight,

subcutaneous], or its respective vehicle for 11 days starting 10 days after the surgery. The samples

for this immunoblotting experiments were obtained from mice exclusively used for

immunoblotting, thereby avoiding interference from in vivo or ex vivo stimulation.

The CC endothelial cells were obtained from health mice and treated with ET-1 (100 nM,

Tocris 1160; diluted in 5% BSA + 95% deionized water), MCC950 (1 μM, Cayman Chemical

17510; diluted in 5% DMSO and 95% deionized water), NS13001 (100 nM, ChemShuttle, CA,

USA; diluted in polyethylene glycol), apamin (100 nM, Latoxan L8407; diluted in deionized

water).

The isolated CC from intact C57BL/6 mice were cleaned from surrounding fat tissue and then

snap-frozen in liquid nitrogen for storage at -80 °C. The samples were later homogenized in a lysis

buffer [50 mM Tris/HCl, 150 mM NaCl, 1% Nonidet P40, 1 mM EDTA, 1 μg/ml leupeptin, 1

μg/ml pepstatin, 1 μg/ml aprotinin, 1 mM sodium orthovanadate, 1 mM phenylmethanesulfonyl

fluoride (PMSF), and 1 mM sodium fluoride] (Fais et al., 2019). Protein concentration was

determined by the Lowry assay. Spectra multicolor broad range protein ladder (10 to 260 KDa)

was used as protein standard. Aliquots with 60 μg of proteins were prepared and separated by

electrophoresis at 100 V for 2 hours at 4° C in 4-20% Criterion TGX Stain-Free gels (bio rad) and

transferred for 1 hour to a nitrocellulose membrane at 100 V at 4° C. After transfer, total protein

was visualized using stain-free technology by exposing the membranes to UV light. Nonspecific

binding sites of the membrane to the primary antibodies were blocked with 0.3% iBlock solution

for 1 hour at room temperature. The primary antibodies described below were incubated for 12

hours at 4° C, and the secondary antibodies were incubated for 1 hour at room temperature. Protein

bands visualization was obtained by chemiluminescence after ECL reaction (Amersham ECL

Prime Western Blotting Detection Reagent) and image capture was performed on ImageQuant 350

gel imager (GE Healthcare, Piscata Way, NJ, USA). The densitometric quantification was

performed by ImageJ® software. Membranes were stripped with Restore 2x Western Blot

Stripping Buffer (Thermo) for 30 minutes at 37° C. The densitometric quantification was

normalized by the total protein in the pre-stained gel. Total protein is considered a more reliable

control than housekeeping proteins [22-24].

The following antibodies were used in the study: caspase-1 (diluted 1:500, Imgenex IMG-

5028. RRID: AB_316670), IL-1β [diluted 1:500, NovusBio NB600-633, RRID: AB_577977],

KCa2.3 (diluted 1:500, NovusBio NBP3-46424. RRID: AB_3738376), β-actin (diluted 1:1000,

Cell Signalling 5125. RRID: AB_1903890. Total protein in each lane was used as an endogenous

control for normalization of all proteins. Membranes were then incubated with the following

secondary antibodies: goat anti-mouse IgG H&L (diluted 1:10.000, ab6789, Abcam. RRID:

AB_955439) and goat anti-rabbit IgG H&L (diluted 1:10.000, ab6721, Abcam. RRID:

AB_955447).

2.10. Statistical analysis

The declared group size is the number of mice. Statistical analysis was undertaken only for

experiments where each group size was at least n=5. The Western blot, CC reactivity, and

immunohistochemistry results were analyzed by one- or two-way ANOVA and followed by Tukey

post-test according to each analysis. The in vivo results were analyzed by repeated-measures two-

way ANOVA and followed by Tukey post-test. The models' assumptions were investigated by

inspecting Q-Q plots and data were logarithmically transformed when necessary to generate a

Gaussian-distributed data set. Relaxation responses were expressed as the percentage change from

pre-contraction induced by phenylephrine. Concentration-effect curves were submitted to

nonlinear regression analysis using the GraphPad Prism program (GraphPad Prism 9.0; GraphPad

Software Inc., San Diego, CA, USA). Relaxant potency and maximal response were expressed as

pEC50 (negative logarithm of molar concentration producing 50% of the maximal response) and

Emax (maximal effect produced by the agonist), respectively. Statistical analysis of the Emax and

pEC50 values was performed using nonlinear regression followed by Student's t-test or one-way

ANOVA. Values of p less than 0.05 were considered statistically significant.

3. Results

Effect of endothelin-1 receptor inhibition on erectile function and NLRP3 inflammasome activity in DOCA/salt model of hypertension

In vivo cavernosal nerve stimulation in mice induced a frequency-dependent increase of ICP

(figure 1b and 1c). When mice were treated with the vehicle (for bosentan), ICP/MAP was reduced

in Deoxycorticosterone acetate (DOCA)/salt mice at the frequencies from 4 to 20 Hz when

compared with unilateral nephrectomized (UNI) mice (Figure 1c). Additionally, DOCA/salt mice

had a gradual increment of the blood pressure from day 6 to 9, but then, it was kept sustainedly

increased until day 21 (Figure 1d). Bosentan, a non-selective endothelin-1 (ET-1) receptor

antagonist, treatment restored the ICP/MAP (Figure 1c) and systolic blood pressure (Figure 1d) in

DOCA/salt mice.

In the mouse DOCA/salt model of arterial hypertension, there was a trend of increased

caspase-1 activity (measured by the caspase-1/pro-caspase-1 ratio) in CC (Figure 1e). IL-1β

activity (obtained by the IL-1β/pro-IL-1β) was increased in the CC from DOCA/salt mice (Figure

1f). On the other hand, $K_{Ca}2.3$ expression was reduced in DOCA/salt mice CC. Additionally,

bosentan prevented the caspase-1 activity trend, IL-1β activity increment, and $K_{Ca}2.3$ expression

reduction in CC from DOCA/salt mice (Figure 1e, 1f, and 1g). Moreover, Bosentan also prevented

the reduction in the pro-IL-1β in the CC of DOCA/salt mice (Figure 1g). No significant alterations

were observed in the expression of pro-caspase-1, caspase-1, and IL-1β (Figure S2a and S2b).

DOCA/salt intervention did not modify the contractility to phenylephrine or the relaxation in

response to electrical field stimulation of CC (Figure 2b & 2c). Bosentan did not change

phenylephrine-induced contraction or the electrical field stimulation-induced relaxation of CC (2d

& 2e, Table 1). However, DOCA/salt reduced endothelium-dependent and endothelium-

independent relaxations to ACh and SNP, respectively, shifting the concentration-response curves

to the right (Figure 2d & 2e, Table 1). Bosentan treatment of DOCA/salt mice prevented these

impairments (2d & 2e, Table 1).

300 Effect of NLRP3 inhibition on caspase-1 and IL-1β activity in DOCA/salt mice CC

In the mouse DOCA/salt model of arterial hypertension, there was an increased caspase-1 and

IL-1β activity in the CC (Figure S1b). On the other hand, $K_{Ca}2.3$ expression was reduced in

DOCA/salt mice CC. Additionally, MCC950 (NLRP3 inhibitor) prevented the caspase-1 and IL-

1β activity increment, and $K_{Ca}2.3$ expression reduction in CC from DOCA/salt mice (Figure S1a,

S1b and S1c). Furthermore, MCC950 also prevented the reduction in the pro-caspase-1 and pro-

IL-1β in the CC of DOCA/salt mice (Figure S2c, S2d). No significant alterations were observed

in the expression of caspase-1 and IL-1β (Figure S2c, S2d).

ET-1 inhibited the KCa2.2/2.3 channels in endothelial cells, impaired the erectile function and activated the NLRP3.

The activity of KCa2.2/2.3 channels using whole cell voltage clamp was measured. The K^+ currents were measured in response to a voltage-step protocol. The total membrane conductance was significantly suppressed in the presence of ET-1 (Figure 3c). The activity of NLRP3 in endothelial cells was evaluated by Western blot for caspase-1 (Figure 3d) and IL-1β activity (Figure 3e), or $K_{Ca}2.3$ expression. These Western blot results suggest that ET-1 administration increased the NLRP3 activity and reduced the KCa.2.3 expression.

The incubation of CC endothelial cells with NS13001, a KCa2.2/2.3 opener, or MCC950 (NLRP3 inhibitor) significantly reversed the ET-1-induced reduction of the background membrane conductance (Figure 3c) and reduced the NLRP3 activity (Figure 3d & 3e) and increased $K_{Ca}2.3$ (Figure 3f). Additionally, the incubation with apamin, a $K_{Ca}2.2/2.3$ blocker, did not alter blockage to the total membrane conductance (Figure 3c) and the NLRP3 activity (Figure 3d & 3e) or $K_{Ca}2.3$ expression (Figure 3f) promoted by ET-1 in endothelial cells from murine CC.

Intracavernosal administration of NS13001 rescued ET-1's reduction in the ICP/MAP responses to submaximal stimuli at 45, 55, and 65 min. Additionally, apamin did not affect ET-1's impairment of ICP/MAP after its administration (Figure S2b). ET-1 did not change the ICP/MAP responses evoked by maximal stimuli (Figure S2a). Additionally, NS13001 rescued ET-1's reduction in the ICP/MAP responses to maximal stimuli (Figure S2c).

Effect of $K_{Ca}2.2/2.3$ opener on ET1-induced erectile dysfunction and NLRP3 activity in DOCA/salt model of hypertension

A selective $K_{Ca}2.2/2.3$ opener NS13001 restored erectile function (Figure 4c) and normalized systolic blood pressure (Figure 4d) in DOCA/salt-treated mice. NS13001 also prevented the increment in the caspase-1 and IL-1β activity in the CC from DOCA/salt mice (Figure 4e and 4f). Additionally, NS13001 prevented the reduction in the $K_{Ca}2.3$ expression (Figure 4g). Moreover, pro-IL-1β expression reduction in the CC from DOCA/salt mice was prevented by NS13001 treatment (Figure S4b). NS13001 impaired electric-field-stimulation-induced relaxation at 2 and 4 Hz (Figure 5c). NS13001 prevented the impaired CC relaxation to ACh and SNP promoted by DOCA/salt (Figure 5d & 5e, Table 1).

The $K_{Ca}2.2/2.3$ blocker apamin did not affect the ET1-induced erectile dysfunction and NLRP3 activity in DOCA/salt model of hypertension

A selective $K_{Ca}2.2/2.3$ blocker, apamin, impaired erectile function (Figure 6c) and elevated the systolic blood pressure in UNI mice (Figure 6d). Curiously, apamin treatment restored erectile function (Figure 6c) and normalized systolic blood pressure (Figure 6d) in DOCA/salt mice. Apamin also prevented caspase-1 and IL-1β increment in the CC from DOCA/salt mice (Figure 6e and f). Pro-caspase-1 expression was reduced by apamin treatment in the CC of DOCA/salt mice (Figure S4c & S4d). Furthermore, apamin prevented $K_{Ca}2.3$ expression reduction in the CC of DOCA/salt mice (Figure 6g). Apamin impaired electric-field-stimulation-induced relaxation at the frequencies 8, 16 and 20 Hz (Figure 7b, Table 1). Apamin also prevented the impaired relaxation to ACh and SNP promoted by DOCA/salt treatment (Figure 7c & 7d, Table 1).

4. Discussion

In this study, we found KCa 2.3 channels play a pivotal role in counteracting ET-1-induced NLRP3 activation and subsequent impairment in vascular and erectile function in DOCA/salt hypertensive mice. This study shows that NS13001, a $K_{Ca}2.3/K_{Ca}2.2$ channel opener, has therapeutic potential by inhibiting the ET-1 and NLRP3 inflammatory pathways and restoring erectile function. It has been shown that NS13001 has a higher affinity for $K_{Ca}2.3$, and there is no strong evidence for $K_{Ca}2.2$ modulation on endothelial and erectile dysfunction[2,25]. The increase in $K_{Ca}2.3$ expression in hypertensive mice highlights its relevance in hypertension. Furthermore, as the ET-1 receptor antagonist, bosentan, normalized blood pressure and erectile function in DOCA/salt mice, it underscores the central role of ET-1 in these dysfunctions, and we suggest that it is modulated through the potassium channels.

In the previous study, we showed that NLRP3 modulates erectile function in mice[4]. In addition, it was shown that ET-1-induced erectile dysfunction and pro-inflammatory effect depends on NLRP3 activation[26]. These findings offer valuable insights into the intricate interplay among ET-1, $K_{Ca}2.2/2.3$ channels, and NLRP3 inflammasome, suggesting potential targets for novel therapeutic interventions for inflammatory-related diseases and the restoration of erectile function.

In accordance with our study, several previous studies reported that DOCA/salt treatment leads to hypertension in rodents [7,8,27]. We found that the DOCA/salt mouse model presented a gradual increase in blood pressure, impairment in erectile function, and reduced relaxation to ACh and SNP, i.e., endothelium-dependent and endothelium-independent relaxations, respectively[1,21]. The detrimental effect of DOCA/salt hypertension in rodents on erectile

dysfunction was reported previously [21,28]. However, the mechanism behind it was not elucidated. We found in this study that it is associated with ET-1 signaling, leading to changes in the expression of $K_{Ca}2.3$ and the NLRP3 inflammasome.

ET-1 treatment leads to endothelial dysfunction, mainly by reducing the function of various populations of potassium channels, such as inward rectifier (Kir)[28], sensitive ATP (KATP)[29,30], voltage-gated (Kv)[31], and calcium-activated (KCa) [15] potassium channels. Moreover, it was already shown that ET-1 exposure releases pro-inflammatory cytokines, like TNF-α [29] and e IL-1β, in monocytes [30] and smooth muscle cells [31]. Accordingly, in this study, treatment with bosentan, an ET-1 type A (ET_A) and B (ET_B) receptor antagonist[32], or MCC950, an NLRP3 inhibitor, restored the $K_{Ca}2.3$ abundance and reduced NLRP3 activity in DOCA/salt mice, suggesting that ET-1 activity in the CC of DOCA/salt mice is mediated via NLRP3 activation[4,26]. Furthermore, treatment with bosentan restored blood pressure, erectile function, and ACh- and SNP-induced relaxation, suggesting that ET-1 signaling contributes to these abnormalities in the DOCA/salt model of hypertension. This is in line with previous reports that administration of ET-1 antagonists reduces the elevation of blood pressure, reverses cardiac hypertrophy and arterial remodeling [16], left ventricular fibrosis [17,18], inflammation, and renal hypertrophy [18].

Accordingly, ET-1 reduces the total membrane conductance in endothelial cells, and our interventions with the $K_{Ca}2.2/2.3$ activator and inhibitor, or NLRP3 inhibitor suggest that this effect is primarily because of the impairment to $K_{Ca}2.2/2.3$ function and NLRP3 inhibition. Notably, the $K_{Ca}2.3$ is expressed explicitly in endothelial but not smooth muscle cells of corpus cavernosum[2]. The $K_{Ca}2.2$ is also expressed in mouse corpus cavernosum, but the role and location of the expression is less known[33] . It has previously been shown that ET-1 leads to

endothelial dysfunction in the pulmonary artery, but mainly, this effect was mediated via reduction

of the intermediate-conductance calcium-activated potassium channels, i.e., $K_{Ca}3.1$[15]. ET-1 was

also shown to inhibit the $K_{Ca}1.1$ in cultured smooth muscle cells from porcine coronary artery [34].

Therefore, this is the first time the inhibitory effect of ET-1 on $K_{Ca}2.3/2.2$ activity has been

reported in endothelial cells.

Administration of the $K_{Ca}2.2/2.3$ modulator, NS13001, or MCC950, restored the $K_{Ca}2.2/2.3$

current in endothelial cells incubated with ET-1. Interestingly, it also reduced the NLRP3 activity

and restored both endothelium-dependent and endothelium-independent relaxations impaired by

exposure to ET-1. $K_{Ca}2.3$ channels were suggested to be involved in endothelium-dependent

vasodilatation in human and rat intracavernous penile arteries [12,13]. $K_{Ca}2.3$ activation regulates

l-arginine uptake through the y+ transporter system and increased NO release[5], which will

counteract the effect of ET-1.

The treatment with NS13001 also reduced caspase-1 and pro-IL-β activity and restored $K_{Ca}2.3$

abundance in DOCA/salt mice CC. It can also be mediated via an anti-inflammatory effect of the

$K_{Ca}2.2/2.3$ channels, mainly, because it is known to inhibit the release of proinflammatory

cytokines such as IL-6, TNF-α, and the migration of immune system cells[35]. Interestingly, the

treatment with apamin reduced $K_{Ca}2.3$ expression in the CC of unilaterally nephrectomized mice

with no effect in DOCA/salt hypertensive mice, possibly because it is already downregulated.

Following our suggestion about the anti-inflammatory effect of the $K_{Ca}2.3$, apamin administration

increased caspase-1 and IL-β activity in DOCA/salt mice CC but not in UNI. Here, it is possible

to suggest that the reduction in the $K_{Ca}2.3$ expression correlates with the impairment of its function,

once the apamin treatment reduces the $K_{Ca}2.3$ expression. Additionally, apamin has a pro-

inflammatory action [36]. Curiously, apamin lowered blood pressure and increased erectile

function in DOCA/salt mice. Apamin has been previously shown to protect against atherosclerosis

by reducing lesion burden, lipid levels, inflammation, and fibrotic markers[37]. Furthermore,

modulating apamin-sensitive SK (small-conductance Ca^{2+}-activated K^+) current, which influences

action potential duration and arrhythmia in failing ventricles, selectively reduces nNOS after-

hyperpolarization, and its effects on SK channels can be modified by nNOS inhibition [38-40].

Indeed, we observed that apamin increased the CC response to non-adrenergic and non-cholinergic

stimulation, suggesting that apamin may be reducing the after-hyperpolarization of the cavernosal

nerve.

Importantly, the MCC950, a specific NLRP3 inhibitor, restores a mix of $K_{Ca}2.3$ and other

ionic currents. In fact, several Ca^{2+}-channels [transient related canonic (TRC) or potential (TRP)

channels] or H^+-channels could also lead to K^+ efflux from the cell, and this mechanism could be

responsible for NLRP3 activation [19]. Then, the MCC950 mechanism of NLRP3 inhibition could

be related to the modulation of these channels. Additionally, an administration of MCC950

prevented the erectile dysfunction induced by ET-1. Previous data from our lab had already shown

that the NLRP3 genetic deletion or pharmacological inhibition prevents the impairment of

endothelium-dependent relaxation evoked by ET-1 in mice CC[26]. Indeed, ET-1, via ET receptor

A or B increase cytosolic Ca^{2+} concentration in vascular smooth muscle cells [41], liver cells and

cardiomyocytes[42,43].

Limitations

Limitations to this study include the possibility of investigating only males. This study focused

on the vascular mechanisms of erectile dysfunction, which opens the possibility to study only

males. In addition, the ICP/MAP ratio was used to measure the erectile function under isoflurane

anesthesia. Isoflurane is known to depress the cardiovascular system not only by direct action but

also by suppression of the sympathetic nervous system **[44,45]**. This depression in sympathetic

activity is closely linked to the increased blood pressure in DOCA/salt mice. In the present study,

we cannot exclude off-target effects of the small molecules used to activate $K_{Ca}2.2/2.3$ channels

and of the endothelin receptor antagonist, bosentan. In previous studies, we examined the effect of

global NLRP3 knockdown in mice, as well as the up- and downregulation of $K_{Ca}2.3$ channels, on

erectile function [2,4]. Other genetic approaches allowing tissue-specific knockout and/or genetic

manipulation by use of Cas/CRISPR may provide additional information on the effect on the

NLRP3 inflammasome by modulation of $K_{Ca}2.3$ channels [46,47].

5. Conclusions

In conclusion, our study shows the role of $K_{Ca}2.3$ channels in ET-1-induced NLRP3 activation

and subsequent impairment in vascular and erectile function in DOCA/salt hypertensive mice. The

novel findings highlight the therapeutic potential of the $K_{Ca}2.2/2.3$ channel modulator, NS13001,

in inhibiting the ET-1 and NLRP3 inflammatory pathways, ultimately leading to the restoration of

erectile function. The observed compensatory increase in the $K_{Ca}2.3$ expression in hypertensive

mice underscores its significance in hypertension. Furthermore, the efficacy of the ET-1 receptor

antagonist, bosentan, in attenuating these effects emphasizes the central role of ET-1 in inducing

vascular and erectile dysfunction through the modulation of potassium channels. Our results

provide valuable insights into the complex interplay between ET-1, $K_{Ca}2.2/2.3$ channels, and

NLRP3 inflammatory-related diseases, paving the way for potential therapeutic interventions

targeting these pathways to ameliorate inflammatory-related diseases and restore erectile function.

Abbreviations

The following abbreviations are used in this manuscript:

Ca^{2+}	Calcium
K^{+}	Potassium
K_{Ca} 2.3	Ca2+-activated K+ channels type 2.3
ET-1	endothelin-1
ICP	intracavernous pressure
CC	corpus cavernosum
IL	interleukin
SK_{Ca}	Small conductance Ca^{2+}-activated K^{+} channels
IK_{Ca}	Intermediate conductance Ca^{2+}-activated K^{+} channels
BK_{Ca}	Large conductance Ca^{2+}-activated K^{+} channels
ED	Erectile dysfunction
NO	Nitric oxide
eNOS	Endothelial NO synthase
TNF-α	Tumor necrosis factor-alpha

K_{ir} Inward-rectifying K^+ channels

K_{ATP} ATP-sensitive K^+ channels

K_v Voltage-gated K^+ channels

DOCA   Deoxycorticosterone acetate

UNI   Unilaterally nephrectomized

SBP   Systolic blood pressure

ICAM2   Intracellular adhesion molecule 2

MAP   mean arterial pressure

ACh   Acetylcholine

SNP   Sodium nitroprusside

465 **Acknowledgments:** We want to express profound gratitude for the support from the funding
466 grants PHAB Level 3, AHA Postdoctoral fellowship (24POST1243617), the Sao Paulo Research
467 Foundation (FAPESP, 2016/11988-5; 2016/07641-0; 2018/05638-7; 2019/19749-8); Novo
468 Nordisk Foundation (NFF130C0007739; NFF60C0023284) and Wilhelm Pedersen and Wife's
469 memorial grant.

References

1. Carneiro, F.S.; Giachini, F.R.; Lima, V.V.; Carneiro, Z.N.; Nunes, K.P.; Ergul, A.; Leite, R.; Tostes, R.C.; Webb, R.C. DOCA-salt treatment enhances responses to endothelin-1 in murine corpus cavernosum. *Can J Physiol Pharmacol* **2008**, *86*, 320-328, doi:10.1139/Y08-031.

2. Comerma-Steffensen, S.; Kun, A.; Hedegaard, E.R.; Mogensen, S.; Aalkjaer, C.; Kohler, R.; Monster Christensen, B.; Simonsen, U. Down-regulation of K(Ca)2.3 channels causes erectile dysfunction in mice. *Sci Rep* **2017**, *7*, 3839, doi:10.1038/s41598-017-04188-5.

3. Comerma-Steffensen, S.G.; Carvacho, I.; Hedegaard, E.R.; Simonsen, U. Small and Intermediate Calcium-Activated Potassium Channel Openers Improve Rat Endothelial and Erectile Function. *Front Pharmacol* **2017**, *8*, 660, doi:10.3389/fphar.2017.00660.

4. Fais, R.S.; Rodrigues, F.L.; Pereira, C.A.; Mendes, A.C.; Mestriner, F.; Tostes, R.C.; Carneiro, F.S. The inflammasome NLRP3 plays a dual role on mouse corpora cavernosa relaxation. *Sci Rep* **2019**, *9*, 16224, doi:10.1038/s41598-019-52831-0.

5. Simonsen, U.; Winther, A.K.; Olivan-Viguera, A.; Comerma-Steffensen, S.; Kohler, R.; Bek, T. Extracellular l-arginine Enhances Relaxations Induced by Opening of Calcium-Activated SKCa Channels in Porcine Retinal Arteriole. *Int J Mol Sci* **2019**, *20*, doi:10.3390/ijms20082032.

6. Kloza, M.; Baranowska-Kuczko, M.; Malinowska, B.; Karpinska, O.; Harasim-Symbor, E.; Kasacka, I.; Kozlowska, H. The influence of DOCA-salt hypertension and chronic administration of the FAAH inhibitor URB597 on K(Ca)2.3/K(Ca)3.1-EDH-type relaxation in rat small mesenteric arteries. *Vascul Pharmacol* **2017**, *99*, 65-73, doi:10.1016/j.vph.2017.10.001.

7. Fhaner, M.J.; Galligan, J.J.; Swain, G.M. Increased catecholamine secretion from single adrenal chromaffin cells in DOCA-salt hypertension is associated with potassium channel dysfunction. *ACS Chem Neurosci* **2013**, *4*, 1404-1413, doi:10.1021/cn400115v.

8. Matsumoto, T.; Szasz, T.; Tostes, R.C.; Webb, R.C. Impaired beta-adrenoceptor-induced relaxation in small mesenteric arteries from DOCA-salt hypertensive rats is due to reduced K(Ca) channel activity. *Pharmacol Res* **2012**, *65*, 537-545, doi:10.1016/j.phrs.2012.02.004.

9. Alexander, S.P.; Mathie, A.; Peters, J.A. Guide to Receptors and Channels (GRAC), 5th edition. *British journal of pharmacology* **2011**, *164 Suppl 1*, S1-324, doi:10.1111/j.1476-5381.2011.01649_1.x.

10. Dong, J.Y.; Zhang, Y.H.; Qin, L.Q. Erectile dysfunction and risk of cardiovascular disease: meta-analysis of prospective cohort studies. *J Am Coll Cardiol* **2011**, *58*, 1378-1385, doi:10.1016/j.jacc.2011.06.024.

11. Jackson, G. Erectile dysfunction and cardiovascular disease. *Arab J Urol* **2013**, *11*, 212-216, doi:10.1016/j.aju.2013.03.003.

12. Kun, A.; Kiraly, I.; Pataricza, J.; Marton, Z.; Krassoi, I.; Varro, A.; Simonsen, U.; Papp, J.G.; Pajor, L. C-type natriuretic peptide hyperpolarizes and relaxes human penile resistance arteries. *J Sex Med* **2008**, *5*, 1114-1125, doi:10.1111/j.1743-6109.2008.00775.x.

13. Schjorring, O.; Kun, A.; Flyvbjerg, A.; Kirkeby, H.J.; Jensen, J.B.; Simonsen, U. Flow-evoked vasodilation is blunted in penile arteries from Zucker diabetic fatty rats. *J Sex Med* **2012**, *9*, 1789-1800, doi:10.1111/j.1743-6109.2012.02743.x.

14. Behringer, E.J.; Hakim, M.A. Functional Interaction among K(Ca) and TRP Channels for Cardiovascular Physiology: Modern Perspectives on Aging and Chronic Disease. *Int J Mol Sci* **2019**, *20*, doi:10.3390/ijms20061380.

15. Peng, W.; Michael, J.R.; Hoidal, J.R.; Karwande, S.V.; Farrukh, I.S. ET-1 modulates KCa-channel activity and arterial tension in normoxic and hypoxic human pulmonary vasculature. *Am J Physiol* **1998**, *275*, L729-739, doi:10.1152/ajplung.1998.275.4.L729.

16. Li, J.S.; Lariviere, R.; Schiffrin, E.L. Effect of a nonselective endothelin antagonist on vascular remodeling in deoxycorticosterone acetate-salt hypertensive rats. Evidence for a role of endothelin in vascular hypertrophy. *Hypertension* **1994**, *24*, 183-188, doi:10.1161/01.hyp.24.2.183.

17. Ammarguellat, F.; Larouche, I.; Schiffrin, E.L. Myocardial fibrosis in DOCA-salt hypertensive rats: effect of endothelin ET(A) receptor antagonism. *Circulation* **2001**, *103*, 319-324, doi:10.1161/01.cir.103.2.319.

18. Ammarguellat, F.Z.; Gannon, P.O.; Amiri, F.; Schiffrin, E.L. Fibrosis, matrix metalloproteinases, and inflammation in the heart of DOCA-salt hypertensive rats: role of ET(A) receptors. *Hypertension* **2002**, *39*, 679-684, doi:10.1161/hy0202.103481.

19. Hafner-Bratkovic, I.; Pelegrin, P. Ion homeostasis and ion channels in NLRP3 inflammasome activation and regulation. *Curr Opin Immunol* **2018**, *52*, 8-17, doi:10.1016/j.coi.2018.03.010.

20. McGrath, J.C.; Lilley, E. Implementing guidelines on reporting research using animals (ARRIVE etc.): new requirements for publication in BJP. *Br J Pharmacol* **2015**, *172*, 3189-3193, doi:10.1111/bph.12955.

21. Carneiro, F.S.; Nunes, K.P.; Giachini, F.R.; Lima, V.V.; Carneiro, Z.N.; Nogueira, E.F.; Leite, R.; Ergul, A.; Rainey, W.E.; Clinton Webb, R.; et al. Activation of the ET-1/ETA pathway contributes to erectile dysfunction associated with mineralocorticoid hypertension. *J Sex Med* **2008**, *5*, 2793-2807, doi:10.1111/j.1743-6109.2008.01009.x.

22. Gilda, J.E.; Gomes, A.V. Stain-Free total protein staining is a superior loading control to beta-actin for Western blots. *Anal Biochem* **2013**, *440*, 186-188, doi:10.1016/j.ab.2013.05.027.

23. Maloy, A.; Alexander, S.; Andreas, A.; Nyunoya, T.; Chandra, D. Stain-Free total-protein normalization enhances the reproducibility of Western blot data. *Anal Biochem* **2022**, *654*, 114840, doi:10.1016/j.ab.2022.114840.

24. Gurtler, A.; Kunz, N.; Gomolka, M.; Hornhardt, S.; Friedl, A.A.; McDonald, K.; Kohn, J.E.; Posch, A. Stain-Free technology as a normalization tool in Western blot analysis. *Anal Biochem* **2013**, *433*, 105-111, doi:10.1016/j.ab.2012.10.010.

25. Kasumu, A.W.; Hougaard, C.; Rode, F.; Jacobsen, T.A.; Sabatier, J.M.; Eriksen, B.L.; Strobaek, D.; Liang, X.; Egorova, P.; Vorontsova, D.; et al. Selective positive modulator of calcium-activated potassium channels exerts beneficial effects in a mouse model of spinocerebellar ataxia type 2. *Chem Biol* **2012**, *19*, 1340-1353, doi:10.1016/j.chembiol.2012.07.013.

26. Sobrano Fais, R.; Menezes da Costa, R.; Carvalho Mendes, A.; Mestriner, F.; Comerma-Steffensen, S.G.; Tostes, R.C.; Simonsen, U.; Silva Carneiro, F. NLRP3 activation contributes to endothelin-1-induced erectile dysfunction. *J Cell Mol Med* **2023**, *27*, 1-14, doi:10.1111/jcmm.17463.

27. Kloza, M.; Baranowska-Kuczko, M.; Malinowska, B.; Karpinska, O.; Harasim-Symbor, E.; Kasacka, I.; Kozlowska, H. The influence of DOCA-salt hypertension and chronic administration of the FAAH inhibitor URB597 on KCa2.3/KCa3.1-EDH-type relaxation in rat small mesenteric arteries. *Vascular pharmacology* **2017**, *99*, 65-73, doi:10.1016/j.vph.2017.10.001.

28. Jin, L.; Lagoda, G.; Leite, R.; Webb, R.C.; Burnett, A.L. NADPH oxidase activation: a mechanism of hypertension-associated erectile dysfunction. *The journal of sexual medicine* **2008**, *5*, 544-551, doi:10.1111/j.1743-6109.2007.00733.x.

29. Browatzki, M.; Schmidt, J.; Kubler, W.; Kranzhofer, R. Endothelin-1 induces interleukin-6 release via activation of the transcription factor NF-kappaB in human vascular smooth muscle cells. *Basic research in cardiology* **2000**, *95*, 98-105.

30. Helset, E.; Sildnes, T.; Seljelid, R.; Konopski, Z.S. Endothelin-1 stimulates human monocytes in vitro to release TNF-alpha , IL-1beta and IL-6. *Mediators of inflammation* **1993**, *2*, 417-422, doi:10.1155/S0962935193000596.

31. Kowalczyk, A.; Kleniewska, P.; Kolodziejczyk, M.; Skibska, B.; Goraca, A. The role of endothelin-1 and endothelin receptor antagonists in inflammatory response and sepsis. *Arch Immunol Ther Exp (Warsz)* **2015**, *63*, 41-52, doi:10.1007/s00005-014-0310-1.

32. Opitz, C.F.; Ewert, R.; Kirch, W.; Pittrow, D. Inhibition of endothelin receptors in the treatment of pulmonary arterial hypertension: does selectivity matter? *Eur Heart J* **2008**, *29*, 1936-1948, doi:10.1093/eurheartj/ehn234.

33. Comerma-Steffensen, S.; Prat-Duran, J.; Mogensen, S.; Fais, R.; Pinilla, E.; Simonsen, U. Erectile Dysfunction and Altered Contribution of KCa1.1 and KCa2.3 Channels in the

Penile Tissue of Type-2 Diabetic db/db Mice. *J Sex Med* **2022**, *19*, 697-710, doi:10.1016/j.jsxm.2022.02.021.

34. Minami, K.; Hirata, Y.; Tokumura, A.; Nakaya, Y.; Fukuzawa, K. Protein kinase C-independent inhibition of the Ca(2+)-activated K+ channel by angiotensin II and endothelin-1. *Biochemical pharmacology* **1995**, *49*, 1051-1056, doi:10.1016/0006-2952(95)98500-9.

35. Dolga, A.M.; Letsche, T.; Gold, M.; Doti, N.; Bacher, M.; Chiamvimonvat, N.; Dodel, R.; Culmsee, C. Activation of KCNN3/SK3/K(Ca)2.3 channels attenuates enhanced calcium influx and inflammatory cytokine production in activated microglia. *Glia* **2012**, *60*, 2050-2064, doi:10.1002/glia.22419.

36. Sobral, F.; Sampaio, A.; Falcao, S.; Queiroz, M.J.; Calhelha, R.C.; Vilas-Boas, M.; Ferreira, I.C. Chemical characterization, antioxidant, anti-inflammatory and cytotoxic properties of bee venom collected in Northeast Portugal. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association* **2016**, *94*, 172-177, doi:10.1016/j.fct.2016.06.008.

37. Kim, S.J.; Park, J.H.; Kim, K.H.; Lee, W.R.; Pak, S.C.; Han, S.M.; Park, K.K. The Protective Effect of Apamin on LPS/Fat-Induced Atherosclerotic Mice. *Evid Based Complement Alternat Med* **2012**, *2012*, 305454, doi:10.1155/2012/305454.

38. Koya, T.; Watanabe, M.; Natsui, H.; Kadosaka, T.; Koizumi, T.; Nakao, M.; Hagiwara, H.; Kamada, R.; Temma, T.; Anzai, T. Pharmacological nNOS inhibition modified small-conductance Ca(2+)-activated K(+) channel without altering Ca(2+) dynamics. *Am J Physiol Heart Circ Physiol* **2022**, *323*, H869-H878, doi:10.1152/ajpheart.00252.2022.

39. Selemidis, S.; Ziogas, J.; Cocks, T.M. Apamin- and nitric oxide-sensitive biphasic non-adrenergic non-cholinergic inhibitory junction potentials in the rat anococcygeus muscle. *J Physiol* **1998**, *513 (Pt 3)*, 835-844, doi:10.1111/j.1469-7793.1998.835ba.x.

40. Shuba, M.F.; Vladimirova, I.A. Effect of apamin on the electrical responses of smooth muscle to adenosine 5'-triphosphate and to non-adrenergic, non-cholinergic nerve stimulation. *Neuroscience* **1980**, *5*, 853-859, doi:10.1016/0306-4522(80)90154-2.

41. Kawanabe, Y.; Nauli, S.M. Involvement of extracellular Ca2+ influx through voltage-independent Ca2+ channels in endothelin-1 function. *Cellular signalling* **2005**, *17*, 911-916, doi:10.1016/j.cellsig.2005.01.001.

42. Jouneaux, C.; Mallat, A.; Serradeil-Le Gal, C.; Goldsmith, P.; Hanoune, J.; Lotersztajn, S. Coupling of endothelin B receptors to the calcium pump and phospholipase C via Gs and Gq in rat liver. *The Journal of biological chemistry* **1994**, *269*, 1845-1851.

43. Zeng, Q.; Li, X.; Zhong, G.; Zhang, W.; Sun, C. Endothelin-1 induces intracellular [Ca2+] increase via Ca 2+ influx through the L-type Ca2+ channel, Ca 2+ -induced Ca2+ release and a pathway involving ET A receptors, PKC, PKA and AT1 receptors in cardiomyocytes. *Sci China C Life Sci* **2009**, *52*, 360-370, doi:10.1007/s11427-009-0046-z.

620   44.   Seagard, J.L.; Hopp, F.A.; Bosnjak, Z.J.; Osborn, J.L.; Kampine, J.P. Sympathetic efferent
621   nerve activity in conscious and isoflurane-anesthetized dogs. *Anesthesiology* **1984**, *61*,
622   266-270, doi:10.1097/00000542-198409000-00006.

623   45.   Constantinides, C.; Mean, R.; Janssen, B.J. Effects of isoflurane anesthesia on the
624   cardiovascular function of the C57BL/6 mouse. *ILAR J* **2011**, *52*, e21-31.

625   46.   Gurevich, E.V.; Gurevich, V.V. Beyond traditional pharmacology: new tools and
626   approaches. *Br J Pharmacol* **2015**, *172*, 3229-3241, doi:10.1111/bph.13066.

627   47.   Mitchison, T.J. Towards a pharmacological genetics. *Chem Biol* **1994**, *1*, 3-6,
628   doi:10.1016/1074-5521(94)90034-5.

Figure Legends:

Graphic abstract. Overview of the K$_{Ca}$2.2/2.3 regulation on the ET-1-induced NLRP3 inflammasome activation in ECs. NLRP3 inflammasome activation in ECs depends on endothelin receptor B. On activation, NLRP3 recruits and forms a complex with ASC as well as procaspase 1. In the final step, the assembled inflammasome platform cleaves pro-caspase-1, and caspase-1 cleaves pro–IL-1β to activate IL-1β. NS13001 activates K$_{Ca}$2.2/2.3, which inhibits ET-1-induced NLRP3 activation. Apamin inhibits K$_{Ca}$2.2/2.3 opening. Bosentan directly inhibits ETB receptors in ECs, preventing the NLRP3 inflammasome activation.

Figure 1. The endothelin receptor antagonist bosentan restores erectile function and inhibits inflammation in DOCA/salt mice. (**a**) Experimental design. (**b**) Original tracings showing the intracavernosal pressure (ICP, lower traces) and mean arterial pressure (MAP, upper traces) in response to cavernosal nerve stimulation assessed over a range of frequencies (0.5-20 Hz) in a unilateral nephrectomized (UNI) and a DOCA/salt (DOCA) hypertensive mice treated with vehicle or bosentan. (**c**) Average responses in ICP/MAP and (**d**) systolic blood pressure of UNI and DOCA hypertensive mice. Immunoblot analysis of (**e**) caspase-1, (**f**) IL-1β, and (**g**) K$_{Ca}$2.3 expression in CC strips of UNI and DOCA hypertensive mice treated with bosentan or vehicle. The arrow indicates when the treatment started (c). Data represent the mean ± SEM values of the groups. * $p<0.05$. n=5-11. The comparison of each value was performed by two-way ANOVA followed by the Holm-Šídák post-test. ICP=intracavernosal pressure; MAP=mean arterial pressure. The entire membranes for pro-caspase-1, caspase-1, pro-IL-1β, IL-1β, and K$_{Ca}$2.3 western blots are shown in figures S7 and S8.

Figure 2. Bosentan restores the impaired endothelium-dependent and -independent relaxations in corpus cavernosum of DOCA/salt mice. (**a**) Experimental design. (**b**) Concentration-response curves for phenylephrine, (**c**) frequency-response curves for electrical-field-stimulation-induced relaxation, and concentration-response curves for (**d**) ACh and (e) SNP in CC strips of DOCA/salt (DOCA) hypertensive and unilaterally nephrectomized (UNI) mice treated with bosentan or vehicle. The values of the graphics represent mean ± SEM. * p <0.05. n = 6-8. The comparison of pEC_{50} and Emax (table 1) and each value was performed by two-way ANOVA followed by the Holm-Šídák post-test.

Figure 3. An opener of $K_{Ca}2.2/2.3$ channels restores the background membrane conductance and inhibits inflammation in primary endothelial cells of corpus cavernosum (CC) after endothelin-1 treatment. (**a**) Experimental design. (**b**) Representative traces from the (**c**) Current evoked by the voltage-step protocol from −100 to +100 mV in endothelial cells in the presence of vehicle or ET-1 (100 nM) and the effect of NS13001 (100 nM) or apamim (100 nM) administration**. The representative traces (upper panel) and the average current-voltage relations (lower panel).** Immunoblotting (upper panel) and expression (lower panel) for (**d**) caspase-1 and (**e**) IL-1β in murine *CC* after the administration of vehicle or ET-1 alone or combined with NS13001 and apamin. The data is presented as means ± SEM. *$p \leq 0.05$. n = 3-6. Each value was compared by two-way ANOVA followed by the Holm-Šídák post-test. The entire membranes for pro-caspase-1, caspase-1, pro-IL-1β, and IL-1β western blots are shown in Figure S9 and S10.

Figure 4. An opener of $K_{Ca}2$ channels, NS13001, restores erectile function and inhibits inflammation in DOCA/salt mice. (**a**) Experimental design. (**b**) Original tracings showing the intracavernosal pressure (ICP, lower traces) and mean arterial pressure (MAP, upper traces) in response to cavernosal nerve stimulation assessed over a range of frequencies (0.5-20 Hz) in unilateral nephrectomized (UNI) and DOCA/salt (DOCA) hypertensive mice treated with vehicle or NS13001. (**c**) Average responses in ICP/MAP and (**d**) systolic blood pressure of UNI and DOCA mice. Analysis of the (**e**) caspase-1, (**f**) IL-1β, **(g)** and $K_{Ca}2.3$ expression in CC strips from DOCA or UNI mice treated with NS13001 or vehicle. Data represent as mean ± SEM values of the groups. * $p<0.05$. n=5-6. The comparison of each value was performed by two-way ANOVA followed by the Holm-Šídák post-test. ICP=intracavernosal pressure; MAP=mean arterial pressure. The entire membranes for pro-caspase-1, caspase-1, pro-IL-1β, IL-1β, and $K_{Ca}2.3$ western blots are shown in figure S11 and S12.

Figure 5. NS13001 restores the impaired endothelium-dependent and -independent relaxations in corpus cavernosum of DOCA/salt mice. (**a**) Experimental design. (**b**) Concentration-response curves for phenylephrine, (**c**) frequency-response curves for electrical-field-stimulation-induced relaxation, and concentration-response curves for (**d**) ACh and (**e**) and SNP in CC strips of DOCA/salt (DOCA) hypertensive and unilaterally nephrectomized (UNI) mice treated with NS13001 or vehicle. The values of the graphics represent as mean ± SEM. * $p <0.05$. n = 6-8. The comparison of pEC_{50} and Emax (table 1) and each value was performed by two-way ANOVA followed by the Holm-Šídák post-test. Figure 5 (b to e) curves share the same UNI and DOCA vehicles as in Figure 3 (b to e).

Figure 6. A blocker of $K_{Ca}2.2/2.3$ channels, apamin, inhibits erectile responses in unilaterally nephrectomized control mice and restore responses in DOCA/salt treated mice. (**a**) Experimental design. (**b**) Original tracings showing the intracavernosal pressure (ICP, lower traces) and mean arterial pressure (MAP, upper traces) in response to cavernosal nerve stimulation assessed over a range of frequencies (0.5-20 Hz) in unilateral nephrectomized (UNI) and DOCA/salt (DOCA) hypertensive mice treated with vehicle or apamin. (**c**) Average responses in ICP/MAP and (**d**) systolic blood pressure of UNI and DOCA hypertensive mice. Analysis of the (e) caspase-1, (**f**) IL-1β, and (**g**) $K_{Ca}2.3$ expression in CC strips of DOCA hypertensive or UNI mice treated with apamin or vehicle. Data represent as mean ± SEM values of the groups. * p<0.05. n=5-6. Comparisons of each value were performed using a two-way ANOVA followed by the Holm-Šídák post hoc test. ICP=intracavernosal pressure; MAP=mean arterial pressure. The entire membranes for pro-caspase-1, caspase-1, pro-IL-1β, IL-1β, and $K_{Ca}2.3$ western blots are shown in figure S13 and 14.

Figure 7. Apamin inhibits relaxation in corpus cavernosum (CC) of unilaterally nephrectomized control mice and has no effect on responses in DOCA/salt mice. (**a**) Experimental design. (**b**) Concentration-response curves for phenylephrine, (**c**) frequency-response curves for electrical-field-stimulation-induced relaxation, and concentration-response curves for (**d**) ACh and (**e**) and SNP in CC strips of DOCA/salt (DOCA) hypertensive and unilaterally nephrectomized (UNI) mice treated with apamin or vehicle. The values of the graphics represent as mean ± SEM. * p <0.05. n = 5-6. The comparison of pEC50 and Emax (table 1) and

729 each value was performed by two-way ANOVA followed by the Holm-Šídák post-test. Figure 8

730 (a to d) curves share the same UNI and DOCA vehicles as Figure 2 (a to d). Only UNI apamin and

731 DOCA apamin curves were statistically compared.

Table 1. Pharmacological inhibition of ET-1 on DOCA/salt-induced impairment of the CC reactivity. Values of Emax and pEC_{50} for the phenylephrine, ACh, and SNP concentration-effect curves in CC of mice. Curves were performed in CC of unilaterally nephrectomized (UNI) or DOCA/salt-treated mice combined with the treatment with vehicle, bosentan or NS13001.

Pharmacological inhibition of ET-1 on DOCA/salt-induced impairment of the CC reactivity.

Drug	UNI	DOCA/salt	UNI bosentan	DOCA/salt bosentan
phenylephrine				
$pEC50$	5.89 ± 0.26	5.85 ± 0.30	6.13 ± 0.18	6.04 ± 0.12
Emax (mN)	634 ± 68	689.9 ± 87.56	406.1 ± 29.16	487.4 ± 22.95
ACh				
pEC_{50}	7.05 ± 0.07	$6.55 \pm 0.11*$	$7.14 \pm 0.07\#$	$7.22 \pm 0.06\#$
Emax(%)	99 ± 3	88 ± 5	97.30 ± 2.80	95.04 ± 2.12
Sodium nitroprusside				
pEC_{50}	7.18 ± 0.05	$6.58 \pm 0.05*$	$7.59 \pm 0.08\#$	$7.31 \pm 0.06\#$

| Emax(%) | 100 ± 5 | 100 ± 3.70 | 100 ± 2.22 | 100 ± 3.42 |

$K_{Ca}2.3$ activity on DOCA/salt-induced impairment in the CC reactivity

Drug	UNI NS13001	DOCA/salt NS13001	**UNI apamin**	**DOCA/salt apamin**
phenylephrine				
pEC_{50}	6.00 ± 0.14	5.98 ± 0.16	5.877 ± 0.12	6.169 ± 0.17
Emax (mN)	473.6 ± 28.05	434.0 ± 28.99	541.7 ± 27.23	513.6 ± 33.95
ACh				
pEC_{50}	7.16 ± 0.13 #	7.42 ± 0.08#	6.828 ± 0.09	6.795 ± 0.11
Emax(%)	100 ± 5	97.33 ± 3.00	90.90 ± 2.36	89.13 ± 2.76
Sodium nitroprusside				
pEC_{50}	7.00 ± 0.06#	7.02 ± 0.08#	6.658 ± 0.10*	6.569 ± 0.10*
Emax(%)	100 ± 3	100 ± 3	100 ± 2.21	100 ± 2.40

750 Values are mean ± SEM (n=5-11 in each group). * $p<0.05$ significantly different from unilateral

751 nephrectomized (UNI). # $p<0.05$ significantly different from DOCA/salt (DOCA). The

752   comparison of pEC_{50} and Emax parameters was performed by two-way ANOVA followed by the

753   Holm-Šídák post-test.

a Male mouse → Tail cuff day 0 — DOCA/Salt 21 Days — Tail cuff day 3, Tail cuff day 7, Tail cuff day 10, Tail cuff day 14, Tail cuff day 21 → Cavernosal tissue contractility

day 11 — Bosentan daily treatment — day 21

b

Contraction to phenylephrine (mN/g of dry tissue) vs phenylephrine Log [M]

- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI bosentan (n=6)
- DOCA bosentan (n=8)

c

Relaxation (%) vs Frequency [Hz]

- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI bosentan (n=5)
- DOCA bosentan (n=7)

d

Relaxation (%) vs acetylcholine Log [M]

- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI Bosentan (n=8)
- DOCA Bosentan (n=8)

* # #

e

Relaxation (%) vs sodium nitroprusside Log [M]

- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI bosentan (n=8)
- DOCA bosentan (n=8)

* # #

a Timeline: Tail cuff day 0 — DOCA/Salt 21 Days — NS13001 daily treatment (day 11 to day 21) — Tail cuff days 3, 7, 10, 14, 21 — Cavernosal tissue contractility

b Contraction to phenylephrine (mN/g of dry tissue) vs phenylephrine Log [M]
- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI NS13001 (n=7)
- DOCA NS13001 (n=6)

c Relaxation (%) vs Frequency [Hz]
- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI NS13001 (n=7)
- DOCA NS13001 (n=6)

d Relaxation (%) vs acetylcholine Log [M]
- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI NS13001 (n=7)
- DOCA NS13001 (n=6)

e Relaxation (%) vs sodium nitroprusside Log [M]
- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI NS13001 (n=6)
- DOCA NS13001 (n=6)

a Timeline: DOCA/Salt 21 Days. Tail cuff day 0, Tail cuff day 3, Tail cuff day 7, Tail cuff day 10, Tail cuff day 14, Tail cuff day 21. day 11 apamin daily treatment day 21. ICP/MAP and Western Blot

b Contraction to phenylephrine (mN/g of dry tissue) vs phenylephrine Log [M]
- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI apamin (n=7)
- DOCA apamin (n=6)

c Relaxation (%) vs Frequency [Hz]
- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI apamin (n=6)
- DOCA apamin (n=6)

d Relaxation (%) vs acetylcholine Log [M]
- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI apamin (n=7)
- DOCA apamin (n=6)

e Relaxation (%) vs sodium nitroprusside Log [M]
- UNI vehicle (n=6)
- DOCA vehicle (n=6)
- UNI apamin (n=7)
- DOCA apamin (n=6)