Original Article

Role of ERK2 dimerization in synaptic plasticity and memory

Santiago Ojea Ramos[1], Candela Medina[1,2,3], María del Carmen Krawczyk[2,3*], Julieta
Millan[3*], María Florencia Acutain[4*], Arturo Gabriel Romano[1,2], María Verónica Baez[4],
Francisco José Urbano[1,2], Mariano Martín Boccia[3], Mariana Feld[1#]

* Equal contribution

[1] Instituto de Fisiología, Biología Molecular y Neurociencias (IFIBYNE), Universidad de Buenos Aires
(UBA) – Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

[2] Departamento de Fisiología, Biología Molecular y Celular "Prof. Héctor Maldonado", Facultad de
Ciencias Exactas y Naturales, UBA.

[3] Laboratorio de Neurofarmacología de Procesos de Memoria, Cátedra de Farmacología, Facultad de
Farmacia y Bioquímica – UBA.

[4] Instituto de Biología Celular y Neurociencias "Prof. Dr. E. de Robertis" (IBCN), CONICET - UBA.

 # Correspondence: Mariana Feld, Email: mfeld@fbmc.fcen.uba.ar, Address: Pabellón
IFIBYNE, Ciudad Universitaria, (1428) CABA, Argentina.

ABSTRACT

Extensive research has focused on extracellular-signal regulated kinase (ERK) 1/2
phosphorylation in different memory and plasticity models. However, the precise mechanism
by which ERK activity leads to memory stabilization and restabilization remains largely elusive,
and there are no reports on the role of ERK1/2 dimerization in those processes. ERK
dimerization is critical for the binding and activation of cytoplasmic targets, many of which
have been strongly associated with these processes. In this study, several key experimental
interventions were performed. First, it was found that reactivation of a weak inhibitory
avoidance (wIA) memory led to a significant decrease in hippocampal ERK2 dimerization.
Additionally, intrahippocampal infusion of DEL-22379 (DEL), an ERK dimerization inhibitor,
after memory reactivation had a bidirectional effect: it blocked the reconsolidation of a strong
inhibitory avoidance (sIA) memory, whereas, in the case of the wIA memory, it enhanced
reconsolidation. Moreover, DEL administration blocked hippocampal ERK2 dimerization in
vivo and impaired high-frequency stimulation-induced long-term potentiation (LTP) in
hippocampal slices. Taken together, these findings reveal for the first time that ERK2
dimerization occurs in the normal environment of the rodent nervous system and plays a
critical role in plasticity and memory processes. Although more research is needed, these
initial findings suggest a relevant role of ERK dimerization in plasticity and memory.

Keywords

INTRODUCTION

Molecular mechanisms involved in learning and memory have been extensively assessed in the last decades using a wide variety of experimental paradigms and techniques. Long-term plasticity is considered the cellular correlate of memory formation[1,2]. In line with this proposal, certain signaling pathways have been associated with both plasticity and memory processes. Particularly, extracellular signal-regulated kinase 1/2 (ERK1/2) phosphorylation has been shown to subserve both of them[3–8]. Pharmacological interventions demonstrated that ERK1/2 inhibition impairs memory formation as well as neuronal plasticity[reviewed in 9]. Mitogen-activated protein kinase (MAPK) family, including the ERK1/2 cascade, is a large group of Ser/Thr protein kinases that were initially associated with development, mitogenesis, and stress response[10]. ERK activation is achieved through dual phosphorylation of residues Thr and Tyr in a common TEY motif (pERK hereafter) by a MAPK/ERK kinase (MEK), which then translates into conformational changes of the enzyme leading to substrate binding and phosphorylation. Although initial results suggested that activation of the protein by phosphorylation led to dimerization and subsequent nuclear translocation, it was later disclosed that in fact dimerization guides extra-nuclear function[11], and contributes to binding and activation of cytosolic targets aided by scaffolding proteins[12]. Consistently, the inhibition of ERK1/2 dimerization selectively disrupted cytoplasmic signaling pathways[13]. Moreover, the multiplicity of ERK1/2 cytosolic targets previously described[14–19] supports the idea that extra-nuclear localization of the kinase is relevant for a wide variety of processes. Herrero and colleagues[13] described a novel compound that inhibits ERK dimerization (DEL-22379, DEL hereafter) and blocked the growth of hyperactive Ras-ERK1/2 signaling tumor cells, without decreasing its effectiveness by resistance mechanisms previously described for phosphorylation inhibitors. Thus, preventing ERK dimerization attenuated cellular proliferation, transformation and tumor development.

The role of ERK activation in synaptic plasticity and memory processes has been extensively studied although there is limited information on specific functions in different subcellular compartments. While oncological research has significantly advanced our understanding of ERK's role in cell function, neuronal activation of this pathway is likely to exhibit distinct characteristics based on subcellular localization and kinetics, particularly in processes such as synaptic plasticity, learning, and memory.

ERK phosphorylation increased at both pre- and post-synaptic sites of pyramidal neurons of the visual cortex after in vivo visual stimulation in dark-reared rats, and its temporal profile was similar to ERK activation in neuronal cell bodies[20]. High-frequency stimulation selectively induced ERK activation, regulating translation at specific subcellular compartments at active synapses in the medial perforant pathway[21]. Moreover, NR2A-induced ERK activation has been shown to play a role in the regulation of dendritic spine density in key brain areas involved in cognition[22]. Additionally, it is associated with neurodegenerative pathologies, depending on the stage of development of the organism, the type of cellular element in which the pathway is activated, and the anatomical neuronal structure[23].

Stability of consolidated memories can be challenged by reactivation, opening a new period of sensitivity named "reconsolidation" and triggering molecular mechanisms involved in re-stabilization[24–26]. Previous work from our group revealed that cytosolic ERK activation is important for the consolidation and reconsolidation of memories in invertebrates[27] as well as

vertebrates[28,29]. ERK cytosolic kinetics following inhibitory avoidance memory reactivation in
mice regulated the reconsolidation, strength and persistence of the memory[28–30]. Moreover,
ERK overactivation in the nuclear compartment was associated with memory impairment at 6
months in a mouse model of Alzheimer's disease, as partial inhibition reverted this retention
deficit[31]. These results evidence a disregarded phenomenon which deserves further attention.

While ERK1/2 phosphorylation has been extensively studied regarding memory and plasticity,
the role of ERK1/2 dimerization in these processes remains unexplored. We hypothesized that
ERK dimerization constitutes a regulatory milestone for long-term memory (LTM)
consolidation and reconsolidation. Here we demonstrate for the first time that ERK2
dimerization can be pharmacologically inhibited in mice hippocampi without affecting its
phosphorylation. DEL administration blocked CA1 hippocampal LTP and affected memory
reconsolidation depending on the strength of the inhibitory avoidance (IA) task. This
represents the first report on the role of ERK dimerization *in vivo* within the nervous system
and its involvement in memory processes. These findings emphasize the significant role of
ERK dimerization in synaptic plasticity and memory processes.

MATERIALS AND METHODS

Animals

Different cohorts of CF-1 male mice aged either 5 weeks (weight range: 21 - 31 g kept at the
Facultad de Ciencias Exactas y Naturales central mouse house, housed in groups of 3-5
individuals in 18.41 x 29.21 x 12.7 cm amber polysulfone cages) or 8-10 weeks old (weight
range: 25 - 35 g, housed in groups of 18 - 22 individuals in stainless-steel 50 x 30 x 15 cm
cages kept at the Laboratorio de Neurofarmacología de Procesos de Memoria) were used
either for electrophysiological experiments or behavioral testing and biochemical analyses,
respectively. The mice had *ad libitum* access to dry food and tap water. They were maintained
under a 12h light/dark cycle (lights on at 6:00 AM) at a controlled temperature of 21-23ºC.
Experiments were conducted during the light cycle. All procedures related to care and
treatments of mice were approved by the local institutional animal care and use committees
(CICUAL - CUDAP # 0044975-2016, Facultad de Farmacia y Bioquímica; CICUAL # 154b,
Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Argentina), and in
compliance with the standard European Union and United States National Institutes of Health
ethical guidelines for Care and Use of Laboratory Animals. All efforts were made to minimize
any possible distress or suffering experienced by mice, as well as to reduce the number of
animals required to generate reliable scientific data.

Drugs

DEL-22379 (DEL), an ERK dimerization inhibitor[13], was purchased from MedChemExpress
(Cat. No.: HY-18932) and dissolved in dimethyl sulfoxide (DMSO) as a vehicle (VEH). For
behavioral experiments and hippocampal ERK2 dimerization assays, 0.5 µl of drug or VEH
solution were injected per hemi-hippocampus (DEL final concentration: 500 nM). Epidermal
growth-factor (EGF) was purchased from Thermo Fisher (Cat. No.: PHG0314) and diluted in
sterile PBS to a final concentration of 100 µM (stock solution).

Hippocampal slices and extracellular field potential recordings

Electrophysiological recordings in slices were performed as described elsewhere[32]. Briefly,
mice were deeply anesthetized with a mix of ketamine (160 mg/kg) and xylazine (8 mg/kg)
before decapitation. Coronal brain slices including dorsal hippocampus (300-400 µm thick)

were obtained by gluing the brain onto a vibratome stage (PELCO, EasiSlicer, Ted Pella Inc.,
CA, USA), submerged in a chamber containing chilled low-sodium/high-sucrose solution (250
mM sucrose, 2.5 mM KCl, 3 mM $MgSO_4$, 0.1 mM $CaCl_2$, 1.25 mM NaH_2PO_4, 0.4 mM ascorbic
acid, 3 mM myo-inositol, 2 mM pyruvic acid, 25 mM D-glucose, and 25 mM $NaHCO_3$).
Sequential slices were obtained in which Schaffer collaterals were transected near CA3 before
being transferred to an incubation chamber and kept at 35 °C for 30 min. Recovery chamber
was containing low Ca^{2+} / high Mg^{2+} normal artificial cerebrospinal fluid (ACSF; 125 mM NaCl,
2.5 mM KCl, 3 mM $MgSO_4$, 0.1 mM $CaCl_2$, 1.25 mM NaH_2PO_4, 0.4 mM ascorbic acid, 3 mM
myo-inositol, 2 mM pyruvic acid, 25 mM d-glucose, and 25 mM $NaHCO_3$ and aerated with 95%
O_2 / 5 % CO_2, pH 7.4). Slices were then transferred to the recording chamber and perfused
continuously with ACSF including bicuculline (20 µM; a GABA-A antagonist).
Recording electrodes were made with borosilicate glass (2–3 MΩ) filled with ACSF +
bicuculline solution and placed in the *stratum radiatum* area of the CA1 region of the
hippocampus. The field excitatory postsynaptic potentials (fEPSP) were recorded as measure
of local excitatory synaptic transmission in the CA1 region. The stimulating electrode was
placed at the Schaffer Collateral fibers and fEPSPs were evoked by test stimuli (0.033 Hz,
0.05-0.2 ms duration) every 30 sec. LTP was induced by applying high frequency stimulation
(HFS; 5 trains, 100 Hz, 10 pulses) after a recorded baseline period of 10 min. Recordings
were maintained for 30 min after LTP induction. The magnitude of LTP was calculated as a
percentage of the fEPSP slope value after HFS compared to baseline fEPSP mean slopes.
Changes in LTP were compared in hippocampal slices incubated with either DEL (30 min.,
1,33 uM) or VEH. Signals were recorded using a MultiClamp 700 amplifier commanded by
pCLAMP 10.0 software (Molecular Devices, CA, USA). Data was filtered at 5 kHz, digitized,
and stored for off-line analysis.

Intra-hippocampal injections

Hippocampal injections were performed under isoflurane anesthesia (induction: 3.5 %,
maintenance: 1 %) using a stereotaxic apparatus (Stoelting) at the indicated time points.
Surgery lasted for no more than 5 min. A small incision was made in the scalp and two holes
were drilled in the skull at the following coordinates: AP: -1.90 mm from bregma, DV: -2.2 mm
from the skull surface and +/-1.2 mm left and right from the midsagittal suture[33] to bilaterally
infuse the drugs. Injections were driven by an automatic pump (MD-1020, Bioanalytical
Systems, Inc.) at a rate of 0.52 µL/min through a 30 gauge stainless steel needle attached to
a 5 µl Hamilton syringe with PE-10 tubing. n. The drug infusion lasted 60 sec and the needle
was left in place for another 30 sec to avoid reflux. Total volume administration was 0.5 µl
(either drug or VEH solution) per hemi-hippocampus. At the end of the procedure, the incision
was sutured and thoroughly disinfected. After the last testing session, animals were
euthanized by cervical dislocation and brains were dissected, fixed by immersion in 4 % PFA
overnight, and sectioned to verify the correct injection site placement. In all cases
administration placement was confirmed in the target area.

Inhibitory Avoidance task

IA behavior was studied using a one-trial step-through protocol, which takes advantage of
mice's natural preference for dark and confined spaces[34,35]. The apparatus consists of a dark
closed compartment (20 x 20 x 15 cm) with a stainless-steel grid floor and a small illuminated
open platform (5 x 5 cm) attached to its entrance. During the training phase (TR), each mouse
was placed on the illuminated platform and received a foot shock upon entry into the dark
compartment. Two different shock intensities were used: a 0.3 mA - 3 sec shock for weak
inhibitory avoidance induction (wIA) and a 0.4 mA - 3 sec shock for strong inhibitory avoidance
(sIA) memory formation[36]. Mice underwent two retention tests after TR. The first retention test,

48 h after TR, followed by a second retention test (TS) 24 h after T1. No shock was delivered
during either test session. Previous work showed that T1 also induces memory reactivation
(RE), and this process was controlled by means of a non-reactivated (NR) group[28,37–39]. The
retention tests were finished either when the mouse stepped into the dark compartment or
failed to enter within 300 sec after the beginning of the test. After the retention test, mice were
returned to their homecage. Step through latencies were manually registered by the
experimenter using a manual chronometer.

Sample preparation and ERK analysis

Neuronal cultures. Primary neuronal cultures were obtained as described before[40] according
to Kaech and Banker´s method[41]. Briefly, Wistar embryos were obtained on day 17 (E17), both
hippocampi were dissected from pups and digested with trypsin (Sigma, Sigma–Aldrich Co.,
St. Louis, MO, United States). Cells were plated onto poly-L-lysine (Sigma)–coated wells and
incubated in neurobasal media (NB; Invitrogen, Life Technologies Corporation, Carlsbad, CA,
United States) supplemented with B27 (Invitrogen) and Glutamax (Invitrogen). Cultures were
maintained in complete NB (NB + B27 + Glutamax) at 37°C and with 5 % CO_2 and one third
of the culture media was replaced every 2 days.
For the induction of ERK activation by EGF, 1 µl of the stock solution was diluted in 1 mL of
culture medium and incubated in the culture for 5 min. Chemical LTP (cLTP) induction was
achieved by a glycine stimulation protocol previously described[42] with some modifications.
Neuronal cultures were incubated with conditioned media containing 200 mM glycine for 3
min. Glycine solution was then removed and neurons were maintained in their media for 2 min
of incubation. Control groups were incubated with conditioned media for 3 min without Glycine
stimulation.
After treatment, neuronal cultures were cooled and homogenized by scraping cells in ice-cold
lysis buffer (20 mM HEPES, pH 7.5, 10 mM ethylene glycol tetra-acetic acid, 40 mM glycerol
2-phosphate, 1 % (w/v) IGEPAL, 2.5 mM $MgCl_2$, 2 mM sodium orthovanadate, 1 mM
dithiothreitol, 0.5 mM PMSF, 50 mM Sodium Fluoride, 1 µg/ml Pepstatin A, 10 µg/mL
Leupeptin, 10 µg/mL Aprotinin) as described previously[43]. Samples were centrifuged at 13000
x g for 10 min at 4 ºC, and the supernatants were collected and conserved at -20 ºC.
205
Hippocampal ERK activation and *tissue* protein extraction

For the induction of hippocampal ERK activation by EGF, two consecutive injections were
performed in each hemi-hippocampus: first either 0.5 µl of phosphate buffer saline (VEH) or
100 µM EGF followed by 0.5 µl of either DMSO (VEH) or 400 nM DEL. At the specified time
points, mice were quickly euthanized by cervical dislocation, decapitated and the hippocampi
were dissected. Each hemi-hippocampi was then homogenized using a Dounce tight
homogenizer and 125 µL of ice-cold lysis buffer (20 mM HEPES, pH 7.5, 10 mM ethylene
glycol tetraacetic acid (EGTA), 40 mM glycerol 2-phosphate, 1 % (w/v) IGEPAL, 2.5 mM
$MgCl_2$, 2 mM orthovanadate, 1 mM dithiothreitol, 0.5 mM PMSF, 50 mM Sodium Fluoride, 1
µg/ml Pepstatin A, 10 µg/mL Leupeptin, 10 µg/mL Aprotinin) by performing 8 strokes with a
tight glass pestle. Samples were centrifuged at 13000 x g for 10 min at 4 ºC, and the cytosolic-
enriched supernatants were collected and conserved at -20 ºC.

Native gel electrophoresis

Native polyacrylamide gel electrophoresis (PAGE) for ERK dimers detection was carried out
as described previously[43]. Briefly, protein levels in lysates were quantified by BCA Assay (Bio-
Rad, USA) and 60 µg of protein were loaded into an 8 % bis-acrylamide gel and run for 150
min at 70 V. Then samples were transferred to a nitrocellulose membrane (General Electric,
USA) overnight at 125 mA and 4 ºC.

SDS-PAGE

ERK phosphorylation levels were assessed by means of denaturing gel electrophoresis as
described before[28]. Briefly, 20 µg of protein were loaded in a 12.5 % bis-acrylamide gel and
run for 90 min at 100 V. The samples were then transferred to a PVDF membrane (Thermo
Scientific, USA) overnight at 300 mA and 4 ºC and stored in TBS until probed with antibodies.

Antibodies used and detection

Initially, membranes were blocked in 4% non-fat dry milk in TBS - 0.05 % Tween-20. A full list
of antibodies used, and dilutions can be found in Table 1. Primary antibodies were incubated
overnight at 4 ºC and secondary antibodies were incubated for 1 h in TTBS at room
temperature. ERK dimers were detected on nitrocellulose membranes by luminol
chemiluminescence kit (Bio Rad ECL, USA) using X-Ray film (Agfa ORTHO CP-GU, cat.
PTNAGCPGUM2, Voxel, Argentina). SDS-PAGE blots were developed using the same
chemiluminescence kit in a GE Amersham Imager.

Table 1.

Target	Supplier	Carrier solution	Dilution
ERK2	Santa Cruz (SC-1647)	4 % milk in TBS - 0.05 % (v/v) Tween	1:1000
pERK1/2	Cell Signaling (9101)		1:2000
ERK1/2	Cell Signaling (9102)	4 % milk in TBS - 1 % (v/v) Tween	1:2000
Rabbit IgG	Santa Cruz (SC-2357)		1:10000
Mouse IgG	Cell Signaling (7076)		1:15000

Data analysis

Data analyses were conducted using R (version 4.2.2,83)[44]. Generalized linear models
(GLMs) were employed using the glmmTMB package (version 1.1.6)[45]. For behavioral data
(latency to step-through), drug (VEH or DEL) and condition (RE or NR) were used as fixed
factors. The model was fitted with a Gamma distribution and log link function. For western blot
data, the model was fitted using a Gaussian distribution and identity link function. In the LTP
experiment, Time was added as a fixed factor in order to account for repeated measures. The
goodness-of-fit for each model was analyzed using the DHARMa package (version 0.4.6)[46].
Visual inspection of residual plots did not reveal any concerning deviations from normality or
homoscedasticity. Anova of the models is reported and, when applicable, post-hoc
comparisons were performed using the emmeans package (version 1.8.5)[47]. All reported p-
values were adjusted for multiple comparisons with the Šidak correction method. Two-tailed
Pearson coefficients were calculated to determine the correlation between phosphorylation
and dimerization at each time point.

RESULTS

ERK dimerization in rodent nervous system and plasticity

The latency to enter the dark compartment in the inhibitory avoidance (IA) task depends in
part on the intensity of the aversive stimulus delivered to the animal during training[30]. Previous
results from our group using the IA task in CF-1 mice showed that reactivation of a
consolidated memory by memory retrieval induced different levels of cytosolic phosphorylated
ERK2 (pERK2) 45 min later in the hippocampus, depending on stimulus strength. Weak IA
(wIA) memory reactivation induced significant pERK2 increase compared to naïve animals,
while strong IA (sIA) memory reactivation significantly decreased it. On the other hand,
contextual exposure alone did not alter pERK2 levels at this time point[29]. Consistently,
intrahippocampal administration of PD098059, a MEK1/2 inhibitor, 45 min after wIA retrieval
strengthened memory when tested 24 h later[28].

Considering that memory reconsolidation induces significant changes in cytoplasmic ERK
activity, and that dimerization regulates its subcellular localization[11], we aimed to investigate
whether this molecular mechanism also influences memory stabilization following reactivation
in mice, specifically using the inhibitory avoidance (IA) task. The small molecule DEL-22379
(DEL hereafter) was shown to inhibit ERK dimerization process, impairing the growth of tumor
cells[13] depending on the specific driver mutation for thyroid cancer onset and progression, and
with no apparent toxicity[48]. However, ERK dimerization has yet to be reported in nervous
tissue. Consequently, our first experiment aimed to characterize ERK2 dimerization in the
rodent brain (fig. 1). To induce ERK2 dimerization, two protocols were used: 1) chemical LTP
(cLTP) induced in rat primary neuronal cultures by glycine stimulation[42] and 2)
intrahippocampal stimulation with EGF. Then, protein-enriched samples from both protocols
were resolved in native polyacrylamide gels. In the first case, cultures were pre-incubated
either with vehicle (VEH) or DEL, and then cLTP was induced with a previously characterized
protocol[42]. As positive control, other cultures were stimulated with EGF, an ERK1/2 pathway
activator that also induces its dimerization[11]. Initial assessment of data showed a tendency to
ERK2 dimerization increase after EGF stimulation (fig. 1A). Furthermore, cLTP also
upregulated ERK2 dimerization and DEL pre-treatment for 30 min completely abolished it (fig.
1A). These results qualitatively validate that ERK dimerization occurs in primary neuronal
cultures.

However, primary neuronal cultures are devoid of the tissue context existing in the mouse
hippocampus; so, we decided to evaluate dimerization *in vivo*. For this purpose, in the second
experiment, we assessed the efficacy of DEL to inhibit ERK2 dimerization in CF-1 mice
hippocampus. We acutely administered either PBS (VEH) or EGF plus either VEH or DEL
solution in each hemi-hippocampus of CF-1 anesthetized mice. Afterwards, mice were
euthanized, and we performed SDS and native gel electrophoresis of protein-enriched
hippocampal samples and western blot. A schematic experimental design is shown on top of
figure 1B. This administration scheme was used in order to perform experimental and control
treatments in the same animal. Consistent with previous reports[49–52], EGF administration
altered hippocampal pERK2 levels 5 (fig. 1B left; $F_{2,13}$ = 5.381, p = 0.048) and 15 min later (fig.
1B right; $F_{2,13}$ = 8.582, p = 0,0100). EGF administration induced a rapid and transient significant
increase in hippocampal pERK2 at 5 min (p = 0.0492) which decreased under VEH baseline
levels at 15 min (p = 0.0100). It is important to point out that DEL treatment did not affect EGF-
induced pERK2 profile at the time points assessed (p = 0.989 at 5 min; p = 0.555 at 15 min in
EGF-VEH vs EGF-DEL contrasts), which were also statistically different from control (p =
0,036 at 5 min; p = 0,009 at 15 min in VEH-VEH vs DEL-EGF contrasts). However, ERK2
dimerization (fig. 1C) showed different kinetics after EGF administration: it did not change at

5 min (fig. 1C left; $F_{2,13}$ = 0.2703; p = 0.7694), but it was significantly altered at 15 min (fig. 1C
right; $F_{2,13}$ = 11.623; p = 0.0044). Consistent with previous results[13], EGF alone induced a
significant rise in ERK2 dimerization (p = 0.014). This effect was prevented by DEL pre-
treatment (p = 0.030) which induced dimer/monomer (dim/mon) levels comparable to VEH-
treated hippocampi (p = 0.915). Therefore, phosphorylation and dimerization of ERK2 showed
a strong negative correlation 15 min after stimulation (fig. 1D, p = 0.049). Thus, based on
these findings, it appears that under our experimental conditions, DEL prevents ERK2
dimerization induced by EGF in CF-1 mice hippocampi without affecting its phosphorylation.

Next, we wondered whether the inhibition of ERK2 dimerization has an impact on hippocampal
plasticity. In order to answer this question, we performed acute LTP experiments on
hippocampal slices. Slices were pre-incubated for 30 min in the presence of VEH or DEL, a
10 min baseline fEPSP was recorded to then extracellularly deliver a HFS protocol on Schaffer
Collaterals (5 trains, 100 Hz, 10 pulses). Neither fEPSP baseline recordings nor immediate
post-HFS fEPSPs were altered by DEL treatment (fig. 1E). However, around 15 min after HFS
application, DEL-treated slices begin to diverge from VEH-treated ones, reaching field
potential levels similar to baseline. In this last part of the experiment, fEPSP recordings were
significantly different between treatments (p = 1.44 x 10^{-10}).

Figure 1. In vivo ERK2 dimerization in the rodent nervous system. **A)** Native PAGE western blot showing ERK2 dimers (dim) and monomers (mon) in samples from rat neuronal primary cultures subjected to either cLTP or EGF stimulation and-or DEL treatment (left). Control and EGF-stimulated HEK-293T culture cells and CF-1 mice hippocampal protein homogenates (right). **B-C)** ERK2 phosphorylation precedes dimerization in EGF-stimulated CF-1 mice hippocampal tissue. Schematic experimental design is shown on **top** of fig. B). Hippocampal ERK2 phosphorylation **(B)** and dimerization **(C)** were assessed 5 (left) or 15 (right) min after *in vivo* local EGF plus VEH or DEL administration. Dimer / monomer (dim/mon) ratios were normalized against "veh+veh" mean value in each case. Circles represent individual mice measurements, single lines represent mean ± s.d., and estimation plots shown on the right of each graph represent the difference between means (black dot) and the bootstrapped distribution of means (grey area) with 95% confidence intervals. *****, p < 0.05; ******, p < 0.01 in Tukey comparisons. n = 4 mice / group. **D)** Pearson correlation between ERK2 phosphorylation and dimerization assessed at 15 min in figures B and C. $R^2 = 0.71$; p < 0.05. **E)** DEL pre-treatment impairs LTP in CA1 hippocampal acute slices. Circles

represent the mean of 4 replicates and single lines represent s.d. Three points of interest are indicated throughout
the recording: immediately before stimulation (i), immediately after stimulation (ii), and approx. 18-20 min after
stimulation (iii). Bars in i), ii) and iii) are the mean ± s.d. of 100 sec recordings. ***, $p < 0.05$ (repeated measures
comparison between VEH- and DEL-treated groups). n = 4 mice / group.

Memory reactivation after Weak IA decreases hippocampal ERK dimerization

Considering that ERK dimerization can modulate hippocampal plasticity, and that IA memory
reconsolidation is fine-tuned by cytoplasmic ERK signaling at 45 min[28], we aimed to
investigate whether its dimerization is regulated by memory reactivation in the hippocampus
at this time point. Therefore, we trained animals in either strong or weak versions of the IA
task and subjected them to memory reactivation (RE) 48 h later. As a control, identically
trained animals were not exposed to reactivation (NR). Animals were euthanized 45 min after
reactivation or the corresponding time for NR groups and their hippocampi were dissected in
order to perform native polyacrylamide gel electrophoresis and ERK dimerization assessment
(fig. 2, scheme).

Figure 2. Hippocampal ERK2 dimerization after memory reactivation. **Top)** Schematic experimental design for
hippocampal ER2 dimerization assessment. **Bottom)** Hippocampal ERK2 dimerization assessment by native
PAGE western blots 45 after either sIA **(A)** or wIA **(B)** memory reactivation. Dim/mon ratios are normalized against
nr mean value for each experiment. Color circles indicate individual values and error bars shown on the right of
each set depict mean ± s.d. values. Estimation plots shown on the right of each graph represent the difference
between means (black dot) and the bootstrapped distribution of means (grey area) with 95% confidence intervals.
Representative western blots are also shown (right). *, $p < 0.05$; **, $p < 0.01$ in Tukey comparisons. n = 7-9 / group.

Reexposure to the IA context 48 h after strong TR did not induce evident changes in ERK2
dimerization levels related to NR control animals 45 min later (fig. 2A; t_{13} = 0.161; p = 0.873).
Meanwhile, wIA memory reactivation showed significant decrease in ERK2 dimerization
relative to the NR control group (fig. 2B; t_{14} = 2.569; p = 0.0223). The observation that memory
reactivation induced differential ERK2 dimerization depending on TR strength suggests the
possibility that this molecular mechanism plays a role in memory reconsolidation, as well.

Post-retrieval hippocampal ERK2 dimerization supports strong IA memories

Given that ERK2 dimerization was differentially regulated upon memory reactivation, we
sought to determine whether memory reconsolidation, a process triggered by reactivation, was
affected by its inhibition. In consequence, we first evaluated whether ERK2 dimerization was
affected by intrahippocampal DEL administration after memory reactivation. For this
experiment, all animals were injected either VEH or DEL in both hippocampi 35 min after RE
or the corresponding time for NR animals (fig. 3A, upper scheme) and 10 min later animals
were euthanized and their hippocampi were removed and homogenized. As shown in figure
3A (lower panel), reactivation did not induce changes in ERK2 dimerization levels ($F_{2,21}$ =
5.559; p = 0.0115; p = 0.336 in NR-VEH vs RE-VEH contrast). However, hippocampal DEL
injection after reactivation significantly reduced ERK2 dimerization (p = 0.0102 in RE-DEL vs
NR-VEH contrast), similar to the levels observed following wIA memory reactivation.
Next, we asked whether basal ERK2 dimerization levels are necessary for strong IA memory
re-stabilization. We aimed to answer this question by administering VEH or DEL 40 min after
strong IA memory reactivation and tested retention 24 h later (fig. 3B, upper panel). All groups
performed similarly during training and reactivated groups expressed strong retention during
reexposure to the training context (fig. 3B, lower panel). However, there was a significant
interaction between reactivation and drug treatment on latencies during the test session ($F_{2,28}$=
5.291; p = 0.029). Thus, while latencies from VEH- and DEL-injected NR animals did not differ
statistically (p = 0.9833), DEL treatment after memory reactivation induced a significant
decrease in latencies compared to the VEH group (p = 0,0287). General response levels
during TS were similar, as latencies from VEH-injected groups were not statistically different
(p = 0.6847). Furthermore, this impairment lasted for at least a week (suppl. fig. 1; p = 0.013).
This long-lasting impairment suggests a specific effect on the reconsolidation process. In
consequence, basal ERK2 dimerization levels would be required for strong IA memory
reconsolidation.

Figure 3. Role of hippocampal ERK2 dimerization after strong IA memory reactivation. **A)** DEL decreases hippocampal ERK2 dimerization after strong IA memory reactivation. Schematic experimental design for hippocampal ERK2 dimerization assessment is shown **(top)**. Hippocampal ERK2 dimerization was assessed 45 min after strong IA memory reactivation and 10 min after VEH or DEL intra-hippocampal administration. Dimer to monomer ratios were normalized against nr mean value for each experiment. Color circles indicate individual values and error bars shown on the right of each dataset depict mean ± s.d. values. Estimation plots shown on the right of each graph represent the difference between means (black dot) and the bootstrapped distribution of means (grey area) with 95% confidence intervals. Representative western blots are also shown (right). *, $p < 0.05$; **, $p < 0.01$ in Tukey adjusted *post hoc* comparisons. n = 8 / group. **B)** Inhibition of hippocampal ERK2 dimerization after strong

400 IA memory reactivation impairs LTM. Schematic experimental design is shown **(top)**. Color circles indicate
401 individual latencies to step through values and error bars shown on the right of each set depict median ± inter
402 quartile range (IQR). Estimation plots shown on the right of ts graph represent the difference between means (black
403 dot) and the bootstrapped distribution of means (grey area) with 95% confidence intervals. *****, p < 0.05; ******, p < 0.01
404 in Tukey adjusted *post hoc* comparisons. n = 8 / group.
405

Inhibiting hippocampal ERK dimerization after reactivation strengthens weak IA memories

We showed that wIA memory reactivation induced a decrease in ERK2 dim/mon ratio in mice
hippocampi (fig. 2B). Consequently, we wondered whether intra-hippocampal DEL
administration after wIA reactivation would have any consequence on LTM retention. We
trained 4 groups of animals in a wIA task and submitted two of them to memory reactivation
(RE) 48 h later. Performance during training was similar in all groups. As expected, RE groups
expressed median latencies of ≈150 sec during the reactivation session. One NR and one RE
groups of animals were administered VEH and the other two groups were injected DEL in the
dorsal hippocampus 40 min after the reactivation session. All groups were tested 24 h later
(fig. 4A). Latencies to step through during the test session exhibited a significant interaction
between reactivation procedure and drug treatment ($F_{2,30}$ = 13.254; p = 0.001). Surprisingly,
in RE-animals, the DEL-injected group showed significantly higher latencies than the VEH-
injected one (p = 0.005), and this effect was specific for the reactivated groups as NR animals
did not show significant differences in latencies between VEH and DEL treatment during test
(p = 0.379). Moreover, VEH-treated groups did not differ statistically in their latencies during
the test session (p = 0.273). Not only did intrahippocampal DEL administration after wIA
memory reactivation improve retention, but it also seemed to have drastically increased the
proportion of animals exhibiting higher latencies during TS compared to RE (suppl. fig.2, right
panel). Notably, a significant positive correlation was observed between RE and TS response
levels in DEL-RE animals (suppl. fig. 2; R^2 = 0.73; p = 0.025), which differed from the VEH-
RE group (suppl. fig. 2; R^2 = 0.42; p = 0.270).

A

B

Figure 4. Role of hippocampal ERK2 dimerization after wIA memory reactivation. **A)** Inhibition of hippocampal
ERK2 dimerization after strong IA memory reactivation improves LTM. Schematic experimental design is shown
on top. Color circles indicate individual latencies to step through values during tr (left), re (center) and ts (right).
Error bars shown on the right of each dataset depict median ± IQR. Estimation plots shown on the right of ts graph
represent the difference between means (black dot) and the bootstrapped distribution of means (grey area) with
95% confidence intervals. *, p < 0.05; **, p < 0.01 in Tukey adjusted *post hoc* comparisons. n = 8 - 9 / group. **B)**
Hippocampal ERK2 dimerization decreases after wIA memory reactivation. Schematic experimental design is
shown on top. Color circles indicate individual values and error bars shown on the right of each dataset depict
mean ± s.d. values. Estimation plots shown on the right of each graph represent the difference between means

(black dot) and the bootstrapped distribution of means (grey area) with 95% confidence intervals. Representative
western blots are also shown (right). *, $p < 0.05$; **, $p < 0.01$ in Tukey adjusted *post hoc* comparisons. n = 8 / group.

Finally, ERK2 dimerization levels were significantly affected by reactivation also in VEH/DEL-
injected animals as well ($F_{2,21}$ = 13.254; p = 0.0002). In contrast to what we observed after sIA
training, wIA memory reactivation led to a significant decrease in ERK2 dimerization (see fig.
2B) which was also observed after VEH-administration in mice hippocampi (fig. 4B; p =
0.0064). However, DEL injection did not significantly affect ERK2 dimerization levels following
memory reactivation (p = 0.3753), though these levels were significantly different from those
in the control group NR-VEH (p = 0.0003).

DISCUSSION

In this work, we show for the first time that ERK2 dimerization constitutes a key regulatory step
involved in plasticity and memory processes. Several lines of evidence indicate that
subcellular localization of a particular intracellular signaling pathway activity is crucial for a
certain biological outcome[12,53,54]. In the central nervous system, for example, it was shown that
NF-kappaB might be regulating different phenomena related to neural plasticity, memory and
learning processes depending on its subcellular localization[55]. Over the past decade,
oncological research has advanced the development of drugs targeting key pathways to
reverse pathological conditions. Interestingly, many of these pathways have also been
implicated in learning and memory processes. However, their regulatory mechanisms within
the nervous system often remain poorly understood. ERK signaling serves as a notable
example. ERK phosphorylation is well known to be essential for neural plasticity and cognitive
processes, and its deregulation often leads to pathological conditions[9]. Interestingly, ERK
dimerization has been proposed to regulate the subcellular location of its activity, thereby
influencing the phosphorylation of specific targets and ultimately the cellular fate[12,13]. However,
despite extensive studies on ERK activation in neural tissue[reviewed in 9], the functional role of
ERK dimerization in the nervous system remains largely unexplored.

Consequently, as an initial step, we demonstrated that ERK dimerization occurs in neural
tissue, both *in vitro* using primary neuronal cultures and *in vivo* within the hippocampus of CF-
1 mice. Primary neuronal cultures stimulated with either EGF or a glycine-based cLTP protocol
showed increased ERK2 dimerization which was inhibited by DEL pre-treatment (fig. 1A).
Moreover, upon EGF stimulation, DEL treatment inhibited *in vivo* hippocampal ERK2
dimerization without affecting its phosphorylation (fig. 1B and 1C). Previous research has
shown that EGF stimulation induces ERK dimerization in cancer cells[13]. In physiological
contexts, the concentration of EGF plays a critical role in determining cellular responses, with
variations in EGF dosage influencing the transition between migratory and proliferative
states[56]. Thus, low EGF concentration, typical of paracrine signaling in early stages of wound
closure, is inadequate to trigger robust proliferative ERK signals in the cytoplasm. However, it
leads to pro-migratory signaling, inducing an increase in cell migration dependent on KSR-
ERK activation at the plasma membrane[56]. The EGF concentration used in our culture
experiments was consistent with that used by Herrero and colleagues[13] for HEK293T cells
and with concentrations promoting proliferation in MCF7 cells[56]. In this study, we replicated
Herrero and colleagues´ experiments, which also served as a control to evaluate ERK2
dimerization in the nervous tissue of rodents (fig. 1A). EGF has been demonstrated to induce

rapid and transient ERK phosphorylation in PC12 cells, which promotes cell proliferation[50].
We observed similar kinetics in our study. As many other tissues, brain cells express EGF
receptors and its activation regulates neurite outgrowth, as well as other outputs related to
neuronal homeostasis[57]. However, it is well known that different ligands are capable of
inducing different ERK activation kinetics and cellular outputs[58]. Recent findings from research
on breast tumors demonstrated that disrupting ERK dimerization, which was ligand-
dependent, hinders cellular motility by impairing the molecular machinery responsible for actin
cytoskeleton rearrangements[59]. It remains to be determined how ERK dimerization is altered
according to ligand specificity and its biological consequences, specifically in neurons.

ERK dimers in plasticity

We have recently discussed the role of ERK signaling in plasticity[9]. Hippocampal plasticity has
been proposed as a cellular correlate of experience-coding leading to memory formation and
storage. Then, a plausible hypothesis suggests that ERK dimerization might regulate memory
processes. Since the initial work by English and Sweatt[3] showing that ERK inhibition blocked
HFS-induced LTP in rat hippocampal area CA1, several authors have provided evidence in
favor of its participation in these processes[5,17,60,61] and, as stated before, different stimuli acting
on the same intracellular signaling pathway might cause alternative cellular fates depending
on the set of scaffolding proteins engaged[12]. As a general hallmark, it has been shown that
HFS protocol applied in the Schaffer Collaterals inputs to CA1 area induced ERK2
phosphorylation in rodents, and blocking ERK phosphorylation previous to LTP induction
prevented longer lasting forms of LTP[3]. In this study, we demonstrated that DEL pre-treatment
on acute slices allowed potentiation induction after HFS, but impaired normal hippocampal
LTP. In combination with previous work, these findings suggest that both ERK phosphorylation
and dimerization are crucial for the maintenance of LTP. Additionally, since ERK dimerization
and phosphorylation target different neuronal compartments, it is possible that at least two
distinct pathways are triggered during ERK activation. Moreover, memory processes have
been shown to involve the activation of multiple receptor systems, which in turn can activate
other molecular pathways that influence ERK signaling ultimately affecting its kinetics and
downstream targets. In consequence, it can be expected that ERK2 dimerization might also
be involved in LTP induction.

ERK dimerization in IA LTM reconsolidation

To this day, ERK signaling has been classically considered a key pathway for LTM
consolidation[5–8,27,61]. It has also been proposed as one of the molecules involved in memory
labilization and reconsolidation, becoming activated in the same set of neurons previously
engaged during conditioning[62]. It is worth noting that ERK's role in plasticity and memory
processes has been attributed almost exclusively to phosphorylation, leaving the potential
involvement of other post-translational modifications in regulating these processes largely
unexplored. Our results show that hippocampal inhibition of ERK dimerization impairs sIA
memory reconsolidation. However, the same manipulation of the pathway induced memory
enhancement when a wIA task was used, suggesting that the same brain structure might
engage opposing mechanisms depending on the strength of the consolidated memory. These
results are consistent with the idea that the restabilization process that takes place after either
sIA or wIA memory retrieval would be extremely specific. Based on this rationale, memory
restabilization might, at some point, quantitatively engage ERK signaling to guide behavioral

output. Consistent with this idea, it has been reported that pharmacological inhibition of ERK
dimerization increases its nuclear signaling[11]. Whether this mechanism might lead to memory
enhancement is still uncertain. Further investigation will either discard or confirm this
possibility. On the other hand, the scenario in which sIA and wIA retrieval/reconsolidation
might recruit distinct neural circuits cannot be discarded. Alternatively, different stimuli (e.g.
ligand identity and/or concentration) may activate distinct components from ERK pathways
which could in turn be transduced in the involvement of spatially distinct ERK pools that
regulate different cellular outputs. For instance, a strong stimulus could trigger protein
translation via ERK activation and promote gene transcription via ERK nuclear translocation.
In this scenario, ERK inhibition would impair memory by disrupting these mechanisms.
Conversely, after a weak stimulus, ERK inhibition might enhance memory by promoting
protein synthesis through alternative pathways. This could compensate for reduced ERK
activation and facilitate memory reconsolidation. Supporting this hypothesis, it was observed
that protein synthesis inhibition after reactivation exerted differential effects on memory
restabilization of an IA task in mice according to the area infused. Thus, protein synthesis
immediately after reactivation was required in the amygdala in order to support memory
reconsolidation while in the hippocampus and mPFC it was necessary to enhance retention[63].
Further experiments are necessary in order to support this proposal.

Our experiments also confirmed that the effect of hippocampal ERK dimerization is specific to
the reconsolidation process by showing that NR animals´ latencies to step through were not
affected by intra-hippocampal DEL injection (figures 3B and 4A). Regarding sIA memory
impairment following DEL administration, we observed persistent impairment after one week
(suppl. fig. 1), suggesting that there was no spontaneous recovery of the original memory
which would have suggested the possibility of the enhancement of an extinction process.

In the present manuscript, we showed for the first time evidence of ERK2 dimerization playing
a critical role in the central nervous system. Our findings reveal that ERK2 dimerization is
essential for hippocampal plasticity and memory reconsolidation, introducing a novel
mechanism of ERK2 regulation in the brain. Future studies will likely elucidate the other
molecular players recruited by this mechanism.

Funding

This work was supported by the following grants: Agencia Nacional de Promoción de la
Investigación, el Desarrollo Tecnológico y la Innovación (ANPCYT PICT 2016 0295, ANPCYT
PICT 2020 01534 and PICT 2015 1199), Consejo Nacional de Investigaciones Científicas y
Técnicas (CONICET PIP 2014-2016 No. 11220130100519CO) and Universidad de Buenos
Aires (UBACYT 2018-2021 - 20020170100390BA and 2014-2017 - 20020130200283BA),
Argentina.

Author Contributions

S.O.R. performed experiments, analyzed data, and contributed to write parts of the
manuscript. C.M. performed behavioral experiments, analyzed data, and reviewed the
manuscript. M.C.K., J.M. and M.F.A. performed experiments and reviewed the manuscript.
M.V.B. performed experiments and contributed to write parts of the manuscript. M.M.B
contributed to design the experiments, performed behavioral experiments and interpreted
data. A.G.R contributed to data analysis and reviewed the manuscript. F.J.U. & S.O.R.

designed and performed electrophysiology experiments in slices, interpreted data. M.F.
proposed the original research project, designed experiments and performed biochemical
assays, interpreted data, and drafted the manuscript.

Conflicts of Interest

The authors declare no conflicts of interest.

REFERENCES

1. Bliss, T. V. P. & Lømo, T. Long-lasting potentiation of synaptic transmission in the

dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J.*

Physiol. **232**, 331–356 (1973).

2. Bliss, T. V. & Collingridge, G. L. A synaptic model of memory: long-term potentiation in

the hippocampus. *Nature* **361**, 31–39 (1993).

3. English, J. D. & Sweatt, J. D. A requirement for the mitogen-activated protein kinase

cascade in hippocampal long term potentiation. *J. Biol. Chem.* **272**, 19103–19106

(1997).

4. Martin, K. C. *et al.* MAP kinase translocates into the nucleus of the presynaptic cell and

is required for long-term facilitation in Aplysia. *Neuron* **18**, 899–912 (1997).

5. Atkins, C. M., Selcher, J. C., Petraitis, J. J., Trzaskos, J. M. & Sweatt, J. D. The MAPK

cascade is required for mammalian associative learning. *Nat. Neurosci.* **1**, 602–609

(1998).

6. Blum, S., Moore, A. N., Adams, F. & Dash, P. K. A mitogen-activated protein kinase

cascade in the CA1/CA2 subfield of the dorsal hippocampus is essential for long-term

spatial memory. *J. Neurosci. Off. J. Soc. Neurosci.* **19**, 3535–3544 (1999).

7. Walz, R. *et al.* Effects of post-training infusions of a mitogen-activated protein kinase

kinase inhibitor into the hippocampus or entorhinal cortex on short- and long-term

retention of inhibitory avoidance. *Behav. Pharmacol.* **10**, 723–730 (1999).

8.   Schafe, G. E. *et al.* Activation of ERK/MAP kinase in the amygdala is required for

memory consolidation of pavlovian fear conditioning. *J. Neurosci. Off. J. Soc. Neurosci.*

20, 8177–8187 (2000).

9.   Ojea Ramos, S., Feld, M. & Fustiñana, M. S. Contributions of extracellular-signal

regulated kinase 1/2 activity to the memory trace. *Front. Mol. Neurosci.* **15**, (2022).

10. Lavoie, H., Gagnon, J. & Therrien, M. ERK signalling: a master regulator of cell

behaviour, life and fate. *Nat. Rev. Mol. Cell Biol.* **21**, 607–632 (2020).

11. Casar, B., Pinto, A. & Crespo, P. Essential role of ERK dimers in the activation of

cytoplasmic but not nuclear substrates by ERK-scaffold complexes. *Mol. Cell* **31**, 708–

721 (2008).

12. Casar, B. & Crespo, P. ERK Signals: Scaffolding Scaffolds? *Front. Cell Dev. Biol.* **4**,

(2016).

13. Herrero, A. *et al.* Small Molecule Inhibition of ERK Dimerization Prevents Tumorigenesis

by RAS-ERK Pathway Oncogenes. *Cancer Cell* **28**, 170–182 (2015).

14. Ahn, N. G. PORE-ing over ERK substrates. *Nat. Struct. Mol. Biol.* **16**, 1004–1005 (2009).

15. Earnest, S., Khokhlatchev, A., Albanesi, J. P. & Barylko, B. Phosphorylation of dynamin

by ERK2 inhibits the dynamin-microtubule interaction. *FEBS Lett.* **396**, 62–66 (1996).

16. Gong, R. & Tang, S.-J. Mitogen-activated protein kinase signaling is essential for

activity-dependent dendritic protein synthesis. *Neuroreport* **17**, 1575–1578 (2006).

17. Kelleher, R. J., 3rd, Govindarajan, A., Jung, H.-Y., Kang, H. & Tonegawa, S.

Translational control by MAPK signaling in long-term synaptic plasticity and memory.

Cell **116**, 467–479 (2004).

18. Kneussel, M. & Wagner, W. Myosin motors at neuronal synapses: drivers of membrane

transport and actin dynamics. *Nat. Rev. Neurosci.* **14**, 233–247 (2013).

19. Schrader, L. A. *et al.* ERK/MAPK regulates the Kv4.2 potassium channel by direct

phosphorylation of the pore-forming subunit. *Am. J. Physiol. Cell Physiol.* **290**, C852-861

(2006).

20. Boggio, E. M., Putignano, E., Sassoè-Pognetto, M., Pizzorusso, T. & Giustetto, M. Visual

stimulation activates ERK in synaptic and somatic compartments of rat cortical neurons

with parallel kinetics. *PloS One* **2**, e604 (2007).

21. Pirbhoy, P. S., Farris, S. & Steward, O. Synaptically driven phosphorylation of ribosomal

protein S6 is differentially regulated at active synapses versus dendrites and cell bodies

by MAPK and PI3K/mTOR signaling pathways. *Learn. Mem. Cold Spring Harb. N* **24**,

341–357 (2017).

22. Gao, C. *et al.* IQGAP1 regulates NR2A signaling, spine density, and cognitive

processes. *J. Neurosci. Off. J. Soc. Neurosci.* **31**, 8533–8542 (2011).

23. Albert-Gascó, H., Ros-Bernal, F., Castillo-Gómez, E. & Olucha-Bordonau, F. MAP/ERK

Signaling in Developing Cognitive and Emotional Function and Its Effect on Pathological

and Neurodegenerative Processes. *Int. J. Mol. Sci.* **21**, (2020).

24. Nader, K., Schafe, G. E. & LeDoux, J. E. The labile nature of consolidation theory. *Nat.*

Rev. Neurosci. **1**, 216–219 (2000).

25. Sara, S. J. Retrieval and reconsolidation: toward a neurobiology of remembering. *Learn.*

Mem. Cold Spring Harb. N **7**, 73–84 (2000).

26. Dudai, Y. & Eisenberg, M. Rites of passage of the engram: reconsolidation and the

lingering consolidation hypothesis. *Neuron* **44**, 93–100 (2004).

27. Feld, M., Dimant, B., Delorenzi, A., Coso, O. & Romano, A. Phosphorylation of extra-

nuclear ERK/MAPK is required for long-term memory consolidation in the crab

Chasmagnathus. *Behav. Brain Res.* **158**, 251–261 (2005).

28. Krawczyk, M. C. *et al.* Memory reconsolidation of an inhibitory avoidance task in mice

involves cytosolic ERK2 bidirectional modulation. *Neuroscience* **294**, 227–237 (2015).

29. Krawczyk, M. C. *et al.* Reconsolidation-induced memory persistence: Participation of late

phase hippocampal ERK activation. *Neurobiol. Learn. Mem.* **133**, 79–88 (2016).

30. Krawczyk, M. C., Millan, J., Blake, M. G., Feld, M. & Boccia, M. M. Relevance of ERK1/2

Post-retrieval Participation on Memory Processes: Insights in Their Particular Role on

Reconsolidation and Persistence of Memories. *Front. Mol. Neurosci.* **12**, 95 (2019).

31. Feld, M. *et al.* Decrease of ERK/MAPK overactivation in prefrontal cortex reverses early

memory deficit in a mouse model of Alzheimer's disease. *J. Alzheimers Dis. JAD* **40**, 69–

82 (2014).

32. Rivero-Echeto, M. C. *et al.* Simultaneous administration of cocaine and caffeine

dysregulates HCN and T-type channels. *Psychopharmacology (Berl.)* **238**, 787–810

(2021).

33. *Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates*. (2019).

34. Boccia, M. M., Acosta, G. B., Blake, M. G. & Baratti, C. M. Memory consolidation and

reconsolidation of an inhibitory avoidance response in mice: effects of i.c.v. injections of

hemicholinium-3. *Neuroscience* **124**, 735–741 (2004).

35. Blake, M. G., Boccia, M. M. & Baratti, C. M. Behavioral differences on memory retrieval

between two variants of step-through inhibitory avoidance task in mice. *Neurosci. Lett.*

444, 102–105 (2008).

36. Blake, M. G., Krawczyk, M. C., Baratti, C. M. & Boccia, M. M. Neuropharmacology of

memory consolidation and reconsolidation: Insights on central cholinergic mechanisms.

J. Physiol. Paris (2014) doi:10.1016/j.jphysparis.2014.04.005.

37. Boccia, M. *et al.* Activation of hippocampal nuclear factor-kappa B by retrieval is required

for memory reconsolidation. *J. Neurosci. Off. J. Soc. Neurosci.* **27**, 13436–13445 (2007).

38. Boccia, M. M., Blake, M. G., Krawczyk, M. C. & Baratti, C. M. Hippocampal α7 nicotinic

receptors modulate memory reconsolidation of an inhibitory avoidance task in mice.

Neuroscience **171**, 531–543 (2010).

39. Krawczyk, M. C., Millan, J., Blake, M. G. & Boccia, M. M. Role of prediction error and the

cholinergic system on memory reconsolidation processes in mice. *Neurobiol. Learn.*

Mem. **185**, 107534 (2021).

40. Acutain, M. F. *et al.* Reduced Expression of Hippocampal GluN2A-NMDAR Increases

Seizure Susceptibility and Causes Deficits in Contextual Memory. *Front. Neurosci.* **15**,

644100 (2021).

41. Kaech, S. & Banker, G. Culturing hippocampal neurons. *Nat. Protoc.* **1**, 2406–2415

(2006).

42. Swanger, S. A., He, Y. A., Richter, J. D. & Bassell, G. J. Dendritic GluN2A Synthesis

Mediates Activity-Induced NMDA Receptor Insertion. *J. Neurosci.* **33**, 8898–8908

(2013).

43. Pinto, A. & Crespo, P. Analysis of ERKs' dimerization by electrophoresis. *Methods Mol.*

Biol. Clifton NJ **661**, 335–342 (2010).

44. R core team. R: The R Project for Statistical Computing. https://www.r-project.org/

(2021).

45. Brooks, M. E. *et al.* glmmTMB Balances Speed and Flexibility Among Packages for

Zero-inflated Generalized Linear Mixed Modeling. *R J.* **9**, 378–400 (2017).

46. Hartig, F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression

models. *DHARMa* http://florianhartig.github.io/DHARMa/ (2024).

47. Lenth, R. V. emmeans: Estimated Marginal Means, aka Least-Squares Means. 1.10.6

https://doi.org/10.32614/CRAN.package.emmeans (2017).

48. Zaballos, M. A. *et al.* Inhibiting ERK dimerization ameliorates BRAF-driven anaplastic

thyroid cancer. *Cell. Mol. Life Sci. CMLS* **79**, 504 (2022).

49. Nguyen, T. T. *et al.* Co-regulation of the mitogen-activated protein kinase, extracellular

signal-regulated kinase 1, and the 90-kDa ribosomal S6 kinase in PC12 cells. Distinct

effects of the neurotrophic factor, nerve growth factor, and the mitogenic factor,

epidermal growth factor. *J. Biol. Chem.* **268**, 9803–9810 (1993).

50. Traverse, S., Gomez, N., Paterson, H., Marshall, C. & Cohen, P. Sustained activation of

the mitogen-activated protein (MAP) kinase cascade may be required for differentiation

of PC12 cells. Comparison of the effects of nerve growth factor and epidermal growth

factor. *Biochem. J.* **288 (Pt 2)**, 351–355 (1992).

51. Marshall, C. J. Specificity of receptor tyrosine kinase signaling: transient versus

sustained extracellular signal-regulated kinase activation. *Cell* **80**, 179–185 (1995).

52. Xing, C. & Imagawa, W. Altered MAP kinase (ERK1,2) regulation in primary cultures of

mammary tumor cells: elevated basal activity and sustained response to EGF.

Carcinogenesis **20**, 1201–1208 (1999).

53. Afinanisa, Q., Cho, M. K. & Seong, H.-A. AMPK Localization: A Key to Differential

Energy Regulation. *Int. J. Mol. Sci.* **22**, 10921 (2021).

54. Formoso, K., Lezoualc'h, F. & Mialet-Perez, J. Role of EPAC1 Signalosomes in Cell

Fate: Friends or Foes? *Cells* **9**, 1954 (2020).

55. Salles, A. *et al.* Hippocampal dynamics of synaptic NF-kappa B during inhibitory

avoidance long-term memory consolidation in mice. *Neuroscience* **291**, 70–80 (2015).

56. Brüggemann, Y., Karajannis, L. S., Stanoev, A., Stallaert, W. & Bastiaens, P. I. H.

Growth factor-dependent ErbB vesicular dynamics couple receptor signaling to spatially

and functionally distinct Erk pools. *Sci. Signal.* **14**, eabd9943 (2021).

57. Romano, R. & Bucci, C. Role of EGFR in the Nervous System. *Cells* **9**, 1887 (2020).

58. Miningou, N. & Blackwell, K. T. The road to ERK activation: Do neurons take alternate

routes? *Cell. Signal.* **68**, 109541 (2020).

59. de la Fuente-Vivas, D. *et al.* ERK1/2 mitogen-activated protein kinase dimerization is

essential for the regulation of cell motility. *Mol. Oncol.* (2024) doi:10.1002/1878-

0261.13732.

60. Impey, S. *et al.* Cross talk between ERK and PKA is required for Ca2+ stimulation of

CREB-dependent transcription and ERK nuclear translocation. *Neuron* **21**, 869–883

(1998).

61. Selcher, J. C. *et al.* A Role for ERK MAP Kinase in Physiologic Temporal Integration in

Hippocampal Area CA1. *Learn. Mem.* **10**, 26–39 (2003).

62. Zamorano, C., Fernández-Albert, J., Storm, D. R., Carné, X. & Sindreu, C. Memory

Retrieval Re-Activates Erk1/2 Signaling in the Same Set of CA1 Neurons Recruited

During Conditioning. *Neuroscience* **370**, 101–111 (2018).

63. Fukushima, H. *et al.* Enhancement of fear memory by retrieval through reconsolidation.

eLife **3**, e02736 (2014).