Engineering Asian elephant TP53: TP53 retrogene knockouts activate common and unique cancer-
relevant pathways

Emil Karpinski* [a], Nikil Badey [b], Esther Mintzer [a], Asaf Ashkenazy-Titelman [a], George M. Church* [a,b].

[a] *Department of Genetics, Harvard Medical School, Boston, MA, 02115, USA*
[b] *Wyss Institute for Biologically Inspired Engineering, Harvard University, Boston, MA 02215,*
USA
7

* Corresponding authors.

E-mail addresses: emil_karpinski@hms.harvard.edu; gchurch@genetics.med.harvard.edu

Abstract

TP53 functions as a central regulator in response to DNA damage and other cell stressors by

inducing the expression of many protective pathways such as cell cycle arrest and apoptosis.

Consequently, this gene is often found disrupted in human cancers. Elephants are a particularly

interesting species for the study of cancer, by virtue of their large number of cell divisions and long lives

yet low incidence of cancer. Elephants also possess multiple retrogene copies of TP53, which have

previously been shown to induce strong cellular responses to DNA damage. However, most previous

studies have largely focused only on African elephant TP53 retrogenes and often in non-native

backgrounds. Here we generated CRISPR-Cas9 knockouts of TP53, all 29 TP53 retrogenes, or both in

combination in Asian elephant fibroblasts. We find that while there is considerable overlap in the DNA

damage responses of the TP53 and retrogene knockouts, there are also many unique pathways enriched

in both. In particular, the retrogene knockouts exhibit strong enrichment of many extracellular pathways

suggesting they may play a large role in the tumor microenvironment and mitigating metastatic growth.

We also find that only a small fraction of these 29 retrogenes appear to be expressed across a variety of

tissues and identify three loci that are likely driving this response. This work shows for the first time the

transcriptomic effect of these retrogenes within their native background and establishes a foundation

for future research into the relative contributions of these genes.

Keywords: Asian elephants; TP53; retrogenes; cancer; CRISPR-Cas9; RNA-seq

1. Introduction

The gene TP53 encodes the transcription factor p53 and acts as one of the central master regulators in response to various cell stresses, including DNA damage, hypoxia, and nutrient deprivation [1,2]. In response to stress, p53 degradation is inhibited, its tetramers are stabilized, and they migrate to the nucleus resulting in the upregulations of many downstream signaling cascades, including those related to cell cycle arrest and apoptosis [1,3]. Consequently, this gene is often implicated in cancer development, and is found mutated in most human cancers, undermining genomic integrity [1,2,4]. Likewise, individuals diagnosed with Li-Fraumeni Syndrome have germline mutations in TP53 and much greater risk and earlier onset of various cancers [2,3]. In addition, TP53 also plays a role in cellular senescence and reprogramming efficiency, making it important in fundamental cellular biology technologies, such as the generation of iPSCs and immortalization [1,2,5].

Peto's paradox is the observation that cancer incidence in a species does not always scale with body mass and/or lifespan [6,7]. Elephants have emerged as a key species in attempts to resolve this paradox due to their large size, long lifespan, and very low cancer risk [7,8]. Genomic analyses have identified the duplication of many tumor suppressor genes across extant elephants, including multiple retrogene copies of TP53 [7,8]. Although there is still considerable uncertainty about how or which of these TP53 retrogenes (RTGs) may confer tumor resistance, previous works have shown that the encoded protein of at least one of these retrogenes, African elephant RTG 9, induces strong cellular responses to DNA damage [7–9]. However, most prior studies have largely focused on their ability to induce apoptosis and have often examined the effect of retrogenes in different genetic backgrounds (i.e. human or mouse cells) [7–9]. Additionally, almost all functional analyses have focused on African elephant retrogenes despite the identification of TP53 retrogene expansion in other extinct and extant proboscideans [7].

Here, we establish a baseline for similar work in Asian elephants. We designed CRISPR-Cas9 guides to disrupt TP53, all TP53 retrogenes, or both in combination in Asian elephant fibroblasts. We then examined the transcriptional profiles of these knockouts with and without DNA damage stress, to resolve the protective roles TP53 retrogenes might play at a cellular level. Additionally, we compare our results with the transcriptomic patterns of these retrogenes across various Asian elephant tissues to identify which may be acting as tumor suppressors and which are likely nonfunctional. The work we present here suggests that these retrogenes may have a much larger and more varied response than previously anticipated and lays the groundwork for future studies focusing on specific retrogene targets.

3. Methods

3.1 – TP53 identification and primer design

We extracted and aligned the canonical TP53 locus as well as all 29 retrogenes (RTGs) from the

Elephas maximus reference genome (GCF_024166365.1). For the RTGs, we trimmed the alignment to

include only the ~1.1kb segment overlapping the exons in TP53. We then designed two primers to

specifically amplify the TP53 gene (Supplementary Table 1). For the RTGs, we designed seven forward

and four reverse primer variations, all amplifying the same locus, but containing all possible binding site

mutations to minimize amplification bias (Supplementary Table 1). These primers were made up to

10μM stock solutions and equal volumes of all forward and all reverse primers combined to produce

master forward and reverse stock solutions.

3.2 – Guide Design

We generated all possible guide RNAs (gRNAs) within TP53 from the start of first exon to the

end of the last exon using the generate_kmery.py script distributed as part of GuideScan2 [10]. All guides

were then screened for predicted off-targets against the entire *E. maximus* reference genome using

GuideScan2 with a maximum mismatch distance of 3 (-m 3), and with the option to output all guides (-t -

1). The output was then further filtered using a custom python script (Cas9_MMFiltering_V1_1.py) and

scoring matrix to filter potential gRNAs based on how many mismatches they possessed relative to

potential off-target sites and where along the spacer those mismatches occur. From the remaining

guides we selected two guides (Supplementary Table 1) that cut within exon 1 and between exon 2 and

3 respectively to generate TP53 knockouts as these disrupt the TAD1 domain and should interfere with

proper splicing resulting in a premature stop codon.

To design guides for the RTG knockouts, we generated all possible guides against the trimmed

RTG alignment for all 29 retrogenes using generate_kmery.py script as before. We then screened all

guides for those that were predicted to have activity to the most possible targets and identified two

gRNAs that were each predicted to have activity against all but three retrogenes, which were only

targetable by either one of the two guides (Supplementary Table 1; Fig. 1). We then subsequently chose

one additional guide that targeted those three remaining retrogenes to ensure that each retrogene had

at least two gRNAs.

3.3 – Cell culturing and media

Fibroblast cells were grown in a custom media formulation with the final concentrations: 50%

FGM-2 (Lonza FBM Basal Media (CC-3131) supplemented with FGM-2 SingleQuots supplements (CC-

4126)); 20% heat-inactivated fetal bovine serum (ThermoFisher, A5670801); and 30% Mem Alpha

(Gibco, 12571063). Cells were grown in an incubator at 37°C with 5% CO_2.

3.4 – Cas9 RNP Electroporation

All gRNAs used for both the TP53 and RTG knockouts were ordered from Synthego and

resuspended to 100μM in 1X TE buffer. Cas9 nuclease was ordered from IDT (catalog # 10007807). All

nucleofections were conducted using the Lonza P4 Primary Cell 96-well Kit (catalog # V4SP-4096) using

code CA-137. Approximately 400-500k cells were used for each transfection in 20μl reactions with 3μM

final concentrations of each guide and 1μM of Cas9 nuclease per guide (i.e. a 3:1 ratio of gRNA:Cas9).

RNPs were pre-complexed in nucleofection solution for 20 minutes at room temperature. Cells were

transfected with either both TP53 gRNAs, all three RTG gRNAs, both TP53 and RTG gRNAs ("All"

condition), or a non-targeting gRNA which has no significant homology to the *E. maximus* genome

(Supplementary Table 1). For the All nucleofection, gRNAs were first combined and then concentrated

using a Savant SpeedVac DNA 120 Concentrator (Thermo Scientific), before being resuspended in

premade nucleofector and supplement solution. Following nucleofection cells were immediately

transferred into 6-well plates containing prewarmed media and allowed to recover for 4 days. Media

was changed 24 hours post nucleofection and ever 2-3 days thereafter. Due to the presence of possible

TP53 mutations that accumulate with prolonging passaging, a second separate nucleofection was

conducted as above prior to the Mitomycin C stressed RNA-seq. In both cases, all treatment conditions

were conducted using three biological replicates.

3.5 – Editing Efficiency

During the first passage (approximately 4 days post nucleofection), about 10% of the cells were

collected and DNA extracted using the DNeasy Blood and Tissue Kit (Qiagen) using the manufacturer's

recommended protocol. From every extraction we amplified both the TP53 locus spanning the expected

knockout deletion, and the retrogene loci. 25μL PCR reactions were run with final concentrations of 1X

Phusion Hot Start Flex 2X Master Mix and 500nM of each primer, using between 1-3μL of template.

Cycling conditions were: 98°C for 3 minutes; 12 cycles of 98°C for 15 seconds, then 69°C for 60 seconds;

25 cycles of 98°C for 15 seconds, 67°C for 30 seconds, then 72°C for 45 seconds; followed by a final

extension at 72°C for 5 minutes. Resulting PCR amplicons were purified over MinElute PCR Purification

columns (Qiagen) using the recommended protocol and visualized on E-gel EX 2% Agarose gels

(Invitrogen) to confirm the presence or absence of any large deletions. We examined the relative

fluorescence of our products using the Image Lab 6.1 (Bio-Rad). We then calculated a minimum editing

efficiency (the relative fluorescence of our expected deleted product, relative to everything else in the

lane) and a maximum editing efficiency (the relative fluorescence of all non-wild-type bands within each

lane).

To confirm efficiency estimates and examine RTG deletion distribution, we sent each of our

amplicon pools for nanopore sequencing using the Premium PCR service at Primordium to get full length

sequences of our DNA fragments. For each sample, raw reads were first filtered using a custom python

script (FastqFilter.py) to between the maximum and minimum expected sizes +/- 10% respectively.

Mapping was carried out using BBMap [11] using default parameters against either the TP53 amplicon, an

RTG consensus amplicon, or a multi-fasta with 26 RTGs (RTG amplicons with identical sequences outside

of the editing region collapsed).

To calculate editing efficiency, we used a custom python script (CigarDel.py) which uses a region

file containing the deletion regions of interest, a minimum deletion proportion (i.e. the proportion of

bases mapping to the deleted region that need to be missing), and the CIGAR string of each read to

evaluate the presence or absence of various deletions in the data. For TP53 we analyzed the presence of

the expected deletion if Cas9 induced a double-strand break at both sgRNA targeting sites. For the RTG

we examined the presence of any of the three possible deletions between two guides. All analyses were

carried out with a minimum deletion proportion of 0.9.

To examine the distribution of deletions at each locus we identified the names of reads

containing deletions and extracted those that mapped to each locus. We then cross-referenced this with

the deletions identified against the RTG consensus to obtain read distributions for reads mapped to all

RTGs or the collapsed RTG loci.

3.6 – Growth Curve

Approximately four days post nucleofection, we seeded 50k cells from each sample into the

wells of four 24-well plate. 36 hours after seeding and every 24 hours thereafter, cells were detached

using TrypLE Express (Gibco) and cell counts,and viability estimates were obtained using a Countess II

cell counter (Thermo Fisher). For each data point we averaged two individual counts. These values were

then imported in GraphPad Prism 10 and analyzed using a two-way ANOVA for significance.

3.7 – RT-qPCR

50k cells were seeded into each well of four 48 well plates. Approximately 36 hours after

seeding, growth media was exchanged for one supplemented with either 0µM, 1µM, 10µM, or 50µM

Mitomycin C (MOLTOX). Cells were incubated for 18 hours in the presence of Mitomycin C-containing

media, after which RNA was immediately extracted using the RNeasy Micro kit (Qiagen) according to the

manufacturer's recommendations. We designed and validated four pairs of RT-qPCR primers for *E.*

maximus GAPDH, ATR, p21, and MDM2 (Supplementary Table 1). 20µL reactions were run with final

concentrations of 1X TaqPath 1-Step Multiplex Master Mix (No ROX), and 500nM of each primer, using

2µL of purified RNA template. Cycling conditions were: 25°C for 2 minutes; 53°C for 10 minutes

seconds; 95°C for 2 minutes; 40 cycles of 95°C for 15 seconds, 60°C for 30 seconds; followed by a final

extension at 60°C for 1 minute, and a melt curve between 60°C and 95°C. All samples for each treatment

condition were analyzed in duplicate, and relative expression standardized to GAPDH, using the $2^{-\Delta\Delta CT}$

method [12]. Data was combined across biological replicates within each treatment. Average results were

imported into GraphPad Prism 10 and analyzed for significance using a two-way ANOVA.

3.8 – Bulk RNA-seq

Approximately 300-500k cells were collected, flash frozen on dry ice, and shipped to Genewiz

for bulk RNA sequencing within ten days of transfection. Expression of transcripts was done using

Salmon (v1.8.0 [13]) using a custom index consisting of all mRNAs and non-coding RNAs annotated within

the *E. maximus* reference genome (GCF_024166365.1), with all 29 retrogene sequences added. The

salmon index was generated using the keepDuplicates flag to avoid collapsing any RTGs. Salmon quant

files were analyzed using DESeq2 [14] as part of a custom Rscript to analyze samples in a pairwise fashion

(i.e. one set of biological replicates against the controls at a time). Differentially expressed genes (DEGs)

were identified with an adjusted p value of less than 0.05 (padj < 0.05). No minimum log2 fold change

was applied for initial analysis of TP53 pathway genes. A minimum absolute log2 fold change of 1 was

used when analyzing DEG overlap between different conditions. Venn diagrams and DEG overlap was

generated using a web based custom venn diagram tool

(https://bioinformatics.psb.ugent.be/webtools/Venn/), and pathway enrichment preformed using

GeneCodis4 [15].

3.9 – Retrogene expression in public datasets

To ensure the expression patterns we observed were not tissue/cell type specific, we examined

the expression of our retrogenes in some preliminary shared data from mesenchymal stem cells and

epithelial cells. These cells contained deletions generated using the same guides as in this study, but the

epithelial cells were also subject to various small molecule treatments. We also downloaded public RNA-

seq datasets for Asian elephant ovary (SRX15554240), salivary glands (SRX15554239), lymph nodes

(SRX15554238), thyroid (SRX15554237), lung (SRX15554236), and lymphocytes (SRX1423033), and

analyzed it as above.

3.10 – Depth plots

To examine where reads from each of the three RTGs showing repeated expression were

originating, we mapped the data for our datasets and the public datasets back to the Asian elephant

reference genome using STAR [16]. We then extracted read depths along the full annotated length of RTGs

LOC126068247, LOC126068267, and LOC126068248 using samtools depth [17]. For our dataset we

averaged expression values within biological replicates and plotted the data in Microsoft Excel.

3.11 – Maximum likelihood phylogeny

We download African elephant retrogene 9 (KF715863.1) to investigate a possible relationship

between this retrogenes and the Asian elephant retrogenes. We then aligned the retrogene to the Asian

elephant TP53 CDS and truncated the RTG 9 sequence to contain only the homologous sequence to

TP53. We then aligned the truncated RTG 9 sequence, Asian elephant TP53, and all 29 Asian elephant

retrogenes using muscle (v5.1; Edgar, 2004). We generated a maximum likelihood phylogeny using IQ-

Tree [19], using ModelFinder to identify the best supported substitution model [20], and 100 bootstraps (-b

100).

3.12 – EMS mutagenesis and clonal expansion

Biological replicates were combined and 50k cells per condition were seeded into 6-well plates.

Cell were grown in media containing 0 µg/mL, 0.5 µg/mL, or 1 µg/mL of EMS (Thermo Scientific) for 16

hours, at which point EMS-containing media was replaced with fresh media [21]. Cells were then grown for

an additional 8 passages, to potentially outgrow any fast-growing and immortalized mutants. Following

8 passages, 1000 cells were seeded onto a 10cm^2 dish and observed over a period of 1-3 weeks. Colonies

were marked and labelled to observe growth. A subset of well-defined circular colonies were removed

by treating with 5µl of TrypLE Express (Gibco) twice, and transferring the detached cells to a 6-well plate

and then to a 10cm^2 dish.

4. Results

4.1 – The Asian elephant genome contains 29 TP53 retrogenes

We searched the annotations of the Asian elephant reference genome and identified 29 retrogene copies of TP53 (RTGs). 27 of these RTGs are found on chromosome 27, while the remaining two are found on chromosomes 24 and 26. By comparison, the canonical copy of TP53 is located on chromosome 19. Most of the RTGs are approximately 1.1 kb in length, roughly the same length as the CDS of TP53 as would be expected of retrogenes, with the exception of three (LOC126068247 – 64.1 kb; LOC126068267 – 32.5 kb; LOC126068248 – 4.9 kb; Fig. S1A).

When aligned and translated in the same frame as TP53, each RTG carries a premature stop codon, which primarily cluster in five main areas across the homology region (Fig. 1A). We then compared the percent identity of the DNA and amino acid sequences of the retrogenes to TP53. We find significantly higher levels of homology at the DNA level than at the amino acid level, irrespective of whether we compare the entire amino acid sequence, or only the truncated peptides (Fig. S1B). Comparatively, across the same homologous region, the RTGs contain significantly higher levels of identity (Fig. S1C).

4.2 – TP53 and RTG editing and validation

We attempted to design paired Cas9 guide RNAs (sgRNAs) to produce large, disruptive deletions in TP53 and all the retrogenes. We managed to identify two sgRNAs that specifically target TP53, which result in the partial loss of exon 1, the entirety of exon 2, and a premature stop codon after read through into intron 2 (Fig. 1A). We could not identify just two guides that were both predicted to have activity across all 29 retrogenes, so we designed a third guide to ensure each RTG was targetable by at least two sgRNAs (Fig. 1B; Fig. S2).

We electroporated fibroblast cells with Cas9 RNPs consisting of either the two TP53 guides, the three RTG guides, all five guides together (All), or a guide predicted to have no activity in the Asian elephant genome (Non Targ.). After approximately four days, we then collected cells, extracted DNA, and amplified a portion of TP53 and RTG genes flanking the expected editing sites. We observed high levels of editing across all three conditions when we examined the resulting amplicons on an agarose gel (Fig. S3). However, due to our PCR reaction amplifying all 29 loci simultaneously, we couldn't assess which and how many of the RTGs were being edited.

To address this issue, we sent both PCR reactions for end-to-end nanopore sequencing. We quantified the number of observed deletions, using a custom script which marks the presence of a

deletion if 90% of the bases within a defined region were absent. We note that this is likely an

underestimate of the true editing efficiency, as this will only identify the perfect deletion product, and

miss any indels and/or incomplete loss of the flanked segment of DNA. We observed high editing

efficiency in all conditions using either the TP53 guides (TP53 KO – 67%; All KO – 66%) or the RTG guides

(RTG KO – 77%; All KO – 82%), and almost no editing in the Non Targ. condition (<0.1% at both loci) (Fig.

1C). We further examined the proportion of each deletion product observed in the RTG PCRs, and noted

a clear pattern, wherein deletions between guides 2 and 3 were the most frequent, in line with

expectations about the targeting scope of those two guides (Fig. 1D).

Next, we wanted to disentangle the editing on a per-locus level for the RTG data. We mapped

the data to each unique RTG (collapsing those with identical sequences outside the editing region) and

identified the proportion of each deletion at each locus. Although we generally observe a mix of all

three deletions at each locus (some of which is likely due to cross mapping between the RTGs), we

generally observe deletions in line with our expectations based on their targeting scope (Fig. 1E). Only

two loci buck this trend, LOC126068338 and LOC126068461, both of which have a single point mutation

at position 10 within the seed region of the guide (Fig. S2). Due to the importance of this region for

guide specificity, we filtered these targets out; however, we note that at least at these two loci the

guides appear to be permissive for editing. When combined with the per-locus reads that were not

identified as containing a deletion (Fig. 1F), we observe an average of 76.8% editing efficiency across all

unique loci.

4.3 – All editing conditions show similar growth and transcriptomic patterns

Approximately four days following transfection, we seeded 50k cells from each condition into

four 24-well plates and observed the growth and viability of the cells over approximately 4.5 days. We

observed no significant difference in the growth (Fig. 2A) or viability (Fig. 2B) of the cells across this time.

However, we do note that the TP53 KO wells appeared to reach confluency a day before the other

conditions suggesting faster growth, which is consistent with our experiences handling these cells in

subsequent passages.

Next, we wanted to see if we could observe a functional effect of the knockouts we generated.

We cultured 50k cells in the presence of increasing concentrations of Mitomycin C (MMC), a DNA

damaging agent [7,22], and examined the expression of p21 (CDKN1A) and MDM2, two direct downstream

targets of p53, and an upstream regulator, ATR (Fig. 2C). We observed significant decreases in

expression of MDM2 and p21 at all non-zero Mitomycin C concentrations across each condition, and

almost no significant differences in ATR. However, we were surprised to see that the response in the

RTG KO was so similar to that of the TP53 and All KOs. We extracted DNA from the same line of cells one

passage later and PCR amplified the TP53 and RTG loci again. We observed a drop in Sanger sequence

quality across the TP53 amplicons in some of the biological replicates (Fig. S4), generally consistent with

the location of the RTG guides. This suggests there may be some off-target editing in the RTG samples

that becomes enriched during routine passaging of the RTG KO cells, particularly if these TP53 mutants

also possess a potential growth advantage as would be suggested by our growth curve.

4.4 – RNA-seq analysis reveals overlapping and unique TP53 and RTG responses

293   To minimize the effect of any off-target mutations on our RNA-seq results, we sent cells soon

294   after nucleofection (within ~10 days). Despite this we observe the presence of off-target edits at the

295   homologous location of the RTG guides within the TP53 transcripts (Fig. S5). Interestingly, we see

296   significant enrichment of these TP53 mutations in our MMC-treated cells. This suggests that these TP53

297   mutations might confer a significant growth advantage to the cells as was also suggested by our growth

298   curve analysis.

299   Given most literature on elephant RTGs has largely focused on how these retrogenes may

interact with TP53 and MDM2, we first decided to look at changes in expression of all direct

downstream targets of TP53 (Fig. 3). We find that in the absence of stress (i.e. 10μM Mitomycin C), we

see relatively small changes in expression within most downstream targets with the exception of MDM2

and CDKN1A in both the TP53 and All KOs. Interestingly, we note that TP53 is not identified as

significantly downregulated in the TP53 KO, suggesting maybe that nonsense-mediated decay is less

pronounced in Asian elephants. In comparison, we observe very few differentially expressed genes

(DEGs) in the RTG KO, and only TP53 that has a log2 fold change greater than 1. This suggests that the

RTGs may play some role in either stabilizing or increasing the abundance of p53, although likely not

directly by acting as a transcription factor as none of the RTGs have an intact DNA-binding domain (Fig.

1).

310   To confirm the results above were genuine and not due to mismapping between the RTGs and

the canonical TP53 gene, we generated a synthetic dataset, consisting of 1,000 2x150 bp reads from

TP53 and each of the RTG loci. We then used this dataset to compare the amount of reads assigned to

each locus when mapped back to the genome using BBMap [11], BWA-MEM [23], or STAR [16], followed by

quantification with Salmon [13], or via direct quantification by Salmon (Fig. S6). We find that all four

methods gave similar results, with BBMap performing the worst, possibly due to how it assigns multi-

mapped reads to the first locus in the reference. In light of these results, we conclude that the decrease

in TP53 expression is likely not a bioinformatics artifact due to poor mapping.

In the presence of Mitomycin C, all three conditions have similar patterns in DEGs (Fig. 3),

although typically less pronounced in the RTG KO. As before, we also observe that TP53 is significantly

less expressed in the RTG KO and not in the TP53 KO. Additionally, IGFBP-3, a gene which can contribute

to cell growth and apoptosis [24,25], is also downregulated in the RTG KO. We also observe significant

downregulation of SFN across all three conditions in the MMC-treated cells, in line with this gene's

known tumor suppressor function of arresting cellular replication in response to DNA damage [26,27].

Next, we wanted to examine the different pathways enriched within each condition. Due to the

large number of genes with small changes we filtered the dataset to a minimum absolute log2 fold-

change of at least 1. We then extracted the DEG lists for each condition and analyzed the overlap

between them in either the non-stressed or MMC-treated cells (Fig. 4A). We performed BioPlanet

pathway enrichment[28] on the 763 DEGs common to all three conditions treated with Mitomycin C. We

observed significant enrichment in p53 signaling pathways, as would be expected of cells exposed to a

DNA damaging agent, as well as enrichment in many DNA replication and cell cycle pathways (Fig. 4B).

We also examined the unique DEGs in the TP53 and RTG KO cells with Mitomycin C treatment

(Fig. 4C). For the TP53 KO, we see enrichment in many pathways that would be expected given the

known checkpoint and cell stress responses coordinated by TP53 (e.g. cell cycle and proliferation, DNA

repair, etc; Fig. 4D) [29–31]. By comparison, unique RTG KO DEGs are enriched in extracellular organization

and signaling/interaction pathways (Fig. 4E), suggesting that RTGs may play a unique protective role in

regulating the tumor microenvironment and limiting metastasis [32]. Additionally, the largest number of

DEGs appear to belong to the apoptotic signaling in response to DNA damage pathway, a function that

has previously been suggested for African elephant RTGs [7,9].

4.5 – Very few RTGs are likely to be expressed or functional

Given the unique function suggested by the RNA-seq of the RTG KO cells, we next wanted to see

if we could investigate which RTGs might be responsible for the changes we observed. In none of the

conditions, irrespective of treatment, did we detect differential expression of any of the RTGs.

Furthermore, we generally do not observe differences in the amount of RTGs expressed in response to

Mitomycin C treatment (Fig. 5A), with the exception of the All KO. We do observe an increase in the

amount of RTGs expressed in the RTG and All KO conditions, irrespective of Mitomycin C treatment

(significance values not shown), although this may be a lingering transcriptional effect caused by double-

strand break repair [33]. However, we do note that all but three RTGs have very low read counts

(Supplementary Table 2).

We next wanted to confirm that this pattern in RTG expression is not due to cell type or tissue

specific. We download public RNA-seq datasets for Asian elephant ovary, salivary glands, lymph nodes,

lung, thyroid, and lymphocytes, as well as examined some preliminary data generated with the same

guides in epithelial and mesenchymal stem cells, and examined RTG expression patterns in these

datasets. We observe the same general pattern, where the only three RTGs which show consistent

expression are LOC126068247, LOC126068267, and LOC126068248 (Supplementary Table 3). However,

these three RTGs are the three largest RTG genes (LOC126068247 – 64.1 kb; LOC126068267 – 32.5 kb;

LOC126068248 – 4.9 kb) based on their annotations.

To examine what was driving the expression of these RTGs, we extracted RNA-seq mapped

reads across each of these three loci and examined the frequency of reads across each locus. For two of

these RTGs, we observe that most of the reads assigned to these RTGs map thousands of base pairs

upstream of the region homologous to the TP53 CDS. In LOC126068247 and LOC126068248, we observe

almost no mapped reads within the TP53 homology region either in our data or the publicly available

datasets (Figs S7-S10). By comparison, LOC126068267 also has peaks of high coverage upstream of the

TP53 homology region but shows similar levels of coverage across the homology region in both datasets

(Figs S11-S12).

Given the lack of expression of most of the annotated retrogenes, we decided to compare those

that might be functional in Asian elephants with African elephant retrogene 9 (RTG 9; KF715863), on

which most previous studies have focused [7–9]. We aligned the TP53 homologous regions of all Asian

elephant RTGs, African elephant RTG 9, and the TP53 CDS, and built a maximum-likelihood phylogeny.

We observe that RTG 9 is most closely related to a clade containing both two of our putative expressed

RTGs, LOC126068267 and LOC126068248 (Fig. 5B), further suggesting that these two RTGs might be

responsible for the changes we observe. However, we do note that due to the high similarity of many of

these RTGs, the bootstrap support values within our phylogeny are quite weak.

4.6 – RTG KO allows for partial clonal expansion

Although compounded by a relative lack of studies and optimization relative to mouse and

human cell lines, elephant cells have proven difficult to immortalize and grow [34,35]. Our own efforts to

produce clonal lines of these cells using established protocols [36] have failed (data not shown). However,

we reasoned that since TP53 mutations often precede cancer development, and many cancers naturally

immortalize, we may be able to exploit this feature in our knockouts to produce isogenic lines.

To examine this, we treated cells with either 0µg/mL, 0.5µg/mL, or 1µg/mL EMS for 16 hours [21],

and then performed eight passages, seeding 10% of the detached cells, to outgrow fast growing and

potentially immortalized cells. All 1µg/mL cells died within the first few passages, as did all the TP53 KO

conditions and the Non Targ. 0µg/mL EMS treatment. After eight passages, the outgrown cells were

collected and 1,000 from each condition were seeded into a 10cm^2 dish, and colonies picked over 1-3

weeks. We observed between 4 and 26 well-defined circular colonies in the RTG and All KO conditions

treated with 0µg/mL or 0.5µg/mL, in comparison to only one in the Non Targ. 0.5µg/mL condition

(Supplementary Table 4). Of these we picked three colonies each from the All 0 µg/mL, All 0.5µg/mL,

and RTG 0 µg/mL plates, two colonies from the RTG 0.5µg/mL, and the only colony from the Non Targ.

0.5µg/mL plate. We attempted to expand colonies, but unfortunately most of the cells in most of these

lines appeared quite sick and exhibited very slow growth. One of the RTG 0.5µg/mL colonies managed to

propagate an additional four passages within the 10cm^2 dishes, but slowed down immensely thereafter

and became senescent.

5. Discussion

TP53 is tremendously important for the cellular response to stress and DNA damage, and has

been implicated in many biological important processes including cellular reprogramming and

immortalization [1,5,37,38]. Additional retrogene copies found in many proboscideans have been implicated

in this taxa's low cancer incidence rate [7,8,39,40], and likely also contribute to the difficulty in working with

and modifying these cells. With the publication of a new high quality Asian elephant reference genome

(as part of the VGP), we wanted to examine the high-level function of these retrogenes, both

independently and in combination with TP53 mutations.

We find that, similar to African elephants, Asian elephant retrogenes all contain premature stop

codons. Additionally, we note the putative encoded proteins are quite dissimilar, suggesting possibly

relaxed selection at these loci and perhaps non-functionality [41]. Despite this, we note that at least three

of these RTGs appear to be stably expressed in various cell types and tissues. Likewise, we observe clear

transcriptional differences between TP53 and RTG KO cells, including a significant downregulation of

TP53 expression in our RTG KO cells, suggesting a novel function for elephant retrogenes. Additionally,

we observe a differential enrichment of pathways when disrupting RTGs versus TP53, indicating that

while there is a high amount of overlap between these genes, they also regulate many unique pathways.

In particular, genes enriched within many extracellular organization and signaling pathways were

observed in the RTG KO population. Some of these (e.g. extracellular matrix organization, angiogenesis)

have been implicated as contributing to the establishment of tumor microenvironments, and are gaining

increased importance as vital to cancer growth and spread [32]. This may also explain the low prevalence

of malignant cancers in Asian elephants [39]. We also show that while these retrogenes are likely not

sufficient for clonal expansion, their knockout appears to permit the generation of single cell colonies.

This is congruent with current work which suggests that the knockdown of these retrogenes is also

important during the production of elephant iPSCs [42].

However, we do note that most of the annotated retrogenes with the Asian elephant genome

are not expressed across an extensive list of tissues, and only a handful may actually be functional.

Although this may be in line with previous estimates as to the amount of tumor suppressor evolution

necessary for the observed incidence rate in elephants [41]. Interestingly, two of the retrogenes we

observed most stably expressed across in different tissues appear to be closely related to African

elephant RTG 9, which has been shown to induce strong cellular responses to DNA damage [7–9] and

apoptosis in mouse fibroblasts and human cancer cell lines [7,9]. Future research should focus on

individual retrogenes, especially LOC126068267, to determine if these three are responsible for the

signal we observe, and possibly what the function of these highly expressed regions upstream of the

TP53 homologous sequences may be. Additionally, by extending the analysis of proboscidean RTGs into

Asian elephants we lay the groundwork for more comparative studies, to examine how much each

response is evolutionary conserved versus uniquely derived along each species' lineage.

6. Author contributions

EK and GMC conceived the study with feedback from EM and AAT. EK and NB designed and

conducted all wet-lab experiments with feedback and help from EM and AAT. EK completed all

bioinformatic analyses. AAT provided additional RNA-seq data for epithelial cell lines. EK wrote the first

draft of the manuscript. All authors helped in the revision of subsequent drafts.

7. Data availability

Bulk RNA-seq data for all conditions with and without mitomycin was deposited on NCBI

(BioProject Accession: PRJNA1147676). Bioinformatic scripts and tools created for this work are

available at: https://github.com/EmilKarpinski/RTG_TP53_Analysis.

8. Declaration of competing interests

For a complete list of G.M.C.'s financial interests, please visit http://arep.med.harvard.edu/gmc/tech.html. All other authors declare no other conflicts of interest.

9. Acknowledgments

We would like to thank Li Li and Zhengkuan Tang for discussions on RNA-seq experimental setup and analysis. We would also like to thank George Chao for guidance on nanopore analysis, and Sergiy Velychko for suggestions and troubleshooting of elephant fibroblast cell culture. Lastly, we would like to thank Roman Teo Oliynyk, Chun-Ting Wu, and Ahmed Mahas, as well as the rest of the Church lab for helpful discussions throughout this project.

EK and AAT were supported by a structured research agreement between Harvard Medical School and Colossal Biosciences.

References

1. Wang, H., Guo, M., Wei, H. & Chen, Y. Targeting p53 pathways: mechanisms, structures, and advances in therapy. *Signal Transduct Target Ther* **8**, 92 (2023).

2. Hainaut, P. & Hollstein, M. p53 and Human Cancer: The First Ten Thousand Mutations. in 81–137 (1999). doi:10.1016/S0065-230X(08)60785-X.

3. Correa, H. Li–Fraumeni Syndrome. *J Pediatr Genet* **05**, 084–088 (2016).

4. Hainaut, P. & Pfeifer, G. P. Somatic TP53 Mutations in the Era of Genome Sequencing. *Cold Spring Harb Perspect Med* **6**, a026179 (2016).

5. Rasmussen, M. A. *et al.* Transient p53 Suppression Increases Reprogramming of Human Fibroblasts without Affecting Apoptosis and DNA Damage. *Stem Cell Reports* **3**, 404–413 (2014).

6. Peto, R., Roe, F. J., Lee, P. N., Levy, L. & Clack, J. Cancer and ageing in mice and men. *Br J Cancer* **32**, 411–426 (1975).

7. Sulak, M. *et al.* TP53 copy number expansion is associated with the evolution of increased body size and an enhanced DNA damage response in elephants. *Elife* **5**, (2016).

8. Abegglen, L. M. *et al.* Potential Mechanisms for Cancer Resistance in Elephants and Comparative Cellular Response to DNA Damage in Humans. *JAMA* **314**, 1850 (2015).

9. Preston, A. J. *et al.* Elephant TP53-RETROGENE 9 induces transcription-independent apoptosis at the mitochondria. *Cell Death Discov* **9**, 66 (2023).

10. Schmidt, H. *et al.* Genome-wide CRISPR guide RNA design and specificity analysis with GuideScan2. (2022) doi:10.1101/2022.05.02.490368.

11. Bushnell, B. BBMap. http://sourceforge.net/projects/bbmap/ (2022).

12. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402–408 (2001).

13. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417–419 (2017).

14. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

15. Garcia-Moreno, A. *et al.* Functional Enrichment Analysis of Regulatory Elements. *Biomedicines* **10**, 590 (2022).

16. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

17. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

18. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**, 1792–1797 (2004).

19. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol* **32**, 268–274
(2015).

20. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**, 587–589 (2017).

21. Munroe, R. & Schimenti, J. Mutagenesis of Mouse Embryonic Stem Cells with
Ethylmethanesulfonate. in 131–138 (2009). doi:10.1007/978-1-59745-471-1_7.

22. Lee, Y.-J., Park, S.-J., Ciccone, S. L. M., Kim, C.-R. & Lee, S.-H. An in vivo analysis of MMC-induced
DNA damage and its repair. *Carcinogenesis* **27**, 446–453 (2006).

23. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**, 1754–1760 (2009).

24. Marzec, K. A., Lin, M. Z., Martin, J. L. & Baxter, R. C. Involvement of p53 in insulin-like growth
factor binding protein-3 regulation in the breast cancer cell response to DNA damage.
Oncotarget **6**, 26583–26598 (2015).

25. Grimberg, A. *et al.* p53-Dependent and p53-Independent Induction of Insulin-Like Growth Factor
Binding Protein-3 by Deoxyribonucleic Acid Damage and Hypoxia. *J Clin Endocrinol Metab* **90**,
3568–3574 (2005).

26. Hermeking, H. The 14-3-3 cancer connection. *Nat Rev Cancer* **3**, 931–943 (2003).

27. Hermeking, H. *et al.* 14-3-3σ Is a p53-Regulated Inhibitor of G2/M Progression. *Mol Cell* **1**, 3–11
(1997).

28. Huang, R. *et al.* The NCATS BioPlanet – An Integrated Platform for Exploring the Universe of
Cellular Signaling Pathways for Toxicology, Systems Biology, and Chemical Genomics. *Front
Pharmacol* **10**, (2019).

29. Dey, G. *et al.* Signaling network of Oncostatin M pathway. *J Cell Commun Signal* **7**, 103–108
(2013).

30. Moldovan, G.-L. & D'Andrea, A. D. How the Fanconi Anemia Pathway Guards the Genome. *Annu
Rev Genet* **43**, 223–249 (2009).

31. Du, W. & Searle, J. The Rb Pathway and Cancer Therapeutics. *Curr Drug Targets* **10**, 581–589
(2009).

32. de Visser, K. E. & Joyce, J. A. The evolving tumor microenvironment: From cancer initiation to
metastatic outgrowth. *Cancer Cell* **41**, 374–403 (2023).

33. Guha, S. & Bhaumik, S. R. Transcription-coupled DNA double-strand break repair. *DNA Repair
(Amst)* **109**, 103211 (2022).

34. Fukuda, T. *et al.* Expression of human cell cycle regulators in the primary cell line of the African
savannah elephant (loxodonta africana) increases proliferation until senescence, but does not
induce immortalization. *In Vitro Cell Dev Biol Anim* **52**, 20–26 (2016).

35.   Pearson, V. R. *et al.* Identification of African Elephant Polyomavirus in wild elephants and the
527   creation of a vector expressing its viral tumor antigens to transform elephant primary cells. *PLoS*
528   *One* **16**, e0244334 (2021).

529   36.   Housden, B. E. *et al.* Identification of potential drug targets for tuberous sclerosis complex by
530   synthetic screens combining CRISPR-based knockouts with RNAi. *Sci Signal* **8**, (2015).

531   37.   Liu, T. M. *et al.* Molecular Basis of Immortalization of Human Mesenchymal Stem Cells by
532   Combination of p53 Knockdown and Human Telomerase Reverse Transcriptase Overexpression.
533   *Stem Cells Dev* **22**, 268–278 (2013).

534   38.   Jha, K. K., Banga, S., Palejwala, V. & Ozer, H. L. SV40-Mediated Immortalization. *Exp Cell Res* **245**,
535   1–7 (1998).

536   39.   Tollis, M. *et al.* Elephant Genomes Reveal Accelerated Evolution in Mechanisms Underlying
537   Disease Defenses. *Mol Biol Evol* **38**, 3606–3620 (2021).

538   40.   Vazquez, J. M. & Lynch, V. J. Pervasive duplication of tumor suppressors in Afrotherians during
539   the evolution of large bodies and reduced cancer risk. *Elife* **10**, 1–45 (2021).

540   41.   Nunney, L. Cancer suppression and the evolution of multiple retrogene copies of TP53 in
541   elephants: A re-evaluation. *Evol Appl* **15**, 891–901 (2022).

542   42.   Appleton, E. *et al.* Derivation of elephant induced pluripotent stem cells. *bioRxiv* (2024)
543   doi:10.1101/2024.03.05.583606.

Figure 1. TP53 Editing. A. Schematic showing an alignment of Asian elephant TP53 and a retrogene (RTG) consensus, with domains labelled based on homology to human TP53. Exons are shown in yellow, with introns indicated by black lines. Missing sequence is indicated with a dashed line in the RTGs. Additionally, we've included the positions of deletions greater than 3bp in the RTG consensus, and the approximate locations of stop codons observed across the 29 Asian elephant retrogenes. The locations and directions of guides are shown for both TP53 and

the retrogenes. **B.** Predicted guide activity against each of the 29 Asian elephant retrogenes, with green indicating the guide is expected to be

functional against that target and red indicating non-functionality. **C.** Observed editing efficiency of the expected deletions based on nanopore

data. **D.** The frequency of deletions observed in all reads mapped against the RTG consensus across different treatment conditions. **E.** The

frequency of each deletion across the individual Asian elephant retrogenes as a proportion of all deletion-containing reads mapped to that locus,

or all mapped read (**F**). Amplicons which were identical across the non-edited regions were collapsed as they are bioinformatically

undistinguishable. Error bars represent standard deviation in all plots.

Figure 2. Cell characterization and expression. A. Growth rates of cells with TP53, RTG, or combined
knockouts, or treated with a non-targeting guide. **B.** Cell viability observed during the growth curve for
each condition. **C.** The expression of two downstream (p21 and MDM2) and one upstream (ATR) genes
within the TP53 pathway. Expression levels for the RTG KO are shown with a dashed line due to the
possible presence of TP53 off-target mutations. Significance levels: $p \leq 0.05$ (*); $p \leq 0.01$ (**); $p \leq 0.001$
(***); $p \leq 0.0001$ (****). Errors bars indicate standard error of the mean in all plots.

Figure 3. Expression of TP53 pathway genes. Log2 fold changes observed for genes immediately
downstream of TP53 in the p53 signaling pathway (KEGG pathway hsa04115). Crosses indicate that no
significant difference in expression was observed for that gene in a given condition. Genes in italics have
an absolute log2 fold change of < 1 and were filtered out of enrichment analyses. Only downstream
genes with a significant difference in at least one condition are shown.

Figure 4. Pathway enrichment. A. Differentially expressed gene (DEG) overlap observed between treatment conditions for cells in normal media or exposed to 10μM Mitomycin C. Genes were filtered for a minimum absolute log2 fold change greater than 1. **B.** BioPlanet pathway enrichment for the 763 DEGs common to all three conditions in the Mitomycin C treatment cells. **C.** DEG overlap between TP53 KO and RTG KO cells in normal media or exposed to 10μM Mitomycin C. **D.** BioPlanet pathway enrichment of unique DEGs in TP53 KO Mitomycin C treated cells. **E.** BioPlanet pathway enrichment of unique DEGs in RTG KO Mitomycin C treated cells.

Figure 5. RTG expression. A. Expression of RTGs in each treatment condition in cells exposed to 10μM
Mitomycin C or not. Errors bars indicate standard deviation. **B**. Maximum likelihood phylogeny showing
the relationship between Asian elephant TP53 and the RTGs, as well as African elephant RTG9. Three
tree was rooted using Asian elephant TP53. Green circles indicate nodes with bootstrap support ≥ 90%,
while numbers above nodes represent bootstrap support for select nodes. The three Asian elephant
RTGs which are most commonly expressed, and African elephant RTG9 are indicated in orange and blue
respectively.

587