CUP-SHAPED COTYLEDON2 activates *MIR319C* transcription and
promotes cell proliferation in Arabidopsis leaf primordia

Naveen Shankar[1#], Abhishek Gupta[1], Somsree Roy[1], Vishwadeep Mane[1], Olivier Hamant[2] and
Utpal Nath[1*]

1 Department of Microbiology and Cell Biology, Indian Institute of Science, Bengaluru, India.
2 Laboratoire de Reproduction et Développement des Plantes, Université de Lyon, ENS de
Lyon, UCBL, INRA, CNRS, 46 Allée d'Italie, 69364 Lyon Cedex 07, France.
Current address: Department of Plant Physiology, Umeå Plant Science Centre, Umeå University,
901 87 Umeå, Sweden.

12
*Corresponding author. Email: utpalnath@iisc.ac.in

Abstract

The microRNA miR319 regulates leaf shape and size in diverse plant species by reducing the

level of the target transcripts that encode JAW-TCP proteins, the transcription factors (TF)

which commit the proliferating pavement cells to differentiation. Expression of *MIR319C*, one

of the three miR319-producing genes in Arabidopsis, is expressed throughout the incipient leaf

primordia, and the expression domain gets restricted to the base at later stages, partly due to

the direct action of JAW-TCPs. However, the factors that activate and maintain *MIR319C*

expression in leaf primordia are yet unknown. Here, we report the CUP-SHAPED

COTYLEDON2 (CUC2) transcription factor as a direct activator of *MIR319C* transcription. In

a yeast one-hybrid (Y1H) screen, we identified several NAC domain TFs as potential regulators

of *MIR319C*. Subsequent ex vivo binding and transactivation assays suggested that CUC2 may

bind to distal promoter region of *MIR319C*. Mutants with compromised *CUC2* and *MIR319C*

activities exhibited smaller leaf areas and reduced pavement cell numbers due to early cell

proliferation-to-differentiation transition. Morphometric analysis of higher order *CUC2* and

MIR319 loss-of-function mutants highlighted the crucial role of the *CUC2-MIR319* module in

promoting leaf cell proliferation and indicated functional redundancy among the three *MIR319*

genes in regulating JAW-TCP levels during leaf growth. Additionally, the phenotypes of

mutants with altered *CUC2* and *MIR319/JAW-TCP* activities demonstrated that *CUC2*

enhances leaf size through the *MIR319C-JAW-TCP* pathway. Overall, our findings uncovered

a novel role for CUC2 in promoting cell proliferation by activating *MIR319C* transcription in

the leaf primordia.

Introduction

Leaf development can be empirically divided into three major stages, namely initiation, primary morphogenesis, and secondary morphogenesis (Bar and Ori, 2014). Initiation of a leaf primordium commences with the specification of a group of founder cells at the periphery of the shoot apical meristem (SAM), a dome-shaped structure at the shoot apex that harbours a constant pool of stem cells throughout plant life (Barton, 2010). Concurrently, a low cell proliferation region, termed as the 'boundary zone' gets established at the interface that separates the primordium from the SAM (Hepworth and Lenhard, 2014; Wang et al., 2016). The basic form of the leaf consisting of the lamina and the petiole is established during primary morphogenesis, wherein leaf growth is primarily sustained by cell proliferation. The end of primary morphogenesis is marked by the switch from cell proliferation to differentiation and the formation of specific cell/ tissue types such as trichomes and stomata (Bar and Ori, 2014). Serrations are also initiated at this stage by outgrowth at specific sites along the base of the leaf margin. During secondary morphogenesis, differentiation of all the cell types is completed, and the leaf grows primarily by cell expansion (Bar and Ori, 2014; Du et al., 2018; Maugarny-Calès and Laufs, 2018).

The shape and size of the mature leaf can be regulated at each of the above three developmental stages by modulating the number of founder cells during initiation, or by regulating the rate, location and duration of cell proliferation and expansion (Gonzalez et al., 2012; Hepworth and Lenhard, 2014; Kalve et al., 2014). Besides, the timing of cell proliferation arrest, which directly affects the number of cells in mature leaves, is a major determinant of the final leaf size in a wide range of plant species (Gázquez and Beemster, 2017). In Arabidopsis leaves, arrest of cell proliferation occurs basipetally; cells towards the tip stop dividing and commit to differentiation (Donnelly et al., 1999; Andriankaja et al., 2012; Fox et al., 2018), while cells at the base continue to proliferate. Subsequently, the diffused boundary between the proliferation and differentiation regions remains at a constant distance from the base before being abruptly terminated (Andriankaja et al., 2012; Fox et al., 2018). Compensation also plays a key role in regulating leaf size, wherein a change in cell proliferation triggers an opposite effect on cell size at maturity, thus maintaining the mature leaf size relatively unaltered (Tsukaya, 2005; Hisanaga et al., 2015).

Genetic factors such as *GROWTH REGULATING FACTORS* (*GRFs*), *ANGUSTI- FOLIA3*
(*AN3*) and the cytochrome P450-encoding gene *KLUH* (*KLU*) promote growth by maintaining
the proliferative state of the cells in the proximal region (Horiguchi et al., 2005; Anastasiou et
al., 2007; Kazama et al., 2010; Vercruyssen et al., 2014). The genetic modules that initiate the
arrest of cell proliferation have also been described to a large extent in Arabidopsis (Gonzalez
et al., 2012; Kalve et al., 2014; Du et al., 2018). Notably, two gene families that encode
antagonistically acting transcription factors (TF) control the transition from cell proliferation
to cell expansion: the *GROWTH REGULATING FACTORs* (*GRFs*) and the *CINCINNATA*-like
TEOSINTE BRANCHED 1, *CYCLOIDEA*, *PROLIFERATING CELL FACTOR1/2* (*CIN-TCPs*)
(Rodriguez et al., 2014; Sarvepalli and Nath, 2018). The spatiotemporal activities of *GRFs* are
regulated by microRNA miR396. miR396 downregulates *GRF* transcripts in the distal region
of the leaf, thereby restricting GRF activities towards the base (Rodriguez et al., 2010). GRFs
interact with GRF-INTERACTING FACTOR 1/ANGUSTIFOLIA 3 (GIF1/AN3) and
upregulate *CYCLIN B* (*CYCB*) expression to delay the proliferation-differentiation transition
(Schneider et al., 2023). Eight *CIN-TCP* genes promote leaf maturation by triggering the
transition of cell proliferation to differentiation in diverse eudicots including Arabidopsis
(Sarvepalli and Nath, 2018). The spatiotemporal pattern of *CIN-TCP* transcripts are regulated
by a combined action of their promoter activity and microRNA miR319 (Palatnik et al., 2003;
Efroni et al., 2008). The miR319 activity prevents the accumulation of transcripts of five *CIN-*
TCPs (*TCP2, 3, 4, 10, 24*), also known as *JAGGED AND WAVY TCPs* (*JAW-TCPs*), in the
proximal region of the young leaf primordia (Palatnik et al., 2003; Nag et al., 2009). Expression
of miR319-resistant form of *JAW-TCPs* with endogenous regulation leads to smaller leaves
with fewer cells due to precocious proliferation arrest, while ectopic expression of miR319
results in large and wavy leaves due to a delay in the transition from proliferation to
differentiation (Palatnik et al., 2003; Efroni et al., 2008; Challa et al., 2019). CIN-TCPs commit
the leaf cells towards differentiation by modulating multiple pathways, such as auxin and
cytokinin response, meristematic activity, and cell cycle progression (Sarvepalli and Nath,
2018).
99
100 The *CUP SHAPED COTYLEDON* (*CUC*) group of genes, which encode the NAM/ ATAF/
101 CUC (NAC) group of TFs, regulate boundary development in diverse angiosperms (Wang et
102 al., 2016). The Arabidopsis *CUC* members *CUC1*, *CUC2*, and *CUC3* are consistently
103 expressed in the boundary zones, and their absence leads to the fusion of adjacent organs such
104 as cotyledons, sepals, stamens, and ovules (Aida et al., 1997; Ishida et al., 2000; Vroemen et

al., 2003). The *CUC* genes are also required for the maintenance of shoot meristem via the activation of the *SHOOT MERISTEMLESS* (*STM*) gene and its homologues (Aida et al., 1999; Hibara et al., 2006; Scofield et al., 2018). In leaf primordia, *CUC2* and *CUC3* are expressed at distinct foci along the margin, where they are necessary for the initiation and growth of serrations/ leaflets through a feedback mechanism involving auxin and the auxin efflux carrier PINFORMED1 (PIN1) (Nikovics et al., 2006; Bilsborough et al., 2011; Hasson et al., 2011; Maugarny-Calès et al., 2019). Interestingly, some of the gain-of-function phenotypes of *JAW-TCPs*, such as impaired SAM function, cotyledon fusion, and smooth leaf margin, are dependent on the downregulation of *CUC* activity (Koyama et al., 2007; Koyama et al., 2010; Koyama et al., 2017; Challa et al., 2019). While TCP3 inhibits *CUC1* and *CUC2* by activating the transcription of the miR164-encoding gene *MIR164A* to promote shoot morphogenesis (Koyama et al., 2007; Koyama et al., 2010), TCP4 protein interacts with CUC and inhibits their homo-dimerization and transactivation abilities to modulate serration formation and outgrowth (Rubio-Somoza et al., 2014).

Expression analysis of the miR319-encoding genes *MIR319A*, *MIR319B*, and *MIR319C* in Arabidopsis suggested their distinct developmental functions due to largely non-overlapping expression pattern (Nag et al., 2009). Whereas *MIR319A* and *MIR319B* promoter activity is not detected in leaves, *MIR319C* is strongly expressed in the leaf primordia (Nag et al., 2009; Shankar et al., 2023). However, the single loss-of-function mutants for miR319a and miR319b show a moderate decrease in leaf serration, and the *miR319a;miR319b* double mutant leaves lack serration, suggesting a role for *MIR319A* and *MIR319B* genes in leaf development (Koyama et al., 2017). However, the *miR319a;miR319b* leaves are as big as wild-type (Koyama et al., 2017), pointing towards a key role of miR319c in downregulating *JAW-TCP* activity during the early leaf growth. We had earlier demonstrated that JAW-TCPs transcriptionally repress the miR319-encoding gene *MIR319C* in the distal region to prevent miR319 accumulation, thereby establishing spatially separated proximal miR319 domain and distal JAW-TCP domain in incipient leaf primordia (Shankar et al., 2023). However, it is still unknown how the *MIR319* genes are transcriptionally activated in the early leaf primordia. Here, we demonstrate that CUC2 transcriptionally activates the *MIR319C* gene, and together with the products *MIR319A* and *MIR319B*, the mature miR319 sustains cell proliferation and promote leaf growth.

Results

A yeast one-hybrid screen identifies transcriptional activators of *MIR319C*

The *MIR319C* promoter activity is primarily detected at the shoot apex, more specifically at the meristem-primordia boundary and in young leaf primordia (Shankar et al., 2023) (**Figs. S1A, S1B**), suggesting that members of the transcriptome subset within this domain activate *MIR319C* transcription. To identify such direct transcriptional regulators of *MIR319C*, we screened a subset of yeast one-hybrid library containing 378 Arabidopsis transcription factors enriched in the shoot apex for their ability to bind to the upstream regulatory region (URR) of *MIR319C* (**Figs. S1C, S1D, Table S3**). Fifty seven TFs were identified that promoted the growth of yeast cells when a 2736 bp URR was used as bait to drive the expression of the *HISTIDINE3* (*HIS3*) selection gene (**Table S4**). Gene ontology analysis revealed that TFs belonging to NAC, TCP, ETHYLENE RESPONSE FACTOR (ERF), BASIC LEUCINE ZIPPER DOMAIN (bZIP), and BASIC HELIX LOOP HELIX (bHLH) families were over-represented among the positive clones identified in the screen (**Fig. 1A**).

Interestingly, 5 out of the 6 NACs recovered from the Y1H screen (**Fig. S1E**) show comparable DNA-binding preferences, the target DNA sequence being TT(G/A)CGT (Lindemose et al., 2014), which is present at the 2619 nucleotide position upstream to the start of pre-*MIR319C* sequence. The binding preference of these 5 NACs is also shared by other members of the NAM, OsNAC7, NAC2, ANAC011, and AtNAC3 subgroup of the AtNAC TF family (Ooka et al., 2003; Lindemose et al., 2014). To identify additional NAC members capable of binding to the *MIR319C URR*, we chose 20 representative members from these 5 AtNAC subgroups (**Fig S2A, S2B**) and tested in yeast for their binding ability to three non-overlapping *MIR319C URR* fragments named as p^{distal} (-2736 to -1866 bp), p^{middle} (-1866 to -1106 bp), and $p^{proximal}$ (-1106 to -1 bp) (**Fig 1B**). Six NAC members including CUC1, CUC2 & CUC3 supported yeast growth when p^{distal} was used as bait (**Fig 1C, Fig. S2B**).

CUC2 activates *MIR319C* in leaf primordia

CUC2, the primary *CUC* member in leaves (Nikovics et al., 2006; Hasson et al., 2011), is active throughout the shoot apex and initiating leaf primordia, overlapping the *MIR319C* domain (**Fig S1A, S1B**). The overlapping expression pattern in early leaf primordia, and the ability of

CUCs to bind to *MIR319C* URR in yeast, led us to investigate if CUCs regulate *MIR319C*

promoter activity in leaves by determining *MIR319C* expression in mutants with altered *CUC2*

levels using a *GUS*-based reporter line for *MIR319C* promoter activity (*pMIR319C::GUS[#1]*)

(Nag et al., 2009; Shankar et al., 2023). We used an Image-J-based method to quantify the

domain of *pMIR319C::GUS* expression relative to the leaf length and termed it as relative GUS

domain (Shankar et al., 2023). The relative GUS domain was significantly reduced (median

~0.35 compared to ~0.6 in Col-0) in *cuc2-1* leaves (Hasson et al., 2011) and was increased

(median ~0.75) in *cuc2-1D* (Larue et al., 2009), a line with elevated CUC2 activity (**Fig 1D,**

1E). The pre-*MIR319C* transcript level was ~1.5-fold lower and >2-fold higher in

pMIR319C::GUS[#1];cuc2-1 and *pMIR319C::GUS[#1];cuc2-1D* seedlings, respectively (**Fig 1F**).

Together, these results suggest that CUC2 is required and sufficient to activate *MIR319C*

expression in leaf primordia.

***CUC2* promotes cell proliferation in miR319-dependent manner in leaves**

Mature miR319 promotes cell proliferation by degrading the transcripts of five *JAW-TCP*s

(Palatnik et al., 2003; Efroni et al., 2008; Koyama et al., 2010; Challa et al., 2019). If CUC

proteins promote miR319 accumulation, inactivation of either *CUC2* or *MIR319C* is expected

to yield smaller leaves with fewer cells. To test this, we analysed the matures leaves of two

existing loss-of-function alleles of *CUC2*, *cuc2-1* and *cuc2-3* (Nikovics et al., 2006; Hasson et

al., 2011), and *miR319c-1*, a new allele of *MIR319* where the level of mature miR319c is

perhaps reduced (**Fig S4A, S4B**). None of these mutants showed any noticeable reduction in

mature leaf area or epidermal cell area (**Fig. 2A-C**, **Fig S3A, S3B**), perhaps because of the

hypomorphic nature of the mutant alleles and due to the presence of their redundant partners.

To test the genetic relationship between *CUC2* and *MIR319C*, we generated the double mutant

line *cuc2-3; miR319c-1* and analysed its leaf size parameters. The double mutant leaves were

nearly half the size of their respective parents (**Fig 2A, 2B, S3**), though the average area of

their pavement cells did not reduce (**Fig. 2C, 2D**). Thus, the *cuc2-3; miR319c-1* leaves were

made up of only ~2000 cells whereas the Col-0 leaves had ~4,500 cells (**Fig. 2E**), suggesting

reduced cell proliferation in the double mutant. The corresponding cell numbers for the

cuc2-3 and *miR319c-1* leaves also reduced compared to the wildtype, but to a much less extent

(**Fig. 2E**). Fewer cells in *cuc2-3* and *miR319c-1* leaves is perhaps partly compensated by

slightly increased cell area (**Fig. 2D**), a phenomenon not observed in the *cuc2-3; miR319c-1*

double mutant leaves.

To determine the timing of this cell proliferation defect during leaf morphogenesis, we

performed confocal imaging of propidium iodide-stained primordia at 1, 3 and 5 days after

initiation (DAI) and analysed the images at the cellular level using morphographX tool. The

analysis revealed that while the cell areas of leaf primordia in all the genotypes at 1 DAI were

nearly homogenous, there was a distinct proximodistal gradient in cell growth at 3 and 5 DAI

marked by smaller cells at the base and larger cells in the distal region of the leaf primordia

(**Fig 3A**). There was a significant increase in average cell areas from 1-5 DAI in all the

genotypes, chiefly contributed by the larger cells in distal region of the primordia (**Figs 3A-C,**

S3C), an indication of onset of differentiation. Interestingly, the average cell area in

cuc2-3;miR319c-1 leaf primordia (median~$125\mu m^2$) was significantly higher than Col-0,

cuc2-3, and *miR319c-1* genotypes at 3 DAI (median~$100\mu m^2$) (**Fig 3A, 3B**) but not at 1 DAI

(**Fig S3C**). The trend was also consistent at 5 DAI where *cuc2-3* and *cuc2-3;miR319c-1* had

higher average cell areas (median~$150\mu m^2$) compared to Col-0 and *miR319c-1*

(median~$125\mu m^2$) (**Fig 3C**). The cellular analysis further revealed that *cuc2-3;miR319c-1* had

a higher proportion of larger size cells ($100\text{-}200\mu m^2$ and $>200\mu m^2$) compared to wild type and

the parental genotypes at 3 DAI (**Fig S3D**), while *cuc2-3* and *cuc2-3;miR319c-1* mutants had

larger population of $200\text{-}500\mu m^2$ and $>500\mu m^2$ cells and a corresponding lower proportion of

$<250\mu m^2$ cells at 5 DAI (**Fig S3E**). Cellular analysis suggested an altered cell growth pattern

in *cuc2-3;miR319c-1* leaf primordia marked by a precocious increase in pavement cell areas.

These results, together with those presented in **Fig. 1**, provides some genetic evidence that

CUC proteins promote cell proliferation in leaf primordia in a miR319c-dependent manner.

To investigate if *MIR319A* and *MIR319B* genes also regulate leaf size, we combined *cuc2-3*

and *miR319ab* (Koyama et al., 2017) and studied the leaf phenotypes in the higher-order

mutants. We observed a significant proportion of seedlings with narrow leaves, fused

cotyledons or arrested growth within the homozygous *cuc2-3;miR319a* and *cuc2-3;miR319b*

population (**Fig 4A**), which was not observed in either of the parent population. While 15.4%

each of seedlings within the *cuc2-3;miR319a* population exhibited narrow leaf and arrested

growth, 3.3% and 26% of the *cuc2-3;miR319b* seedlings had narrow leaves and arrested

growth/fused cotyledons, respectively (**Fig 4A**). We also observed a similar range of

phenotypes in the *cuc2-3;miR319a/+;miR319b/+* and *cuc2-3;miR319a/+;miR319b* mutant

population (**Fig 4A**). For simplicity, we shall denote *cuc2-3;miR319a/+;miR319b/+* and

cuc2-3;miR319a/+;miR319b mutants as *cuc2-3;miR319ab* from hereon. Next, we measured

the leaf areas, pavement cell area and number in matured first pair of leaves from the seedlings

having narrow leaves within the *cuc2-3;miR319a*, *cuc2-3;miR319b*, and *cuc2-3;miR319ab*

population. The leaf areas of *cuc2-3;miR319a* and *cuc2-3;miR319b* mutants were significantly

lower (2 to 3-fold) than the wild-type and the parent genotypes (**Fig 4B, 4C**). The leaf areas in

cuc2-3;miR319ab were reduced to the same extent of *cuc2-3;miR319a* and *cuc2-3;miR319b*

double mutants (**Fig 4B, 4C**). Interestingly, the lower leaf areas in *cuc2-3;miR319ab* mutants

was due to the reduction in both pavement cell areas and cell numbers (**Figs 4D, S5A, S5B**).

While the pavement cell shapes of *cuc2-3;miR319a/+;miR319b/+* mutant was similar to the

wild-type, *cuc2-3;miR319a/+;miR319b* had less lobed and more elongated cells (**Fig S5B**),

indicating an alteration in cell growth pattern.

We speculated that low miR319 levels in the *cuc2-3;miR319a* and *cuc2-3;miR319b* mutants in

would cause an increase in *JAW-TCP* activity during embryogenesis, resulting in cotyledon

fusion and seedling arrest phenotypes. To test this possibility, we established *cuc2-1;GR#2* by

crossing *cuc2-1* mutant to Col-0;*GR#2* (Shankar et al., 2023), a line where miR319-resistant

dexamethasone (DEX) inducible form of TCP4 protein is expressed under its endogenous

promoter in the wild-type background. 12/583 (2%) seedlings obtained from the seeds of the

DEX-treated *cuc2-1;GR#2* flowering plants (**See *Methods***) exhibited cotyledon separation

defects while all the seedlings from the ethanol-treated Col-0;*GR#2* and *cuc2-1;GR#2* plants or

the DEX-treated Col-0;*GR#2* had well-separated cotyledons (**Fig S4C**). The above results

suggest that the *CUC2*-mediated activation of miR319c also contributes to the downregulation

of *JAW-TCP* activity during the establishment of the SAM-cotyledon boundary.

To further explore the functional redundancy among the three *MIR319* genes in leaf cell

proliferation, we determined the final leaf area, pavement cell area and number in the first pair

of leaves in the triple mutants obtained by combining *miR319ab* and *miR319c-1* genotypes.

The   higher   order   *miR319abc*   mutants,   i.e.,   *miR319a;miR319b;miR319c-1/+*,

miR319a;miR319b/+;miR319c-1, and *miR319a;miR319b;miR319c-1* had highly reduced leaf

areas (median 5-8 mm^2) compared to the wild-type (median~22 mm^2) (**Fig 5A, 5B**). The

reduction in leaf areas was only due to significantly lower cell numbers (~4-fold reduction)

(**Fig 4D**) but not cell areas (**Fig S5A, S5B**). Together, the above results suggests functional

redundancy among three *MIR319* genes and a critical role for *CUC2-MIR319* pathway in

promoting leaf size via cell proliferation in leaf primordia.

CUC2 binds and activates *MIR319C* transcription

We earlier showed that CUCs specifically bind to the distal URR of *MIR319C* containing the putative CUC binding site in yeast (**Fig 1E**). To study the importance of the CUC binding site for *MIR319C* promoter activity, we generated a *GUS*-based reporter line for *MIR319C* promoter carrying mutations in the CUC binding site (*mCUCBSpMIR319C::GUS*) (**Fig 5C**) and determined the relative GUS domain in the *mCUCBSpMIR319C::GUS* expressing leaf primordia. The relative GUS domain was significantly reduced in the leaf primordia expressing *mCUCBSpMIR319C::GUS* transgene as compared to the wild-type *pMIR319C::GUS*[#2] (Shankar et al., 2023) expressing leaf primordia (**Fig 5D-F**), suggesting that the CUC binding site is required for the *MIR319C* promoter activity.

Next, to determine if CUC2 directly activates *MIR319C* transcription, we induced CUC2 activity in 10-day-old *pRPS5A::CUC2-GR-HA* seedlings (Tian et al., 2014) by treating them with dexamethasone (DEX) for two hours in the presence of the protein synthesis inhibitor cycloheximide (CHX) and measured pre-*MIR319C* levels by RT-qPCR. CUC2 was able to significantly elevate pre-*MIR319C* transcript levels (~2-fold), similar to transcripts of its known direct target *CUC3*, both in the presence and the absence of CHX (**Fig 5G**), suggesting that *MIR319C* could be a direct target of CUC2. The above results indicate that CUC2 binds and activates *MIR319C* transcription in early leaf primordia.

***CUC-MIR319C-JAW-TCP* module regulates leaf size**

The gain-of-function line for *CUC2* activity, *cuc2-1D*, has larger than wild-type leaf areas due to increased cell numbers (Larue et al., 2009). Since *MIR319C* expression was elevated in *cuc2-1D* (**Fig 1D-F**), we investigated if the increased leaf size in *cuc2-1D* is due to reduced *JAW-TCP* activities. We crossed the *TCP4* overexpression line Col-0;*GR*[#1] (Challa et al., 2016) to *cuc2-1D* and studied leaf areas in ethanol and DEX-treated F1 progeny. Although Col-0;*GR*[#1] line contains a DEX-inducible miR319-resistant TCP4, it resembles wild-type under non-inductive conditions (**Fig 6A**), as reported earlier (Challa et al., 2016). The average first leaf areas of ethanol-treated Col-0;*GR*[#1]x *cuc2-1D* F1 individuals were significantly higher (median~32 mm^2) than ethanol-treated Col-0;*GR*[#1]x Col-0 F1 control (median~23 mm^2) (**Fig 6A, 6B**). Induction of *rTCP4* activity in both Col-0;*GR*[#1]x Col-0 F1 and Col-0;*GR*[#1]x *cuc2-1D* F1 seedlings resulted in the formation of smaller leaves with similar areas (median~14

mm^2) (**Fig 6A, 6B**), suggesting that *CUC2* promotes leaf size via miR319-mediated downregulation of *JAW-TCPs*.

Overexpression of miR319 leads to the formation of larger than wild-type leaves with deeper serrations (**Fig 6C**) (Efroni et al. 2008; Challa et al. 2019). The lack of *CUC2* activity in the miR319 overexpression background only affected the serration formation but did not alter the leaf area (**Fig 6C, 6D**). The similar leaf areas of *pBLS>>miR319a* and *cuc2-1;pBLS>>miR319a* mutants strongly suggests that the *MIR319-JAW-TCP* module regulates leaf size downstream of *CUC2*. These results also demonstrate that the *CUC-MIR319-JAW-TCP* module regulates leaf size and serration pattern by distinct mechanisms.

CUC2 and JAW-TCP activities establish MIR319C expression domain in early leaf primordia

We previously demonstrated that JAW-TCPs downregulate *MIR319C* expression in the distal region of the leaf primordia (Shankar et al., 2023). A previous study showed that *JAW-TCPs* transcriptionally inhibit the expression of *CUCs* during early shoot organogenesis (Koyama et al., 2007). To investigate if *JAW-TCPs* transcriptionally downregulate *CUC2* and thereby *MIR319C* expression in leaf primordia, we studied the domain of expression of the promoter-reporter line for *CUC2*, i.e., *pCUC2::GUS* (Nikovics et al., 2006) in knockdown line for all five *JAW-TCPs*, i.e., *jaw-D* (Palatnik et al., 2003). The expression pattern of *CUC2* was highly similar in Col-0 and *jaw-D* leaf primordia (**Fig 7A**). To support the observation that *JAW-TCPs* do not transcriptionally downregulate *CUC2* expression during early stages of leaf primordia, we studied the domain of *pCUC2::GUS* expression in seedlings heterozygous for Col-0;*GR*[#1] transgene, i.e., Col-0 x Col-0;*GR*[#1] F1. The expression pattern of *pCUC2::GUS* did not vary significantly between ethanol and DEX-treated early Col-0 x Col-0;*GR*[#1] F1 leaf primordia (**Fig S5C**), suggesting *JAW-TCPs* do not affect *CUC2* transcription in early leaf primordia. JAW-TCPs also suppress *CUC2* and *CUC3* activities at the post-transcriptional level via miR164a upregulation (Koyama et al., 2010) and at the protein level through physical interaction (Rubio-Somoza et al., 2014). Hence, to test if JAW-TCPs downregulate *MIR319C* expression at least partly by modulating *CUC2* activity, we studied *MIR319C* promoter activity in mutants with altered *CUC2* and *JAW-TCP* levels.

First, we looked at the *MIR319C* expression domain in Col-0;*GR^#2* (Shankar et al., 2023) and

cuc2-1D;GR^#2 leaf primordia where *rTCP4* activity can be induced by DEX treatment in wild-

type and *cuc2-1D* genetic backgrounds, respectively. Under mock conditions, the

pMIR319C::GUS domain was distally expanded in *cuc2-1D;GR^#2* leaf primordia compared to

Col-0;*GR^#2* control (median~0.55 in *cuc2-1D;GR^#2* mock compared to ~0.4 in Col-0;*GR^#2*

mock) (**Fig 7B, 7C**). While *rTCP4* induction greatly restricted the *MIR319C* expression

proximally in the Col-0 background (median~0.25), the *pMIR319C::GUS* domain was only

restricted to near wild-type level (median~0.35) when *rTCP4* activity was induced in the

cuc2-1D background (**Fig 7B, 7C**), indicating that *MIR319C* expression domain is established

by *CUC2* and *JAW-TCPs* via independent pathways.

We also looked at *MIR319C* expression domain in mutants where *miR319a* is overexpressed

in *cuc2-1* and *cuc2-1D* mutant backgrounds. Similar to earlier results in (**Fig 1D, 1E**), the loss

of *CUC2* activity (*cuc2-1*) alone reduced the domain of *MIR319C* expression (median~0.3)

compared to wild-type (median~0.45) (**Fig 7D, 7E**). In contrast, the downregulation of all the

JAW-TCPs (*pBLS>>miR319a*) or elevation of *CUC2* activity (*cuc2-1D*) resulted in the distal

expansion of the *MIR319C* expression domain (median~0.5) (**Fig 7D, 7E**). The *MIR319C*

expression domain in *cuc2-1;pBLS>>miR319a* leaf primordia (median~0.4) was similar to that

of the wild-type. In addition, *cuc2-1D;pBLS>>miR319a* leaf primordia did not show a further

increase in *MIR319C* expression domain compared to either of its parent genotypes (**Fig 7D,**

7E). These results support the earlier conclusion that *CUC2* and *JAW-TCPs* independently

regulate the *MIR319C* expression domain in early leaf primordia.

Discussion

The miR319-CIN-TCP module regulates leaf morphology in diverse eudicots by affecting the

timing of pavement cell proliferation to differentiation transition (Palatnik et al., 2003; Efroni

et al., 2008; Shleizer-Burko et al., 2011; Challa et al., 2019). Although the molecular

mechanisms for miR319-CIN-TCP mediated regulation of leaf morphology had been described

(Sarvepalli and Nath, 2018; Challa et al., 2019; Challa and Rath et al., 2021), the genetic basis

for the transcriptional activation of *MIR319* and *CIN-TCP* genes during leaf growth was

unknown. In this study, we provide evidence to suggest that CUCs bind to the *MIR319C*

promoter and activate transcription in leaf primordia. All three CUCs specifically bound to the

MIR319C distal promoter fragment that harboured a unique CUC binding site, TTACGT, in

yeast (**Fig 1C**). Mutations in the CUC binding site resulted in more proximal restriction of

MIR319C expression domain (**Fig 5D-F**), a phenotype that was also observed in the *cuc2-1*

mutant (**Fig 1D, 1E**). While *cuc2-1* had lower pre-*MIR319C* levels, transient induction of

CUC2 activity in presence of protein synthesis inhibitor was sufficient to increase the cellular

pre-*MIR319C* transcript levels (**Fig 1F, 5G**), suggesting CUC2 activity is both required and

sufficient to promote *MIR319C* transcription. However, it remains to be clarified whether

CUC2 directly binds to the cis element on the *MIR319C* promoter in planta and if CUC2

requires additional factors to activate *MIR319C* expression. Nevertheless, the evidence from

this study suggests a model where ubiquitous *CUC2* expression in the initiating leaf primordia

activates *MIR319C* transcription which results in high miR319c activity, whereas during later

stages of primordia growth, *CUC2* expression and hence miR319c activity gets restricted

towards the SAM-leaf boundary and proximal leaf margins (**Fig S6**).

What is the significance of *CUC2*-mediated activation of *MIR319C* in leaf primordia?

In developing leaves, miR319-directed cleavage of *JAW-TCP* transcripts is required for proper

timing of the transition between cell division and differentiation (Palatnik et al., 2003; Efroni

et al., 2008; Challa et al., 2019). Interestingly, loss of *CUC2* barely affects cell proliferation in

leaves, suggested by a small yet consistent decrease in leaf size chiefly due to reduced cell

number (**Figs 2, 4, S3A, S3B**). The negative effect of *CUC2* loss on leaf cell proliferation is

likely due to reduced miR319c levels as a result of partial downregulation of *MIR319C*

promoter activity (**Fig 1D, 1E**). The role of CUC2-mediated activation of *MIR319C* in leaf cell

proliferation was further revealed by the strong reduction in cell numbers in mature leaves of

the *cuc2-3;miR319c-1* double mutant (**Fig 2**). *miR319c-1* has a T-DNA insertion 528 bp

upstream of the pre-*MIR319C* transcript (**Fig S4A**). *GUS*-based *MIR319C* promoter deletion

reporter line carrying only 533 bp showed reduced promoter activity as compared to 1866 bp

and full length (2736 bp) promoter lines (**Fig S4B**). Thus, the presence of 4.7 kb T-DNA in the

MIR319C promoter is likely to negatively impact the transactivation ability of TFs (including

CUC2) that bind to distal region, leading to reduced miR319c levels. We speculate that

cuc2-3;miR319c-1 mutant has lower levels of miR319c than the individual parent genotypes

due to additive loss of *CUC2* and other TF-mediated activation of *MIR319C* transcription. The

cellular analysis during early stages of *cuc2-3;miR319c-1* leaf primordia growth revealed an

increased proportion of enlarged cells as early as 3 DAI, which persisted even at 5 DAI

(**Fig 3**). These observations suggest a scenario where reduced miR319c levels leads to a
precocious transition of pavement cells from proliferation into differentiation due to elevated
JAW-TCP activity, particularly at the base of the leaf primordia. The hypothesis is further
supported by epistatic interaction observed between *CUC2* and *JAW-TCPs*, where miR319
overexpression led to increased leaf size even in *cuc2-1* background and ectopic *JAW-TCP*
activity resulted in smaller than wild-type leaves in *CUC2* overexpression and wild-type
backgrounds (**Fig 6**). In sum, we demonstrate *CUC2* as an activator of *MIR319C* expression in
leaf primordia and propose a novel function for *CUC2* in promoting pavement cell division
during early stages of leaf growth via *MIR319C* activation. Using Y1H approach, we
uncovered other potential activators of *MIR319C* transcription such as members of YABBY,
LBD, and WOX TF families (**S4 Table**), which show highly overlapping expression patterns
with *MIR319C* and are known to have roles in leaf growth (Wu et al., 2008; Sarojam et al.,
2010; Nakata et al., 2012). It will be an interesting future prospect to investigate whether these
TFs directly associate with *MIR319C* promoter and activate its expression in planta.

Role of *CUC2-MIR319-JAW-TCP* module in pavement cell division and growth

miR319a;miR319b and *miR319c-1* mutants had wild-type leaf areas whereas the *miR319abc*
triple mutants displayed smaller leaves with fewer cells of unchanged areas (**Figs 5A, 5B, 4D,**
S5A, and S5B). The leaf and cell phenotypes of *miR319abc* were similar to *rTCP4*
overexpression lines, supporting the conclusion from earlier kinematic studies that *miR319-*
JAW-TCP module regulates pavement cell proliferation to differentiation transition, but has
little effect on post-mitotic cell expansion (Challa et al., 2019). Together the above
observations strongly suggest a functional redundancy among *MIR319* genes in suppressing
JAW-TCP activity at the base of the primordia. While *miR319b* mutant completely lacks
miR319b activity (Koyama et al., 2017), both *miR319a* and *miR319c-1* lines are likely to retain
low levels of miR319a and miR319c activities, respectively (Nag et al., 2009; Koyama et al.,
2017) (**Fig S4A, S4B**). We used knockdown alleles of *miR319a* and *miR319c* to study genetic
interaction with *CUC2* and among *MIR319* genes as *miR319* knockouts are expected to have
high levels of ectopic *JAW-TCP* activity which may lead to embryo lethality and/or seedling
arrest (Palatnik et al., 2003; Koyama et al., 2017). The F2 population from the cross between
cuc2-3/cuc2-1 and *miR319ab* exhibited a range of phenotypes that included narrow leaves and
fused cotyledons (**Fig S3A, S3F**), suggesting a role for *CUC2-MIR319* pathway in leaf growth
and cotyledon separation. To our surprise, even the homozygous *cuc2-3;miR319a* and

cuc2-3;miR319b population and the *cuc2-3;miR319ab* triple mutants displayed similar range of leaf and cotyledon defects (**Fig 4**), probably due to penetrance in miR319 activity. The roles of the *CUC2-MIR319* module in SAM establishment and cotyledon separation have been demonstrated earlier (Koyama et al., 2007; Koyama et al., 2010; Koyama et al., 2017), but its role in pavement cell proliferation has not been explored. Interestingly, unlike the *miR319abc* mutants which had fewer cells of wild-type areas, the *cuc2-3;miR319ab* leaves had lower cell numbers and cell areas (**Fig S5**). What could be the underlying cause for the difference in cell growth pattern between *miR319abc* and *cuc2-3;miR319ab* mutants? Previous studies have demonstrated that severe defects in leaf cell proliferation often induces post-mitotic cell expansion leading to larger than wild-type cells. This phenomenon of coordinated cell growth during leaf development is termed as compensation (Tsukaya, 2005). Recent genetic and kinematic studies suggests that compensation is not a passive result of low cell proliferation but rather a genetically regulated active mechanism (Hisanaga et al., 2015; Tsukaya, 2019). Moreover, compensation is highly dependent on the extent of changes in causal gene activity and can involve both cell autonomous and non-autonomous mechanisms (Hisanaga et al., 2015; Vercruysse et al., 2021). We speculate that *CUC2* may promote compensation in leaves independent of *MIR319C* activation. Thus, the lack of *CUC2* and *MIR319A/B* activities would lead to premature proliferation arrest due to ectopic *JAW-TCP* activity and impaired compensation via an unknown genetic pathway, resulting in leaves with reduced cell numbers and areas. Previous studies have demonstrated that JAW-TCPs repress the activities of all three *CUCs* at the mRNA and protein levels (Koyama et al., 2010; Rubio-Somoza et al., 2014). Hence, ectopic *JAW-TCP* activity would feedback to further downregulation of *CUC* activity and accentuate the cell proliferation and expansion defects in *cuc2-3;miR319ab* mutants during leaf growth.

How is the *MIR319C* expression domain established and maintained in leaf primordia?

Here, we demonstrate that CUC2 activates *MIR319C* expression throughout the initiating leaf primordia and at the base during later stages of leaf growth. We previously showed that JAW-TCPs bind to *MIR319C* and repress its transcription in the distal region of the early leaf primordia (Shankar et al., 2023). Interestingly, *CUC2* is initially expressed throughout the primordia but restricted towards the base around the 150 μm stage coinciding with the onset of distal *TCP4* expression (**Fig 7**). This raised a possibility that these two processes are linked and

lead to the restriction of *MIR319C* expression. However, we show that JAW-TCPs repress *MIR319C* expression independent of *CUC2* activity (**Fig 7**), suggesting that *CUC2*-mediated activation and *JAW-TCP*-mediated repression establish the basal *MIR319C* expression domain via independent mechanisms. Although *MIR319C* expression highly coincided with the cell proliferation zone, we previously demonstrated that *MIR319C* expression domain remained unchanged in mutants with altered cell proliferation pattern (Shankar et al., 2023). This evidence suggests that changes in the proliferation status of cells may not feedback to regulate *MIR319C* expression. Previous modelling-based studies on leaf morphogenesis suggest that cell growth induced mechanical forces could feedback and regulate growth and gene expression (Kuchen et al., 2012; Fox et al., 2018). It will be interesting to investigate if such feedback mechanisms play role in maintenance of basal *MIR319C* expression domain. Additionally, some of the outstanding questions regarding the establishment of *MIR319C* expression domain are as follows: Which genetic factors specifically promote *JAW-TCP* expression in the distal region of the 150 μm primordia? How is *CUC2* expression, which is ubiquitous during leaf initiation, later restricted towards the SAM-leaf boundary and proximal margins of the leaf primordia? Answers to these questions will advance our understanding of the molecular mechanisms that divide the uniformly proliferating primordia into two contrasting domains of cell proliferation and differentiation and subsequently coordinate growth across these zones during leaf morphogenesis.

Together, our data is consistent with a model where *CUC2* upregulates *MIR319C* expression to maintain high miR319c levels, which together with miR319a and miR319b, restricts JAW-TCP activity towards the distal region of the growing leaf primordia. High JAW-TCP protein levels due to transcriptional activation leads to repression of *MIR319C* promoter activity in the distal region (Shankar et al., 2023). CUC2-mediated activation of *MIR319C* and the miR319-JAW-TCP feedback loop contributes to the establishment of spatially separated growth domains of basal cell proliferation and distal cell differentiation in leaf primordia (**Fig S6**). We propose that the *CUC2-MIR319-JAW-TCP* module is critical for spatiotemporal patterning of cell proliferation in Arabidopsis leaf primordia. As the *CUC*, *MIR319*, and *JAW-TCP* genes are conserved across plant evolution, it will be worthwhile to investigate if *CUC2-MIR319-JAW-TCP* module regulates leaf growth in other plant species, especially those that exhibit basipetal mode of growth.

Context-dependent effect of *CUC2* on cell proliferation during organ morphogenesis

During embryogenesis, all three *CUCs* are expressed at the junction between the SAM and the cotyledons, where they redundantly repress cell proliferation to establish a boundary zone and allow for the separation and subsequent outgrowth of the cotyledons (Vroemen et al., 2003; Hibara et al., 2006; Scofield et al., 2018). Later on during leaf morphogenesis, CUC2 protein accumulates at certain foci (termed sinus) along the leaf margin and induces serration formation by inhibiting cell proliferation locally at the sinus region (Nikovics et al., 2006; Bilsborough et al., 2011; Hasson et al., 2011). In addition, CUC2 promotes serration outgrowth (termed tooth) via a non-cell autonomous mechanism involving *CUC3*, cytochrome P450 protein encoding *KLUH/CYP78A5*, and auxin response (Maugarny-Calès et al., 2019). These studies point towards a dual role for CUC2 during serration formation and outgrowth where it initially represses cell proliferation at the sinus region but promotes cell proliferation-induced tooth outgrowth during later stages. Interestingly, expression of *miR164*-resistent *CUC2* in the boundary zone during flower development led to a significant enlargement of boundaries between sepal primordia due to increased cell proliferation (Laufs et al., 2004; Sieber et al., 2007). Here, we show that *CUC2* promotes and maintains pavement cell proliferation via the miR319-JAW-TCP pathway in early leaf primordia. Our data supports the notion that *CUC2* can either activate or repress cell proliferation depending on the developmental stage and the type of tissue it is expressed and regulate organ morphology. So far, only a few direct downstream targets of CUCs have been identified (Takeda et al., 2011; Li et al., 2020; Aida et al., 2020). By integrating spatial transcriptomics, whole-genome binding, and proteomics experiments, future research could unveil the molecular mechanisms by which CUC2 exerts context-dependent activation and repression.

Do *CUCs* act as growth polarity organisers during leaf morphogenesis?

Evidence from previous kinematics, genetics and modelling-based studies suggest that morphogenesis of a simple cone-like primordium to complex planar mature leaf involves highly coordinated spatiotemporal regulation of key parameters such as cell growth and division rates, growth orientation and tissue deformation (Kuchen et al., 2012; Fox et al., 2018; Sarvepalli et al., 2019). Growth polarity is apparent in early primordia where the proximal region grows at a faster rate due to high cell proliferation, whereas distal region displays lower growth rates and transitions into differentiation (Kuchen et al., 2012; Fox et al., 2018). Previous

studies proposed *CUC* as a growth polarity organizer, which, according to growth prediction models is a factor that is highly expressed at the base and drives differential growth by upregulating factors that promote growth and proliferation (Kuchen et al., 2012; Sarvepalli et al., 2019). As mentioned earlier, CUCs are highly expressed at the boundaries and can regulate growth at leaf margin by modulating auxin transport, activating factors that either promote or repress cell proliferation (Bilsborough et al., 2011; Maugarny-Calès et al., 2019). It remains to be seen if CUC-mediated promotion of pavement cell proliferation also involves some of these mechanisms that exist in the context of serration outgrowth. Here, we demonstrate a novel role of CUC2 in upregulating miR319c levels that results in low proximal JAW-TCP activity in leaf primordia. JAW-TCPs are known to inhibit cell proliferation factors including GROWTH REGULATION FACTOR (GRF) and GRF- INTERACTING FACTOR (GIF) which promote growth and proliferation via activation of several cell cycle genes, including *CYCLINB1;1* (Horiguchi et al., 2005; Schneider et al., 2024). Overall, CUC2-mediated activation of *MIR319C* would enhance growth and proliferation in the proximal region, thus providing proof for the role of *CUCs* as growth polarity organizer during leaf morphogenesis. In animal models, it has been well established that boundaries function as morphogen providers to pattern organ growth during embryogenesis (Meinhardt, 2009; Kicheva and Briscoe, 2023). The role of CUC2 as a growth polarity organizer also raises an interesting possibility that boundaries can serve as growth organizers during organ morphogenesis in plants. Future research could explore mechanisms that link boundary gene networks with growth regulatory modules to test this possibility.

Materials and Methods

In silico screening for SAM and leaf primordia-specific transcription factors

To identify SAM and leaf primordia-specific TFs, we took advantage of studies that generated cellular-resolution maps of genes whose mRNA is actively translated in different domains within the SAM and leaf primordia (Tian et al., 2014; 2019). We considered genes that are active in the *CLV3*, *WUS*, *UFO* expression domains of the SAM, *LAS* expression domain of the SAM-leaf boundary, and the *AS1* expression domain of the leaf primordia. We overlapped the 4426 SAM and leaf primordia-specific genes with the list of 1956 genome-wide Arabidopsis TFs (Pruneda-Paz et al., 2014) and obtained 343 TFs expressed explicitly in the SAM and leaf primordia domains **S1 Table**. We also scanned the 2.7 kb URR of *MIR319C* for

TF binding sites using the publicly available transcription factor binding site (TFBS) prediction tool (PlantPAN 3.0 - *http://plantpan.itps.ncku.edu.tw*). We recovered several potential DNA-binding motifs on the *MIR319C* promoter, including TCP, KNOX, YABBY, NAC and HD-like TF-binding consensus. We listed 49 representatives from the top 5 TF families recovered from the TFBS prediction software **S2 Table**. Next, we combined this list with the previously obtained 343 SAM and leaf primordia-specific TFs to get a final list of 378 TF candidates for the yeast one-hybrid screen **S3 Table**.

Yeast transformation

Yeasts were transformed with the desired DNA constructs (plasmids) by the high-efficiency LiAc-PEG method described by Walhout and Vidal, 2001, with minor modifications. A single colony of host strain was inoculated in 10 ml YPAD (Bacto yeast extract-1g, Tryptone-1g, Dextrose-2g, and Adenine sulphate-10mg per 100ml) and incubated overnight at 30ºC with constant shaking. The next day, the suitable culture volume was inoculated into 200 ml of fresh YPAD solution to an initial OD of 0.2 and kept at 30ºC with constant shaking until an OD of 0.6-0.8 was reached. Cells were pelleted down at 700 g for 10 min and washed twice with 5 ml of sterile water. The cells were washed with 2.5 ml of LiAc-TE (0.1 M LiAc, Tris- 10 mM, EDTA- 1mM) solution. Cells were finally resuspended in 2 ml of LiAc-TE solution. A mixture containing 280 μL of 50% PEG (Mol wt 3350, Sigma-Aldrich, USA), 36 μL of 1 M LiAc in TE, 6 μL of 10 mg/ml boiled carrier DNA (D-3159, Sigma-Aldrich, USA) was prepared and added to 1.5 ml microfuge tubes containing 60 μL cells and 100 ng of DNA construct. The cell suspension was vortexed briefly and incubated at 30ºC with constant shaking for 30 min. Cells were transferred into a water bath maintained at 42ºC for 30 min. Cells were pelleted, washed twice with sterile water, resuspended in 100 μL sterile water, and plated onto appropriate media.

For the Y1H screen, transformations were carried out in 96 well-format described by Walhout and Vidal, 2001. Yeast cells were prepared as described in the high-efficiency protocol. 20 μL of cell suspension (containing LiAc-PEG-carrier DNA) were transferred to each well of a sterile U-bottom 96-well plate (3795, Corning, USA). 100 ng of plasmid and 100 μL of TE/LiAc/PEG solution (50% PEG and 0.1 M LiAc in TE) were individually added and resuspended carefully using a multichannel pipette. 96-well plate was incubated at 30ºC with constant shaking for 30 min. Cells were transferred to a water bath maintained at 42ºC for 30 min. Cells were pelleted at 700 g for 5 min. To each well, 120 μL of sterile water was added,

608   and 100 μL was removed directly to get rid of PEG in the supernatant. Finally, cells were

609   resuspended in the remaining 20 μL water and spotted on appropriate media.

Yeast bait, prey, and diploid generation

The full length (2.7 kb) and the non-overlapping promoter fragments, i.e., -2694-1866 (p^{distal}),

-1866-1107 (p^{middle}), and -1107+35 ($p^{proximal}$) of *MIR319C* were amplified from *pMIR319C-*

P4P1R entry clone using Taq DNA polymerase and cloned into pENTR/D-TOPO vector

(K240020, Invitrogen, USA) according to the manufacturer's instructions. The entry clones

were confirmed by Sanger sequencing before cloning upstream to the *minimalPro::HIS3*

cassette of *pMW#2* (Gift from Ram Yadav, IISER Mohali, INDIA) by LR reaction. Y1H baits

were generated according to the protocol described by Fuxman Bass et al., 2016. Briefly, the

pMIR319C containing *pMW#2* reporter constructs were linearized with *Xho*I (R0146, New

England Biolabs, USA) and integrated into the mutant *HIS3* locus of the *YM4271* host strain

(Gift from Ram Yadav, IISER Mohali, INDIA) using the high-efficiency LiAc-PEG

transformation method described by Walhout and Vidal, 2001. Following transformation, yeast

cells were plated on the synthetic complete media lacking histidine (Sc-His) media (Walhout

and Vidal, 2001) and incubated at 30ºC till colonies appeared. Post growth, genomic DNA was

extracted individually from 3-4 colonies, and the integration of the *MIR319C* promoter

fragment was confirmed by PCR. All primer sequences used for cloning are mentioned in **S5**

Table.

To generate the prey strains for the Y1H screen, each of the 384 TF-encoding plasmids was

individually purified from corresponding glycerol stocks of the 1956 DEST-U Arabidopsis TF

ORFeome collection (CD4-89, ABRC, USA) using QIAprep Spin Miniprep Kit. 100 ng of

purified plasmid were individually transformed into Yalpha1867 strain (Gift from Dr. Ram

Yadav, IISER Mohali, INDIA) using the high throughput 96-well format transformations

protocol described earlier. Following transformation, the cells were spotted on synthetic

complete media lacking tryptophan (Sc-Trp) media and incubated at 30ºC till colonies

appeared.

To generate the diploids used in the Y1H assay, prey TF-encoding plasmids was individually

purified from corresponding glycerol stocks of the 1956 DEST-U Arabidopsis TF ORFeome

collection (CD4-89, ABRC, USA) using QIAprep Spin Miniprep Kit. 100 ng of purified

plasmid were individually transformed into *pMIR319C* baits using the high-efficiency LiAc-

PEG method described earlier. Following transformation, the cells were spotted on synthetic

complete media lacking histidine and tryptophan (Sc-His-Trp) media and incubated at 30ºC till

colonies appeared.

Yeast one-hybrid (Y1H) screen/ assay

The auto activity of yeast baits on Sc-His due to minimal *HIS3* expression was determined

by spotting them on Sc-His media containing 0-30 mM 3-Amino-1,2,4-triazole (3-AT)

(A8056, Sigma-Aldrich, USA). The minimum concentration of 3-AT that completely inhibits

the growth of the bait was considered for the Y1H screen/ assay. For the mating-based Y1H

screen, the diploids generated by mating were resuspended in sterile 96-well plates containing

Sc-Trp-His media and allowed to grow in shaking incubators for 24 hrs at 30ºC. After 24 hrs

of growth, OD was measured. Volumes corresponding to equal OD were spotted on Sc-Trp-

His+3AT media, incubated at 30ºC, and scored for growth on days 3-5 post-incubation. Spots

containing at least 5 colonies were considered positive for growth. For the Y1H screen, spots

showing growth on all three concentrations of 3-AT, i.e., 20, 25, and 30 mM by day 5, or

growth on 20 and 25 mM 3-AT by day 3 were considered positive for binding. For the co-

transformation-based Y1H assay, yeast baits were transformed with TF encoding plasmids and

selected on Sc-Trp-His media. Following selection, the cells were resuspended in Sc-Trp-His

media and allowed to grow in shaking incubators for 24 hrs at 30ºC. After 24 hrs of growth,

OD was measured. Volumes corresponding to equal OD were spotted on Sc-Trp-His media

with respective concentrations of 3-AT and incubated at 30ºC.The plates were inspected for

growth and imaged over days 3- 5 post-incubation.

Plant materials

Arabidopsis thaliana ecotype Columbia-0 (Col-0) was used as the wild-type control. All the

lines used in this study are in Col-0 genetic background. The genotypes used in this study are

described in **S6 Table**. The genotypes *cuc2-1* and *pBLS>>miR319a* (Originally in L*er*; Efroni

et al., 2008; Hasson et al., 2011) was brought into the Col-0 background by back crossing three

times. *pMIR319C::GUS*[#1] (Nag et al., 2009) was established in various mutant backgrounds by

crossing and was confirmed by GUS assays in the F3/F4 generation. *cuc2-3;miR319c-1*,

cuc2-3;miR319ab and *miR319abc* mutants were generated by crossing respective mutant lines

cuc2-3 (Nikovics et al., 2006), *miR319ab* (Koyama et al., 2017), *miR319c-1* (SAIL_647_B03/

CS827925) and confirmed by genotyping in F3 generation. *cuc2-1;GR*[#2] was established by

crossing *cuc2-1* and Col-0;*GR*[#2] (Shankar et al., 2023) mutants and confirmed by phenotyping

in F3 generation. All genotyping primers are listed in **S5 Table**.

Plant growth condition and chemical treatments

For GUS assays and phenotyping experiments, seeds were surface-sterilized with 0.05% SDS

containing 70% ethanol solution for 10 min, followed by wash with 100% ethanol twice and

sown on 1X Murashige and Skoog (MS), 0.8% agar medium (PT011, HiMedia Laboratories,

India) containing 0.025% ethanol (Mock) or 6 μM dexamethasone (DEX) (D4902, Sigma-

Aldrich, USA). After sowing, seeds were kept for stratification for three days in the dark at

4°C and then shifted to a plant growth chamber (I-41LL, Percival Scientific, USA) set at 22°C,

60% relative humidity, long-day conditions (16h/8h light/dark period, 100 $\mu mol.m^{-2}.s^{-1}$). 14-

day old seedlings were transplanted onto a soil mixture containing Soilrite and Perlite (Keltech

Energies, India) in a 3:1 ratio. Plants were grown till maturity in growth rooms with the

conditions as described for growth chamber.

For transient DEX and cycloheximide (CHX) treatments, 9 day-old mock (1X MS, 0.8% agar

medium containing 0.025% ethanol) grown seedlings were transferred onto petri dishes filled

with 0.5X MS medium, supplemented with 1% sucrose, 1X Gamborg B5 vitamins (PL018,

HiMedia Laboratories, India), 12 μM DEX, and/or 10 μM CHX (C7698, Sigma-Aldrich,

USA), incubated for 2 hours and harvested under liquid N2 prior to RNA isolation. In the case

of phenotyping experiment to study embryonic and cotyledon defects in *cuc2-1;GR*[#2]

seedlings, 14 day old mock-grown *cuc2-1;GR*[#2] seedlings were transferred onto soil mixture

(as described earlier) and incubated in growth rooms until bolting stage (growth conditions as

described for growth chamber). Upon bolting, plants were sprayed with 12 μM DEX solution

on alternate days until seed harvest.

For confocal imaging experiments, seeds were stratified in the dark for two days at 4°C to

synchronize germination. Plants were grown on petri dishes filled with 0.5X MS medium,

supplemented with 0.8% agar, 1% sucrose, and 0.1% Plant Protective Mixture (Plant Cell

Technology) in a growth chamber under long-day conditions (16h/8h light/dark period, 95

$\mu mol.m^{-2}.s^{-1}$) at 22 ± 1°C with 60-70% relative humidity.

DNA constructs and transgenic line generation

For the generation of *pMIR319C* deletion and *mCUCBSpMIR319C::GUS* constructs, a 2.7 kb genomic fragment corresponding to the *MIR319C* URR from the start of pre-*MIR319C* was amplified from Col-0 genomic DNA using Phusion High Fidelity DNA polymerase (M0530, New England Biolabs, USA) and cloned into modified *P4P1R* entry vector (a kind gift from Ram Yadav, IISER Mohali, India) by gateway BP reaction (11789020, Invitrogen, USA) to generate the *pMIR319C-P4P1R* construct. To introduce mutation at the CUC-binding site, we utilized a PCR-based site directed mutagenesis approach. Briefly, forward, and reverse primers carrying mutations in CUC-binding site (mCUCBS) was first used in a PCR reaction with *pMIR319C-P4P1R* template using Phusion DNA polymerase. Post PCR reaction, the product was digested with *Dpn*I (R0176, New England Biolabs, USA) and transformed into *E. coli* and selected on LB agar containing 100 mg/l ampicillin. Plasmids were purified from resulting colonies using QIAprep Spin Miniprep Kit (Qiagen, Germany) and mCUCBS was confirmed by Sanger sequencing. The entry clone *mCUCBSpMIR319C-P4P1R* was recombined into *R4L1pGWB532* destination vector (Gift from Tsuyoshi Nakagawa, Shimane University, Japan) by LR reaction (11791020, Invitrogen, USA) to generate the *mCUCBSpMIR319C::GUS* construct. Sequence information of primers used to generate the constructs are provided in the **S5 Table**.

mCUCBSpMIR319C::GUS construct was integrated into Col-0 background by the *Agrobacterium tumefaciens*-mediated floral dip method (Clough and Bent, 1998) except Silwet L-77 was used at a concentration of 50 µl/litre. Following transformation, seeds were harvested and screened for the ability to grow on MS media in presence of 20 mg/l hygromycin. At least five independent lines were established for each construct and confirmed by genotyping in T3 generation. Sequence information of primers used for genotyping are provided in the **S5 Table**.

β-glucuronidase (GUS) assay

GUS assays were performed as described earlier (Shankar et al., 2023). Briefly, samples were collected in ice-cold 90% acetone, incubated at room temperature for 20 min, followed by washing with ice-cold staining buffer (50 mM sodium phosphate, pH 7.0, 0.2% Triton X-100, 5 mM potassium ferrocyanide, and 5 mM potassium ferricyanide). Fresh staining buffer was added with 2 mM X-Gluc (R0851, Thermo Scientific, USA) and incubated overnight at 37°C in dark. Staining buffer was replaced with 70% ethanol, and 70% ethanol was changed 3-4

times in regular intervals to clear the tissue. Shoot apices and leaf primordia were dissected and mounted on a glass slide containing lactic acid under the microscope and were imaged using Olympus BX51 trinocular DIC microscope fitted with ProgResC3 camera and ProgResCapture Pro2.6 software (Jenoptik, Germany) with the following settings, exposure-65ms, brightness-10 units, contrast- 35 units, colour temperature- 5000K, image resolution-(1040 x 770 2X Bin), while keeping all other settings in default mode.

GUS domain, leaf area and pavement cell area quantification

The extent of reporter gene expression in the *pMIR319C::GUS* and *mCUCBSpMIR319C::GUS* transgenic lines were determined using a pixel density quantification method reported in Shankar et al., 2023. Images of 6 and 9-day old seedlings were captured using LEICA EZ4 E stereomicroscope equipped with LAS EZ software. Areas of the dissected mature first pair of leaves from 25-day old plants were determined using the polygon selection tool of the ImageJ software. For pavement cell area measurements, dissected mature first pair of leaves were first incubated in 7:1 absolute ethanol: glacial acetic acid clearing solution. After 24 hours, clearing solution was replaced with 1M KOH solution for 30 minutes, washed twice with sterile water and incubated in 37% D-lactic acid solution until tissue became completely transparent. The clarified leaf samples were mounted on a glass slide using D-lactic acid and imaged using Olympus BX51 trinocular DIC microscope fitted with ProgResC3 camera and ProgResCapture Pro2.6 software (Jenoptik, Germany) with similar settings as described earlier. Cell areas were determined using the polygon selection tool of the ImageJ software.

Confocal microscopy and imaging

For imaging of the first pair of leaf, one cotyledon was removed using fine tweezers or an injection needle from two-day-old plants to expose the initiating leaf primordium for imaging of leaf at 1 day after initiation (DAI). The seedlings were stained with 1 mg/mL Propidium Iodide (PI) solution for 2-3 minutes and then were rinsed with water. The dissected plants then were placed horizontally in Ø60 mm Petri dishes filled with 0.5X MS medium, supplemented with 1.5% agar, 1% sucrose, and 0.1% Plant Protective Mixture (Plant Cell Technology). Plants were immersed in water containing 0.1% PPM for imaging. Three to four replicates were imaged using the PI staining and at least half of the abaxial leaf surface was imaged at 1 DAI, 3 DAI and 5 DAI respectively.

All confocal imaging was performed with an LSM980 upright confocal microscope using a
long-distance water-dipping objective (AP 20X/1.0; Zeiss) at the confocal facility of
MechanoDevo team (RDP, ENS de Lyon). Excitation was performed using a diode laser with
561 nm for PI. Images were collected at 580-650 nm for PI with a GAsP detector and a beam
splitter operating at 488 and 561 nm. Confocal stacks were acquired at 512x512 resolution and
16-bit image depth, with 1.4 µm distance in Z-dimension. For samples larger than the field of
view, multiple overlapping stacks were obtained and later stitched together using
MorphoGraphX (Barbier de Reuille et al., 2015; Strauss et al., 2022).
777

Confocal Image analysis

Cellular growth and expression quantifications were conducted using MorphoGraphX. Stacks
were processed as described previously (Le Gloanec et al., 2022). After digitally removing the
trichomes when necessary, the organ surface was detected with the 'Edge detect' tool with a
threshold of 20000 with 'closing' to fill holes, followed by the 'Edge detect angle' (10000). An
initial 5 µm cube size mesh was then created and subdivided three times before projecting the
membrane signal (2-5 µm). Cell segmentation was performed manually. Metrics such as area
cell size were computed as described before (Barbier de Reuille et al., 2015; Sapala et al.,
2018).
787

RNA isolation and RT-qPCR

Total RNA was extracted using the TRIzol method described earlier (Challa et al., 2016). 10
µg of RNA was used for DNase (EN0521, Thermo Scientific, USA) treatment. 1.5 µg DNase-
treated RNA was converted to cDNA using RevertAid Reverse Transcriptase (EP0441,
Thermo Scientific, USA) and 2.5 µM oligo dT according to the manufacturer's instructions.
Quantitative PCR reactions with 25 ng cDNA and 0.4 µM primers were set up using the
DyNAmo Flash SYBR Green RT-qPCR kit (F415L, Thermo Scientific, USA), according to
the manufacturer's instructions. 10 µl qPCR reactions were carried out in QuantStudio™ 6
Flex Real-Time PCR System, 384-well format (Applied Biosystems, USA). Results were
analyzed using the inbuilt QuantStudio6 Pro software, and $\Delta\Delta CT$ values were determined after
normalization with *PROTEIN PHOSPHATASE 2A SUBUNIT A3 (PP2AA3)*. Relative fold
changes in transcript levels were calculated using the formula $2^{-\Delta\Delta CT}$. All primer sequences are
provided in **S5 Table**.
801

Statistical analysis

All graphical representations and statistical analysis were performed using GraphPad Prism 9 software (GraphPad, USA). First, samples were tested for Gaussian distribution using the Shapiro-Wilk normality test. If samples were normally distributed, pairwise comparisons were performed using unpaired *t*-test. Welch's correction was applied if variances were different among the samples. Statistical differences among multiple genotypes were determined by ordinary One-Way ANOVA followed by Tukey's multiple comparisons test or One-Way ANOVA followed by Dunnet's T3 post-hoc test, depending on whether variances were significantly different among the genotypes. If the samples failed the normality test, then pairwise comparisons were carried out by Mann Whitney test. Statistical differences among multiple genotypes were determined by One-Way ANOVA followed by Dunn's post-hoc test. To compare statistical differences among multiple genotypes subjected to ethanol (Mock) or dexamethasone (DEX) treatments, we used Two-Way ANOVA followed by Tukey's post-hoc test. *p*-values obtained from the respective tests are indicated above the data bars in several plots. For plots with a large number of comparisons, statistical differences among samples are indicated by different alphabets.

References

Aida M, Ishida T, Fukaki H, Fujisawa H, Tasaka M (1997) Genes involved in organ separation in Arabidopsis: An analysis of the cup-shaped cotyledon mutant. Plant Cell **9**: 841–857

Aida M, Ishida T, Tasaka M (1999) Shoot apical meristem and cotyledon formation during Arabidopsis embryogenesis: interaction among the CUP-SHAPED COTYLEDON and SHOOT MERISTEMLESS genes. Development **126**: 1563–1570

Aida M, Tsubakimoto Y, Shimizu S, Ogisu H, Kamiya M, Iwamoto R, Takeda S, Karim MR, Mizutani M, Lenhard M, et al (2020) Establishment of the embryonic shoot meristem involves activation of two classes of genes with opposing functions for meristem activities. Int J Mol Sci **21**: 1–16

Anastasiou E, Kenz S, Gerstung M, MacLean D, Timmer J, Fleck C, et al (2007) Control of Plant Organ Size by KLUH/CYP78A5-Dependent Intercellular Signaling. Dev Cell 13: 843–56

Andriankaja M, Dhondt S, DeBodt S, Vanhaeren H, Coppens F, DeMilde L, Mühlenbock P, Skirycz A, Gonzalez N, Beemster GTS, et al (2012) Exit from Proliferation during Leaf Development in Arabidopsis thaliana: A Not-So-Gradual Process. Dev Cell **22**: 64–78

Bar M, Ori N (2014) Leaf development and morphogenesis. Development **141**: 4219–4230

Barbier de Reuille P, Routier-Kierzkowska A.L, Kierzkowski D, Bassel GW, Schüpbach T, Tauriello G, Bajpai N, Strauss S, Weber A, Kiss A et al (2015). MorphoGraphX: A platform for quantifying morphogenesis in 4D. Elife **4**: 05864

Barton MK (2010) Twenty years on: The inner workings of the shoot apical meristem, a developmental dynamo. Dev Biol **341**: 95–113

Bilsborough GD, Runions A, Barkoulas M, Jenkins HW, Hasson A, Galinha C, Laufs P, Hay A,
Prusinkiewicz P, Tsiantis M (2011) Model for the regulation of Arabidopsis thaliana leaf margin
development. Proc Natl Acad Sci U S A **108**: 3424–3429
Challa KR, Aggarwal P, Nath U (2016) Activation of *YUCCA5* by the Transcription Factor TCP4
Integrates Developmental and Environmental Signals to Promote Hypocotyl Elongation in
Arabidopsis. Plant Cell **28**: 2117–2130
Challa KR, Rath M, Nath U (2019) The CIN-TCP transcription factors promote commitment to
differentiation in Arabidopsis leaf pavement cells via both auxin-dependent and independent
pathways. PLoS Genet **15**: 1–30
Challa KR, Rath M, Sharma AN, Bajpai AK, Davuluri S, Acharya KK, Nath U (2021) Active
suppression of leaflet emergence as a mechanism of simple leaf development. Nat Plants **7**: 1264–
1275
Clough SJ, Bent AF (1998). Floral dip: A simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J **16**: 735–43
Donnelly PM, Bonetta D, Tsukaya H, Dengler RE, Dengler NG (1999) Cell cycling and cell
enlargement in developing leaves of Arabidopsis. Dev Biol **215**: 407–419
Du F, Guan C, Jiao Y (2018) Molecular Mechanisms of Leaf Morphogenesis. Mol Plant **11**: 1117–
1134
Efroni I, Blum E, Goldshmidt A, Eshed Y (2008) A Protracted and Dynamic Maturation Schedule
Underlies Arabidopsis Leaf Development. the Plant Cell Online **20**: 2293–2306
Fox S, Southam P, Pantin F, Kennaway R, Robinson S, Castorina G, Sánchez-Corrales YE,
Sablowski R, Chan J, Grieneisen V, et al (2018) Spatiotemporal coordination of cell division and
growth during organ morphogenesis. PLoS Biol **16**: e2005952
Fuxman Bass JI, Pons C, Kozlowski L, Reece-Hoyes JS, Shrestha S, Holdorf AD, et al (2016). A
gene-centered C. elegans protein–DNA interaction network provides a framework for functional
predictions. Mol Syst Biol **12**: 884
Gázquez A, Beemster GTS (2017) What determines organ size differences between species? A meta-
analysis of the cellular basis. New Phytologist **215**: 299–308
Gonzalez N, Vanhaeren H, Inzé D (2012) Leaf size control: Complex coordination of cell division
and expansion. Trends Plant Sci **17**: 332–340
Hasson A, Plessis A, Blein T, Adroher B, Grigg S, Tsiantis M, Boudaoud A, Damerval C, Laufs
P (2011) Evolution and diverse roles of the CUP-SHAPED COTYLEDON genes in Arabidopsis leaf
development. Plant Cell **23**: 54–68
Hepworth J, Lenhard M (2014) Regulation of plant lateral-organ growth by modulating cell number
and size. Curr Opin Plant Biol **17**: 36–42
Hibara KI, Karim MR, Takada S, Taoka KI, Furutani M, Aida M, Tasaka M (2006) Arabidopsis
CUP-SHAPED COTYLEDON3 regulates postembryonic shoot meristem and organ boundary
formation. Plant Cell **18**: 2946–2957
Hisanaga T, Kawade K, Tsukaya H (2015) Compensation: A key to clarifying the organ-level
regulation of lateral organ size in plants. J Exp Bot **66**: 1055–1063
Horiguchi G, Kim GT, Tsukaya H (2005) The transcription factor AtGRF5 and the transcription
coactivator AN3 regulate cell proliferation in leaf primordia of Arabidopsis thaliana. Plant Journal **43**:
68–78
Ishida T, Aida M, Takada S, Tasaka M (2000) Involvement of CUP-SHAPED COTYLEDON
genes in gynoecium and ovule development in Arabidopsis thaliana. Plant Cell Physiol **41**: 60–67
Kalve S, De Vos D, Beemster GTS (2014) Leaf development: a cellular perspective. Front Plant Sci
5: 1–25

Kazama T, Ichihashi Y, Murata S, Tsukaya H (2010). The mechanism of cell cycle arrest front
progression explained by a KLUH/CYP78A5-dependent mobile growth factor in developing leaves of
arabidopsis thaliana. Plant Cell Physiol **51**:1046–54

Kicheva A, Briscoe J (2023) Control of Tissue Development by Morphogens. Annu Rev Cell Dev
Biol 2023 **39**: 2023

Koyama T, Furutani M, Tasaka M, Ohme-Takagi M (2007) TCP Transcription Factors Control the
Morphology of Shoot Lateral Organs via Negative Regulation of the Expression of Boundary-Specific
Genes in Arabidopsis. Plant Cell **19**: 473–484

Koyama T, Mitsuda N, Seki M, Shinozaki K, Ohme-Takagi M (2010) TCP transcription factors
regulate the activities of ASYMMETRIC LEAVES1 and miR164, as well as the auxin response,
during differentiation of leaves in Arabidopsis. Plant Cell **22**: 3574–3588

Koyama T, Sato F, Ohme-Takagi M (2017) Roles of miR319 and TCP transcription factors in leaf
development. Plant Physiol **175**: 874–885

Larue CT, Wen J, Walker JC (2009) A microRNA-transcription factor module regulates lateral
organ size and patterning in Arabidopsis. Plant Journal **58**: 450–463

Laufs P, Peaucelle A, Morin H, Traas J (2004) MicroRNA regulation of the CUC genes is required
for boundary size control in Arabidopsis meristems. Development **131**: 4311–4322

Le Gloanec C, Collet L, Silveira SR, Wang B, Routier-Kierzkowska AL, and Kierzkowski D
(2022). Cell type-specific dynamics underlie cellular growth variability in plants. Development **149**:
dev200783.

Li Y, Xia T, Gao F, Li Y (2020) Control of plant branching by the CUC2/CUC3-DA1-UBP15
regulatory module. Plant Cell **32**: 1919–1932

Lindemose S, Jensen MK, De Velde J Van, O'Shea C, Heyndrickx KS, Workman CT,
Vandepoele K, Skriver K, Masi F De (2014) A DNA-binding-site landscape and regulatory network
analysis for NAC transcription factors in Arabidopsis thaliana. Nucleic Acids Res **42**: 7681–7693

Maugarny-Calès A, Cortizo M, Adroher B, Borrega N, Gonçalves B, Brunoud G, Vernoux T,
Arnaud N, Laufs P (2019) Dissecting the pathways coordinating patterning and growth by plant
boundary domains. PLoS Genet **15**: 1–30

Maugarny-Calès A, Laufs P (2018) Getting leaves into shape: A molecular, cellular, environmental
and evolutionary view. Development **145**: dev161646

Meinhardt H (2009) Models for the generation and interpretation of gradients. Cold Spring Harb
Perspect Biol **1**: a001362

Nag A, King S, Jack T (2009) miR319a targeting of TCP4 is critical for petal growth and
development in Arabidopsis. Proceedings of the National Academy of Sciences **106**: 22534–22539

Nakata M, Matsumoto N, Tsugeki R, Rikirsch E, Laux T, Okada K (2012) Roles of the middle
domain-specific WUSCHEL-RELATED HOMEOBOX genes in early development of leaves in
Arabidopsis. Plant Cell **24**: 519–535

Nikovics K, Blein T, Peaucelle A, Ishida T, Morin H, Aida M, Laufs P (2006) The balance
between the MIR164A and CUC2 genes controls leaf margin serration in Arabidopsis. Plant Cell **18**:
2929–2945

Ooka H, Satoh K, Doi K, Nagata T, Otomo Y, Murakami K, Matsubara K, Osato N, Kawai J,
Carninci P, et al (2003) Comprehensive Analysis of NAC Family Genes in Oryza sativa and
Arabidopsis thaliana. DNA Research **10**: 239–247

Palatnik JF, Allen E, Wu X, Schommer C, Schwab R, Carrington JC, Weigel D (2003) Control of
leaf morphotgenesis by microRNAs. Nature **425**: 257–263

Pruneda-Paz, JL, Breton, G, Nagel, DH, Kang, SE, Bonaldi, K, Doherty, CJ, Ravelo, S, Galli, M,
Ecker, JR, and Kay, SA (2014). A Genome-Scale Resource for the Functional Characterization of
Arabidopsis Transcription Factors. Cell Rep. **8**: 622–632

Rodriguez RE, Mecchia MA, Debernardi JM, Schommer C, Weigel D, Palatnik JF (2010)
Control of cell proliferation in Arabidopsis thaliana by microRNA miR396. Development 137:103–12
Rodriguez RE, Debernardi JM, Palatnik JF (2014) Morphogenesis of simple leaves: Regulation of
leaf size and shape. Wiley Interdiscip Rev Dev Biol 3: 41–57
Rubio-Somoza I, Zhou CM, Confraria A, Martinho C, Von Born P, Baena-Gonzalez E, Wang
JW, Weigel D (2014) Temporal control of leaf complexity by miRNA-regulated licensing of protein
complexes. Current Biology **24**: 2714–2719
Sapala, A, Runions A, Routier-Kierzkowska AL, Gupta MD, Hong L, Hofhuis H, Verger S,
Mosca G, Li CB, Hay A et al (2018). Why plants make puzzle cells, and how their shape emerges.
Elife **7**: e32794.
Sarojam R, Sappl PG, Goldshmidt A, Efroni I, Floyd SK, Eshed Y, Bowmana JL (2010)
Differentiating Arabidopsis shoots from leaves by combined YABBY activities. Plant Cell **22**: 2113–
2130
Sarvepalli K, Das Gupta M, Challa KR, Nath U (2019) Molecular cartography of leaf
development — role of transcription factors. Curr Opin Plant Biol **47**: 22–31
Sarvepalli K, Nath U (2018) CIN-TCP transcription factors: Transiting cell proliferation in plants.
IUBMB Life **70**: 718–731
Schneider M, Van Bel M, Inzé D, Baekelandt A (2024) Leaf growth – complex regulation of a
seemingly simple process. Plant Journal **117**: 1018–1051
Scofield S, Murison A, Jones A, Fozard J, Aida M, Band LR, Bennett M, Murray JAH (2018)
Coordination of meristem and boundary functions by transcription factors in the SHOOT
MERISTEMLESS regulatory network. Development **145**: dev157081
Shankar N, Sunkara P, Nath U (2023) A double-negative feedback loop between miR319c and
JAW-TCPs establishes growth pattern in incipient leaf primordia in Arabidopsis thaliana. PLoS Genet
19: 1–25
Shleizer-Burko S, Burko Y, Ben-Herzel O, Ori N (2011) Dynamic growth program regulated by
LANCEOLATE enables flexible leaf patterning. Development **138**: 695–704
Sieber P, Wellmer F, Gheyselinck J, Riechmann JL, Meyerowitz EM (2007) Redundancy and
specialization among plant microRNAs: Role of the MIR164 family in developmental robustness.
Development **134**: 1051–1060
Strauss, S, Runions A, Lane B, Eschweiler D, Bajpai N, Trozzi N, Routier-Kierzkowska AL,
Yoshida S, Rodrigues da Silveira S, Vijayan A et al (2022). Using positional information to
provide context for biological image analysis with MorphoGraphX 2.0. Elife **11**: e72601
Takeda S, Hanano K, Kariya A, Shimizu S, Zhao L, Matsui M, Tasaka M, Aida M (2011) CUP-
SHAPED COTYLEDON1 transcription factor activates the expression of LSH4 and LSH3, two
members of the ALOG gene family, in shoot organ boundary cells. Plant Journal **66**: 1066–1077
Tian C, Zhang X, He J, Yu H, Wang Y, Shi B, Han Y, Wang G, Feng X, Zhang C, et al (2014).
An organ boundary-enriched gene regulatory network uncovers regulatory hierarchies underlying
axillary meristem initiation. Mol Syst Biol **10**: 755
Tian, C., Wang, Y., Yu, H., He, J., Wang, J., Shi, B., Du, Q., Provart, N.J., Meyerowitz, E.M.,
and Jiao, Y (2019). A gene expression map of shoot domains reveals regulatory mechanisms. Nat.
Commun **10**: 1–12.
Tsukaya H (2005) Leaf shape: Genetic controls and environmental factors. International Journal of
Developmental Biology **49**: 547–555
Tsukaya H (2019) Re-examination of the role of endoreduplication on cell-size control in leaves. J
Plant Res **132**: 571–580
Vercruysse J, Baekelandt A, Gonzalez N, Inzé D (2021) Molecular networks regulating cell
division during Arabidopsis leaf growth. J Exp Bot **71**: 2365–2378

Vercruyssen L, Verkest A, Gonzalez N, Heyndrickx KS, Eeckhout D, Han SK, et al (2014)
ANGUSTIFOLIA3 binds to SWI/SNF chromatin remodeling complexes to regulate transcription
during Arabidopsis leaf development. Plant Cell 26: 210–29
Vroemen CW, Mordhorst AP, Albrecht C, Kwaaitaal MACJ, De Vries SC (2003) The CUP-
SHAPED COTYLEDON3 gene is required for boundary and shoot meristem formation in
Arabidopsis. Plant Cell **15**: 1563–1577
Wang Q, Hasson A, Rossmann S, Theres K (2016) Divide et impera: Boundaries shape the plant
body and initiate new meristems. New Phytologist **209**: 485–498
Wu G, Lin W-C, Huang T, Scott Poethig R, Springer PS, Kerstetter RA (2008) KANADI1
regulates adaxial-abaxial polarity in Arabidopsis by directly repressing the transcription of
ASYMMETRIC LEAVES2. Proc Natl Acad Sci U S A **105**:16392-7

995
996 **Figures**
997
998

999
1000

Figure 1: ***MIR319C*** **expression in leaf primordia with altered** ***CUC2*** **levels**. **(A)** Distribution of TF
families (in %) recovered from the yeast one hybrid screen (Y1H) with 378 TFs and ~2.736 kb
MIR319C URR. **(B)** Schematic representation of truncated *MIR319C* promoter fragments used for Y1H
assay with NAC TFs. Grey box corresponds to *HIS3* CDS. The three horizontal lines below the URR
scheme represent the truncated promoter regions, i.e., p^{distal} (-2736-1866), p^{middle} (-1866-1106) and
$p^{proximal}$ (-1106), respectively. **(C)** Images of yeast growth harboring three different truncated *MIR319C*
promoter regions upstream to the *HIS3* reporter gene and corresponding NAC TF preys as indicated,
grown in the presence of inhibitory concentration of 3-AT. NAC027 served as negative control.
(D) Bright field images of 10-day shoot apices (top panel) or 6[th] leaf primordia (bottom panel) of the
indicated genotypes expressing *pMIR319C::GUS* transgene. Numbers below the leaf primordia images
indicate leaf lengths and GUS domain/leaf length ratios are indicated within the parenthesis. Scale
bar,100 μm. **(E)** Violin plots displaying the quantification of *pMIR319C::GUS* domain/leaf length (Y-

axis) in the genotypes indicated on the X-axis. Differences among samples are indicated by *p*-values on
top of the comparisons, One-way ANOVA, Dunnett's post hoc test. N=20-32. **(F)** Quantification of
GUS, pre-*MIR319C* and *CUC2* transcript in genotypes indicated on the X-axis. Error bars represent
SEM of four biological replicates. Multiple comparisons were performed using One-way ANOVA
followed by Dunnett's post hoc test. *p*-values are shown above the data bars.

1018
1019
1020

1021
1022

Figure 2: Role of *CUC2-MIR319C* module in leaf cell proliferation. (A) Representative images of
mature first pair of leaves from 25-day old plants of the indicated genotypes. Scale, 5 mm. **(B)** Areas
(Y-axis) of mature first pair of leaves of the indicated genotypes on the X-axis. N=15-44 leaves.
Significant differences among the samples are indicated by different alphabets. p<0.0001. One-way
ANOVA, followed by Dunn's multiple comparison test are performed to determine the significant
differences among samples. **(C)** Representative images for pavement cell areas of mature first pair of
leaves from 25-day old plants of the indicated genotypes. Pavement cells and stomata are outlined in
black. Scale bar, 200 μm. **(D)** Distribution of pavement cell areas of mature first pair of leaves from 25-
day old plants of the indicated genotypes (X-axis). N= 115-333 cells from 3-4 leaves. Significant
differences among the samples are indicated by different alphabets. p<0.01. One-way ANOVA,
followed by Dunn's multiple comparison test are performed to determine the significant differences
among samples. **(E)** Cell numbers (Y-axis) in mature first pair of leaves of the indicated genotypes (X-
axis). Cell number was determined as the ratio of leaf area by the average cell area for 3-4 leaves per
genotype. Significant differences among the samples are indicated by different alphabets. p<0.01. One-
way ANOVA, followed by Tukey's multiple comparison test are performed to determine the significant
differences among samples.

1039

1040
1041

Figure 3: Cell proliferation pattern in leaf primordia with altered *CUC2* and *MIR319C* activities.
(A) Heat maps of cell areas in the first pair leaf primordia at 1, 3, and 5 days after initiation (DAI) of the indicated genotypes. Scale, 100 μm. **(B)** Distribution of pavement cell areas of leaf primordia at 3 DAI of the indicated genotypes (X-axis). N= 380-1079 cells from 2-3 leaves. Significant differences among the samples are indicated by different alphabets. p<0.05. One-way ANOVA, followed by Dunn's multiple comparison test are performed to determine the significant differences among samples. **(C)** Distribution of pavement cell areas of leaf primordia at 5 DAI of the indicated genotypes (X-axis). N= 2435-3280 cells from 2-3 leaves. Significant differences among the samples are indicated by different alphabets. p<0.01. One-way ANOVA, followed by Dunn's multiple comparison test are performed to determine the significant differences among samples.

Figure 4: Genetic interaction between *CUC2* and *MIR319* genes in regulating pavement cell number during leaf growth. (A) Representative images of 9-day old seedlings for each of the three categories of phenotypes, i.e., wild-type, narrow leaves, and seedling arrest/cotyledon fusion observed in the *cuc2;miR319a;miR319b* mutant combinations. Numbers above the images denote the proportion (in %) of seedlings displaying the indicated phenotype. **(B)** Representative images of mature first pair of leaves from 25-day old plants of the indicated genotypes. Scale, 5 mm. **(C)** Areas (Y-axis) of mature first pair of leaves of the indicated genotypes (X-axis). N=6-31 leaves. Significant differences among the samples are indicated by different alphabets. p<0.01. One-way ANOVA, followed by Dunnett's T3 multiple comparison test are performed to determine the significant differences among samples. **(D)** Cell numbers (Y-axis) in mature first pair of leaves of the indicated genotypes (X-axis). Cell number was determined as the ratio of leaf area by the average cell area for 3-4 leaves per genotype. Significant differences among the samples are indicated by different alphabets. p<0.05. One-way ANOVA, followed by Tukey's multiple comparison test are performed to determine the significant differences among samples.

Figure 5: Genetic interaction between *MIR319* genes in regulating leaf size. (A) Representative images of mature first pair of leaves from 25-day old plants of the indicated genotypes. Scale, 5mm. **(B)** Areas (Y-axis) of mature first pair of leaves of the indicated genotypes (X-axis). N=4-10 leaves. Significant differences among the samples are indicated by different alphabets. p<0.0001. One-way ANOVA, followed by Dunnett's T3 multiple comparison test are performed to determine the significant differences among samples. **(C)** Schematic representation of the *MIR319C* promoter *GUS* construct with mutated bases on the CUC-binding site at the -2617, -2618, and -2620[th] nucleotide positions indicated in red. The *GUS* region is indicated by grey box. **(D-E)** Bright field images of shoot apices and 250, and 675 µm long leaf primordia on 6[th], and 4[th] nodes respectively, expressing *GUS*-based reporter transgene under **(D)** *MIR319C* promoter or **(E)** *MIR319C* promoter carrying mutations on the CUC-binding site as indicated in **(C)**, respectively. **(F)** Violin plot displaying the quantification of *pMIR319C::GUS* or *mCUCBSpMIR319C::GUS* domain/leaf length in the 50-750 µm long leaf primordia on 4[th]-8[th] nodes. N=25 leaf primordia for *pMIR319C::GUS#2* and 31 leaf primordia for *mCUCBSpMIR319C::GUS*. Differences among samples are indicated by *p*-values on top of the comparisons, Unpaired *t*-test with Welch's correction for differences in variance. **(G)** Quantification of pre-*MIR319C*, *CUC1,* and *CUC3* transcript in 10-day old *pRPS5A::CUC2-GR-HA* seedlings treated for 2 hours with ethanol (Mock) or 12 µM dexamethasone (DEX) or 10 µM cycloheximide (CHX) or 10 µM cycloheximide and 12 µM dexamethasone (CHX+DEX) as indicated in the plot. Error bars represent SEM of three biological replicates. Pairwise comparisons were performed using unpaired *t*-test. *p*-values are shown above the data bars.

Figure 6: Effect of altered *CUC2* and *JAW-TCP* levels on leaf size. (A) Representative images of
mature first leaves from 25-day old plants of the indicated genotypes and chemical treatments. Scale, 5
mm. **(B)** Areas (Y-axis) of mature first pair of leaves of the indicated genotypes and treatments (X-
axis). N=18-35 leaves. Significant differences among samples are indicated by different lower-case
alphabets. p<0.01, two-way ANOVA, Tukey's post hoc test was performed. **(C)** Representative images
of first mature leaves from 25-day old plants of the indicated genotypes. Scale, 5 mm. **(D)** Areas (Y-
axis) of mature first pair of leaves of the indicated genotypes (X-axis). N=12-25 leaves. Significant
differences among the samples are indicated by different alphabets. p<0.05. One-way ANOVA,
followed by Dunnett's T3 multiple comparison test are performed to determine the significant
differences among samples.

Figure 7: Regulation of *MIR319C* expression by *CUC2* and *JAW-TCP* genes in leaf primordia.
(A) Bright field images of shoot apices (top panel) and 6th leaf primordia (bottom panel) from 10-day old Col-0 and *jaw-D* seedlings expressing *pCUC2::GUS* transgene. Scale bar, 50 μm. **(B)** Bright-field images of shoot apices (top panel) and 6th leaf primordia (bottom panel) dissected from 10-day old seedlings of indicated genotypes expressing the *pMIR319C::GUS* transgene. Scale bar, 50 μm. Numbers below the images represent leaf lengths in μm and those inside parenthesis denote the length of GUS domain relative to the leaf length. **(C)** Length of *pMIR319C::GUS* domain relative to total leaf length in 50-350 μm leaf primordia are plotted for the indicated genotypes. Differences among samples are indicated by different lower-case alphabets. $p < 0.05$. N = 33-114. Two-way ANOVA, followed by Tukey's post hoc test are performed to determine the significant differences among samples. **(D)** Bright-field images of shoot apices (top panel) and 6th leaf primordia (bottom panel) dissected from 10-day old seedlings of indicated genotypes expressing the *pMIR319C::GUS* transgene. Scale bar, 50 μm. Numbers below the images represent leaf lengths in μm and those inside parenthesis denote the length of GUS domain relative to the leaf length. **(E)** Length of *pMIR319C::GUS* domain relative to total leaf length in 50-350 μm leaf primordia are plotted for the indicated genotypes. N = 18-43. Significant differences among the samples are indicated by different alphabets. $p < 0.05$. One-way ANOVA, followed by Dunnett's post hoc test are performed to determine the significant differences among samples.