Title:

MIR319C activation by CUP-SHAPED COTYLEDON2 delays cell proliferation-to-

differentiation transition in Arabidopsis leaf primordia

Short title:

CUC2–MIR319C module controls leaf cell growth

Authors and affiliations:

Naveen Shankar[1φ], Abhishek Gupta[1#], Vishwadeep Mane[1#], Somsree Roy[1], Ahana

Jayanarayan Perumadaraya[1ε], Avani V[1], Olivier Hamant[2] and Utpal Nath[1*]

1 Department of Microbiology and Cell Biology, Indian Institute of Science, Bengaluru,

India.

2 Laboratoire de Reproduction et Développement des Plantes, Université de Lyon, ENS de

Lyon, UCBL, INRA, CNRS, 46 Allée d'Italie, 69364 Lyon Cedex 07, France.

# These authors contributed equally

φ Current address: Department of Plant Physiology, Umeå Plant Science Centre, Umeå

University, 901 87 Umeå, Sweden.

ε Current address: Institut de biologie moléculaire des plantes, CNRS, Université de

Strasbourg, Strasbourg, France, 67000.

* Corresponding author. Email: utpalnath@iisc.ac.in

Abstract

The microRNA miR319 regulates leaf size in diverse plant species by reducing the level of the target transcripts that encode JAW-TCPs, the transcription factors (TF) that restrict leaf size by committing the proliferating pavement cells to differentiation. *MIR319C*, one of the three miR319-producing genes in Arabidopsis, is expressed throughout the incipient leaf primordia, and its expression domain gets restricted to the base at later stages, partly due to its transcriptional repression by JAW-TCPs. However, the factors that activate and maintain *MIR319C* expression in leaf primordia are yet unknown. Here, we identify the CUP-SHAPED COTYLEDON2 (CUC2) TF as a direct activator of *MIR319C* transcription. Using a yeast one-hybrid (Y1H) screen, we identified several NAC domain TFs as potential regulators of *MIR319C*. Subsequent ex vivo binding and transactivation assays suggested that CUC2 binds to a distal promoter region of the *MIR319C* locus and activates its transcription. Mutants with compromised *CUC2* and *MIR319C* activities resulted in smaller leaves with fewer cells. Detailed morphometric analysis of higher order *CUC2* and *MIR319* loss-of-function mutants highlighted the crucial role of the *CUC2-MIR319* module in maintaining the duration of cell proliferation in leaf primordia. Additionally, the phenotype of mutants with altered *CUC2* and *MIR319/JAW-TCP* activities demonstrated that *CUC2* enhances leaf size through the *MIR319C-JAW-TCP* pathway. Overall, our findings uncovered a novel role for CUC2 in sustaining cell division by activating *MIR319C* transcription in the leaf primordia.

Introduction

Leaf development can be empirically divided into three major stages namely initiation, primary morphogenesis, and secondary morphogenesis (Bar and Ori 2014). Initiation of a leaf primordium commences with the specification of a group of founder cells at the periphery of the shoot apical meristem (SAM), a dome-shaped structure at the shoot apex that harbours a constant pool of stem cells throughout plant life (Barton 2010). Concurrently, a low cell proliferation region, termed as the 'boundary zone', gets established at the interface that separates the primordium from the SAM (Hepworth and Lenhard 2014; Wang et al. 2016). The basic form of the leaf consisting of the lamina and the petiole is established in eudicots during primary morphogenesis, wherein leaf growth is primarily sustained by cell proliferation. The end of primary morphogenesis is marked by the switch from cell proliferation to differentiation and the formation of specific cell types such as trichomes and

stomata (Bar and Ori 2014). During secondary morphogenesis, differentiation of all the cell

types is completed, and the leaf grows primarily by cell expansion (Bar and Ori 2014; Du et

al. 2018; Maugarny-Calès and Laufs 2018).

The shape and size of the mature leaf are regulated at each of the three developmental stages

by modulating the number of founder cells during initiation, and by regulating the rate,

location and duration of cell proliferation and expansion (Gonzalez et al. 2012; Hepworth and

Lenhard 2014; Kalve et al. 2014). The timing of cell proliferation arrest, which directly

influences the number of cells in mature leaves, plays a critical role in determining the final

leaf size across many plant species (Gázquez and Beemster 2017), suggesting that the

transition from cell division to expansion is a key cellular mechanism shaped by organ size

evolution. In Arabidopsis leaves, arrest of cell proliferation occurs basipetally; cells towards

the tip stop dividing and commit to differentiation early (Donnelly et al. 1999; Andriankaja et

al. 2012; Fox et al. 2018), while the cells towards the base continue to proliferate. Despite the

critical role of the timing in the transition from cell proliferation to differentiation, its

molecular basis remains less understood. Among the factors essential for this process, two

microRNA-transcription factor modules control the transition from cell proliferation to cell

expansion: miR396 and *GROWTH REGULATING FACTORs* (*GRFs*), and miR319 and five

CINCINNATA-like *TEOSINTE BRANCHED 1*, *CYCLOIDEA*, *PROLIFERATING CELL*

FACTOR1/2 (*CIN-TCPs*) (Rodriguez et al. 2014; Sarvepalli and Nath 2018). miR396

downregulates seven *GRF* transcripts in the distal region of the leaf, thereby restricting GRF

activities towards the base (Rodriguez et al. 2010). GRFs interact with GRF-INTERACTING

FACTOR 1/ANGUSTIFOLIA 3 (GIF1/AN3) transcriptional coactivators and upregulate

CYCLIN B (*CYCB*) expression to delay the proliferation-to-differentiation transition

(Schneider et al. 2023). Eight *CIN-TCP* genes promote leaf maturation by triggering the

transition of cell proliferation-to-differentiation in diverse eudicots including Arabidopsis

(Sarvepalli and Nath 2018). The spatiotemporal pattern of *CIN-TCP* transcripts is regulated

by a combined action of their promoter activity and miR319 activity (Palatnik et al. 2003;

Efroni et al. 2008), which prevents the accumulation of transcripts of five *CIN-TCPs* (*TCP2,*

3, 4, 10, 24), also known as *JAGGED AND WAVY TCPs* (*JAW-TCPs*), in the proximal region

of the young leaf primordia (Palatnik et al. 2003; Nag et al. 2009). Expression of miR319-

resistant form of *JAW-TCPs* under endogenous regulation leads to smaller leaves with fewer

cells due to precocious proliferation arrest, while ectopic expression of miR319 results in

larger leaves with wavy margin due to a delay in the transition from proliferation to

differentiation (Palatnik et al. 2003; Efroni et al. 2008; Challa et al. 2019). CIN-TCPs commit

the leaf cells towards differentiation by modulating multiple pathways, such as auxin and

cytokinin response, meristematic activity, and cell cycle progression (Sarvepalli and Nath

2018).

The *CUP SHAPED COTYLEDON* (*CUC*) group of genes, which encode the NAM/ ATAF/

CUC (NAC) group of TFs, regulate boundary development in diverse angiosperms (Wang et

al. 2016). The Arabidopsis *CUC* members *CUC1*, *CUC2*, and *CUC3* are consistently

expressed in the boundary zones, and their absence leads to the fusion of adjacent organs

such as cotyledons, sepals, stamens, and ovules (Aida et al. 1997; Ishida et al. 2000;

Vroemen et al. 2003). The *CUC* genes are also required for the maintenance of shoot

meristem via the activation of the *SHOOT MERISTEMLESS* (*STM*) gene and its homologues

(Aida et al. 1999; Hibara et al. 2006; Scofield et al. 2018). *CUC1* transcript is not detected in

leaves, while *CUC2* and *CUC3* are broadly expressed along the SAM-leaf boundary and leaf

margins (Nikovics et al. 2006; Hasson et al. 2011). However, *CUC3* shows weaker

expression, and its activity is dependent on *CUC2* (Hasson et al. 2011). In leaf primordia,

CUC2 and CUC3 proteins accumulate at distinct foci along the margins where they are

necessary for the initiation and growth of serrations through a feedback mechanism involving

auxin and the auxin efflux carrier protein PINFORMED1 (PIN1) (Nikovics et al. 2006;

Bilsborough et al. 2011; Hasson et al. 2011; Maugarny-Calès et al. 2019). Interestingly, some

of the gain-of-function phenotypes of *JAW-TCPs* such as impaired SAM function, cotyledon

fusion, and smooth leaf margin are dependent on the downregulation of *CUC* activity

(Koyama et al. 2007; Koyama et al. 2010; Koyama et al. 2017; Challa et al. 2019). While

TCP3 inhibits *CUC1* and *CUC2* by activating the transcription of the miR164-encoding gene

MIR164A to promote shoot morphogenesis (Koyama et al. 2007; Koyama et al. 2010), TCP4

interacts with CUC and inhibits their homo-dimerization and transactivation abilities to

modulate serration formation and outgrowth (Rubio-Somoza et al. 2014).

Expression analysis of the miR319-encoding genes *MIR319A*, *MIR319B*, and *MIR319C* in

Arabidopsis suggested their distinct developmental functions due to largely non-overlapping

expression pattern (Nag et al. 2009). Whereas *MIR319A* and *MIR319B* promoter activity is

not detected in leaves, *MIR319C* is strongly expressed in the leaf primordia (Nag et al. 2009;

Shankar et al. 2023). The single loss-of-function mutants for miR319a and miR319b show a

moderate decrease in leaf serration, and the *miR319a;miR319b* double mutant leaves lack leaf

serration, suggesting a role for *MIR319A* and *MIR319B* genes in leaf development (Koyama

et al. 2017). However, the *miR319a;miR319b* leaves are as big as wild-type (Koyama et al.

2017), pointing towards a key role of miR319c in downregulating *JAW-TCP* activity during

the early leaf growth. We had earlier demonstrated that JAW-TCPs transcriptionally repress

the miR319-encoding gene *MIR319C* in the distal region to prevent miR319 accumulation,

thereby establishing spatially separated proximal miR319 domain and distal JAW-TCP

domain in incipient leaf primordia (Shankar et al. 2023). However, it is still unknown how

the *MIR319* genes are transcriptionally activated in early leaf primordia. Here, we

demonstrate that the CUC2 protein transcriptionally activates the *MIR319C* gene and,

together with the miR319 product of *MIR319A* and *MIR319B*, promotes cell proliferation and

leaf size by downregulating *JAW-TCP* level.

Results

Transcriptional activators of *MIR319C* identified in a yeast one-hybrid screen

The *MIR319C* promoter activity is primarily detected at the shoot apex, more specifically at

the meristem-primordia boundary and in young leaf primordia and is excluded from the

central domain of apical meristem (Shankar et al. 2023) (**Fig. S1A, S1B**), suggesting that

proteins encoded by the transcriptome subset within this domain include factors that activate

MIR319C transcription. To identify these regulators, we screened a subset of yeast one-

hybrid library containing 378 Arabidopsis transcription factors, enriched in the shoot apex,

for their ability to bind to the upstream regulatory region (URR) of *MIR319C* (See methods,

Fig. S1C, S1D, Suppl Data Set. 1). Fifty-seven TFs were identified that promoted the

growth of yeast cells when a 2736 bp URR was used as bait to drive the expression of the

HISTIDINE3 (*HIS3*) selection gene (**Suppl Data Set. 2**). Gene ontology analysis revealed

that TFs belonging to NAC, TCP, ETHYLENE RESPONSE FACTOR (ERF), BASIC

LEUCINE ZIPPER (bZIP), and BASIC HELIX LOOP HELIX (bHLH) families were over-

represented among the positive clones identified in the screen (**Fig. 1A**).

Five out of the 6 NAC prey proteins recovered from the Y1H screen (**Fig. S1E**) show

comparable DNA-binding preferences to the target DNA TT(G/A)CGT (Lindemose et al.

2014), a sequence element present at the 2616-2621 nucleotide position upstream to the start

site of pre-*MIR319C* sequence (**Fig. 1B**). The binding preference of these 5 NACs is also

shared by other members of the NAM, OsNAC7, NAC2, ANAC011, and AtNAC3 subgroups

of the AtNAC TF family (Ooka et al. 2003; Lindemose et al. 2014). To identify additional

NAC members capable of binding to the *MIR319C* URR in a targeted, small-scale Y1H

screen, we tested the binding ability of 20 representative members from these 5 AtNAC

subgroups (**Fig. S2A, S2B**) to three non-overlapping *MIR319C* URR fragments named as

p^{distal} (-2736 to -1866 bp), p^{middle} (-1866 to -1106 bp), and $p^{proximal}$ (-1106 to -1 bp) (**Fig. 1B**).

Five additional NAC members including CUC1, CUC2 & CUC3 supported yeast growth

when p^{distal}, which includes the putative NAC-binding DNA element, was used as bait (**Fig.**

1C).

To test which URR region is important for *MIR319C* promoter activity, we generated

transgenic lines expressing the *GUS* reporter construct under truncated (-533 bp or -1866 bp

long) or full-length (-2736 bp long) promoter fragments, and studied *pMIR319C::GUS*

expression pattern in young leaves. The *MIR319C* activity was significantly weaker in the

line carrying -1866 bp promoter region, and barely detectable in the line carrying -533 bp

promoter region, compared to the full length promoter (**Fig. S3B**). This suggests that the

distal (-2736 to -1866 bp) and the middle (-1866 to -1106 bp) promoter regions contain cis

elements required for *MIR319C* expression.

CUC2 activates *MIR319C* in leaf primordia

CUC2, the primary *CUC* member in leaves (Nikovics et al. 2006; Hasson et al. 2011), is

active throughout the shoot apex, nascent leaf primordia, and at the base of young leaves,

overlapping the *MIR319C* expression domain (**Fig. S1A, B**). As the CUC2 protein binds to

MIR319C URR in yeast (**Fig. 1**), we explored whether CUC2 regulates *MIR319C* promoter

activity in developing leaves. Comparative GUS reporter activity analysis in leaves

expressing the *pMIR319C::GUS^#1* construct (Nag et al. 2009; Shankar et al. 2023) revealed

that the *MIR319C* promoter activity is altered in the *CUC2* mutants compared to wild type

(**Fig. 1D, 1E**). The domain of *pMIR319C::GUS* expression relative to the leaf length was

proximally restricted (median ~0.35) in the *cuc2-1* loss-of-function leaves (Hasson et al.

2011), and distally extended (median ~0.75) in the *cuc2-1D* gain-of-function leaves (Larue et

al., 2009), compared to wild type (median ~0.6). Corresponding to the change in the GUS

domain in the mutant leaves, there was a ~2-fold reduction in the level of *GUS* transcript in

the *pMIR319C::GUS^#1;cuc2-1* seedlings (p value, 0.005), and ~2.5-fold increase in the

pMIR319C::GUS#1;cuc2-1D seedlings (p value, 0.06), as analyzed by RT-qPCR (**Fig. 1F**),

suggesting that CUC2 activates *MIR319C* promoter. This was further supported by the

observation that the pre-*MIR319C* transcript level was ~1.5-fold lower in

pMIR319C::GUS#1;cuc2-1 (p value, 0.004) and >2-fold higher in *pMIR319C::GUS#1;cuc2-*

1D seedlings (p value, 0.279; **Fig. 1F**). Together, these results suggest that CUC2 is required

and sufficient to activate *MIR319C* expression in leaf primordia.

CUC2 binds to *MIR319C* locus and activates its transcription

To determine the functional significance of the predicted CUC-binding cis element within the

MIR319C URR, we generated a GUS-based reporter line of *MIR319C* promoter carrying

mutations in the CUC-binding site (**Fig. 2A**) and determined the GUS activity in the resultant

mCUCBS-pMIR319C::GUS leaves. The relative GUS domain was much reduced in the leaf

primordia expressing *mCUCBS-pMIR319C::GUS* transgene compared to the wild-type

pMIR319C::GUS#2 transgene (Shankar et al. 2023) (**Fig. 2B-D**), suggesting that the CUC-

binding site is required for the *MIR319C* promoter activity.

To further test whether CUC2 directly activates *MIR319C* transcription, we induced CUC2

activity in 10-day-old *pRPS5A::CUC2-GR-HA* seedlings (Tian et al. 2014) by treating them

with dexamethasone (DEX) for two hours in the presence of the protein synthesis inhibitor

cycloheximide (CHX) and measured pre-*MIR319C* level by RT-qPCR. CUC2 induction

increased the transcript of *CUC3* (**Fig. 2E**), one of its known direct targets (Maugarny-Calès

et al. 2019), but not of *CUC1*, in the presence of CHX. Similar to positive control, CUC2

induction also elevated the pre-*MIR319C* transcript level by ~2-fold in the presence of CHX,

suggesting that no new protein synthesis is required for activating *MIR319C* transcription by

CUC2. These results indicate that CUC2 binds to the *MIR319C* locus and activates its

transcription in leaf primordia.

CUC2 **promoters leaf cell proliferation and leaf size in miR319-dependent manner**

The mature miR319 produced by the *MIR319A*, *MIR319B* & *MIR319C* genes promotes cell

proliferation and leaf size by degrading the transcripts of five *JAW-TCP*s (Palatnik et al.

2003; Efroni et al. 2008; Koyama et al. 2010; Challa et al. 2019). If CUC2 promotes miR319

accumulation, inactivation of either *CUC2* or *MIR319C* is expected to yield smaller leaves

with fewer cells. To test this, we analysed mature leaves of the *CUC2* loss-of-function allele
cuc2-3 (Nikovics et al. 2006), and a *MIR319C* loss-of-function allele available in the
Arabidopsis stock centre (https://arabidopsis.info/; established as a line and named here as
miR319c-1) (**Fig. S3A**). While *miR319c-1* mutant did not show any noticeable reduction in
mature leaf area (**Fig. 2F, 2G**) or epidermal cell area (**Fig. S4A, S4B**), mature leaf size of
cuc2-3 was reduced by ~20% compared to Col-0 (**Fig. 2F, 2G**). The weak phenotypes of
single mutants are perhaps because of the hypomorphic nature of the mutant alleles or due to
the presence of their redundant partners (Nag et al. 2009; Hasson et al. 2011). To test the
former possibility for *MIR319C*, we generated novel knockout mutants of *MIR319C* using
CRISPR zCas9i editing approach (See methods; Grützner et al. 2021, Stuttmann et al. 2021),
and recovered the *miR319c-2* allele with complete, and the *miR319c-3* allele with partial
deletion of the region that produces mature miR319c (**Fig. S5**). Both these alleles produced
mature leaves with nearly 40% smaller area compared to wild type (**Fig. 2F, 2G**) without a
noticeable change in average cell area (**Fig. S4A, S4B**), resulting in reduced cell number in
the mutants (**Fig. 2H**). These results show that miR319c activity is required for leaf cell
proliferation.
253
To explore the genetic relationship between *CUC2* and its target *MIR319C* in regulating cell
proliferation, we generated the double mutant line *cuc2-3;miR319c-1* by crossing weak
alleles of these two genes and analysed its leaf size. Mature *cuc2-3;miR319c-1* leaves were
nearly half the size of their respective parents (**Fig. 2F, 2G**), though the average area of their
pavement cells remained unaltered (**Fig. S4A, S4B**). The *cuc2-3;miR319c-1* leaf blade was
made up of ~2000 pavement cells whereas the Col-0 leaves had ~4500 cells (**Fig. 2H**),
demonstrating that smaller area of the double mutant leaves is due to reduced cell
proliferation. The corresponding cell numbers for the *cuc2-3* and *miR319c-1* leaves also
reduced compared to the wildtype (~3100 for *cuc2-3* and ~3500 for *miR319c-1*), but to a
much less extent (**Fig. 2H**). In combination with the biochemical and expression analysis
described above, we interpreted results of these genetic analysis as CUC2 promoting cell
proliferation in leaf primordia in a miR319c-dependent manner.
266
During leaf morphogenesis, cell proliferation commences throughout the incipient primordia,
followed by concurrent proliferation and expansion in distinct zones, with all cells finally
ceasing to proliferate and undergoing secondary expansion (Donnelly et al. 1999;
Andriankaja et al. 2012; Fox et al. 2018). To determine the stage at which cell proliferation in

the   *cuc2-3;miR319c-1* leaves diverges from wild type, we performed confocal

imaging of propidium iodide-stained leaf primordia at 1, 3, and 5 days after initiation (DAI)

and analysed their cellular parameters using MorphographX software (Barbier de Reuille et

al. 2015; Strauss et al. 2022). The basipetal growth pattern of *cuc2-3;miR319c-1* leaves were

comparable with that of Col-0 at all the early growth stages studied (**Fig. 3A**). Further, the

total number and average size of the pavement cells also remained unaltered in the mutant

leaf primordia (**Fig. S6A-D**), suggesting that loss of *CUC2* or *MIR319C* does not affect cell

proliferation or its spatial pattern at early growth stage.

To examine whether the *CUC2-MIR319C* module promotes cell proliferation at later stages,

we analysed leaf and cell growth kinematics by measuring leaf area, pavement cell area, and

total cell number of Col-0 and *cuc2-3;miR319c-1* leaves at 10, 15, 20, and 25 DAI. The

analysis revealed that average leaf area of *cuc2-3;miR319c-1* was reduced to nearly half at

later stages of growth, i.e., 15 DAI onwards (**Fig. 3B, 3C**), with concomitant and

corresponding decrease in total cell number at all stages (**Fig. 3D**). The average area of the

cuc2-3;miR319c-1 epidermal cells, though increased with time, remained comparable to that

of Col-0 (**Fig. 3E**), suggesting that the reduction of leaf size in the mutant was primarily due

to reduced cell proliferation. Though average cell size remained same, the mutant leaves had

higher proportion of larger cells at all growth stages measured (**Fig. S7**), pointing at

precocious differentiation. Together, these results demonstrate that the *CUC2-MIR319C*

module promote cell proliferation and delays differentiation in leaf primordia.

MIR319A and *MIR319B*, the two homologues of *MIR319C*, are required for leaf serration and

loss of function of both the genes produces smooth margin (Koyama et al. 2017). To examine

whether *MIR319A* and *MIR319B* also regulate cell proliferation in leaf primordia along with

MIR319C, we generated higher order mutants of these three genes by crossing

miR319a;miR319b double mutant with *miR319c-1* and determined their leaf area and

pavement cell parameters. Mature *miR319a;miR319b;miR319c-1/+* and

miR319a;miR319b/+;miR319c-1 mutant plants produced leaves that are less than one-third

the size of the leaves of their respective parents and of Col-0, while the

miR319a;miR319b;miR319c-1 triple homozygous leaves were ~4.5 times smaller than Col-0

leaves (**Fig. 4A, 4C**). Reduction in cell number alone accounted for smaller leaf area (**Fig.**

4D), with the average size of the pavement cells remaining nearly unaltered (**Fig. S8A, S8B**).

Together, these results suggest a functional redundancy among three *MIR319* genes in
promoting leaf cell proliferation.

To further investigate a possible genetic interaction between *MIR319A/ MIR319B* and *CUC2*
in regulating leaf cell proliferation, we combined *miR319a;miR319b* genotype with *cuc2-3*
and *cuc2-1* and studied the leaf phenotype in higher-order mutants. Whereas most (~96%) F2
individuals of the *miR319a;miR319b* x *cuc2-3* genetic cross displayed parental phenotype, a
small fraction of them (32 out of 821) showed aberrant phenotypic features including smaller
leaves and fused cotyledons (**Fig. S9A**), the latter being characteristic to *cuc* loss of function
mutants (Aida et al. 1997; Hibara et al. 2006). Similar phenotypic aberrations of varying
proportions were also noticed among the F2 individuals of the *miR319a;miR319b* x *cuc2-1*
cross (**Fig. S9A**). More importantly, nearly one third individuals of the genetically
established   *cuc2-3;miR319a* & *cuc2-3;miR319b* homozygous populations, and of
cuc2-3;miR319a/+;miR319b/+ & *cuc2-3;miR319a/+;miR319b* populations also showed
smaller leaves (2-4 folds) and fused cotyledons (**Fig. 4E**, **Fig. S9B**), suggesting a genetic
interaction among these genes. Reduced leaf area in *cuc2-3;miR319a/+;miR319b/+* &
cuc2-3;miR319a/+;miR319b lines was accompanied with >2-fold decrease in the number of
pavement cells (**Fig. 4D**) without any increase in their size (**Figs. S8A, S8B**). Thus, the
CUC2-MIR319 pathway promotes cell proliferation in early leaf primordia.

***CUC-MIR319C-JAW-TCP* module regulates leaf size**

Since mature miR319 suppresses the function of five *JAW-TCP* genes by degrading their
transcripts, CUC2-mediated *MIR319* activation and promotion of cell proliferation is
expected to be dependent of JAW-TCP activity. We tested this by carrying out genetic
interaction studies between loss and gain-of-function lines of *CUC2* and *JAW-TCP*s. The
gain-of-function line of *CUC2*, *cuc2-1D*, forms larger leaves with increased cell number
(Larue et al. 2009), perhaps due to enhanced *MIR319C* level (**Fig. 1D-F**) and a consequential
decrease in *JAW-TCP* level. Activation of an miR319-resistant, dominant form of TCP4
within its endogenous expression domain in the *Col-0;GR*[#1] x Col-0 F1 plants reduces leaf
size by limiting cell proliferation (**Fig. 5A, 5C**, Challa et al. 2016). TCP4 activation in the
Col-0;GR[#1] x *cuc2-1D* F1 plants reduced the size of the bigger *cuc2-1D* leaves to the same
level as in the Col-0 background (**Fig. 5A, 5C**), thus suggesting that *JAW-TCP*s function
downstream to *CUC2*. Overexpression of miR319 under a CUC2-insentitive promoter in the

pBLS>>MIR319A leaves expectedly led to larger lamina (**Fig. 5B, 5D;** Efroni et al. 2008;

Challa et al. 2019). However, additionally mutating *CUC2* in this genetic background did not

further alter the leaf size (**Fig. 5B, 5D**). Taken together, these results support the conclusion

that *CUC2* promotes leaf size by reducing *JAW-TCP* level in a miR319-dependent manner.

***CUC2* and *JAW-TCP* activities together establish *MIR319C* expression domain in early**

leaf primordia

Members of the *JAW-TCP* clade establish early expression patterns of *MIR319C* by

suppressing its transcription in the distal incipient leaf primordia, and of the *CUC* genes by

restricting them to the proximal domain of the early shoot lateral organs (Koyama et al. 2007;

Challa & Rath et al. 2021; Shankar et al. 2023). To examine whether JAW-TCPs modulate

CUC2 transcription in establishing the endogenous *MIR319C* expression pattern during leaf

growth, we investigated *pCUC2::GUS* activity (Nikovics et al. 2006) in *jaw-D*, a knockdown

line of *JAW-TCP*s (Palatnik et al. 2003), and found that the *CUC2* expression domain

remained unaltered in the *jaw-D* leaf primordia (**Fig. 6A**). Further, activation of a dominant

TCP4 by dexamethasone treatment in the Col-0 x Col-0;*GR*[#1] F1 seedlings did not change the

pCUC2::GUS domain either (**Fig. S8C**), suggesting that JAW-TCPs do not affect *CUC2*

transcription during the growth of leaf primordia.

To further investigate how JAW-TCPs and CUC2 together establish *MIR319C* expression

pattern, we determined *MIR319C* promoter activity in leaf primordia where the level of both

CUC2 and *JAW-TCP*s were altered. As was reported earlier, activation of a dominant TCP4

protein in the Col-0;*GR*[#2] leaves, where an miR319-resistant TCP4 is induced by

dexamethasone treatment in the wildtype background (Shankar et al. 2023),

pMIR319C::GUS domain was reduced and restricted more proximally (**Fig. 6B, 6D**).

However, the extent of restriction was less when TCP4 was induced in the *cuc2-1D;GR*[#2] leaf

primordia, indicating that the endogenous *MIR319C* expression domain is established by

independent and opposing activities *CUC2* and *JAW-TCPs*. In support of this,

downregulation of the *JAW-TCP*s rescued the reduced *MIR319C* expression of the *cuc2-1*

leaves (**Fig. 6C, 6E**). However, downregulation of *JAW-TCP*s failed to further increase the

already extended *MIR319C* domain in   *cuc2-1D* (**Fig. 6C, 6E**), perhaps

suggesting that either *CUC2* activation or *JAW-TCP* inactivation results in the maximum

extent of *MIR319C* domain at the leaf growth stages studied here.

Discussion

Regulation of organ size is a fundamental question in developmental biology. In angiosperm
species such as Arabidopsis, leaves grow initially via cell proliferation, followed by a
combination of proliferation and expansion that are spatially separated along the length of the
primordia. Finally, all pavement cells attain their maximum and fairly uniform size
throughout the leaf lamina in a mature leaf. While the timing of the transition from cell
proliferation to differentiation has been shown to influence final cell number of leaf size, its
molecular basis remains somewhat unclear. Here, we report that the *CUC2-MIR319-JAW-*
TCP module plays a key role in promoting pavement cell proliferation by delaying its
transition to differentiation, thereby impacting leaf size. Our data is consistent with the model
(**Fig. 7A**) where CUC2 activates *MIR319C* transcription by binding to a cis element in the
MIR319C upstream regulatory region (**Figs. 1 and 2**), leading to an elevated level of
miR319c activity in the incipient leaf primordia. As the primordia grow, *CUC2* expression
becomes progressively restricted towards leaf-SAM boundary where it continues to activate
MIR319C expression. The miR319c activity at the leaf base prevents JAW-TCP
accumulation and sustains cell proliferation. JAW-TCPs first accumulate in the distal cells of
the growing primordia due to a yet unknown mechanism, and feedback to repress *MIR319C*
transcription within their expression domain (Shankar et al. 2023). The opposing activities of
CUC2 and JAW-TCPs *MIR319C* transcription independently establish the endogenous
miR319 domain in early leaf primordia (**Fig. 6**). Further experiments are needed to test
whether CUC2 directly binds to this element in planta or if additional factors are required for
CUC2 binding on the *MIR319C* locus.

Role of the *CUC2-MIR319C* module in spatiotemporal regulation of leaf cell
proliferation

In our study, loss of *CUC2* results in only a modest reduction in leaf size, primarily due to a
decrease in cell number (**Fig. 2**). This may in part be attributed to the partial downregulation
of miR319c, as evidenced by reduced promoter activity in the *cuc2* loss-of-function mutant
(**Fig. 1**). The strong reduction in cell number observed in the *cuc2-3;miR319c-1*, and in the
miR319c-2 or *miR319c-3* null mutants, support this hypothesis (**Fig. 2**). We speculate that the
cuc2-3;miR319c-1 mutant has lower levels of miR319c than the individual parental alleles

that are perhaps hypomorphic. Since *MIR319C* expression is restricted to the basal half of leaf primordia by 3 DAI (~250 μm long primordia; **Fig. S1**), and JAW-TCPs are active at the distal end by this stage (Shankar et al. 2023), we investigated if the cell proliferation pattern is altered in *cuc2-3;miR319c-1* during early leaf development. However, no significant changes in cell number, cell areas or cell size distribution were observed up to 5 DAI (~500 μm long primordia; **Fig. 3**). Difference in leaf area and cell number started appearing only by 10 DAI onwards (**Fig. 3**), a stage when *CUC2* and *MIR319C* expressions are restricted to the base of the petiole (**Fig. S1**). These observations suggest that the *CUC2-MIR319-JAW-TCP* module does not influence cell division rate or the establishment of the initial leaf growth pattern but rather affects the duration of pavement cell proliferation by delaying the transition.

JAW-TCPs are known to promote the proliferation-to-differentiation transition via auxin dependent and independent genetic pathways (Challa et al. 2019). The reduction in miR319 level in *cuc2-3;miR319c-1* leaves is likely to enhance JAW-TCP level, leading to premature differentiation of cells that would otherwise continue to proliferate in the wild-type primordia. Over time, this effect compounds, ultimately leading to a detectable reduction in cell number at later stages, leading to smaller leaves. This is further supported by our kinematic analysis, which revealed that *cuc2-3;miR319c-1* leaves consistently have fewer cells at 10 DAI and at subsequent stages of growth (**Fig. 3**), indicating an early transition from proliferation to differentiation. This interpretation corresponds with the increased proportion of larger pavement cells in the mutant leaves as early as at 10 DAI (**Fig. S7**). Additional evidence comes from the genetic interactions between the mutants of *CUC2* and *JAW-TCPs*, where ectopic JAW-TCP activity reduced the size of *CUC2*-overexpressing leaves, and miR319 overexpression rescued smaller-size phenotype of the *cuc2-1* leaves (**Fig. 5**). Overall, our findings support the model (**Fig. 7B**) where CUC2 activation of *MIR319C* maintains the proliferation potential of leaf cells from the base to the transition zone in growing leaves by downregulating JAW-TCP activity. In the absence of this regulation, the proliferation phase is shortened, resulting in fewer cells and smaller leaves in *cuc2-3;miR319c-1*.

Redundancy among *MIR319* genes in regulating leaf size

Individual role of the three *MIR319* genes in regulating leaf cell proliferation were not clear

until now due to functional redundancy among them and due to the availability of null

mutants. Defects in leaf area and cell number in *miR319c-2; miR319c-3* plants (**Fig. 2**)

revealed a key role for all the three *MIR319C* genes in promoting cell proliferation via

downregulation of JAW-TCP level. Reduced leaf area and cell number in the triple

miR319a;miR319b;miR319c-1 mutant but not in the *miR319a;miR319b* mutant or in

miR319c-1 mutant (**Fig. 4**) suggest that *MIR319A* and *MIR319B* also play a role in

maintaining cell proliferation. Moreover, *MIR319A/ MIR319B* displayed a strong genetic

interaction with *CUC2* in promoting cell number and leaf size (**Fig. 4**). This is likely due to

the additional reduction in the mature miR319 pool in   *cuc2-3;miR319a;miR319b*

leaves resulting from the combined loss of all three *MIR319* genes. It is also possible that

CUC2 activates *MIR319A/ MIR319B* genes in leaf primordia, in addition to *MIR319C*.

Interestingly, the *cuc2-3;miR319a; miR319b* mutants exhibited reduced cell area and, in

some cases, distinctly altered pavement cell morphology (**Fig. S8**), suggesting varying

degrees of contribution from *MIR319A/ MIR319B* genes to leaf growth. This variation may

result from spatiotemporal differences in *MIR319* expression during leaf development. In

future, detailed analysis of *MIR319A/ MIR319B* expression pattern (Nag et al. 2009)

combined with growth kinematic study of their mutant leaves , will help clarify these

possibilities.

Context-dependent role of *CUC2* in cell proliferation

CUC2 perhaps plays a context-dependent role in regulating cell proliferation and establishing

growth polarity during organ morphogenesis. During embryogenesis , *CUC2* suppresses cell

proliferation at the boundary between SAM and emerging cotyledon primordia to ensure

cotyledon separation (Aida et al. 1997; Ishida et al. 2000). Similarly, *CUC2* also inhibits cell

proliferation in at the sinus region of leaf primordia and thereby promotes serration

development (Nikovics et al. 2006; Maugarny-Calès et al. 2019; Hasson et al. 2011), a

boundary function parallel to its effect on cotyledon separation during embryogenesis. Here,

we demonstrate that *CUC2* rather promotes cell proliferation in the medial region of the

lamina base through the *miR319-JAW-TCP* pathway. This dual role of *CUC2* as both a

suppressor and a promoter of cell proliferation could be influenced by developmental context

and tissue-specific factors. Additionally, our results support the idea that *CUC2* functions as a

growth polarity organizer (Kuchen et al. 2012; Sarvepalli et al. 2019), where it drives

differential growth between the proximal and the distal regions of the leaf primordia via

MIR319C activation. This also raises an interesting possibility that tissue boundaries can

serve as growth organizers during organ morphogenesis in plants (Bhatia et al. 2023; Hu et

al. 2024). Future research could explore mechanisms that link boundary gene network and

growth regulatory modules to test this possibility.

Methods

In silico screening for SAM and leaf primordia-specific transcription factors

To identify SAM and leaf primordia-specific TFs, we took advantage of studies that

generated cellular-resolution maps of genes whose mRNA is actively translated in different

domains within the SAM and leaf primordia (Tian et al. 2014; 2019). We considered genes

that are active in the *CLV3*, *WUS*, *UFO* expression domains of the SAM, *LAS* expression

domain of the SAM-leaf boundary, and the *AS1* expression domain of the leaf primordia. We

overlapped the 4426 SAM and leaf primordia-specific genes with the list of 1956 genome-

wide Arabidopsis TFs (Pruneda-Paz et al. 2014) and obtained 343 TFs expressed explicitly in

the SAM and leaf primordia domains (**Suppl Data Set. 3**). We also scanned the 2.7 kb URR

of *MIR319C* for TF binding sites using the publicly available transcription factor binding site

(TFBS) prediction tool (PlantPAN 3.0 - *http://plantpan.itps.ncku.edu.tw*). We recovered

several potential DNA-binding motifs on the *MIR319C* promoter, including TCP, KNOX,

YABBY, NAC and HD-like TF-binding consensus. We listed 49 representatives from the top

5 TF families recovered from the TFBS prediction software (**Suppl Data Set. 4**). Next, we

combined this list with the previously obtained 343 SAM and leaf primordia-specific TFs to

get a final list of 378 TF candidates for the yeast one-hybrid screen (**Suppl Data Set. 1**).

Yeast transformation

Yeasts were transformed with the desired DNA constructs (plasmids) by the high-efficiency

LiAc-PEG method described by Walhout and Vidal 2001, with minor modifications. A single

colony of host strain was inoculated in 10 ml YPAD (Bacto yeast extract-1g, Tryptone-1g,

Dextrose-2g, and Adenine sulphate-10mg per 100ml) and incubated overnight at 30ºC with

constant shaking. The next day, the suitable culture volume was inoculated into 200 ml of

fresh YPAD solution to an initial OD of 0.2 and kept at 30ºC with constant shaking until an

OD of 0.6-0.8 was reached. Cells were pelleted down at 700 g for 10 min and washed twice

with 5 ml of sterile water. The cells were washed with 2.5 ml of LiAc-TE (0.1 M LiAc, Tris-

10 mM, EDTA- 1mM) solution. Cells were finally resuspended in 2 ml of LiAc-TE solution.

A mixture containing 280 μL of 50% PEG (Mol wt 3350, Sigma-Aldrich, USA), 36 μL of 1

M LiAc in TE, 6 μL of 10 mg/ml boiled carrier DNA (D-3159, Sigma-Aldrich, USA) was

prepared and added to 1.5 ml microfuge tubes containing 60 μL cells and 100 ng of DNA

construct. The cell suspension was vortexed briefly and incubated at 30ºC with constant

shaking for 30 min. Cells were transferred into a water bath maintained at 42ºC for 30 min.

Cells were pelleted, washed twice with sterile water, resuspended in 100 μL sterile water, and

plated onto appropriate media.

For the Y1H screen, transformations were carried out in 96 well-format described by

Walhout and Vidal, 2001. Yeast cells were prepared as described in the high-efficiency

protocol. 20 μL of cell suspension (containing LiAc-PEG-carrier DNA) were transferred to

each well of a sterile U-bottom 96-well plate (3795, Corning, USA). 100 ng of plasmid and

100 μL of TE/LiAc/PEG solution (50% PEG and 0.1 M LiAc in TE) were individually added

and resuspended carefully using a multichannel pipette. 96-well plate was incubated at 30ºC

with constant shaking for 30 min. Cells were transferred to a water bath maintained at 42ºC

for 30 min. Cells were pelleted at 700 g for 5 min. To each well, 120 μL of sterile water was

added, and 100 μL was removed directly to get rid of PEG in the supernatant. Finally, cells

were resuspended in the remaining 20 μL water and spotted on appropriate media.

Yeast bait, prey, and diploid generation

The full length (2.7 kb) and the non-overlapping promoter fragments, i.e., -2694-1866 (p^{distal}),

-1866-1107 (p^{middle}), and -1107+35 ($p^{proximal}$) of *MIR319C* were amplified from *pMIR319C-*

P4P1R entry clone using Taq DNA polymerase and cloned into pENTR/D-TOPO vector

(K240020, Invitrogen, USA) according to the manufacturer's instructions. The entry clones

were confirmed by Sanger sequencing before cloning upstream to the *minimalPro::HIS3*

cassette of *pMW#2* (Gift from Ram Yadav, IISER Mohali, INDIA) by LR reaction. Y1H

baits were generated according to the protocol described by Fuxman Bass et al. 2016. Briefly,

the *pMIR319C* containing *pMW#2* reporter constructs were linearized with *Xho*I (R0146,

New England Biolabs, USA) and integrated into the mutant *HIS3* locus of the *YM4271* host

strain (Gift from Ram Yadav, IISER Mohali, INDIA) using the high-efficiency LiAc-PEG

transformation method described by Walhout and Vidal, 2001. Following transformation,

yeast cells were plated on the synthetic complete media lacking histidine (Sc-His) media

(Walhout and Vidal, 2001) and incubated at 30ºC till colonies appeared. Post growth,

genomic DNA was extracted individually from 3-4 colonies, and the integration of the

MIR319C promoter fragment was confirmed by PCR. All primer sequences used for cloning

are mentioned in **Suppl Data Set. 5**.

To generate the prey strains for the Y1H screen, each of the 384 TF-encoding plasmids was

individually purified from corresponding glycerol stocks of the 1956 DEST-U Arabidopsis

TF ORFeome collection (CD4-89, ABRC, USA) using QIAprep Spin Miniprep Kit. 100 ng

of purified plasmid were individually transformed into Yalpha1867 strain (Gift from Dr. Ram

Yadav, IISER Mohali, INDIA) using the high throughput 96-well format transformations

protocol described earlier. Following transformation, the cells were spotted on synthetic

complete media lacking tryptophan (Sc-Trp) media and incubated at 30ºC till colonies

appeared.

To generate the diploids used in the Y1H assay, prey TF-encoding plasmids was individually

purified from corresponding glycerol stocks of the 1956 DEST-U Arabidopsis TF ORFeome

collection (CD4-89, ABRC, USA) using QIAprep Spin Miniprep Kit. 100 ng of purified

plasmid were individually transformed into *pMIR319C* baits using the high-efficiency LiAc-

PEG method described earlier. Following transformation, the cells were spotted on synthetic

complete media lacking histidine and tryptophan (Sc-His-Trp) media and incubated at 30ºC

till colonies appeared.

Yeast one-hybrid (Y1H) screen/assay

The auto activity of yeast baits on Sc-His due to minimal *HIS3* expression was determined

by spotting them on Sc-His media containing 0-30 mM 3-Amino-1,2,4-triazole (3-AT)

(A8056, Sigma-Aldrich, USA). The minimum concentration of 3-AT that completely inhibits

the growth of the bait was considered for the Y1H screen/ assay. For the mating-based Y1H

screen, the diploids generated by mating were resuspended in sterile 96-well plates

containing Sc-Trp-His media and allowed to grow in shaking incubators for 24 hrs at 30ºC.

After 24 hrs of growth, OD was measured. Volumes corresponding to equal OD were spotted

on Sc-Trp-His+3AT media, incubated at 30ºC, and scored for growth on days 3-5 post-

incubation. Spots containing at least 5 colonies were considered positive for growth. For the

Y1H screen, spots showing growth on all three concentrations of 3-AT, i.e., 20, 25, and 30

mM by day 5, or growth on 20 and 25 mM 3-AT by day 3 were considered positive for

binding. For the co-transformation-based Y1H assay, yeast baits were transformed with TF

encoding plasmids and selected on Sc-Trp-His media. Following selection, the cells were

resuspended in Sc-Trp-His media and allowed to grow in shaking incubators for 24 hrs at

30ºC. After 24 hrs of growth, OD was measured. Volumes corresponding to equal OD were

spotted on Sc-Trp-His media with respective concentrations of 3-AT and incubated at

30ºC.The plates were inspected for growth and imaged over days 3- 5 post-incubation.

Plant materials

Arabidopsis thaliana ecotype Columbia-0 (Col-0) was used as the wild-type control. All the

lines used in this study are in Col-0 genetic background. The genotypes used in this study are

described in **Suppl Data Set. 6**. The genotypes *cuc2-1* and *pBLS>>miR319a* (Originally in

L*er*; Efroni et al. 2008; Hasson et al. 2011) was brought into the Col-0 background by back

crossing three times. *pMIR319C::GUS[#1]* (Nag et al. 2009) was established in various mutant

backgrounds by crossing and was confirmed by GUS assays in the F3/F4 generation.

cuc2-3;miR319c-1, *cuc2-3;miR319ab* and *miR319a;miR319b;miR319c-1* mutants were

generated by crossing respective mutant lines *cuc2-3* (Nikovics et al. 2006), *miR319ab*

(Koyama et al. 2017), *miR319c-1* (SAIL_647_B03/ CS827925) and confirmed by genotyping

in F3 generation. *cuc2-1D;GR[#2]* was established by crossing *cuc2-1D* (Larue et al. 2009) and

Col-0;*GR[#2]* (Shankar et al. 2023) mutants and confirmed by phenotyping in F3 generation.

All genotyping primers are listed in **Suppl Data Set. 5**.

Plant growth condition and chemical treatments

For GUS assays and phenotyping experiments, seeds were surface-sterilized with 0.05% SDS

containing 70% ethanol solution for 10 min, followed by wash with 100% ethanol twice and

sown on 1X Murashige and Skoog (MS), 0.8% agar medium (PT011, HiMedia Laboratories,

India) containing 0.025% ethanol (Mock) or 6 μM dexamethasone (DEX) (D4902, Sigma-

Aldrich, USA). After sowing, seeds were kept for stratification for three days in the dark at

4°C and then shifted to a plant growth chamber (I-41LL, Percival Scientific, USA) set at

22°C, 60% relative humidity, long-day conditions (16h/8h light/dark period, 100 $\mu mol.m^{-2}.s^{-1}$

^{1}). 14-day old seedlings were transplanted onto a soil mixture containing Soilrite and Perlite

(Keltech Energies, India) in a 3:1 ratio. Plants were grown till maturity in growth rooms with

the conditions as described for growth chamber.

For transient DEX and cycloheximide (CHX) treatments, 9 day-old mock (1X MS, 0.8% agar

medium containing 0.025% ethanol) grown seedlings were transferred onto petri dishes filled

with 0.5X MS medium, supplemented with 1% sucrose, 1X Gamborg B5 vitamins (PL018,

HiMedia Laboratories, India), 12 µM DEX, and/or 10 µM CHX (C7698, Sigma-Aldrich,

USA), incubated for 2 hours and harvested under liquid N2 prior to RNA isolation.

For confocal imaging experiments, seeds were stratified in the dark for two days at 4°C to

synchronize germination. Plants were grown on petri dishes filled with 0.5X MS medium,

supplemented with 0.8% agar, 1% sucrose, and 0.1% Plant Protective Mixture (Plant Cell

Technology) in a growth chamber under long-day conditions (16h/8h light/dark period, 95

$\mu mol.m^{-2}.s^{-1}$) at $22 \pm 1°C$ with 60-70% relative humidity.

DNA constructs and plant transformation

For the generation of *pMIR319C* deletion and *mCUCBSpMIR319C::GUS* constructs, a 2.7 kb

genomic fragment corresponding to the *MIR319C* URR from the start of pre-*MIR319C* was

amplified from Col-0 genomic DNA using Phusion High Fidelity DNA polymerase (M0530,

New England Biolabs, USA) and cloned into modified *P4P1R* entry vector (a kind gift from

Ram Yadav, IISER Mohali, India) by gateway BP reaction (11789020, Invitrogen, USA) to

generate the *pMIR319C-P4P1R* construct. To introduce mutation at the CUC-binding site, we

utilized a PCR-based site directed mutagenesis approach. Briefly, forward, and reverse

primers carrying mutations in CUC-binding site (mCUCBS) was first used in a PCR reaction

with *pMIR319C-P4P1R* template using Phusion DNA polymerase. Post PCR reaction, the

product was digested with *Dpn*I (R0176, New England Biolabs, USA) and transformed into

E. coli and selected on LB agar containing 100 mg/l ampicillin. Plasmids were purified from

resulting colonies using QIAprep Spin Miniprep Kit (Qiagen, Germany) and mCUCBS was

confirmed by Sanger sequencing. The entry clone *mCUCBSpMIR319C-P4P1R* was

recombined into *R4L1pGWB532* destination vector (Gift from Tsuyoshi Nakagawa, Shimane

University, Japan) by LR reaction (11791020, Invitrogen, USA) to generate the

mCUCBSpMIR319C::GUS construct. Sequence information of primers used to generate the

constructs are provided in the **Suppl Data Set. 5**. *miR319c* targeting CRISPR constructs were

generated based on zCas9i modular cloning approach (Grützner et al. 2021, Stuttmann et al.

2021). The zCas9i cloning kit: MoClo-compatible CRISPR/Cas9 cloning kit with intronized

Cas9 was a gift from Sylvestre Marillonnet and Johannes Stuttmann (Addgene kit

#1000000171). The guide RNAs- sgRNA1 (5'-AGTGACTGTGTGAATGATG-3') and

sgRNA2   (5'-TATCTGTGTTTGGACTGAA-3') were

cloned into pDGE332 and pDGE334 shuttle vectors and assembled into pDGE347 vector

according to cloning manual v5 of the zCas9i cloning kit.

mCUCBSpMIR319C::GUS and the *miR319c* CRISPR constructs were integrated into Col-0

background by the *Agrobacterium tumefaciens*-mediated floral dip method (Clough and Bent,

1998). For *mCUCBSpMIR319C::GUS* line generation, seeds were harvested and screened

for the ability to grow on MS media in presence of 20 mg/l hygromycin. At least five

independent lines were established for each construct and confirmed by genotyping in T3

generation. Sequence information of primers used for genotyping are provided in the **Suppl**

Data Set. 5. For generating CRISPR knockouts of *miR319c*, seeds were harvested and

screened for red fluorescence and for the ability to grow on MS media in presence of 10 mg/l

BASTA. T2 seedlings from ten independent T1 plants were screened for deletions in

miR319c region by Sanger sequencing. Two independent lines, *miR319c-2* and *miR319c-3*,

carrying deletions in the miR319c region were established in T3 generation after counter

selecting for lack of red fluorescence in T2 population.

β-glucuronidase (GUS) assay

GUS assays were performed as described earlier (Shankar et al. 2023). Briefly, samples were

collected in ice-cold 90% acetone, incubated at room temperature for 20 min, followed by

washing with ice-cold staining buffer (50 mM sodium phosphate, pH 7.0, 0.2% Triton X-100,

5 mM potassium ferrocyanide, and 5 mM potassium ferricyanide). Fresh staining buffer was

added with 2 mM X-Gluc (R0851, Thermo Scientific, USA) and incubated overnight at 37°C

in dark. Staining buffer was replaced with 70% ethanol, and 70% ethanol was changed 3-4

times in regular intervals to clear the tissue. Shoot apices and leaf primordia were dissected

and mounted on a glass slide containing lactic acid under the microscope and were imaged

using Olympus BX51 trinocular DIC microscope fitted with ProgResC3 camera and

ProgResCapture Pro2.6 software (Jenoptik, Germany) with the following settings, exposure-

65ms, brightness-10 units, contrast- 35 units, colour temperature- 5000K, image resolution-

(1040 x 770 2X Bin), while keeping all other settings in default mode.

GUS domain, leaf area and pavement cell area quantification

The extent of reporter gene expression in the *pMIR319C::GUS* and

mCUCBSpMIR319C::GUS transgenic lines were determined using a pixel density

quantification method reported in Shankar et al., 2023. Images of 6 and 9-day old *cuc2-*

3;miR319ab seedlings were captured using LEICA EZ4 E stereomicroscope equipped with
LAS EZ software. Areas of the dissected mature first pair of leaves from 25-day old plants
were determined using the polygon selection tool of the ImageJ software. For pavement cell
area measurements, dissected mature first pair of leaves were first incubated in 7:1 absolute
ethanol: glacial acetic acid clearing solution. After 24 hours, clearing solution was replaced
with 1M KOH solution for 30 minutes, washed twice with sterile water and incubated in 37%
D-lactic acid solution until tissue became completely transparent. The clarified leaf samples
were mounted on a glass slide using D-lactic acid and imaged at 20x magnification on a
Olympus BX51 trinocular DIC microscope fitted with ProgResC3 camera and
ProgResCapture Pro2.6 software (Jenoptik, Germany). Cell areas were determined using the
polygon selection tool of the ImageJ software. The average cell number was derived using
the leaf area and cell area of leaves taken for the experiment. Silhouettes of cleared leaves
were used for representative purposes using an exact scale of the image captured.

Confocal microscopy and imaging

For imaging of the first pair of leaf, one cotyledon was removed using fine tweezers or an
injection needle from two-day-old plants to expose the initiating leaf primordium for imaging
of leaf at 1 day after initiation (DAI). The seedlings were stained with 1 mg/mL Propidium
Iodide (PI) solution for 2-3 minutes and then were rinsed with water. The dissected plants
then were placed horizontally in Ø60 mm Petri dishes filled with 0.5X MS medium,
supplemented with 1.5% agar, 1% sucrose, and 0.1% Plant Protective Mixture (Plant Cell
Technology). Plants were immersed in water containing 0.1% PPM for imaging. Three to
four replicates were imaged using the PI staining and at least half of the abaxial leaf surface
was imaged at 1 DAI, 3 DAI and 5 DAI respectively.

All confocal imaging was performed with an LSM980 upright confocal microscope using a
long-distance water-dipping objective (AP 20X/1.0; Zeiss) at the confocal facility of
MechanoDevo team (RDP, ENS de Lyon). Excitation was performed using a diode laser with
561 nm for PI. Images were collected at 580-650 nm for PI with a GaAsP detector and a
beam splitter operating at 488 and 561 nm. Confocal stacks were acquired at 512x512
resolution and 16-bit image depth, with 1.4 µm distance in Z-dimension. For samples larger
than the field of view, multiple overlapping stacks were obtained and later stitched together
using MorphoGraphX (Barbier de Reuille et al., 2015; Strauss et al., 2022).

Confocal Image analysis

Cellular growth and expression quantifications were conducted using MorphoGraphX. Stacks were processed as described previously (Le Gloanec et al. 2022). After digitally removing the trichomes when necessary, the organ surface was detected with the 'Edge detect' tool with a threshold of 20000 with 'closing' to fill holes, followed by the 'Edge detect angle' (10000). An initial 5 µm cube size mesh was then created and subdivided three times before projecting the membrane signal (2-5 µm). Cell segmentation was performed manually. Metrics such as area cell size were computed as described before (Barbier de Reuille et al. 2015; Sapala et al. 2018).

RNA isolation and RT-qPCR

Total RNA was extracted using the TRIzol method described earlier (Challa et al. 2016). 10 µg of RNA was used for DNase (EN0521, Thermo Scientific, USA) treatment. 1.5 µg DNase-treated RNA was converted to cDNA using RevertAid Reverse Transcriptase (EP0441, Thermo Scientific, USA) and 2.5 µM oligo dT according to the manufacturer's instructions. Quantitative PCR reactions with 25 ng cDNA and 0.4 µM primers were set up using the DyNAmo Flash SYBR Green RT-qPCR kit (F415L, Thermo Scientific, USA), according to the manufacturer's instructions. 10 µl qPCR reactions were carried out in QuantStudio™ 6 Flex Real-Time PCR System, 384-well format (Applied Biosystems, USA). Results were analyzed using the inbuilt QuantStudio6 Pro software, and $\Delta\Delta$CT values were determined after normalization with *PROTEIN PHOSPHATASE 2A SUBUNIT A3 (PP2AA3)*. Relative fold changes in transcript levels were calculated using the formula $2^{-\Delta\Delta CT}$. All primer sequences are provided in **Suppl Data Set. 5**.

Statistical analysis

All graphical representations and statistical analysis were performed using GraphPad Prism 9 software (GraphPad, USA). First, samples were tested for Gaussian distribution using the Shapiro-Wilk normality test. If samples were normally distributed, pairwise comparisons were performed using unpaired *t*-test. Welch's correction was applied if variances were different among the samples. Statistical differences among multiple genotypes were determined by ordinary One-Way ANOVA followed by Tukey's multiple comparisons test or One-Way ANOVA followed by Dunnet's T3 post-hoc test, depending on whether variances were significantly different among the genotypes. If the samples failed the normality test,

then pairwise comparisons were carried out by Mann Whitney test. Statistical differences

among multiple genotypes were determined by One-Way ANOVA followed by Dunn's post-

hoc test. To compare statistical differences among multiple genotypes subjected to ethanol

(Mock) or dexamethasone (DEX) treatments, we used Two-Way ANOVA followed by

Tukey's post-hoc test. p-values obtained from the respective tests are indicated above the

data bars in several plots. For plots with a large number of comparisons, statistical

differences among samples are indicated by different letters. Raw data and statistics summary

of all the plots are provided in **Suppl Data Set. 7**.

List of Supplementary Data Files

Suppl Data Set 1. List of 378 candidate TFs used in yeast one hybrid screen (TF mini library)

Suppl Data Set 2. List of 57 TF positives from yeast one hybrid screen

Suppl Data Set 3. List of SAM and leaf primordia-specific TFs (343)

Suppl Data Set 4. List of TFs from TFBS analysis on *MIR319C* promoter (49)

Suppl Data Set 5. List of primers used in this study (5' to 3')

Suppl Data Set 6. List of genotypes used in this study

Suppl Data Set 7. Raw data and statistics summary

Accession numbers

TAIR (The Arabidopsis Information Resource, www.arabidopsis.org) accession numbers of

the significant genes used in this study are *AT4G23713* (*MIR319A*), *AT5G41663* (*MIR319B*),

AT2G40805 (*MIR319C*), *AT3G15170* (*CUC1*), *AT5G53950* (*CUC2*), *AT1G76420* (*CUC3*),

AT3G15030 (*TCP4*).

Competing Interests

The authors declare no conflict of interest.

Funding

This work was supported by fellowships from the Ministry of Education (N.S., V.M.),

Department of Biotechnology (S.R.), Council of Scientific and Industrial Research (A.G.),

and Anusandhan National Research Foundation (JC Bose Fellowship to U.N.) from

Government of India. V.M. was also supported by EMBO Scientific exchange grant 10242.
Authors thank the Department of Science & Technology for Improvement of S&T
Infrastructure (DST-FIST, No. SR/FST/LSII-044/2016 dated 15.12.2016) and the
Department of Biotechnology (DBT)-IISc Partnership Program Phase-II at IISc (sanction No.
BT/PR27952/INF/22/212/ 2018) for the funding and infrastructure support. U.N. also
acknowledges the financial support from the Ministry of Education, Govt. of India, through a
STARS grant (MoE-STARS/STARS-2/2023-0930). The funders had no role in study design,
data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We sincerely thank Ram Yadav (IISER Mohali, India) for vectors and strains used in Y1H
assays, Koyama Tomotsugu (Bioorganic Research Institute, Kyoto, Japan) for the *miR319ab*
seeds, Patrick Laufs (Institut Jean Pierre Bourgin, INRA Versailles, France) for the
pCUC2::GUS line, Yuling Jiao (Peking University, Beijing, China) for *pRPS5A::CUC2-GR-*
HA line. We thank Sachin Kotak, Yadukrishnan Premachandran and Anurag Sharma of IISc
Bangalore, India, for helpful discussions during the preparation of this manuscript.
783

Author contributions

N.S. initiated the project, designed & performed most experiments, analysed & interpreted
the results, organized the figures, wrote the first draft of the manuscript and contributed to its
finalization. V.M. performed confocal microscopy and growth kinematics experiments,
collected data and assisted in organizing the figures. A.G. assisted in some of the RT-qPCR
experiments and imaged mature leaves, young seedlings for some of the phenotyping
experiments. V.M. and A.G. performed most of the cell area measurements. S.R. generated
miR319c-2, and *miR319c-3* lines, performed some of the GUS expression experiments. A.J.P.
established the *miR319c-1* line. A.V. made CRISPR constructs for generation of *miR319c-2*
and *miR319c-3* lines. O.H. contributed to data interpretation and conclusions, corrected the
manuscript and provided the infrastructure for confocal imaging and analysis. U.N
contributed to designing experiments and data interpretation, guided the first six co-authors,
corrected and finalised the manuscript.

References

Aida M, Ishida T, Fukaki H, Fujisawa H, Tasaka M Genes involved in organ separation in

Arabidopsis: An analysis of the cup-shaped cotyledon mutant. Plant Cell 1997: **9**: 841–857

Aida M, Ishida T, Tasaka M Shoot apical meristem and cotyledon formation during

Arabidopsis embryogenesis: interaction among the CUP-SHAPED COTYLEDON and

SHOOT MERISTEMLESS genes. Development. 1999:**126**: 1563–1570

Andriankaja M, Dhondt S, DeBodt S, Vanhaeren H, Coppens F, DeMilde L,

Mühlenbock P, Skirycz A, Gonzalez N, Beemster GTS, et al Exit from Proliferation during

Leaf Development in Arabidopsis thaliana: A Not-So-Gradual Process. Dev Cell. 2012:**22**:

64–78

Bar M, Ori N Leaf development and morphogenesis. Development. 2014: **141**: 4219–4230

Barbier de Reuille P., Routier-Kierzkowska A.L., Kierzkowski D., Bassel GW.,

Schüpbach T., Tauriello G., Bajpai N, Strauss S, Weber A, Kiss A et al. MorphoGraphX:

A platform for quantifying morphogenesis in 4D. Elife. 2015: 4: 05864.

doi:10.7554/eLife.05864

Barton MK Twenty years on: The inner workings of the shoot apical meristem, a

developmental dynamo. Dev Biol. 2010: **341**: 95–113

Bhatia N, Wilson-Sánchez D, Strauss S, Vuolo F, Pieper B, Hu Z, Rambaud-Lavigne L,

and Tsiantis M. Interspersed expression of CUP-SHAPED COTYLEDON2 and REDUCED

COMPLEXITY shapes Cardamine hirsuta complex leaf form. Current Biology.

2023:33(14):2977-2987.e6. https://doi.org/10.1016/j.cub.2023.06.037

Bilsborough GD, Runions A, Barkoulas M, Jenkins HW, Hasson A, Galinha C, Laufs P,

Hay A, Prusinkiewicz P, Tsiantis M. Model for the regulation of Arabidopsis thaliana leaf

margin development. Proc Natl Acad Sci U S A. 2011: **108**: 3424–3429

Challa KR, Aggarwal P, Nath U. Activation of *YUCCA5* by the Transcription Factor TCP4

Integrates Developmental and Environmental Signals to Promote Hypocotyl Elongation in

Arabidopsis. Plant Cell. 2016: **28**: 2117–2130

Challa KR, Rath M, Nath U. The CIN-TCP transcription factors promote commitment to differentiation in Arabidopsis leaf pavement cells via both auxin-dependent and independent pathways. PLoS Genet. 2019: **15**: 1–30

Clough SJ, Bent AF. Floral dip: A simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J. 1998: **16**: 735–43

Donnelly PM, Bonetta D, Tsukaya H, Dengler RE, Dengler NG. Cell cycling and cell enlargement in developing leaves of Arabidopsis. Dev Biol. 1999: **215**: 407–419

Du F, Guan C, Jiao Y. Molecular Mechanisms of Leaf Morphogenesis. Mol Plant. 2018: **11**: 1117–1134

Efroni I, Blum E, Goldshmidt A, Eshed Y. A Protracted and Dynamic Maturation Schedule Underlies Arabidopsis Leaf Development. the Plant Cell Online. 2008: **20**: 2293–2306

Fox S, Southam P, Pantin F, Kennaway R, Robinson S, Castorina G, Sánchez-Corrales YE, Sablowski R, Chan J, Grieneisen V, et al. Spatiotemporal coordination of cell division and growth during organ morphogenesis. PLoS Biol. 2018: **16**: e2005952

Fuxman Bass JI, Pons C, Kozlowski L, Reece-Hoyes JS, Shrestha S, Holdorf AD, et al. A gene-centered C. elegans protein–DNA interaction network provides a framework for functional predictions. Mol Syst Biol. 2016: **12**: 884

Gázquez A, Beemster GTS. What determines organ size differences between species? A meta-analysis of the cellular basis. New Phytologist. 2017: **215**: 299–308

Gonzalez N, Vanhaeren H, Inzé D. Leaf size control: Complex coordination of cell division and expansion. Trends Plant Sci. 2012: **17**: 332–340

Grutzner R, Martin P, Horn C, Mortensen S, Cram EJ, Lee-Parsons CWT, Stuttmann J, Marillonnet S. High-efficiency genome editing in plants mediated by a Cas9 gene containing multiple introns. Plant Commun. 2021:2:100135.

Hasson A, Plessis A, Blein T, Adroher B, Grigg S, Tsiantis M, Boudaoud A, Damerval C, Laufs P. Evolution and diverse roles of the CUP-SHAPED COTYLEDON genes in Arabidopsis leaf development. Plant Cell. 2011: **23**: 54–68

Hepworth J, Lenhard M. Regulation of plant lateral-organ growth by modulating cell number and size. Curr Opin Plant Biol. 2014: **17**: 36–42

Hibara KI, Karim MR, Takada S, Taoka KI, Furutani M, Aida M, Tasaka M. Arabidopsis CUP-SHAPED COTYLEDON3 regulates postembryonic shoot meristem and organ boundary formation. Plant Cell. 2006:**18**: 2946–2957

Hu ZL, Wilson-Sánchez D, Bhatia N, Rast-Somssich MI, Wu A, Vlad D, McGuire L, Nikolov LA, Laufs P, Gan X, et al. A CUC1/auxin genetic module links cell polarity to patterned tissue growth and leaf shape diversity in crucifer plants. Proc Natl Acad Sci U S A. 2024:121(26). https://doi.org/10.1073/pnas.2321877121

Ishida T, Aida M, Takada S, Tasaka M. Involvement of CUP-SHAPED COTYLEDON genes in gynoecium and ovule development in Arabidopsis thaliana. Plant Cell Physiol. 2000: **41**: 60–67

Kalve S, De Vos D, Beemster GTS. Leaf development: a cellular perspective. Front Plant Sci . 2014:**5**: 1–25

Koyama T, Furutani M, Tasaka M, Ohme-Takagi M.TCP Transcription Factors Control the Morphology of Shoot Lateral Organs via Negative Regulation of the Expression of Boundary-Specific Genes in Arabidopsis. Plant Cell. 2007: **19**: 473–484

Koyama T, Mitsuda N, Seki M, Shinozaki K, Ohme-Takagi M. TCP transcription factors regulate the activities of ASYMMETRIC LEAVES1 and miR164, as well as the auxin response, during differentiation of leaves in Arabidopsis. Plant Cell. 2010: **22**: 3574–3588

Koyama T, Sato F, Ohme-Takagi M. Roles of miR319 and TCP transcription factors in leaf development. Plant Physiol. 2017: **175**: 874–885

Kuchen EE, Fox S, De Reuille PB, Kennaway R, Bensmihen S, Avondo J, et al. Generation of Leaf Shape Through Early Patterns of Growth and Tissue Polarity. 2012; 335:1092–6

Larue CT, Wen J, Walker JC. A microRNA-transcription factor module regulates lateral organ size and patterning in Arabidopsis. Plant Journal. 2009: **58**: 450–463

Le Gloanec C, Collet L, Silveira SR, Wang B, Routier-Kierzkowska AL, and Kierzkowski D. Cell type-specific dynamics underlie cellular growth variability in plants. Development. 2022: **149**: dev200783.

Lindemose S, Jensen MK, De Velde J Van, O'Shea C, Heyndrickx KS, Workman CT, Vandepoele K, Skriver K, Masi F De. A DNA-binding-site landscape and regulatory

network analysis for NAC transcription factors in Arabidopsis thaliana. Nucleic Acids Res.
2014: **42**: 7681–7693

Maugarny-Calès A, Cortizo M, Adroher B, Borrega N, Gonçalves B, Brunoud G,
Vernoux T, Arnaud N, Laufs P. Dissecting the pathways coordinating patterning and
growth by plant boundary domains. PLoS Genet. 2019: **15**: 1–30

Maugarny-Calès A, Laufs P. Getting leaves into shape: A molecular, cellular, environmental
and evolutionary view. Development. 2018: **145**: dev161646

Nag A, King S, Jack T. miR319a targeting of TCP4 is critical for petal growth and
development in Arabidopsis. Proceedings of the National Academy of Sciences. 2009: **106**:
22534–22539

Nikovics K, Blein T, Peaucelle A, Ishida T, Morin H, Aida M, Laufs P. The balance
between the MIR164A and CUC2 genes controls leaf margin serration in Arabidopsis. Plant
Cell. 2006: **18**: 2929–2945

Ooka H, Satoh K, Doi K, Nagata T, Otomo Y, Murakami K, Matsubara K, Osato N,
Kawai J, Carninci P, et al. Comprehensive Analysis of NAC Family Genes in Oryza sativa
and Arabidopsis thaliana. DNA Research. 2003: **10**: 239–247

Palatnik JF, Allen E, Wu X, Schommer C, Schwab R, Carrington JC, Weigel D. Control
of leaf morphotgenesis by microRNAs. Nature. 2003: **425**: 257–263

Pruneda-Paz, JL, Breton, G, Nagel, DH, Kang, SE, Bonaldi, K, Doherty, CJ, Ravelo, S,
Galli, M, Ecker, JR, and Kay, SA. A Genome-Scale Resource for the Functional
Characterization of Arabidopsis Transcription Factors. Cell Rep. 2014: **8**: 622–632

Rodriguez RE, Mecchia MA, Debernardi JM, Schommer C, Weigel D, Palatnik JF.
Control of cell proliferation in Arabidopsis thaliana by microRNA miR396. Development.
2010: 137:103–12

Rodriguez RE, Debernardi JM, Palatnik JF. Morphogenesis of simple leaves: Regulation
of leaf size and shape. Wiley Interdiscip Rev Dev Biol. 2014: 3: 41–57

Rubio-Somoza I, Zhou CM, Confraria A, Martinho C, Von Born P, Baena-Gonzalez E,
Wang JW, Weigel D. Temporal control of leaf complexity by miRNA-regulated licensing of
protein complexes. Current Biology. 2014: **24**: 2714–2719

Sapala, A, Runions A, Routier-Kierzkowska AL, Gupta MD, Hong L, Hofhuis H,

Verger S, Mosca G, Li CB, Hay A et al. Why plants make puzzle cells, and how their shape

emerges. Elife. 2018: **7**: e32794.

Sarvepalli K, Das Gupta M, Challa KR, Nath U. Molecular cartography of leaf

development — role of transcription factors. Curr Opin Plant Biol. 2019: **47**: 22–31

Sarvepalli K, Nath U. CIN-TCP transcription factors: Transiting cell proliferation in plants.

IUBMB Life. 2018: **70**: 718–731

Schneider M, Van Bel M, Inzé D, Baekelandt A. Leaf growth – complex regulation of a

seemingly simple process. Plant Journal. 2023: **117**: 1018–1051

Scofield S, Murison A, Jones A, Fozard J, Aida M, Band LR, Bennett M, Murray JAH.

Coordination of meristem and boundary functions by transcription factors in the SHOOT

MERISTEMLESS regulatory network. Development. 2018: **145**: dev157081

Shankar N, Sunkara P, Nath U. A double-negative feedback loop between miR319c and

JAW-TCPs establishes growth pattern in incipient leaf primordia in Arabidopsis thaliana.

PLoS Genet. 2023: **19**: 1–25

Strauss S, Runions A, Lane B, Eschweiler D, Bajpai N, Trozzi N, Routier-Kierzkowska

AL, Yoshida S, Rodrigues da Silveira S, Vijayan A et al. Using positional information to

provide context for biological image analysis with MorphoGraphX 2.0. Elife. 2022: **11**:

e72601

Stuttmann J, Barthel K, Martin P, Ordon J, Erickson, J L, Herr R, Ferik F, Kretschmer

C, Berner T, Keilwagen J.et al. Highly efficient multiplex editing: one-shot generation of

8× Nicotiana benthamiana and 12× Arabidopsis mutants. Plant J. 2021: **106,** 8-22**.**

Tian C, Zhang X, He J, Yu H, Wang Y, Shi B, Han Y, Wang G, Feng X, Zhang C, et al.

An organ boundary enriched gene regulatory network uncovers regulatory hierarchies

underlying axillary meristem initiation. Mol Syst Biol. 2014: **10**: 755

Tian, C., Wang, Y., Yu, H., He, J., Wang, J., Shi, B., Du, Q., Provart, N.J., Meyerowitz,

E.M., and Jiao, Y. A gene expression map of shoot domains reveals regulatory mechanisms.

Nat. Commun. 2019: **10**: 1–12.

Vroemen CW, Mordhorst AP, Albrecht C, Kwaaitaal MACJ, De Vries SC. The CUP-
SHAPED COTYLEDON3 gene is required for boundary and shoot meristem formation in
Arabidopsis. Plant Cell. 2003: **15**: 1563–1577

Wang Q, Hasson A, Rossmann S, Theres K. Divide et impera: Boundaries shape the plant
body and initiate new meristems. New Phytologist. 2016: **209**: 485–498

Main figures

Figure 1: Identification of CUC proteins as activators of *MIR319C* transcription. **(A)** Abundance of transcription factor (TF) family members (in %) recovered from the yeast one hybrid screen (Y1H) with 378 TFs as prey proteins and 2736 bp *MIR319C* URR fragment as bait. **(B)** A schematic representation of truncated *MIR319C* promoter fragments used for Y1H assay with selected NAC TFs as prey proteins. Grey box corresponds to *HIS3 CDS*. The three bold horizontal lines below the URR scheme represent the promoter regions with truncation (dotted lines), i.e., p^{distal} (-2736-1866 bp), p^{middle} (-1866-1106 bp) and $p^{proximal}$ (-1106 bp), respectively. The consensus CUC-binding sequence (BS) TTACGT is located between -2616 bp to -2621 bp positions. **(C)** Images of yeast growth harboring three different truncated *MIR319C* promoter regions upstream to the *HIS3* reporter gene and corresponding NAC TF preys as indicated, grown in the presence of inhibitory concentration of 3-AT. NAC027 served as a negative control. **(D)** Bright field images of shoot apices (top panel) or 6[th] leaf primordia (bottom panel) of 10-day old seedlings, processed for GUS-staining, of indicated genotypes expressing *pMIR319C::GUS* transgene. Black lines highlight the boundaries. Numbers below the images in lower panel indicate leaf lengths; GUS domain length relative to the leaf length (Relative GUS domain) is indicated within parentheses. Scale bar, 100 μm. **(E)** Violin plots displaying the relative GUS domain (Y-axis) in the genotypes indicated on the X-axis. Differences among samples are indicated by *p*-values on top of the data bars; one-way ANOVA, Dunnett's post hoc test was used. N, 20-32. **(F)** Level of *GUS,* pre-*MIR319C* and *CUC2* transcripts quantified by RT-qPCR in genotypes indicated on the X-axis. Error bars represent SEM of four biological replicates. Multiple comparisons were performed using One-way ANOVA followed by Dunnett's post hoc test. *p*-values are shown above the data bars.

Figure 2: Role of *CUC2-MIR319C* module in leaf cell proliferation. (A) Schematic representation of the *pMIR319C::GUS* construct with mutated bases (red) on the putative CUC-binding site (TTACGT). Numbers indicate nucleotide positions upstream to transcription start site. The *GUS* coding sequence is indicated by a grey box. **(B-C)** Bright field images of processed shoot apices (left) and leaf primordia expressing *GUS*-based reporter transgene under wildtype (*pMIR319C::GUS*, **B**) or mutated (*mCUCBSpMIR319C::GUS*, **C**) *MIR319C* promoters. Numbers below the images indicate leaf length (larger leaf primordia are considered for the apices); relative GUS domain values are indicated within parentheses. Black lines highlight the boundaries. **(D)** Violin plot representations of relative GUS domain in the 50-750 µm long leaf primordia (N, 25-31) on 4^{th}-8^{th} nodes of indicated genotypes. Differences between samples are indicated by *p*-value above the data points; unpaired *t*-test with Welch's correction for differences in variance was used. **(E)** Transcript level of genes indicated below the X-axis labels in 10-day old *pRPS5A::CUC2-GR-HA* seedlings treated for 2 hours with ethanol (Mock) or 12 µM dexamethasone (DEX) and/or 10 µM cycloheximide (CHX). Error bars represent SEM of three biological replicates. Pairwise comparisons were performed using unpaired *t*-test; *p*-values are shown above the data bars. **(F)** Representative images of mature first pair of leaves from 25-day old plants of indicated genotypes. Scale bar, 5 mm. **(G)** Areas (Y-axis) of mature first pair of leaves of indicated genotypes. N, 15-44. Significant differences among the samples are indicated by lower-case letters. $p < 0.0001$. One-way ANOVA, followed by Dunn's multiple comparison test were performed to determine the significant differences among samples. **(H)** Total number of pavement cells in mature first pair of leaves of the indicated genotypes. Cell number was determined by dividing leaf area by cell area (100-120 cells per leaf) averaged over 3-6 leaves per genotype. Significant differences among samples are indicated by lower-case letters. $p < 0.01$; one-way ANOVA, followed by Tukey's multiple comparison test were performed.

Figure 3: Growth kinematics of *cuc2-3;miR319c-1* leaves. **(A)** Heat maps of cell areas in the first pair of Col-0 and *cuc2-3;miR319c-1* leaves at 1, 3, and 5 days after initiation (DAI). Scale bar, 100 μm. **(B)** Impressions of Col-0 and *cuc2-3;miR19c-1* leaves at indicated DAI. Scale bar, 5 mm. **(C-E)** Violin plot representations of leaf area **(C)**, total pavement cell number **(D)**, and average pavement cell area **(E)** of mature first leaf pair of Col-0 and *cuc2-3;miR319c-1* leaves at DAI indicated on the X-axes. Bold horizontal lines within the datapoints represent median. Differences among samples are indicated by *p*-values above data points; unpaired *t*-test with Welch's correction for differences in variance **(C, D)** or Mann Whitney U test **(E)** were used. N, 5-15 leaves **(C, D)** or 199-447 cells **(E)**.

Figure 4: Genetic interaction between *CUC2* and *MIR319* genes in regulating leaf cell proliferation. (A, B) Representative images of mature first pair of leaves dissected from 25-day old plants of the indicated genotypes. Scale bar, 5 mm. **(C)** Areas of mature first pair of leaves of the indicated genotypes. N, 5-10 leaves. Significant differences among the samples are indicated by lower-case letters. p<0.0001; one-way ANOVA, followed by Dunnett's T3 multiple comparison test were performed to determine the significant differences among samples. **(D)** Total number of pavement cells in mature first pair of leaves of the indicated genotypes, as derived from dividing leaf area by cell area (75-100 cells per leaf) averaged from 3-4 leaves per genotype. Significant differences among the samples are indicated by lower-case letters. p<0.05; one-way ANOVA, followed by Tukey's multiple comparison test was performed to determine the significant differences among samples. **(E)** Areas of mature first pair of leaves of the indicated genotypes. N, 6-31 leaves. Significant differences among the samples are indicated by lower-case letters. p<0.01; one-way ANOVA, followed by Dunnett's T3 multiple comparison test was performed to determine the significant differences among samples.

Figure 5: Genetic interaction between *CUC2* and *JAW-TCP* genes in regulating leaf size. (A, B) Representative images of mature first leaves from 25-day old plants of the indicated genotypes. Scale bar, 5 mm. **(C, D)** Areas of mature first pair of leaves of indicated genotypes. N, 13-35. Significant differences among samples are indicated by different lower-case letters. $p<0.01$; two-way ANOVA, followed by Tukey's post hoc test (**C**) or Dunnett's T3 multiple comparison test (**D**) were performed to determine the significant differences among samples. Mock and DEX (**A**, **C**) indicate continuous treatment with solvent (Mock) or 12 μM dexamethasone (DEX).

Figure 6: Regulation of *MIR319C* expression by *CUC2* and *JAW-TCP* genes in leaf primordia. (A) Bright field images of cleared shoot apices (top panel) and 6th leaf primordia (bottom panel) from 10-day old Col-0 and *jaw-D* seedlings expressing *pCUC2::GUS* transgene. Black lines highlight borders of the images. Scale bar, 50 μm. **(B, C)** Bright-field images of cleared shoot apices (top panel) and 6th leaf primordia (bottom panel) dissected from 10-day old seedlings of indicated genotypes expressing the *pMIR319C::GUS* transgene. Black lines highlight borders. Scale bar, 50 μm. Numbers below the images represent leaf lengths in μm and relative GUS domain values (in parentheses). Mock and DEX in **(B, D)** indicate continuous treatment with solvent (Mock) or 12 μM dexamethasone (DEX). **(D, E)** Relative *pMIR319C::GUS* domain values in 50-350 μm leaf primordia plotted for the indicated genotypes. Differences among samples are indicated by lower-case letters. $p < 0.05$. N varied from 33-114 (**D**) or 18-43 (**E**). Two-way ANOVA followed by Tukey's post hoc test (**D**), or one-way ANOVA followed by Dunnett's post hoc test (**E**) were performed to determine the significant differences among samples.

Figure 7: Schematic representation of proposed model for CUC2-miR319c mediated upregulation of cell proliferation in leaf primordia. (A) CUC2 (purple wavy lines) expression throughout the incipient leaf primordium (on the right) where it directly activates *MIR319C* expression (green). The mature miR319c ensures cell proliferation by preventing accumulation of *JAW-TCP* transcripts (bold black T-line). As the primordium grows out of the shoot apical meristem (SAM), CUC2 expression gets restricted towards the SAM-leaf boundary (primordium on the left), where it maintains miR319c activity that continues to repress *JAW-TCP*s, thus establishing the basal cell proliferation domain (left panel). At this stage, JAW-TCP proteins accumulate at the distal region (magenta on the left panel), and directly repress *MIR319C* transcription (inverted magenta triangle) (Shankar et al. 2023), thus promoting proliferation-to-differentiation transition in distal cells by distinct mechanisms (Challa et al. 2019). Thus, the *MIR319C* expression domain is established via the antagonistic regulation by JAW-TCP and CUC2 proteins. Image credits: Hand-drawn outline of the shoot apex adapted from Du et al. 2018 (https://doi.org/10.1016/j.molp.2018.06.006) **(B)** In Col-0 (upper panel), high miR319c level (green line) inhibit JAW-TCP (magenta line), ensuring proliferation (green triangle) at the base of early leaf primordia. As primordia develop, miR319c domain declines, whereas JAW-TCP domain increases concurrently, committing proliferating cells to differentiation (magenta triangle). JAW-TCP activity remains high in the growing leaf, promoting increased cell differentiation and expansion-driven growth. As leaves approach maturity, JAW-TCP level decline, and all cells complete differentiation and growth by expansion. In *cuc2;miR319c* leaf primordia (bottom panel), reduced miR319c levels lead to an early rise in JAW-TCP activity, causing a premature transition from proliferation to differentiation and expansion-driven growth. This reduction in proliferation potential decreases the final cell number, ultimately resulting in a smaller mature leaf with fewer cells.