α5-GABA-A Receptor Positive Allosteric Modulation prevents neuronal atrophy and cognitive decline independently of p-Tau accumulation in the PS19 mouse model

Ravinder N. Dharavath[1+], Ashley M. Bernardo[1+], Cassandra Marceau-Linhares[1+], Michael Marcotte[1], Carla Mezo-Gonzalez[1], Kayla Wong[1], Celeste Pina-Leblanc[1], Adrien Bouchet[1], Dishary Sharmin[2], Kamal P. Pandey[2], James M. Cook[2], Thomas D. Prevot[1,3,4*], Etienne Sibille[1,3,4*]

[1]Campbell Family Mental Health Research Institute of CAMH, 250 college street, Toronto, ON, M5T 1R8 Canada

[2]Department of Chemistry and Biochemistry, University of Wisconsin–Milwaukee, 3210 N Cramer Street, 53211, WI, USA

[3]Department of Psychiatry, University of Toronto, 250 college street, Toronto, ON, M5T 1R8 Canada

[4]Department of Pharmacology and Toxicology, University of Toronto, Medical Sciences Building, 1 King's College Circle Room 4207, Toronto, ON, M5S 1A8, Canada

[+]These authors equally contributed to the study

[*]Corresponding Authors:

Etienne Sibille, Ph.D, CAMH, 250 College Street, room 134, Toronto, ON M5T 1R8, Canada

E-mail: Etienne.sibille@camh.ca

Thomas D. Prevot, Ph.D., CAMH, 250 College Street, room 131, Toronto, ON M5T 1R8, Canada

Tel: 416-535-8501, ext 33785; E-mail: thomas.prevot@camh.ca

ABSTRACT

Background: Dysregulated Tau phosphorylation (p-Tau) is a hallmark of neurodegenerative

disorders such as Alzheimer's disease (AD) or frontotemporal dementia (FTD), resulting in

neurofibrillary tangle accumulation, neuronal atrophy and cognitive impairment. Impaired

somatostatin (SST) expression and reduced SST-expressing GABAergic neurons significantly

contributes to AD-related pathophysiology and correlates with cognitive impairment. SST+

interneurons inhibit the dendrites of excitatory neurons in cortical layers and hippocampus,

primarily through α5-GABA-A receptors, regulating cognitive function. Leveraging a newly

developed small molecule that targets the α5-GABA-A receptors via positive allosteric

modulation (α5-PAM), we assessed its effects on p-Tau-related neuronal morphology, cognitive

deficits and protein expression.

Methods: In the PS19 transgenic mouse model, we administered the α5-PAM, GL-II-73, either

acutely or chronically at 3 and 6 months, corresponding to early and advanced stage of p-Tau

accumulation. Golgi staining analyzed dendritic morphology and spine density in mice

chronically exposed to α5-PAM. Western blotting was used to quantify p-tau and Tau

expression. Spatial working memory was assessed using the Y-maze.

Results: Chronic treatment at 3 and 6 months mitigated p-Tau-induced loss of spine density and

reduced dendritic length. α5-PAM treatment did not affect p-tau levels. α5-PAM effectively

reversed spatial working memory deficits induced by p-tau accumulation both acutely and

chronically.

Conclusions: α5-GABA-A receptor positive allosteric modulation displayed neurotrophic (spine

and dendritic pathology) and procognitive (working memory) effects in the PS19 model,

independently of p-Tau burden. This suggests a novel therapeutic strategy for p-Tau-related

pathologies with both symptomatic and disease-modifying potential.

INTRODUCTION

Tauopathies are a group of progressive neurodegenerative disorders marked by the pathological accumulation of hyperphosphorylated tau protein (p-Tau), which aggregates into neurofibrillary tangles (NFTs) (*1*), and lead to neuronal dysfunctions, cognitive deficits, and brain atrophy (*2*). Alzheimer's disease (AD) and frontotemporal dementia (FTD) are amongst the most common tauopathies and share these pathological hallmarks (*3, 4*). The accumulation of p-Tau correlates with the progression of neuronal atrophy and cognitive deficits progression, contributing substantially to disease burden and progression (*5*).

Despite clear understanding of the role of p-Tau aggregates and NFTs in the pathophysiology of AD and other tauopathy-based disorders, pharmacotherapies targeting Tau have had limited success and interest, likely due to the failures in the AD field, when targeting Tau or amyloid β (*6-8*). However, the recent approval of antibody-based therapies targeting the soluble and insoluble amyloid protein (*9*), raised interest in revisiting both amyloid β-targeting and Tau-targeting interventions. Therapies targeting the Tau-pathology include Zagotenemab (a humanized anti-Tau antibody(*10*)), LMTX (TauRx, an aggregation inhibitor (*11*)), TPI-287 (a microtubule stabilizer, (*12*)) and others (*13*), but so far none have demonstrated efficacy at resolving neuronal atrophy or cognitive decline in patients.

Although p-Tau accumulation is a major contributor to AD and FTD pathophysiologies, neurodegenerative disorders are complex and multifactorial diseases that may require a polypharmacopeia approach to reduce the burden of the disease and improve patients' quality of life (*14*). Aβ- and Tau-targeting therapies can help with clearing or limiting accumulation of plaques and tangles, but no studies demonstrated better neuronal function after Aβ clearance or tangle accumulation inhibition following these therapies. Neuronal hyperexcitability is

consistently reported in the hippocampus (*15*) and prefrontal cortex (*16*) (brain regions highly

involved in cognitive functions (*17*)) of AD patients, and proposed to be associated with Tau-

accumulation (*18*). This hyperexcitability is thought to be due in part to a loss of GABAergic

inhibition (*19*). Evidence supports the contribution of GABAergic inhibition in memory (*20, 21*)

and demonstrated that altered GABAergic signaling is linked to poorer cognitive functions in

elderly (*22*), in AD (*23*) and in FTD (*24*) patients. Therefore, targeting the GABAergic system is

emerging as a valid approach for reducing biological changes in AD pathology, potentially

providing disease-modifying effects and symptom relief.

Recently, the α5-GABAA receptors (α5-GABAA-R) have been proposed as relevant targets for

cognitive regulation (*25, 26*), aligned with their discrete distribution in the hippocampus and

prefrontal cortex. α5-GABAA-Rs are localized synaptically and extrasynaptically in distal

dendrites of pyramidal neurons. In patients with AD, α5-GABAA-R binding was reduced by

~30% reduction in the hippocampus (*27*), the associated gene (*GABRA5*) was transcriptional

downregulated in middle temporal gyrus (*28*). An especially promising treatment involves the

exploration of compounds selectively modulating α5-GABAA-R (*26*). Previous studies in our

group showed that treatment with GL-II-73, a novel α5-GABAA-R positive allosteric modulator

(α5-PAM), reverses cognitive deficits in animal models of stress (*29, 30*), aging (*31*), and

amyloid deposition (*32*). In addition, we showed that chronic GL-II-73 treatment reversed the

loss of dendritic lengths, spine counts and spine density in the prefrontal cortex (PFC) and

hippocampus of animals in the same three models of these conditions (*30-32*).

In the present study, we hypothesize that chronic α5-PAM treatment prevents and/or reverses

neuronal deficits during the early, and potentially later stage, of p-Tau accumulation, and that

these effects would be associated with improved cognitive functions. To test this hypothesis, we

investigated the potential of GL-II-73 to mitigate cognitive deficits and neuronal loss in the PS19

mouse model, characterized by the progressive accumulation of p-Tau. The PS19 model

expresses the P301S mutant form of human microtubule-associated protein Tau (MAPT), under

the control of the mouse prion protein promoter (Prnp) (*33*). These mice exhibit increased

accumulation of p-Tau, leading to cognitive decline and neuronal damage (*34*). We conducted

assessments on both male and female mice at 3 and 6 months of age, representing early and later

stages of Tau tangle deposition.

METHODS

Ethical Statement

All animal procedures adhered to the Ontario Animals for Research Act (RSO 1990, Chapter

A.22) and the Canadian Council on Animal Care (CCAC). The study was approved by the

Institutes Animal Care Committees (Protocol #846).

Animals

PS19 breeders, expressing the P301S mutant form of human microtubule-associated protein Tau

(MAPT), were obtained from Jackson Laboratories (Cat #008169-JAX) on a C57BL/6xC3H

background and paired with WT breeders to generate experimental groups' parents. Four cohorts

of PS19 heterozygous (HET) and WT littermates were produced: Acute studies at 3 months (N=8

per group, 3 groups 50% male, **Fig. 1A**) and 6 months (N=8-9 per group, 3 groups, ~50%

female, **Fig. 1B**); Chronic studies at 2-3 months (N=12-13 per group, 3 groups, 50% female, **Fig.**

1C) and 5-6 months (N=12-13 per group, 3 groups, 50% female, **Fig. 1D**). Animals were group-

housed until the study commenced, then transitioned to single housing to facilitate drug

administration and monitoring. They had ad libitum access to food and water and were

maintained on a 12-hour light/dark cycle. Animal handling followed the technique outlined in

Marcotte et al (*35*) to minimize reactivity to the experimenter.

Drug Preparation and Administration

GL-II-73, an α5-GABAA receptor (α5-GABAA-R) positive allosteric modulator (PAM) (*29*),

was synthesized by Dr. Cook's group (University of Wisconsin-Milwaukee). For acute

administration via intraperitoneal (IP) injection, GL-II-73 was dissolved in a vehicle solution

(85% distilled H_2O, 14% propylene glycol (Sigma Aldrich), 1% Tween 80 (Sigma Aldrich)).

Acute studies at 3 months and 6 months utilized GL-II-73 at 5mg/kg and 10mg/kg in the Y maze

via IP injection, administered 30 minutes before testing. For chronic studies, GL-II-73 was

dissolved in tap water and orally administered to mice at 30mg/kg in the drinking water, based

on daily water intake calculations as in (*36*). Drug solution dissolved in drinking water was

prepared fresh every other day and was provided ad libitum for 4 weeks.

Y-Maze

The Y maze, adapted from Faucher et al. (*37*), uses spontaneous alternation to assess spatial

working memory. It consists of three arms—two "goal" arms and one "starting" arm—separated

by 120°, with sliding doors for the goal arms and a start box at the end of the starting arm. Mice

undergo a 2-day habituation phase, exploring the maze for 10 minutes each day. On the third

day, the training phase begins. Mice are placed in the start box, and after a 30-second interval,

the sliding door opens, allowing them to choose a goal arm. After selection, the goal arm's door

closes, and the mouse stays there for 30 seconds before being returned to the start box. This is

repeated for seven trials. On the fourth day, the procedure is repeated with a 60-second interval

between trials, except for the eighth trial, where a 10-second interval tests motivation to

alternate. Mice that do not alternate in the eighth trial are excluded to prevent motivational bias.

Mean % alternation is calculated as [(Number of alternations/number of trials) * 100]. The maze

is cleaned with 70% ethanol between trials to remove olfactory cues.

Golgi Staining and Analysis

After euthanasia and brain extraction, brains from six mice per group (three males, three

females) were placed in Golgi Solution for two weeks, selected pseudo-randomly based on Y

maze performance. They were then coded for blinded analysis and sent to NeuroDigitech (San

Diego, CA, USA). Remaining brains were dissected for PFC and hippocampus extraction for

western blot analysis.

For Golgi-stained brains, 100µm coronal slices were made using a cryostat and mounted on glass

slides. Basal and apical dendrites of pyramidal neurons in layers II/III of the PFC and CA1 were

identified as regions of interest (ROIs). Dendritic features were quantified using NeuroLucida

v10 on a Zeiss Axioplan 2 microscope with an Optronics MicroFire CCD camera. Motorized X,

Y, and Z focus enabled high-resolution imaging. Low magnification identified ROIs, while

higher magnifications (40x-60x) were used for detailed analysis. Immersion oil and a Zeiss 100x

lens facilitated 3D imaging and spine counting. Neurons (n=6 per animal) were selected based on

complete dendritic filling and clear 3D visualization. Neurons with incomplete staining or

visualization were excluded, and spine sampling focused on well-characterized spines orthogonal

to the dendritic shaft. Data were analyzed using NeuroExplorer following (*38*).

Protein measurement via Western Blot

Brain tissue from the chronic 3- and 6-month groups (each consisting of 10 mice, with

approximately 50% female) was prepared for protein analysis. The prefrontal cortex (PFC) and

whole hippocampus regions were sonicated in APL buffer from Qiagen AllPrep RNA/Protein

extraction kit (Qiagen, Germany; Cat # 80404). Proteins were extracted from the brain

homogenate. Protein concentrations were determined using the Bicinchoninic Acid (BCA) assay

(Thermo Fisher Scientific, USA). Bovine serum albumin (BSA) standards were used to generate

a standard curve, and absorbance was measured at 562 nm with a microplate reader.

Equal amounts of protein (20 μg) from each sample, as determined by the BCA assay, were

separated on 10% SDS-PAGE stain-free gels (Bio-Rad Laboratories, #5678094). Following

electrophoresis, the gels were imaged using the ChemiDoc system (Bio-Rad Laboratories, USA)

to verify equal loading and quantify total protein quantification. Proteins were then transferred

onto nitrocellulose membranes using a semi-dry transfer system. Membranes were blocked with

5% BSA in 1X TBST (Tris-buffered saline with 0.1% Tween 20) for 1 hour at room temperature.

Membranes were incubated overnight at 4°C with primary antibodies targeting p-Tau or Tau (see

Supplementary Table 1 for details). After washing, membranes were incubated with HRP-

conjugated secondary antibodies (Vector, #PI-1000 for rabbit and #PI-2000 for mouse) for 1

hour at room temperature. Signal detection was performed using an enhanced

chemiluminescence (ECL) substrate (Thermo Fisher Scientific, USA) and the membrane was

imaged in the ChemiDoc. Quantitative analysis of total protein and band intensity was conducted

using ImageLab software.

Study Design

Heterozygote PS19 mice and their WT littermate were studied at 3 months of age (corresponding

to an early stage of phosphorylation of Tau; **Fig.1AB**) or at 6 months of age (corresponding to a

more advanced stage of phosphorylation of Tau; **Fig.1CD**). At each age, the effect of acute or

chronic (4 weeks) administration of GL-II-73 was investigated on working memory

performances in the Y maze. In the chronic studies, brains were extracted for downstream

analyses including morphology investigation using a Golgi staining technique, or western blot, 7

days after completion of the Y maze.

Figure 1. *Study Design*

Statistics

GraphPad Prism (v10.3.1) was employed for statistical analyses, and the results were graphically

presented to convey mean ± SEM (**Supplementary Tables 2-4**). Morphological Golgi stain

results were analyzed by Neurodigitech through two-way ANOVAs with Tukey's multiple

comparisons test. The Y-maze data underwent analysis via Kruskal-Wallis test, followed by

uncorrected Dunnett's multiple comparisons test. The p-tau/Tau ratio was calculated from the

western blot relative expression of pTau and Tau, with the limitation that pTau levels in WT

were not detectable. Therefore, relative expression was calculated using the HET group as a

reference.

RESULTS

Chronic GL-II-73 treatment prevents neuronal shrinkage and reduced spine density in the

PFC and CA1 of PS19 mice at 3 and 6 months of age.

Morphological changes were quantified on pyramidal neurons located in the prefrontal cortex

(PFC) and the CA1 of the hippocampus (HIP) using a Golgi-Cox staining technique on mice

from the chronic studies at 3 and 6 months.

At 3 months of age, untreated HET mice showed a significant reduction in spine density in the

PFC and CA1, corresponding to 80-81.5% of WT control levels (**Figure 2G,** top panel). This

reduction was partially prevented by chronic administration (1 month) of GL-II-73 in both brain

regions (**Supplementary Table 2**). The spine density of PS19 HET treated with GL-II-73

(HET+GL) were at 92.7% and 92.4% of WT levels in PFC and CA1 respectively, significantly

higher than untreated HET mice, but still lower than WT mice. Similar reductions of spine

density in untreated HET mice were observed at 6 months of age (79.3% and 69.4% of WT

levels in PFC and CA1, respectively), which were in large part prevented by chronic GL-II-73

treatment (95.6% and 93.7% in PFC and CA1 of WT mice, respectively) (**Figure 2H;**

Supplementary Table 3). The reduction in spine density in HET mice and partial reversal with

GL-II-73 treatment was observed in apical and basal segments of the pyramidal neurons

(**Supplementary Figure 1**) in both the PFC and CA1.

Similar results were obtained for the analyses of dendritic length and spine counts (**Figure 1H**

and **G**, middle and lower panels). Significant reductions of dendritic length were observed,

corresponding to 82.6% and 79.3% of WT levels at 3 months and 80.2% and 77.5% of WT

levels at 6 months in the PFC and CA1, respectively, which were reversed or prevented by GL-

II-73 treatment. Dendritic length of mice treated with GL-II-73 were not significantly different

from WT levels at 3 months (92.6% and 95.8% of WT levels in the PFC and CA1), and at 6

months (99.7% and 100% of WT levels in the PFC and CA1, respectively). Similarly, spine

count reductions were observed in untreated HET mice at both age, and in both brain regions

(66.2% and 64.3% of WT levels in the PFC and CA1 at 3 months, and 64.7% and 62.2% of WT

levels in the PFC and CA1 at 6 months), which were prevented or reversed by GL-II-73

treatment (86.2% and 88.5% of WT levels in the PFC and CA1 at 3 months, and 95% and 93.6%

of WT levels in the PFC and CA1 at 6 months).

Figure 2. *Chronic treatment with GL-II-73 reverses the reduced spine density, dendritic length and spine loss in the PFC and CA1*

of PS19 mice at 3 and 6 months of age.

Chronic GL-II-73 does not affect the accumulation of phosphorylated Tau.

The PFC and HIP from 9-10 animals that were not used for Golgi staining were collected to measure Tau and phospho-Tau levels after chronic administration of GL-II-73 at 3 and 6 months of age (**Supplementary Figure 2, and Supplementary Table 4**). There was no group differences in expression of Tau in the PFC or HIP at 3 and 6 months. However, there was a significant increase in pTau levels in HET mice and in GL-II-73-treated HET mice, compared to WT. Taking the native expression of Tau and the expression of pTau in the same samples, a ratio between pTau/Tau was calculated as an index of disease progression (**Figure 3**). HET mice exhibited a significant increase in pTau/Tau ratio (driven by increase in pTau level; **Supplementary Figure 2**) in the PFC and HIP at 3 months and at 6 months. Mice receiving chronic administration of GL-II-73 also exhibited higher ratio of pTau/Tau compared to WT, which were nominally lower in the 6-months of age group but not significantly different from untreated HET mice in both brain regions and at both ages.

Acute and chronic GL-II-73 treatment reverses working memory deficits.

Mice were trained in the alternation task in the Y maze and tested at 3 months and 6 months,

after either acute administration of GL-II-73, 30 minutes prior to testing, or after chronic

administration in the drinking water for 4 weeks. In the groups of mice participating in the acute

studies, PS19 HET mice showed a significant deficit in alternation compared to WT mice at 3

and 6 months (**Figure 4, and Supplementary Table 5**). Acute administration of GL-II-73

displayed a dose-dependent response with significant reversal of alternation deficits at 10 mg/kg

and intermediate results at a lower 5 mg/k dose. Neither of the acute doses had significant effects

at 6 months.

Figure 4. Effects of acute and chronic administration of GL-II-73 on alternation rate in PS19 HET mice

In the chronic administration studies (**Figure 5, and Supplementary Table 5**), alternation rates

in the 3 months-old mice were not significantly different between groups. At 6 months, PS19

HET mice demonstrated a significant deficit in alternation rate deficit compared to WT, which

was reversed by chronic administration of GL-II-73.

DISCUSSION

In this study, we show that chronic administration of GL-II-73, a positive allosteric modulator of the α5-GABAAR subtype (*39*), prevents and reverses the loss of spine density in the prefrontal cortex and hippocampus of the PS19 mice in early and advanced stages of p-Tau accumulation. These effects occurred despite accumulation of p-Tau. In addition, improvement of cognitive functions and reversal of p-Tau-induced deficits, more specifically working memory, were observed, in particular at 3 months when given acutely, and 6 months when given chronically. Importantly, the beneficial effects on neuronal morphology and cognitive functions was independent of Tau-levels, since the chronic treatment did not significantly reduce the p-Tau accumulation.

These findings align with previous findings in other models, demonstrating that targeting the α5-GABAA-Rs with GL-II-73 has neurotrophic effects in the PFC and CA1, as well as cognitive benefits. Indeed, our group demonstrated that chronic administration of GL-II-73 prevents or reverses neuronal atrophy in animal models of chronic stress (*30*), aging (*31*), and amyloid deposition (*32*). Chronic treatment also provided relief of cognitive deficits in these models. This new finding in the context of p-Tau accumulation in mice further suggests that intervention with GL-II-73 in populations with tauopathies would provide neuronal support and would contribute to prevention or reversal of spine density loss and cognitive decline in conditions such as AD or FTD.

Importantly, the PS19 mouse model only expresses the 1N4R isoform, which represents 20-25% of the tauopathies in human, but is not the only isoform present in AD (*40*). A complete model of AD would require including the other isoforms reported to be present in AD (*40*), such as the various 3R isoforms. Interestingly, the MAPT 1N4R isoform expressed in the PS19 mice is

present in other disorders with tauopathy, such as progressive supranuclear palsy (*41*),

corticobasal degeneration (*42*), globular glial tauopathy (*43*) and FTD (*44*), which suggest they

may be of relevance for intervention with α5-PAM for cognitive deficits and neurodegeneration

in these disorders.

Targeting Tau pathologies has been of great interest over the years in the context of AD (*45*), as

studies suggested that Tau shows a stronger correlation with symptom severity than does

amyloid β (*46*). Interventions focusing on Tau protein modification failed in clinical trials over

the years, like Salsalate (*47*), MR-8719, or recently LY3372689 that failed meeting primary

endpoint in Phase 2, BIIB113 which was discontinued or ASN-51 Phase 2 trials halted in March

2025. Tau accumulation inhibitors are also investigated in clinical trials (LMTX also known as

TRx0237) but their efficacy remains to be demonstrated. Immunotherapies are now the most

prominent approaches used to target Tau pathologies in clinical trials (*8, 46, 48, 49*), with

promising avenues (see (*45*) and (*46*) for reviews). Most trials are currently assessing safety and

tolerability of such interventions. Overall, their impact on neuronal health or cognitive functions

remains unclear. But since the Tau pathologies in AD is only one component of the disease, one

can predict that targeting other aspects such as neuronal dysfunction can provide additional

benefits than solely targeting Tau pathologies (*50*).

Targeting GABAergic neurotransmission, particularly through α5-GABAA R, offers a promising

avenue in neurodegenerative disease research, such as AD and FTD. These receptors are

predominantly expressed in hippocampal regions, brain areas critical for memory and cognition

(*26*) and severely affected in AD. Dysregulation of GABAergic signaling in AD has been linked

to cognitive decline, and modulating α5 GABAA-R activity could help restore inhibitory balance

and improve cognitive function (*26*). Tauopathies were shown to generate hyperexcitability (*51*).

GL-II-73 may restore part of the excitation/inhibition balance by increasing GABAergic activity

onto glutamatergic cells, thus stabilizing neuronal activity, altogether helping with cognitive

functions and neuronal exhaustion and atrophy. Moreover, combining this approach with anti-

Tau therapies could provide synergistic effects, as p-Tau accumulation contributes to

neurodegeneration and cognitive impairments. By simultaneously addressing p-Tau

accumulation, GABAergic dysfunction and neuronal loss, combination therapies have the

potential to enhance treatment efficacy and slow disease progression.

Interestingly, the effects observed in this study were independent from p-Tau levels, as chronic

treatment did not alter p-Tau levels in mice, neither at 3 months nor at 6 months. This

observation can be due to the constitutive aspect of the model, where p-Tau is produced

continuously in the PS19 mouse brain. However, despite this continuous accumulation of p-Tau,

treatment with GL-II-73 was able to prevent spine density reduction and cognitive decline,

suggesting strong efficacy despite the presence of the traditional hallmark of AD, similarly to

what was previously demonstrated in a amyloid accumulation model (*32*).

This study has some limitations. As mentioned before, the model used only expresses the 1N4R

isoform of Tau, while there are other isoforms present in AD and in other tauopathy. Therefore,

confirming such effect in other models of tauopathy expressing the other isoforms, or a

combination would further validate the efficacy of GL-II-73. However, in view of similar effects

in models of stress (*29, 30*), aging (*31*), and amyloid deposition (*32*), the likelihood of

generalization effects in other models is high. From a cognitive assessment standpoint, we only

investigated the impact of working memory, when AD and other brain disorders associated with

tauopathy are characterized by more cognitive impairments, such as episodic memory, executive

function, attention and others. Investigating efficacy in these domains would represent an

immediate next step to strengthen the potential of GL-II-73 as a pro-cognitive drug. In addition,

the neurotrophic effect was observed in both regions investigated. Without a negative control, it

is possible that the effect observed with the treatment is generalized throughout the brain, but

similar findings were reported in the context of stress (*30*), aging (*31*), and amyloid deposition

(*32*), suggesting that this effect on spine density is robust.

To conclude, we showed that chronic treatment with GL-II-73 demonstrated robust neurotrophic

effects in the PS19 model of tauopathy in both early (2 months) and advanced (5 months) stages

of p-Tau pathology, together supporting its clinical testing across stages of tauopathy, as stand-

alone or combination therapy, and potentially prophylactically, to delay onset or slow down

disease progression.

FUNDING

This work has been funded by the Weston Brain Institute (TR192043).

ACKNOWLEDGEMENTS

Authors thank Mehrab Ali for support with administrative tasks throughout the study, the CAMH

animal facility staff for the caring for our animals over the study duration, the members of

NeuroDigitech for their contribution to data generation, and all the animals that contributed to

these studies.

DISCLOSURES

JMC, MM, ES and TP are listed inventors on patents covering syntheses and use of the

compound. ES is co-founder of Damona Pharmaceuticals, a biopharma dedicated to bring novel

GABAergic compounds to the clinic. TP provides consulting services to Damona related to

preclinical Research and Development. All other authors declare no conflicts of interest.

References

357   1.   S. Muralidar, S. V. Ambi, S. Sekaran, D. Thirumalai, B. Palaniappan, Role of tau protein in
358   Alzheimer's disease: The prime pathological player. *International Journal of Biological*
359   *Macromolecules* **163**, 1599-1617 (2020).
360   2.   G. G. Kovacs, Tauopathies. *Handb Clin Neurol* **145**, 355-368 (2017).
361   3.   R. Medeiros, D. Baglietto-Vargas, F. M. LaFerla, The role of tau in Alzheimer's disease and
362   related disorders. *CNS Neurosci Ther* **17**, 514-524 (2011).
363   4.   C. M. Moloney, V. J. Lowe, M. E. Murray, Visualization of neurofibrillary tangle maturity in
364   Alzheimer's disease: A clinicopathologic perspective for biomarker research. *Alzheimer's &*
365   *Dementia* **17**, 1554-1574 (2021).
366   5.   A. Bejanin *et al.*, Tau pathology and neurodegeneration contribute to cognitive impairment in
367   Alzheimer's disease. *Brain* **140**, 3286-3300 (2017).
368   6.   Y. Zhang, H. Chen, R. Li, K. Sterling, W. Song, Amyloid β-based therapy for Alzheimer's disease:
369   challenges, successes and future. *Signal Transduction and Targeted Therapy* **8**, 248 (2023).
370   7.   G. P. Morris, I. A. Clark, B. Vissel, Inconsistencies and Controversies Surrounding the Amyloid
371   Hypothesis of Alzheimer's Disease. *Acta Neuropathologica Communications* **2**, 135 (2014).
372   8.   E. E. Congdon, E. M. Sigurdsson, Tau-targeting therapies for Alzheimer disease. *Nat Rev Neurol*
373   **14**, 399-415 (2018).
374   9.   B. Dunn, P. Stein, P. Cavazzoni, Approval of aducanumab for Alzheimer disease—the FDA's
375   perspective. *JAMA internal medicine* **181**, 1276-1278 (2021).
376   10.   B. A. Willis *et al.*, Safety, Tolerability, and Pharmacokinetics of Zagotenemab in Participants with
377   Symptomatic Alzheimer's Disease: A Phase I Clinical Trial. *J Alzheimers Dis Rep* **7**, 1015-1024
378   (2023).
379   11.   D. Seripa *et al.*, Tau-directed approaches for the treatment of Alzheimer's disease: focus on
380   leuco-methylthioninium. *Expert Review of Neurotherapeutics* **16**, 259-277 (2016).
381   12.   S. A. Goldlust *et al.*, Phase 1 trial of TPI 287, a microtubule stabilizing agent, in combination with
382   bevacizumab in adults with recurrent glioblastoma. *Neuro-Oncology Advances* **6**, (2024).
383   13.   S. Jadhav *et al.*, A walk through tau therapeutic strategies. *Acta Neuropathologica*
384   *Communications* **7**, 22 (2019).
385   14.   K. Iqbal, I. Grundke-Iqbal, Alzheimer's disease, a multifactorial disorder seeking multitherapies.
386   *Alzheimers Dement* **6**, 420-424 (2010).
387   15.   S. E. Setti, H. C. Hunsberger, M. N. Reed, Alterations in hippocampal activity and Alzheimer's
388   disease. *Translational issues in psychological science* **3**, 348 (2017).
389   16.   N. Choudhury, L. Chen, L. Al-Harthi, X.-T. Hu, Hyperactivity of medial prefrontal cortex pyramidal
390   neurons occurs in a mouse model of early-stage Alzheimer's disease without β-amyloid
391   accumulation. *Frontiers in Pharmacology* **14**, (2023).
392   17.   E. T. Rolls, G. Deco, C.-C. Huang, J. Feng, The effective connectivity of the human hippocampal
393   memory system. *Cerebral Cortex* **32**, 3706-3725 (2022).
394   18.   W. Huijbers *et al.*, Tau Accumulation in Clinically Normal Older Adults Is Associated with
395   Hippocampal Hyperactivity. *The Journal of Neuroscience* **39**, 548-556 (2019).
396   19.   J. Jiménez-Balado, T. S. Eich, GABAergic dysfunction, neural network hyperactivity and memory
397   impairments in human aging and Alzheimer's disease. *Seminars in Cell & Developmental Biology*
398   **116**, 146-159 (2021).
399   20.   L. Topolnik, S. Tamboli, The role of inhibitory circuits in hippocampal memory processing. *Nature*
400   *Reviews Neuroscience* **23**, 476-492 (2022).

21.   B. Bie, J. Wu, F. Lin, M. Naguib, J. Xu, Suppression of hippocampal GABAergic transmission
impairs memory in rodent models of Alzheimer's disease. *European Journal of Pharmacology*
917, 174771 (2022).
22.   J. Jiménez-Balado *et al.*, Reduced Hippocampal GABA+ Is Associated With Poorer Episodic
Memory in Healthy Older Women: A Pilot Study. *Frontiers in Behavioral Neuroscience* **15**,
(2021).
23.   Y. Li *et al.*, Reversible GABAergic dysfunction involved in hippocampal hyperactivity predicts
early-stage Alzheimer disease in a mouse model. *Alzheimer's Research & Therapy* **13**, 114
(2021).
24.   A. G. Murley *et al.*, GABA and glutamate deficits from frontotemporal lobar degeneration are
associated with disinhibition. *Brain* **143**, 3449-3462 (2020).
25.   T. C. Jacob, Neurobiology and Therapeutic Potential of alpha5-GABA Type A Receptors. *Frontiers*
in molecular neuroscience **12**, 179 (2019).
26.   T. Prevot, E. Sibille, Altered GABA-mediated information processing and cognitive dysfunctions
in depression and other brain disorders. *Mol Psychiatry* **26**, 151-167 (2021).
27.   O. Howell, J. R. Atack, D. Dewar, R. M. McKernan, C. Sur, Density and pharmacology of alpha5
subunit-containing GABA(A) receptors are preserved in hippocampus of Alzheimer's disease
patients. *Neuroscience* **98**, 669-675 (2000).
28.   K. Govindpani, C. Turner, H. J. Waldvogel, R. L. M. Faull, A. Kwakowsky, Impaired Expression of
GABA Signaling Components in the Alzheimer's Disease Middle Temporal Gyrus. *Int J Mol Sci* **21**,
8704 (2020).
29.   T. D. Prevot *et al.*, Novel Benzodiazepine-Like Ligands with Various Anxiolytic, Antidepressant, or
Pro-Cognitive Profiles. *Mol Neuropsychiatry* **5**, 84-97 (2019).
30.   A. Bernardo *et al.*, Symptomatic and neurotrophic effects of GABAA receptor positive allosteric
modulation in a mouse model of chronic stress. *Neuropsychopharmacology* **47**, 1608-1619
(2022).
31.   T. D. Prevot *et al.*, Reversal of Age-Related Neuronal Atrophy by alpha5-GABAA Receptor
Positive Allosteric Modulation. *Cereb Cortex* **31**, 1395-1408 (2021).
32.   A. M. Bernardo *et al.*, Procognitive and neurotrophic benefits of α5-GABA-A receptor positive
allosteric modulation in a β-amyloid deposition mouse model of Alzheimer's disease pathology.
Neurobiology of Aging **147**, 49-59 (2025).
33.   M. Z. Zhong, T. Peng, M. L. Duarte, M. Wang, D. Cai, Updates on mouse models of Alzheimer's
disease. *Molecular Neurodegeneration* **19**, 23 (2024).
34.   S. Zampar, O. Wirths, Characterization of a Mouse Model of Alzheimer's Disease Expressing Aβ4-
42 and Human Mutant Tau. *International Journal of Molecular Sciences* **22**, 5191 (2021).
35.   M. Marcotte *et al.*, Handling Techniques to Reduce Stress in Mice. *J Vis Exp*, (2021).
36.   T. D. Prevot *et al.*, Reversal of Age-Related Neuronal Atrophy by α5-GABAA Receptor Positive
Allosteric Modulation. *Cereb Cortex* **31**, 1395-1408 (2021).
37.   P. Faucher, N. Mons, J. Micheau, C. Louis, D. J. Beracochea, Hippocampal Injections of
Oligomeric Amyloid β-peptide (1-42) Induce Selective Working Memory Deficits and Long-lasting
Alterations of ERK Signaling Pathway. *Front Aging Neurosci* **7**, 245 (2015).
38.   C. C. Wu *et al.*, Selective vulnerability of dentate granule cells prior to amyloid deposition in
PDAPP mice: digital morphometric analyses. *Proc Natl Acad Sci U S A* **101**, 7141-7146 (2004).
39.   T. D. Prevot *et al.*, Residual avoidance: A new, consistent and repeatable readout of chronic
stress-induced conflict anxiety reversible by antidepressant treatment. *Neuropharmacology* **153**,
98-110 (2019).

40.   S. Buchholz, H. Zempel, The six brain-specific TAU isoforms and their role in Alzheimer's disease
and related neurodegenerative dementia syndromes. *Alzheimers Dement* **20**, 3606-3628 (2024).
41.   A. Martínez-Maldonado *et al.*, Molecular Processing of Tau Protein in Progressive Supranuclear
Palsy: Neuronal and Glial Degeneration. *J Alzheimers Dis* **79**, 1517-1531 (2021).
42.   W. Zhang *et al.*, Novel tau filament fold in corticobasal degeneration. *Nature* **580**, 283-287
(2020).
43.   D. C. Chung *et al.*, Tau exhibits unique seeding properties in globular glial tauopathy. *Acta
Neuropathol Commun* **7**, 36 (2019).
44.   L. A. A. Giannini *et al.*, Presymptomatic and early pathological features of MAPT-associated
frontotemporal lobar degeneration. *Acta Neuropathologica Communications* **11**, 126 (2023).
45.   J. L. Cummings *et al.*, The therapeutic landscape of tauopathies: challenges and prospects.
Alzheimer's Research & Therapy **15**, 168 (2023).
46.   E. E. Congdon, C. Ji, A. M. Tetlow, Y. Jiang, E. M. Sigurdsson, Tau-targeting therapies for
Alzheimer disease: current status and future directions. *Nature Reviews Neurology* **19**, 715-736
(2023).
47.   L. VandeVrede *et al.*, Open-Label Phase 1 Futility Studies of Salsalate and Young Plasma in
Progressive Supranuclear Palsy. *Mov Disord Clin Pract* **7**, 440-447 (2020).
48.   A. Boutajangout, J. Ingadottir, P. Davies, E. M. Sigurdsson, Passive immunization targeting
pathological phospho-tau protein in a mouse model reduces functional decline and clears tau
aggregates from the brain. *J Neurochem* **118**, 658-667 (2011).
49.   N. Karimi, F. Bayram Çatak, E. Arslan, A. Saghazadeh, N. Rezaei, Tau immunotherapy in
Alzheimer's disease and progressive supranuclear palsy. *Int Immunopharmacol* **113**, 109445
(2022).
50.   M. M. Martins, P. S. Branco, L. M. Ferreira, Enhancing the Therapeutic Effect in Alzheimer's
Disease Drugs: The role of Polypharmacology and Cholinesterase inhibitors. *ChemistrySelect* **8**,
e202300461 (2023).
51.   J. Brown *et al.*, Tau in cerebrospinal fluid induces neuronal hyperexcitability and alters
hippocampal theta oscillations. *Acta Neuropathologica Communications* **11**, 67 (2023).

FIGURE CAPTIONS

Figure 1. Study Design

Heterozygote PS19 mice (HET) and their wildtype (WT) littermates were included in these studies,

assessing the effect of GL-II-73 on working memory and neuronal morphology. Acute treatment (**A-B**)

was provided to the animals via intraperitoneal administration at 5mg/kg or 10mg/kg at 3 (**A**) or 6 (**B**)

months of age. Chronic administration of GL-II-73 was also investigated at 3 (**C**) and 6 (**D**) months, with

4 weeks of administration in the drinking water at 30 mg/kg. One week after completion of the behavioral

screen, mice were euthanized by rapid cervical dislocation. Brains from 4 mice per group were collected

for Golgi staining, while the remaining 8 per group were collected for Western Blot analyses.

Figure 2. Chronic treatment with GL-II-73 reverses the reduced spine density, dendritic length and

spine loss in the PFC and CA1 of PS19 mice at 3 and 6 months of age.

Four brains per group were collected after chronic administration of GL-II-73 at 3 and 6 months. Six

pyramidal neurons in the PFC (illustrated in **A-F**) and 6 principal cells of the CA1 of the HIP were

isolated per animal and quantified for spine density, dendritic length and spine counts (PFC (**G**) and CA1

(**H**)). Untreated HET mice showed a significant reduction in spine density at 3 months in the PFC and

CA1 which were partially reversed by chronic GL-II-73 (**G**). Similar effects were observed at 6 months

(**H**). ***p<0.001, ****p<0.0001. Mean ±SEM.

Figure 3. pTau/Tau ratio in the PFC and HIP from PS19 mice treated with GL-II-73 compared to

untreated PS19 and WT controls.

Using Western Blot targeting Tau and p-Tau, a p-Tau/Tau ration was calculated. HET mice at 3 months

of age showed a higher ratio compared to WT (as expected) in the PFC (**A**) and HIP (**B**). HET mice

receiving chronic GL-II-73 administration also showed higher ratio compared to WT, not different from

untreated HET mice. At 6 months, the increased ratio in untreated HET compared to WT was confirmed in the PFC (**C**) and HIP (**D**). HET mice receiving GL-II-73 showed ratio not different from WT or HET in the PFC, and significantly higher than WT in the HIP. $**p<0.01$, $***p<0.001$, $****p<0.0001$. Mean ±SEM.

Error! Reference source not found.

Alternation rate in the Y maze were measured after acute administration of GL-II-73 at 3 months (5 and 10 mg/kg dose) (**A**) and 6 months (10 mg/kg) (**B**). At 3 months, HET mice showed a reduction in alternation rate compared to WT. Acute treatment with 10mg/kg dose reversed this deficit, while 5 mg/kg did not significantly change alternation rate. At 6 months, untreated HET mice showed a significantly reduced alternation rate compared to WT. HET mice treated with GL-II-73 did not show significantly improved performances compared to untreated HET mice. Similarly, alternation rate in the Y maze were measured after chronic administration of GL-II-73 at 3 months (**C**) and 6 months (**D**). At 3 months, there was no difference between groups. At 6 months, untreated HET mice showed a significantly reduced alternation rate compared to WT. On the other hand, HET mice treated with GL-II-73 showed significantly higher alternation rate than untreated HET and comparable to WT levels. $*p<0.05$, $**p<0.01$. Mean ±SEM.