Endophilin A1 facilitates organization of the GABAergic postsynaptic machinery

to maintain excitation-inhibition balance

Xue Chen[1,2], Deng Pan[1,2], Jia-Jia Liu[1,2*] and Yanrui Yang[1*]

[1]State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and

Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China

[2]College of Life Sciences, University of Chinese Academy of Sciences, Beijing

100039, China

*Correspondence to Yanrui Yang: yryang@genetics.ac.cn; Jia-Jia Liu:

jjliu@genetics.ac.cn.

Abstract

The assembly and operation of neural circuits in the brain rely on the coordination and

balance of excitatory and inhibitory activities. Inhibitory synapses are key regulators

of the functional balance of neural circuits. However, due to the diversity of inhibitory

presynaptic neurons, the complex composition of postsynaptic receptor subunits and

the lack of typical postsynaptic dense structure, there are relatively few studies on the

regulatory mechanisms for inhibitory synaptic structure and function, and insufficient

understanding of the cellular and molecular abnormalities of inhibitory synapses in

neurological and neuropsychiatric disorders. Here, we report a crucial role for

endophilin A1 in inhibitory synapses. We show that endophilin A1 directly interacts

with the inhibitory postsynaptic scaffold protein gephyrin in excitatory neurons, and

promotes organization of the inhibitory postsynaptic density and synaptic

recruitment/stabilization of the γ-aminobutyric acid type A receptors via its plasma

membrane association and actin polymerization promoting activities. Loss of

endophilin A1 by gene knockout in mouse hippocampal CA1 pyramidal cells weakens

inhibitory synaptic transmission and causes imbalance in the excitatory/inhibitory

function of neural circuits, leading to increased susceptibility to epilepsy. Our findings

identify endophilin A1 as an iPSD component and provide new insights into the

organization and stabilization of inhibitory postsynapses to maintain E/I balance as

well as the pathogenesis of epilepsy.

Key words: endophilin A1, E/I balance, epilepsy, $GABA_AR$, gephyrin, inhibitory

synapse

Introduction

The excitatory and inhibitory synaptic activities of neural circuits are tightly

orchestrated in the central nervous system (Sakimoto et al., 2021). The imbalance

between excitatory and inhibitory neuronal activity is implicated in many neurological

and neuropsychiatric diseases including schizophrenia, autism spectrum disorder and

epilepsy (Canitano and Pallagrosi, 2017; Canitano and Palumbi, 2021; Foss-Feig et al.,

2017; Fritschy, 2008). In the brain, in contrast to asymmetric excitatory synapses

formed between axonal boutons and dendritic postsynaptic structures with thick

disc-shaped PSD, inhibitory synapses are symmetric and are constituted of the

presynaptic terminal from diverse interneurons and the postsynaptic plasma

membrane with thin sheet-like PSD where receptor complexes are anchored (Bai et al.,

2021; Favuzzi et al., 2019; Liu et al., 2020; Mody and Pearce, 2004; Tao et al., 2018).

Although the formation, maturation and plasticity of both excitatory and inhibitory

synapses are vitally important for neural circuit functioning, much more research

efforts have been focused on the regulatory mechanisms governing excitatory

synaptic structure and function. The mechanisms by which inhibitory synapses are

constructed in the brain and fine-tuned in response to neural activity remain largely

unexplored.

The γ-aminobutyric acid type A receptors ($GABA_ARs$), major targets of

antiseizure drugs, are heteropentameric chloride channels that mediate fast synaptic

transmission at inhibitory synapses in the brain (Scott and Aricescu, 2019). Several

neuropathologies are characterized by alterations in $GABA_A$ receptor trafficking,

synaptic E/I ratio and neuronal excitability (Sakimoto et al., 2021). At the inhibitory

postsynaptic site, synaptic localization and functioning of $GABA_ARs$ requires

trans-synaptic cell adhesion molecules, submembranous scaffold proteins and

intracellular signaling proteins. Gephyrin is a master organizer of iPSD, which self-assembles into planar lattices beneath the plasma membrane and anchors GABA$_A$Rs or glycine receptors (GlyRs) to the postsynaptic sites (George et al., 2021; Kasaragod and Schindelin, 2019; Kirsch et al., 1991; Tyagarajan and Fritschy, 2014). Collybistin, a membrane lipid-binding protein, functions in recruitment of gephyrin and GABA$_A$Rs to the plasma membrane by interacting with gephyrin, the synaptic adhesion molecule neuroligin-2 (NL-2) and the α2 subunit of GABA$_A$Rs (Kalscheuer et al., 2009; Kins et al., 2000; Poulopoulos et al., 2009; Saiepour et al., 2010). The transmembrane protein Shisa7 binds GABA$_A$Rs and regulates their trafficking, synaptic abundance and functional properties via binding to protein(s) other than gephyrin (Han et al., 2019; Liu et al., 2020). Nevertheless, mechanisms governing the assembly, recruitment and maintenance of various inhibitory synaptic components remain largely enigmatic.

The majority of inhibitory synapses are formed on the dendritic shaft, or around the cell body of excitatory principal neurons where both actin and microtubule cytoskeletons are present underneath the plasma membrane. Through direct interaction with polymerized tubulin, gephyrin organizes the iPSD by anchoring the GlyRs and GABA$_A$Rs to the microtubule cytoskeleton (Parato and Bartolini, 2021). Notably, although small F-actin patches are observed within the dendritic shaft where the majority of inhibitory synapses are assembled (van Bommel et al., 2019), role(s) of the actin cytoskeleton in the organization and maintenance of inhibitory synapses is still unclear.

Endophilin A1 (EEN1) is a member of the endophilin A protein family characterized by an amino (N)-terminal membrane-binding N-BIN/amphiphysin/Rvs (BAR) domain and a carboxyl (C)-terminal Src homology 3 (SH3) domain. The gene encoding endophilin A1 (*EEN1*, gene name *sh3gl2*) is almost exclusively expressed in the brain (Ringstad et al., 1997), and has been implicated in epilepsy, schizophrenia, depression, Parkinson's disease, and Alzheimer's disease (Chang et al., 2017; Corponi et al., 2019; Germer et al., 2019; Ren et al., 2008; Yu et al., 2018a; Yu et al., 2018b).

In excitatory synapses, endophilin A1 functions in synaptic vesicle recycling and autophagosome formation at the axon terminal (Bademosi et al., 2023; Milosevic et al., 2011; Murdoch et al., 2016; Ringstad et al., 1999; Schuske et al., 2003; Soukup et al., 2016; Verstreken et al., 2003; Watanabe et al., 2018), and regulates the morphogenesis, maturation, and acute structural plasticity of dendritic spines by promoting actin polymerization via its effector protein p140Cap (Yang et al., 2021; Yang et al., 2018; Yang et al., 2015).

Previous studies reported that the double knockout (KO) mice of *EEN1* and *EEN2* exhibit spontaneous epileptic seizures (Milosevic et al., 2011). However, *EEN1* depletion only causes a slight decrease in excitatory synaptic transmission in hippocampal CA1 neurons (Yang et al., 2021; Yang et al., 2018). As E/I imbalance is caused by changes in either excitatory or inhibitory activity, or both, we were prompted to investigate the biological function of endophilin A1 in the inhibitory synapses. In this study, we show that endophilin A1 interacts with gephyrin and localizes to GABAergic postsynapses. Via its membrane-binding and actin polymerization-promoting activities, endophilin A1 facilitates organization of the inhibitory postsynaptic machinery. Loss of endophilin A1 causes impairment of inhibitory synaptic transmission, E/I imbalance and aggravation of epileptic susceptibility.

Results

Ablation of endophilin A1 causes E/I imbalance and increases epilepsy susceptibility

Previous studies found that *EEN1* and *EEN2* double knockout (KO) mice exhibit spontaneous epileptic seizures (Milosevic et al., 2011), suggesting that endophilin A1 and A2 are involved in maintaining the E/I balance. However, another study reported that the protein levels of endophilin A1 were higher in temporal neocortex neurons of temporal lobe epilepsy (TLE) patients (Yu et al., 2018b). Moreover, in a chemical kindling mouse model of epilepsy, in which mice are chronically induced with the

GABA$_A$R antagonist pentylenetetrazole (PTZ) (Dhir, 2012), endophilin A1 was also

upregulated in the hippocampus and adjacent temporal cortex , and silencing its

expression in the hippocampal CA3 region reduced PTZ-induced seizure

susceptibility and severity (Yu et al., 2018b). To investigate the role of endophilin A1

in epilepsy, we determined the impact of *EEN1* gene ablation on chemical kindling of

pan-neural *EEN1* KO mice (Yang et al., 2018) with PTZ at a sub-convulsive dose of

25 mg/kg. Compared with the wild-type, PTZ-induced epileptic behaviors were

significantly exacerbated in *EEN1* KO mice (Figure 1A-C and Video S1).

We previously found that *EEN1* KO caused a slight decrease in the amplitude of

evoked excitatory postsynaptic currents (eEPSC) and a significant attenuation of

long-term potentiation (Yang et al., 2018). As imbalance between excitatory and

inhibitory signals in neuronal circuits is implicated in epilepsy, we then determined

whether loss of endophilin A1 in excitatory neurons affects the E/I ratio by

electrophysiological analysis. To this end, we eliminated the *EEN1* gene in pyramidal

neurons by stereotaxic injection of adeno-associated virus (AAV) expressing

CaMKIIα promoter-driven Cre recombinase into the hippocampal CA1 region of

EEN1$^{fl/fl}$ mice (Figure 1D). Whole-cell patch clamp recording of CA1 pyramidal cells

indicated that, while the eEPSC and its paired-pulse ratio were not significantly

changed, the evoked inhibitory postsynaptic current (eIPSC), not its PPR, was

decreased in the *EEN1* KO neurons, resulting in a significant increase in the E/I ratio

(Figure 1E-I). Of note, neither the frequency nor the amplitude of miniature EPSCs

(mEPSC) or miniature IPSCs (mIPSC) was altered in *EEN1* KO neurons (Figure

1J-O). Moreover, neither fast mIPSCs (perisomatic events: < 2.8 ms) nor slow

mIPSCs (distal dendritic events: > 2.8 ms) (Miles et al., 1996) were affected by *EEN1*

KO (Figure S1), indicating that presynaptic quantal release of soma- or

dendrite-targeting inhibitory synapses is normal. Taken together, these data indicate

that endophilin A1 in hippocampal pyramidal neurons functions in inhibitory synaptic

transmission to maintain E/I balance.

Endophilin A1 is required for the formation and stabilization of inhibitory synapses

Diminished eIPSC in *EEN1*^{-/-} pyramidal neurons prompted us to determine the role of endophilin A1 in inhibitory synapses by immunofluorescence staining of brain sections from *EEN1*^{-/-} mice. Analysis of confocal microscopy images revealed that, compared with the wild-type, there was a significant decrease in fluorescent signals of both VGAT and gephyrin, the inhibitory pre- and post-synaptic markers, and PSD95, the excitatory postsynaptic marker, in all subregions of the hippocampus of the *EEN1*^{-/-} mice on postnatal day 21 (P21) (Figure 2A,C,D,F, and S2A,S2C), suggesting defects in synapse development. The decreases in inhibitory but not excitatory postsynaptic protein were also detected in the *EEN1*^{-/-} mice at P60 (Figure 2B,C,E,F, and S2B,S2C), indicating that the assembly and/or stabilization of inhibitory postsynapses was impaired throughout brain development. Notably, gephyrin signals in all layers of the hippocampal subregions were reduced in the *EEN1*^{-/-} mice (Figure S2D and S2E). As different types of interneurons synapse with hippocampal excitatory neurons at distinct subcellular compartments (soma, axon initial segment, proximal and distal dendrites) indicated by the layered structures of the hippocampus (Spruston, 2008), these data suggest that endophilin A1 functions in the assembly/stabilization of inhibitory synapses formed between various presynaptic neurons and postsynaptic neurons.

Next we analyzed the GABA$_A$ receptors and found that, consistent with changes in inhibitory postsynaptic marker, signals of the GABA$_A$R γ2 subunit were decreased in the hippocampi of both juvenile and adult *EEN1*^{-/-} mice (Figure 2G-I). These data collectively indicate that endophilin A1 orchestrates the assembly of inhibitory synapses, the architecture of inhibitory postsynapses in particular, and functioning of GABAergic synapses.

Endophilin A1 localizes to the iPSD and augments the organization of the inhibitory postsynaptic machinery

To substantiate the role of postsynaptic endophilin A1 in the assembly of inhibitory

synapses and exclude the impacts of presynaptic endophilin A1 and its depletion on

early embryonic development, we determined the effects of endophilin A1KO on

inhibitory and excitatory synapses in cultured hippocampal pyramidal neurons during

late developmental stages. As the expression levels of GABA$_A$R α1 are highly

heterogeneous in individual cultured neurons (Figure S3), we determined the effect of

Cre-mediated *EEN1* KO on surface expression of GABA$_A$R γ2 in *EEN1*$^{fl/fl}$

hippocampal neurons by immunofluorescence staining. Whilst no changes in surface

GluA1 was detected, the fluorescence signal intensity and number of surface

GABA$_A$R γ2 clusters were largely lower in dendrites of *EEN1* KO neurons, which

was rescued by re-expression of endophilin A1 (Figure 3A-D). We further examined

the total and surface levels of synaptic proteins by immunoblotting. Biotinylation

assay of surface proteins revealed that, compared with wild-type, although there was

no change in total levels of synaptic proteins in *EEN1* KO pyramidal neurons, the

surface expression of GABA$_A$Rs but not NL2 or the NR1 subunit of the NMDA-type

glutamate receptor was significantly reduced (Figure 3E,F). Collectively, these data

indicate that postsynaptic endophilin A1 enhances the surface clustering of GABA$_A$Rs

and facilitates the assembly of inhibitory postsynapses in pyramidal neurons.

Next we sought to determine the molecular function of endophilin A1 in

inhibitory synapses at the cellular level. By immunofluorescence staining and

super-resolution microscopy of hippocampal neurons in dissociated culture, we

observed partial colocalization of endophilin A1 with gephyrin as well as surface

GABA$_A$R γ2 in dendrites of hippocampal neurons, and juxtaposition or partial

colocalization of endophilin A1 and VGAT puncta (Figure 3G-I and Video S2-4),

indicating that endophilin A1 localizes to not only ePSD but also iPSD. Therefore, we

speculated that endophilin A1 acted as a component of iPSD. Previous studies

reported that synaptic GABA$_A$Rs contain the γ2 subunit along with α and β subunits

(e.g., α1β2γ2) (Scott and Aricescu, 2019). The γ2 subunit is essential for receptor

clustering at the synapse, mediated by interactions with scaffold proteins (Essrich et

al., 1998). Indeed, co-immunoprecipitation (coIP) of endogenous proteins in membrane fractions from mouse brain lysates with antibodies against GABA$_A$R α1 detected not only the known iPSD components γ2, NL2 and gephyrin, but also endophilin A1, and vice versa (Figure 3J). Of note, loss of endophilin A1 caused decreases in levels of GABA$_A$R γ2 and gephyrin but not NL2 in anti-GABA$_A$R α1 immunoprecipitates (Figure 3K). Further, immunoblotting of synaptosome fractions revealed that the levels of GABA$_A$R γ2 and gephyrin were reduced at synapses in adult *EEN1* KO mice (Figure 3L). Together, these data support that endophilin A1 functions at the iPSD to promote the assembly and/or stabilization of the GABAergic postsynaptic machinery.

Endophilin A1 directly interacts with the scaffold protein gephyrin and promotes its clustering and localization of GABA$_A$Rs to iPSD

We subsequently sought to investigate the mechanistic role of endophilin A1 in the clustering of inhibitory postsynaptic proteins. To this end, we performed coIP from mouse brain lysates with antibodies against endophilin A1 (Figure 4A) and analyzed the immunoprecipitates by mass spectrometry. Gene Ontology (GO) analysis revealed that, in addition to proteins participating in known cellular processes involving endophilin A1, such as synaptic vesicle recycling, actin cytoskeleton remodeling, endocytosis and long-term potentiation (Gad et al., 2000; Ringstad et al., 1999; Schuske et al., 2003; Verstreken et al., 2003; Yang et al., 2021; Yang et al., 2018; Yang et al., 2015), proteins related to not only glutamatergic but also dopaminergic and GABAergic synapses were also identified, including gephyrin (Figure 4B and Table S1).

Gephyrin is the major scaffold protein which interacts with the intracellular domains of GABA$_A$Rs and anchors them at the inhibitory postsynaptic membrane (Essrich et al., 1998; Kneussel et al., 1999; Mukherjee et al., 2011; Tretter et al., 2008). The cell adhesion molecule NL2 drives postsynaptic assembly at perisomatic inhibitory synapses via interaction with gephyrin and the membrane lipid-binding

protein collybistin (Poulopoulos et al., 2009). We speculated that endophilin A1

functions in the assembly of iPSD via interaction with the scaffold and/or intercellular

adhesion proteins. To determine which inhibitory synaptic protein(s) interacts with

endophilin A1, we performed coIP from HEK293T cells transiently expressing

endophilin A1 and NL2, or gephyrin, and detected interaction of endophilin A1with

gephyrin but not NL2 (Figure 4C and 4D). Moreover, coIP from mouse brain lysates

corroborated interaction between endophilin A1 and gephyrin but not $GABA_AR$ $\alpha1$ or

$\gamma2$ (Figure 4E). Further, neither endophilin A2 nor A3 interacted with gephyrin

(Figure 4F).

Notably, although the expression levels of gephyrin were not affected by ablation

of *EEN1*, its distribution in the PSD fraction was decreased (Figure 3L). Consistently,

immunofluorescence staining revealed a decrease in the clustering of gephyrin across

the soma, the axon initial segment (AIS), and dendrites (Figure S4). The gephyrin

clustering phenotype was rescued by expression of endophilin A1 but not endophilin

A2 or A3 (Figure 4G,H), suggesting that their functions at the inhibitory postsynapses

are not redundant. Conversely, overexpression of endophilin A1 caused an increase in

the mean signal intensity and number of gephyrin puncta in the soma and dendrites

(Figure 4I,J). Collectively, these data demonstrate that endophilin A1 interacts with

the scaffold protein gephyrin and facilitates its clustering at inhibitory postsynapses.

At the submembraneous postsynaptic density, gephyrin self assembles via the

N-terminal G and C-terminal E domains to provide scaffold for inhibitory

postsynapses and anchors $GABA_ARs$ to the postsynaptic sites. Consistent with the

effect of *EEN1* KO on the clustering of gephyrin (Figure 4G,H) and $GABA_AR$ $\gamma2$

(Figure 3A-D), immunofluorescence staining and confocal microscopy analysis

showed that, depletion of endophilin A1 also caused a decrease in the localization of

$GABA_AR$ $\gamma2$ to gephyrin puncta in *EEN1* KO neurons (Figure 5A-C). Moreover,

overexpression of endophilin A1 also enhanced clustering of $GABA_AR$ $\gamma2$ in the soma

and dendrites (Figure 5D,E). Together these data indicate that endophilin A1 promotes

anchoring of $GABA_ARs$ to the iPSD.

To further understand the mechanism by which endophilin A1 regulates the

assembly and function of inhibitory postsynapses, next we attempted domain mapping

of the endophilin A1-gephyrin interaction. To this end, we generated a series of

truncation and internal deletion mutants for endophilin A1 (Figure 5F). For those

which did not express in mammalian cells, we determined their interaction with

gephyrin by in vitro binding assays with recombinant proteins. None of the truncation

or internal deletion mutants lost the ability to bind to gephyrin, and deletion of neither

the BAR nor the SH3 domain weakened the binding (Figure 5G,H), suggesting that

they are involved in the interaction with gephyrin. Indeed, the decreases in clustering

of either gephyrin or GABA$_A$R γ2 in *EEN1* KO neurons were not rescued by

expression of endophilin A1 fragments (Figure 5I-L), indicating that all functional

domains of endophilin A1 are required for the assembly of the GABAergic

postsynaptic machinery.

Actin polymerization is required for stabilization of inhibitory postsynapses

Endophilin A1 functions in the morphogenesis and structural plasticity of dendritic

spines by promoting actin polymerization underneath the plasma membrane (Yang et

al., 2021; Yang et al., 2018; Yang et al., 2015). Of note, previous studies reported

controversial results about the role of the actin and microtubule cytoskeleton in

gephyrin clustering (Allison et al., 2000; Charrier et al., 2006; Kirsch and Betz, 1995).

To further investigate the mechanism underlying endophilin A1-mediated assembly of

inhibitory postsynapses, we asked whether actin polymerization contributes to the

stabilization of iPSD and GABA$_A$R clustering in dendrites. Consistent with previous

studies (van Bommel et al., 2019), we observed large patches of F-actin labeled by

LifeAct-mCherry in spines and small puncta in dendritic shafts that exhibited partial

colocalization with or adjoined gephyrin in cultured hippocampal neurons by

super-resolution microscopy (Figure S5A and Video S5). We then determined the

impact of F-actin depolymerization on gephyrin-labeled iPSD and γ2-containing

GABA$_A$Rs in cultured hippocampal neurons. One-hour or two-hour treatment of

mature neurons with nocodazole, a microtubule depolymerizing reagent, caused

decreases in the number of both PSD95 and gephyrin puncta (Figure 6A-D), which

was in good agreement with previous findings that microtubule dynamics regulates

the spatial organization of both excitatory and inhibitory postsynaptic structures (Gu

et al., 2008; Jaworski et al., 2009; Kirsch and Betz, 1995; Kirsch et al., 1991).

Short-term or long-term incubation with latrunculin A, which disrupts actin

polymerization, also caused decreases in the number of both ePSD and iPSD puncta

(Figure 6A-D), indicating that actin polymerization also contributes to the

stabilization of the iPSD. We then tested the effect of actin depolymerization on

$GABA_AR$ clustering and found that, long-term but not short-term latrunculin A

application caused a reduction in the number of $GABA_AR$ γ2 puncta in dendrites

(Figure 6E-H), this finding supported the observation that the decrease in gephyrin

cluster preceded the reduction in $GABA_AR$ γ2 receptor cluster following latrunculin A

treatment (Figure 6I). In addition, colocalization analysis revealed less recruitment of

γ2 to gephyrin puncta even after short-term drug application, without changes in γ2

clustering (Figure 6J,K). Together, these data demonstrate that actin polymerization is

essential for the stabilization of iPSD and anchoring of $GABA_ARs$ to the postsynaptic

sites.

Both the membrane-binding and p140Cap-binding capacities of endophilin A1

are indispensable for its function at the inhibitory postsynapses

At the excitatory synapse, the plasma membrane-associated endophilin A1 regulates

the morphogenesis, maturation and structural plasticity of dendritic spines by

promoting branched actin polymerization underneath the postsynaptic membrane via

its interaction with the cytoskeleton regulator p140Cap (Yang et al., 2021; Yang et al.,

2018; Yang et al., 2015). Having established the requirement for both endophilin A1

and the actin cytoskeleton in the assembly and stabilization of the inhibitory

postsynaptic structure, we speculated that regulation of inhibitory postsynapse

assembly by endophilin A1 necessitated its plasma membrane-binding and actin

polymerization-promoting activities. Silencing p140Cap expression in postsynaptic

CA1 pyramidal cells by shRNA-mediated knockdown caused decreases in the

clustering of gephyrin puncta in dendrites, which was restored by overexpression of

p140Cap (Figure S5B,S5C), suggesting that p140Cap-mediated actin polymerization

is required for the assembly of iPSD in postsynaptic neurons. However, compared

with mere depletion of endophilin A1, p140Cap knockdown did not further attenuate

the clustering of gephyrin in *EEN1* KO neurons (Figure S5D,S5E). Moreover,

overexpression of p140Cap did not rescue the iPSD phenotype in *EEN1* KO neurons

(Figure 7A,B), suggesting that p140Cap-enhanced actin polymerization promotes the

organization of iPSD through endophilin A1-mediated pathway. Indeed,

overexpression of neither the membrane-binding deficient (KKK-EEE) nor the

p140Cap-binding deficient (Y343A) mutant of endophilin A1 (Yang et al., 2021)

could rescue the iPSD defect (Figure 7A,B). Consistently, neither p140Cap nor the

endophilin A1 loss-of-function mutants could rescue the GABA$_A$R γ2 clustering

phenotype in *EEN1* KO neurons (Figure 7C,D). Collectively, these data indicate that

both plasma membrane association and actin polymerization-promoting activities are

required for endophilin A1 to promote the assembly of the inhibitory postsynaptic

machinery.

Next, we determined the mechanistic role of endophilin A1 in vivo by rescuing

the epilepsy susceptibility caused by *EEN1* KO. Indeed, when AAVs expressing the

Cre recombinase and endophilin A1 were co-injected into the hippocampal CA1

region of *EEN1*$^{\text{fl/fl}}$ mice, only the wild-type but not the membrane-binding or

p140Cap-binding deficient mutant of endophilin A1 rescued the early onset and

severity of PTZ-induced seizures (Figure 7E,F and Video S6). Consistently,

electrophysiological recording in acute brain slices indicated that only wild-type

endophilin A1 but not the mutants restored the amplitude of eIPSC and E/I balance in

EEN1 KO neurons (Figure 7G-K). All together, these data indicate that endophilin A1

contributes to the structure and function of inhibitory synapses by promoting the

submembrane assembly of iPSD and synaptic localization of the GABA$_A$Rs, thereby

maintaining the balance of excitatory and inhibitory activities in neural circuits

(Figure 8).

Discussion

Although collybistin is implicated in plasma membrane recruitment and clustering of

gephyrin and GABA$_A$R α2 (Saiepour et al., 2010), studies of collybistin KO mice

showed that the GABA$_A$R α3 subunit promotes gephyrin-mediated formation of

inhibitory synapses in the absence of collybistin (Wagner et al., 2021), raising the

possibility that formation of a subset of GABAergic synapses requires

gephyrin-interacting membrane-associated proteins other than collybistin. In the

present study, we identify endophilin A1 as a novel binding partner for gephyrin and

demonstrate that it promotes the assembly of inhibitory synapses and clustering of

GABA$_A$Rs at the iPSD. Loss of endophilin A1 activity causes impairment of

inhibitory synaptic transmission in neuronal circuits and increases susceptibility to

seizures. In the central nervous system, inhibitory interneurons are highly diverse in

morphologies, molecular profiles and physiological functions. Presynaptic terminals

from different types of interneurons target distinct subcellular compartments in

postsynaptic pyramidal neurons where GABA$_A$ receptors of specific subunit

compositions are located. Parvalbumin-expressing (PV+) interneurons usually contact

the soma and proximal dendrites, whereas somatostatin-expressing (SST+)

interneurons mainly innervate the dendrites of pyramidal cells (Klausberger and

Somogyi, 2008). Although different inhibitory synapses localized in different

subcellular compartments of postsynaptic neurons have distinct properties and express

a diversity of synaptic proteins, gephyrin remains a core postsynaptic component

(Favuzzi et al., 2019). Our study revealed that endophilin A1 promotes the clustering

of gephyrin in all subcellular compartments of pyramidal neurons, indicating that

endophilin A1 functions in the organization of inhibitory postsynapses innervated by

various presynaptic interneurons.

Among the endophilin A proteins, endophilin A1 is the most abundant in the

brain (Milosevic et al., 2011). Although they have redundant roles in physiological or

pathological functions (Milosevic et al., 2011), endophilin A1, but not A2 or A3,

promotes spine morphogenesis and excitatory postsynaptic organization (Yang et al.,

2018; Yang et al., 2015; Milosevic et al., 2011). Here, we uncover a previously

unanticipated role for endophilin A1 at GABAergic synapses. By deleting endophilin

As in specific neuron subtypes, we found that endophilin A1 regulates GABAergic

postsynaptic compartments. Endophilin A1, rather than A2 or A3, interacts with

gephyrin and modulates the inhibitory postsynaptic machinery. Although endophilin

A1 might function redundantly with other endophilin As in synaptic vesicle recycling

at the presynaptic site, it participates in the organization or reorganization of

postsynaptic structures through specific binding to postsynaptic proteins, a function

that cannot be fully replaced by other endophilin As at both excitatory and inhibitory

postsynapses.

The membrane lipid-binding protein collybistin interacts with the trans-synaptic

cell adhesion molecule NL-2 to recruit gephyrin and $GABA_AR$ α2 to the postsynaptic

membrane (Kalscheuer et al., 2009; Kins et al., 2000; Poulopoulos et al., 2009;

Saiepour et al., 2010). Adamtsl3 secreted from presynapses binds to the

transmembrane protein DCC (Deleted in Colorectal Cancer) and controls its synaptic

localization, which is required for the maintenance of GABAergic synapses in the

adult brain (Cramer et al., 2023). Anchoring of the α5 subunit of $GABA_AR$ to the

actin cytoskeleton through radixin, an actin-binding protein, facilitates its clustering

(Loebrich et al., 2006). These findings underscore the necessity for investigation of

mechanism(s) regulating the plasma membrane-iPSD-actin cytoskeleton interaction.

As membrane-binding and actin polymerization-promoting activities of endophilin A1

are both required for its function in enhancing iPSD formation and γ2–containing

$GABA_AR$ clustering to iPSD, we propose that membrane-bound endophilin A1

promotes postsynaptic assembly by coordinating the plasma membrane tethering of

the postsynaptic protein complex and its stabilization with the actin cytomatrix

(Figure 8). In contrast, clustering of β2/3 subunit-containing $GABA_ARs$ is

independent of the actin or microtubule cytoskeleton in hippocampal neurons (Allison

et al., 2000; Kasaragod et al., 2022). Given that synaptic and extrasynaptic GABA$_A$Rs

are composed of distinct subunits (Kasaragod et al., 2022; Koh et al., 2023), these

findings indicate that the actin cytoskeleton plays distinct roles in the clustering and

possibly dynamic changes in the subunit composition and subcompartmental

localization of different types of GABA$_A$Rs. Of note, the subcellular distribution of

gephyrin, which has a tubulin-binding motif, is not affected by depolymerization of

F-actin or microtubule (Allison et al., 2000). Overall, the discrepancy in data about

the involvement of the actin cytoskeleton in inhibitory synapses may be due to

differences in drug concentration, treatment time, receptor subtypes and neuronal cell

types (Mody and Pearce, 2004; Sexton et al., 2021).

The number of synaptic GABA$_A$ receptors is subject to alterations not only

during development and neuronal plasticity but also during pathological processes

such as epilepsy (Nusser et al., 1998). Synaptic GABA$_A$ receptors are recruited

directly from extrasynaptic pools following exocytosis, and their endocytosis occurs

exclusively at extrasynaptic sites (Bogdanov et al., 2006). Therefore, lateral diffusion

of GABA$_A$Rs between synaptic and extrasynaptic sites and their iPSD anchoring are

very important for their synaptic distribution (Thomas et al., 2005). In good

agreement with the finding that GABA$_A$Rs stabilization at synapses depends on

gephyrin clustering (Jacob et al., 2005), we found that decreases in gephyrin

clustering precedes the reduction in surface GABA$_A$Rs following F-actin

depolymerization (Figure 6). As the core scaffold protein in iPSD, gephyrin is highly

enriched at the inhibitory postsynaptic sites and drives the formation of iPSD sheets

via phase separation (Bai et al., 2021). As binding of gephyrin by its partner proteins

inhibits the iPSD condensate formation via autoinhibition of the multivalent

interactions between its E domain and GABA$_A$Rs (Bai et al., 2021; Hoffmann and

Milovanovic, 2021; Liu et al., 2020), it is conceivable that endophilin A1 binding not

only tethers the scaffold protein to the plasma membrane, but also releases it from the

autoinhibitory conformation to facilitate the phase separation of the gephyrin-receptor

complexes. Whether the activity of endophilin A1 is regulated by neuronal activity to
participate in the structural and functional remodeling of iPSD awaits further
investigation.

At the presynaptic site of excitatory synapses, endophilin A1 functions in
synaptic vesicle recycling by recruiting the clathrin-uncoating factor synaptojanin
(Milosevic et al., 2011; Schuske et al., 2003; Verstreken et al., 2003). At the excitatory
postsynapse, plasma membrane-associated endophilin A1 facilitates the
morphogenesis, stabilization, maturation and plasticity of dendritic spines by
promoting actin polymerization via its interaction with p140Cap (Yang et al., 2021;
Yang et al., 2018; Yang et al., 2015). In this study, we demonstrate that endophilin A1
functions upstream of p140Cap in not only excitatory but also inhibitory postsynapses
of excitatory neurons. Intriguingly, recent studies reported that p140Cap is present in
the inhibitory presynaptic compartment and regulates GABAergic synaptogenesis and
neurotransmission (Russo et al., 2019), raising the possibility that endophilin A1 and
p140Cap also coordinate to regulate the organization and activity of GABAergic
synapses at the inhibitory presynaptic site. Remodeling of inhibitory synapses and
dendritic spines are spatially clustered by sensory inputs in the central nervous system
(Chen et al., 2012). Notably, the turnover of inhibitory synapses outpace that of
excitatory synapses (Chen et al., 2012). Therefore, it is conceivable that deficits in
inhibitory synapses are much more severe in *EEN1* KO mice, leading to perturbation
of the E/I balance and increased epileptic vulnerability.

Materials and Methods
Ethics statement
All animal experiments were approved by and performed in accordance with the
guidelines of the Animal Care and Use Committee of Institute of Genetics and
Developmental Biology, Chinese Academy of Sciences (Approval code: AP2022008).
All animals were housed in standard mouse cages at 22-24°C on a 12 h light/dark
cycle with access to food and water freely.

Animals

Generation of *EEN1*^{fl/fl} and *EEN1*^{-/-} (KO first) mice on the C57BL/6J background were as previously described. Genotyping of mouse lines was performed by genomic PCR of tail prep DNA from offspring with the following primer pairs: loxPF/loxPR: 5'-CAAGGACTCCCAGAGACCTAGCATC-3' and 5'-GAGATGGCGCAACGCAATTAAT-3' (A PCR product of 375 base pairs in *EEN1* KO first mice but none in wild-type mice). zptF/zptR: 5'-gtaagcggctctagcgcatgttct-3' and 5'-GCAGGGGCATGTAGGTGGCTCAAC-3' (A PCR product of 466 base pairs in wild-type mice, none in *EEN1*^{-/-} mice, and of 627 base pairs in *EEN1*^{fl/fl} mice).

Constructs and viruses

pCMV-Tag2B-EEN1, pCMV-Tag2B-p140Cap, pCMV-Tag2B-EEN1 Y343A, pCMV-Tag2B-EEN1 KKK-EEE, pCMV-Tag2B-EEN1 (aa172-352), pGEX-4T-1-SH3 (aa295-352) and LifeAct-mCherry constructs were described previously (Yang et al., 2021; Yang et al., 2015). pAOV-CAMKIIα-mCherry-2A-3FLAG were purchased from OBiO Technology (Shanghai) Corp. Ltd., (Shanghai, China). pNICE-NL2(-) (Plasmid #15246) and pCR3-FLAG-gephyrin P1 (Plasmid #68816) were purchased from Addgene (Chih et al., 2006; Tyagarajan et al., 2013). The pCMV-Tag3B-GPN construct was generated by cloning gephyrin cDNA amplified from pCR3-FLAG-gephyrin into pCMV-Tag3B (Clontech Laboratories, Inc.). The pGEX-4T-1 EEN1 ΔBAR (Δaa6-242) and pGEX-4T-1 EEN1 ΔSH3 (Δaa1-352) constructs were generated by cloning fragments amplified from pCMV-Tag2B-EEN1 into pGEX-4T-1. All constructs were verified by DNA sequencing. Viral particles of adeno-associated virus (AAV) carrying mCherry, EGFP, the Cre recombinase, EEN1 WT or mutants were purchased from OBiO Technology (Shanghai) Corp. Ltd., (Shanghai, China).

Antibodies

The following antibodies were obtained from commercial sources: rabbit anti-endophilin A1 (159 002, Synaptic Systems GmbH, Germany), mouse anti-Gephyrin (147 111, Synaptic Systems), guinea pig anti-gephyrin (147 318, Synaptic Systems), mouse anti-neuroligin 2 (129 511, Synaptic Systems), mouse anti-GluA1 (MAB2263, Millipore), rabbit anti-GABA$_A$R α1 (06-868, Millipore Sigma), rabbit anti-GABA$_A$R γ2 (224 003, Synaptic Systems), mouse anti-SYP (D-4) (sc-17750, Santa Cruz), rabbit anti-FLAG (F3165, Sigma-Aldrich), rabbit and mouse anti-RFP which recognizes DsRed and mCherry (PM005 and M165-3, Medical & Biological Laboratories, Naka-ku Nagoya, Japan); mouse anti-α-tubulin (T9026, Sigma-Aldrich), mouse anti-PSD95 (610495, BD Biosciences, San Diego, CA) and mouse anti-GST (M071-3, MBL) for western blotting (WB) and rabbit anti-VGAT (131 004, Synaptic Systems), mouse anti-PSD95 (75-028, NeuroMab, Davis, CA), guinea pig anti-Ankyrin G (386 004, Synaptic Systems), guinea pig anti GABA$_A$R γ2 (224 004, Synaptic Systems) for immunofluorescence staining; Rabbit anti-p140Cap was described previously (Yang et al., 2015). Alexa Fluor dye-conjugated secondary antibodies for immunofluorescence staining were from Molecular Probes (Invitrogen, Carlsbad, CA, USA).

Pentylenetetrazole (PTZ)-kindling and seizure behavior scoring

To evaluate seizure susceptibility, 8-10-week-old male C57BL/6N mice were intraperitoneally administered with PTZ (25 mg/kg; K0250, Sigma-Aldrich) or saline (control) for 8-12 days, and the resulting seizure behaviors were video-recorded for the next 30 min. Seizure susceptibility was measured by rating seizures on a scale of 0 to 7 as follows (Van Erum et al., 2019): no abnormal behavior (0), reduced motility and prostate position (1), facial jerking (Schulz et al.), neck ierks (Huo et al.), clonic seizure (sitting) (4), clonic, tonic-clonic seizure (lving on belly) (5), clonic, tonic-clonic seizure (lving on side) and wild jumping (6), tonic extension, possibly leading to respiratory arrest and death (7).

Stereotaxic injection

For virus injection, 9-10-week-old naive male and female mice were anesthetized and stereotactically injected with viral particles in the hippocampal CA1 region as described (Yang et al., 2018). Briefly, mice were weighed and anesthetized with isoflurane (2% 2,2,2-tribromoethanol, Sigma), and secured in a stereotaxic apparatus. The scalp of mouse was cut along the midline between the ears after sterilized with iodophors and 75% (vol/vol) alcohol. Holes for inserting needle were drilled in the bilateral skull. Viral particles carrying pAOV-CAMKIIa-MCS-mCherry-3FLAG or pAOV-CAMKIIa-mCherry-2A-Cre and pAAV-CaMKIIα-EGFP-2A-MCS-3FLAG or pAAV-CaMKIIα-EGFP-2A-MCS-3FLAG-EEN1 WT or mutants (1.2 μL, 2.0×10^{12} viral genomes/mL) (OBiO Technology Corp., Ltd) were injected bilaterally into the hippocampal CA1 regions using the following coordinates relative to bregma: 2.0 mm posterior, 1.8 mm lateral, and 1.4 mm ventral at a rate of 0.125 μL/min using a microinjection system (World Precision Instruments). The needle was kept in place for 2 min before withdrawal, and the skin was sutured with surgical line. The virus-injected mice were tested for behavior two weeks later.

Immunofluorescence staining of brain sections

Mice (P21 or P60) were anesthetized and immediately perfused, brains were dissected out, fixed overnight in 4% (vol/vol) paraformaldehyde, incubated overnight with 30% sucrose in phosphate-buffered saline (PBS), and then sliced into 30 μm-thick coronal sections using a cryotome (Model CM-1950; Leica Biosystems). For immunostaining of brain sections, floating 30 μm-thick slices were rinsed with PBS and permeabilized in 0.4% Triton X-100 in 0.01M PBS for 30 min. Cyrosections were blocked with 2% BSA in PBS containing 0.4% Triton X-100 for 2 h at room temperature (RT), then incubated with primary antibodies overnight at 4 °C. Secondary antibodies conjugated with Alexa Fluor 488 or 555 were used for detection. Sections were then incubated with DAPI (Roche, Grenzach-Wyhlen, Germany) for nuclear staining for 3 min at RT. Following rinsing, cyrosections were mounted on gelatin-coated slides and covered

with coverslip with mounting medium. Confocal images were collected using the

Spectral Imaging Confocal Microscope Digital Eclipse A1Si (Nikon, Tokyo, Japan)

with a 40× Plan Fluo (NA 1.30) oil objective (Yang et al., 2018).

Cell culture, transfection and drug treatment

Primary hippocampal neurons were cultured as previously described (Yang et al.,

2021). Briefly, Hippocampi were rapidly dissected under sterile conditions, kept in

cold DMEM (4°C) with high glucose. Tissue was then digested with 0.25% trypsin

solution at 37°C for 18-20 min, triturated in DMEM supplemented with 10% F12

(11765054, Gibco) and 10% FBS (Gibco, Carlsbad, CA, USA). Then the tissue was

mechanically dissociated into single cells by gentle pipetting up and down. Neurons

were plated at a density of ~3.0-3.5× 10^5 cells/well in poly-D-lysine (P6407,

Sigma-Aldrich) -coated 6-well plate and at a density of 2.5–3.0×10^4 cells/well in

24-well plate. Four hours later, media was changed to Neurobasal A (NB-A) growth

medium supplemented with 2% B27 supplement (17504-044, Gibco), 1% GlutaMAX

(35050-061, Gibco) and 33 mM glucose. Culture media were changed by half volume

with Neurobasal maintenance media once a week. The cultures were maintained at

37°C with 5% CO_2. Experiments were performed at DIV 14-16 unless otherwise

stated. For neuronal transfection, hippocampal neurons (DIV11) were transfected

using Lipofectamine-LTX (15338100, Invitrogen, ThermoFisher Scientific). For 12

mm coverslips in individual wells of a 24 well plate, 0.5 μg of DNA was mixed with

0.5 μL PLUS reagent in 50 μL of NB-A and incubated for 5 min at RT, then gently

mixed with 1 μL Lipofectamine LTX in 50 μL NB-A and incubated for 20 min at RT.

The DNA-Lipofectamine LTX mixture was then added to the neurons in NB-A at

37°C in 5% CO_2 for 1 h. Neurons were then rinsed with NB-A and incubated in the

original medium at 37°C in 5% CO_2 for 3-4 days.

For HEK293T cell culture, DMEM supplemented with 10% FBS were used. Cell

transfections were performed using Lipofectamine-2000 (11668019, Invitrogen,

ThermoFisher Scientific) according to the manufacturer's instructions. Cells were

harvested 24-48 h post transfection.

For inhibitor treatment, primary neurons cultured on coverslips were

pre-incubated with Latrunculin A (1 μM, 76343-93-6, Sigma-Aldrich), Nocodazole (1

μM, 31430-18-9, Sigma-Aldrich) for 60 min or 120 min.

Immunofluorescence staining of cultured neurons

Hippocampal neurons were fixed in 4% paraformaldehyde/4% sucrose in PBS at RT

for 15 min, then permeabilized in 0.4% Triton X-100 in PBS for 10 min. After

blocking with 2% BSA in PBS containing 0.4% Triton X-100 for 30 min at RT,

neurons were incubated with primary antibodies for 1 h at RT or overnight at 4°C, and

appropriate secondary antibodies conjugated with Alexa Fluor 488, Alexa Fluor 555,

or Alexa Fluor 647 were used for detection. The primary antibody was diluted in a

PBS-2% BSA solution and incubated overnight at 4°C. After washing with PBS three

times, samples were incubated with Alexa Fluor secondary antibodies (1:2000,

Thermo Fisher) for 1 h at RT, and mounted without bubbles.

Biotinylation of cell surface proteins

Surface biotinylation experiments were performed as previously described (Guo et al.,

2022). Briefly, Neurons were washed with ice cold extracellular solution (ECS) (in

mM: 125 NaCl, 2 CaCl$_2$, 2.5 KCl, 5 HEPES, 33 glucose, pH, 8.0) and incubated for

30 min with 0.8 mg/ml NHS-SS-Biotin (21327, Thermo Fisher Scientific) in ECS

buffer at 37°C. After incubation, cells were washed once with PBS and the unbound

biotin was quenched via two 8 min incubations with quenching buffer (20 mM

glycine in PBS). Lysis was performed by mechanical scraping in ice-cold lysis buffer

(50 mM Tris-Cl pH 7.4, 150 mM NaCl, 1% TritonX-100, 0.1% SDS). After rotating

for 30 min at 4°C, lysates were centrifuged for 10 min at 14,000× *g* at 4°C. Balanced

Pierce Streptavidin Agarose beads (25 μL, 20353, Thermo Fisher Scientific) were

added to the supernatant. The mixture of lysates and beads were rotated at 4°C

overnight and then washed twice with ice-cold lysis buffer. Beads were incubated

with 2× SDS sample buffer for 10 min at 37°C and biotinylated proteins were analyzed by immunoblotting.

Protein expression and purification

GST-tagged EEN1 or gephyrin fragments was expressed in *E. coli* BL21 (DE3). Cells were grown at 37°C in LB (g/L: tryptone 10, yeast extract 5, NaCl 10) supplemented with ampicillin. Cells were induced at OD600 of ~0.6 with 0.4 mM isopropyl-D-1-thiogalactopyranoside (IPTG, A600168-0005, Bio Basic Inc) for 4 h at 30°C or 16 h at 16°C. Cell pellets were resuspended in PBS supplemented with 0.2% Triton X-100 and 0.1 mM PMSF. Proteins were purified with Glutathione Sepharose 4B (GE17-0756-01, Sigma-Aldrich, St. Louis, MO, USA) according to the manufacturer's instructions.

GST-pull down, co-immunoprecipitation (IP) and immunoisolation

For GST-pull down assays, GST-fused protein EEN1 and its fragments were expressed and purified from *E. coli*. Myc-tagged GPN constructs were expressed in HEK293 cells for 48 h and lysates were prepared in a buffer (in mM: Tris-HCl 50, pH 7.4, NaCl 150, 1% NP-40, EDTA 2) plus protease inhibitors cocktail. Cell lysates were centrifuged at 12,000× *g* for 15 min at 4°C and the supernatant was incubated with individual GST-fused proteins conjugated with Glutathione-Sepharose beads for overnight at 4°C. Beads were washed five times with lysis buffer and boiled in 2× SDS sample buffer. Bound proteins were analyzed by immunoblotting.

For co-IP from brain lysates, the whole brain from 8-10-week-old WT and KO mice were dissected and homogenized in the pre-chilled lysis buffer (in mM: Tris-HCl 20, pH 7.4, NaCl 150, 1% NP-40, EDTA 5) plus protease inhibitor cocktail, followed by incubation on ice for 15 min. The lysates were then centrifuged at 12,000× *g* for 15 min and the supernatant was collected and incubated with anti-EEN1 antibody (2 μg) or control IgG overnight at 4°C, followed by incubation with 20 μL Dynabeads

638 Protein A (ThermoFisher, 10007D for rabbit antibody) for 2 h at 4°C. After washing

639 the beads with the same lysis buffer four times, the bound proteins were incubated 2×

640 SDS sample buffer at 37°C for 10 min and subjected to SDS-PAGE and Western blot

641 with indicated antibodies.

642  For co-IP of proteins in HEK293T cells, cells were lysed with lysis buffer (in

643 mM: 0.5% NP-40, Tris-HCl 50, pH 7.4, NaCl 150, EDTA 5) supplemented with

644 protease inhibitors. Lysates were then centrifuged at $16,000\times g$ for 15 min at 4°C. Cell

645 lysates were incubated with anti-Flag Affinity Gel (A2220, Sigma-Aldrich) overnight

646 at 4°C on a roller mixer. Immunoprecipitates were washed four times in lysis buffer

647 and boiled in 2× SDS sample buffer, then subjected to SDS-PAGE and

648 immunoblotting.

649  For immunoisolation of membrane proteins, hippocampi were homogenized with

650 lysis buffer (in mM: Tris-HCl 20, pH 7.4, HEPES 10, pH 7.4, NaCl 150, sucrose 250,

651 EDTA 0.5, pH 8.0) supplemented with protease inhibitors and centrifuged at $800\times g$

652 for 15 min. The supernatants were collected and subjected to high-speed

653 centrifugation at $100,000\times g$ for 1 h (TLS-55 rotor, OptimaTMMAX Ultracentrifuge;

654 Beckman Coulter, USA). The supernatants (S100) and pellets (P100, the membrane

655 fraction) resuspended in lysis buffer were subjected to immunoisolation with

656 Dynabeads Protein A (100-4D, Invitrogen, Carlsbad, CA, USA) coupled with 2 μg of

657 rabbit anti-EEN1 or rabbit anti-GABA$_A$R α1 antibody. Bound proteins were eluted by

658 incubation with 2× SDS loading buffer at 37°C for 10 min and subjected to

659 SDS-PAGE and immunoblotting.

661 **PSD fractionation of brain lysates**

662 PSD fractions from mouse brain were prepared as previously described (Yang et al.,

663 2015). In brief, hippocampi were homogenized on ice with 20 strokes of a

664 Teflon-glass homogenizer in 1 mL of HEPES-buffered sucrose (0.32 M sucrose, 4

665 mM HEPES, pH 7.4) containing freshly added protease inhibitors, then centrifuged at

666 $800\times g$ at 4°C to remove the pelleted nuclear fraction (P1). Supernatant (S1) was

centrifuged at 10,000× g for 15 min to yield the crude synaptosomal pellet (P2) and

the pellet was washed once in 1 mL HEPES-buffered sucrose. P2 was lysed by

hypoosmotic shock in 900 μL ice-cold 4 mM HEPES, pH 7.4 plus protease inhibitors,

homogenized by pipetting and rotating for 30 min at 4 °C. The lysate was centrifuged

at 25,000× g for 20 min to yield supernatant (S3, crude synaptic vesicle fraction) and

pellet (P3, lysed synaptosomal membrane fraction). To prepare the PSD fraction, P3

was resuspended in 900 μL of ice-cold 50 mM HEPES, pH 7.4, 2 mM EDTA, plus

protease inhibitors and 0.5% Triton X-100, rotated for 15 min at 4 °C and centrifuged

at 32,000× g for 20 min to obtain the PSD pellet. PSD pellets were resuspended in 50

μL ice-cold 50 mM HEPES, pH 7.4, 2 mM EDTA plus protease inhibitors. Proteins

were subjected to SDS-PAGE and immunoblotting.

Protein identification with mass spectrometry and pathway analysis

For identification of EEN1-interacting proteins, mouse brain was homogenized with

lysis buffer (in mM: Tris-HCl 20, pH 7.4, HEPES 10, pH 7.4, NaCl 150, EDTA 1, 1%

NP-40) supplemented with protease inhibitors. Lysates were centrifuged at 12,000× g

for 15 min. The supernatants were incubated with rabbit anti-EEN1 antibody

overnight at 4°C. Beads were rinsed with lysis buffer for five times and bound

proteins were resolved by SDS-PAGE. Proteins in the gels were reduced with 10 mM

DTT at 37°C for 1 h, then alkylated with 25 mM iodoacetamide at RT for 1 h in the

dark before digestion with trypsin (T1426, Sigma-Aldrich; enzyme-to-substrate ratio

1:50) in 25 mM ammonium bicarbonate at 37°C overnight. Tryptic peptides were

extracted from gel by sonication with a buffer containing 5% trifluoroacetic acid and

50% acetonitrile. The liquid was dried by SpeedVac, and peptides were resolubilized

in 0.1% formic acid and filtered with 0.45 μm centrifugal filters before analysis with a

TripleTOF 5600 mass spectrometer (AB SCIEX, Canada) coupled to an Eksigent

nanoLC. Proteins were identified by searching the MS/MS spectra against the *Mus*

musculus SwissProt database using the ProteinPilot 4.2 software.

Carbamidomethylation of cysteine was set as the fixed modification. Trypsin was

specified as the proteolytic enzyme with a maximum of 2 miss cleavages. Mass

tolerance was set to 0.05 Da and the false discovery rates for both proteins and

peptides were set at 1%. For enrichment pathway analysis: Only proteins with

intensity to IgG ratio of > 1.5 were considered in the analysis. These genes were

analyzed by Gene Ontology (GO) Enrichment Analysis (http://bioinfo.org/kobas).

Electrophysiology

To knockout *EEN1* in the brain, *EEN1*$^{fl/fl}$ mice were injected with high-titer AAV

stock carrying pAOV-CaMKIIα-mCherry-2A-Cre (AAV-mCherry-2A-Cre) (1~5 ×

10^{13} IU/ml) in the hippocampal CA1 region within 24 h after birth. Newborn mice

were anesthetized on ice for 4-5 min and then mounted in a custom ceramic mold to

align the head along the X and Y axes, with lambda set as (X, Y) = (0, 0). The zero

point of the Z axis was the position at which the injecting needle penetrated the skin.

Approximately 10 nL of the viral solution was injected at each of the seven sites

(coordinates in mm, [X, Y, Z] = [1.2, 1.2, 1.4/1.0/0.6] and [1.5, 1.0, 1.7/1.3/0.9/0.5])

targeting the hippocampus in each cerebral hemisphere using a microsyringe (Sutter

Instrument, Novato, CA, USA) and beveled glass injection pipette (Yang et al., 2018).

For the rescue experiment, P0 *EEN1*$^{fl/fl}$ mice were injected with a mixture of two

high-titer AAV stocks: AAV-EGFP and AAV-mCherry or AAV-mCherry-2A-Cre;

AAV-mCherry-2A-Cre   and   AAV-EGFP-2A-FLAG-EEN1,

AAV-EGFP-2A-FLAG-EEN1 Y343A or AAV-EGFP-2A-FLAG-EEN1 KKK-EEE.

Approximately 20 nL of the viral solution was injected at each of the seven sites. The

injected pups were returned to the home cage and used for recording 2 weeks later.

To prepare acute brain slices for electrophysiological analyses, mice were

anesthetized with tribromoethanol (330-390 mg/kg) (T48402, Sigma-Aldrich, St.

Louis, MO, USA) and decapitated. Whole brains were removed promptly and placed

in ice-cold oxygenated (95% O_2/5% CO_2) low-Ca^{2+}/high-Mg^{2+} high-sucrose cutting

solution (in mM: KCl 2.5, $NaHCO_3$ 25, NaH_2PO_4 1.25, glucose 10, sucrose 210,

Na-Ascorbate 1.3, $CaCl_2$ 0.5, $MgSO_4$ 7). Coronal 300 μm slices were obtained using a

vibrating tissue slicer (Leica VT1200S; Leica Biosystems, Wetzlar, Deutschland). Slices were immediately transferred to standard artificial cerebrospinal fluid (ACSF) (in mM: NaCl 119, KCl 2.5, $MgSO_4$ 1.3, $CaCl_2$ 2.5, glucose 11, $NaHCO_3$ 26.2, NaH_2PO_4 1, pH 7.3-7.4) at 33°C and continuously bubbled with 95% O_2/5% CO_2. After 30 min of incubation, slices were transferred to a recording chamber with the same extracellular buffer at room temperature (RT, ~25°C).

For whole-cell patch clamp recordings, brain slices were continuously perfused with standard ACSF at RT under visual control with differential interference contrast illumination on an upright microscope (BX51WI, Olympus, Tokyo, Japan) with a 40× water-immersion objective and a digital camera. The mCherry- and/or EGFP-positive hippocampal CA1 pyramidal neurons were identified by an epifluorescence microscope (pE-300, CoolLED Ltd, Andover, UK). Whole-cell recordings were performed on an AAV-infected neuron and a neighboring uninfected neuron. Patch clamp data were obtained using a Multiclamp 700B amplifier (Molecular Devices, San Jose, CA, USA), and signals were low-pass filtered at 2 kHz and digitized at 10 kHz using an Axon Digidata 1550B analog-to-digital board (Molecular Devices, San Jose, CA, USA).

To analyze evoked excitatory postsynaptic currents (eEPSCs), evoked inhibitory postsynaptic currents (eIPSCs), and paired-pause ratio, cells were recorded with 3-7 $M\Omega$ borosilicate glass patch pipettes filled with the intracellular solution (in mM: $CsMeSO_3$ 135, HEPES 10, EGTA 1, QX-314 3.3, MgATP 4, NaGTP 0.3, and Na_2-phosphocreatine 8, pH 7.3) (Horn and Nicoll, 2018), following stimulation of Schaffer collaterals with a bipolar stimulation electrode placed in the stratum radiatum of the CA1 region. EPSCs were measured at a holding potential of -70 mV, and IPSCs were recorded at +10 mV. Paired-pulse ratios of EPSCs were measured at a holding potential of -70 mV by applying two electrical stimulations at a 50 ms interval and calculating the ratio of the peak amplitudes. Paired-pulse ratios of IPSCs were measured at a holding potential of +10 mV by applying two electrical stimulations at a 200 ms interval and calculating the ratio of the peak amplitudes. E/I

ratios were measured by applying the same stimulation current and then calculating

the ratio of the absolute values of the peak amplitudes of EPSCs and IPSCs. The

miniature EPSCs and IPSCs were measured in the ACSF bath solution containing 1

μM tetrodotoxin (TTX, MSS0002, BGB Analytik AG, BL, Switzerland) at -70 mV

and +10 mV, respectively, in the voltage-clamp mode. Data were collected and

analyzed using the pClamp 10.6 software (Molecular Devices, San Jose, CA, USA).

All electrophysiological recordings were carried out blindly and all n numbers in

the data figures indicate the number of recorded neurons. All chemicals were

purchased from Sigma-Aldrich except otherwise noted.

Confocal and three-dimensional structured illumination microscopy (3D-SIM) image acquisition and analysis

Confocal images were collected using the Spectral Imaging Confocal Microscope

Digital Eclipse C1Si (Nikon, Tokyo, Japan) with a 100× Plan Apochromat VC (NA

1.40) oil objective (4× optical zoom). Images were z projections of images taken at

0.25-μm step intervals. The number of planes, typically 5-7, was chosen to cover the

entire dendrite from top to bottom. Images were captured using a 1024×1024 pixels

screen for neuronal cells. Laser power, digital gain, and offset settings were identical

in each experiment. Images were processed with the NIH ImageJ software and

exported into Adobe Photoshop. Only brightness and contrast were adjusted for the

whole frame, and no part of a frame was enhanced or modified in any way. Control

and experimental group neurons which were to be directly compared were imaged

with the same acquisition parameters.

For 3D-SIM, secondary antibodies conjugated with Alexa Fluor 568 were used

and images were captured as described by Niu et al (Niu et al., 2017) on the

DeltaVision OMX V4 imaging system (Applied Precision Inc, USA) with a 100× 1.4

oil objective (Olympus UPlanSApo), solid state multimode lasers (488, 593 and 642

nm) and electron-multiplying charge-coupled device (CCD) cameras (Evolve 512×

512, Photometrics, USA). Serial Z-stack sectioning was done at 125 nm intervals. The

microscope is routinely calibrated with 100 nm fluorescent spheres to calculate both

the lateral and axial limits of image resolution. SIM image stacks were reconstructed

using softWoRx 5.0 (Applied Precision) with the following settings: pixel size 39.5

nm; channel-specific optical transfer functions; Wiener filter 0.001000; discard

Negative Intensities background; drift correction with respect to first angle; custom

K0 guess angles for camera positions. Pixel registration was corrected to be less than

1 pixel for all channels using 100 nm Tetraspeck beads. Three dimensional images

were reconstructed from z stacks. For clarity of display, small linear changes of

brightness and contrast were performed on three-dimensional reconstructions.

For quantitative analysis of immunofluorescent signals in neurons, maximal

projection images were created with the ImageJ software (https://imagej.net.) from

6-7 serial optical sections. Background was subtracted by using the "subtract

background" function and the background level was held identical for all cells within

each experiment. To examine the number and fluorescence intensity of signal puncta

in soma or dendrites, the EGFP- or DsRed-labeled soma or dendrites were outlined

automatically by using the "Wand Tool" function. The region of interest (ROI) in

cultured neurons was defined along a segment of the dendrite according to the

fluorescence signal distinguished from the background. Average values of

fluorescence intensities in ROI (the total fluorescent intensity divided by the total area

of a dendritic segment) were calculated by Analyze > measure. Gephyrin clusters

were thresholded and confirmed visually to select appropriate clusters following a

minimal size cut-off by using the "Analyze Particles" plugin of the ImageJ software.

For colocalization quantifications, NIS-Elements Software was used as follows:

region of interest (ROI) of dendritic segments was manually chosen, and thresholded

using the "Threshold" function to remove background signals. Pearson's correlation

coefficient and Manders overlap coefficient were analyzed by using the "Analyze

Control > colocalization" function.

Statistical analysis

For all biochemical, cell biological and electrophysiological recordings, at least three

independent experiments were performed (independent cultures, transfections or

different mice). Statistical analysis was performed in GraphPad Prism 9.0 (GraphPad

Software, La Jolla, CA). Data sets were tested for normality and direct comparisons

between two groups were made using two-tailed Student's t test (t test, for normally

distributed data) as indicated. To evaluate statistical significance of three or more

groups of samples, one-way analysis of variance with a Tukey test was used. The

statistical significance was defined as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$,

respectively. All data are presented as mean ± S.E.M.

Acknowledgments

We thank Drs Shiva Tyagarajan (Addgene #68816) and Peter Scheiffele (Addgene #

15246) for constructs. This work was supported by funding from the National Natural

Science Foundation of China (NSFC 32271005 and 32070785 to Y. Yang, 31921002,

32270730 and 91954126 to J-J. Liu) and the State Key Laboratory of Molecular

Developmental Biology (2022-MDB-KF-13).

The authors declare no competing financial interests.

Author contributions: J-J. Liu and Y. Yang conceived the project. Y. Yang and J-J. Liu

designed and supervised research. X. Chen, D. Pan and Y. Yang performed

experiments and analyzed the data. Y. Yang and J-J. Liu wrote the manuscript with

help from all other authors.

References

Allison, D.W., Chervin, A.S., Gelfand, V.I., and Craig, A.M. (2000). Postsynaptic scaffolds of

excitatory and inhibitory synapses in hippocampal neurons: maintenance of core components

independent of actin filaments and microtubules. The Journal of neuroscience : the official

journal of the Society for Neuroscience *20*, 4545-4554.

Bademosi, A.T., Decet, M., Kuenen, S., Calatayud, C., Swerts, J., Gallego, S.F., Schoovaerts,

N., Karamanou, S., Louros, N., Martin, E.*, et al.* (2023). EndophilinA-dependent coupling

between activity-induced calcium influx and synaptic autophagy is disrupted by a

Parkinson-risk mutation. Neuron.

Bai, G., Wang, Y., and Zhang, M. (2021). Gephyrin-mediated formation of inhibitory

postsynaptic density sheet via phase separation. Cell research *31*, 312-325.

Bogdanov, Y., Michels, G., Armstrong-Gold, C., Haydon, P.G., Lindstrom, J., Pangalos, M.,

and Moss, S.J. (2006). Synaptic GABAA receptors are directly recruited from their

extrasynaptic counterparts. The EMBO journal *25*, 4381-4389.

Canitano, R., and Pallagrosi, M. (2017). Autism Spectrum Disorders and Schizophrenia

Spectrum Disorders: Excitation/Inhibition Imbalance and Developmental Trajectories.

Frontiers in psychiatry *8*, 69.

Canitano, R., and Palumbi, R. (2021). Excitation/Inhibition Modulators in Autism Spectrum

Disorder: Current Clinical Research. Frontiers in neuroscience *15*, 753274.

Chang, D., Nalls, M.A., Hallgrímsdóttir, I.B., Hunkapiller, J., van der Brug, M., Cai, F., Kerchner,

G.A., Ayalon, G., Bingol, B., Sheng, M.*, et al.* (2017). A meta-analysis of genome-wide

association studies identifies 17 new Parkinson's disease risk loci. Nature genetics *49*,

1511-1516.

Charrier, C., Ehrensperger, M.V., Dahan, M., Lévi, S., and Triller, A. (2006). Cytoskeleton

regulation of glycine receptor number at synapses and diffusion in the plasma membrane. J

Neurosci *26*, 8502-8511.

Chen, J.L., Villa, K.L., Cha, J.W., So, P.T., Kubota, Y., and Nedivi, E. (2012). Clustered

dynamics of inhibitory synapses and dendritic spines in the adult neocortex. Neuron *74*,

361-373.

Chih, B., Gollan, L., and Scheiffele, P. (2006). Alternative splicing controls selective

trans-synaptic interactions of the neuroligin-neurexin complex. Neuron *51*, 171-178.

Corponi, F., Bonassi, S., Vieta, E., Albani, D., Frustaci, A., Ducci, G., Landi, S., Boccia, S.,

Serretti, A., and Fabbri, C. (2019). Genetic basis of psychopathological dimensions shared

between schizophrenia and bipolar disorder. Prog Neuropsychopharmacol Biol Psychiatry *89*,

23-29.

Cramer, T.M.L., Pinan-Lucarre, B., Cavaccini, A., Damilou, A., Tsai, Y.C., Bhat, M.A.,

Panzanelli, P., Rama, N., Mehlen, P., Benke, D.*, et al.* (2023). Adamtsl3 mediates DCC

signaling to selectively promote GABAergic synapse function. Cell reports *42*, 112947.

Dhir, A. (2012). Pentylenetetrazol (PTZ) kindling model of epilepsy. Current protocols in

neuroscience *Chapter 9*, Unit9.37.

Essrich, C., Lorez, M., Benson, J.A., Fritschy, J.M., and Luscher, B. (1998). Postsynaptic

clustering of major GABAA receptor subtypes requires the gamma 2 subunit and gephyrin.

Nature neuroscience *1*, 563-571.

Favuzzi, E., Deogracias, R., Marques-Smith, A., Maeso, P., Jezequel, J., Exposito-Alonso, D.,

Balia, M., Kroon, T., Hinojosa, A.J., E, F.M.*, et al.* (2019). Distinct molecular programs regulate

synapse specificity in cortical inhibitory circuits. Science *363*, 413-417.

Foss-Feig, J.H., Adkinson, B.D., Ji, J.L., Yang, G., Srihari, V.H., McPartland, J.C., Krystal, J.H.,

Murray, J.D., and Anticevic, A. (2017). Searching for Cross-Diagnostic Convergence: Neural

Mechanisms Governing Excitation and Inhibition Balance in Schizophrenia and Autism

Spectrum Disorders. Biol Psychiatry *81*, 848-861.

Fritschy, J.-M., Harvey, R.J., and Schwarz, G. (2008). Gephyrin: where do we stand, where do

we go? Trends in neurosciences *31*, 257-264.

Fritschy, J.M. (2008). Epilepsy, E/I Balance and GABA(A) Receptor Plasticity. Frontiers in

molecular neuroscience *1*, 5.

Gad, H., Ringstad, N., Low, P., Kjaerulff, O., Gustafsson, J., Wenk, M., Di Paolo, G., Nemoto,

Y., Crun, J., Ellisman, M.H.*, et al.* (2000). Fission and uncoating of synaptic clathrin-coated

vesicles are perturbed by disruption of interactions with the SH3 domain of endophilin. Neuron

27, 301-312.

George, S., Chiou, T.T., Kanamalla, K., and De Blas, A.L. (2021). Recruitment of Plasma

Membrane GABA-A Receptors by Submembranous Gephyrin/Collybistin Clusters. Cellular

and molecular neurobiology.

Germer, E.L., Imhoff, S., Vilariño-Güell, C., Kasten, M., Seibler, P., Brüggemann, N., Klein, C.,

and Trinh, J. (2019). The Role of Rare Coding Variants in Parkinson's Disease GWAS Loci.

Frontiers in neurology *10*, 1284.

Gu, J., Firestein, B.L., and Zheng, J.Q. (2008). Microtubules in dendritic spine development. J

Neurosci *28*, 12120-12124.

Guo, Z., Jiang, C.H., Tong, C., Yang, Y., Wang, Z., Lam, S.M., Wang, D., Li, R., Shui, G., Shi,

Y.S.*, et al.* (2022). Activity-dependent PI4P synthesis by PI4KIIIα regulates long-term synaptic

potentiation. Cell Rep *38*, 110452.

Han, W., Li, J., Pelkey, K.A., Pandey, S., Chen, X., Wang, Y.X., Wu, K., Ge, L., Li, T.,

Castellano, D*., et al.* (2019). Shisa7 is a GABA(A) receptor auxiliary subunit controlling

benzodiazepine actions. Science *366*, 246-250.

Hoffmann, C., and Milovanovic, D. (2021). Gephyrin: a scaffold that builds a phase at the

inhibitory postsynapses. Cell research *31*, 245-246.

Horn, M.E., and Nicoll, R.A. (2018). Somatostatin and parvalbumin inhibitory synapses onto

hippocampal pyramidal neurons are regulated by distinct mechanisms. Proceedings of the

National Academy of Sciences *115*, 589-594.

Huo, J.Z., Cortez, M.A., and Snead, O.C., 3rd (2009). GABA receptor proteins within lipid rafts

in the AY-9944 model of atypical absence seizures. Epilepsia *50*, 776-788.

Jacob, T.C., Bogdanov, Y.D., Magnus, C., Saliba, R.S., Kittler, J.T., Haydon, P.G., and Moss,

S.J. (2005). Gephyrin regulates the cell surface dynamics of synaptic GABAA receptors. J

Neurosci *25*, 10469-10478.

Jaworski, J., Kapitein, L.C., Gouveia, S.M., Dortland, B.R., Wulf, P.S., Grigoriev, I., Camera, P.,

Spangler, S.A., Di Stefano, P., Demmers, J*., et al.* (2009). Dynamic microtubules regulate

dendritic spine morphology and synaptic plasticity. Neuron *61*, 85-100.

Kalscheuer, V.M., Musante, L., Fang, C., Hoffmann, K., Fuchs, C., Carta, E., Deas, E.,

Venkateswarlu, K., Menzel, C., Ullmann, R*., et al.* (2009). A balanced chromosomal

translocation disrupting ARHGEF9 is associated with epilepsy, anxiety, aggression, and

mental retardation. Human mutation *30*, 61-68.

Kasaragod, V.B., Mortensen, M., Hardwick, S.W., Wahid, A.A., Dorovykh, V., Chirgadze, D.Y.,

Smart, T.G., and Miller, P.S. (2022). Mechanisms of inhibition and activation of extrasynaptic

alphabeta GABA(A) receptors. Nature *602*, 529-533.

Kasaragod, V.B., and Schindelin, H. (2019). Structure of Heteropentameric GABAA Receptors

and Receptor-Anchoring Properties of Gephyrin. Frontiers in molecular neuroscience *12*, 191.

Kins, S., Betz, H., and Kirsch, J. (2000). Collybistin, a newly identified brain-specific GEF,

induces submembrane clustering of gephyrin. Nat Neurosci *3*, 22-29.

Kirsch, J., and Betz, H. (1995). The postsynaptic localization of the glycine

receptor-associated protein gephyrin is regulated by the cytoskeleton. The Journal of

neuroscience : the official journal of the Society for Neuroscience *15*, 4148-4156.

Kirsch, J., Langosch, D., Prior, P., Littauer, U.Z., Schmitt, B., and Betz, H. (1991). The 93-kDa

glycine receptor-associated protein binds to tubulin. J Biol Chem *266*, 22242-22245.

Klausberger, T., and Somogyi, P. (2008). Neuronal diversity and temporal dynamics: the unity

of hippocampal circuit operations. Science *321*, 53-57.

Kneussel, M., Brandstätter, J.H., Laube, B., Stahl, S., Müller, U., and Betz, H. (1999). Loss of

Postsynaptic GABAAReceptor Clustering in Gephyrin-Deficient Mice. J Neurosci *19*,

9289-9297.

Kneussel, M., Haverkamp, S., Fuhrmann, J.C., Wang, H., Wässle, H., Olsen, R.W., and Betz,

H. (2000). The gamma-aminobutyric acid type A receptor (GABAAR)-associated protein

GABARAP interacts with gephyrin but is not involved in receptor anchoring at the synapse.

Proc Natl Acad Sci U S A *97*, 8594-8599.

Koh, W., Kwak, H., Cheong, E., and Lee, C.J. (2023). GABA tone regulation and its cognitive

functions in the brain. Nature reviews Neuroscience *24*, 523-539.

Liu, Y.T., Tao, C.L., Zhang, X., Xia, W., Shi, D.Q., Qi, L., Xu, C., Sun, R., Li, X.W., Lau, P.M.*,*

et al. (2020). Mesophasic organization of GABAA receptors in hippocampal inhibitory

synapses. Nat Neurosci *23*, 1589-1596.

Loebrich, S., Bahring, R., Katsuno, T., Tsukita, S., and Kneussel, M. (2006). Activated radixin

is essential for GABAA receptor alpha5 subunit anchoring at the actin cytoskeleton. The

EMBO journal *25*, 987-999.

Miles, R., Tóth, K., Gulyás, A.I., Hájos, N., and Freund, T.F. (1996). Differences between

somatic and dendritic inhibition in the hippocampus. Neuron *16*, 815-823.

Milosevic, I., Giovedi, S., Lou, X., Raimondi, A., Collesi, C., Shen, H., Paradise, S., O'Toole, E.,

Ferguson, S., Cremona, O., *et al.* (2011). Recruitment of endophilin to clathrin-coated pit

necks is required for efficient vesicle uncoating after fission. Neuron *72*, 587-601.

Mody, I., and Pearce, R.A. (2004). Diversity of inhibitory neurotransmission through GABA(A)

receptors. Trends in neurosciences *27*, 569-575.

Mukherjee, J., Kretschmannova, K., Gouzer, G., Maric, H.M., Ramsden, S., Tretter, V., Harvey,

K., Davies, P.A., Triller, A., Schindelin, H., *et al.* (2011). The residence time of GABA(A)Rs at

inhibitory synapses is determined by direct binding of the receptor α1 subunit to gephyrin. J

Neurosci *31*, 14677-14687.

Murdoch, J.D., Rostosky, C.M., Gowrisankaran, S., Arora, A.S., Soukup, S.F., Vidal, R.,

Capece, V., Freytag, S., Fischer, A., Verstreken, P., *et al.* (2016). Endophilin-A Deficiency

Induces the Foxo3a-Fbxo32 Network in the Brain and Causes Dysregulation of Autophagy and

the Ubiquitin-Proteasome System. Cell reports *17*, 1071-1086.

Niu, Y., Dai, Z., Liu, W., Zhang, C., Yang, Y., Guo, Z., Li, X., Xu, C., Huang, X., Wang, Y., *et al.*

(2017). Ablation of SNX6 leads to defects in synaptic function of CA1 pyramidal neurons and

spatial memory. eLife *6*.

Nothdurfter, C., Tanasic, S., Di Benedetto, B., Uhr, M., Wagner, E.M., Gilling, K.E., Parsons,

C.G., Rein, T., Holsboer, F., Rupprecht, R.*, et al.* (2013). Lipid raft integrity affects GABAA

receptor, but not NMDA receptor modulation by psychopharmacological compounds. The

international journal of neuropsychopharmacology *16*, 1361-1371.

Nusser, Z., Hajos, N., Somogyi, P., and Mody, I. (1998). Increased number of synaptic

GABA(A) receptors underlies potentiation at hippocampal inhibitory synapses. Nature *395*,

172-177.

Parato, J., and Bartolini, F. (2021). The microtubule cytoskeleton at the synapse.

Neuroscience letters *753*, 135850.

Poulopoulos, A., Aramuni, G., Meyer, G., Soykan, T., Hoon, M., Papadopoulos, T., Zhang, M.,

Paarmann, I., Fuchs, C., Harvey, K.*, et al.* (2009). Neuroligin 2 drives postsynaptic assembly

at perisomatic inhibitory synapses through gephyrin and collybistin. Neuron *63*, 628-642.

Ren, Y., Xu, H.W., Davey, F., Taylor, M., Aiton, J., Coote, P., Fang, F., Yao, J., Chen, D., Chen,

J.X.*, et al.* (2008). Endophilin I expression is increased in the brains of Alzheimer disease

patients. The Journal of biological chemistry *283*, 5685-5691.

Ringstad, N., Gad, H., Low, P., Di Paolo, G., Brodin, L., Shupliakov, O., and De Camilli, P.

(1999). Endophilin/SH3p4 is required for the transition from early to late stages in

clathrin-mediated synaptic vesicle endocytosis. Neuron *24*, 143-154.

Ringstad, N., Nemoto, Y., and De Camilli, P. (1997). The SH3p4/Sh3p8/SH3p13 protein family:

binding partners for synaptojanin and dynamin via a Grb2-like Src homology 3 domain. Proc

Natl Acad Sci U S A *94*, 8569-8574.

Russo, I., Gavello, D., Menna, E., Vandael, D., Veglia, C., Morello, N., Corradini, I., Focchi, E.,

Alfieri, A., Angelini, C.*, et al.* (2019). p140Cap Regulates GABAergic Synaptogenesis and

Development of Hippocampal Inhibitory Circuits. Cerebral Cortex *29*, 91-105.

Saiepour, L., Fuchs, C., Patrizi, A., Sassoè-Pognetto, M., Harvey, R.J., and Harvey, K. (2010).

Complex role of collybistin and gephyrin in GABAA receptor clustering. J Biol Chem *285*,

29623-29631.

Sakimoto, Y., Oo, P.M., Goshima, M., Kanehisa, I., Tsukada, Y., and Mitsushima, D. (2021).

Significance of GABAA Receptor for Cognitive Function and Hippocampal Pathology. Int J Mol

Sci *22*.

Schulz, A., Miehl, C., Berry, M.J., 2nd, and Gjorgjieva, J. (2021). The generation of cortical

novelty responses through inhibitory plasticity. eLife *10*.

Schuske, K.R., Richmond, J.E., Matthies, D.S., Davis, W.S., Runz, S., Rube, D.A., van der

Bliek, A.M., and Jorgensen, E.M. (2003). Endophilin is required for synaptic vesicle

endocytosis by localizing synaptojanin. Neuron *40*, 749-762.

Scott, S., and Aricescu, A.R. (2019). A structural perspective on GABA(A) receptor

pharmacology. Current opinion in structural biology *54*, 189-197.

Sexton, C.A., Penzinger, R., Mortensen, M., Bright, D.P., and Smart, T.G. (2021). Structural

determinants and regulation of spontaneous activity in GABAA receptors. Nature

communications *12*, 5457.

Soukup, S.F., Kuenen, S., Vanhauwaert, R., Manetsberger, J., Hernandez-Diaz, S., Swerts, J.,

Schoovaerts, N., Vilain, S., Gounko, N.V., Vints, K.*, et al.* (2016). A LRRK2-Dependent

EndophilinA Phosphoswitch Is Critical for Macroautophagy at Presynaptic Terminals. Neuron

92, 829-844.

Spruston, N. (2008). Pyramidal neurons: dendritic structure and synaptic integration. Nature

reviews Neuroscience *9*, 206-221.

Tao, C.L., Liu, Y.T., Sun, R., Zhang, B., Qi, L., Shivakoti, S., Tian, C.L., Zhang, P., Lau, P.M.,

Zhou, Z.H.*, et al.* (2018). Differentiation and Characterization of Excitatory and Inhibitory

Synapses by Cryo-electron Tomography and Correlative Microscopy. The Journal of

neuroscience : the official journal of the Society for Neuroscience *38*, 1493-1510.

Thomas, P., Mortensen, M., Hosie, A.M., and Smart, T.G. (2005). Dynamic mobility of

functional GABAA receptors at inhibitory synapses. Nature neuroscience *8*, 889-897.

Tretter, V., Jacob, T.C., Mukherjee, J., Fritschy, J.M., Pangalos, M.N., and Moss, S.J. (2008).

The clustering of GABA(A) receptor subtypes at inhibitory synapses is facilitated via the direct

binding of receptor alpha 2 subunits to gephyrin. The Journal of neuroscience : the official

journal of the Society for Neuroscience *28*, 1356-1365.

Tyagarajan, S.K., and Fritschy, J.M. (2014). Gephyrin: a master regulator of neuronal function?

Nat Rev Neurosci *15*, 141-156.

Tyagarajan, S.K., Ghosh, H., Yévenes, G.E., Imanishi, S.Y., Zeilhofer, H.U., Gerrits, B., and

Fritschy, J.M. (2013). Extracellular signal-regulated kinase and glycogen synthase kinase 3β

regulate gephyrin postsynaptic aggregation and GABAergic synaptic function in a

calpain-dependent mechanism. The Journal of biological chemistry *288*, 9634-9647.

van Bommel, B., Konietzny, A., Kobler, O., Bar, J., and Mikhaylova, M. (2019). F-actin patches

associated with glutamatergic synapses control positioning of dendritic lysosomes. EMBO J *38*,

e101183.

Van Erum, J., Van Dam, D., and De Deyn, P.P. (2019). PTZ-induced seizures in mice require

a revised Racine scale. Epilepsy & behavior : E&B *95*, 51-55.

Verstreken, P., Koh, T.W., Schulze, K.L., Zhai, R.G., Hiesinger, P.R., Zhou, Y., Mehta, S.Q.,

Cao, Y., Roos, J., and Bellen, H.J. (2003). Synaptojanin is recruited by endophilin to promote

synaptic vesicle uncoating. Neuron *40*, 733-748.

Wagner, S., Lee, C., Rojas, L., Specht, C.G., Rhee, J., Brose, N., and Papadopoulos, T.

(2021). The α3 subunit of GABA(A) receptors promotes formation of inhibitory synapses in the

absence of collybistin. J Biol Chem *296*, 100709.

Watanabe, S., Mamer, L.E., Raychaudhuri, S., Luvsanjav, D., Eisen, J., Trimbuch, T.,

Sohl-Kielczynski, B., Fenske, P., Milosevic, I., Rosenmund, C.*, et al.* (2018). Synaptojanin and

Endophilin Mediate Neck Formation during Ultrafast Endocytosis. Neuron *98*, 1184-1197

e1186.

Yang, Y., Chen, J., Chen, X., Li, D., He, J., Wang, S., Zhao, S., Yang, X., Deng, S., Tong, C.*,*

et al. (2021). Endophilin A1 drives acute structural plasticity of dendritic spines in response to

Ca2+/calmodulin. The Journal of cell biology *220*.

Yang, Y., Chen, J., Guo, Z., Deng, S., Du, X., Zhu, S., Ye, C., Shi, Y.S., and Liu, J.J. (2018).

Endophilin A1 Promotes Actin Polymerization in Dendritic Spines Required for Synaptic

Potentiation. Frontiers in molecular neuroscience *11*, 177.

Yang, Y., Wei, M., Xiong, Y., Du, X., Zhu, S., Yang, L., Zhang, C., and Liu, J.J. (2015).

Endophilin A1 regulates dendritic spine morphogenesis and stability through interaction with

p140Cap. Cell research *25*, 496-516.

Yu, Q., Wang, Y., Du, F., Yan, S., Hu, G., Origlia, N., Rutigliano, G., Sun, Q., Yu, H., Ainge, J.*,*

et al. (2018a). Overexpression of endophilin A1 exacerbates synaptic alterations in a mouse

model of Alzheimer's disease. Nature communications *9*, 2968.

Yu, X., Xu, T., Ou, S., Yuan, J., Deng, J., Li, R., Yang, J., Liu, X., Li, Q., and Chen, Y. (2018b).

Endophilin A1 mediates seizure activity via regulation of AMPARs in a PTZ-kindled epileptic

mouse model. Experimental neurology *304*, 41-57.

1064

1065

Figure legends

Figure 1. Ablation of endophilin A1 in CA1 neurons disrupts the E/I balance and increases epilepsy susceptibility.

(A) Experimental protocol of PTZ kindling. Saline serves as negative control for PTZ.

(B) Representative images of convulsive behavior in each group.

(C) Seizures were monitored and scored after each injection. Nine 10-week-old mice were used in each condition. Seizure scores are shown as means ± S.E.M. Statistical test: * $p < 0.05$ and ** $p < 0.01$ compared with $EEN1^{+/+}$_PTZ.

(D) The representative mCherry-positive hippocampal CA1 neuron used for electrophysiological recording.

(E-F) Representative traces of evoked excitatory postsynaptic currents (eEPSC) and quantification of mean eEPSC amplitude (E), and paired-pulse recording (F) from 10 control (Ctrl, mCherry alone) and 10 *EEN1* KO (mCherry and Cre) pyramidal neurons.

(G-H) Representative traces of evoked inhibitory postsynaptic currents (eIPSCs) and quantification of mean eIPSC amplitude (G), and paired-pulse recording (H) from 9 control and 9 *EEN1* KO pyramidal neurons. Statistical test: * $p < 0.05$.

(I) Effect of *EEN1* KO on the E/I ratio from 9 control and 9 *EEN1* KO pyramidal neurons. Statistical test: * $p < 0.05$.

(J) Representative recordings of miniature excitatory postsynaptic currents (mEPSCs) traces (+10 mV) from control or *EEN1* KO CA1 pyramidal cells.

(K) Cumulative probability and quantification of mEPSC amplitude from 7 control and 7 *EEN1* KO pyramidal neurons.

(L) Cumulative probability and quantification of mEPSC frequency from 7 control and 7 *EEN1* KO pyramidal neurons.

(M) Representative recordings of miniature inhibitory postsynaptic currents (mIPSCs) traces (–70 mV) from control or *EEN1* KO CA1 pyramidal cells.

(N) Cumulative probability and quantification of mIPSC amplitude from 7 control and 7 *EEN1* KO pyramidal neurons.

(O) Cumulative probability and quantification of mIPSC frequency from 7 control and 7 *EEN1* KO pyramidal neurons. Statistical test: ns, non-significant. All data shown are means ± S.E.M.

Figure 2. *EEN1* KO causes defects in the formation and stabilization of inhibitory synapses.

(A-B) Immunofluorescence staining and confocal microscopy of gephyrin (GPN) in the hippocampal region of 3-week-old (P21) and 10-week-old (P60) $EEN1^{+/+}$ or $EEN1^{-/-}$ mouse brains. Scale bar, 50 μm.

(C) Quantification of the signal intensity of GPN in (A) and (B). n = 6 mice for each genotype.

(D-E) Same as (A-B) except that slices were immunostained for VGAT, (D) for P21, (E) for P60. Scale bar, 50 μm.

(F) Quantification of VGAT signal intensity in (D) and (E). n = 6 mice for each genotype.

(G-H) Same as (A-B) except that slices were immunostained for GABA$_A$R γ2 (γ2). (G) for P21, (H) for P60. Scale bar, 50 μm.

(I) Quantification of γ2 signal intensity in (G) and (H). n = 6 mice for each genotype. Statistical test: * p < 0.05, ** p < 0.01, and *** p < 0.001; ns, non-significant. All data are means ± S.E.M. Abbreviations: CA1-3, cornus ammonis region 1-3; DG, dentate gyrus; DAPI staining of cell nuclei is shown in blue.

Figure 3. Endophilin A1 is distributed in the iPSD and regulates the assembly of the inhibitory postsynaptic machinery.

(A) Cultured *EEN1$^{fl/fl}$* hippocampal neurons were transfected with construct expressing GFP or GFP-2A-Cre on DIV12, fixed and immunostained for surface GluA1 (sGluA1) or GABA$_A$R γ2 (sγ2) on DIV16. Shown are representative confocal microscopy images. Lower panels are magnification of boxed regions in the upper panels. Scale bar, 10 μm in the upper panels and 5 μm in the lower panels.

(B) Quantification of the signal intensity (left) and cluster number (right) of GPN puncta in dendrites in (A). GFP: 56 segments of dendrites from 15 neurons; Cre: 70 segments of dendrites from 18 neurons.

(C-D) Cultured *EEN1$^{fl/fl}$* hippocampal neurons were co-transfected with constructs for GFP or GFP-2A-Cre and Flag vector or Flag-endophilin A1 (Flag-EEN1) on DIV12, and immunostained for surface γ2 on DIV16. Shown are representative confocal images (C) and quantification of surface γ2 in dendrites (D). Scale bar, 5 μm. GFP: 60 segments of dendrites from 18 neurons; Cre: 59 segments of dendrites from 19 neurons; Cre + EEN1: 60 segments of dendrites from 18 neurons.

(E-F) *EEN1$^{fl/fl}$* hippocampal neurons were infected with adeno-associated virus (AAV) carrying mCherry or mCherry-2A-Cre on DIV3 and surface biotinylated on DIV16. Total and surface proteins were detected by SDS-PAGE and immunoblotting. N = 3. LE: long exposure.

(G) Hippocampal neurons were transfected with construct expressing GFP to label neuronal morphology on DIV12, then immunostained with antibodies against EEN1 and GPN on DIV16. Shown are representative z stack maximum projection images captured by structured illumination microscopy (SIM). Lower panels are insets of boxed regions in the upper panel. Dendrites and spines are outlined in white. Scale bars, 2 μm in upper panels and 1 μm in magnified images.

(H) Same as (G) except that neurons were immunostained for EEN1 and surface γ2.

(I) Same as (G) except that neurons were immunostained for EEN1 and VGAT.

(J) Immunoisolation from the membrane fraction of mouse brain tissues with antibodies
against GABA$_A$R α1 (α1) or EEN1.

(K) Immunoisolation from *EEN1$^{+/+}$* and *EEN1$^{-/-}$* brain lysates with antibodies to GABA$_A$R α1.
Shown are representative immunoblot (left) and quantification of neuroligin 2 (NL2), GPN
and GABA$_A$R γ2 co-immunoisolated with α1 (right). N = 4 independent experiments.

(L) PSD fractions were isolated by differential centrifugation from mouse brain lysates and
analyzed by immunoblotting with antibodies to EEN1 and synaptic proteins. S1, homogenates;
PSD, postsynaptic density. SYP, synaptophysin. N = 4 independent experiments. Statistical
test: * p < 0.05; ** p < 0.01; *** p < 0.001. ns: not significant. All data represent means ±
S.E.M or scatterplots with means ± S.E.M.

1153

Figure 4. Endophilin A1 interacts with gephyrin and promotes its clustering in
postsynaptic neurons.

(A) Silver staining of proteins co-immunoprecipitated from *EEN1$^{+/+}$* mouse brain lysates with
antibodies to endophilin A1.

(B) Gene Ontology (GO) enrichment analysis of EEN1-interacting proteins identified by mass
spectrometry.

(C-D) HEK293T cells co-transfected with constructs expressing Myc-tagged GPN or
HA-tagged NL2 and Flag-tagged EEN1 for 48 h were lysed for Flag IP. Input and bound
proteins were analyzed by immunoblotting with antibodies against Myc, HA and Flag.

(E) Co-IP from brain lysates with antibodies to EEN1.

(F) HEK293T cells overexpressing Flag-tagged endophilin As and Myc-tagged GPN were
lysed for co-immunoprecipitation.

(G) *EEN1$^{fl/fl}$* hippocampal neurons were co-transfected with constructs expressing GFP and
Flag vector, or GFP-2A-Cre and Flag vector, Flag-EEN1, Flag-EEN2 or Flag-EEN3 on
DIV12, fixed on DIV16 and immunostained for GPN, GFP and Flag. Shown are
representative confocal microscopy images. Scale bar, 5 μm.

(H) Quantification of GPN signal intensity (left) and cluster number (right) in dendrites in (G).
52 dendrites from 14 neurons for GFP + vector; 58 dendrites from 15 neurons for Cre +
vector; 59 dendrites from 16 neurons for Cre + EEN1; 44 dendrites from 14 neurons for Cre +
EEN2; 45 dendrites from 11 neurons for Cre + EEN3.

(I) Hippocampal neurons were co-transfected with constructs expressing GFP and vector or
Flag-EEN1 on DIV12, fixed and immunostained for GPN on DIV16. Shown are
representative confocal microscopy images of GPN in the soma (left) and dendrites (right).
Scale bars, 5 μm.

(J) Quantification of GPN signal intensity (left) and cluster number (right) in the soma and
dendrites. 65 dendrites from 25 neurons for GFP + vector; 66 dendrites from 25 neurons for

GFP + EEN1. Statistical test: ** p < 0.01; *** p < 0.001. All data represent scatterplots with

means ± S.E.M.

1182

Figure 5. Both BAR and SH3 domains are required for endophilin A1-mediated organization of iPSD.

(A) *EEN1^{fl/fl}* hippocampal neurons expressing GFP or Cre were immunostained for GPN, surface γ2 and GFP on DIV16. Shown are representative confocal microscopy images. Scale bars, 5 μm in left panels and 1 μm in magnified images.

(B-C) Quantification of the mean intensity of GPN (green) and surface γ2 (red) and their colocalization in (A). 18 neurons for GFP, 20 neurons for Cre.

(D) DIV12 hippocampal neurons were co-transfected with constructs expressing GFP and vector or Flag-EEN1, fixed on DIV16 and immunostained for surface γ2. Shown are representative confocal microscopy images of surface γ2 in the soma (left) and dendrites (right). Scale bars, 5 μm.

(E) Quantification of surface γ2 mean intensity (left) and cluster number (right) in the soma and dendrites. 68 dendrites from 20 neurons for GFP + vector, 61 dendrites from 20 neurons for GFP + EEN1.

(F) Schematic representation of EEN1 domain structure and diagrams of EEN1 fragments used for interaction mapping.

(G) GST-tagged EEN1 fragments conjugated to Glutathione-Sepharose beads were incubated with lysates of HEK293T cells overexpressing Myc-tagged GPN. Input and bound proteins were analyzed by immunoblotting with antibodies against Myc. Bottom panel: Coomassie blue-stained SDS-PAGE gel showing GST fusion proteins. FL, full length.

(H) HEK293T cells co-transfected with constructs encoding Myc-tagged GPN and Flag-tagged EEN1 fragments were lysed for Flag IP. Input and bound proteins were analyzed by immunoblotting with antibodies against Myc and Flag.

(I) DIV12 *EEN1^{fl/fl}* hippocampal neurons were co-transfected with constructs expressing GFP and Flag vector or GFP-2A-Cre, and Flag vector or Flag-EEN1 FL, Flag-EEN1 ΔSH3, Flag-EEN1 aa172-352, fixed and immunostained for GPN on DIV16. Shown are representative confocal microscopy images of GPN. Scale bar, 5 μm.

(J) Quantification of mean intensity (left) and cluster number (right) in dendrites are shown for GPN in (I). Scale bar, 5 μm. 60 dendrites from 14 neurons for GFP + vector; 62 dendrites from 15 neurons for Cre + vector; 60 dendrites from 13 neurons for Cre + FL; 50 dendrites from 12 neurons for Cre + ΔSH3; 42 dendrites from 10 neurons for Cre + aa172-352.

(K) Same as (I) except that neurons were immunostained for surface γ2. Scale bar, 5 μm.

(L) Quantification of mean intensity (left) and cluster number (right) in dendrites are shown for surface γ2 in (K). 59 dendrites from 16 neurons for GFP + vector; 58 dendrites from 14 neurons for Cre + vector; 60 dendrites from 14 neurons for Cre + FL; 45 dendrites from 12 neurons for Cre + ΔSH3; 42 dendrites from 10 neurons for Cre + aa172-352. Statistical test: *

$p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. All data represent scatterplots with means ± S.E.M.

1220

Figure 6. Actin polymerization is required for stabilization of inhibitory postsynapses.

(A-D) DIV12 cultured hippocampal neurons transiently expressing GFP were treated with

DMSO (vehicle control), 1 μM Latrunculin A (LatA) or 1 μM Nocodazole (Noco) for 60 min

(A) or 120 min (C) on DIV16, respectively. Changes in postsynaptic proteins GPN (red) and

PSD95 (green) were detected by confocal microscopy. Quantitative data shown were the

mean intensity (left) and cluster number (right) of GPN and PSD95 at 60 min (B) and 120

min (D). Ctrl: 49 dendrites from 13 neurons for 60 min; 49 dendrites from 12 neurons for 120

min; LatA: 45 dendrites from 11 neurons for 60 min; 48 dendrites from 15 neurons for 120

min; Noco: 49 dendrites from 15 neurons for 60 min; 51 dendrites from 12 neurons for 120

min. Scale bar, 5 μm.

(E-H) Same as (A-D) except that neurons were immunostained for GPN (red) and surface γ2

(green). Changes in GPN and surface γ2 at 60 min (E) or 120 min (G) were detected by

confocal microscopy. Quantitative data shown were the mean intensity (left) and cluster

number (right) of GPN and surface γ2 at 60 min (F) and 120 min (H). Ctrl: 51 dendrites from

15 neurons for 60 min; 37 dendrites from 12 neurons for 120 min; LatA: 51 dendrites from 15

neurons for 60 min; 45 dendrites from 15 neurons for 120 min; Noco: 49 dendrites from 15

neurons for 60 min; 36 dendrites from 11 neurons for 120 min. Scale bar, 5 μm.

(I) Levels of synaptic proteins normalized to control neurons following LatA or Noco

treatment in (A,C,E,G).

(J-K) The colocalization between surface γ2 and GPN signals in dendrites in (E) and (G).

Statistical test: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. All data are scatterplots with means ±

S.E.M.

1243

Figure 7. Both the membrane-binding and p140Cap-binding capacities of

endophilin A1 are required for iPSD assembly and E/I balance.

(A) Cultured hippocampal neurons from *EEN1*$^{fl/fl}$ mice were co-transfected with construct

expressing GFP and Flag vector or GFP-2A-Cre and Flag vector, Flag-EEN1 wild-type (WT),

mutants (KKK-EEE or Y343A) or p140Cap, and analyzed on DIV16 by immunofluorescence

staining for GPN (green) and GFP (red). Scale bar, 5 μm.

(B) Quantification of GPN mean intensity (left) and cluster number (right) in (A). 65

dendrites from 14 neurons for GFP vector; 67 dendrites from 13 neurons for Cre + vector; 48

dendrites from 13 neurons for Cre + WT; 50 dendrites from 15 neurons for Cre + KKK-EEE;

52 dendrites from 14 neurons for Cre + Y343A; 62 dendrites from 14 neurons for Cre +

p140Cap.

(C) Same as (A) except that neurons were immunostained for surface γ2 on DIV16. Scale bar,

5 μm.

(D) Quantification of surface γ2 mean intensity (left) and number (right) in (C). 40 dendrites

from 11 neurons for GFP + vector; 38 dendrites from 10 neurons for Cre + vector; 40

dendrites from 11 neurons for Cre + WT; 45 dendrites from 14 neurons for Cre + KKK-EEE;

35 dendrites from 11 neurons for Cre + Y343A.

(E) Experimental scheme for seizure scoring of *EEN1$^{fl/fl}$* mice stereotactically co-injected with

AAVs carrying mCherry and Cre, and EGFP and EEN1 or its mutants (KKK-EEE and

Y343A) in the hippocampal CA1 region. Mice were intraperitoneally administered PTZ for

12 days after AAV injections, then seizures were scored.

(F) PTZ-induced seizures in mice injected with the indicated AAVs were scored for 30 min.

Ctrl, 7 mice; Cre, 7 mice; Cre + WT, 7 mice; Cre + KKK-EEE, 6 mice; Cre + Y343A, 7 mice.

(G-K) AAVs expressing Cre (AAV-mCherry-2A-Cre) and GFP (AAV-EGFP) or Cre and

EEN1 WT or mutant (AAV-EGFP-2A-EEN1, AAV-EGFP-2A-EEN1 KKK-EEE,

AAV-EGFP-2A-EEN1 Y343A) were stereotaxically injected into the CA1 regions of *EEN1$^{fl/fl}$*

mouse brain at P0. Acute hippocampal slices were prepared on P14-P21 for dual recording of

evoked excitatory postsynaptic currents (eEPSC) and inhibitory postsynaptic currents (eIPSC).

Shown are representative traces and pairwise comparisons of noninfected (Ctrl) and infected

neurons in the same slice. Scale bar = 50/50, 50/50, 50/50, 50/50, 40/20 (pA/ms) in (G); Scale

bar = 100/100, 100/400, 100/100, 100/200, 300/100 (pA/ms) in (H); Scale bar = 50/50, 50/50,

50/50, 50/50, 50/50 (pA/ms) in (I); Scale bar = 100/100, 200/100, 100/100, 100/200, 100/200

(pA/ms) in (J). Statistical test: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; red color is for Cre *vs*

Ctrl, green color is for Cre + WT *vs* Cre; ns, not significant; All data are means ± S.E.M or

scatterplots with means ± S.E.M.

1279

Figure 8. Model for the role of endophilin A1 at the inhibitory postsynapses and E/I

balance.

(A) Endophilin A1 directly interacts with the inhibitory postsynaptic scaffold protein gephyrin

to promote organization of iPSD and synaptic recruitment/stabilization of the GABA$_A$Rs via

its plasma membrane association and actin polymerization-promoting activities. **(B)** Loss of

endophilin A1 impairs inhibitory synaptic neurotransmission and disrupts the balance of

excitatory/inhibitory activities in the neural circuits, leading to enhanced epileptic seizure

susceptibility.

1288

1289   **Figure S1. Loss of endophilin A1 *in vivo* does not affect fast mIPSCs (perisomatic events:**

1290   **< 2.8 ms) and slow mIPSCs (distal dendritic events: >2.8 ms). Related to Figure 1**

1291   **(A)** Quantification of somatic and dendritic amplitudes in Ctrl and *EEN1* KO neuron. n = 7

1292   WT and 9 *EEN1* KO neurons. Data are mean ± S.E.M.

(B) Frequency of somatic and dendritic events from Ctrl and *EEN1* KO mice. WT: 7 neurons; *EEN1* KO: 9 neurons. Data are mean ± S.E.M.

Figure S2. Effects of *EEN1* knockout on excitatory and inhibitory synapses in the hippocampus. Related to Figure 2

(A-B) Immunofluorescence analysis of PSD95 clusters in the hippocampal region of 3-week-old (P21) (left) and 10-week-old (P60) (right) *EEN1*$^{+/+}$ and *EEN1*$^{-/-}$ mouse brains. Scale bar, 50 μm.

(C) Quantification of the mean intensity of PSD95 in the hippocampal region of 3-week-old (P21) and 10-week-old (P60) *EEN1*$^{+/+}$ and *EEN1*$^{-/-}$ mouse brains. n = 3 mice for each genotype.

(D) Immunofluorescent signals of GPN in distinct layers of mouse hippocampus. Scale bar, 50 μm.

(E) Quantification of gephyrin (GPN) mean intensity in distinct layers of hippocampus. n = 6 mice for each genotype. Abbreviations: CA1-3, cornus Ammonis region 1-3; DG, dentate gyrus; slm, stratum lacunosum-moleculare; so, stratum oriens; sp, stratum pyramidale; sr, stratum radiatum. DAPI staining of cell nuclei is shown in blue. Statistical test: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; ns, non-significant. Data represent mean ± S.E.M.

Figure S3. Expression of GABA$_A$R α1 in cultured hippocampal neurons. Related to Figure 3

Cultured hippocampal neurons were fixed on DIV16 and immunostained for surface GABA$_A$R α1. Shown are representative confocal microscopy images. Scale bar, 10 μm.

Figure S4. Effect of *EEN1* knockout on gephyrin clustering in cultured hippocampal neurons. Related to Figure 4

(A-C) *EEN1*$^{fl/fl}$ hippocampal neurons were transfected with construct expressing mCherry or mCherry-2A-Cre on DIV12 and immunostained for GPN and mCherry on DIV16. Shown are representative confocal microscopy images for GPN in the soma, axon initial segment (AIS) and dendrites 30 μm, 60 μm, 90 μm or120 μm from the soma, respectively. Scale bars, 5 μm.

(D) Quantification of GPN mean intensity normalized to the mCherry group in the soma (A), AIS (B) and dendrites at various distances from the soma (C). Number: 20 (mCherry), 20 (Cre) for soma; 25 (mCherry), 25 (Cre) for AIS; 94 (mCherry), 76 (Cre) for dendrites 30 μm; 80 (mCherry), 51 (Cre) for dendrites 60 μm; 72 (mCherry), 61 (Cre) for dendrites 90 μm; 57 (mCherry), 41 (Cre) for dendrites 120 μm. Ankyrin-G: Axon initial segment (AIS) labeling marker. Statistical test: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. All data represent means ± S.E.M for each group.

Figure S5. p140Cap promotes gephyrin clustering via endophilin A1-mediated pathway. Related to Figure 6 and 7

(A) Cultured hippocampal neurons transiently expressing GFP and LifeAct-mCherry were immunostained for GPN, GFP and mCherry on DIV16. Shown are representative z stack maximum projection images captured by structured illumination microscopy (SIM). Right panels are insets of boxed regions in the left panel. Scale bars, 2 μm in left panels and 1 μm in magnified images.

(B) Cultured hippocampal neurons were co-transfected with Ctrl shRNA and vector or p140Cap shRNA and vector or RNAi-resistant Flag-tagged p140Cap on DIV 11, followed by immunostaining with antibodies to GPN, Flag (blue) and RFP on DIV 16. Scale bar, 5 μm.

(C) Quantification of GPN mean intensity and puncta number in (B). 82 dendrites from 18 neurons for shCtrl + vector; 92 dendrites from 20 neurons for shp140Cap + vector; 88 dendrites from 22 neurons for shp140Cap + p140Cap.

(D) $EEN1^{fl/fl}$ hippocampal neurons coexpressing GFP and Ctrl shRNA or p140Cap shRNA, or GFP-2A-Cre and Ctrl shRNA or p140Cap shRNA were immunostained for GPN on DIV16. Shown are representative confocal microscopy images. Scale bar, 5 μm.

(E) Quantification of GPN mean intensity and puncta number in (D). 59 dendrites from 14 neurons for GFP + shCtrl; 53 dendrites from 15 neurons for GFP + shp140Cap; 46 dendrites from 14 neurons for Cre + shCtrl; 63 dendrites from 17 neurons for Cre + shp140Cap. Statistical test: *** p < 0.001, ns, non-significant. All data shown are means ± S.E.M.

Table S1. Proteins identified by mass spec analysis of endophilin A1 co-immunoprecipitates and their abundance. Related to Figure 4

Video S1. PTZ kindling behaviors of $EEN1^{+/+}$ and $EEN1^{-/-}$ mice. Related to Figure 1

Video S2. The spatial relationship between EEN1 and GPN. **Related to Figure 3G**. 3D-SIM movie of an enlarged region of interest (ROI) from a hippocampal neuron dendrite shows the association of EEN1 (red) and GPN (green).

Video S3. The spatial relationship between endophilin A1 and surface GABA$_A$R γ2. **Related to Figure 3H**. 3D-SIM movie of an enlarged region of interest from a hippocampal neuron dendrite shows the association of EEN1 (red) and surface GABA$_A$R γ2 (green).

Video S4. The spatial relationship between endophilin A1 and VGAT. **Related to Figure 3I**. 3D-SIM movie of an enlarged region of interest from a hippocampal neuron dendrite shows the signals of EEN1 (red) and VGAT (green).

1369   **Video S5. The spatial relationship between gephyrin and LifeAct. Related to Figure 6**

1370   **and Figure S5A**. 3D-SIM movie of an enlarged region of interest from a hippocampal neuron

1371   dendrite shows the signals of LifeAct (red) and GPN (green).

1372

1373   **Video S6. PTZ kindling behaviors of *EEN1^{fl/fl}* mice injected with AAVs expressing Cre**

1374   **and EEN1 WT or mutants. Related to Figure 7**

1375

A

GABAergic terminal

Gephyrin
F-actin
microtubule
Endophilin A1
Neuroligin 2
GABA$_A$ receptor
Neurexin
p140cap

B

EEN1$^{+/+}$

E   I

EEN1$^{-/-}$

Epilepsy

E   I

Impaired inhibitory
neurotransmission