Discovery of novel quinoline papain-like protease inhibitors for COVID-19 through topology constrained molecular generative model

Yongzhi Lu[1#], Ting Ran[1#], Qi Yang[1#], Guihua Zhang[1#], Jiayi Chen[1,2], Peiqi Zhou[1], Wenqi Li[1,3], Minyuan Xu[1], Jielin Tang[1], Minxian Dai[1], Jinpeng Zhong[1], Hua Chen[1], Pan He[1], Anqi Zhou[1], Bao Xue[1], Jiyun Zhang[1,2], Kunzhong Wu[1,4], Xinyu Wu[1,5], Miru Tang[1], Xinwen Chen[1*], Hongming Chen[1*] and Jinsai Shang[1,6*]

Affiliations

[1]State Key Laboratory of Respiratory Disease, The First Affiliated Hospital of Guangzhou Medical University, School of Basic Medical Sciences, Guangzhou National Laboratory, Guangzhou Medical University, Guangzhou 511436, China

[2]School of Pharmaceutical Sciences, Sun Yat-Sen University, Guangzhou 510006, China

[3]College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan 430074, China

[4]State Key Laboratory of Medicinal Chemical Biology, College of Life Sciences, Nankai University, Tianjin 300071, China

[5]State Key Laboratory of Virology, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan, 430071, China

[6]Lead contact

[#] These authors contributed equally to this work

* Correspondence: Xinwen Chen; Hongming Chen; Jinsai Shang
Email: chen_xinwen@gzlab.ac.cn; chen_hongming@gzlab.ac.cn; shang_jinsai@gzlab.ac.cn

Abstract

The rapid emergence of drug-resistant SARS-CoV-2 variants poses a persistent challenge to current antiviral strategies. Mutations in key viral targets, including RNA-dependent RNA polymerase (RdRp), main protease ($3CL^{pro}$), and papain-like protease (PL^{pro}), have been shown to markedly reduce the efficacy of approved therapeutics, highlighting the urgent need for next-generation antivirals capable of overcoming resistance. Here, we report the discovery of a novel class of PL^{pro} inhibitors through an AIDD strategy based on a topology-constrained molecular generative model (Tree-Invent) integrated with structure-guided optimization. Scaffold hopping from a previously reported lead enabled the identification of a quinoline-based chemical series with substantially improved metabolic stability and antiviral potency. Structure-guided optimization yielded compound GZNL-2016, which exhibited potent enzymatic inhibition of PL^{pro} (IC_{50} = 10.5 nM), robust antiviral activity against multiple SARS-CoV-2 variants, including Omicron BA.5 and XBB.1, and favorable pharmacokinetic properties following oral administration. Notably, GZNL-2016 retained substantial inhibitory activity against the clinically relevant drug-resistant mutant PL^{pro} E167K (IC_{50} = 480.2 nM; K_i = 439.3 nM), in contrast to previously reported inhibitors that exhibit markedly reduced potency. In a SARS-CoV-2 infection mouse model, oral administration of GZNL-2016 significantly reduced pulmonary viral titers, demonstrating in vivo antiviral efficacy. Collectively, this study establishes an AI-enabled strategy for rapid antiviral discovery and identifies GZNL-2016 as a promising lead compound to address the threat of coronavirus infections caused by drug-resistant mutant SARS-CoV-2 variants.

Keywords: SARS-CoV-2, Papain-Like Protease, AIDD, DUBs, Drug Design.

48 **Graphical abstract**

49
50 We leveraged an AI generative model to discover a novel PL^{pro} inhibitor with excellent liver
51 stability, low CYP, hERG inhibition and reasonable oral PK properties. At the same time, the
52 compound **16** exhibits high efficacy for the resistance mutation E167K, which resulted in severe
53 resistance to the previously reported inhibitor Jun12682 and PF-07957472.

Introduction

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) is the infectious virus responsible for coronavirus disease 2019 (COVID-19). It was initially discovered in China in December 2019 and has rapidly evolved into a global epidemic with a significant cost to human society. Over the past five years, scientists worldwide have been exploring various approaches to treat infected individuals and reduce the severity of the disease. To date, several COVID-19 vaccines have been developed and granted emergency authorization for market distribution[1]. All these vaccines target the spike protein to prevent the virus from entering the host cells. However, the high mutation rate of the spike protein, especially within the receptor-binding domain (RBD), diminishes the long-term protective effectiveness of the vaccines[2]. This underscores the need to develop alternative therapeutic modalities, particularly small molecule antiviral drugs targeting more conserved viral proteins.

Currently, the FDA has approved three small-molecule drugs, including remdesivir, molnupiravir and nirmatrelvir[3]. They all target relatively conserved proteins across different SARS-CoV-2 variants. The first two drugs are viral RNA-dependent RNA polymerase (RdRp) inhibitors, while the third one is an inhibitor of the viral main protease (M^{pro}), also known as 3CL protease ($3CL^{pro}$). Remdesivir was the first drug authorized for emergency use, but its administration is limited to intravenous infusion for hospitalized patients with severe symptoms[4]. Its clinical efficacy has been found to be sub-optimal, as demonstrated by several clinical trials[5]. Similar to remdesivir, molnupiravir is a nucleotide analogue and functions as a mutagen, inhibiting SARS-CoV-2 replication by increasing the viral mutation rate[6]. However, there are concerns that molnupiravir may also induce similar mutations in the host[7]. Nirmatrelvir, combined with a cytochrome P450 3A (CYP3A) inhibitor (*i.e.,* ritonavir), is an oral medication and suitable for non-hospitalized patients[8]. Ensitrelvir, a M^{pro} inhibitor approved in Japan[9], has demonstrated high efficacy in clinical trials. However, it is also a potent CYP3A inhibitor, which may lead to severe adverse drug-drug interactions with other medications[10]. Therefore, new antivirals with alternative mechanisms are still urgently needed.

The papain-like protease (PL^{pro}) is one of the two viral proteases responsible for cleaving the polyproteins encoded by the SARS-COV-2 genome[11]. Additionally, it can remove the ISG-15 and ubiquitin modifications on host proteins[12]. This dual enzymatic function allows it to both promote virus replication and suppress the host's immune response. Moreover, PL^{pro} is a highly conserved protein across SARS-CoV-2 variants[13], making it a promising drug target for anti-COVID-19 therapy. The catalytic site of the protein is situated within a small buried pocket, adjacent to an induced substrate-binding cavity by the reshaping of the BL2 loop, a flexible beta turn/loop region (Gly266–Gly271)[14].

GRL0617 was originally discovered as a non-covalent anti-SARS-CoV PL^{pro} inhibitor binding at the induced substrate-binding cavity and later validated to have inhibitory potency against SARS-COV-2 PL^{pro} at the micromolar level[14]. Analogs of **GRL0617** were developed with moderate enzymatic and cellular activities, such as **Jun9-84-3**[15], **XR8-23**[16], **Jun11273**[17] and covalent inhibitors **CP7**[18]. Recently, a few compounds[19] (such as **Jun12862, PF-07957472,**

GZNL-P4) were reported to have in vivo efficacy in SARS-CoV-2 infected mice, along with oral
bio-availability. **GZNL-P4** is a potent compound identified in our previous study, but structural
optimization is needed to improve its low liver stability. The development of PL^{pro} inhibitors at
the clinical stage remains rare. To date, only one PL^{pro} inhibitor has been reported to have entered
clinical trials in China, and its structure has yet to be disclosed[20]. Therefore, the development of
PL^{pro} inhibitors remains necessary to explore their potential as a COVID-19 therapy.

The high mutation frequency of coronaviruses has given rise to a severe and increasingly
pressing challenge of drug resistance. During the COVID-19 pandemic, vaccines and antibody
drugs played a crucial role; however, later-emerging Omicron subvariants (e.g., BA.1, BA.2, BA.5,
BQ.1.1, XBB, XBB.1.5, etc.) exhibited high or complete neutralization escape against nearly all
authorized monoclonal antibodies, leading to their successive withdrawal from clinical use [21].
Similarly, mutations in Mpro (e.g., Q192R, H172Y, H172Y/Q189E, L50F/E166V, and
L50F/E166A/L167V) have been demonstrated to reduce susceptibility to nirmatrelvir and ritonavir,
with the decrease ranging from moderate (25–99-fold) to high (> 100-fold) [21-22]. *In vitro* studies
further indicate that multiple mutations in RdRp (including E802D, V166A, N198S, S759A,
V792I, C799F/R, etc.) can decrease sensitivity to remdesivir, with the magnitude of reduction
varying from several-fold to several hundred-fold [23]. Several key mutations in PL^{pro} have been
confirmed to significantly impair the efficacy of existing inhibitors, which constitutes a core
challenge for the development of PL^{pro} inhibitors. Tan et al. reported that residues E167, Y268,
and Q269 are mutation hotspots associated with resistance to existing PL^{pro} inhibitors [24].
Especially, the mutations of E167K and Y268N in PL^{pro} dramatically decrease the inhibitory
activity of Jun2682 and PF-02957472 with the Ki from 26.51 nM and 6.85 nM to greater than 10
μM and 8.036 μM, respectively [24]. Although no PL^{pro} inhibitor has reached clinical approval to
date, drug resistance remains a significant concern. Consequently, developing PL^{pro} inhibitors that
can overcome drug resistance is critical for future pandemics.

Artificial intelligence (AI) technology has shown great potential to accelerate drug
discovery[25]. Among these models, deep molecular generation models have garnered particular
attention due to their ability to autonomously produce novel molecular structures. Early molecular
generation models primarily involved random sampling from the learned continuous molecular
chemical space to generate a large number of molecular structures[26]. However, due to the vastness
of chemical space, the probability of discovering active molecules through sampling these models
is relatively low. In recent years, there has been an increasing focus on achieving goal-directed
molecular design through the development of conditional molecular generation models. Key
approaches include altering the form of model input[27], utilizing conditional deep neural networks
[28], applying transfer learning[29], and employing reinforcement learning[30], etc. Some of the latest
methods directly decode the learned three-dimensional structural features of protein pockets into
molecular chemical structures[31]. Although numerous methods have been developed, only a few
of them have been reported with application in real-world scenarios.

In earlier research, we developed Tree-Invent[32], a graph-based molecular generation model
with topology constraints. This model is versatile across different drug design scenarios and excels

in achieving precise control over structural topology during molecular generation. In this model, a
molecular graph is represented as a topological tree comprising ring nodes, non-ring nodes, and
edges connecting the nodes. Molecule generation is achieved by carrying out operations of node
addition, ring generation, and node connection in an autoregressive manner. Reinforcement
learning (RL)[33] is also integrated within the Tree-Invent model to facilitate the exploration of
chemical space toward desired properties. We applied this method to conduct scaffold hopping
exercise and discovered a novel quinolone chemical series that exhibits nanomolar potency in
enzymatic assay and excellent anti-viral potency in cellular experiments. Our best compound
exhibits nanomolar level inhibition activity to PL^{pro} E167K and demonstrates reasonable in vitro
and in vivo oral PK properties. The significant selectivity over normal human cells and weak
inhibition on the CYP and hERG proteins indicate its promising safety profile for antiviral
treatments.
151

Results

Scaffold hopping through topology constrained molecular generation with Tree-Invent

So far, exploration of the structure-activity relationship (SAR) based on the structure of **GRL-**
0617 has resulted in several potent PL^{pro} inhibitors (Fig. S1A). In current study, we utilized the
Tree-Invent model (Fig. S2) to design novel PL^{pro} inhibitors, using the crystal structure of PL^{pro}
enzymatic domain in complex with known inhibitor GRL0617 (PDB ID: 7JRN), which exhibits
potency at micromolar range (Fig. S1B), where the naphthalene ring serves as the head group, is
located at the BL2 groove sandwiched between the aromatic residue Y269 and the hydrophobic
residues P247 and P248. The aniline group in the tail group is located at the P3 sub-pocket and
extends into the P2 sub-pocket near L162 through a methyl group. The linker group is located at
the P4 sub-pocket, where the hydrogen bonds with D164 and Q269 are formed by the amide group.
Accordingly, the structure of **GRL0617** can be divided into three parts, namely the head, linker
and tail part, corresponding to three sub-pockets in the substrate binding site (Fig. S1C) [14].

In our previous work, we identified a potent compound, **GZNL-P4**, which features a tricyclic
dihydroacenaphthylene (DHAN) scaffold. The co-crystal structure of **GZNL-P4** with PL^{pro} (PDB
ID: 8YX2) reveals that its tail group extends into the solvent-exposed region at the P3 sub-pocket.
The positively charged piperazine nitrogen in this group can form salt bridge with E167, an
interaction absent in the **GRL0617** complex structure (Fig. 1A). However, **GZNL-P4** has poor
liver metabolic stability (rat liver microsome stability t$_{1/2}$ 7.95 min), which is believed to be
associated with the head DHAN group, as observed in the metabolism of **GRL0617** [34], so finding
other head groups is needed to address the stability issue.

Herein, we employed Tree-Invent model to explore the chemical space of head group. The
structure of **GZNL-P4** was transformed to a topology tree with four nodes connected by single
bonds (Fig. 1B). To focus on the generation of the head group, node N1 was predefined as the
substructure comprising the linker and tail part of **GZNL-P4** and kept unchanged in the process
of molecule generation. Node N2 was designated as a changeable node grown from node A to
replace the head group of **GZNL-P4**. Analysis of the binding modes of **GZNL-P4** and **GRL0617**

indicates that the aromatic ring system of the head group is important for binding through π-π
stacking interaction with Y268 and hydrophobic interaction with P247 and P248 [14]. To mimic
this binding mode, node N2 was confined to contain 5,6- or 6,6-fused bicyclic systems with at
least one aromatic ring. Furthermore, two nodes, *i.e.,* N3 and N4, were additionally appended to
node N2 as R-group decorations. However, their generation was non-essential and entirely
contingent upon the model's decision regarding node addition. With these topological constraints,
Tree-Invent was implemented with RL to generate molecules (Fig. 1C), and two rounds of RL
were conducted for the generation of 5,6- and 6,6-fused bicyclic molecules. The crystal structure
of **GRL0617** was chosen as the protein model for carrying out docking during the generative
process and the docking score was transformed into reward score during the RL process, aiming
to generate molecules with high affinity.

Through 1000 epochs of RL, the learning curves of the Tree-Invent model converge (Fig.
S3A), suggesting the model has learned how to generate high-score molecules with 5,6- and 6,6-
fused bicyclic systems. Combining the generation sets of the two RL runs, 2817 unique molecules
were obtained. Among them, 1270 molecules have 5,6-fused ring structure and 1547 molecules
have 6,6-fused ring structure (Fig. S3B). Moreover, the substructures corresponding to node N1 in
these molecules are kept the same as in **GZNL-P4**. Half of the molecules have a naked ring
structure in the head group, while the others have no more than two substitutes on the ring structure
as expected (Fig. S3C). This demonstrates that the Tree-Invent model is capable of generating
molecules satisfying the topological constraints. It was also found that the naphthalene group of
GRL0617 can be reproduced which proves the capability of this method in identifying active
molecules. Subsequently, all generated molecules were redocked to the binding pocket using the
Glide docking program under the standard precision mode (Glide-SP)[35], resulting to 367
molecules with similar binding mode to **GZNL-P4** and **GRL0617** and docking scores lower than
-7.0 kcal/mol, a cutoff to distinguish suitable molecules from all generated molecules (Fig. S3D).
Moreover, these remaining molecules are all compliant with the Lipinski rules[36]. Chemical space
analysis based on the PCA analysis [37] using the Morgan fingerprint [38] reveals that these
molecules can be grouped into several categories, and each cluster has some molecules with good
docking scores (Fig. 1D). The chemical space of molecules filtered based on docking scores was
displayed using tSNE plot [39] (Fig. 1D).

Identification of novel scaffold of PL^{pro} inhibitors

Considering synthesis feasibility, nine molecules were chosen for synthesis. The synthetic
route was illustrated as Scheme 1 in the supplemental text. The structures of the synthesized
compounds are shown in Fig. 2A. Their docking poses can be found in Figure S4. Then, they were
tested using a fluorescence-based biochemical assay[40] to evaluate their inhibitory activities on
PL^{pro} (Fig. 2B). Compounds **1** and **2**, containing a quinoline ring in the head group, are the most
active compounds. Their aromatic nitrogen atoms in the quinoline ring are situated at the para-
position to the linker substituent. Chlorination on the quinoline ring in compound **1** shows a minor
impact on the activity, while compound **2** is suggested to have better binding affinity (Fig. 2C).
The binding affinity of compound **2** with PL^{pro} measured by isothermal titration calorimetry (ITC)

and biolayer interferometry (BLI) are 285 nM and 187 nM (Fig. S5A, S5C), respectively.
Compound **6** with benzothiophene exhibits somewhat lower activity than compound **2**. Other
compounds having 5,6 ring systems such as indole, benzopyrazole and 1H-pyrrolopyridine have
much worse activity, regardless of whether the head group is a five-membered or six-membered
ring. Among them, compound **8** with azaindole is the weakest. N-methyl substitution on the indole
ring of compound **9** improves the activity, probably because it makes additional van der Waal
interaction with the residue P248. Compound **5** with furopyridine has a special substitution
position on the head group, indicating that the direction of substitution on the head group is very
important for activity. Through the AI-driven R-group generation process, we successfully
identified several PL^{pro} inhibitors with new scaffolds having reasonable potency. The liver
metabolic stability of compound **2** was measured, and to our delight, its half-life in liver microsome
has been largely improved compared to **GZNL-P4** (Fig. 5E).

To better understand the binding mode of compound **2**, we resolved its co-crystal structure in
complex with PL^{pro} (Fig. 3A). The residues surrounding the binding pocket maintain nearly
identical conformations to those observed in the co-crystal structure of **GRL0617**, and there is a
good alignment between the binding poses of the two compounds (Fig. 3E). The binding mode
analysis shows that the quinoline ring of compound **2** is overlapped with the head group of
GRL0617. As expected, the quinoline ring makes π-π stacking interaction with Y268, and extends
the interaction with P247 and P248. The nitrogen atom on the quinoline ring is exposed to solvent,
making no obvious binding with protein residues. The cyclopropyl group binds at the bottom of
the pocket surrounding Y264 and T301, and the amide linker group makes hydrogen bonds with
D164, Y268 and Q269. The tail group, i.e., piperazine-substituted benzyl group, is exposed to
solvent, and the positively charged piperazine nitrogen in this group can form a salt bridge with
E167, an interaction absent in the GRL0617 complex structure, while the benzyl group extends
into the sub-pocket near L162 through a methyl group. This binding mode is consistent with that
of **GZNL-P4**, which provides clear guidance for structural optimization.

Structural optimization of compound 2

To further improve the potency of compound **2**, we explored substituents at the R^2 position
to optimize its interaction with E167. We selected commercially available amine-containing
reagents to replace the N-methylpiperazine group of compound **2** and designed a virtual library of
2347 compounds containing positively charged amine groups (Fig. 2D). Molecular docking using
the co-crystal structure of compound **2** as docking model was then carried out, compounds whose
docking pose is similar to the binding pose of compound **2** and simultaneously maintains the salt
bridge with E167 in an appropriate distance (the cutoff is set to 4.0 Å) were selected. At last, 29
top-ranked compounds (by Glide docking score) were selected to synthesize (Scheme 2 in the
supplemental text) and their structures are listed in Table 1. They can be divided into two types in
terms of the ring count in the R^2 group.

Following SAR analysis reveals that the R^2 group greatly influences the inhibitory activity of
PL^{pro} (Table 1 and Fig. S6). Several compounds show superior activity to compound **2**, and three
compounds demonstrate IC$_{50}$ values lower than 0.015 μM. Among type I molecules with a single

ring in the R^2 group, compound **10** with piperazine substitute has worse potency than compound
2 having a methylpiperazine group, which means that the activity is sensitive to the position of
methyl substituent on the piperazine ring. Compound **2** with 4-methylpiperazine exhibits much
higher activity than compounds **14** and **15** which contain the substitutes of 2- and 3-
methylpiperazine. The compounds with hydroxyalkyl groups substituted on the 4-position of
piperazine have slightly worse activity than compound **2**. However, substituting with
dimethylaminoalkyl groups, as seen in compound **12**, negatively impacts activity. N-sulfonylation
of piperazine significantly reduces activity, likely due to the disruption of the salt bridge formed
between piperazine and the acid group in E167. It is worthwhile noting that the bridged piperazine
significantly enhances activity. Compound **16** (also named as GZNL-2016) is the most potent in
this series with IC_{50} of 10.5 nM. Compounds bearing similar cycloamine groups, including the
ones with spiral and fused rings, also demonstrate reasonable activity, although they don't show
improvement compared to compound **2**. These groups can mimic the salt bridge interaction of
piperazine. Nevertheless, the introduction of seven-member-ring results in a reduction of activity,
which may be due to the sub-optimal salt bridge interaction of the secondary amine in the enlarged
ring. Compared to compound **2** with a Kd of 285 nM (Fig. S5A), compound **16** exhibits higher
binding affinity with PL^{pro}. The measured Kd of compound **16** by isothermal titration calorimetry
(ITC) and biolayer interferometry (BLI) are 68.6 nM and 78.9 nM (Fig. S5B, S5D), respectively.

Type II molecules contain the R^2 substituents involving two cycloamine groups linked by a
single bond. Interestingly, the majority of compounds exhibited comparable activity to compound
2, except for compounds **35** and **36**. The reduced activity of compound **35** is attributed to the N-
methyl-piperazine substituent, which may form a sub-optimal salt bridge with E167. The weak
activity observed for compound **36** indicates that piperidine is worse than piperazine for the
substitution on the benzene ring, as evidenced by the decrease in the activity of compound **20**. On
the other hand, compounds featuring azetidines substituted with pentacyclic/hexacyclic amines
demonstrated the highest activity within the sub-group. Notably, compounds **30** (also named as
GZNL-2030) and **31** exhibit potency comparable to compound **16**, making them also promising
candidates for further profiling. Corresponding to the activity, the binding affinity of compound
30 with PL^{pro} measured by biolayer interferometry (BLI) is 447 nM (Fig. S5E). Most compounds
among **10-36** can improve the thermal stability of PL^{pro} with a delta Tm larger than 3 ℃ (Fig. S5F).
These compounds with larger delta Tm exhibit better potency for PL^{pro} (Fig. S5G).
Correspondingly, compound **16** exhibits the largest delta Tm (Fig. S5F), which is consistent with
its most potent inhibitory activity against PL^{pro} (Table 1).

The binding poses of the most potent compounds from the crystal structures (compounds **16**, **30**)
and docking (compound **31**) are consistent with that of compound **2**, which is characterized by the
sandwiched conformation of the quinoline ring, the hydrogen bonds with D164, Y268 and Q269,
and the salt-bridge with E167 (Fig. 3A-D). Q269 formed a hydrogen bond with compound **2**, but
a π-π stacking interaction with compound **16** (Fig. 3B). In the binding pose of compound **16**, the
hydrogen bond network composed of D164, E167 and compound **16** mediated by two water
molecules W1 and W2 may lower energy status, which may explain the superior performance of
compound **16** to other compounds.

The enzymatic inhibition of compound 16 on the PL^{pro} mutations

Mutations at residues E167 and Y268 of PL^{pro} lead to a dramatic decrease in sensitivity to the existing PL^{pro} inhibitors Jun12682 and PF-07957472, with a nearly 1000-fold increase in Ki [24]. To assess the susceptibility of compound **16** to distinct PLpro mutations, we evaluated its inhibitory activity against a panel of PLpro mutants, including E167K, E167S, Q269H, Y268N, Y268H, and the double mutant E167G/Q269H (Fig. 4A, S7). The results demonstrated that compound **16** essentially abolished its inhibitory activity against Y268N, while its potency was also significantly diminished against E167S and E167G/Q269H double mutant with IC$_{50}$ of 4.421 μM and 6.883 μM, respectively. Encouragingly, compound **16** retained relatively high inhibitory activity against E167K, Q269H, and Y268H, with IC$_{50}$ values of 480.2 nM, 357.7 nM, and 570.9 nM, respectively. For compound **16**, the calculated Ki for PL^{pro} E167K using the Cheng-Prusoff equation [41] was 439.3 nM. In stark contrast, both Jun12682 and PF-07957472 exhibited dramatically declined inhibition activity with Ki around 10 μM against E167K [24]. The binding affinity Kd to PL^{pro} E167K determined using ITC was 256 nM, 1.33 μM, and 1.02 μM, respectively (Fig. 4B-E). This may account for the reason why Compound **16** still retains favorable inhibitory activity against E167K.

Antiviral activity, liver stability and PK profiling of compound 16

The antiviral activities of several potent compounds were measured using a cell-based assay. The antiviral potency with 50% effective concentration (EC$_{50}$) of compound **2** against wild-type SARS-CoV-2 and two epidemic variants Omicron BA.5 and XBB.1 in infected Vero E6 cells are 0.4954 ± 0.1486 μM, 0.6199 ± 0.1061 μM and 0.6159 ± 0.0345 μM (Fig. 5A-C). At the same time, compound **2** shows rather low toxicity to the normal human cell line HEK293T (CC$_{50}$ > 200 μM) (Fig. 5D). Structural optimization on compound **2** resulted in three more potent compounds, compound **16**, **30** and **31**, which have enhanced anti-viral activity and a large enough safety window (CC$_{50}$/EC$_{50}$ for all variants: >379, >292, and >348 folds for compounds **16**, **30**, and **31**, respectively). Notably, compound **16** has the best antiviral activity compared to compounds **30** and **31** on all three tested virus variants (Fig. 5A-C). And, the variant of Omic-BA.5 is the least sensitive to all three PL^{pro} inhibitors.

Compared to the previously reported compound **GZNL-P4** (human and rat liver microsome stability t$_{1/2}$ of 3.8 min and 7.95 min, respectively[19b]), the liver stability of the Quinoline series significantly improved the metabolism stability in both rat and human liver microsomes (Fig. 5E). In rat liver microsomal stability assay, compound **16** displayed a prolonged half-life (t$_{1/2}$ = 120.12 min) and a low CLint value (20.77 mL/min/kg), which were 2.66-fold, 2.01-fold, 2.86-fold longer in t$_{1/2}$ and 2.66-fold, 2.01-fold, 2.86-fold lower in CLint than those of compound **2** (t$_{1/2}$ = 45.20 min, CLint = 55.20 mL/min/kg), compound **30** (t$_{1/2}$ = 59.70 min, CLint = 41.80 mL/min/kg), and compound **31** (t$_{1/2}$ = 41.98 min, CLint = 59.40 mL/min/kg), respectively. In human liver microsomal stability assay, compound **16** maintained favorable metabolic performance compared with compounds **2** and **31,** but slightly worse than compound **30**.

In vivo PK studies were performed in mice via oral gavage at a single dose of 20 mg/kg, and
the key PK parameters were quantified and compared among compounds **16**, **30**, and **31** (Fig. 5F).
Compound **16** exhibited significant superiority in systemic exposure, as evidenced by its area
under the concentration-time curve from time 0 to the last measurable concentration (AUC_{0-t}) was
1862.128 h·ng/mL. This value was 19.6% and 52.8% higher than that of compound **30** (AUC_{0-t} =
1556.770 h·ng/mL) and **31** (AUC_{0-t} = 1219.430 h·ng/mL), respectively, indicating significantly
enhanced oral bioavailability. For the elimination half-life ($T_{1/2}$), compound **16** exhibited a
moderate value of 0.757 h, which was comparable to those of compound **30** ($T_{1/2}$ = 0.699 h) and
31 ($T_{1/2}$ = 0.930 h). The time to reach peak plasma concentration (Tmax) of compound **16** was
0.667 h, slightly longer than that of compound **30** (Tmax = 0.417 h) and **31** (Tmax = 0.333 h).
However, it still ensured rapid oral absorption and subsequent onset of pharmacological effects.
The peak plasma concentration (Cmax) of compound **16** was 750.218 ng/mL, which was lower
than that of compound **30** (Cmax = 1095.919 ng/mL) and **31** (Cmax = 1245.280 ng/mL) but still
maintained a sufficiently high effective plasma concentration level to elicit the desired therapeutic
effects.
In general, compound **16** has the best inhibition activity against the enzyme and virus as well
as excellent liver metabolic stability. Based on these favorable properties, compound **16** was
selected as the lead compound for further profiling. The CYP and hERG inhibition at 10 μM and
30 μM for compound **16** were summarized in Figure 5F, the results indicate this compound has a
low risk of inhibiting selected CYP isoforms and hERG channel. All of these favorable
characteristics indicate that compound **16** is a promising lead compound suitable for further
optimization.

***In vivo* antiviral efficacy of compound 16**

To assess the *in vivo* anti-viral activity of compound **16**, we treated the model mice infected
with SARS-CoV-2 EG.5 by oral administration (Fig. 6A). K18-hACE2 transgenic mice aged 8
weeks were used as our mouse model. Thirty-two female hACE2 transgenic mice were divided
into four groups with eight mice in each group to evaluate the efficacy of mock, vehicle, positive
control PF-07321332 of 500 milligrams per kilogram (mpk), and compound **16** of 25 mpk in the
therapeutic treatment. The weight loss plot shows that the weight loss is less than 5% for all groups
(Fig. 6B). The lung live viral titers for the group treated with PF-07321332 at 500 mpk and
compound **16** at 25 mpk significantly decreased at 3 days post-infection (Fig. 6C).

Conclusion and Discussion

The continuous mutation of SARS-CoV-2 enables the virus to escape neutralization by
antibody therapies[21]. Furthermore, resistance to small-molecule drugs is emerging due to
mutations in key viral targets such as RdRp, 3CLpro, and PLpro, posing a major challenge to
pandemic control. The mutation of 3CL[pro] at E166V dramatically decreased the susceptibility to
Nirmatrelvir (IC_{50} increased 100-fold) [42]. Gandhi et al. reported a De novo emergence of a
remdesivir resistance mutation during treatment [23a]. Although no PL[pro] inhibitors are clinically
available, a number of PL[pro] inhibitors have been reported, such as Jun12682, PF-07957472 [19a,

[19c] and GZNL-P36 [19b]. In addition, drug-resistant mutations of PL^{pro} have also been documented,
among which E167 and Y268 are two key drug-resistant mutation sites. In particular, E167G,
E167A, E167S, E167K and Y268N confer strong drug resistance to Jun12682 and PF-07957472.
Specifically, the Ki value of PF-07957472 increases by more than 130-fold against E167K and
969-fold against Y268N; the Ki value of Jun12682 increases by more than 130-fold against E167K
and 385-fold against Y268N. Thus, the drug development targeting the E167 and Y268 mutations
of PL^{pro} will help address the potential drug resistance that may arise in the future.

In the current study, we leveraged an AI generative model to discover a novel series of PL^{pro}
inhibitors targeting the SARS-CoV-2 virus. Using the generative model, Tree-Invent, we
successfully identified a novel scaffold of PL^{pro} inhibitors exhibiting comparable activity to a
known reference compound **GZNL-P4**, though it has much improved liver metabolic stability.
Further optimization work on this series results in a lead compound **16** with an improved enzyme
potency and cellular-level antiviral activity. This compound demonstrates excellent liver stability,
low CYP and hERG inhibition and it also demonstrates reasonable oral PK properties. At the same
time, unlike the previously reported PL^{pro} inhibitors Jun12682 and PF-07957472 [24], compound
16 exhibits strong activity against E167K and can effectively overcome the drug resistance caused
by the E167K mutation. We believe that this study offers valuable insights for the future
development of PL^{pro} inhibitors as a potential treatment for COVID-19 disease, especially provides
a scaffold compound and points the way forward for overcoming E167K-mediated resistance.

Methods and Materials

397 Detailed descriptions of the in vitro pharmacology studies, *in vivo* pharmacology studies,
398 transcriptomics studies, X-ray crystallography, computational study, and synthetic methods can be
399 found in Supplementary Information.

Data analysis

401 The data in the figures represent mean ± SD. All data were analyzed using GraphPad Prism
402 9.0 software. Statistical comparison between different groups was performed using the
403 corresponding statistical analysis labeled in the figure legends combining several experiments. The
404 binding mode of ligands with protein was illustrated by PyMOL2.6.0a0 (open source).

Data Availability

406 Crystal structures generated during the current study are available in the Protein Data Bank (PDB)
407 under accession codes 8ZSE (PL^{pro} bound to compound **2**, here renamed as GZNL-2002), 21MS
408 (PL^{pro} bound to compound **16**, here renamed as GZNL-2016), and 21MU (PL^{pro} bound to
409 compound **30**, here renamed as GZNL-2030). The supplementary note for synthetic compounds
410 can be found in Supplementary Information.

Acknowledgments

412 This work is supported by the fundings: Overseas Experts Supporting Programs under
413 National Research Platform (WGZJ22-001); National Natural Science Foundation of China
414 (82170473); The Major Program of Guangzhou National Laboratory (GZNL2024A01005,

GZNL2023A01008); Guangdong Natural Science Foundation (2021QN020451, 2021CX020227, 2024A1515011589); Basic and Applied Basic Research Projects of Guangzhou Basic Research Program (2023A04J0161). We thank the staff at Guangzhou Laboratory Core Facility and Animal Center. We also thank the staff at BL10U2/BL02U1/BL17B1/BL18U1/BL19U1 beamlines at Shanghai Synchrotron Radiation Facility (SSRF) of the National Facility for Protein Science in Shanghai (NFPS), Shanghai Advanced Research Institute, Chinese Academy of Sciences, for providing technical support in X-ray diffraction data collection and analysis.

Author Contributions

J.S, Y.L., Q.Y., W.L., M.D., J.T., J.C., J.Z., K.W., and X.W. performed the cellular and biochemical assays. Y.L. and J.S. performed crystallography. G.Z., P.Z., H.C., P.H., and T.R. performed the medicinal chemistry. T.R., M.X., M.T., and C.H. performed the computational studies. Y.L., W.L., M.D., J.C., J.Z., K.W., and X.W. performed cloning, protein purification and biophysical analysis. Y.L., Q.Y., T.R., G.Z., X.C., C.H., and J.S. conceived the experiments and wrote the manuscript with input from all authors.

Conflict of Interest

The authors declare no competing interests.

References

[1] M. Li, H. Wang, L. Tian, Z. Pang, Q. Yang, T. Huang, J. Fan, L. Song, Y. Tong, H. Fan, *Signal transduction and targeted therapy* **2022**, *7*, 146.

[2] W. T. Harvey, A. M. Carabelli, B. Jackson, R. K. Gupta, E. C. Thomson, E. M. Harrison, C. Ludden, R. Reeve, A. Rambaut, S. J. Peacock, D. L. Robertson, *Nature reviews. Microbiology* **2021**, *19*, 409-424.

[3] G. Li, R. Hilgenfeld, R. Whitley, E. De Clercq, *Nature reviews. Drug discovery* **2023**, *22*, 449-475.

[4] aA. Amstutz, B. Speich, F. Mentré, C. S. Rueegg, D. Belhadi, L. Assoumou, C. Burdet, S. Murthy, L. E. Dodd, Y. Wang, K. A. O. Tikkinen, F. Ader, M. Hites, M. Bouscambert, M. A. Trabaud, M. Fralick, T. C. Lee, R. Pinto, A. Barratt-Due, F. Lund-Johansen, F. Müller, O. P. O. Nevalainen, B. Cao, T. Bonnett, A. Griessbach, A. Taji Heravi, C. Schönenberger, P. Janiaud, L. Werlen, S. Aghlmandi, S. Schandelmaier, Y. Yazdanpanah, D. Costagliola, I. C. Olsen, M. Briel, *The Lancet. Respiratory medicine* **2023**, *11*, 453-464; bR. L. Gottlieb, C. E. Vaca, R. Paredes, J. Mera, B. J. Webb, G. Perez, G. Oguchi, P. Ryan, B. U. Nielsen, M. Brown, A. Hidalgo, Y. Sachdeva, S. Mittal, O. Osiyemi, J. Skarbinski, K. Juneja, R. H. Hyland, A. Osinusi, S. Chen, G. Camus, M. Abdelghany, S. Davies, N. Behenna-Renton, F. Duff, F. M. Marty, M. J. Katz, A. A. Ginde, S. M. Brown, J. T. Schiffer, J. A. Hill, *The New England journal of medicine* **2022**, *386*, 305-315.

[5] M. Yasir, C. R. Lankala, P. Kalyankar, A. Ishak, M. Mekhail, C. Sestacovschi, E. Kima, *Cureus* **2023**, *15*, e43060.

[6] F. Kabinger, C. Stiller, J. Schmitzová, C. Dienemann, G. Kokic, H. S. Hillen, C.
Höbartner, P. Cramer, *Nature structural & molecular biology* **2021**, *28*, 740-746.
[7] M. D. Waters, S. Warren, C. Hughes, P. Lewis, F. Zhang, *Environmental and molecular*
mutagenesis **2022**, *63*, 37-63.
[8] D. R. Owen, C. M. N. Allerton, A. S. Anderson, L. Aschenbrenner, M. Avery, S. Berritt,
B. Boras, R. D. Cardin, A. Carlo, K. J. Coffman, A. Dantonio, L. Di, H. Eng, R. Ferre, K.
S. Gajiwala, S. A. Gibson, S. E. Greasley, B. L. Hurst, E. P. Kadar, A. S. Kalgutkar, J. C.
Lee, J. Lee, W. Liu, S. W. Mason, S. Noell, J. J. Novak, R. S. Obach, K. Ogilvie, N. C.
Patel, M. Pettersson, D. K. Rai, M. R. Reese, M. F. Sammons, J. G. Sathish, R. S. P.
Singh, C. M. Steppan, A. E. Stewart, J. B. Tuttle, L. Updyke, P. R. Verhoest, L. Wei, Q.
Yang, Y. Zhu, *Science (New York, N.Y.)* **2021**, *374*, 1586-1593.
[9] Y. Unoh, S. Uehara, K. Nakahara, H. Nobori, Y. Yamatsu, S. Yamamoto, Y. Maruyama,
Y. Taoda, K. Kasamatsu, T. Suto, K. Kouki, A. Nakahashi, S. Kawashima, T. Sanaki, S.
Toba, K. Uemura, T. Mizutare, S. Ando, M. Sasaki, Y. Orba, H. Sawa, A. Sato, T. Sato,
T. Kato, Y. Tachibana, *Journal of medicinal chemistry* **2022**, *65*, 6499-6512.
[10] R. Shimizu, T. Matsuzaki, R. Oka, T. Sonoyama, T. Fukuhara, A. Kuwata, Y. Matsuo, R.
Kubota, *Journal of clinical pharmacology* **2023**, *63*, 918-927.
[11] B. H. Harcourt, D. Jukneliene, A. Kanjanahaluethai, J. Bechill, K. M. Severson, C. M.
Smith, P. A. Rota, S. C. Baker, *Journal of virology* **2004**, *78*, 13600-13612.
[12] D. Shin, R. Mukherjee, D. Grewe, D. Bojkova, K. Baek, A. Bhattacharya, L. Schulz, M.
Widera, A. R. Mehdipour, G. Tascher, P. P. Geurink, A. Wilhelm, G. J. van der Heden
van Noort, H. Ovaa, S. Müller, K. P. Knobeloch, K. Rajalingam, B. A. Schulman, J.
Cinatl, G. Hummer, S. Ciesek, I. Dikic, *Nature* **2020**, *587*, 657-662.
[13] S. Kandwal, D. Fayne, *Virology* **2023**, *581*, 97-115.
[14] J. Osipiuk, S. A. Azizi, S. Dvorkin, M. Endres, R. Jedrzejczak, K. A. Jones, S. Kang, R.
S. Kathayat, Y. Kim, V. G. Lisnyak, S. L. Maki, V. Nicolaescu, C. A. Taylor, C. Tesar,
Y. A. Zhang, Z. Zhou, G. Randall, K. Michalska, S. A. Snyder, B. C. Dickinson, A.
Joachimiak, *Nature communications* **2021**, *12*, 743.
[15] C. Ma, M. D. Sacco, Z. Xia, G. Lambrinidis, J. A. Townsend, Y. Hu, X. Meng, T. Szeto,
M. Ba, X. Zhang, M. Gongora, F. Zhang, M. T. Marty, Y. Xiang, A. Kolocouris, Y.
Chen, J. Wang, *ACS central science* **2021**, *7*, 1245-1260.
[16] Z. Shen, K. Ratia, L. Cooper, D. Kong, H. Lee, Y. Kwon, Y. Li, S. Alqarni, F. Huang, O.
Dubrovskyi, L. Rong, G. R. J. Thatcher, R. Xiong, *Journal of medicinal chemistry* **2022**,
65, 2940-2955.
[17] P. Jadhav, B. Huang, J. Osipiuk, X. Zhang, H. Tan, C. Tesar, M. Endres, R. Jedrzejczak,
B. Tan, X. Deng, A. Joachimiak, J. Cai, J. Wang, *European journal of medicinal*
chemistry **2024**, *264*, 116011.
[18] B. C. Sanders, S. Pokhrel, A. D. Labbe, Mathews, II, C. J. Cooper, R. B. Davidson, G.
Phillips, K. L. Weiss, Q. Zhang, H. O'Neill, M. Kaur, J. G. Schmidt, W. Reichard, S.
Surendranathan, J. Parvathareddy, L. Phillips, C. Rainville, D. E. Sterner, D. Kumaran,
B. Andi, G. Babnigg, N. W. Moriarty, P. D. Adams, A. Joachimiak, B. L. Hurst, S.
Kumar, T. R. Butt, C. B. Jonsson, L. Ferrins, S. Wakatsuki, S. Galanie, M. S. Head, J. M.
Parks, *Nature communications* **2023**, *14*, 1733.
[19] aR. M. Garnsey MR, Nguyen LT, Cardin R, Tillotson J, Mashalidis E, Yu A,
Aschenbrenner L, Balesano A, Behzadi A, Boras B, Chang JS, Eng H, Ephron A, Foley
T, Ford KK, Frick JM, Gibson S, Hao L, Hurst B, Kalgutkar AS, Korczynska M,

Lengyel-Zhand Z, Gao L, Meredith HR, Patel NC, Polivkova J, Rai D, Rose CR, Rothan H, Sakata SK, Vargo TR, Qi W, Wu H, Liu Y, Yurgelonis I, Zhang J, Zhu Y, Zhang L, Lee AA. , *Sci. Adv.* **2024**, *10*, eado4288; bY. Lu, Q. Yang, T. Ran, G. Zhang, W. Li, P. Zhou, J. Tang, M. Dai, J. Zhong, H. Chen, P. He, A. Zhou, B. Xue, J. Chen, J. Zhang, S. Yang, K. Wu, X. Wu, M. Tang, W. K. Zhang, D. Guo, X. Chen, H. Chen, J. Shang, *Nature communications* **2024**, *15*, 10169; cB. Tan, X. Zhang, A. Ansari, P. Jadhav, H. Tan, K. Li, A. Chopra, A. Ford, X. Chi, F. X. Ruiz, E. Arnold, X. Deng, J. Wang, *Science (New York, N.Y.)* **2024**, *383*, 1434-1440.

[20]   W. He, L. Liu, B. Li, L. Xu, H. Li, J. Zheng, S. Wan, C. Wu, T. Wang, M. Wang, China, **2022**.

[21]   M. Vukovikj, A. Melidou, P. Nannapaneni, T. Normark, A. Kraus, E. K. Broberg, *Euro Surveill* **2025**, *30*.

[22]   E. Heilmann, F. Costacurta, S. A. Moghadasi, C. Ye, M. Pavan, D. Bassani, A. Volland, C. Ascher, A. K. H. Weiss, D. Bante, R. S. Harris, S. Moro, B. Rupp, L. Martinez-Sobrido, D. von Laer, *Sci Transl Med* **2023**, *15*, eabq7360.

[23]   aS. Gandhi, J. Klein, A. J. Robertson, M. A. Peña-Hernández, M. J. Lin, P. Roychoudhury, P. Lu, J. Fournier, D. Ferguson, S. A. K. Mohamed Bakhash, M. Catherine Muenker, A. Srivathsan, E. A. Wunder, Jr., N. Kerantzas, W. Wang, B. Lindenbach, A. Pyle, C. B. Wilen, O. Ogbuagu, A. L. Greninger, A. Iwasaki, W. L. Schulz, A. I. Ko, *Nature communications* **2022**, *13*, 1547; bL. J. Stevens, A. J. Pruijssers, H. W. Lee, C. J. Gordon, E. P. Tchesnokov, J. Gribble, A. S. George, T. M. Hughes, X. Lu, J. Li, J. K. Perry, D. P. Porter, T. Cihlar, T. P. Sheahan, R. S. Baric, M. Gotte, M. R. Denison, *Sci Transl Med* **2022**, *14*, eabo0718.

[24]   H. Tan, Q. Zhang, K. Georgiou, S. Zhang, K. Li, G. Lambrinidis, A. Kolocouris, X. Deng, J. Wang, *Nature communications* **2025**, *16*, 4548.

[25]   V. Chenthamarakshan, S. C. Hoffman, C. D. Owen, P. Lukacik, C. Strain-Damerell, D. Fearon, T. R. Malla, A. Tumber, C. J. Schofield, H. M. E. Duyvesteyn, W. Dejnirattisai, L. Carrique, T. S. Walter, G. R. Screaton, T. Matviiuk, A. Mojsilovic, J. Crain, M. A. Walsh, D. I. Stuart, P. Das, *Science advances* **2023**, *9*, eadg7865.

[26]   R. Gómez-Bombarelli, J. N. Wei, D. Duvenaud, J. M. Hernández-Lobato, B. Sánchez-Lengeling, D. Sheberla, J. Aguilera-Iparraguirre, T. D. Hirzel, R. P. Adams, A. Aspuru-Guzik, *ACS central science* **2018**, *4*, 268-276.

[27]   Y. Yang, S. Zheng, S. Su, C. Zhao, J. Xu, H. Chen, *Chemical science* **2020**, *11*, 8312-8322.

[28]   S. Kang, K. Cho, *Journal of chemical information and modeling* **2019**, *59*, 43-52.

[29]   S. Amabilino, P. Pogány, S. D. Pickett, D. V. S. Green, *Journal of chemical information and modeling* **2020**, *60*, 5699-5713.

[30]   M. Olivecrona, T. Blaschke, O. Engkvist, H. Chen, *Journal of cheminformatics* **2017**, *9*, 48.

[31]   O. Zhang, T. Wang, G. Weng, D. Jiang, N. Wang, X. Wang, H. Zhao, J. Wu, E. Wang, G. Chen, Y. Deng, P. Pan, Y. Kang, C. Y. Hsieh, T. Hou, *Nature computational science* **2023**, *3*, 849-859.

[32]   M. Xu, H. Chen, *Journal of chemical information and modeling* **2023**, *63*, 7067-7082.

[33]   Y. Matsuo, Y. LeCun, M. Sahani, D. Precup, D. Silver, M. Sugiyama, E. Uchibe, J. Morimoto, *Neural networks : the official journal of the International Neural Network Society* **2022**, *152*, 267-275.

[34] H. Cho, Y. J. Kim, J. W. Chae, M. R. Meyer, S. K. Kim, C. S. Ryu, *Frontiers in*
pharmacology **2023**, *14*, 1067408.
[35] R. A. Friesner, J. L. Banks, R. B. Murphy, T. A. Halgren, J. J. Klicic, D. T. Mainz, M. P.
Repasky, E. H. Knoll, M. Shelley, J. K. Perry, D. E. Shaw, P. Francis, P. S. Shenkin,
Journal of medicinal chemistry **2004**, *47*, 1739-1749.
[36] C. A. Lipinski, F. Lombardo, B. W. Dominy, P. J. Feeney, *Advanced drug delivery*
reviews **2001**, *46*, 3-26.
[37] M. Ringnér, *Nature biotechnology* **2008**, *26*, 303-304.
[38] H. Zhou, J. Skolnick, *The journal of physical chemistry. B* **2024**, *128*, 5363-5370.
[39] G. C. Linderman, S. Steinerberger, *SIAM journal on mathematics of data science* **2019**, *1*,
313-332.
[40] K. Ratia, S. Pegan, J. Takayama, K. Sleeman, M. Coughlin, S. Baliji, R. Chaudhuri, W.
Fu, B. S. Prabhakar, M. E. Johnson, S. C. Baker, A. K. Ghosh, A. D. Mesecar,
Proceedings of the National Academy of Sciences of the United States of America **2008**,
105, 16119-16124.
[41] Y. Cheng, W. H. Prusoff, *Biochem Pharmacol* **1973**, *22*, 3099-3108.
[42] S. Iketani, H. Mohri, B. Culbertson, S. J. Hong, Y. Duan, M. I. Luck, M. K. Annavajhala,
Y. Guo, Z. Sheng, A. C. Uhlemann, S. P. Goff, Y. Sabo, H. Yang, A. Chavez, D. D. Ho,
Nature **2023**, *613*, 558-564.
566

  Figures and Tables

Figure 1. Scaffold hopping of the GZNL-P4 structure. A) Binding mode of **GZNL-P4**. The crystal structures of **GZNL-P4** (orange) and **GRL-0617** (yellow) are overlapped. Ligands and residues surrounding the binding pockets are shown as sticks. Protein secondary structure is shown as transparent cartoons. Green and red dashes represent hydrogen bond and salt bridge interactions. B) Transformation of **GZNL-P4** to a topological tree. The solid and dashed circles represent essential and inessential nodes for molecular generation, respectively. The dashed lines represent the bonds to the inessential nodes. C) Workflow of scaffold hopping of the **GZNL-P4** structure based on the Tree-Invent model using the RL strategy. The blue and red circles represent prefixed and changeable nodes. D) Chemical space analysis based on the PCA analysis [37] using the Morgan fingerprint [38]. tSNE plot [39] was used to display the chemical space of molecules filtered based on docking scores (represented by the color of the heatmap) and their pharmacophore model. The labels I-X indicate ten diverse structures with reasonable docking scores.

Figure 2. Lead compound identification. A) Structures of nine synthesized compounds **1-9**. R^1 groups are shown at the top. B) Dose-dependent inhibitory activity of compounds **1-9** against PL^{pro} as determined by the fluorescence-based biochemical assay. Half maximum inhibitory concentration (IC_{50}) is shown in the legend. Data are presented as mean ± s.d. of three technical replicates. C) Differential scanning fluorimetry assay analysis of SARS-CoV-2 PL^{pro} with compounds. The data are presented as the mean of three replicates. D) Scheme of library synthesis for compound **10-36**. Synthesis routes were shown in the supplemental text. The inhibition activity (IC_{50}) and the delta melt temperature (ΔTm) of lead compounds **2**, **16**, **30**, and **31** were shown. The inhibition activity of IC_{50} is presented as mean ± s.d. of three technical replicates, but the ΔTm is presented as the mean of three replicates.

Figure 3. Binding mode of active compounds. A-C) The binding mode of compounds **2** (A)**, 16** (B)**, 30** (C) were determined by X-ray diffraction. Compounds **2, 16, 30** were shown as orange sticks. Residues around the binding pocket are shown as gray sticks. Protein secondary structure is shown as gray cartoons. Black dashes represent hydrogen bonds and salt bridges. The contour level of ligand map was set as 1.0. D) Docking poses of compound **31**. Ligands and residues are shown as orange and yellow sticks, respectively. Black dash represents either hydrogen bond or salt bridge interactions labeled with the distance. E) Alignment of the crystal structures with compound **2** (yellow) and **GRL0617** (green, PDB code: 7JRN). Protein secondary structure is shown as cartoons and residues around the binding pocket are shown as sticks. Ligands are specifically shown as orange sticks.

607
608 **Figure 4. The binding and inhibition activity of compound 16 to PL^{pro}E167K mutant.** A) The
609 inhibition activity of compound 16 for the PL^{pro}E167K mutant conferring drug resistance. The
610 error bars are mean ± s.d based on three technical replicates. B-E) The binding affinity obtained
611 by ITC of compound 16, Jun12682, and PF-07957472 to PL^{pro}E167K by ITC.

E

parameter	2	16	30	31
Rat Livermicrosome stability $t_{1/2}$ (min)	45.20	120.12	59.70	41.98
Rat Clint (mL/min/kg)	55.20	20.77	41.80	59.40
Human Livermicrosome stability $t_{1/2}$ (min)	30.29	39.55	47.90	35.30
Human Clint (mL/min/kg)	52.90	40.53	33.50	45.50

F

parameter	16	30	31
Mouse $T_{1/2}$ (h) PO@20mg/kg	0.757	0.699	0.930
Mouse Tmax (h) PO@20mg/kg	0.667	0.417	0.333
Mouse Cmax (ng/mL) PO@20mg/kg	750.218	1095.919	1245.280
Mouse AUC(0-t) (h * ng/mL) PO@20mg/kg	1862.128	1556.770	1219.430
inh%CYP 2C9@10μM	4.33	NT	NT
inh% CYP 2D6@10μM	37.15	NT	NT
inh%CYP 3A4@10μM (Midazolam)	56.88	NT	NT
inh%CYP 3A4@10μM (Testosteron)	37.13	NT	NT
hERG inh%@30μM	38.94	NT	NT

notes:
NT: No testing; PO: peros

Figure 5. Cellular activity and PK profiling of the compounds 2, 16, 30, and 31. A-C) Antiviral

activity of compounds **2**, **16**, **30**, **31** against SARS-CoV-2 wild type (WT) (A), variants Omicron

BA.5 (B), and XBB.1 (C). The EC_{50} was measured on virus infected Vero E6 cells. The error bars

are mean ± s.d based on three technical replicates. D) Toxicity to normal HEK293T cell lines. Data

are presented as the mean of two technical replicates. E) Table for liver stability in rat and human

liver microsomes. F) Table for in vivo PK characteristics, CYP inhibition and hERG toxicity.

620
Figure 6. *In vivo* antiviral activity of PL^pro inhibitor compound 16. A) Experimental design for
the 3-day experiment in K18-ACE2 mouse. B) Body weight loss of mice from different groups (n
$= 8$ per group). C) Live viral titers in lungs collected at 3 d.p.i. ($n = 8$ per group). The error bars
are mean $\pm$ s.d. Statistical differences were determined by Ordinary one-way ANOVA with
Dunnett multiple comparisons test for the body weight on day 3 in (B) and the live viral titer in
(C). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$; ns, not significant.
627

628   Table 1. SAR exploration of R² group

Compound	R²	PL^{pro} IC$_{50}$ (µM)	Compound	R²	PL^{pro} IC$_{50}$ (µM)
2		0.0238±0.0011	23		0.0992±0.0217
10		0.0327±0.0016	24		0.0563±0.0059
11		0.0288±0.0018	25		0.0156±0.0018
12		2.2330±0.2485	26		0.2861±0.0224
13		1.0300±0.1295	27		0.3016±0.0287
14		0.0453±0.0069	28		0.0376±0.0055
15		11.0700±14.6380	29		0.0256±0.0050
16		0.0105±0.0006	30		0.0108±0.0036
17		0.0316±0.0065	31		0.0131±0.0015
18		0.0520±0.0088	32		0.0103±0.0028
19		0.0284±0.0022	33		0.0494±0.0019
20		0.3836±0.0510	34		0.0519±0.0107
21		0.0483±0.0107	35		0.1817±0.0131
22		0.0560±0.0090	36		1.0540±0.1536

629   Data are presented as mean ± s.d. of three technical replicates.