1

Predicting speech-in-noise ability with static and dynamic auditory figure-ground

analysis using structural equation modelling

Xiaoxuan Guo[1*], Ester Benzaquén[1], Emma Holmes[2], Joel I Berger[3], Inga Brühl[4],

William Sedley[1], Steven P Rushton[5,6], Timothy D Griffiths[1,6]

[1]Biosciences Institute, Newcastle University, Newcastle upon Tyne, NE2 4HH,

United Kingdom

[2]Department of Speech Hearing and Phonetic Sciences, UCL

[3]Human Brain Research Laboratory, Department of Neurosurgery, University of Iowa

Hospitals and Clinics

[4]Maastricht University, Netherlands

[5]School of Natural and Environmental Sciences, Newcastle University, Newcastle

upon Tyne, NE1 7RU, United Kingdom

[6]Co-senior authors contributed equally to this work.

*All correspondence should be addressed to: Xiaoxuan Guo, Auditory Cognition Lab,

Newcastle University Medical School, Framlington Place, Newcastle upon Tyne NE2

4HH. Email: x.guo20@newcastle.ac.uk.

Abstract

Auditory figure-ground paradigms assess the ability to extract a foreground figure

from a random background, a crucial part of central hearing. Previous studies have

shown that the ability to extract static figures (with fixed frequencies) predicts real-life

listening: speech-in-noise ability. In this study we assessed both fixed and dynamic

figures: the latter comprised component frequencies that vary over time like natural

speech. 159 participants (aged 18-79) with a range of peripheral hearing sensitivity

were studied. We used hierarchal linear regression and structural equation modelling

to examine how well speech-in-noise ability (for words and sentences) could be

predicted by age, peripheral hearing, and static and dynamic figure-ground.

Regression demonstrated that in addition to the audiogram and age, the low-

frequency dynamic figure-ground accounted for significant variance of speech-in-

noise, higher than the static figure-ground. The structural models showed that a

combination of all types of figure-ground tasks predicted speech-in-noise with a

higher effect size than the audiogram or age. Age influenced word perception in

noise directly but sentence perception indirectly via effects on peripheral and central

hearing. Overall, this study demonstrates that dynamic figure-ground explains more

variance of real-life listening than static figure-ground, and the combination of both

predicts real-life listening better than hearing sensitivity or age.

Key words: speech-in-noise perception, auditory figure-ground, pitch perception,

structural equation modelling, pure-tone audiogram, new hearing measure

Introduction

Tracking a target sound in a complex auditory scene is one of the core tasks that the

auditory system performs and forms an important part of hearing ability. Complaints

about understanding speech in noisy environments are frequently encountered in

audiology clinics but are difficult to assess because the pure-tone audiogram does

not fully reflect this ability (Merten et al., 2022; Besser et al., 2015; George et al.,

2007). Sentences-in-noise and word-in-noise tests have been developed to simulate

real-life speech-in-noise (SIN) situations, but responses are influenced by other

factors, such as levels of education, accent, and language experience as much as

hearing and sound segregation. This has motivated work to develop non-speech

measures of figure-ground analysis. A non-speech auditory target-in-noise task has

been developed called the stochastic figure-ground test (SFG) or fixed-frequency

auditory figure ground (AFG-Fixed, Teki et al., 2013). Modelling suggests that sound

segregation is achieved based on temporal coherence of the figure (Teki et al.,

2013), requiring auditory cortex. Brain studies implicate a network including high-

level auditory cortex in humans (Teki et al., 2016; Teki & Griffiths, 2016) and in a

primate model (Schneider et al., 2018). Holmes & Griffiths (2019) demonstrated that

an AFG discrimination task requiring detection of a gap in auditory figure predicted

SIN performance and explained variance in SIN independent of that explained by the

pure-tone audiogram (PTA).

SIN perception recruits acoustic features to better segregate sounds in noise. One of

the key features is the fundamental frequency corresponding to the perception of

pitch in sentences. In this study we assess a type of AFG task in which the

frequency components vary over time following the pitch contour of natural speech.

This makes the stimulus more speech-like, whilst retaining the overall advantage of

the AFG task as a 'pure' measure of grouping relevant to real-life listening without

linguistic confounds. Two previous studies have investigated AFG with changing

frequency patterns (Holmes & Griffiths, 2019; O'Sullivan et al., 2015). Holmes &

Griffiths (2019) generated figures that followed the formants of spoken stimuli. While

figures generated from the first three formants of speech did not significantly predict

SIN, a stimulus based on the first formant and additional components that changed

over time coherently with the first formant did predict SIN. We assess similar stimuli

here with multiple frequency components that change coherently based on the pitch

contours of speech sentences, rather than the frequency of the first formant. We test

how well the detection of these frequency components predicts SIN perception.

Natural voiced speech contains multiple harmonics related to the fundamental

frequency and is associated with pitch. Harmonicity aids hearing in noise

(McPherson et al., 2022). Pitch contributes to SIN processing, especially for people

with higher language or hearing competence (Llanos et al., 2021; Shen & Souza,

2018; Huang et al., 2017). In this study, we generated figures related to the harmonic

structure of speech, in contrast to the non-harmonic figures used in previous work.

We extracted the fundamental frequency from naturally spoken sentences and

developed a new type of dynamic auditory figure-ground stimulus using harmonic

complexes based on this. We call this the dynamic figure-ground stimulus (AFG-

Dynamic). The harmonic features make the auditory figure-ground more speech-like

from an acoustic perspective, without incorporating high-level linguistic cues.

We created harmonic complexes in different frequency ranges to explore the

importance of the frequency range of the figure. Previous studies suggest that high-

frequency hearing sensitivity based on the audiogram may be an important

determinant of SIN ability (Polspoel et al., 2022; Zadeh et al., 2019; Holmes &

Griffiths, 2019), but have not examined complex figures in different frequency

ranges. We constrained the frequency range of the AFG-Dynamic stimuli to low-

frequency AFG (AFG -Low) and high-frequency AFG (AFG-High) components to

explore how grouping ability in different frequency ranges contributes to SIN

perception.

Predictive measures of Speech-in-Noise Perception

The first aim of the study was to investigate if the new dynamic auditory figure-

ground tests are predictive of SIN measures. We hypothesised that both versions of

AFG-Dynamic tests (AFG-Low and AFG-High) can predict SiN perception and

explain extra variance of SIN independent of the PTA or the prototypical AFG-Fixed.

As speech is dynamic in its frequency profile whereas single words have relatively

static frequency pattern, we predicted that the fixed-frequency AFG-Fixed better

predicts word-level segregation whereas the AFG-Dynamic tests with the changing

pitch pattern better predicts sentence-level sound segregation. Three metrics that

reflected the SIN ability were used as the outcome measures, including a word-in-

noise test (WiN, Guo et al., 2024; Geller et al., 2021), a sentence-in-babble test (SiB,

Holmes & Griffiths, 2019), and a subjective self-report measure (The Speech, Spatial

and Qualities of Hearing Scale, 'SSQ'; Gatehouse & Noble, 2004). To capture the

word-level perception, the British-ITCP test was used in this study, which has been

demonstrated to have a close association with sentence-in-babble performance

(Geller et al., 2021). Sentence-level perception is an important aspect of SIN

perception and is most commonly assessed as a measure of functional hearing. SiB

tests have long been developed to capture patient speech perception in noise

(Elliott, 1995; Nilsson et al., 1994; Killion et al., 2004; Wilson et al., 2012). The

sentence-in-babble test used here consists of the Oldenburg sentences and babble

noise and has been used previously as an assessment of SIN perception (Holmes &

Griffiths, 2019). We further included a subjective measure (SSQ) to incorporate

participant experience of real-life listening to the overall assessment of SIN ability as

well.

Modelling the Relationships Among Auditory Figure-Ground Perception, Age,

and Speech-in-Noise Perception

The second aim of the study was to describe the relationships among the

psychoacoustic measures used in the study and identify the contribution of different

factors to SIN perception in a multivariate model using structural equation modelling

(SEM). The current study had a complex design investigating different measures of

hearing thresholds, auditory figure-ground, and speech-in-noise. This type of design

favours the use of SEM compared to regression as it allows having multiple

observed variables indicating one latent variable (hypothetical constructs that are not

directly measured but can be inferred by their observed variables) and reflects the

relative importance of indirect effects, such as the interaction between covariates on

outcomes. Three conceptual models based on different outcome variables were

therefore constructed with assumptions on the direction of causality according to

existing literature. The three outcome variables were: the word-in-noise measure, the

sentence-in-babble measure, and the two measures combined. As the word- and

sentence-level SIN analysis and the self-reported SIN ability tap into different

domains of SIN perception, models predicting the three SIN measures separately

should provide additional information on the differences of the three domains of SIN

analysis when interacting with AFG, PTA and Age. We also investigated the domain-

general SIN by combining the three measures into one latent variable.

The fixed-frequency AFG test has been shown to predict SIN perception (Holmes &

Griffiths, 2019). In this study, we added the two additional dynamic AFG measures

with high and low frequencies to form an AFG latent variable that predicts SIN

perception.

In terms of the exogenous variables, PTA and the participant's age were taken into

account. PTA has been shown to predict SIN ability (George et al., 2007; Wong et

al., 2008; Besser et al., 2015; Bochner et al., 2015; Holmes & Griffiths, 2019). Age

has also been recognised as a key factor impacting both hearing and SIN perception

(see Billings & Madsen, 2018 for a review on the topic). Deterioration of the auditory

periphery – including hair cell and cochlear nerve loss, as well as cochlear

synaptopathy (Dias et al., 2024; Liu et al., 2024; Xie et al., 2024) could all lead to

decreased real-life listening ability, and these peripheral deteriorations are all tied

with aging (Chadha et al., 2021). Holmes and Griffiths[8] found a relationship between

age and both AFG and SIN performance. Altered auditory peripheral function could

result in lowered frequency and temporal resolution, which would inevitably impact

the central sound segregation ability measured by the AFG tests. However, the

relationships between AFG and SIN perception retained after accounting for age and

PTA, which indicates that figure-ground measures may also index PTA- and age-

independent deficits in SIN perception. Thus, we hypothesised that PTA and Age

would predict both SIN and AFG with Age impacting PTA, and that AFG can

independently predict SIN after accounting for Age and PTA.

Methods

Participants

A total of 170 participants were recruited, of whom 11 were excluded due to data

quality as per criteria described later. The final sample size used for analysis was

159. The sample had a wide range of age (mean = 45.95, SD = 18.47, range = 18 –

79) as well as hearing thresholds (mean = 13.30 dB, SD = 9.90 dB, see Figure 1 for

more detailed audiogram results), with 105 female participants. All participants were

neurotypical native English speakers with no history of auditory disorders, no history

of speech and language disorders, and who were not currently taking any

psychotropic drugs. Informed consent was obtained from participants before the

experiments. The study was approved by the Newcastle University Ethics Committee

(46225/2023).

Figure 1. The distribution of hearing sensitivity of 250 – 8000 Hz for the left and the right ear separately

of all participants. The x axis shows the frequencies measured, and the y axis shows hearing thresholds

measured in decibels. The coloured lines with circles plot individual audiogram results, and the thicker black line

with circles and error bars show the averaged group PTA. The error bars display the standard deviation.

Stimuli and Tasks

Fixed-Frequency Auditory Figure-Ground Gap Discrimination Task

The AFG-Fixed gap discrimination test was the same as that developed by Holmes

& Griffiths (2019); an auditory figure was composed of temporally coherent pure-tone

elements (each 50 ms duration) repeating over time. Each figure was 42 chords long

with 3 figure components per chord (i.e. coherence level of 3). The figure was

superimposed on an auditory ground, which is a tone cloud made of pure-tone

elements (also 50 ms duration each) of randomised (or stochastic) frequencies

between 180 – 7246 Hz. In each trial, two figure-ground stimuli were presented to

the participants, sequentially with an inter-stimulus interval of 400 ms. A gap (6

chords long) was present in either figure. Although, importantly, the ground tones

continued through the gap, so participants needed to have segregated the figure

from the ground to perform this task. The participants were instructed to choose

which of the two figure-ground stimuli contained a gap in the figure. The test used a

1-up 1-down adaptive procedure, starting at signal-to-noise ratio (SNR) of 6 dB and

varied systematically across trials. The step size started at 2 dB and went down to

0.5 dB after 3 reversals. Two runs were presented to each participant with different

exemplars, with both runs terminating after 10 reversals. The median of the last 6

reversals for both runs were taken and averaged as a measure of performance.

Higher SNR scores indicate worse performance.

Dynamic Auditory Figure-Ground Pattern Discrimination Task

In contrast to the prototype AFG-Fixed that has a fixed-frequency pattern over time,

the novel dynamic AFG contains pitch information akin to speech. The pitch contours

were extracted from the English Oldenburg sentences read by a male British

speaker (Holmes & Griffiths, 2019), using Praat version 6.2.09 with a time step of

0.75/75 Hz (100 pitch values per second), and had a frequency range of 74.94 –

295.44 Hz (M=131.59, SD=15.61). The low-frequency noise (below 10 Hz) and

artificial high frequencies (above 300 Hz) introduced by the Praat periodicity analysis

were removed to obtain pitch trajectories (see Figure 2(a) for an example). There are

gaps in natural pitch tracks as shown in Figure 2 (a). To avoid the participants using

these gaps, the natural speech gaps and stops were first removed from the pitch

contour. An example of the conjoined signal is shown in Figure 2(b). As

demonstrated in the plot, the new signal has a general downwards trend and some

sharp transitions caused by the removal of the gaps and linear interpolation. To

remove the drift from individual signals, we demeaned the signal and applied

detrending to the demeaned signal. Low-pass filtering (minimum-order filter with a

stopband attenuation of 60 dB) with a 2000 Hz cutoff frequency was then carried out

to remove the artificial spikes. The trend and the mean were then added back to the

filtered signal to keep the final signal as similar to the original pitch trajectory as

possible. An example of the final pitch signal is plotted in red in Figure 2(b).

Figure 2. The figure shows the extraction of pitch contour (Figure 2(a) (b)) and the AFG stimuli with the

pitch contour embedded (Figure 2(c) (d)). Figure 2(a) and (b) shows that the pitch information extracted from the

sentence "Alan brought four small desks". The x axis plots the time in seconds and the y axis shows the

frequencies in Hz. Figure 2 (a) shows the raw pitch contour plotted against time. Figure 2 (b) shows the pitch

trajectory after being processed. The blue line is the pitch contour with the gaps removed. The red line shows the

final processed signal. The dotted plots illustrate examples of the two different types of AFG-Dynamic stimuli.

Figure 2 (c) shows the lower-frequency dynamic AFG. Figure 2 (d) on the right side is the high-frequency

dynamic AFG. The x axis shows the time in milliseconds and the y axis shows the frequency in Hz. Figure

elements are depicted in orange while ground elements are depicted in grey.

After processing the pitch signals, the resultant frequency profiles were grouped into

50-ms long segments by computing the average to form the figure elements. The

second, third, and fourth harmonics of these fundamental components were used to

construct the remaining elements of each chord (see Figure 2(c)) for the AFG-Low.

The tones were gated with a 10 ms raised-cosine ramp to smooth the onset and

offset of the sounds. The high-frequency figure (AFG-High) retained the pitch

trajectories of the low-frequency version, but the components were the fundamental

frequencies multiplied by 5, 10, 20, and 30. The top frequency of each figure was

checked not to exceed the masking frequencies. Like the AFG-Fixed stimuli, the

auditory ground was comprised of randomised pure-tone segments on a logarithmic

scale. Although, while ground tones for the AFG-High stimuli used the same range of

frequencies as the AFG-Fixed (180 Hz–7246 Hz), AFG-Low stimuli used ground

tones with a lower frequency range (90 – 3623 Hz; in other words, half of the upper

and lower frequency values from the AFG-High stimuli) to achieve a better masking

effect. See Figure 2 (c) (d) for an illustrated example of the two types of stimuli.

The two AFG-Dynamic tests (AFG-Low and AFG-High) had the same paradigm and

were counter-balanced across participants. Within each test, both the figure and the

ground stimuli were presented every trial, either with the same or different figure

pattern. In case of a trial with different figure patterns, the durations of the figures

were matched but the frequency elements were based on different pitch trajectories.

The ground elements were tailored to different figures. The tests used a two-

alternative forced-choice task, which required the participants to hold the sounds in

memory and decide whether or not the second figure had the same pattern as the

first one. The inter-stimulus (within each trial) interval was 0.2 seconds. A two-down

one-up staircase procedure was used with a total of 22 reversals. The initial SNR

was 12 dB with a step size of 2 dB, which then changed to 0.5 dB after 7 reversals.

The final score was calculated by averaging the dB SNR of the last 6 reversals and a

higher SNR would indicate poorer performance. The sound pairs, which were

matched for duration, and the trial orders were kept the same across participants.

The same design was used for both the low-frequency and the high-frequency

versions of the AFG-Dynamic test.

Speech-in-Noise

Word-in-Noise Test

The WiN test was the British Iowa Test of Consonant Perception (B-ITCP) adapted

from the Iowa Test of Consonant Perception ((Guo et al., 2024)). Target speech

sounds were monosyllabic words with a consonant-vowel-consonant (CVC) or

CVCC structure. The test used British accents including one female and one male

voice. The babble noise was an 8-talker babble, presented at -2 dB signal-to-noise

ratio (SNR). The onset of the auditory target was between 0.5 and 1.0 s, randomly

positioned from the babble onset. The babble segment of each trial was randomly

selected from a 15-seconds babble stimulus. The length of the words varied from

0.304 – 0.757 s (mean: 0.508 s, SD: 0.086 s). Participants were asked to choose the

word they heard out of a list of 4 words displayed on the screen balanced for

difficulty and phonetic contrasts. Percent correct across trials was taken as the

outcome measure for the WiN test. This is the only test that was scored differently,

and a higher score for the WiN test indicates better performance.

Sentence-in-Babble Test

Details of the SiB test are reported in Holmes & Griffiths (2019). The target

sentences were English Oldenburg sentences consisting of 5 words following a

structure of Name-Verb-Number-Adjective-Noun (e.g., "Alan brought four small

desks"). The sentences were masked by 16-talker babble. The target sentences

appeared 500 ms after babble onset and ended 500 ms before babble offset.

Participants were shown a 5*10 matrix on the screen, where each word in the

sentence had 10 options. The test used a one-down one-up adaptive paradigm with

the starting SNR at 0 dB. The total number of reversals was 10 and the step size

began at 2 dB and decreased to 0.5 dB after 3 reversals. The task had two runs

interleaved. The target sentences were different in each run. The final score was

calculated by averaging the dB SNRs of the last 6 reversals across the 2 runs. A

lower score in this test indicates better performance.

Speech, Spatial and Qualities of Hearing Scale

The subjective self-report SIN ability was assessed using the Speech, Spatial and

Qualities of Hearing Scale-speech hearing (SSQ, Gatehouse & Noble, 2004). Two of

the questions were removed from the shortened speech-hearing questionnaire due

to their ambiguity. See Appendix 1 for the full list of questions used in this

questionnaire. Each item has a score from 0 to 10 with the higher score indicating

more difficulty in hearing.

Procedure

An audiometry test was carried out, followed by the 5 computer tasks, which were

presented in a fixed order for all participants, except that the order of the AFG-High

and the AFG -Low tests were counterbalanced across participants. The tasks were

presented in the following order: (1) SiB, (2) AFG-Dynamic test (AFG -High or AFG-

Low, determined by counterbalancing across participants), (3) SSQ, (4) WiN, (5)

second version of the AFG-Dynamic test (AFG- High or AFG -Low, whichever they

had not already completed), (6) AFG-Fixed. Participants were asked to sit in the front

of a computer monitor (Dell Inc.) used to present the tasks. The auditory stimuli were

presented through headphones (Sennheiser HD 380 Pro) linked to a sound card

(RME FireFace UC).

Data Analysis

Test of Correlation for AFG-Dynamic

The outcome measures of SiB and the AFG tests were the medians of the last 6

reversals. The performance was considered stable if the performance differences of

the last 6 reversals were smaller than ± 5 dB. Participants who did not show stable

performance were excluded from the final analysis. Bivariate correlations and

hierarchical regressions were carried out to explore the relationship between AFG-

Dynamic and SIN tests. Tests of normality (Kolmogorov-Smirnov) showed that AFG-

High and WiN were not normally distributed, so Spearman's rho was used to

examine the hypotheses regarding the relationships between the three speech

measures with AFG-Dynamic (low and high version), AFG-Fixed, PTA and age.

Holm-Bonferroni correction was applied to correct for multiple comparisons. As linear

regression is a more tolerant measure for non-normality, stepwise regression was

conducted to check if there were predictive relationships between SIN and AFG as

well as specifying the variance explained by individual predictors. The order of

entering the independent variables is based on how close their associations were

according to the correlation results (Figure 4). These tests were performed using

SPSS version 28 and visualised with MATLAB R2021a.

Modelling the Inter-Relations of Predictors of SIN

To account for the inter-relationships of the indicator variables, structural equation

modelling (SEM) was conducted using the lavaan (version 0.6-15) package in R

(version 4.2.1). Maximum likelihood estimation was used with nonnormality

correction based on the Satorra-Bentler scaled test statistic. Robust measures were

reported in this study (Brosseau-Liard et al., 2012; Brosseau-Liard & Savalei, 2014).

Initial conceptual models (Model 1 and 2) are illustrated in Figure 3 (a). Model 1&2

were devised to explore word-level and sentence-level SIN analysis separately.

Model 3 illustrates a combined model of all three SIN measures. In all three models,

the SIN measures were predicted by AFG indicated by the AFG-Fixed, and two

AFG-Dynamic measures. PTA and Age also predicted SIN as well as AFG. To

decide the latent variable structure, a confirmatory factor analysis (CFA) was used to

examine the measurement quality on a subset of the data (101 participants) before

conducting the final analysis. Figure 3 (b) demonstrates the CFA models of the two

latent constructs in the three models: SIN and AFG. While there are no rules of

thumb defining the acceptable thresholds of a factor loading, SSQ as a measure of

functional hearing should predict a large variance of SIN tests similar to the other two

SIN indicators. The SSQ however had a visibly weak connection to SIN and thus

was removed from further analysis. All three AFG indicators seemed to be

acceptable to be entered to the final model.

Figure 3. Figure 3 (a) shows the conceptual models of WiN(Model 1), SiB perception (Model 2), and SIN

with combined word and sentence perception (Model 3). The shaded ovals represent latent variables, the

rectangles represent observed variables, and the diamonds with striped shading are exogenous variables. The

arrowed circle of each variable represents the error (the size of the circle is not proportional to the radius). The

indicators have arrows pointing to them from the latent variables. External variables point the arrows to the latent

variables to suggest a causal effect on the latent variables. Figure 3 (b) shows the CFA of SIN and AFG. Shaded

ovals represent the latent variables, rectangular boxes are the indicators, and the circles associated with each

variable are the residuals. Latent variables are connected to their indicators through arrows pointing to the

indicators. The error for SIN and AFG is 1 as they are not subject to any causal influences in this limited model.

The results of this analysis were used to guide the selection of scaling variable

(Bollen et al., 2022). Scaling variables are used to assign scales to latent variables,

which is essential when identifying a model. The method used in lavaan is the Fixed

Marker (FM) scaling that fixes the loading of the chosen scaling variable to

1(*Lavaan.Org - Model Syntax 2*, n.d.). The path coefficients (abbreviated as β) can

be interpreted as: one SD of variable A increase leads to a β SD increase of variable

B while all other relevant connections were held constant. The residual or

measurement error of the indicators represents variance unexplained by the

measure "due to random measurement error, or score unreliability" (Kline, 2015,

pp9).

The WiN measure was chosen as the scaling variable based on its high path

coefficient connecting to the SIN latent variable. WiN was the only test measured by

percentage, which resulted in a difference in scale of the outcome compared to the

other tests. This was re-scaled via z-scoring (removing the mean and dividing the

results by the standard deviation (SD) of the original scores of WiN). Importantly,

contrary to the measures assessed with SNR, a higher score of WiN indicated better

performance. This means one SD increase from the mean in WiN would lead to a β

SD decrease in SIN. However, since WiN was used as the scaling variable, the SIN

latent variable took the scale of WiN, SiB instead showed a negative path coefficient.

The different interpretation of SNR- and percent correct -based scoring would further

influence other factors connecting to SIN. To avoid confusion and simplify results

interpretation, the WiN results were multiplied by -1 so a higher score would indicate

worse performance.

AFG-High, AFG-Low, and AFG-Fixed are the indicators of the latent variable AFG.

Similarly, the AFG-Fixed was chosen as the scaling variable due to its close

connection with the AFG latent variable. AFG-High and AFG-Low were made with

similar parameters except for the frequency range and should tap into very similar

mechanisms, hence they covary. The three SEM models further consisted of Age

and PTA as exogenous variables, which were both configured to predict SIN and

AFG.

The model fit was evaluated by a set of criteria (Hu & Bentler, 1999; Kline, 2015).

These includes the Bentler comparative fit index (CFI) and Tucker-Lewis Index (TLI),

the root-mean-square error of approximation (RMSEA), the standardised root mean

squared residual (SRMR), and the chi-square test. Both RMSEA and SRMR are

absolute measures of the estimated discrepancy between the predicted and

observed models. The SRMR is a measure of the mean absolute correlation residual

measuring the differences between the original correlations (observed) and the

implied correlations by the model. RMSEA ≤ 0.06 and SRMR ≤ 0.08 have been

suggested to indicate a close model fit (Hu & Bentler, 1999). RMSEA up to 0.10 is

considered fair fit, but above 0.10 is generally unacceptable (Browne & Cudeck,

1992). CFI and TLI, on the other hand, are incremental indices that reflect the

relative improvement of the model fit compared to a baseline model (Kline, 2015).

TLI is non-normed so it can fall outside the 0-1 range whereas CFI is normed, but

the cutoff for both of them is above 0.95 for a good fit (Hu & Bentler, 1999). The chi-

square (χ^2) result was also reported (Kline, 2015). The null hypothesis for the chi-

square test is that the predicted model reflects the true data perfectly. Thus, a

nonsignificant chi-square would indicate a good model fit.

Results

The descriptive statistics are reported in Table 1. WiN had the narrowest range of

data distribution (Table 1), with an accuracy of around 67% correct. All other tests

were measured in signal-to-noise ratio, with the AFG tests showing a wider spread of

performance than SiB (Table 1). The two AFG-Dynamic tests had positive thresholds

whereas the fixed-frequency AFG elicited negative thresholds, indicating that the

AFG-Dynamic tests were more challenging.

	Mean	Standard Deviation
SiB	-0.880	2.114
WiN	0.673	0.107
AFG-Low	8.991	10.897
AFG-High	7.252	10.447
AFG-Fixed	-14.542	8.200

Table 1. The mean and standard deviation of the participants' performance on the five computer tasks.

Relationships between SIN measures and AFG-Dynamic

Both the sentence-level and the word-level SIN tests showed moderate to strong

correlations with all the AFG measures, hearing, and age (Figure 4). After correction,

all p-values remained highly significant. SSQ, however, did not show any significant

correlation with other speech measures (p > 0.34) and was removed from further

analysis.

Figure 4. Bivariate correlation among variables. The p values shown here were corrected. The lines of

best fit were plotted as straight lines in the figure with shaded error bars. The x axis for the left panel showed the

WiN results as proportion correct (number of correct answers divided by the overall number of trials), and the y

axes of the figures on the left panel from top to down were the two AFG tasks measured in dB SNR and Age and

PTA. On the right panel were three figures of the same variables plotted against the SiB test results measured in

dB SNR.

The hierarchical regression predicting SiB performance gave three significant

predictors, revealing that PTA, AFG-Low, and AFG-Fixed performance significantly

predicted SiB performance ($F_{(3, 155)} = 39.879$, $p < .001$). The model accounted for

43.56 % of the variance in SiB. Table 2 specifies the variance explained by the

significant predictors. For the WiN model, four significant predictors were significant:

age, PTA, AFG-Low, and AFG-Fixed ($F_{(4, 154)} = 62.560$, $p < .001$). The model

accounted for 61.90% of the variance in WiN. Table 2 specifies the variance

explained by each predictor. For SiB, PTA was the best predictor explaining about

31% of the model with the AFG-Low adding 9.9% to the model. Whereas, for WiN,

Age seemed to be the strongest predictor. Of the significant predictors, AFG-Fixed

added the least variance to both SiB and WiN, which was about 1%~2% after

accounting for the other variables.

SiB	Standardised Coefficients Beta	R^2	p	WiN	Standardised Coefficients Beta	R^2	p
PTA	0.368	0.314	< 0.001	Age	-0.409	0.498	< 0.001
+ AFG-Low	0.253	0.413	< 0.001	+AFG-Low	-0.217	0.580	< 0.001
+ AFG-Fixed	0.197	0.436	0.015	+ PTA	-0.218	0.609	0.003

+AFG-High	0.126	-	0.125	+ AFG-Fixed	-0.134	0.619	0.049
+Age	0.074	-	0.400	+AFG-High	-0.121	-	0.075

Table 2 This table displays the R² values and p values of models including an increasing number of predictors that add significant variance to the models predicting either SiB or WiN.

Structural Equation Model of SIN, AFG, Hearing, and Age

The fit indices for the three models are shown in Table 3, and path coefficients are plotted in Figure 5. All fit indices for Model 1 and Model 2 were within our criteria. The path coefficients in Model 1 were all significant. Model 2 had mostly significant paths with a nonsignificant one of Age to SiB.

Fit Index	Model 1	Model 2	Model 3
χ^2	9.547 (p=0.92)	7.910 (p=0.91)	20.617 (p=0.940)
RMSEA	0.079	0.065	0.092
CFI	0.990	0.992	0.978
TLI	0.969	0.975	0.949
SRMR	0.028	0.029	0.036

Table 3 Fit indices for Model 1, 2 and 3.

As expected, in both Models 1 & 2, all three AFG indicators showed significant

contributions to the AFG indicator, and the AFG-High and AFG-Low shared

significant covariance. AFG-Fixed contributed to AFG with the largest path

coefficient ($|\beta| = 0.82$) followed by the two dynamic AFG measures. The latent AFG

variable predicted WiN and SiB significantly, with the largest variance compared to

PTA and Age in both models. PTA had a significant but smaller contribution to each

SIN measure. Age only has a significant direct impact on WiN and not on SiB.

Model 3 followed the conceptual model structure shown in Figure 3 but had the path

connecting SSQ to SIN removed as it was not significant. This model met most of

the set criteria for an excellent model fit except for the RMSEA (Table 3). RMSEA

incorporates model complexity and models with smaller degrees of freedom tend to

obtain a poorer RMSEA (Kenny et al., 2015). This pattern of results is similar to that

obtained for Models 1 and 2, which also had excellent fit based on most of the

indicators but poorer than expected RMSEA. However, as the combined results of

other indicators all showed that the model fits the data very well, we deem that this

model is acceptable.

Similar to Models 1 and 2, AFG explained the largest variance of the latent SIN

variable (β = 0.56) in Model 3 (Figure 5), compared to Age and PTA. Both SiB and

WiN showed significant contributions to the latent SIN variable, but WiN had a

numerically greater contribution ($|\beta|$ = 0.86 for WiN compared to $|\beta|$ = 0.71 for SiB).

Age was the second largest predictor of SIN, after AFG, and PTA had a smaller (but

nevertheless significant) direct impact on SIN. Although, both PTA and Age had a

significant indirect impact on SIN through AFG.

508   **Discussion**

Predicting SIN Perception with Dynamic AFG in the Linear Models

This study showed a moderate to strong correlation between all AFG measures and

SIN, both on the word and the sentence level. The low-frequency AFG came out as

a significant predictor of WiN and SiB, explaining the largest variance in both models

after accounting for demographic factors (Age or PTA). It is unexpected that even for

the WiN model the AFG-Low explained more variance than the static AFG. The

dynamic AFG was designed to carry the fundamental frequency patterns and so

should better predict sentence-level sound segregation than word-level. AFG-Fixed,

on the other hand, had no frequency change over time, which was considered more

similar to WiN perception. Based on the regression results, however, it seems that

adding the speech pitch pattern to the AFG stimuli only improved its predictive power

of SIN in general, not specifically to sentence-level perception. This general

improvement could be the reason the AFG-Fixed did not explain a higher portion of

the variance of SIN as well. Considering that AFG-Low combined both the

mechanism of segregating static figure from ground employing the figure's temporal

coherence feature, and speech-like frequency pattern to aid SIN perception, it is

reasonable to see higher variance obtained by AFG-Low in a linear model.

One possible explanation for the relationship between AFG-Low and both word and

sentence-level SIN is its harmonic structure. AFG-Fixed differed from AFG-Dynamic

in two ways: it is both static and inharmonic. Some of the AFG-Fixed stimuli might

contain near-harmonic figures by chance, but most of the stimuli were inharmonic,

which elicit weaker pitch perception (Micheyl et al., 2012). Pitch plays an important

role in SIN perception (Meha-Bettison et al., 2018; Shen & Souza, 2018; Binns &

Culling, 2007; Carroll & Zeng, 2007), the mechanism of which was reviewed by

Oxenham (2008). This includes not only its strong association with the stress contour

of a whole sentence but also other linguistic features such as phonemes in words.

The pitch information embedded in the AFG-Low can help with differentiating the

envelope fluctuations of the target sound from the background sound, which is key

for speech intelligibility. Thus, the stronger pitch strength could be the reason that

AFG-Low, while sharing the same basic principles with the static AFG, predicts SiB

or WiN better.

The high-frequency dynamic AFG had a numerically weaker correlation as was

hypothesised and did not explain additional variance in WiN or SiB after accounting

for other tasks. This could be because AFG-High shared a high covariance with

AFG-Low due to the similar parameters used for these two tests. The AFG-Low

more closely resembles the speech stimuli used in this study with its frequency

range being closely configured to the pitch range of speech formants, which might be

the reason that AFG-Low outperformed AFG-High in predicting SIN.

The SSQ measure did not correlate with either of the speech measures, which was

not a unique finding (Oberfeld & Klöckner-Nowotny, 2016; Ertürk et al., 2023). This

could be because the shorter SSQ version does not have enough sensitivity to

capture SIN perception as only a few questions were related to speech

comprehension in human speech noise.

Predicting SIN Perception in a Multivariate Model

The linear regression models displayed the core contribution of the new dynamic

AFG measure as well as the static measure. However, the stepwise procedure did

not account for the interaction between variables. The SEM model provided a more

comprehensive picture of the experiment that went beyond ranking the important

predictors of SIN measures.

Firstly, Models 1 and 2 showed that all three AFG predictors have an impact on the

SIN performance. This means that when accounting for the interaction and

covariance shared between the indicators, all of the AFG predictors should be

considered a necessary part of the auditory figure-ground analysis. Interestingly,

while the linear measures showed a tighter relationship between AFG-Low and

WiN/SiB, AFG-Fixed in the SEM models contributes the most to the AFG latent

variable. This suggest that as the 'prototype' AFG, the static AFG that assesses

people's ability to pick up the temporally coherent figure from the tone cloud, is still

the core of the AFG analysis process. Combined with the regression results, it shows

that the dynamic pitch pattern does add an important aspect to AFG and should be

used in combination with AFG-Fixed. Based on their individual predictive value of the

regression results, in a linear model, when using both measures are not possible,

AFG-Low should be a better test to measure SIN ability compared to AFG-Fixed.

The combined AFG measures explained the largest variance (43%, 52%) of both

speech measures in Model 1 & 2, compared to Age and PTA. This also differs from

the linear regression results, where PTA or Age was found to be the greatest

predictor. This difference suggests that AFG tasks have greater predictive power in

combination as various ways to assess scene segregation, whereas each AFG task

individually assesses slightly different abilities that are weaker individually than the

influence of the demographic factors. The lower path coefficient of PTA compared to

AFG indicates that the ability to process speech (either single-word utterances or

sentences) in a noisy environment directly relies more on segregating auditory

streams and tracking the pattern of the target sounds over time than simply picking

up acoustic signals as measured by PTA. However, PTA also had a mediation effect

on WiN/SiB through a large path coefficient to AFG. This means that in addition to a

relatively small direct impact on SIN perception, deteriorated peripheral hearing

could alter functional hearing through modifying central sound processing, which is

consistent with our hypothesis.

A mediation effect was also evident with the Age-driven impact on SIN perception.

Age led to 71% SD change in PTA in this study, meaning the PTA variance was

largely dominated by age-related hearing loss. Age also decreased central sound

processing measured by AFG here by 34%, consistent with the results reported by

Holmes and Griffiths (2019). However, while Age showed a significant correlation

with SiB, it did not modify SiB performance directly in the SEM model, which is

consistent with the regression results. WiN is a harder task for people who are older

or with higher hearing thresholds. This is because less in the way of compensatory

mechanisms can be employed for hearing a short word compared to a sentence that

contains a legitimate syntactic structure. While normal ageing can result in

deteriorated hearing sensitivity and the perception of other acoustic properties (fine

structure or harmonicity), language perception skills are generally preserved (Burke

& Mackay, 1997).

The interaction among predictors in Model 3 did not change much after combining

the WiN and SiB into one latent variable. WiN and SiB had a similar level of

contribution to the SIN latent factor and the small residual term of SIN suggests that

WiN and SiB together provide a holistic measurement of SIN, with a small effect of

unmeasured sources of unique variance on the latent variable. It is important to

highlight however, that combining the measures into a latent variable could hide the

different effects of other predictors such as Age, like in Models 1 and 2.

Limitations and Future Direction

The sample size of the current study might have caused the fit to be suboptimal.

There is no golden rule in terms of determining an appropriate sample size for SEM.

Researchers have suggested a variety of standards based on the number of

observations (N) per statistical estimates (q) ranging from 20:1 to 5:1 depending on

the complexity of the model (Bentler & Chou, 1987; Kline, 2015) or an absolute

sample size of 250 if using the Satorra-Bentler scaled method (Hu & Bentler, 1999).

The current study has around 8:1 N:q, which is sufficient to find a good solution to

meet the convergence criteria, but not optimal. Further studies are needed to

validate the model with a larger sample size.

This study also focussed mainly on individuals without symptomatic hearing

imapirment. The new dynamic measures will need to be tested on different

populations such as hearing impaired or patients with cochlear implants, to see if the

results can be replicated with people who struggle with SIN perception. Indeed,

recent data suggest that AFG-fixed does predict SIN performance in CI users (Choi

et al., 2023), so it is plausible that AFG-dynamic in CI users may explain even further

variance. This then can potentially be used for clinical practice to assess patients'

dynamic sound segregation. Future research can also incorporate other aspects of

SIN perception (e.g., subcortical sound analysis, language ability) and cognitive

measures (general intelligence, auditory working memory) to test if the effect of AFG

on SIN holds when accounting for these other factors.

In conclusion, these data show that an adequate model of SIN perception needs to

account for age, peripheral auditory function, and measures of grouping that we

have previously demonstrated to have a brain basis. We introduced new measures

of central grouping in this work that incorporate harmonicity and a pitch trajectory

taken from natural speech. These measures have improved the prediction of speech

in noise in the multivariate model.

References

Bentler, P. M., & Chou, C.-P. (1987). *Practical Issues in Structural Modeling*.

648   https://journals.sagepub.com/doi/10.1177/0049124187016001004

649   Besser, J., Festen, J. M., Goverts, S. T., Kramer, S. E., & Pichora-Fuller, M. K.

650   (2015). Speech-in-speech listening on the LiSN-S test by older adults with

651   good audiograms depends on cognition and hearing acuity at high

652   frequencies. *Ear and Hearing*, *36*(1), 24–41.

653   https://doi.org/10.1097/AUD.0000000000000096

654   Billings, C., & Madsen, B. (2018). A perspective on brain-behavior relationships and

655   effects of age and hearing using speech-in-noise stimuli. *Hearing Research*,

656   *369*. https://doi.org/10.1016/j.heares.2018.03.024

657   Binns, C., & Culling, J. F. (2007). The role of fundamental frequency contours in the

658   perception of speech against interfering speech. *The Journal of the Acoustical*

659   *Society of America*, *122*(3), 1765–1776. https://doi.org/10.1121/1.2751394

660   Bochner, J. H., Garrison, W. M., & Doherty, K. A. (2015). The NTID speech

661   recognition test: NSRT(®). *International Journal of Audiology*, *54*(7), 490–498.

662   https://doi.org/10.3109/14992027.2014.991976

663   Bollen, K. A., Lilly, A. G., & Luo, L. (2022). Selecting scaling indicators in structural

664   equation models (sems). *Psychological Methods*.

665   https://doi.org/10.1037/met0000530

666   Brosseau-Liard, P. E., & Savalei, V. (2014). Adjusting Incremental Fit Indices for

667   Nonnormality. *Multivariate Behavioral Research*, *49*(5), 460–470.

668   https://doi.org/10.1080/00273171.2014.933697

669   Brosseau-Liard, P. E., Savalei, V., & Li, L. (2012). An Investigation of the Sample

670   Performance of Two Nonnormality Corrections for RMSEA. *Multivariate*

Behavioral Research, *47*(6), 904–930.

https://doi.org/10.1080/00273171.2012.715252

Browne, M. W., & Cudeck, R. (1992). Alternative Ways of Assessing Model Fit.

Sociological Methods & Research, *21*(2), 230–258.

https://doi.org/10.1177/0049124192021002005

Burke, D. M., & Mackay, D. G. (1997). Memory, language, and ageing. *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences,

352(1363), 1845–1856. https://doi.org/10.1098/rstb.1997.0170

Carroll, J., & Zeng, F.-G. (2007). Fundamental frequency discrimination and speech

perception in noise in cochlear implant simulations. *Hearing Research*, *231*(1–

2), 42–53. https://doi.org/10.1016/j.heares.2007.05.004

Chadha, S., Kamenov, K., & Cieza, A. (2021). The world report on hearing, 2021.

Bulletin of the World Health Organization, *99*(4).

https://doi.org/10.2471/BLT.21.285643

Choi, I., Gander, P. E., Berger, J. I., Woo, J., Choy, M. H., Hong, J., Colby, S.,

McMurray, B., & Griffiths, T. D. (2023). Spectral Grouping of Electrically

Encoded Sound Predicts Speech-in-Noise Performance in Cochlear

Implantees. *Journal of the Association for Research in Otolaryngology: JARO*,

24(6), 607–617. https://doi.org/10.1007/s10162-023-00918-x

Dias, J., McClaskey, C., Alvey, A., Lawson, A., Matthews, L., Dubno, J. R., & Harris,

K. (2024). Effects of age and noise exposure history on auditory nerve

response amplitudes: A systematic review, study, and meta-analysis. *Hearing*

Research, *447*. https://doi.org/10.1016/j.heares.2024.109010

Elliott, L. L. (1995). Verbal auditory closure and the speech perception in noise

(SPIN) Test. *Journal of Speech and Hearing Research*, *38*(6), 1363–1376.

https://doi.org/10.1044/jshr.3806.1363

Ertürk, P., Aslan, F., & Türkyılmaz, M. D. (2023). Listening to speech-in-noise with

hearing aids: Do the self-reported outcomes reflect the behavioral speech

perception task performance? *European Archives of Oto-Rhino-Laryngology*.

https://doi.org/10.1007/s00405-023-08193-5

Gatehouse, S., & Noble, W. (2004). The Speech, Spatial and Qualities of Hearing

Scale (SSQ). *International Journal of Audiology*, *43*(2), 85–99.

Geller, J., Holmes, A., Schwalje, A., Berger, J. I., Gander, P. E., Choi, I., &

McMurray, B. (2021). Validation of the Iowa Test of Consonant Perception.

The Journal of the Acoustical Society of America, *150*(3), 2131–2153.

https://doi.org/10.1121/10.0006246

George, E. L. J., Zekveld, A. A., Kramer, S. E., Goverts, S. T., Festen, J. M., &

Houtgast, T. (2007). Auditory and nonauditory factors affecting speech

reception in noise by older listeners. *The Journal of the Acoustical Society of*

America, *121*(4), 2362–2375. https://doi.org/10.1121/1.2642072

Guo, X., Benzaquén, E., Holmes, E., Choi, I., McMurray, B., Bamiou, D.-E., Berger,

J. I., & Griffiths, T. D. (2024). *British Version of the Iowa Test of Consonant*

Perception (p. 2024.09.04.611204). bioRxiv.

https://doi.org/10.1101/2024.09.04.611204

Holmes, E., & Griffiths, T. D. (2019). 'Normal' hearing thresholds and fundamental

auditory grouping processes predict difficulties with speech-in-noise

perception. *Scientific Reports*, *9*(1), Article 1. https://doi.org/10.1038/s41598-

019-53353-5

Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure

analysis: Conventional criteria versus new alternatives. *Structural Equation*

Modeling: A Multidisciplinary Journal, *6*(1), 1–55.

https://doi.org/10.1080/10705519909540118

Huang, Y. T., Newman, R. S., Catalano, A., & Goupell, M. J. (2017). Using prosody

to infer discourse prominence in cochlear-implant users and normal-hearing

listeners. *Cognition*, *166*, 184–200.

https://doi.org/10.1016/j.cognition.2017.05.029

Kenny, D. A., Kaniskan, B., & McCoach, D. B. (2015). The Performance of RMSEA

in Models With Small Degrees of Freedom. *Sociological Methods & Research*,

44(3), 486–507. https://doi.org/10.1177/0049124114543236

Killion, M. C., Niquette, P. A., Gudmundsen, G. I., Revit, L. J., & Banerjee, S. (2004).

Development of a quick speech-in-noise test for measuring signal-to-noise

ratio loss in normal-hearing and hearing-impaired listeners. *The Journal of the*

Acoustical Society of America, *116*(4 Pt 1), 2395–2405.

https://doi.org/10.1121/1.1784440

Kline, R. B. (2015). *Principles and Practice of Structural Equation Modeling, Fourth*

Edition: Fourth Edition (4th edition). Guilford Press.

lavaan.org—Model syntax 2. (n.d.). Retrieved 8 March 2024, from

https://lavaan.ugent.be/tutorial/syntax2.html

Liu, J., Stohl, J., & Overath, T. (2024). Hidden hearing loss: Fifteen years at a

glance. *Hearing Research*, *443*, 108967.

https://doi.org/10.1016/j.heares.2024.108967

Llanos, F., German, J. S., Gnanateja, G. N., & Chandrasekaran, B. (2021). The

neural processing of pitch accents in continuous speech. *Neuropsychologia*,

158, 107883. https://doi.org/10.1016/j.neuropsychologia.2021.107883

McPherson, M. J., Grace, R. C., & McDermott, J. H. (2022). Harmonicity aids

hearing in noise. *Attention, Perception & Psychophysics*, *84*(3), 1016–1042.

https://doi.org/10.3758/s13414-021-02376-0

Meha-Bettison, K., Sharma, M., Ibrahim, R. K., & Mandikal Vasuki, P. R. (2018).

Enhanced speech perception in noise and cortical auditory evoked potentials

in professional musicians. *International Journal of Audiology*, *57*(1), 40–52.

https://doi.org/10.1080/14992027.2017.1380850

Merten, N., Boenniger, M. M., Herholz, S. C., & Breteler, M. M. B. (2022). The

Associations of Hearing Sensitivity and Different Cognitive Functions with

Perception of Speech-in-Noise. *Ear and Hearing*, *43*(3), 984–992.

https://doi.org/10.1097/AUD.0000000000001154

Micheyl, C., Ryan, C. M., & Oxenham, A. J. (2012). Further evidence that

fundamental-frequency difference limens measure pitch discrimination. *The*

Journal of the Acoustical Society of America, *131*(5), 3989–4001.

https://doi.org/10.1121/1.3699253

Nilsson, M., Soli, S. D., & Sullivan, J. A. (1994). Development of the Hearing in Noise

Test for the measurement of speech reception thresholds in quiet and in

noise. *The Journal of the Acoustical Society of America*, *95*(2), 1085–1099.

https://doi.org/10.1121/1.408469

Oberfeld, D., & Klöckner-Nowotny, F. (2016). Individual differences in selective

attention predict speech identification at a cocktail party. *eLife*, *5*, e16747.

https://doi.org/10.7554/eLife.16747

O'Sullivan, J. A., Shamma, S. A., & Lalor, E. C. (2015). Evidence for Neural

Computations of Temporal Coherence in an Auditory Scene and Their

Enhancement during Active Listening. *The Journal of Neuroscience: The*

Official Journal of the Society for Neuroscience, *35*(18), 7256–7263.

https://doi.org/10.1523/JNEUROSCI.4973-14.2015

Oxenham, A. J. (2008). Pitch Perception and Auditory Stream Segregation:

Implications for Hearing Loss and Cochlear Implants. *Trends in Amplification*,

12(4), 316–331. https://doi.org/10.1177/1084713808325881

Schneider, F., Dheerendra, P., Balezeau, F., Ortiz-Rios, M., Kikuchi, Y., Petkov, C.

I., Thiele, A., & Griffiths, T. D. (2018). Auditory figure-ground analysis in

rostral belt and parabelt of the macaque monkey. *Scientific Reports*, *8*(1),

17948. https://doi.org/10.1038/s41598-018-36903-1

Shen, J., & Souza, P. E. (2018). On Dynamic Pitch Benefit for Speech Recognition in

Speech Masker. *Frontiers in Psychology*, *9*, 1967.

https://doi.org/10.3389/fpsyg.2018.01967

Teki, S., Barascud, N., Picard, S., Payne, C., Griffiths, T. D., & Chait, M. (2016).

Neural Correlates of Auditory Figure-Ground Segregation Based on Temporal

Coherence. *Cerebral Cortex (New York, N.Y.: 1991)*, *26*(9), 3669–3680.

https://doi.org/10.1093/cercor/bhw173

Teki, S., Chait, M., Kumar, S., Shamma, S., & Griffiths, T. D. (2013). Segregation of

complex acoustic scenes based on temporal coherence. *eLife*, *2*, e00699.

https://doi.org/10.7554/eLife.00699

Teki, S., & Griffiths, T. D. (2016). Brain Bases of Working Memory for Time Intervals

in Rhythmic Sequences. *Frontiers in Neuroscience*, *10*, 239.

https://doi.org/10.3389/fnins.2016.00239

792 Wilson, R. H., McArdle, R., Watts, K. L., & Smith, S. L. (2012). The Revised Speech

793   Perception in Noise Test (R-SPIN) in a multiple signal-to-noise ratio paradigm.

794   *Journal of the American Academy of Audiology*, *23*(8), 590–605.

795   https://doi.org/10.3766/jaaa.23.7.9

796 Wong, L. L. N., Cheung, C., & Wong, E. C. M. (2008). Comparison of hearing

797   thresholds obtained using pure-tone behavioral audiometry, the Cantonese

798   Hearing in Noise Test (CHINT) and cortical evoked response audiometry.

799   *Acta Oto-Laryngologica*, *128*(6), 654–660.

800   https://doi.org/10.1080/00016480701642189

801 Xie, R., Wang, M., & Zhang, C. (2024). Mechanisms of age-related hearing loss at

802   the auditory nerve central synapses and postsynaptic neurons in the cochlear

803   nucleus. *Hearing Research*, *442*, 108935.

804   https://doi.org/10.1016/j.heares.2023.108935

Author contributions

Xiaoxuan Guo: conceptualisation, study design, data collection, data analysis,

manuscript writing

Ester Benzaquén: conceptualisation, study design

Emma Holmes: study design

Joel I Berger: study design

Inga Brühl: data collection

William Sedley: conceptualisation, study design

Steven Rushton (co-senior author): data analysis

Timothy D Griffiths (co-senior author): conceptualisation, study design, data analysis,

funding.

All authors have edited the manuscript and approved its submission to this journal.

Data availability: Data will be available upon request.

Additional Information: This work was supported by the MRC [grant number

MR/T032553/1]. There are no conflicts of interest, financial, or otherwise.

821