A can of worms: estimating the global number of earthworm species

Running title: How many earthworm species?

Abstract

Estimating the overall species number for a given taxon is a central issue in ecology and conservation biology. This is especially topical for soil organisms, which comprise the majority of known species but whose taxonomy remains largely understudied. We propose here an estimate of the global number of earthworm species based on the Joppa approach, which models taxonomic effort over time to estimate the total number of known and as yet unknown species in a given taxa. Our Bayesian estimation of the Joppa model suggests a global diversity of the order of 30,000 species, suggesting that the 5,679 earthworm species and subspecies already described only represent around 20% of the actual global species diversity. However, the uncertainty around this estimate is considerable due to severe undersampling and as the model cannot unambiguously decide whether we are describing few species because of a small pool of as yet unknown species, or because of a lack of taxonomic efficiency. Considering the current rate of new species description, we calculate that it would take at least 120 years to describe all the earthworm species existing on Earth. Different strategies should be developed to accelerate the facilitate and accelerate the discovery and naming of species new to science and to reduce the magnitude of this taxonomic deficit.

Keywords

Soil biodivtersity; global species number estimates; taxonomic deficit; French Guiana; temperate France

INTRODUCTION

Estimating the total number of species of a given taxon, whether on a global scale or for a given region or ecosystem, is a question of great practical and symbolic importance that has long captivated and challenged scientists (May, 1992; Stork, 1993; Pimm, 2001; Dirzo & Raven, 2003). It represents an indispensable step towards a comprehensive description of the World's biodiversity and is also necessary for the definition of efficient conservation strategies (Joppa *et al.*, 2011; Caley *et al.*, 2014). In this context, soil biota represents one of the most diversified and least studied compartments in terrestrial ecosystems. They are among the most diverse, comprising up to 59% of currently known living species (Decaëns *et al.*, 2006; Anthony *et al.*, 2023). They are involved in regulating a large number of ecological processes that are essential to the provision of key ecosystem services for human populations (Brussaard *et al.*, 1997; Wardle *et al.*, 2004; Bardgett & van der Putten, 2014). This has been particularly well documented for emblematic groups of soil ecosystem engineers, such as earthworms (Lavelle *et al.*, 2006, 2016). Paradoxically, even these well-studied groups still suffer from a glaring lack of basic studies on their biodiversity patterns (Cameron *et al.*, 2018; Phillips *et al.*, 2019, 2021), and questions as fundamental as "how many species are there?" still remain largely unanswered (Decaëns, 2010). This knowledge gap is of particular concern, as it limits our capacity to characterize the dynamics of soil biodiversity in the face of ongoing global change and to assess its implications for ecosystem functioning.

Earthworms are an emblematic illustration of this paradox. Some 5,700 earthworm species have been described to date (Misirlioğlu *et al.*, 2023), but the proportion of the actual diversity of these organisms that this figure represents is still poorly documented. Estimates vary, depending on the author, from a minimum of 7,000 to some 20,000 species (Reynolds, 1994; Lavelle *et al.*, 2006; Blakemore, 2009; Anthony *et al.*, 2023), but most of these figures are based on expert opinion, and no mathematical formalisation has yet been used to obtain a consolidated prediction of the number of as-yet undiscovered species. Some studies have

also highlighted the existence of potentially high and as yet largely undescribed regional

diversity throughout the tropics (Phillips *et al.*, 2019; Goulpeau *et al.*, 2024), which could

challenge these previously proposed estimates of global species number.

Various methods can be used to estimate the number of heretofore unknown species within

a given taxon. One popular approach is to use rarefaction and extrapolation curves,

combined with non-parametric estimators, to predict the taxonomic deficit and the number of

species at different spatial scales (Chao, 1984; Colwell *et al.*, 2004; Chao *et al.*, 2005; Chao

& Jost, 2012; Cazzolla-Gatti *et al.*, 2022). However, to be effective, this approach requires a

sufficiently dense dataset on a global scale, which is lacking for many soil organisms such

as earthworms. Another method consists of using Arrhenius' power law to estimate the

number of species for large areas on the basis of a known number of species in a smaller

area (Arrhenius, 1921; Krishnamani *et al.*, 2004; Marcon *et al.*, 2024). The procedure

involves estimating a power parameter (z) based on the negative linear relationship between

the log of compositional similarity and the log of geographical distance between small

equivalent areas. It therefore requires a network of plots located at very different distances,

which must be supplemented by a continuous inventory over an area corresponding to the

smallest scale of the power law. Apart from the fact that this type of data is rare, log

transformation is impossible for pairs of plots with a compositional similarity of 0 (i.e. no

species in common), which is common in the case of earthworms, particularly in the tropics

(Goulpeau *et al.*, 2024). Finally, Joppa *et al.* (2011) propose to use global species checklists

to model the rate at which taxonomists describe new species as a function of time, and to

estimate the number of species that will have been reached when this rate will drop to zero

due to the exhaustion of the pool of as yet unknown species. This approach has the

advantage of not being dependent on the existence of comprehensive field datasets, but

requires the establishment of reliable species checklists that can be challenged by

taxonomic issues.

In this work, we propose a global estimate of the number of earthworm species based on the method of Joppa *et al.* (2011) and the most recent global checklist of earthworms (Brown *et al.*, 2023; Misirlioğlu *et al.*, 2023). In parallel, we analysed three datasets obtained from field inventories for regions equivalent in surface area in tropical and temperate regions (Bouché, 1972; Goulpeau *et al.*, 2024; Gérard *et al.*, 2024). We used these data to quantify the extent of the taxonomic deficit and estimate the actual number of species for each region. We then used these regional estimates to discuss the range of results obtained from the Joppa method.

MATERIAL AND METHODS

Estimating the number of species from the global checklist

For the Joppa approach, we used the checklist recently published by Misirlioğlu *et al.* (2023) and Brown *et al.* (2023). This list includes 5,738 species/subspecies (5,406 species and 332 unique subspecies, i.e. not counting nomino-typical subspecies) belonging to 23 families (including one non-crassiclitellate family: Moniligastridae). We chose to consider subspecies as taxa in their own right in the analysis, because the choice of whether to consider them as infraspecific divisions or to elevate them to species rank is a matter of taxonomists' points of view. For the sake of simplicity, we referred to all the taxa in the list as "species". After deleting certain duplicates in the table, we obtained a consolidated list of 5,752 taxa names published between 1758 and 2023, containing for each of them the authors and the year of description. For each 5-year period, we then counted the number of unique species described and the total number of authors (i.e. all the co-authors of the species, after standardising the names of authors with accents and taking account of homonyms, wherever possible).

Joppa *et al.* (2011) expected the number of species (S_i) described in a given interval ($Y_i = i$, numbered sequentially from 1) to depend on the taxonomic effort, i.e. the number of taxonomists (T_i) actively describing species during that period, and the unknown number of species remaining to be described, that is the total number of species (S_T) minus those

already discovered ($\sum_{j=0}^{i} S_j$ where S_0 is the number of known species before Y_1). Over time,

taxonomists were expected to become more efficient, i.e. the discovery rate of unknown

species per taxonomist was assumed to increase linearly. The original model was thus

$S_i \sim T_i(\beta_1 + \beta_2 Y_i)\left(S_T - \sum_{j=0}^{i} S_j\right)$,

where β_1 and β_2 are unknown parameters. The model was log-transformed and estimated

with a normal error of variance σ^2. The parameters to be estimated were thus β_1, β_2, S_T and

σ, the total number of species S_T being our main target.

The increase in efficiency accounted for in the Joppa model is not obvious as earthworms

are concerned. Methodological advancements, such as the development of integrative

taxonomy, are likely offset by the increased difficulty of sampling in remote areas, namely

tropical rainforests. Moreover, contemporary scientists rarely focus exclusively on taxonomy,

a discipline whose publications are generally undervalued, leaving them with less time to

describe new species. Consequently, we studied an alternative model with no efficiency

progress (hereafter referred to as the "no-progress" model), where β_2 equals zero.

We chose Bayesian inference to obtain the distribution of the estimates of the parameters

which revealed to be as important as the most probable values. We chose non-informative

priors, uniformly distributed between 0 and numerically possible values β_1, β_2 and σ.

According to expert opinion, the prior of S_T was forced between 6,000, i.e. the number of

already discovered species, and 100,000. The parameter space was explored by a Monte-

Carlo Markov Chain with 10^7 steps, 10% considered as burn-in. The analysis began with

time period 1822-1827, with a previously known number of species (S_0) equal to 1 (i.e.

Lumbricus terrestris Linnaeus, 1758). The last period is 2018-2022; species published in

2023 and 2024 were excluded from the analysis.

Estimating regional diversity and the taxonomic deficit

Compared to the checklist of angiosperms used by Joppa *et al.* (2011), that of earthworms is very short: data is scarce. We endeavoured to supplement the global approach with three more exhaustive regional studies.

For the analyses of regional diversity, we used the data of Goulpeau *et al.* (2022, 2024) in French Guiana (supplemented by a small number of as yet unpublished data), and, for Southern and Northern France, those of Bouché (1972) updated for taxonomy by Gérard *et al.* (2024). The choice of these three regions allows us to consider contrasting situations in terms of earthworm diversity. French Guiana is considered to be a diversity hotspot, with a high proportion of species with restricted distribution and a large taxonomic deficit (Decaëns *et al.*, 2016, 2024; Maggia *et al.*, 2021; Goulpeau *et al.*, 2024). By comparison, mainland France has a lower *a priori* species diversity, but the South is nevertheless characterised by a high proportion of endemic species, some of which continue to be discovered regularly, whereas the North is characterised by great compositional uniformity (Bouché, 1972; Mathieu & Davies, 2014; Gérard *et al.*, 2023; Marchán *et al.*, 2023).

Three 300 x 400 km^2 zones, each subdivided into twelve 100 x 100 km^2 cells, were delimited in each region. In each 100 x 100 km^2 cell, we counted the species present in order to obtain three occurrence tables comprising 12 rows and as many columns as species present. In French Guiana, we used the operational taxonomic units (OTUs) proposed by Goulpeau *et al.* (2022) as a substitute for species in order to overcome the problem posed by the taxonomic deficit. For each region separately, we plotted incidence-based rarefaction and extrapolation curves in order to represent the accumulation of species richness as a function of the number of 100 x 100 km^2 cells sampled in the study areas. The 95% confidence intervals of the interpolated and extrapolated richness values were obtained by a bootstrap procedure with 200 replications. We also calculated the non-parametric Chao 2 index, which estimates the lower limit of expected asymptotic species richness (Gotelli & Chao, 2013).

The taxonomic deficit, i.e. the proportion of undiscovered species, was calculated using the

formula:

$\text{TDef} = ((S_{EST} - S_{OBS}) \times 100 / S_{EST}$,

with $TDef$ the taxonomic deficit, S_{OBS} the number of species currently known for a given

region, and S_{EST} the richness estimated by the Chao 2 index.

For French Guiana, the taxonomic deficit was calculated by first considering S_{OBS} as the set

of OTUs delimited by Goulpeau *et al.* (2022), and then by restricting S_{OBS} to the number of

species formally described to date (i.e. 55 species taking into account the 18 new

descriptions by Decaëns *et al.* (2024). These analyses were performed using the *'iNEXT'*

package for R (Hsieh *et al.*, 2019; R Core Team, 2024).

RESULTS

During the period from 1822 to 2022, the number of species described per 5-year period

showed two peaks: the first from the end of the 19th to the middle of the 20th century, the

other from the 1970s to the present day (Fig. 1A). Over the same period, the total number of

authors of original species descriptions has increased exponentially (i.e. roughly linearly on a

logarithmic scale; Fig. 1A). The ratio of numbers of species described per taxonomist thus

shows a very marked peak in the first half of the 20th century, a period during which it

generally exceeds 15 species described per taxonomist per time period, whereas

everywhere else it is generally below 10 species per taxonomist (Fig. 1B).

The Joppa model yields the median posterior S_T equal to 6,814 species, with a 95% higher

posterior density interval (HPDI) between 6,000 and 11,148. The no-progress model leads to

quite different estimations: 26,283 species (HDPI between 6,003 and 87,425). Posterior

probabilities are shown in Fig. 2.

The rarefaction curves obtained for the three study regions show a decrease in beta diversity

from tropical to temperate study regions (Fig. 3). In French Guiana, the slope of the curve is

steep, showing that the asymptote is far from being reached, whereas it has been or is about

to be reached in the two study regions in temperate France. The diversity metrics logically

follow the same trend: the cumulative number of species observed is 305 in French Guiana,

66 in Southern France, and 31 in Northern France, whereas the Chao 2 index predicts a

diversity of 1,680 species in French Guiana, 99 in Southern France, and 36 in Northern

France (Fig. 4A, 4B). Consequently, the taxonomic deficit is of the order of 80% in French

Guiana, 33% in the south of France, and 16% in the north of France (Fig. 4C). In French

Guiana, this 80% taxonomic deficit was obtained by considering as observed richness all of

the OTUs mentioned in Goulpeau *et al.* (2022), most of which are new to science and have

not yet been named. If we consider only the 55 named species actually known from French

Guiana (Fig. 4A), the estimated taxonomic deficit rises to 97% (Fig. 4C).

DISCUSSION

Earthworm taxonomy experienced a golden age from the end of the 19th century to the

middle of the 20th century, followed by a revival from the end of the 1970s. The first of these

two periods can be credited to the great explorations that enabled zoologists of the time to

explore the world and describe its biological diversity (Csuzdi & Szlávecz, 2016). It was also

made possible by the exceptional productivity of the few taxonomists of the time, who were

able to describe dozens of species in a short time. By comparison, the contemporary period

is characterised by a larger community of taxonomists, but with comparatively lower output.

Overall, we do not observe the monotonic decline in the number of species described per

taxonomist as described for plants and other animal taxa (Pimm *et al.*, 2010; Joppa *et al.*,

2011; Fisher *et al.*, 2018), which is assumed to reflect a depletion of the pool of new species

to be discovered. Rather, with the exception of the notable peak in the first half of the 20th

century, the rate of description of new earthworm species remained modest and constant

over time, at around 5 species per taxonomist per 5-year period. Even in the early 2000s, we

did not observe any clear effect from the development of molecular tools such as DNA

barcoding and metabarcoding (*sensu* Hebert *et al.*, 2003; Taberlet *et al.*, 2012). This may

seem surprising, given that their usefulness in the field of earthworm taxonomy has been

widely acknowledged (Rougerie *et al.*, 2009; Decaëns *et al.*, 2013; Marchán *et al.*, 2022),

and their advent has been touted as the possible catalyst of a new golden age of taxonomy

(Hebert & Gregory, 2005; Hajibabaei, 2012; Pennisi, 2019). Over the last few decades, the

increase in the number of new earthworm species described per time period seems to be

simply correlated with the number of active taxonomists.

Our estimates of the Joppa and the no-progress model tell us two different stories. Assuming

a constant progress of the efficiency of taxonomists, the global number of species would be

below 7,000, i.e. around 1,000 species remaining to be described. On the other hand,

assuming constant efficiency, the global number of species would rather be around 30,000,

i.e. the 5,679 species of our checklist are only the beginning of the story.

In both cases, the confidence intervals around our median estimates are considerable,

motivating us not to provide a single significant figure in our estimates. This can be

explained by two main causes of uncertainty. First, we lack data: more publications of new

species would have allowed smaller, i.e. more, time intervals, thus more statistical power.

Also, the Joppa model does not allow to completely disentangle the effects of the efficiency

and the number of species: in the no-progress model, the estimates are highly correlated

(Pearson correlation of the posteriors: -0,73), and the correlation is -0,55 between β_2 and S_T

in the original model. This issue could not be detected by Joppa *et al.* (2011) who relied on a

frequentist estimation of the most likely number of species.

The analysis of regional diversity in French Guiana points in the direction of a high number

of species: we estimate that the taxonomic deficit there is of the order of 97%, if only

currently named species are considered in the calculation. An unknown thousand species

may well all be there: a small area (actually around 1%) of Amazonia may thus be enough to

find the species missing from the Joppa model. If we consider that around half of the

earthworm species currently known are tropical (*i.e.* around 2,800 species), the

generalisation of such a taxonomic deficit would lead us to estimate that more than 90,000

species could exist throughout the tropics. Of course, such a generalisation is highly

debatable, but it does provide an argument for considering that the global number of

unknown earthworm species may not be only a few thousands.

As expected, we calculate a lower taxonomic deficit in temperate regions, which have been

better studied and host a more modest diversity. However, our results indicate that this

deficit can still be significant—ranging from 15 to 30% in mainland France—suggesting that

even in temperate regions, a substantial number of species remains undiscovered. The

South of France, known for its high diversity and richness in endemic earthworms,

exemplifies this, as evidenced by the recent descriptions of several new species, including

some of large body size (Marchán *et al.*, 2023). The persistence of a significant taxonomic

deficit in Northern France is more unexpected. It may reflect a gap in our understanding of

species distribution in the region (i.e. the Wallacean shortfall according to Lomolino, 2004),

or the persistence of taxonomic confusion that can conceal the presence of certain species.

A striking example of this is the discovery of the existence of two cryptic species that were

mistakenly considered synonymous under the name *L. terrestris* (James *et al.*, 2010).

These considerations allow us to propose a conservative estimate of the global number of

earthworm species of the order of 30,000 species. This is above the upper range of

commonly accepted predictions, i.e. from 10,000 to 20,000 species (Reynolds, 1994;

Decaëns *et al.*, 2006; Blakemore, 2009; Anthony *et al.*, 2023). Our expert opinion is that

there is still a large pool of undiscovered species, but that we lack the efficiency to discover

them or describe them once they have been discovered. First, a low rate of new species

description per taxonomist, compared with those active at the beginning of the 20th century,

is expected due the fact that taxonomy is no longer considered a 'competitive' discipline in

modern science (Wilson, 2004; Agnarsson & Kuntner, 2007). The pernicious effect of this

situation is that contemporary authors are rarely pure taxonomists, as they are encouraged

by the academic system to devote most of their research time to more 'valuable' areas of

research, such as evolution or ecology. Added to this is the fact that taxonomists generally

lack the time and financial resources for field sampling, particularly in the most diverse and

yet least well-studied tropical regions (Guerra *et al.*, 2020). The field scientists among us all

have in mind regions that they know contain a diversity of as yet unknown species, but which

they do not have the time or the resources to explore. Similarly, taxonomists have a

multitude of species in their cupboards waiting to be described, but which they do not have

the time to process.

Overall, our results suggest that at best we have only described one fifth of the global

number of earthworm species. If we consider the current community of taxonomists (an

average of 45 active taxonomists per 5-year period over the last 20 years), and the number

of species they have managed to describe (an average of 4.5 species per taxonomist per

time step), we calculate that it would take at least 120 years to describe all the species

existing on Earth. Obviously, such an objective is certainly unattainable within a reasonable

timeframe, but we can nonetheless think about the strategies that could be developed to

reduce the magnitude of the taxonomic deficit. This will require maintaining a major

taxonomic effort, through the training of young taxonomists, but also by raising the profile of

taxonomic work in the scientific corpus. Agnarsson & Kuntner (2007) have proposed that

original descriptions, and their underlying taxonomic hypotheses, should be formally cited in

articles that refer to them – which is not systematically the case at present. This would allow

the impact metrics of taxonomic journals to improve while more accurately reflecting their

real scientific impact, thus encouraging scientists to invest more in taxonomy. We will also

need to improve taxonomic efficiency, first by increasing our ability to discover as yet

unknown species where they occur. This means stepping up sampling efforts, particularly in

the tropics where most of this undiscovered diversity is to be found. Our results confirm the

existence of a regional peak in diversity at tropical latitudes, essentially explained by a

strong beta component (Phillips *et al.*, 2019; Goulpeau *et al.*, 2024), while emphasising that

it mirrors a considerable taxonomic deficit. Finally, once sampling coverage has been

improved, new taxonomic approaches based on turbotaxonomy (i.e. the rapid description of

species new to science allowed by automated procedures and the use of molecular tools;

Butcher *et al.*, 2012; Fernandez-Triana, 2022) and integrative taxonomy (i.e. the integrated

use of multiple, independent lines of evidence—e.g. morphology, genetics, ecology, and

behavior—for species delimitation and description; Dayrat, 2005) will be necessary to

improve our ability to describe new species at a consistent rate (Padial *et al.*, 2010;

Fernandez-Triana, 2022).

In conclusion, we propose an updated estimate of the global number of earthworm species,

as well as arguments for considering earthworms as a potentially hyperdiverse group of

organisms, which has hitherto not been considered as such due to a strong taxonomic

knowledge deficit (i.e the Linnean shortfall as defined in Lomolino, 2004). This is particularly

remarkable when considering that earthworms are among the most widely publicised soil

macro-organisms, due to their acknowledged importance in soil functioning. It is likely that

this situation could also apply to other less well-studied soil taxa, leading to a considerable

reassessment of the contribution of these organisms to global biodiversity.

REFERENCES

Agnarsson, I. & Kuntner, M. (2007) Taxonomy in a changing world: Seeking solutions for a

science in crisis. *Systematic Biology*, **56**, 531–539.

Anthony, M.A., Bender, S.F. & van der Heijden, M.G.A. (2023) Enumerating soil biodiversity.

Proceedings of the National Academy of Sciences of the United States of America, **120**,

e2304663120.

Arrhenius, O. (1921) Species and area. *Journal of Ecology*, **9**, 95–99.

Bardgett, R.D. & van der Putten, W.H. (2014) Belowground biodiversity and ecosystem

functioning. *Nature*, **515**, 505–511.

Blakemore, R.J. (2009) *Chapter 14. Cosmopolitan earthworms – a global and historical*

perspective. Annelids as model systems in the biological sciences (ed. by D.H. Shain), pp.

257–283. John Wiley & Sons, Inc., New York.

Bouché, M.B. (1972) *Lombriciens de France*, INRA, Paris.

Brown, G.G., James, S.W., Csuzdi, C., Lapied, E., Decaëns, T., Reynolds, J.W., Misirlioğlu,

M., Stovanic, M., Trakić, T., Sekulić, J., Phillips, H.R.P. & Cameron, E.K. (2023) A checklist

of megadrile earthworm (Annelida: Clitellata) species and subspecies of the world. *Zedono*,

https://doi.org/10.5281/zenodo.8348860.

Brussaard, L., Behan-Pelletier, V.M., Bignell, D.E., Brown, V.K., Didden, W., Folgarait, P.,

Fragoso, C., Freckman, D.W., Gupta, V., Hattori, T., Hawksworth, D.L., Klopatek, C.,

Lavelle, P., Malloch, D.W., Rusek, J., Soderstrom, B., Tiedje, J.M. & Virginia, R.A. (1997)

Biodiversity and ecosystem functioning in soil. *AMBIO*, **26**, 563–570.

Butcher, B.A., Smith, M.A., Sharkey, M.J. & Quicke, D.L.J. (2012) A turbo-taxonomic study

of Thai Aleiodes (Aleiodes) and Aleiodes (Arcaleiodes) (Hymenoptera: Braconidae:

Rogadinae) based largely on COI barcoded specimens, with rapid descriptions of 179 new

species. *Zootaxa*, 1–232.

Caley, M.J., Fisher, R. & Mengersen, K. (2014) Global species richness estimates have not

converged. *Trends in Ecology & Evolution*, **29**, 187–188.

Cameron, E.K., Martins, I.S., Lavelle, P., Mathieu, J., Tedersoo, L., Gottschall, F., Guerra,

C.A., Hines, J., Patoine, G., Siebert, J., Winter, M., Cesarz, S., Delgado-Baquerizo, M.,

Ferlian, O., Fierer, N., Kreft, H., Lovejoy, T.E., Montanarella, L., Orgiazzi, A., Pereira, H.M.,

Phillips, H.R.P., Settele, J., Wall, D.H. & Eisenhauer, N. (2018) Global gaps in soil

biodiversity data. *Nature Ecology & Evolution*, **2**, 1042–1043.

Cazzolla-Gatti, R., Reich, P.B., Gamarra, J.G.P., Crowther, T., Hui, C., Morera, A., Bastin,

J.-F., de-Miguel, S., Nabuurs, G.-J., Svenning, J.-C., Serra-Diaz, J.M., Merow, C., Enquist,

B., Kamenetsky, M., Lee, J., Zhu, J., Fang, J., Jacobs, D.F., Pijanowski, B., Banerjee, A.,

Giaquinto, R.A., Alberti, G., Almeyda Zambrano, A.M., Alvarez-Davila, E., Araujo-Murakami,

A., Avitabile, V., Aymard, G.A., Balazy, R., Baraloto, C., Barroso, J.G., Bastian, M.L.,

Birnbaum, P., Bitariho, R., Bogaert, J., Bongers, F., Bouriaud, O., Brancalion, P.H.S.,

Brearley, F.Q., Broadbent, E.N., Bussotti, F., Castro da Silva, W., Gomes Cesar, R., Cesljar,

G., Moscoso, V.C., Chen, H.Y.H., Cienciala, E., Clark, C.J., Coomes, D.A., Dayanandan, S.,

Decuyper, M., Dee, L.E., Del Aguila Pasquel, J., Derroire, G., Djuikouo, M.N.K., Tran, V.D.,

Dolezal, J., Dordevic, I.D., Engel, J., Fayle, T.M., Feldpausch, T.R., Fridman, J.K., Harris,

D.J., Hemp, A., Hengeveld, G., Herault, B., Herold, M., Ibanez, T., Jagodzinski, A.M.,

Jaroszewicz, B., Jeffery, K.J., Johannsen, V.K., Jucker, T., Kangur, A., Karminov, V.N.,

Kartawinata, K., Kennard, D.K., Kepfer-Rojas, S., Keppel, G., Khan, M.L., Khare, P.K.,

Kileen, T.J., Kim, H.S., Korjus, H., Kumar, A., Kumar, A., Laarmann, D., Labriere, N., Lang,

M., Lewis, S.L., Lukina, N., Maitner, B.S., Malhi, Y., Marshall, A.R., Martynenko, O.V.,

Mendoza, A.L.M., Ontikov, P.V., Ortiz-Malavasi, E., Camacho, N.C.P., Paquette, A., Park,

M., Parthasarathy, N., Peri, P.L., Petronelli, P., Pfautsch, S., Phillips, O.L., Picard, N., Piotto,

D., Poorter, L., Poulsen, J.R., Pretzsch, H., Ramirez-Angulo, H., Correa, Z.R., Rodeghiero,

M., Gonzales, R.D.P.R., Rolim, S.G., Rovero, F., Rutishauser, E., Saikia, P., Salas-Eljatib,

C., Schepaschenko, D., Scherer-Lorenzen, M., Seben, V., Silveira, M., Slik, F., Sonke, B.,

Souza, A.F., Sterenczak, K.J., Svoboda, M., Taedoumg, H., Tchebakova, N., Terborgh, J.,

Tikhonova, E., Torres-Lezama, A., van der Plas, F., Vasquez, R., Viana, H., Vibrans, A.C.,

Vilanova, E., Vos, V.A., Wang, H.-F., Westerlund, B., White, L.J.T., Wiser, S.K., Zawila-

Niedzwiecki, T., Zemagho, L., Zhu, Z.-X., Zo-Bi, I.C. & Liang, J. (2022) The number of tree

species on Earth. *Proceedings of the National Academy of Sciences of the United States of*

America, **119**, e2115329119.

Chao, A. (1984) Nonparametric estimation of the number of classes in a population.

Scandinavian Journal of Statistics, **11**, 265–270.

Chao, A., Chazdon, R.L., Colwell, R.K. & Shen, T.J. (2005) A new statistical approach for

assessing similarity of species composition with incidence and abundance data. *Ecology*

Letters, **8**, 148–159.

Chao, A. & Jost, L. (2012) *Diversity measures. Encyclopedia of Theoretical Ecology* (ed. by

A. Hastings) and L. Gross), pp. 203–207. University of California Press, Berkeley, USA.

Colwell, R.K., Mao, C.X. & Chang, J. (2004) Interpolating, extrapolating, and comparing

incidence-based species accumulation curves. *Ecology*, **85**, 2717–2727.

Csuzdi, C. & Szlávecz, K. (2016) Earthworm (Clitellata: Megadrili) taxonomy in the last 200

years: A homage to András Zicsi (1928–2015). *Opuscula Zoologica (Budapest)*, **47**, 1–7.

Dayrat (2005) Towards integrative taxonomy. *Biological Journal of the Linnean Society*, **85**,

407–417.

Decaëns, T. (2010) Macroecological patterns in soil communities. *Global Ecology and*

Biogeography, **19**, 287–302.

Decaëns, T., Bartz, M.L.C., Goulpeau, A., Marchan, D.F., Maggia, M.-E., Lapied, E., Feijoo,

A.M. & James, S.W. (2024) Earthworms (Oligochaeta: Clitellata) of the Mitaraka range

(French Guiana): commented checklist with description of one genus and eighteen species

new to science. *Zoosystema*, **46**, 195–244.

Decaëns, T., Jiménez, J.J., Gioia, C., Measey, J. & Lavelle, P. (2006) The values of soil

animals for conservation biology. *European Journal of Soil Biology*, **42**, S23–S38.

Decaëns, T., Porco, D., James, S.W., Brown, G.G., Chassany, V., Dubs, F., Dupont, L.,

Lapied, E., Rougerie, R., Rossi, J.-P. & Roy, V. (2016) DNA barcoding reveals diversity

patterns of earthworm communities in remote tropical forests of French Guiana. *Soil Biology*

& Biochemistry, **92**, 171–183.

Decaëns, T., Porco, D., Rougerie, R., Brown, G.G. & James, S.W. (2013) Potential of DNA

barcoding for earthworm research in taxonomy and ecology. *Applied Soil Ecology*, **65**, 35–

42.

Dirzo, R. & Raven, P. (2003) Global state of biodiversity and loss. *Annu Rev Environ*

Resour, **28**, 137–167.

Fernandez-Triana, J.L. (2022) Turbo taxonomy approaches: lessons from the past and

recommendations for the future based on the experience with Braconidae (Hymenoptera)

parasitoid wasps. *Zootaxa*, **1087**.

Fisher, M.A., Vinson, J.E., Gittleman, J.L. & Drake, J.M. (2018) The description and number

of undiscovered mammal species. *Ecology and Evolution*, **8**, 3628–3635.

Gérard, S., Decaëns, T., Marchán, D.F., Beauchesne, M., Berlioz, L., Capowiez, Y., Clause,

J., Decaëns, L., Dellavedova, R., Duhaut, C.-B., Garnier-Fière, C., Goulpeau, A.,

Goussopoulos, J., Iannelli, M., Marsden, C., Navarro, A., Orrière, S., Revertégat, C.,

Vannier, A., Versavel, C. & Hedde, M. (2024) #Vers2022: a dataset of species composition

and individual mass of earthworm communities resampled 50 years after Bouché's historical

work. *Ecology*, **in press**.

Gérard, S., Marchán, D.F., Martinez Navarro, A., Hedde, M. & Decaëns, T. (2023)

Resampling Bouché's historical localities reveals three new species and helps identifying a

new genus of earthworms (Oligochaeta, Hormogastridae and Lumbricidae) in Southeastern

France. *Zoosystema*, **45**, 749–768.

Gotelli, N. & Chao, A. (2013) *Measuring and Estimating Species Richness, Species*

Diversity, and Biotic Similarity from Sampling Data. Encyclopedia of Biodiversity (ed. by S.A.

Levin), pp. 195–211. Academic Press, Waltham, MA.

Goulpeau, A., Hedde, M., Maggia, M.-E., Gérard, S., Lapied, E. & Decaëns, T. (2024) A

dataset on earthworm communities in French Guiana. *Soil Organisms*, **92**, 79–90.

Goulpeau, A., Penel, B., Maggia, M.-E., Marchán, D.F., Steinke, D., Hedde, M. & Decaëns,

T. (2022) OTU delimitation with earthworm DNA barcodes: a comparison of methods.

Diversity, **14**, 866.

Guerra, C.A., Heintz-Buschart, A., Sikorski, J., Chatzinotas, A., Guerrero-Ramirez, N.,

Cesarz, S., Beaumelle, L., Rillig, M.C., Maestre, F.T., Delgado-Baquerizo, M., Buscot, F.,

Overmann, J., Patoine, G., Phillips, H.R.P., Winter, M., Wubet, T., Kuesel, K., Bardgett,

R.D., Cameron, E.K., Cowan, D., Grebenc, T., Marin, C., Orgiazzi, A., Singh, B.K., Wall,

D.H. & Eisenhauer, N. (2020) Blind spots in global soil biodiversity and ecosystem function

research. *Nature Communications*, **11**, 3870.

Hajibabaei, M. (2012) The golden age of DNA metasystematics. *Trends in Genetics*, **28**,

535–537.

Hebert, P.D.N., Cywinska, A., Ball, S.L. & DeWaard, J.R. (2003) Biological identifications

through DNA barcodes. *Proceedings of the Royal Society B-Biological Sciences*, **270**, 313–

321.

Hebert, P.D.N. & Gregory, T.R. (2005) The promise of DNA barcoding for taxonomy.

Syestematic Biology, **54**, 852–859.

Hsieh, T.C., Ma, K.H. & Chao, A. (2019) iNEXT: iNterpolation and EXTrapolation for species

diversity. R package version 2.0.19.

James, S.W., Porco, D., Decaens, T., Richard, B., Rougerie, R. & Erseus, C. (2010) DNA

Barcoding Reveals Cryptic Diversity in Lumbricus terrestris L., 1758 (Clitellata): Resurrection

of L. herculeus (Savigny, 1826). *Plos One*, **5**, e15629.

Joppa, L.N., Roberts, D.L. & Pimm, S.L. (2011) How many species of flowering plants are

there? *Proceedings of the Royal Society B*, **278**, 554–559.

Krishnamani, R., Kumar, A. & Harte, J. (2004) Estimating species richness at large spatial

scales using data from small discrete plots. *Ecography*, **27**, 637–642.

Lavelle, P., Decaens, T., Aubert, M., Barot, S., Blouin, M., Bureau, F., Margerie, P., Mora, P.

& Rossi, J.-P. (2006) Soil invertebrates and ecosystem services. *European Journal of Soil*

Biology, **42**, S3–S15.

Lavelle, P., Spain, A., Blouin, M., Brown, G.G., Decaëns, T., Grimaldi, M., Jiménez, J.J.,

McKey, D., Mathieu, J., Velasquez, E. & Zangerlé, A. (2016) Ecosystem engineers in a self-

organized soil: A review of concepts and future research questions. *Soil Science*, **18**, 91–

109.

Linnaeus, C. (1758) *Systema naturae*, 10th edition. Salvi: Holmiae, Stockholm.

Lomolino, M.V. (2004) *Conservation biogeography. Frontiers of biogeography: new*

directions in the geography of nature (ed. by M.V. Lomolino) and L.R. Heaney), pp. 293–

296. Sunderland, MA.

Maggia, M.-E., Decaëns, T., Lapied, E., Dupont, L., Roy, V., Schimann, H., Orivel, J.,

Murienne, J., Baraloto, C., Cottenie, K. & Steinke, D. (2021) At each site its diversity: DNA

barcoding reveals remarkable earthworm diversity in neotropical rainforests of French

Guiana. *Applied Soil Ecology*, **164**, 103932.

Marchán, D.F., Decaëns, T., Domínguez, J. & Novo, M. (2022) Perspectives in earthworm

phylogeny: recent advances in Lumbricoidea and standing questions. *Diversity*, **19**, 22.

Marchán, D.F., Novo, M., Domínguez, J., da Silva, A., Martínez, A. & Decaëns, T. (2023)

The best is yet to come: six new species within a large-bodied earthworm genus

(Scherotheca, Lumbricidae) in a densely sampled region (Southwestern France). *Zoological*

Journal of the Linnean Society, **20**, 1–19.

Marcon, E., Mirabel, A., Molino, J.-F. & Sabatier, D. (2024) Estimation of the number of tree

species in French Guiana by extrapolation of permanent plots richness. *Journal of Tropical*

Ecology, **in press**.

Mathieu, J. & Davies, T.J. (2014) Glaciation as an historical filter of below-ground

biodiversity. *Journal of Biogeography*, **41**, 1204–1214.

May, R.M. (1992) How many species inhabit the earth? *Scientific American*, **267**, 18–24.

Misirlioğlu, M., Reynolds, J.W., Stojanović, M., Trakić, T., Sekulić, J., James, S.W., Csuzdi,

C., Decaëns, T., Lapied, E., Phillip, H., Cameron, E.K. & Brown, G.G. (2023) Earthworms

(Clitellata, Megadrili) of the world: an updated checklist of valid species and families, with

notes on their distribution. *Zootaxa*, **5255**, 417–438.

Padial, P.J., Miralles, A., de la Riva, I. & Vences, M. (2010) The integrative future of

taxonomy. *Frontiers in Zoology*, **7**, 16.

Pennisi, E. (2019) DNA barcodes jump-start search for new species. *Science*, **364**, 920–921.

Phillips, H.R.P., Bach, E.M., Bartz, M.L.C., Bennett, J.M., Beugnon, R., Briones, M.J.I.,

Brown, G.G., Ferlian, O., Gongalsky, K.B., Guerra, C.A., Koenig-Ries, B., Krebs, J.J.,

Orgiazzi, A., Ramirez, K.S., Russell, D.J., Schwarz, B., Wall, D.H., Brose, U., Decaens, T.,

Lavelle, P., Loreau, M., Mathieu, J., Mulder, C., van der Putten, W.H., Rillig, M.C., Thakur,

M.P., de Vries, F.T., Wardle, D.A., Ammer, C., Ammer, S., Arai, M., Ayuke, F.O., Baker,

G.H., Baretta, D., Barkusky, D., Beausejour, R., Bedano, J.C., Birkhofer, K., Blanchart, E.,

Blossey, B., Bolger, T., Bradley, R.L., Brossard, M., Burtis, J.C., Capowiez, Y., Cavagnaro,

T.R., Choi, A., Clause, J., Cluzeau, D., Coors, A., Crotty, F.V., Crumsey, J.M., Davalos, A.,

Diaz Cosin, D.J., Dobson, A.M., Dominguez, A., Esteban Duhour, A., van Eekeren, N.,

Emmerling, C., Falco, L.B., Fernandez, R., Fonte, S.J., Fragoso, C., Franco, A.L.C.,

Fusilero, A., Geraskina, A.P., Gholami, S., Gonzalez, G., Gundale, M.J., Gutierrez Lopez,

M., Hackenberger, B.K., Hackenberger, D.K., Hernandez, L.M., Hirth, J.R., Hishi, T.,

Holdsworth, A.R., Holmstrup, M., Hopfensperger, K.N., Huerta Lwanga, E., Huhta, V.,

Hurisso, T.T., Iannone, B.V., Iordache, M., Irmler, U., Ivask, M., Jesus, J.B., Johnson-

Maynard, J.L., Joschko, M., Kaneko, N., Kanianska, R., Keith, A.M., Kernecker, M.L., Kone,

A.W., Kooch, Y., Kukkonen, S.T., Lalthanzara, H., Lammel, D.R., Lebedev, I.M., Le Cadre,

E., Lincoln, N.K., Lopez-Hernandez, D., Loss, S.R., Marichal, R., Matula, R., Minamiya, Y.,

Moos, J.H., Moreno, G., Moron-Rios, A., Motohiro, H., Muys, B., Neirynck, J., Norgrove, L.,

Novo, M., Nuutinen, V., Nuzzo, V., Rahman, P.M., Pansu, J., Paudel, S., Peres, G., Perez-

Camacho, L., Ponge, J.-F., Prietzel, J., Rapoport, I.B., Rashid, M.I., Rebollo, S., Rodriguez,

M.A., Roth, A.M., Rousseau, G.X., Rozen, A., Sayad, E., van Schaik, L., Scharenbroch, B.,

Schirrmann, M., Schmidt, O., Schroeder, B., Seeber, J., Shashkov, M.P., Singh, J., Smith,

S.M., Steinwandter, M., Szlavecz, K., Antonio Talavera, J., Trigo, D., Tsukamoto, J., Uribe-

Lopez, S., de Valenca, A.W., Virto, I., Wackett, A.A., Warren, M.W., Webster, E.R., Wehr,

N.H., Whalen, J.K., Wironen, M.B., Wolters, V., Wu, P., Zenkova, I.V., Zhang, W., Cameron,

E.K. & Eisenhauer, N. (2021) Global data on earthworm abundance, biomass, diversity and

corresponding environmental properties. *Scientific Data*, **8**, 136.

Phillips, H.R.P., Guerra, C.A., Bartz, M.L.C., Briones, M.J.I., Brown, G., Crowther, T.W.,

Ferlian, O., Gongalsky, K.B., van den Hoogen, J., Krebs, J., Orgiazzi, A., Routh, D.,

Schwarz, B., Bach, E.M., Bennett, J., Brose, U., Decaens, T., Koenig-Ries, B., Loreau, M.,

Mathieu, J., Mulder, C., van der Putten, W.H., Ramirez, K.S., Rillig, M.C., Russell, D.,

Rutgers, M., Thakur, M.P., de Vries, F.T., Wall, D.H., Wardle, D.A., Arai, M., Ayuke, F.O.,

Baker, G.H., Beausejour, R., Bedano, J.C., Birkhofer, K., Blanchart, E., Blossey, B., Bolger,

T., Bradley, R.L., Callaham, M.A., Capowiez, Y., Caulfield, M.E., Choi, A., Crotty, F.V.,

Davalos, A., Diaz Cosin, D.J., Dominguez, A., Duhour, A.E., van Eekeren, N., Emmerling,

C., Falco, L.B., Fernandez, R., Fonte, S.J., Fragoso, C., Franco, A.L.C., Fugere, M.,

Fusilero, A.T., Gholami, S., Gundale, M.J., Gutierrez Lopez, M., Hackenberger, D.K.,

Hernandez, L.M., Hishi, T., Holdsworth, A.R., Holmstrup, M., Hopfensperger, K.N., Lwanga,

E.H., Huhta, V., Hurisso, T.T., Iannone, B.V., Iordache, M., Joschko, M., Kaneko, N.,

Kanianska, R., Keith, A.M., Kelly, C.A., Kernecker, M.L., Klaminder, J., Kone, A.W., Kooch,

Y., Kukkonen, S.T., Lalthanzara, H., Lammel, D.R., Lebedev, I.M., Li, Y., Jesus Lidon, J.B.,

Lincoln, N.K., Loss, S.R., Marichal, R., Matula, R., Moos, J.H., Moreno, G., Moron-Rios, A.,

Muys, B., Neirynck, J., Norgrove, L., Novo, M., Nuutinen, V., Nuzzo, V., Rahman, M.P.,

Pansu, J., Paudel, S., Peres, G., Perez-Camacho, L., Pineiro, R., Ponge, J.-F., Rashid, M.I.,

Rebollo, S., Rodeiro-Iglesias, J., Rodriguez, M.A., Roth, A.M., Rousseau, G.X., Rozen, A.,

Sayad, E., van Schaik, L., Scharenbroch, B.C., Schirrmann, M., Schmidt, O., Schroeder, B.,

Seeber, J., Shashkov, M.P., Singh, J., Smith, S.M., Steinwandter, M., Talavera, J.A., Trigo,

D., Tsukamoto, J., de Valenca, A.W., Vanek, S.J., Virto, I., Wackett, A.A., Warren, M.W.,

Wehr, N.H., Whalen, J.K., Wironen, M.B., Wolters, V., Zenkova, I.V., Zhang, W., Cameron,

E.K. & Eisenhauer, N. (2019) Global distribution of earthworm diversity. *Science*, **366**, 480–

485.

Pimm, S.L. (2001) *The World according to Pimm: A Scientist Audits the Earth*, McGraw-Hill.,

New York, USA.

Pimm, S.L., Jenkins, C.N., Joppa, L.N., Roberts, D.L. & Russell, G.J. (2010) How Many

Endangered Species Remain to be Discovered in Brazil? *Natureza & Conservação*, **08**, 71–

77.

R Core Team (2024) R: a language and environment for statistical computing. Version 4.4.1.

Reynolds, J.W. (1994) Earthworms of the world. *Global biodiversity*, **4**, 11–16.

Rougerie, R., Decaens, T., Deharveng, L., Porco, D., James, S.W., Chang, C.-H., Richard,

B., Potapov, M., Suhardjono, Y. & Hebert, P.D.N. (2009) DNA barcodes for soil animal

taxonomy. *Pesquisa Agropecuaria Brasileira*, **44**, 789–802.

Stork, N. (1993) How many species are there? *Biodiversity and Conservation*, **2**, 215–232.

Taberlet, P., Coissac, E., Hajibabaei, M. & Rieseberg, L.H. (2012) Environmental DNA.

Molecular Ecology, **21**, 1789–1793.

Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., van der Putten, W.H. & Wall, D.H.

(2004) Ecological linkages between aboveground and belowground biota. *Science*, **304**,

1629–1633.

Wilson, E.O. (2004) Taxonomy as a fundamental discipline. *Philosophical Transactions of

the Royal Society B - Biological Sciences*, **359**, 739–739.

Data Accessibility Statement

538   All the data and scripts for R used in this work are available on GitHub:

539   https://github.com/EricMarcon/A_Can_Of_Worms-Appendix.

Figure legends

Figure 1. Fluctuations over time, for each 5-year period from 1822 to 2017, in the numbers of described earthworm species and taxonomists (A), and in the number of species described per taxonomist (B). The loess models fitted to the data are shown in colour, with the grey area corresponding to the 95% confidence interval.

Figure 2. Results of Bayesian estimation of the global number of earthworm species according to the original Joppa model (A) and the no-progress model assuming the taxonomists' efficiency constant (B). The diagram shows the probability density of the estimated ST value. The red line represents the median and the dashed lines are the limits of the highest posterior density interval.

Figure 3. Incidence-based rarefaction and extrapolation curves showing how earthworm species richness increases as a function of sampling effort (number of 100 x 100 km2 windows) in the three regions studied. Solid lines represent rarefaction curves (interpolated values); dashed lines represent extrapolation curves; shaded areas represent 95% confidence intervals obtained with a bootstrap of 200 replications.

Figure 4. Analysis of earthworm diversity in the three study regions: A) cumulative number of observed species, B) estimated number of species (Chao 2 index) and C) taxonomic deficit (i.e. the proportion of estimated diversity that has not been observed). In A, the grey hatched part corresponds to species from French Guiana that have already been named, while the non-hatched part of the bar corresponds to species that have been discovered but not yet described; in B, the error bars represent the 95% confidence interval obtained using a bootstrap with 200 replications; in C, the hatched part corresponds to the inflation of the taxonomic deficit in French Guiana as a result of taking into account only species that have been already named, the non-hatched part of the bar corresponding to the deficit calculated by also taking into account species that have been discovered but not yet described.

567   *Figure 1.*

572   *Figure 2.*

575 **_Figure 3._**

578 **Figure 4.**