A can of worms: estimating the global number of earthworm species

Thibaud Decaëns[1*], George G. Brown[2], Erin K. Cameron[3], Csaba Csuzdi[4], Nico Eisenhauer[5,6], Sylvain Gérard[7], Arnaud Goulpeau[8], Mickaël Hedde[9], Samuel W. James[10], Emmanuel Lapied[11], Marie-Eugénie Maggia[12], Daniel F. Marchán[13], Jérôme Mathieu[14], Helen R. P. Phillips[15], Eric Marcon[16]

[1] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France. ORCID ID: 0000-0003-0915-3882.

[2] EMBRAPA-Floresta, Curitiba, Paraná, Brazil. ORCID ID: 0000-0001-9550-6909

[3] Saint Mary's University, Halifax, NS, Canada. ORCID ID: 0000-0002-3374-9830

[4] Dept. of Zoology, Eszterházy Károly Catholic University, Eger, Hungary. ORCID ID: 0000-0002-0319-7836

[5] German Centre for Integrative Biodiversity Research (iDiv), Leipzig, Germany. ORCID ID: 0000-0002-0371-6720

[6] Leipzig University, Leipzig, Germany

[7] Eco&Sols, INRAE, IRD, CIRAD, l'Institut Agro Montpellier, Univ Montpellier, Montpellier, France. ORCID ID: 0009-0009-6836-6475

[8] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France. ORCID ID: 0009-0007-7489-1722

[9] &Sols, INRAE, IRD, CIRAD, l'Institut Agro Montpellier, Univ Montpellier, Montpellier, France. ORCID ID: 0000-0002-6733-3622

[10] Maharishi International University, Department of Regenerative Agriculture, Fairfield, IA, USA. ORCID ID: 0000-0001-6758-5726

[11] Taxonomia Biodiversity Fund, 7 rue Beccaria, 72012, Paris, France. ORCID ID: 0000-0001-7242-6703

[12] Department of Integrative Biology, University of Guelph, Ontario, Canada. ORCID ID: 0000-0002-0970-6494

[13] Biodiversity, Ecology and Evolution Department, Faculty of Biology, Universidad Complutense de Madrid, Madrid, Spain. ORCID ID: 0000-0001-7971-4329

[14] Sorbonne Université, CNRS, UPEC, INRAE, IRD, Institut d'Ecologie et des Sciences de l'Environnement de Paris, Paris, France. ORCID ID: 0000-0002-9106-6106

[15] University of Helsinki, Helsinki, Finland ORCID: 0000-0002-7435-5934

[16] AgroParisTech, UMR AMAP, CIRAD, CNRS, INRAE, IRD, Univ Montpellier, Montpellier, France. ORCID ID: 0000-0002-5249-321X`

* Corresponding author

Correspondence: thibaud.decaens@cefe.cnrs.fr

ABSTRACT

Estimating the overall species number for a given taxon is a central issue in ecology and conservation biology. This is especially topical for soil organisms, which comprise most known species but whose taxonomy remains largely understudied. Here, we estimated the global number of earthworm species based on the Joppa approach, which models taxonomic effort over time to estimate the total number of known and as yet unknown species in a given taxa. Our Bayesian estimation of the Joppa model suggests a global diversity of the order of 30,000 species, suggesting that the 5,679 earthworm taxa already described only represent around 20% of the actual global species diversity. However, the uncertainty around this estimate is considerable due to severe undersampling and as the model cannot unambiguously decide whether we are describing few species because of a small pool of as yet unknown species, or because of a lack of taxonomic efficiency. Considering the current rate of new species description, we calculate that it would take at least 120 years to describe all the earthworm species existing on Earth, and we discuss thedifferent strategies that should be developed to facilitate and accelerate the discovery and naming of species new to science.

Keywords: Soil biodiversity; global species number estimates; taxonomic deficit; French Guiana; temperate France

Introduction

Estimating the total number of species of a given taxon, whether on a global scale or for a
given region or ecosystem, is a question of great practical and symbolic importance that has long
captivated and challenged scientists (May, 1992; Stork, 1993; Pimm, 2001; Dirzo & Raven,
2003). It represents an indispensable step towards a comprehensive description of the World's
biodiversity and is also necessary for the definition of efficient conservation strategies (Joppa et
al., 2011; Caley et al., 2014). In this context, soil biota represents one of the most diversified and
least studied compartments in terrestrial ecosystems. They are among the most diverse,
comprising up to 59% of currently known living species (Decaëns et al., 2006; Anthony et al.,
2023). They are involved in regulating a large number of ecological processes that are essential
to the provision of key ecosystem services for human populations (Brussaard et al., 1997;
Wardle et al., 2004; Bardgett & van der Putten, 2014). This has been particularly well
documented for emblematic groups of soil ecosystem engineers, such as earthworms (Lavelle et
al., 2006, 2016). Paradoxically, even these well-studied groups still suffer from a glaring lack of
basic studies on their biodiversity patterns (Cameron et al., 2018; Phillips et al., 2019, 2021), and
questions as fundamental as "how many species are there?" still remain largely unanswered
(Decaëns, 2010). This knowledge gap is of particular concern, as it limits our capacity to
characterize the dynamics of soil biodiversity in the face of ongoing global change and to assess
its implications for ecosystem functioning.

Earthworms are an emblematic illustration of this paradox. Some 5,700 earthworm species
have been described to date (Misirlioğlu et al., 2023), but the proportion of the actual diversity of
these organisms that this figure represents is still poorly documented. Estimates vary, depending
on the author, from a minimum of 7,000 to some 20,000 species (Reynolds, 1994; Lavelle et al.,
2006; Blakemore, 2009; Anthony et al., 2023), but most of these figures are based on expert
opinion, and no mathematical formalisation has yet been used to obtain a consolidated prediction
of the number of as-yet undiscovered species. Some studies have also highlighted the existence
of potentially high and as yet largely undescribed regional diversity throughout the tropics
(Phillips et al., 2019; Goulpeau et al., 2024), which could challenge these previously proposed
estimates of global species number.

Various methods can be used to estimate the number of heretofore unknown species within a
given taxon. One popular approach is to use rarefaction and extrapolation curves, combined with
non-parametric estimators, to predict the taxonomic deficit and the number of species at different
spatial scales (Chao, 1984; Colwell et al., 2004; Chao et al., 2005; Chao & Jost, 2012; Cazzolla-
Gatti et al., 2022). However, to be effective, this approach requires a sufficiently dense dataset
on a global scale, which is lacking for many soil organisms such as earthworms. Another method
consists of using Arrhenius' power law to estimate the number of species for large areas on the
basis of a known number of species in a smaller area (Arrhenius, 1921; Krishnamani et al., 2004;
Marcon et al., 2024). The procedure involves estimating a power parameter (z) based on the
negative linear relationship between the log of compositional similarity and the log of
geographical distance between small equivalent areas. It therefore requires a network of plots
located at very different distances, which must be supplemented by a continuous inventory over
an area corresponding to the smallest scale of the power law. Apart from the fact that this type of
data is rare, log transformation is impossible for pairs of plots with a compositional similarity of 0
(i.e. no species in common), which is common in the case of earthworms, particularly in the
tropics (Goulpeau et al., 2024). Finally, Joppa *et al.* (2011) propose to use global species
checklists to model the rate at which taxonomists describe new species as a function of time, and
to estimate the number of species that will have been reached when this rate will drop to zero
due to the exhaustion of the pool of as yet unknown species. This approach has the advantage
of not being dependent on the existence of comprehensive field datasets, but requires the
establishment of reliable species checklists that can be challenged by taxonomic issues.

In this work, we propose a global estimate of the number of earthworm species based on the
method of Joppa *et al.* (2011) and the most recent global checklist of earthworms (Brown et al.,
2023; Misirlioğlu et al., 2023). In parallel, we analysed three datasets obtained from field
inventories for regions equivalent in surface area in tropical and temperate regions (Bouché,
1972; Goulpeau et al., 2024; Gérard et al., 2024). We used these data to quantify the extent of

the taxonomic deficit and estimate the actual number of species for each region. We then used
these regional estimates to discuss the range of results obtained from the Joppa method.

Material and methods

Estimating the number of species from the global checklist

For the Joppa approach, we used the checklist recently published by Misirlioğlu *et al.* (2023)
and Brown *et al.* (2023). This list includes 5,738 species/subspecies (5,406 species and 332
unique subspecies, i.e. not counting nomino-typical subspecies) belonging to 23 families
(including one non-crassiclitellate family: Moniligastridae). We chose to consider subspecies as
taxa in their own right in the analysis, because the choice of whether to consider them as
infraspecific divisions or to elevate them to species rank is a matter of taxonomists' points of
view. For the sake of simplicity, we referred to all the taxa in the list as "species". After deleting
certain duplicates in the table, we obtained a consolidated list of 5,752 taxa names published
between 1758 and 2023, containing for each of them the authors and the year of description. For
each 5-year period, we then counted the number of unique species described and the total
number of authors (i.e. all the co-authors of the species, after standardising the names of authors
with accents and taking account of homonyms, wherever possible).

Joppa *et al.* (2011) expected the number of species (S_i) described in a given interval ($Y_i = i$,
numbered sequentially from 1) to depend on the taxonomic effort, i.e. the number of taxonomists
(T_i) actively describing species during that period, and the unknown number of species remaining
to be described, that is the total number of species (S_T) minus those already discovered ($\sum_{j=0}^{i} S_j$
where S_0 is the number of known species before Y_1). Over time, taxonomists were expected to
become more efficient, i.e. the discovery rate of unknown species per taxonomist was assumed
to increase linearly. The original model was thus

$$S_i \sim T_i(\beta_1 + \beta_2 Y_i)\left(S_T - \sum_{j=0}^{i} S_j\right),$$

where β_1 and β_2 are unknown parameters. The model was log-transformed and estimated
with a normal error of variance σ^2. The parameters to be estimated were thus β_1, β_2, S_T and
$\sigma,$ the total number of species S_T being our main target.

The increase in efficiency accounted for in the Joppa model is not obvious as earthworms are
concerned. Methodological advancements, such as the development of integrative taxonomy,
are likely offset by the increased difficulty of sampling in remote areas, namely tropical
rainforests. Moreover, contemporary scientists rarely focus exclusively on taxonomy, a discipline
whose publications are generally undervalued, leaving them with less time to describe new
species. Consequently, we studied an alternative model with no efficiency progress (hereafter
referred to as the "no-progress" model), where β_2 equals zero.

We chose Bayesian inference to obtain the distribution of the estimates of the parameters
which revealed to be as important as the most probable values. We chose non-informative priors,
uniformly distributed between 0 and numerically possible values β_1, β_2 and σ. According to expert
opinion, the prior of S_T was forced between 6,000, i.e. the number of already discovered species,
and 100,000. The parameter space was explored by a Monte-Carlo Markov Chain with 10^7 steps,
10% considered as burn-in. The analysis began with time period 1822-1827, with a previously
known number of species (S_0) equal to 1 (i.e. *Lumbricus terrestris* Linnaeus, 1758). The last
period is 2018-2022; species published in 2023 and 2024 were excluded from the analysis.

Estimating regional diversity and the taxonomic deficit

Compared to the checklist of angiosperms used by Joppa *et al.* (2011), that of earthworms is
very short: data is scarce. We endeavoured to supplement the global approach with three more
exhaustive regional studies.

For the analyses of regional diversity, we used the data of Goulpeau *et al.* (2022, 2024) in
French Guiana (supplemented by a small number of as yet unpublished data), and, for Southern
and Northern France, those of Bouché (1972) updated for taxonomy by Gérard *et al.* (2024). The
choice of these three regions allows us to consider contrasting situations in terms of earthworm

diversity. French Guiana is considered to be a diversity hotspot, with a high proportion of species with restricted distribution and a large taxonomic deficit (Decaëns et al., 2016, 2024; Maggia et al., 2021; Goulpeau et al., 2024). By comparison, mainland France has a lower *a priori* species diversity, but the South is nevertheless characterised by a high proportion of endemic species, some of which continue to be discovered regularly, whereas the North is characterised by great compositional uniformity (Bouché, 1972; Mathieu & Davies, 2014; Gérard et al., 2023; Marchán et al., 2023).

Three 300 x 400 km^2 zones, each subdivided into twelve 100 x 100 km^2 cells, were delimited in each region. In each 100 x 100 km^2 cell, we counted the species present in order to obtain three occurrence tables comprising 12 rows and as many columns as species present. In French Guiana, we used the operational taxonomic units (OTUs) proposed by Goulpeau *et al.* (2022) as a substitute for species in order to overcome the problem posed by the taxonomic deficit. For each region separately, we plotted incidence-based rarefaction and extrapolation curves in order to represent the accumulation of species richness as a function of the number of 100 x 100 km^2 cells sampled in the study areas. The 95% confidence intervals of the interpolated and extrapolated richness values were obtained by a bootstrap procedure with 200 replications. We also calculated the non-parametric Chao 2 index, which estimates the lower limit of expected asymptotic species richness (Gotelli & Chao, 2013). The taxonomic deficit, i.e. the proportion of undiscovered species, was calculated using the formula:

$$\text{TDef} = ((S_{EST} - S_{OBS}) \times 100 / S_{EST} \, ,$$

with $TDef$ the taxonomic deficit, S_{OBS} the number of species currently known for a given region, and S_{EST} the richness estimated by the Chao 2 index.

For French Guiana, the taxonomic deficit was calculated by first considering S_{OBS} as the set of OTUs delimited by Goulpeau *et al.* (2022), and then by restricting S_{OBS} to the number of species formally described to date (i.e. 55 species taking into account the 18 new descriptions by Decaëns *et al.* (2024). These analyses were performed using the *'iNEXT'* package for R (Hsieh et al., 2019; R Core Team, 2024).

Figure 1. Fluctuations over time, for each 5-year period from 1822 to 2017, in the numbers of described earthworm species and taxonomists (A), and in the number of species described per taxonomist (B). The loess models fitted to the data are shown in colour, with the grey area corresponding to the 95% confidence interval.

202   **Figure 2.** Results of Bayesian estimation of the global number of earthworm
203   species according to the original Joppa model (A) and the no-progress model
204   assuming the taxonomists' efficiency constant (B). The diagram shows the
205   probability density of the estimated ST value. The red line represents the median
206   and the dashed lines are the limits of the highest posterior density interval.

Results

During the period from 1822 to 2022, the number of species described per 5-year period showed two peaks: the first from the end of the 19th to the middle of the 20th century, the other from the 1970s to the present day (Fig. 1A). Over the same period, the total number of authors of original species descriptions has increased exponentially (i.e. roughly linearly on a logarithmic scale; Fig. 1A). The ratio of numbers of species described per taxonomist thus shows a very marked peak in the first half of the 20th century, a period during which it generally exceeds 15 species described per taxonomist per time period, whereas everywhere else it is generally below 10 species per taxonomist (Fig. 1B).

Figure 3. Incidence-based rarefaction and extrapolation curves showing how earthworm species richness increases as a function of sampling effort (number of 100 x 100 km2 windows) in the three regions studied. Solid lines represent rarefaction curves (interpolated values); dashed lines represent extrapolation curves; shaded areas represent 95% confidence intervals obtained with a bootstrap of 200 replications.

The Joppa model yields the median posterior ST equal to 6,814 species, with a 95% higher posterior density interval (HPDI) between 6,000 and 11,148. The no-progress model leads to quite different estimations: 26,283 species (HDPI between 6,003 and 87,425). Posterior probabilities are shown in Fig. 2.

The rarefaction curves obtained for the three study regions show a decrease in beta diversity from tropical to temperate study regions (Fig. 3). In French Guiana, the slope of the curve is steep, showing that the asymptote is far from being reached, whereas it has been or is about to be reached in the two study regions in temperate France. The diversity metrics logically follow the same trend: the cumulative number of species observed is 305 in French Guiana, 66 in Southern France, and 31 in Northern France, whereas the Chao 2 index predicts a diversity of 1,680 species in French Guiana, 99 in Southern France, and 36 in Northern France (Fig. 4A, 4B). Consequently, the taxonomic deficit is of the order of 80% in French Guiana, 33% in the south of France, and 16% in the north of France (Fig. 4C). In French Guiana, this 80% taxonomic deficit

was obtained by considering as observed richness all of the OTUs mentioned in Goulpeau et al.
(2022), most of which are new to science and have not yet been named. If we consider only the
55 named species actually known from French Guiana (Fig. 4A), the estimated taxonomic deficit
rises to 97% (Fig. 4C).

Figure 4. Analysis of earthworm diversity in the three study regions: A) cumulative
number of observed species, B) estimated number of species (Chao 2 index) and
C) taxonomic deficit (i.e. the proportion of estimated diversity that has not been
observed). In A, the grey hatched part corresponds to species from French Guiana
that have already been named, while the non-hatched part of the bar corresponds
to species that have been discovered but not yet described; in B, the error bars
represent the 95% confidence interval obtained using a bootstrap with 200
replications; in C, the hatched part corresponds to the inflation of the taxonomic
deficit in French Guiana as a result of taking into account only species that have
been already named, the non-hatched part of the bar corresponding to the deficit
calculated by also taking into account species that have been discovered but not
yet described.

Discussion

Earthworm taxonomy experienced a golden age from the end of the 19th century to the
middle of the 20th century, followed by a revival from the end of the 1970s. The first of these two
periods can be credited to the great explorations that enabled zoologists of the time to explore
the world and describe its biological diversity (Csuzdi & Szlávecz, 2016). It was also made
possible by the exceptional productivity of the few taxonomists of the time, who were able to
describe dozens of species in a short time. By comparison, the contemporary period is
characterised by a larger community of taxonomists, but with comparatively lower output.
Overall, we do not observe the monotonic decline in the number of species described per
taxonomist as described for plants and other animal taxa (Pimm et al., 2010; Joppa et al., 2011;
Fisher et al., 2018), which is assumed to reflect a depletion of the pool of new species to be

discovered. Rather, with the exception of the notable peak in the first half of the 20th century, the
rate of description of new earthworm species remained modest and constant over time, at
around 5 species per taxonomist per 5-year period. Even in the early 2000s, we did not observe
any clear effect from the development of molecular tools such as DNA barcoding and
metabarcoding (sensu Hebert et al., 2003; Taberlet et al., 2012). This may seem surprising,
given that their usefulness in the field of earthworm taxonomy has been widely acknowledged
(Rougerie et al., 2009; Decaëns et al., 2013; Marchán et al., 2022), and their advent has been
touted as the possible catalyst of a new golden age of taxonomy (Hebert & Gregory, 2005;
Hajibabaei, 2012; Pennisi, 2019). Over the last few decades, the increase in the number of new
earthworm species described per time period seems to be simply correlated with the number of
active taxonomists.

Our estimates of the Joppa and the no-progress model tell us two different stories. Assuming
a constant progress of the efficiency of taxonomists, the global number of species would be
below 7,000, i.e. around 1,000 species remaining to be described. On the other hand, assuming
constant efficiency, the global number of species would rather be around 30,000, i.e. the 5,679
species of our checklist are only the beginning of the story.

In both cases, the confidence intervals around our median estimates are considerable,
motivating us not to provide a single significant figure in our estimates. This can be explained by
two main causes of uncertainty. First, we lack data: more publications of new species would have
allowed smaller, i.e. more, time intervals, thus more statistical power. Also, the Joppa model
does not allow to completely disentangle the effects of the efficiency and the number of species:
in the no-progress model, the estimates are highly correlated (Pearson correlation of the
posteriors: -0,73), and the correlation is -0,55 between β_2 and S_T in the original model. This
issue could not be detected by Joppa et al. (2011) who relied on a frequentist estimation of the
most likely number of species.

The analysis of regional diversity in French Guiana points in the direction of a high number of
species: we estimate that the taxonomic deficit there is of the order of 97%, if only currently
named species are considered in the calculation. An unknown thousand species may well all be
there: a small area (actually around 1%) of Amazonia may thus be enough to find the species
missing from the Joppa model. If we consider that around half of the earthworm species currently
known are tropical (i.e. around 2,800 species), the generalisation of such a taxonomic deficit
would lead us to estimate that more than 90,000 species could exist throughout the tropics. Of
course, such a generalisation is highly debatable, but it does provide an argument for
considering that the global number of unknown earthworm species may not be only a few
thousands.

As expected, we calculate a lower taxonomic deficit in temperate regions, which have been
better studied and host a more modest diversity. However, our results indicate that this deficit
can still be significant—ranging from 15 to 30% in mainland France—suggesting that even in
temperate regions, a substantial number of species remains undiscovered. The South of France,
known for its high diversity and richness in endemic earthworms, exemplifies this, as evidenced
by the recent descriptions of several new species, including some of large body size (Marchán et
al., 2023). The persistence of a significant taxonomic deficit in Northern France is more
unexpected. It may reflect a gap in our understanding of species distribution in the region (i.e. the
Wallacean shortfall according to Lomolino, 2004), or the persistence of taxonomic confusion that
can conceal the presence of certain species. A striking example of this is the discovery of the
existence of two cryptic species that were mistakenly considered synonymous under the name L.
terrestris (James et al., 2010).

These considerations allow us to propose a conservative estimate of the global number of
earthworm species of the order of 30,000 species. This is above the upper range of commonly
accepted predictions, i.e. from 10,000 to 20,000 species (Reynolds, 1994; Decaëns et al., 2006;
Blakemore, 2009; Anthony et al., 2023). Our expert opinion is that there is still a large pool of
undiscovered species, but that we lack the efficiency to discover them or describe them once
they have been discovered. First, a low rate of new species description per taxonomist,
compared with those active at the beginning of the 20th century, is expected due the fact that
taxonomy is no longer considered a 'competitive' discipline in modern science (Wilson, 2004;

Agnarsson & Kuntner, 2007). The pernicious effect of this situation is that contemporary authors
are rarely pure taxonomists, as they are encouraged by the academic system to devote most of
their research time to more 'valuable' areas of research, such as evolution or ecology. Added to
this is the fact that taxonomists generally lack the time and financial resources for field sampling,
particularly in the most diverse and yet least well-studied tropical regions (Guerra et al., 2020).
The field scientists among us all have in mind regions that they know contain a diversity of as yet
unknown species, but which they do not have the time or the resources to explore. Similarly,
taxonomists have a multitude of species in their cupboards waiting to be described, but which
they do not have the time to process.

Overall, our results suggest that at best we have only described one fifth of the global number
of earthworm species. If we consider the current community of taxonomists (an average of 45
active taxonomists per 5-year period over the last 20 years), and the number of species they
have managed to describe (an average of 4.5 species per taxonomist per time step), we
calculate that it would take at least 120 years to describe all the species existing on Earth.
Obviously, such an objective is certainly unattainable within a reasonable timeframe, but we can
nonetheless think about the strategies that could be developed to reduce the magnitude of the
taxonomic deficit. This will require maintaining a major taxonomic effort, through the training of
young taxonomists, but also by raising the profile of taxonomic work in the scientific corpus.
Agnarsner & Kuntner (2007) have proposed that original descriptions, and their underlying
taxonomic hypotheses, should be formally cited in articles that refer to them – which is not
systematically the case at present. This would allow the impact metrics of taxonomic journals to
improve while more accurately reflecting their real scientific impact, thus encouraging scientists
to invest more in taxonomy. We will also need to improve taxonomic efficiency, first by increasing
our ability to discover as yet unknown species where they occur. This means stepping up
sampling efforts, particularly in the tropics where most of this undiscovered diversity is to be
found. Our results confirm the existence of a regional peak in diversity at tropical latitudes,
essentially explained by a strong beta component (Phillips et al., 2019; Goulpeau et al., 2024),
while emphasising that it mirrors a considerable taxonomic deficit. Finally, once sampling
coverage has been improved, new taxonomic approaches based on turbotaxonomy (i.e. the
rapid description of species new to science allowed by automated procedures and the use of
molecular tools; Butcher et al., 2012; Fernandez-Triana, 2022) and integrative taxonomy (i.e. the
integrated use of multiple, independent lines of evidence—e.g. morphology, genetics, ecology,
and behavior—for species delimitation and description; Dayrat, 2005) will be necessary to
improve our ability to describe new species at a consistent rate (Padial et al., 2010; Fernandez-
Triana, 2022).

In conclusion, we propose an updated estimate of the global number of earthworm species,
as well as arguments for considering earthworms as a potentially hyperdiverse group of
organisms, which has hitherto not been considered as such due to a strong taxonomic
knowledge deficit (i.e the Linnean shortfall as defined in Lomolino, 2004). This is particularly
remarkable when considering that earthworms are among the most widely publicised soil macro-
organisms, due to their acknowledged importance in soil functioning. It is likely that this situation
could also apply to other less well-studied soil taxa, leading to a considerable reassessment of
the contribution of these organisms to global biodiversity.

Acknowledgments and funding

The French Guiana dataset was acquired as part of different projects: DIADEMA (Dissecting
Amazonian Diversity by Improving a Multiple Taxonomic Group Approach), DIAMOND
(Dissecting and Monitoring Amazonian Diversity) and Wormbank (DNA barcoding earthworms in
biodiversity hot spots of French Guiana) projects funded by "Investissement d'Avenir" grants
managed by the Agence Nationale de la Recherche (CEBA: ANR-10-LABX-25-01; TULIP: ANR-
10-LABX-41); "Our Planet Reviewed" French Guiana 2015 expedition (Touroult et al., 2018)
organised by the Muséum national d'Histoire naturelle (MNHN, Paris) and Pro-Natura
international in collaboration with the Amazonian Park of French Guiana, and financed by the
European Fund for Regional Development (FEDER), the Regional Council of French Guiana, the
General Council of French Guiana, the Direction de l'Environnement, de l'Aménagement et du

Logement and by the Ministère de l'Éducation nationale, de l'Enseignement supérieur et de la
Recherche. Additional support was received from the Réserve naturelle des Nouragues
(Nouragues grants 2010 and 2011), the Réserve naturelle de la Trinité and the Parc Amazonien
de Guyane.

The dataset from temperate France was obtained through a project called GloWorms, mostly
funded by the 'AgroEcoSystems' Scientific Department of INRAE (National Institute for Research
in Agriculture, Environment and Food, Paris, France). Samplings in Normandy were funded by
the INPN (National Inventory of Natural Heritage, Paris, France). Sylvain Gérard's PhD thesis
was funded by a thesis grant from the Ecole Normale Supérieure (Paris, France). Daniel F.
Marchán's contribution was funded by the Make Our Planet Great Again programme (Campus
France, Paris), via a postdoctoral grant.

JM and GGB acknowledge the FaunaServices group jointly funded by the synthesis center
CESAB   of   the   French   Foundation   for   Research   on   Biodiversity   (FRB;
www.fondationbiodiversite.fr)", CEBA, Sinbiose, Fapesp and CNPq (Brazil). NE gratefully
acknowledges the support of iDiv, which is funded by the German Research Foundation (DFG –
FZT 118, 202548816), as well as by the DFG (Ei 862/29-1, Ei 862/31-1). HRPP was funded by
the European Union's Horizon 2020 research and innovation program under the Marie
Skłodowska-Curie grant agreement No. 101033214 (GloSoilBio). GGB acknowledges project
support by the Brazilian National Council for Scientific and Technological Development (CNPq
Nos. 312824/2022-0, 441930/2020-4 and 404191/2019-3).

Conflict of interest disclosure

395 The authors declare that they comply with the PCI rule of having no financial conflicts of
396 interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

398 Data and scripts are available online on GitHub:
399 https://github.com/EricMarcon/A_Can_Of_Worms-Appendix.

References

Agnarsson I, Kuntner M (2007) Taxonomy in a changing world: Seeking solutions for a science in
crisis. *Systematic Biology*, **56**, 531–539. https://doi.org/10.1080/10635150701424546

Anthony MA, Bender SF, van der Heijden MGA (2023) Enumerating soil biodiversity.
Proceedings of the National Academy of Sciences of the United States of America, **120**,
e2304663120.

Arrhenius   O   (1921)   Species   and   area.   *Journal   of   Ecology*,   **9**,   95–99.
https://doi.org/10.2307/2255763

Bardgett RD, van der Putten WH (2014) Belowground biodiversity and ecosystem functioning.
Nature, **515**, 505–511. https://doi.org/10.1038/nature13855

Blakemore RJ (2009) Chapter 14. Cosmopolitan earthworms – a global and historical
perspective. In: *Annelids as model systems in the biological sciences* (ed Shain DH), pp.
257–283. John Wiley & Sons, Inc., New York.

Bouché MB (1972) *Lombriciens de France*. INRA, Paris.

Brown GG, James SW, Csuzdi C, Lapied E, Decaëns T, Reynolds JW, Misirlioğlu M, Stovanic M,
Trakić T, Sekulić J, Phillips HRP, Cameron EK (2023) A checklist of megadrile earthworm
(Annelida: Clitellata)   species   and   subspecies   of   the   world.   *Zedono*,
https://doi.org/10.5281/zenodo.8348860. https://doi.org/10.5281/zenodo.8348860

Brussaard L, Behan-Pelletier VM, Bignell DE, Brown VK, Didden W, Folgarait P, Fragoso C,
Freckman DW, Gupta V, Hattori T, Hawksworth DL, Klopatek C, Lavelle P, Malloch DW,

Rusek J, Soderstrom B, Tiedje JM, Virginia RA (1997) Biodiversity and ecosystem functioning in soil. *AMBIO*, **26**, 563–570.

Butcher BA, Smith MA, Sharkey MJ, Quicke DLJ (2012) A turbo-taxonomic study of Thai Aleiodes (Aleiodes) and Aleiodes (Arcaleiodes) (Hymenoptera: Braconidae: Rogadinae) based largely on COI barcoded specimens, with rapid descriptions of 179 new species. *Zootaxa*, 1–232.

Caley MJ, Fisher R, Mengersen K (2014) Global species richness estimates have not converged. *Trends in Ecology & Evolution*, **29**, 187–188.

Cameron EK, Martins IS, Lavelle P, Mathieu J, Tedersoo L, Gottschall F, Guerra CA, Hines J, Patoine G, Siebert J, Winter M, Cesarz S, Delgado-Baquerizo M, Ferlian O, Fierer N, Kreft H, Lovejoy TE, Montanarella L, Orgiazzi A, Pereira HM, Phillips HRP, Settele J, Wall DH, Eisenhauer N (2018) Global gaps in soil biodiversity data. *Nature Ecology & Evolution*, **2**, 1042–1043. https://doi.org/10.1038/s41559-018-0573-8

Cazzolla-Gatti R, Reich PB, Gamarra JGP, Crowther T, Hui C, Morera A, Bastin J-F, de-Miguel S, Nabuurs G-J, Svenning J-C, Serra-Diaz JM, Merow C, Enquist B, Kamenetsky M, Lee J, Zhu J, Fang J, Jacobs DF, Pijanowski B, Banerjee A, Giaquinto RA, Alberti G, Almeyda Zambrano AM, Alvarez-Davila E, Araujo-Murakami A, Avitabile V, Aymard GA, Balazy R, Baraloto C, Barroso JG, Bastian ML, Birnbaum P, Bitariho R, Bogaert J, Bongers F, Bouriaud O, Brancalion PHS, Brearley FQ, Broadbent EN, Bussotti F, Castro da Silva W, Gomes Cesar R, Cesljar G, Moscoso VC, Chen HYH, Cienciala E, Clark CJ, Coomes DA, Dayanandan S, Decuyper M, Dee LE, Del Aguila Pasquel J, Derroire G, Djuikouo MNK, Tran VD, Dolezal J, Dordevic ID, Engel J, Fayle TM, Feldpausch TR, Fridman JK, Harris DJ, Hemp A, Hengeveld G, Herault B, Herold M, Ibanez T, Jagodzinski AM, Jaroszewicz B, Jeffery KJ, Johannsen VK, Jucker T, Kangur A, Karminov VN, Kartawinata K, Kennard DK, Kepfer-Rojas S, Keppel G, Khan ML, Khare PK, Kileen TJ, Kim HS, Korjus H, Kumar A, Kumar A, Laarmann D, Labriere N, Lang M, Lewis SL, Lukina N, Maitner BS, Malhi Y, Marshall AR, Martynenko OV, Mendoza ALM, Ontikov PV, Ortiz-Malavasi E, Camacho NCP, Paquette A, Park M, Parthasarathy N, Peri PL, Petronelli P, Pfautsch S, Phillips OL, Picard N, Piotto D, Poorter L, Poulsen JR, Pretzsch H, Ramirez-Angulo H, Correa ZR, Rodeghiero M, Gonzales RDPR, Rolim SG, Rovero F, Rutishauser E, Saikia P, Salas-Eljatib C, Schepaschenko D, Scherer-Lorenzen M, Seben V, Silveira M, Slik F, Sonke B, Souza AF, Sterenczak KJ, Svoboda M, Taedoumg H, Tchebakova N, Terborgh J, Tikhonova E, Torres-Lezama A, van der Plas F, Vasquez R, Viana H, Vibrans AC, Vilanova E, Vos VA, Wang H-F, Westerlund B, White LJT, Wiser SK, Zawila-Niedzwiecki T, Zemagho L, Zhu Z-X, Zo-Bi IC, Liang J (2022) The number of tree species on Earth. *Proceedings of the National Academy of Sciences of the United States of America*, **119**, e2115329119. https://doi.org/10.1073/pnas.2115329119

Chao A (1984) Nonparametric estimation of the number of classes in a population. *Scandinavian Journal of Statistics*, **11**, 265–270.

Chao A, Chazdon RL, Colwell RK, Shen TJ (2005) A new statistical approach for assessing similarity of species composition with incidence and abundance data. *Ecology Letters*, **8**, 148–159. https://doi.org/10.1111/j.1461-0248.2004.00707.x

Chao A, Jost L (2012) Diversity measures. In: *Encyclopedia of Theoretical Ecology* (eds Hastings A, Gross L), pp. 203–207. University of California Press, Berkeley, USA.

Colwell RK, Mao CX, Chang J (2004) Interpolating, extrapolating, and comparing incidence-based species accumulation curves. *Ecology*, **85**, 2717–2727. https://doi.org/10.1890/03-0557

Csuzdi C, Szlávecz K (2016) Earthworm (Clitellata: Megadrili) taxonomy in the last 200 years: A homage to András Zicsi (1928–2015). *Opuscula Zoologica (Budapest)*, **47**, 1–7.

Dayrat (2005) Towards integrative taxonomy. *Biological Journal of the Linnean Society*, **85**, 407–417. https://doi.org/10.1111/j.1095-8312.2005.00503.x

Decaëns T (2010) Macroecological patterns in soil communities. *Global Ecology and Biogeography*, **19**, 287–302.

Decaëns T, Bartz MLC, Goulpeau A, Marchan DF, Maggia M-E, Lapied E, Feijoo AM, James SW (2024) Earthworms (Oligochaeta: Clitellata) of the Mitaraka range (French Guiana): commented checklist with description of one genus and eighteen species new to science. *Zoosystema*, **46**, 195–244.

Decaëns T, Jiménez JJ, Gioia C, Measey J, Lavelle P (2006) The values of soil animals for conservation biology. *European Journal of Soil Biology*, **42**, S23–S38.

Decaëns T, Porco D, James SW, Brown GG, Chassany V, Dubs F, Dupont L, Lapied E, Rougerie R, Rossi J-P, Roy V (2016) DNA barcoding reveals diversity patterns of earthworm communities in remote tropical forests of French Guiana. *Soil Biology & Biochemistry*, **92**, 171–183. https://doi.org/10.1016/j.soilbio.2015.10.009

Decaëns T, Porco D, Rougerie R, Brown GG, James SW (2013) Potential of DNA barcoding for earthworm research in taxonomy and ecology. *Applied Soil Ecology*, **65**, 35–42. https://doi.org/10.1016/j.apsoil.2013.01.001

Dirzo R, Raven P (2003) Global state of biodiversity and loss. *Annu Rev Environ Resour*, **28**, 137–167.

Fernandez-Triana JL (2022) Turbo taxonomy approaches: lessons from the past and recommendations for the future based on the experience with Braconidae (Hymenoptera) parasitoid wasps. *Zootaxa*, **1087**. https://doi.org/10.3897/zookeys.1087.76720

Fisher MA, Vinson JE, Gittleman JL, Drake JM (2018) The description and number of undiscovered mammal species. *Ecology and Evolution*, **8**, 3628–3635.

Gérard S, Decaëns T, Marchán DF, Beauchesne M, Berlioz L, Capowiez Y, Clause J, Decaëns L, Dellavedova R, Duhaut C-B, Garnier-Fière C, Goulpeau A, Goussopoulos J, Iannelli M, Marsden C, Navarro A, Orrière S, Revertégat C, Vannier A, Versavel C, Hedde M (2024) #Vers2022: a dataset of species composition and individual mass of earthworm communities resampled 50 years after Bouché's historical work. *Ecology*, **in press**.

Gérard S, Marchán DF, Martinez Navarro A, Hedde M, Decaëns T (2023) Resampling Bouché's historical localities reveals three new species and helps identifying a new genus of earthworms (Oligochaeta, Hormogastridae and Lumbricidae) in Southeastern France. *Zoosystema*, **45**, 749–768.

Gotelli N, Chao A (2013) Measuring and Estimating Species Richness, Species Diversity, and Biotic Similarity from Sampling Data. In: *Encyclopedia of Biodiversity* (ed Levin SA), pp. 195–211. Academic Press, Waltham, MA.

Goulpeau A, Hedde M, Maggia M-E, Gérard S, Lapied E, Decaëns T (2024) A dataset on earthworm communities in French Guiana. *Soil Organisms*, **92**, 79–90.

Goulpeau A, Penel B, Maggia M-E, Marchán DF, Steinke D, Hedde M, Decaëns T (2022) OTU delimitation with earthworm DNA barcodes: a comparison of methods. *Diversity*, **14**, 866.

Guerra CA, Heintz-Buschart A, Sikorski J, Chatzinotas A, Guerrero-Ramirez N, Cesarz S, Beaumelle L, Rillig MC, Maestre FT, Delgado-Baquerizo M, Buscot F, Overmann J, Patoine G, Phillips HRP, Winter M, Wubet T, Kuesel K, Bardgett RD, Cameron EK, Cowan D, Grebenc T, Marin C, Orgiazzi A, Singh BK, Wall DH, Eisenhauer N (2020) Blind spots in global soil biodiversity and ecosystem function research. *Nature Communications*, **11**, 3870. https://doi.org/10.1038/s41467-020-17688-2

Hajibabaei M (2012) The golden age of DNA metasystematics. *Trends in Genetics*, **28**, 535–537. https://doi.org/10.1016/j.tig.2012.08.001

Hebert PDN, Cywinska A, Ball SL, DeWaard JR (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society B-Biological Sciences*, **270**, 313–321. https://doi.org/10.1098/rspb.2002.2218

Hebert PDN, Gregory TR (2005) The promise of DNA barcoding for taxonomy. *Syestematic Biology*, **54**, 852–859. https://doi.org/10.1080/10635150500354886

Hsieh TC, Ma KH, Chao A (2019) iNEXT: iNterpolation and EXTrapolation for species diversity.
R package version 2.0.19.

James SW, Porco D, Decaens T, Richard B, Rougerie R, Erseus C (2010) DNA Barcoding
Reveals Cryptic Diversity in Lumbricus terrestris L., 1758 (Clitellata): Resurrection of L.
herculeus (Savigny, 1826). *Plos One*, **5**, e15629.
https://doi.org/10.1371/journal.pone.0015629

Joppa LN, Roberts DL, Pimm SL (2011) How many species of flowering plants are there?
Proceedings of the Royal Society B, **278**, 554–559.

Krishnamani R, Kumar A, Harte J (2004) Estimating species richness at large spatial scales
using data from small discrete plots. *Ecography*, **27**, 637–642. https://doi.org/10.1111/j.0906-
7590.2004.03790.x

Lavelle P, Decaens T, Aubert M, Barot S, Blouin M, Bureau F, Margerie P, Mora P, Rossi J-P
(2006) Soil invertebrates and ecosystem services. *European Journal of Soil Biology*, **42**, S3–
S15. https://doi.org/10.1016/j.ejsobi.2006.10.002

Lavelle P, Spain A, Blouin M, Brown GG, Decaëns T, Grimaldi M, Jiménez JJ, McKey D, Mathieu
J, Velasquez E, Zangerlé A (2016) Ecosystem engineers in a self-organized soil: A review of
concepts and future research questions. *Soil Science*, **18**, 91–109.

Linnaeus C (1758) *Systema naturae*. Salvi: Holmiae, Stockholm.

Lomolino MV (2004) Conservation biogeography. In: *Frontiers of biogeography: new directions in*
the geography of nature (eds Lomolino MV, Heaney LR), pp. 293–296. Sunderland, MA.

Maggia M-E, Decaëns T, Lapied E, Dupont L, Roy V, Schimann H, Orivel J, Murienne J, Baraloto
C, Cottenie K, Steinke D (2021) At each site its diversity: DNA barcoding reveals remarkable
earthworm diversity in neotropical rainforests of French Guiana. *Applied Soil Ecology*, **164**,
103932.

Marchán DF, Decaëns T, Domínguez J, Novo M (2022) Perspectives in earthworm phylogeny:
recent advances in Lumbricoidea and standing questions. *Diversity*, **19**, 22.

Marchán DF, Novo M, Domínguez J, da Silva A, Martínez A, Decaëns T (2023) The best is yet to
come: six new species within a large-bodied earthworm genus (Scherotheca, Lumbricidae) in
a densely sampled region (Southwestern France). *Zoological Journal of the Linnean Society*,
20, 1–19.

Marcon E, Mirabel A, Molino J-F, Sabatier D (2024) Estimation of the number of tree species in
French Guiana by extrapolation of permanent plots richness. *Journal of Tropical Ecology*, **in**
press.

Mathieu J, Davies TJ (2014) Glaciation as an historical filter of below-ground biodiversity. *Journal*
of Biogeography, **41**, 1204–1214. https://doi.org/10.1111/jbi.12284

May RM (1992) How many species inhabit the earth? *Scientific American*, **267**, 18–24.

Misirlioğlu M, Reynolds JW, Stojanović M, Trakić T, Sekulić J, James SW, Csuzdi C, Decaëns T,
Lapied E, Phillip H, Cameron EK, Brown GG (2023) Earthworms (Clitellata, Megadrili) of the
world: an updated checklist of valid species and families, with notes on their distribution.
Zootaxa, **5255**, 417–438.

Padial PJ, Miralles A, de la Riva I, Vences M (2010) The integrative future of taxonomy. *Frontiers*
in Zoology, **7**, 16.

Pennisi E (2019) DNA barcodes jump-start search for new species. *Science*, **364**, 920–921.
https://doi.org/10.1126/science.364.6444.920

Phillips HRP, Bach EM, Bartz MLC, Bennett JM, Beugnon R, Briones MJI, Brown GG, Ferlian O,
Gongalsky KB, Guerra CA, Koenig-Ries B, Krebs JJ, Orgiazzi A, Ramirez KS, Russell DJ,
Schwarz B, Wall DH, Brose U, Decaens T, Lavelle P, Loreau M, Mathieu J, Mulder C, van der
Putten WH, Rillig MC, Thakur MP, de Vries FT, Wardle DA, Ammer C, Ammer S, Arai M,
Ayuke FO, Baker GH, Baretta D, Barkusky D, Beausejour R, Bedano JC, Birkhofer K,

Blanchart E, Blossey B, Bolger T, Bradley RL, Brossard M, Burtis JC, Capowiez Y, Cavagnaro TR, Choi A, Clause J, Cluzeau D, Coors A, Crotty FV, Crumsey JM, Davalos A, Diaz Cosin DJ, Dobson AM, Dominguez A, Esteban Duhour A, van Eekeren N, Emmerling C, Falco LB, Fernandez R, Fonte SJ, Fragoso C, Franco ALC, Fusilero A, Geraskina AP, Gholami S, Gonzalez G, Gundale MJ, Gutierrez Lopez M, Hackenberger BK, Hackenberger DK, Hernandez LM, Hirth JR, Hishi T, Holdsworth AR, Holmstrup M, Hopfensperger KN, Huerta Lwanga E, Huhta V, Hurisso TT, Iannone BV, Iordache M, Irmler U, Ivask M, Jesus JB, Johnson-Maynard JL, Joschko M, Kaneko N, Kanianska R, Keith AM, Kernecker ML, Kone AW, Kooch Y, Kukkonen ST, Lalthanzara H, Lammel DR, Lebedev IM, Le Cadre E, Lincoln NK, Lopez-Hernandez D, Loss SR, Marichal R, Matula R, Minamiya Y, Moos JH, Moreno G, Moron-Rios A, Motohiro H, Muys B, Neirynck J, Norgrove L, Novo M, Nuutinen V, Nuzzo V, Rahman PM, Pansu J, Paudel S, Peres G, Perez-Camacho L, Ponge J-F, Prietzel J, Rapoport IB, Rashid MI, Rebollo S, Rodriguez MA, Roth AM, Rousseau GX, Rozen A, Sayad E, van Schaik L, Scharenbroch B, Schirrmann M, Schmidt O, Schroeder B, Seeber J, Shashkov MP, Singh J, Smith SM, Steinwandter M, Szlavecz K, Antonio Talavera J, Trigo D, Tsukamoto J, Uribe-Lopez S, de Valenca AW, Virto I, Wackett AA, Warren MW, Webster ER, Wehr NH, Whalen JK, Wironen MB, Wolters V, Wu P, Zenkova IV, Zhang W, Cameron EK, Eisenhauer N (2021) Global data on earthworm abundance, biomass, diversity and corresponding environmental properties. *Scientific Data*, **8**, 136. https://doi.org/10.1038/s41597-021-00912-z

Phillips HRP, Guerra CA, Bartz MLC, Briones MJI, Brown G, Crowther TW, Ferlian O, Gongalsky KB, van den Hoogen J, Krebs J, Orgiazzi A, Routh D, Schwarz B, Bach EM, Bennett J, Brose U, Decaens T, Koenig-Ries B, Loreau M, Mathieu J, Mulder C, van der Putten WH, Ramirez KS, Rillig MC, Russell D, Rutgers M, Thakur MP, de Vries FT, Wall DH, Wardle DA, Arai M, Ayuke FO, Baker GH, Beausejour R, Bedano JC, Birkhofer K, Blanchart E, Blossey B, Bolger T, Bradley RL, Callaham MA, Capowiez Y, Caulfield ME, Choi A, Crotty FV, Davalos A, Diaz Cosin DJ, Dominguez A, Duhour AE, van Eekeren N, Emmerling C, Falco LB, Fernandez R, Fonte SJ, Fragoso C, Franco ALC, Fugere M, Fusilero AT, Gholami S, Gundale MJ, Gutierrez Lopez M, Hackenberger DK, Hernandez LM, Hishi T, Holdsworth AR, Holmstrup M, Hopfensperger KN, Lwanga EH, Huhta V, Hurisso TT, Iannone BV, Iordache M, Joschko M, Kaneko N, Kanianska R, Keith AM, Kelly CA, Kernecker ML, Klaminder J, Kone AW, Kooch Y, Kukkonen ST, Lalthanzara H, Lammel DR, Lebedev IM, Li Y, Jesus Lidon JB, Lincoln NK, Loss SR, Marichal R, Matula R, Moos JH, Moreno G, Moron-Rios A, Muys B, Neirynck J, Norgrove L, Novo M, Nuutinen V, Nuzzo V, Rahman MP, Pansu J, Paudel S, Peres G, Perez-Camacho L, Pineiro R, Ponge J-F, Rashid MI, Rebollo S, Rodeiro-Iglesias J, Rodriguez MA, Roth AM, Rousseau GX, Rozen A, Sayad E, van Schaik L, Scharenbroch BC, Schirrmann M, Schmidt O, Schroeder B, Seeber J, Shashkov MP, Singh J, Smith SM, Steinwandter M, Talavera JA, Trigo D, Tsukamoto J, de Valenca AW, Vanek SJ, Virto I, Wackett AA, Warren MW, Wehr NH, Whalen JK, Wironen MB, Wolters V, Zenkova IV, Zhang W, Cameron EK, Eisenhauer N (2019) Global distribution of earthworm diversity. *Science*, **366**, 480–485. https://doi.org/10.1126/science.aax4851

Pimm SL (2001) *The World according to Pimm: A Scientist Audits the Earth*. McGraw-Hill., New York, USA.

Pimm SL, Jenkins CN, Joppa LN, Roberts DL, Russell GJ (2010) How Many Endangered Species Remain to be Discovered in Brazil? *Natureza & Conservação*, **08**, 71–77. https://doi.org/10.4322/natcon.00801011

R Core Team (2024) R: a language and environment for statistical computing. Version 4.4.1.

Reynolds JW (1994) Earthworms of the world. *Global biodiversity*, **4**, 11–16.

Rougerie R, Decaëns T, Deharveng L, Porco D, James SW, Chang C-H, Richard B, Potapov M, Suhardjono Y, Hebert PDN (2009) DNA barcodes for soil animal taxonomy. *Pesquisa Agropecuaria Brasileira*, **44**, 789–802. https://doi.org/10.1590/S0100-204X2009000800002

Stork N (1993) How many species are there? *Biodiversity and Conservation*, **2**, 215–232.

Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH (2012) Environmental DNA. *Molecular*
Ecology, **21**, 1789–1793. https://doi.org/10.1111/j.1365-294X.2012.05542.x

Touroult J, Pollet M, Pascal O (2018) Overview of Mitaraka survey: research frame, study site
and   field   protocols.   *Zoosystema*,   **40**,   327–365.
https://doi.org/10.5252/zoosystema2018v40a13

Wardle DA, Bardgett RD, Klironomos JN, Setälä H, van der Putten WH, Wall DH (2004)
Ecological linkages between aboveground and belowground biota. *Science*, **304**, 1629–1633.
https://doi.org/10.1126/science.1094875

Wilson EO (2004) Taxonomy as a fundamental discipline. *Philosophical Transactions of the*
Royal Society B - Biological Sciences, **359**, 739–739. https://doi.org/10.1098/rstb.2003.1440