High early lactational synchrony within baboon groups predicts increased infant mortality

Authors

Jack C. Winans[1,2,3,a], Niki H. Learn[4], I. Long'ida Siodi[5], J. Kinyua Warutere[5], Elizabeth A. Archie[6], Jenny Tung[7,8,9,10,11], Susan C. Alberts[8,9,*], A. Catherine Markham[12,*]

Author Affiliations

[1]Interdepartmental Doctoral Program in Anthropological Sciences, Stony Brook University, Stony Brook, NY, USA

[2]Department for the Ecology of Animal Societies, Max Planck Institute of Animal Behavior, Konstanz, Germany

[3]Department of Biology, University of Konstanz, Konstanz, Germany

[4]Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA

[5]Amboseli Baboon Research Project, Amboseli National Park, Kajiado, Kenya

[6]Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA

[7]Department of Primate Behavior and Evolution, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

[8]Department of Evolutionary Anthropology, Duke University, Durham, NC, USA

[9]Department of Biology, Duke University, Duke University, Durham, NC, USA

[10]Canadian Institute of Advanced Research, Toronto, Canada

[11]Faculty of Life Sciences, University of Leipzig, Leipzig, Germany

[12]Department of Anthropology, Stony Brook University, Stony Brook, NY, USA

*These authors contributed equally to this work.

[a]To whom correspondence may be addressed (email: jwinans@ab.mpg.de).

Author Contributions

JCW, ACM, and SCA designed research; SCA, EAA, and JT administered the long-term research project; SCA, EAA, JT, JCW, and ACM acquired funding; JCW, ILS, JKW, and NHL performed research; JCW analyzed data; JCW, ACM, SCA, EAA, JT, ILS, JKW, and NHL wrote the paper.

Competing Interest Statement

The authors declare no competing interest.

Classification

Biological Sciences: Anthropology

Keywords

female-female competition; reproductive synchrony; female infanticide

Abstract
Social group composition can have fitness implications for group members by
determining opportunities for affiliative and competitive interactions. Female-female
competition may be particularly acute when many groupmates have young infants at the same
time, with potential consequences for infant survival. Here, we used decades of data on wild
baboons (*Papio* sp.) in Amboseli, Kenya, to examine the effects of 'early lactational synchrony'
(here, the proportion of females in a group with an infant <90 days old) on female-female
agonistic interactions and infant survival. Because early lactation is an energetically demanding
time for mothers and a risky time for infants, we expected early lactational synchrony to produce
intensified competition for food and/or male protectors, resulting in more frequent female-female
agonistic interactions and high infant mortality. In support of these predictions, we found that the
frequency of female-female agonistic interactions increased with increasing early lactational
synchrony. Reproductive state affected this relationship: while females in all states (cycling,
pregnant, and postpartum amenorrhea) initiated more agonistic interactions when early
lactational synchrony was high, only females in postpartum amenorrhea (including, but not
limited to, females in early lactation) received more agonistic interactions. Furthermore, while
high early lactational synchrony was rare, it strongly predicted infant mortality. This association
may result from both aggression among adult females and infanticidal behavior by peripubertal
females. These findings provide novel evidence that social dynamics may shape reproductive
phenology in a nonseasonal breeder. Specifically, both competition among reproductive females
and harassment from nonreproductive females may select against synchronous reproduction.

64
Significance Statement
Competition between females is a powerful force shaping the behavior and fitness of wild
animals, but how it is exacerbated (or mitigated) by the overall composition of social groups is
understudied. We asked if female-female competition is enhanced when more females in baboon
social groups have young infants, a time that is particularly energetically demanding for mothers
and particularly dangerous for their offspring. In these settings, we found that both adult female
competitive behavior and infant mortality risk increased. Strikingly, almost all recorded cases of
infanticide by adolescent, nulliparous females in our study population occurred on days when
many mothers had young infants. These results highlight how social interactions may shape the
timing of reproduction in wild social mammals.
75
Background
Competition between conspecifics is an important selective force acting on social
behavior, morphology, and life history (1-3). In many vertebrates, females compete with each
other for food, mates, social partners, and other resources, with tactics ranging from mating
interference to infanticide (4-10). This competition can have important fitness effects. For
example, in many species, high dominance rank or increased aggressive behavior is associated
with higher female reproductive success (11-16). Intense female-female competition can even be
sufficient to prevent the emergence of group living (17), while moderate female-female
competition can shape many aspects of sociality among group-living species, from the
differentiation of individual relationships to the structuring of groups according to kinship or
dominance hierarchies (18, 19).
For group-living species, group demography is expected to shape the dynamics of
female-female competition. Because female reproduction is energetically costly and food

resources are often limited, group size may affect the level of female competition and its fitness
consequences (19, 20). Indeed, studies both within and across species report increased rates of
agonistic interactions among females with increasing group size (21-26). Other studies have
implicated female-female competition as a cause of depressed reproductive success or survival in
larger groups (14, 27-29). Although large group size can also benefit females (e.g., via
cooperative breeding (30, 31) or advantages in intergroup competition (32, 33)), its relationship
with female-female competition highlights one potential "top-down" pathway linking group
demography to fitness.
However, group size is not the sole determinant of the intensity of competition: group age
structure, sex ratio, and female reproductive state can also modulate competitive intensity
because they contribute to body size- and state-dependent variation in energetic demands (34,
35). Among female social mammals, reproductive state is a particularly salient source of
variation in competitive landscapes because a female's resource and social requirements vary
depending on whether she is cycling, pregnant, lactating, or reproductively quiescent (36). For
example, female bottlenose dolphins (*Tursiops aduncus*) avoid male kin in favor of unrelated
males while sexually receptive, presumably to minimize the risk of inbreeding (37), but avoid
unrelated males while lactating, when the risk of infanticide by unrelated males is high (38).
Accordingly, shared requirements can promote association by females in similar reproductive
states (39), but shared requirements can also increase competition for food, mates, and/or
allomaternal care (5).
The importance of female reproductive state in shaping female-female competition is
supported by the observation that conspicuous female-female competition is more frequent
among species that exhibit systematically high reproductive synchrony, as expected if females
compete over reproductive state-specific resources (3, 5). For example, in topi antelopes
(*Damaliscus lunatus*) more than 90% of calves are conceived within an annual 1.5-month period
that is marked by aggressive mating interruptions. Similarly, female red deer (*Cervus elaphus*),
female feral horses (*Equus caballus*), and female golden snub-nosed monkeys (*Rhinopithecus
roxellana*) display increased rates of aggression and/or disruption of other females' copulations
during their short breeding seasons (40-43). Even in nonseasonal breeders, individuals can
sometimes experience high reproductive synchrony and competition with groupmates. For
instance, in chacma baboons, sexually receptive females experience heightened aggression from
other sexually receptive females, and also from females who have recently conceived, pointing
to competition over mates and paternal investment when females cycle relatively synchronously
(6, 44). However, the overall effects of reproductive synchrony on female competition in
nonseasonal breeders, and especially the fitness consequences of competition resulting from such
synchrony, remain poorly understood. Addressing this gap is important for understanding the
relationship between resource availability, resource competition, and reproductive timing, which
can vary even between closely related, ecologically similar species (45, 46).
In this study, we investigated how the fitness consequences of female-female competition
are affected by one form of reproductive synchrony—early lactational synchrony—in wild
baboons (*Papio* sp.), where the importance of female-female competition is well-established.
Female-female competition for resources and mates has been studied closely in several baboon
species, including chacma (*P. ursinus*), anubis (*P. anubis*), and yellow baboons (*P.
cynocephalus*). These species live in large, multimale, multifemale groups characterized by
female philopatry and male dispersal (47). Females form stable, linear dominance hierarchies
and typically experience nepotistic 'inheritance' of maternal dominance rank (48-50). Rank-

mediated competition for food is likely a prominent force shaping the social behavior of female anubis and yellow baboons because agonistic behaviors are more common among females that experience greater energetic burdens and during periods when food availability is thought to be lower (51-53). Females of both yellow and chacma baboons also compete for mates and/or protective male associates who buffer mothers and their infants from harassment by other adult and immature females (6, 44, 54, 55). Reproductive state influences the rate at which female baboons initiate and receive aggression (6, 44, 51, 52, 56), and previous studies of reproductive synchrony in baboons suggest that the composition of female reproductive states in baboon groups may be an important predictor of competitive behavior (45, 57, 58). Nonetheless, the ultimate consequences of reproductive synchrony for mothers or their infants in baboons and other nonseasonal breeders remains unclear.

Our study subjects were members of multiple social groups in a population of wild baboons (*P. cynocephalus* with significant *P. anubis* admixture: 59-60) in the Amboseli basin of southern Kenya, which has been continuously monitored since 1971 (61). Females in this population give birth year-round, and the first 90 days after birth (hereafter, 'early lactation') constitutes a period of both high energetic demands for mothers and increased risk of harassment from other females towards newborn infants (62, 63). We defined early lactational synchrony as the proportion of adult females in a social group with dependent infants <90 days old. Our main hypothesis was that groups exhibiting high early lactational synchrony would experience intensified intragroup competition, and that intensified competition would lead to negative fitness-related outcomes for both mothers and their dependent offspring. We tested two predictions generated by this hypothesis. First, we predicted that female-female agonistic interactions would increase in frequency with higher lactational synchrony, either for all adult females or for females in specific reproductive states (considering ovarian cycling, pregnant, and postpartum amenorrhea). Second, we predicted that infant survival would be compromised when early lactational synchrony was higher, because increased competition between mothers could negatively impact their dependent offspring. We also examined the most prevalent sources of infant mortality during periods of high early lactational synchrony to gain insight into the specific risks imposed by female-female competition.

Results

Characterizing early lactational synchrony across the study period. We began by measuring the proportion of females in each social group who were in early lactation on each day (i.e., in the first 90 days of postpartum amenorrhea with a living infant), across all group-days of observation from November 1976 to December 2021. This proportion varied considerably across the study period, from group-days on which no females (0.000) were in early lactation to group-days on which more than 70% (0.714) of females were in early lactation (Fig S1). This distribution was heavily skewed toward zero, such that days of high early lactational synchrony were rare, as expected for this nonseasonally breeding species. The mean proportion of females in early lactation on a given group-day was 0.121 (SD: 0.102; median = 0.105, 75th percentile = 0.176 of the group's females per group-day; 95th percentile = 0.316 of the group's females per group-day). On average, groups had 1.96 total females in early lactation on any given day (SD: 1.74 females; median = 2 females, 75th percentile = 3 females, 95th percentile = 5 females; group sizes ranged in our data set from 3 to 33 adult females per group).

Female agonistic behavior. Using behavioral data collected from 1999-2021 (when we
could control for observer effort, see Methods), we found that the relative frequency of adult
female-female agonistic interactions increased in months when a higher mean proportion of the
females in a group were in early lactation. Adult females both initiated relatively more agonistic
interactions with other adult females and received relatively more agonistic interactions from
other adult females during months when their group had higher mean early lactational synchrony
(generalized linear mixed models, GLMMs; agonistic interactions initiated: $\beta = 0.90$, 95% CI:
[0.71, 1.08], Tables S1-2; agonistic interactions received: $\beta = 0.66$, 95% CI: [0.49, 0.84], Tables
S3-4).
Notably, females of all reproductive states initiated agonistic interactions with other
females more frequently when a higher proportion of females were in early lactation (Fig. 1a).
Specifically, when we partitioned the dataset by female reproductive state (ovarian cycling,
pregnant, or postpartum amenorrhea), we found that females in all three reproductive states
initiated more agonistic interactions with all other adult females during group-months of higher
mean early lactational synchrony (GLMM for cycling females: $\beta = 0.74$, 95% CI: [0.39, 1.08];
Tables S5-6; for pregnant females: $\beta = 1.48$, 95% CI: [1.11, 1.84]; Tables S7-8; for females in
postpartum amenorrhea: $\beta = 0.51$, 95% CI: [0.23, 0.79]; Tables S9-10). For instance, in group-
months when mean early lactational synchrony was in the 75th percentile (a mean of 16.5% of
females had infants in the first 90 days of life in that month), cycling females were predicted to
initiate 8.51% more agonistic interactions compared to when mean early lactational synchrony
was in the 25th percentile (a mean of 5.4% of females had infants in the first 90 days of life in
that month). Similarly, pregnant females were predicted to initiate 17.8% more agonistic
interactions, and females in postpartum amenorrhea were predicted to initiate 5.81% more
agonistic interactions, when mean early lactational synchrony was in the 75th versus the 25th
percentile.
However, higher early lactational synchrony only predicted an increased frequency of
received agonistic interactions for females who were in postpartum amenorrhea (GLMM for
females in postpartum amenorrhea: $\beta = 1.10$, 95% CI: [0.83, 1.37]; Tables S11-12; for cycling
females: $\beta = 0.02$, 95% CI: [-0.16, 0.20]; Tables S13-14; for pregnant females: $\beta = 0.26$, 95% CI:
[-0.18, 0.71]; Tables S15-16, Fig. 1b). Specifically, females in postpartum amenorrhea were
predicted to receive 13.0% more agonistic interactions in months when mean early lactational
synchrony was in the 75th compared to the 25th percentile. Cycling and pregnant females did not
receive significantly more or fewer agonistic interactions during months of high versus low mean
early lactational synchrony.
We also tested for a relationship between early lactational synchrony and the number of
agonistic interactions exchanged within dyads of females in the same reproductive state. In
group-months with higher mean early lactational synchrony, pairs of cycling females did not
exchange significantly more or fewer agonistic interactions (GLMM; $\beta = 0.18$, 95% CI: [-0.28,
0.64]; Table 1 and S17). However, dyads of pregnant females and females in postpartum
amenorrhea exchanged more agonistic interactions in group-months with higher mean early
lactational synchrony (GLMM for pregnant females: $\beta = 1.02$, 95% CI: [0.55, 1.48]; for females
in postpartum amenorrhea: $\beta = 0.66$, 95% CI: [0.39, 0.92]; Table 1 and S18-19). For instance,
when group-months were in the 75th versus the 25th percentile of mean early lactational
synchrony, pairs of pregnant females were predicted to exchange 11.92% more agonistic
interactions and pairs of females in postpartum amenorrhea were predicted to exchange 7.56%
more agonistic interactions.

226
Infant survival. Using data collected from 1976-2021, we next employed a time-varying
Cox proportional hazards model to test whether infant survival was compromised when early
lactational synchrony was high. Controlling for other covariates thought to explain infant
mortality (maternal death, maternal rank, maternal age, group size, and rainfall), early lactational
synchrony strongly predicted infants' daily mortality risk during the first year of life (β = 4.45,
95% CI: [3.46, 5.44]; Table 2 and S20, Fig. 2b). For instance, on days when a group was in the
75th percentile of early lactational synchrony (17.6% of females in a group were in early lactation
that day), infants were predicted to have a 75.2% increased risk of mortality relative to days
when the group was in the 25th percentile (5.0% of females in a group were in early lactation that
day). We also tested the same covariates in a Cox proportional hazards model predicting daily
infant survival up to 90 days after birth (coincident with the period in which their mothers would
be in early lactation). This model produced qualitatively similar results (Tables S21-22).
239
Infanticide. To understand which sources of infant mortality were most prevalent during
periods of high early lactational synchrony, we reviewed the most probable causes of death for
infants included in the survival analysis (i.e., from 1976-2021). Deaths from infanticide were
proportionally more common when early lactational synchrony was high (16.4% of infant deaths
on days in the top quartile of early lactational synchrony versus 7.27% of all infant deaths; Figs.
2c and S2). Further, of the 37 potentially infanticidal deaths that occurred during this period, 22
occurred on days with very high early lactational synchrony (>80th percentile for all group-days;
Table S23). When we limit these potential infanticides to those that were directly observed or
strongly suspected with corroborating evidence (n = 24; see Methods) and were performed by
females in the same group as the mother and infant (n = 14), 12 (85.7%) occurred on days when
groups were in the top quartile of early lactational synchrony overall. This was significantly
greater than the proportion of all infant deaths that occurred in the top quartile of early lactational
synchrony (109/300 deaths, 36.3%; Fisher's exact test: odds ratio = 10.4, 95% CI: [2.3, 97.8], *p*
= 0.0003). The vast majority of these within-group infanticides by females (11 of 14; 78.6 %)
were attributed to female "kidnapping" events, in which a female separated an infant from its
mother for an extended period of time and the infant subsequently died from starvation,
dehydration, and/or rough handling. The remaining three within-group infanticides by females
(21.4%) were thought to have occurred during episodes of severe, multiparty aggression that
preceded a permanent social group fission. Although infanticide by females accounts for a
substantial fraction of infant mortality during high synchrony periods, all 11 of the fatal female
kidnappings were performed by peripubertal, nulliparous females (Table S23). Hence, these fatal
kidnappings are not examples of direct competition among adult females for immediate
resources. In contrast, the three infanticides thought to be the result of severe, multiparty
aggression probably involved fully adult females, although the identities of the perpetrators are
unknown (Table S23).
265
Discussion
These results provide considerable evidence that group composition, specifically the
proportion of adult females in early lactation, influences female-female competition and infant
survival in baboons. Higher levels of early lactational synchrony were linked to more frequent
agonistic interactions between adult females, suggesting that early lactational synchrony
increases intragroup, female-female competition over food and/or access to protective male

associates. High early lactational synchrony predicted an increased frequency of agonistic
interactions initiated by adult females in all reproductive states. However, high early lactational
synchrony only predicted a higher frequency of received agonistic interactions for females in
postpartum amenorrhea, suggesting that lactating females disproportionately bear the costs of
high synchrony. At the dyadic level, pairs of pregnant females and pairs of females in
postpartum amenorrhea also exchanged more agonistic interactions when early lactational
synchrony was higher, but pairs of cycling females did not.

Early lactational synchrony also strongly predicted infant mortality. Notably, infanticides,
especially those performed by peripubertal females, were overrepresented in periods of high
early lactational synchrony relative to infant deaths overall. Taken together, these results suggest
that infant baboons are more likely to die in highly synchronous periods. This increased risk may
be caused by two phenomena. First, the increased risk of kidnapping by young, nonreproductive
females during periods of high early lactational synchrony contributed substantially to the
mortality we measured, either immediately or on future days due to the costs of prolonged
maternal separation. Second, the increased rates of aggression directed towards females in
postpartum amenorrhea when early lactational synchrony was high may decrease the viability of
dependent offspring. These two explanations are not mutually exclusive: increased rates of
aggression overall may also grant peripubertal females more opportunities to succeed in
kidnapping attempts. Therefore, the negative fitness consequences of group-level synchrony in
lactation may select against strong breeding seasonality, even in an environment characterized by
highly seasonal resource fluctuations.

Female agonistic behavior. Our finding that agonistic interactions between adult females
increased with early lactational synchrony corroborates previous findings in female chacma
baboons in Tsaobis, Namibia, and female yellow baboons in Mikumi, Tanzania (44, 57).
Researchers in Tsaobis have argued that a positive correlation between aggression and group-
level reproductive synchrony reflects competition for mates and access to paternal care (44, 45).
On the other hand, researchers in Mikumi cited the reduction of the future competition faced by
offspring as the most likely explanation for reproductive synchrony-related multiparty attacks
(57, 58). In our study, increased aggression by pregnant and cycling females toward females in
postpartum amenorrhea supports some of the predictions of the "future competition reduction"
hypothesis. However, we believe that competition over current food resources or available male
protectors are more parsimonious explanations for the observed patterns in Amboseli. Because
high early lactational synchrony should increase group-level energetic demand and therefore
average daily group travel distances (34, 64), females of all reproductive states will incur the
energetic costs of high early lactational synchrony. Additionally, because protective male-female
associations often form around the time of conception, peak in early lactation, and can persist
until offspring independence (65), the patterns of aggression we observed also conform to
expectations of competition for male social partners. As Baniel et al. (44) highlight, pregnant,
lactating, and ovulating female baboons all potentially benefit from close association with males.
This observation, combined with the peak in male-female association during early lactation, may
explain why females in postpartum amenorrhea are the most frequent recipients of synchrony-
related agonistic behavior from other adult females. Such an explanation avoids the need to
invoke more complex cognitive mechanisms involving the projection of future competitive
landscapes.

Similarly, our dyadic analysis revealed that increased aggression in periods of high early lactational synchrony is not strictly linked to competition over resources sought exclusively by new mothers. Rather, dyads of both pregnant females and females in postpartum amenorrhea exchanged more agonistic interactions in these periods. This observation lends support to the hypothesis of competition over food, as both pregnant females and females in postpartum amenorrhea must invest energy in growing offspring. However, it does not rule out the hypothesis of competition over access to male associates: unlike females in other reproductive states, fertile, cycling females may attract males without engaging in much active competition. Nevertheless, we note that the potential hypotheses (i.e., competition over current vs. future resources; competition over food vs. male associates) are not mutually exclusive and may be dependent on other characteristics (e.g., rank, social bonds).

Infant survival and female infanticide. An unexpected and striking observation was the rise in infanticidal behavior by peripubertal females during periods of high early lactational synchrony. We suggest that synchrony-related infanticidal behavior by peripubertal females is not directly related to competition. If infanticide serves to reduce current competition, we would expect parous females rather than peripubertal (i.e., non-reproductive, nulliparous) females to commit infanticide when early lactational synchrony is high, especially because nulliparous females are unlikely to conceive in close temporal proximity to the births of the infants they target. Contrary to this expectation, parous females rarely engage in kidnapping, even when they have no current infants of their own. Furthermore, if infanticide serves to reduce future competition, neither peripubertal nor parous females should restrict their infanticidal attacks to periods of high early lactational synchrony, because the intensity of competition in the future is unlikely to be predictable at any given time and because female baboons compete with more individuals than their immediate birth cohort.

One explanation for the increased rates of kidnapping by peripubertal females during periods of high early lactational synchrony is that the unusual abundance of young infants may spur higher-than-usual interest in infants on the part of peripubertal females. If so, the probability that interactions with infants escalate to kidnapping may increase. A second possibility is that, because infants of lower-ranking mothers are likely more easily kidnapped than infants of higher-ranking mothers, high early lactational synchrony may simply increase the probability that any given peripubertal female has access to infants of lower-ranking mothers. However, a *post hoc* analysis casts some doubt on this second possibility: during periods of high early lactational synchrony, an early lactating mother was *less* likely to have a higher-ranking peripubertal female as her nearest neighbor than during other periods (see Supplementary Methods; GLMM: β = -1.29, 95% CI: [-2.37, -0.21], Tables S24-S25). This pattern suggests that mothers avoid the most probable kidnappers in their group when early lactational synchrony is high, especially since early lactational synchrony did not predict whether early lactating mothers were likely to have *lower*-ranking peripubertal females as their nearest neighbors (GLMM: β = -0.12, 95% CI: [-0.77, 0.52], Tables S26-S27).

While the drive to kidnap infants in peripubertal females is not likely directly related to competition among adult females, adult female competition could interact with the interest of peripubertal females in infants if the increased rates of agonistic interactions between adult females make kidnapping attempts more likely to be successful. Furthermore, the relationship between infant kidnapping and early lactational synchrony may be compounded if some mother-

362 infant pairs are competitively excluded from receiving male protection because of the unusually
363 high number of new mothers per male in the group.
364
Conclusions. The patterns of female aggression, infant survival, and female infanticide
reported here provide new evidence that group-level synchrony in female reproductive state can
negatively impact breeding success at the individual level, even in a non-seasonal breeder. First,
an increase in the proportion of females with young infants was associated with higher relative
frequencies of adult female-female agonistic behavior, indicating intensified competition.
Additionally, this synchrony predicted higher infant mortality, likely due to a combination of
infanticidal behavior by peripubertal females and the downstream effects of female-female
competition and aggression. These results set the stage for future studies testing the proximate
factors shaping female aggression, the underlying mechanisms motivating female interactions
with non-offspring infants, and the roles of inter- and intrasexual social interactions in shaping
the timing of reproduction.
376
Methods
Study site and subjects. This population of wild baboons lives in the Amboseli basin, a
semi-arid short-grass savannah ecosystem at the northern base of Mt. Kilimanjaro in southern
Kenya. The basin experiences an annual dry season from June to October when virtually no rain
falls, followed by a highly variable wet season from November to May when the amount of
rainfall varies greatly from year to year (67). The baboon population has been continuously
monitored by the Amboseli Baboon Research Project (ABRP) since 1971 (61). Over the course
of the study's history, the number of habituated social groups monitored at any one time has
ranged from one to six, and the groups have undergone multiple fissions and fusions since the
late 1980s. See Supplementary Methods for additional details on daily demographic, behavioral,
and ecological monitoring by ABRP.
388
Female reproductive states. In our analyses, we considered both individual female
reproductive states and a group-level metric of reproductive state similarity as important
predictors of individual behavior. We identified four female reproductive states (see
Supplementary Methods for details, also (66)). (i) Ovarian cycling lasted from the first day a
female's sex skin was turgescent after menarche or her last live birth to either the next onset of
pregnancy or death. (ii) Pregnancy lasted from the first day a female's sex skin was deturgescent
in her last sexual cycle preceding a known pregnancy to the day before she gave birth,
experienced a spontaneous abortion, or died. (iii) Postpartum amenorrhea lasted from the day a
female gave birth or experienced a spontaneous abortion to the next day her sex skin was
turgescent or she died. (iv) Early lactation refers to a subset of the period of postpartum
amenorrhea, specifically the first 90 days of postpartum amenorrhea with a live infant. Any
female who was in early lactation was, by definition, in postpartum amenorrhea, but postpartum
amenorrhea typically extends long past the first 90 days (mean ± SD length of postpartum
amenorrhea: 322 ± 87 days, (66)). Throughout, we use the term early lactational synchrony to
refer to the proportion of adult females in a group that were in early lactation, i.e., had an infant
<90 days old. See the Supplementary Methods for further information on how early lactational
synchrony was calculated.
406

Statistical approach. We took an information theoretic approach to all analyses, which
were conducted in R v4.0.3 (68). We began by constructing "full models" that included all likely
covariates. We then used the *dredge* function in the *MuMIn* R package (69) to acquire Akaike's
information criterion (AIC_c) values for all models with all possible combinations of fixed effects
included in the full model (70, 71). All candidate models within 2 AIC_c values of the candidate
model with the lowest AIC_c value were selected as the best-fitting models. We used *MuMIn*'s
model.avg function to obtain a weighted average of the best-fitting models (those within 2 AIC_c
of the best model) and estimate effect sizes for individual fixed effects. If a fixed effect was left
out of a particular candidate model, then a weighted effect size of zero was included in the
calculation of the overall weight-averaged effect size for that fixed effect. We considered fixed
effects significant when the 95% confidence interval of their weight-averaged effect size did not
overlap zero. Cox proportional hazards models were constructed using the *coxph* function from
the package *survival* (72) and GLMMs were constructed using the package *glmmTMB* (73).

Female agonistic behavior. To test the prediction that early lactational synchrony is
associated with increased adult female-female competition, we aggregated data collected from
October 1999 to December 2021 on all intragroup dyadic agonistic interactions occurring
between adult females in which either one female displayed aggressive behaviors and the other
displayed submissive behaviors or in which one female spatially displaced the other. We then
tallied, for each female in each month of her adult life (each "female-month"), the number of
agonistic interactions that she initiated (i.e., interactions in which she displayed aggressive
behavior or she spatially displaced the other female). Next we tallied, for each female in each
month of her adult life, the number of agonistic interactions she received (i.e., interactions in
which she displayed submissive behavior in response to aggression or she was spatially
displaced). We restricted our behavioral analysis to 1999-2021 because this was the period of
time during which we were best able to control for observer effort. Specifically, we calculated
these tallies only for female-months in which our records of observation times were of the
highest quality, and during which observers were with the female's group for at least 30 hours.
We also assigned each female-month to one of three reproductive states (cycling, pregnant, or
postpartum amenorrhea), as described above. When a female transitioned from one reproductive
state to another or changed social group membership within a calendar month, we split that
female-month at the date of transition, and counted agonistic interactions before the transition as
belonging to the first state or social group, and those after the transition to the second state or
social group. Thus, each count is informative of a female in one social group and in one
reproductive state. For counts of agonistic interactions both initiated and received, our sample
sizes reflected 21,321 unique female-reproductive state-group-months for 278 individual females
in 14 social groups.
We then used these counts of agonistic interactions initiated and received as response
variables in two negative binomial GLMMs with log link functions. We chose to model these
counts as negative binomial distributions because they were overdispersed. In both models, we
included the following fixed effects: (i) the average proportion of adult females in the group who
were in early lactation for that month, (ii) the female's proportional dominance rank, (iii) the
female's age, (iv) the female's reproductive state (cycling, pregnant, or postpartum amenorrhea),
(v) season, and (vi) a measure of observer effort to control for variation in observation intensity
across the study period (see Supplementary Methods). In both models, we controlled for
variation in the duration of female reproductive state-group-months (e.g., due to transitions in

reproductive state) by including an offset term which was calculated as the the log-transformed
number of days in the month in which the female was in her present group and reproductive
state. We included individual ID and group ID as random effects to control for repeated sampling
of females and groups. In all models presented in the main text, we did not include a fixed effect
of group size because group size is strongly negatively correlated with our measure of observer
effort. However, to assess the potential role of group size, we built an alternative model structure
which instead incorporated observer effort into the offset term and included group size as a fixed
effect (see Supplementary Methods and Results), which produced similar results (Tables S28-
49).
To determine if females in different reproductive states experienced different
consequences of early lactational synchrony, we partitioned the agonistic interaction dataset by
the reproductive state of the female in question, resulting in 6,861 female-months for 273 cycling
females in 14 social groups, 6,095 female-months for 254 pregnant females in 14 social groups,
and 8,365 female-months for 245 postpartum amenorrhea females in 14 social groups. For each
of these three datasets, we again modeled the monthly counts of agonistic interactions either
initiated or received using negative binomial GLMMs with log link functions (see also
Supplementary Methods), but removing the female reproductive state term. We included all
other fixed and random effects that were included in the overall agonism models, as well as the
log-transformed offset term for the number of days in the month in which the female was a
member of her group and in her present reproductive state.
To investigate the relationship between early lactational synchrony and agonistic
interactions at the dyadic level, we identified all possible pairs of females that were in the same
reproductive state in each social group and each month and tallied the total number of agonisms
they exchanged in that month (cycling: N=23,547 dyad-months across 3,703 dyads in 14 groups;
pregnant: N=18,677 dyad-months across 2,746 dyads in 14 groups; postpartum amenorrhea:
N=35,975 dyad-months across 2,891 dyads in 14 groups). We constructed Poisson GLMMs with
a log link function predicting the total number of agonistic interactions exchanged in each dyad-
group-month for each of these three reproductive states (see Supplementary Methods). We chose
to model these counts using a Poisson distribution because, unlike in the previous agonism
models, the count data for agonistic interactions exchanged within dyads were not overdispersed.
As fixed effects in all three models we included: (i) the average proportion of adult females in
the group who were in early lactation for that month, (ii) the absolute value of the difference
between the two females' proportional ranks, (iii) season, and (iv) observer effort. We also
included the logged number of days in the month that both females were present in the group and
in the same state as an offset term, as well as dyad ID and group ID as random effects to control
for repeated sampling of dyads and groups.
Because only one observer typically collects observational data on any given day, we are
not able to record all true occurrences of agonistic interactions. Observed interaction counts per
capita are therefore typically low, especially in large groups (mean ± SD of agonistic interactions
initiated = 1.82 ± 3.06; mean ± SD of agonistic interactions received = 1.82 ± 2.64). Therefore,
while interpreting the results of these models, even small effect sizes for our count response
variables can indicate meaningful differences in the relative frequencies of agonistic interactions.
495
496   *Infant survival.* To test our prediction that high early lactational synchrony is associated
497 with an increased risk of infant death, we used a Cox proportional hazards model with time
498 varying covariates to predict the daily risk of infant death in the first year of life (see

Supplementary Methods). Our covariates were: (i) the proportion of adult females in early lactation on each day of the infant's life; (ii) group size; (iii) whether or not the infant's mother was dead; (iv) maternal proportional rank; maternal age as a (v) linear and (vi) quadratic term; and the (vii) cumulative total rainfall that fell in the last 90 days. If an infant lived at least a day after its mother's death (n = 19 infants), then the covariate data for maternal rank and age were entered as the mother's rank and age on her death date. These post-maternal death data points were rare, constituting 0.05% of the total dataset. In total, 1,323 infants born between November 1976 and December 2021 were included in this analysis, of whom 300 died in the first year of life, and 87 were right censored because observations on them ceased before they reached their first birthday. We restricted our survival analyses to this period because rainfall data were not recorded before November 1976. To determine if the effect of early lactational synchrony on infant survival differed when infants themselves had early lactating mothers, we ran another Cox proportional hazards model predicting the daily risk of infant death from the day of birth until the infant was 90 days old using the same covariates. This model included 1,304 infants born between November 1976 and December 2021 (140 deaths in the first 90 days of life; 30 infants were right censored).

Infanticide risk. After observing a negative relationship between early lactational synchrony on infant survival, we sought to investigate the causes of death for infants who died during periods of high early lactational synchrony (see Supplementary Methods for information on how we assigned causes of death and confidence in the assigned causes). Of the 300 infant deaths included in our survival analysis, causes of death with our highest confidence level could be assigned to 75 individuals.

Following this assignment process, we visualized the relative proportion of each cause of death for (i) all infants, (ii) those infants who died on days in the lowest quartile of early lactational synchrony across all infant death days (< 0.076 of females in the group in early lactation, n = 72 deaths), and (iii) those infants who died on days in the highest quartile of early lactational synchrony across all infant death days (> 0.222 of females in the group in early lactation, n = 75 deaths, Fig. S1). In this assessment, any infant whose cause of death was not assigned with our highest level of confidence was considered to have died of an unknown cause. Infanticide deaths appeared to be overrepresented during the periods of highest early lactational synchrony, relative to all deaths and relative to deaths that occurred in the lowest quartile of early lactational synchrony. This observation led us to review field notes related to each infanticide death to determine the most likely actors (male or female) and contexts (within-group or between-group) of infanticide when early lactational synchrony was high (Table S21). In assigning causes of death, we use the definition of infanticide proposed by Digby (74) and more recently invoked by Lukas and Huchard (7): "an act that makes a direct or significant contribution to the immediate or imminent death of conspecific young." This definition includes acts of physical aggression toward infants and acts such as kidnappings that cause infants to die via "enforced neglect" (7). It thus focuses on the ultimate consequences of interactions with conspecific young, regardless of proximate motivation.

Acknowledgments

We thank Jeanne Altmann for her pivotal role in stewarding the Amboseli Baboon Research Project and in designing many of the ABRP's long-term data collection protocols. We thank the Max Planck Institute for Evolutionary Anthropology, Duke University, Princeton

University, the University of Notre Dame, and Stony Brook University for financial and logistical support. In Kenya, our research was approved by the Wildlife Research Training Institute (WRTI), Kenya Wildlife Service (KWS), the National Commission for Science, Technology, and Innovation (NACOSTI), and the National Environment Management Authority (NEMA). We also thank the University of Nairobi, the Kenya Institute of Primate Research (KIPRE), the National Museums of Kenya, the members of the Amboseli-Longido pastoralist communities, the Enduimet Wildlife Management Area, Ker & Downey Safaris, AirKenya, and Safarilink for their cooperation and assistance in the field. We are particularly grateful to Raphael Mututua, Serah Sayialel, and Lilian Musembei (members of the ABRP long-term field team), to Tim Wango and Vivian Oudu for assistance in Nairobi, and to Jacob Gordon and William Wilbur who expertly manage the baboon project database, BABASE. Database design and programming are provided by Karl Pinc. We also thank Andreas Koenig and Heather Lynch who provided feedback on an earlier version of this manuscript, as well as Jacob Feder and the Behavioral Ecology Group at Stony Brook University for helpful discussion at an early stage of this analysis. This research was approved by the IACUCs at Duke University, University of Notre Dame, and Princeton University, and the Ethics Council of the Max Planck Society and adhered to all the laws and guidelines of Kenya.

Funding

Long-term data collection in Amboseli has been supported by the National Science Foundation and National Institutes of Health, most recently through R01AG071684, R01AG075914, R01AG053308, P01AG031719, and R61AG078470. Current support for field-based data collection also comes from the Max Planck Institute for Evolutionary Anthropology. Further support for this specific project came from the National Science Foundation (DDRIG 2141839, GRFP 1839287) and the Wenner-Gren Foundation (10355). Please see https://amboselibaboons.nd.edu/acknowledgements/ for a complete list of funding acknowledgments.

References

1. A. J. Wilson, Competition as a source of constraint on life history evolution in natural populations. *Heredity* **112**, 70-78 (2014).

2. K. A. Judge, V. L. Bonano, Male weaponry in a fighting cricket. *PLoS One* **3**, e3980 (2008).

3. S. T. Emlen, L. W. Oring, Ecology, sexual selection, and the evolution of mating systems. *Science* **197**, 215-223 (1977).

4. T. Clutton-Brock, E. Huchard, Social competition and its consequences in female mammals. *Journal of Zoology* **289**, 151-171 (2013).

5. P. Stockley, J. Bro-Jørgensen, Female competition and its evolutionary consequences in mammals. *Biological Reviews* **86**, 341-366 (2011).

6. A. Baniel, G. Cowlishaw, E. Huchard, Jealous females? Female competition and reproductive suppression in a wild promiscuous primate. *Proceedings of the Royal Society B: Biological Sciences* **285**, 20181332 (2018).

7. D. Lukas, E. Huchard, The evolution of infanticide by females in mammals. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**, 20180075 (2019).

590   8.   K. A. Rosvall, Intrasexual competition in females: evidence for sexual selection?
591   *Behavioral Ecology* **22**, 1131-1140 (2011).

592   9.   K. Summers, Sexual selection and intra-female competition in the green poison-dart frog,
593   *Dendrobates auratus*. *Animal Behaviour* **37**, 797-805 (1989).

594   10.   M. M. F. Lutnesky, R. K. Kosaki, Female-female competition in a coral reef fish and a
595   test of the temporal threshold model of polygynous mating. *The American Naturalist* **146**,
596   832-847 (1995).

597   11.   A. Pusey, J. Williams, J. Goodall, The influence of dominance rank on the reproductive
598   success of female chimpanzees. *Science* **277**, 828-831 (1997).

599   12.   D. von Holst *et al.*, Social rank, fecundity and lifetime reproductive success in wild
600   European rabbits (*Oryctolagus cuniculus*). *Behavioral Ecology and Sociobiology* **51**,
601   245-254 (2002).

602   13.   K. E. Holekamp, L. Smale, M. Szykman, Rank and reproduction in the female spotted
603   hyaena. *Journal of Reproduction and Fertility* **108**, 229-237 (1996).

604   14.   J. Altmann, S. C. Alberts, "Intraspecific variability in fertility and offspring survival in a
605   nonhuman primate: behavioral control of ecological and social sources" in Offspring: the
606   biodemography of fertility and family behavior*,* K. W. Wachter, R. A. Bulatao, Eds.
607   (National Academy Press, Washington, DC, 2003), pp. 140-169.

608   15.   K. A. Rosvall, Sexual selection on aggressiveness in females: evidence from an
609   experimental test with tree swallows. *Animal Behaviour* **75**, 1603-1610 (2008).

610   16.   D. L. Sinn, G. M. While, E. Wapstra, Maternal care in a social lizard: links between
611   female aggression and offspring fitness. *Animal Behaviour* **76**, 1249-1257 (2008).

612   17.   C. Schradin, B. König, N. Pillay, Reproductive competition favours solitary living while
613   ecological constraints impose group-living in African striped mice. *Journal of Animal
614   Ecology* **79**, 515-521 (2010).

615   18.   J. B. Silk, The adaptive value of sociality in mammalian groups. *Philosophical
616   Transactions of the Royal Society B: Biological Sciences* **362**, 539-559 (2007).

617   19.   E. H. M. Sterck, D. P. Watts, C. P. Van Schaik, The evolution of female social
618   relationships in nonhuman primates. *Behavioral Ecology and Sociobiology* **41**, 291-309
619   (1997).

620   20.   A. Koenig, Competition for resources and its behavioral consequences among female
621   primates. *International Journal of Primatology* **23**, 759-783 (2002).

622   21.   B. C. Wheeler, C. J. Scarry, A. Koenig, Rates of agonism among female primates: a
623   cross-taxon perspective. *Behavioral Ecology* **24**, 1369-1380 (2013).

624   22.   D. T. Blumstein, C. S. Evans, J. C. Daniel, An experimental study of behavioural group
625   size effects in tammar wallabies, *Macropus eugenii*. *Animal Behaviour* **58**, 351-360
626   (1999).

627   23.   J. D. Goss-Custard, Competition for food and interference among waders. *Ardea* **55**, 31-
628   52 (1980).

629   24.   C. P. Van Schaik, M. A. Van Noordwijk, R. J. de Boer, I. den Tonkelaar, The effect of
630   group size on time budgets and social behaviour in wild long-tailed macaques (*Macaca
631   fascicularis*). *Behavioral Ecology and Sociobiology* **13**, 173-181 (1983).

632   25.   R. Slotow, Aggression in white-crowned sparrows: effects of distance from cover and
633   group size. *The Condor* **98**, 245-252 (1996).

26. C. A. Johnson, J. W. A. Grant, L.-A. Giraldeau, Thhe effect of patch size and competitor
number on aggression among foraging house sparrows. *Behavioral Ecology* **15**, 412-418
(2004).
27. S. Creel, N. M. Creel, Opposing effects of group size on reproduction and survival in
African wild dogs. *Behavioral Ecology* **26**, 1414-1422 (2015).
28. T. H. Clutton-Brock, S. D. Albon, F. E. Guinness, Competition between female relatives
in a matrilocal mammal. *Nature* **300**, 178-180 (1982).
29. C. Borries, E. Larney, A. Lu, K. Ossi, A. Koenig, Costs of group size: lower
developmental and reproductive rates in larger groups of leaf monkeys. *Behavioral*
Ecology **19**, 1186-1191 (2008).
30. D. L. Cram, J. D. Blount, A. J. Young, The oxidative costs of reproduction are group-size
dependent in a wild cooperative breeder. *Proceedings of the Royal Society B* **282**,
20152031 (2015).
31. H. Kokko, R. A. Johnstone, T. H. Clutton-Brock, The evolution of cooperative breeding
through group augmentation. *Proceedings of the Royal Society B* **268**, 187-196 (2001).
32. A. C. Markham, S. C. Alberts, J. Altmann, Intergroup conflict: ecological predictors of
winning and consequences of defeat in a wild primate population. *Animal Behaviour* **84**,
399-403 (2012).
33. S. Lemoine *et al.*, Group dominance increases territory size and reduces neighbor
pressure in wild chimpanzees. *Royal Society Open Science* **7** (2020).
34. A. C. Markham, L. R. Gesquiere, Costs and benefits of group living in primates: an
energetic perspective. *Philosophical Transactions of the Royal Society B* **372**, 20160239
(2017).
35. B. T. Hirsch, Within-group spatial position in ring-tailed coatis: balancing predation,
feeding competition, and social competition. *Behavioral Ecology and Sociobiology* **65**,
391-399 (2011).
36. F. H. Bronson, Mammalian reproduction: an ecological perspective. *Biology of*
Reproduction **32**, 1-26 (1985).
37. M. M. Wallen, E. Krzyszczyk, J. Mann, Mating in a bisexually philopatric society:
bottlenose dolphin females associate with adult males but not adult sons during estrous.
Behavioral Ecology and Sociobiology **71**, 153 (2017).
38. Q. A. Gibson, J. Mann, The size, composition and function of wild bottlenose dolphin
(*Tursiops* sp.) mother-calf groups in Shark Bay, Australia. *Animal Behaviour* **76**, 389-405
(2008).
39. L. Möller, R. G. Harcourt, Shared reproductive state enhances female associations in
dolphins. *Research Letters in Ecology*, 1-5 (2008).
40. J. Bro-Jørgensen, Overt female mate competition and preference for central males in a
lekking antelope. *Proceedings of the National Academy of Sciences* **99**, 9290-9293
(2002).
41. N. Bebié, A. G. McElligott, Female aggression in red deer: Does it indicate competition
for mates? *Mammalian Biology* **71**, 347-355 (2006).
42. D. M. Powell, Female-female competition or male mate choice? Patterns of courtship and
breeding behavior among feral horses (*Equus caballus*) on Assateague Island. *Journal of*
Ethology **26**, 137-144 (2008).

43. X.-G. Qi *et al.*, Sexual interference in the golden snub-nosed monkey (Rhinopithecus
roxellana): A test of the sexual competition hypothesis in a polygynous species.
American Journal of Primatology **73**, 366-377 (2011).
44. A. Baniel, G. Cowlishaw, E. Huchard, Context dependence of female reproductive
competition in wild chacma baboons. *Animal Behaviour* **139**, 37-49 (2018).
45. J. Dezeure *et al.*, Evolutionary determinants of nonseasonal breeding in wild chacma
baboons. *The American Naturalist* **201**, 106-124 (2023).
46. A. R. E. Sinclair, S. A. R. Mduma, P. Arcese, What determines phenology and synchrony
of ungulate breeding in Serengeti? *Ecology* **81**, 2100-2111 (2000).
47. J. Fischer *et al.*, Insights into the evolution of social systems and species from baboon
studies. *eLife* **8**, e50989 (2019).
48. G. Hausfater, J. Altmann, S. Altmann, Long-term consistency of dominance relations
among female baboons (*Papio cynocephalus*). *Science* **217**, 752-755 (1982).
49. J. A. Johnson, Dominance rank in juvenile olive baboons, *Papio anubis*: the influence of
gender, size, maternal rank and orphaning. *Animal Behaviour* **35**, 1694-1708 (1987).
50. J. B. Silk, D. L. Cheney, R. M. Seyfarth, The structure of social relationships among
female savanna baboons in Moremi Reserve, Botswana. *Behaviour* **136**, 679-703 (1999).
51. S. K. Patterson, S. C. Strum, J. B. Silk, Resource competition shapes female–female
aggression in olive baboons, *Papio anubis*. *Animal Behaviour* **176**, 23-41 (2021).
52. J. B. Silk, Correlates of agonistic and competitive interactions in pregnant baboons.
American Journal of Primatology **12**, 479-495 (1987).
53. R. A. Barton, A. Whiten, Feeding competition among female olive baboons, *Papio
anubis*. *Animal Behaviour* **46**, 777-789 (1993).
54. E. A. Archie, J. Tung, M. Clark, J. Altmann, S. C. Alberts, Social affiliation matters: both
same-sex and opposite-sex relationships predict survival in wild female baboons.
Proceedings of the Royal Society B: Biological Sciences **281**, 20141261 (2014).
55. N. Nguyen, R. C. Van Horn, S. C. Alberts, J. Altmann, "Friendships" between new
mothers and adult males: adaptive benefits and determinants in wild baboons (*Papio
cynocephalus*). *Behavioral Ecology and Sociobiology* **63**, 1331-1344 (2009).
56. E. A. Archie, J. Altmann, S. C. Alberts, Costs of reproduction in a long-lived female
primate: injury risk and wound healing. *Behavioral Ecology and Sociobiology* **68**, 1183-
1193 (2014).
57. S. K. Wasser, A. K. Starling, Proximate and ultimate causes of reproductive suppression
among female yellow baboons at Mikumi National Park, Tanzania. *American Journal of
Primatology* **16**, 97-121 (1988).
58. R. J. Rhine, S. K. Wasser, G. W. Norton, Eight-year study of social and ecological
correlates of mortality among immature baboons of Mikumi National Park, Tanzania.
American Journal of Primatology **16**, 199-212 (1988).
59. S. C. Alberts, J. Altmann, Immigration and hybridization patterns of yellow and anubis
baboons in and around Amboseli, Kenya. *American Journal of Primatology* **53**, 139-154
(2001).
60. T. P. Vilgalys *et al.*, Selection against admixture and gene regulatory divergence in a
long-term primate field study. *Science* **377**, 635-641 (2022).
61. S. C. Alberts, J. Altmann, "The Amboseli Baboon Research Project: 40 years of
continuity and change" in Long-term field studies of primates, P. Kappeler, D. P. Watts,
Eds. (Springer, 2012), pp. 261-287.

62.   J. Altmann, *Baboon mothers and infants* (Harvard University Press, Cambridge, 1980).

63.   S. Kleindorfer, S. K. Wasser, Infant handling and mortality in yellow baboons (*Papio*
cynocephalus): evidence for female reproductive competition? *Behavioral Ecology and*
Sociobiology **56**, 328-337 (2004).

64.   C. Carbone, G. Cowlishaw, N. J. B. Isaac, J. M. Rowcliffe, How far do animals go?
Determinants of day range in mammals. *The American Naturalist* **165**, 290-297 (2005).

65.   A. Baniel, G. Cowlishaw, E. Huchard, Stability and strength of male-female associations
in a promiscuous primate society. *Behavioral Ecology and Sociobiology* **70**, 761-775
(2016).

66.   L. R. Gesquiere, J. Altmann, E. A. Archie, S. C. Alberts, Interbirth intervals in wild
baboons: Environmental predictors and hormonal correlates. *American Journal of*
Physical Anthropology **166**, 107-126 (2018).

67.   S. C. Alberts *et al.*, "Seasonality and long term change in a savannah environment" in
Primate seasonality: Implications for human evolution*, D. K. Brockman, C. P. van
Schaik, Eds. (Cambridge University Press, Cambridge, 2005), pp. 157-196.

68.   R Core Team (2020) R: A language and environment for statistical computing. (Vienna,
Austria).

69.   K. Bartoń (2018) Package 'MuMIn'. in *Model selection and model averaging based on*
information criteria (AICc and alike). R package, p R package.

70.   K. P. Burnham, D. R. Anderson, *Model selection and multimodel inference: A practical*
information-theoretic approach. (Springer-Verlag, New York, 1998).

71.   K. P. Burnham, D. R. Anderson, K. P. Huyvaert, AIC model selection and multimodel
inference in behavioral ecology: some background, observations, and comparisons.
Behavioral Ecology and Sociobiology **65**, 23-35 (2011).

72.   T. Therneau (2022) A package for survival analysis in R.

73.   M. E. Brooks *et al.*, *glmmTMB* balances speed and flexibility among packages for zero-
inflated generalized linear mixed modeling. *The R Journal* **9**, 378-400 (2017).

74.   L. Digby, "Infanticide by female mammals: implications for the evolution of social
systems" in Infanticide by males and its implications*, C. P. van Schaik, C. H. Janson,
Eds. (Cambridge University Press, Cambridge, UK, 2000), pp. 178-197.

Tables & Figures

Table 1. GLMM results for the models predicting monthly agonistic interactions observed within dyads of cycling females (n = 23,547 dyad-months), pregnant females (n = 18,677 dyad-months), and females in postpartum amenorrhea (n = 35,975 dyad-months). Dyad ID and group ID were included as random effects in all three models. The results for cycling dyads and pregnant dyads reflect model averaging of candidate models that were within 2 AIC_c units of the top candidate model, but no other candidate models were within 2 AIC_c units of the top model for postpartum amenorrhea dyads (see Tables S17-19 for the top 20 candidate models for all three reproductive states). Significant fixed effects are indicated in bold. Model predictions are presented to aid interpretation of estimates and were calculated for the extreme values[1] of each significant fixed effect, while holding values for other fixed effects constant at their observed median value.

Dyad type	Fixed effect	Estimate	Std. Error	95 % CI	Model predictions
Cycling – cycling	(intercept)	-5.59	0.16	[-5.91, -5.27]	–
	Season (dry vs. wet)	**-0.10**	**0.03**	**[-0.17, -0.03]**	Dyads of cycling females exchanged 0.22 observed agonistic interactions per month in the dry season vs. 0.20 observed agonistic interactions per month in the wet season
	Rank difference	**0.55**	**0.09**	**[0.37, 0.72]**	Dyads of cycling females exchanged 0.18 observed agonistic interactions per month when they had the least rank difference vs. 0.31 observed agonistic interactions per month when they had the greatest rank difference
	Early lactational synchrony	0.18	0.24	[-0.28, 0.64]	–
	Observer effort	**0.54**	**0.04**	**[0.47, 0.61]**	Dyads of cycling females exchanged 0.14 observed agonistic interactions per month when observer effort was least intense vs. 4.06 observed agonistic interactions when observer effort was most intense
Pregnant – pregnant	(intercept)	-5.30	0.17	[-5.62, -4.98]	–
	Season (dry vs. wet)	-0.02	0.04	[-0.10, 0.05]	–
	Rank difference	**0.22**	**0.10**	**[0.04, 0.41]**	Dyads of pregnant females exchanged 0.20 observed agonsistic interactions per month when they had the least rank difference vs.

					0.25 observed agonistic interactions per month when they had the greatest rank difference
	Early lactational synchrony	1.02	0.24	[0.55, 1.48]	Dyads of pregnant females exchanged 0.20 observed agonistic interactions per month when early lactational synchrony was the lowest vs. 0.32 observed agonistic interactions per month when early lactational synchrony was the highest
	Observer effort	0.29	0.04	[0.21, 0.36]	Dyads of pregnant females exchanged 0.18 observed agonistic interactions per month when observer effort was least intense vs. 0.94 observed agonistic interactions when observer effort was most intense
Postpartum amenorrhea – postpartum amenorrhea	(intercept)	-5.41	0.13	[-5.66, -5.16]	–
	Season (dry vs. wet)	-0.19	0.02	[-0.23, -0.14]	Dyads of PPA females exchanged 0.23 observed agonistic interactions per month in the dry season vs. 0.19 observed agonistic interactions per month in the wet season
	Rank difference	0.55	0.07	[0.41, 0.70]	Dyads of PPA females exchanged 0.19 observed agonistic interactions per month when they had the least rank difference vs. 0.33 observed agonistic interactions per month when they had the greatest rank difference
	Early lactational synchrony	0.66	0.13	[0.39, 0.92]	Dyads of PPA females exchanged 0.21 observed agonistic interactions per month when early lactational synchrony was the lowest vs. 0.29 observed agonistic interactions per month when early lactational synchrony was the highest
	Observer effort	0.40	0.03	[0.35, 0.45]	Dyads of PPA females exchanged 0.18 observed agonistic interactions per month when observer effort was least intense vs. 1.77 observed agonistic interactions when observer effort was most intense

767 [1]The minimum and maximum values for each significant fixed effect that we observed in our dataset were: i) season = [dry, wet] (binary), ii) rank difference =

768 [0.03, 1.00], iii) mean early lactational synchrony = [0, 0.47], iv) observer effort = [0.03, 6.27].

Table 2. Model-averaged estimates for the best-fitting models ($\Delta AIC_c < 2$) predicting the daily risk of all-cause infant mortality up to one year after birth (Cox proportional hazards model, total infants = 1,323; died = 300, censored = 87; top 20 candidate models shown in Table S20). Significant covariates are indicated in bold. Model predictions are presented to aid interpretation of estimates and were calculated for the extreme observed values[1] for each significant predictor variable for a 10-day-old infant, while holding values for other fixed effects constant at their observed median value.

Fixed effects	Estimate (hazard ratio)[2]	Std. Error	95 % CI	Model predictions
Maternal death	**5.47 (236.51)**	**0.16**	**[5.14, 5.79]**	Infants had a 0.008% chance of dying on a day when their mother was alive vs. a 1.963% chance on a day when she was dead
Early lactational synchrony	**4.45 (85.63)**	**0.51**	**[3.46, 5.44]**	Infants had a 0.005% chance of dying on a day when early lactational synchrony was lowest vs. a 0.122% chance on a day when it was highest
Maternal age	**-0.42 (0.65)**	**0.07**	**[-0.57, -0.28]**	Infants had a 0.008% chance of dying on a day when their mother was the median age vs. a 0.053% chance on a day when she was the oldest age
Maternal age²	**0.016 (1.02)**	**0.003**	**[0.011, 0.021]**	Infants had a 0.008% chance of dying on a day when their mother was the median age vs. a 0.024% chance on a day when she was the youngest age
Maternal rank	**-0.37 (0.69)**	**0.18**	**[-0.72, -0.02]**	Infants had a 0.010% chance of dying on a day when their mother was the lowest-ranking female vs. a 0.007% chance of dying on a day when she was the highest-ranking female
Group size	-0.004 (1.00)	0.003	[-0.010, 0.003]	–
Cumulative rainfall	-0.00009 (1.00)	0.00035	[-0.00078, 0.00061]	–

[1]The minimum and maximum values for each significant predictor variable that we observed in our dataset were: i) maternal death = [mother alive, mother dead] (binary), ii) early lactational synchrony = [0.00, 0.71], iii) maternal age = [4.75, 24.50], iv) maternal rank= [0.00, 1.00].

[2]A positive coefficient or a hazard ratio >1 means that higher values of the predictor (or the presence of the predictor in the case of maternal death) are associated with higher mortality hazard (lower infant survival), and negative coefficients or a hazard ratio <1 mean that higher values of the predictor are associate with lower mortality hazard (higher infant survival).

786
787
Figure 1. (A) Females in all reproductive states initiated agonistic interactions with other adult
females at higher rates when early lactational synchrony (ELS) was high, but (B) only females in
postpartum amenorrhea received agonistic interactions from other adult females at higher rates
when early lactational synchrony was high. Red diamonds depict the β coefficient estimates for
the fixed effect of mean early lactational synchrony from negative binomial generalized linear
mixed models predicting the relative frequencies of agonistic interactions (A) initiated by all
adult females, cycling females, pregnant females, and females in postpartum amenorrhea (PPA
females) and (B) received by all adult females, cycling females, pregnant females, and PPA
females. Horizontal black bars indicate the 95% confidence intervals of these estimates. We
interpret effects as significant if their 95% confidence intervals do not overlap zero (indicated by
the vertical dotted line). Note that while 'early lactational synchrony' on the x-axis labels refers
to the proportion of adult females in a group who had a live infant <90 days old, the term 'PPA
females' on the y-axes refers to all females in postpartum amenorrhea, not only those with live
infants <90 days old (see Methods).
802

Figure 2. The relationship between early lactational synchrony (ELS), infant mortality, and infanticide. A) The distribution of early lactational synchrony across the study period and in subsets of days when infants died or were killed by infanticide. The y-axis shows the daily proportion of adult females in early lactation as a function of different partitions of our data on the x-axis: all group-days between 1976 and 2021, group-days on which infants less than 1 year old died, group-days on which infants less than 1 year old were killed by a groupmate, group-days on which infants less than 1 year old were killed by a female groupmate, and group-days on which infants less than 1 year old were killed by a male groupmate. Violin plots depict probability densities of the data on each category of group-day, while the overlaid boxplots depict the median values and interquartile ranges. Infanticides were only included in counts for this figure if they were directly observed or strongly suspected with corroborating evidence (see Methods). B) Lactational synchrony predicts infant mortality risk. Monthly mortality rates per 1000 infants are depicted with values calculated across all infants (green line), for infants in groups-months in which mean early lactational synchrony was in the highest quartile relative to

other infants in the same month of life (orange line), and for infants in groups-months in which
mean early lactational synchrony was in the highest decile relative to other infants in the same
month of life (purple line). Early lactational synchrony was modeled as a continuous, daily, time-
varying variable in our survival analysis; monthly mean values and quartiles are used for
visualization purposes. C) The distribution of early lactational synchrony is higher on days when
infants are killed by females than on days when infants die by any cause. This quantile-quantile
plot compares the distribution of early lactational synchrony on female infanticide days with the
distribution of early lactational synchrony across all infant death days. The solid, black line
depicts the expected relationship if female infanticide was equally likely on all infant death days.
Almost all points fall above the line, indicating that days on which infants are killed by females
tend to have higher early lactational synchrony relative to days on which infants die from any
cause.
832
833