A guanidine-based coronavirus replication inhibitor which targets the nsp15 endoribonuclease and combines a direct antiviral effect with subtle boosting of interferon

Benjamin Van Loy[1], Eugènia Pujol[2,3], Kenichi Kamata[4#], Xiao Yin Lee[4], Nikolai Bakirtzoglou[1], Ria Van Berwaer[1], Julie Vandeput[1], Leentje Persoons[1], Brent De Wijngaert[1], Quinten Goovaerts[1], Sam Noppen[1], Kourosh Ahmadzadeh[1], Juan Martín-López[2,3], Celia Escriche[2,3], Bert Vanmechelen[1], Besir Krasniqi[5†], Abhimanyu K. Singh[1‡], Piet Maes[1], Patrick Matthys[1], Wim Dehaen[5], Jef Rozenski[6], Kalyan Das[1], Arnout Voet[4], Santiago Vázquez[2,3], Lieve Naesens[1*] and Annelies Stevaert[1]

[1]Rega Institute, Department of Microbiology, Immunology and Transplantation, KU Leuven, Leuven, Belgium

[2]Laboratori de Química Farmacèutica (Unitat Associada al Consejo Superior de Investigaciones Científicas), Facultat de Farmàcia i Ciències de l'Alimentació, Universitat de Barcelona, Barcelona, Spain

[3]Institute of Biomedicine of the University of Barcelona (IBUB), Universitat de Barcelona, Barcelona, Spain

[4]Biochemistry, Molecular and Structural Biology, Department of Chemistry, KU Leuven, Leuven, Belgium

[5]Sustainable Chemistry for Metals and Molecules, Department of Chemistry, KU Leuven, Leuven, Belgium

[6]Rega Institute, Department of Pharmaceutical and Pharmacological Sciences, KU Leuven, Leuven, Belgium

[#]Current address: Yokohama City University, Yokohama, Japan

[†]Current address: Vlaamse Instelling voor Technologisch Onderzoek (VITO), Mol, Belgium

[‡]Current address: Mahatma Gandhi Medical Advanced Research Institute, Sri Balaji Vidyapeeth, Puducherry, India

*Corresponding author

E-mail: lieve.naesens@kuleuven.be (LN)

Short title

Coronavirus replication inhibitor targeting the nsp15 endoribonuclease

Abstract

The approval of COVID-19 vaccines and two classes of antiviral drugs has been crucial in addressing the global health crisis caused by SARS-CoV-2. However, to prepare for future outbreaks from drug-resistant variants and novel zoonotic coronaviruses (CoVs), additional therapeutics with a distinct antiviral mechanism are needed. Here, we report a novel guanidine-substituted diphenylurea compound that suppresses CoV replication by interfering with the uridine-specific endoribonuclease (EndoU) activity of the viral non-structural protein-15 (nsp15). This compound, designated EPB-113, exhibits strong and selective cell culture activity against human coronavirus 229E (HCoV-229E) and also suppresses the replication of SARS-CoV-2. HCoV-229E viruses selected for resistance to EPB-113 carried mutations at or near the catalytic residue His250 in the nsp15 EndoU domain. EndoU is known to reduce the levels of viral dsRNA and avoid induction of type I interferon (IFN-I). Accordingly, the mutant viruses selected under EPB-113 were unable to replicate in human macrophages and were readily outcompeted by the wild-type virus upon co-infection of human fibroblast cells. This highlights the critical role of EndoU in maintaining CoV replication. By targeting nsp15, EPB-113 combines a direct antiviral effect with subtle boosting of IFN-I. Using a combination of biophysical and enzymatic assays with the recombinant nsp15 proteins from HCoV-229E and SARS-CoV-2, we discovered that EPB-113 reduces the thermal stability of hexameric nsp15, while enhancing its EndoU cleavage activity. This EndoU-stimulating mechanism offers an explanation why the virus escapes EPB-113 by becoming EndoU-deficient. Our findings suggest that targeting nsp15 as an antiviral strategy may require a mechanism that alters the conformation of this protein and disrupts its EndoU activity and other functions related to immune evasion and viral replication. Based on the appealing pharmacological profile of EPB-113, we conclude that nsp15 is a challenging but highly relevant drug target.

Author summary

Despite significant progress in controlling the global COVID-19 crisis through vaccines and antiviral drugs, SARS-CoV-2 infection remains a serious health threat for vulnerable individuals, such as elderly and immunocompromised patients. These populations could benefit from innovative treatments that target the virus from multiple angles. In this study, we focus on the nsp15 endoribonuclease (EndoU), a poorly understood coronavirus protein that aids in viral evasion from the host innate immune response and is assumed to regulate viral RNA synthesis. Our research centers around a newly identified, first-in-class coronavirus inhibitor, designated EPB-113, which specifically targets nsp15. This compound has the unique property to suppress the virus in two complementary ways via subtle induction of interferon combined with a direct impact on viral RNA synthesis. EPB-113 selects for escape mutants that carry substitutions in the core of EndoU, leading to a severe replication defect. When incubated with nsp15 protein, EPB-113 induces a structural change and enhances its EndoU enzyme activity. Hence, our study introduces a novel drug concept that disrupts the multifunctional nsp15 protein and suppresses coronavirus replication on multiple fronts.

Introduction

In the five years since COVID-19 was first identified, Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has caused an estimated 776 million human infections and 7 million deaths worldwide [1]. The fast roll-out of innovative vaccines was crucial in addressing the global health crisis when SARS-CoV-2 suddenly emerged in the human population. Today, annual COVID-19 vaccination remains recommended, as the vaccines provide only short-term immunity and the virus continues to undergo antigenic drift. Serious complications from COVID-19 are common in vulnerable individuals, such as the elderly or immunocompromised patients. For those with mild-to-moderate COVID-19 who are at high risk for progression to severe disease, antiviral medication is recommended, specifically ritonavir-boosted nirmatrelvir [which targets the coronavirus main protease (M^{pro})] or remdesivir [a nucleotide inhibitor of the viral RNA-dependent RNA polymerase (RdRp)] [2].

To prepare for future outbreaks caused by drug-resistant variants or novel zoonotic coronaviruses (CoVs), exploring alternative antiviral drug targets is essential. Many ongoing efforts are focused on one of the 16 non-structural proteins (nsps) that make up the viral RNA transcription-replication complex (RTC) [3-5]. Several nsps help the virus to evade the host innate immune response that aims to halt viral replication [6]. This dual function also applies to nsp15, a uridine-specific endoribonuclease (EndoU) that is highly conserved among all CoVs [7, 8]. As part of the RTC, nsp15 is believed to cleave specific sites in viral RNA to regulate viral RNA synthesis [9]. Additionally, EndoU prevents the accumulation of viral dsRNA intermediates, thereby avoiding their detection by dsRNA sensors of the innate immune response [10-12]. This may be accomplished by EndoU cleavage of the 5´-poly(U) extension in viral negative-sense RNA [12] or the CACA sequence near the 3´-poly(A) tail of the positive strand [9]. Evidence also suggests that nsp15 interacts with cellular proteins involved in the production of type I interferon (IFN-I) [13, 14] and may protect infected cells against apoptosis [10, 15], autophagy [16], cell cycle arrest [17] or the formation of stress granules [18-20].

The immune-evading activity of EndoU has been demonstrated in cell culture studies with different CoVs [10, 11, 21, 22], including human nasal epithelial cells infected with SARS-CoV-2 [23]. EndoU-deficient mutants of mouse hepatitis virus (MHV) and porcine epidemic diarrhea virus (PEDV) proved significantly less pathogenic than their wild-type (WT) counterparts in mice and piglets, respectively [10, 21].

However, an EndoU-deficient mutant of transmissible gastroenteritis virus exhibited comparable

virulence to WT in piglets, while the analogous mutant of feline infectious peritonitis virus was more

virulent than WT in cats [24]. Hence, many questions remain about the functions of EndoU in viral

replication and immune evasion.

The potential of nsp15 as a novel drug target has spurred recent efforts to identify inhibitors in enzymatic

EndoU assays [25-30] and resolve the protein structure of nsp15 in complex with a mono- or

oligonucleotide ligand [31-35]. For nearly all CoVs, including SARS-CoV-2, nsp15 adopts a

characteristic hexameric structure consisting of a dimer of trimers, with the six EndoU domains facing

outwards of the complex [36]. Each monomer contains an N-terminal domain (NTD), middle domain and

C-terminal domain harboring the EndoU site. Nsp15 hexamerization is regulated by the NTD and crucial

for EndoU enzymatic activity [37]. A recent review covers the biochemical properties, structure and

assumed functions of nsp15 in greater detail and can be found elsewhere [38].

Our interest in developing nsp15 inhibitors started several years ago, following the identification of a

betulonic acid-based inhibitor (encoded 5h in reference [39]) that exhibits submicromolar activity against

human coronavirus 229E (HCoV-229E), low cytotoxicity and a mechanism related to nsp15. The finding

that viral resistance to this compound was mapped to an inter-protomer region in the hexameric nsp15

protein of HCoV-229E, allowed us to create a docking model and predict the binding mode of 5h [39].

Shortly after, we identified compound EPB-113, another inhibitor of HCoV-229E that is structurally

unrelated to 5h. Intrigued by the discovery that viral resistance to EPB-113 was linked to mutations in

the catalytic center of nsp15 EndoU, we conducted the present study to uncover the mechanism of this

antiviral molecule. Through cell-based assays, we explored the association between EndoU and the

innate immune response, as well as other presumed functions of nsp15 beyond EndoU. We also

thoroughly profiled EPB-113-resistant nsp15-mutant viruses. Furthermore, biochemical assays with

recombinant nsp15 protein revealed that EPB-113 enhances the EndoU activity, which was opposite to

our initial expectations. Taking together, our findings suggest that EPB-113 acts through a previously

unrecognized pharmacological mechanism, by binding to nsp15, altering its conformation and disrupting

both its EndoU activity and other functions in viral replication and immune evasion.

Materials and methods

Compounds

In S1 Appendix, we provide all details on the chemical synthesis, analysis and full characterization of diphenylurea compounds EPB-113, EPB-102 and JMLV-061. The synthesis of 5h, a closely related 1,2,3-triazolo-fused betulonic acid derivatives, was described previously [39].

The following compounds were purchased from a commercial supplier: GS-441524 (Carbosynth); E64d (MedChemExpress); nirmatrelvir (Chemat); and K22 (ChemDiv). Recombinant human IFN-β was purchased from PeproTech.

Cells and media

Human embryonic lung fibroblast cells [HEL299; American Type Culture Collection (ATCC) No. CCL-137] were cultured in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal calf serum (FCS); 10 mM HEPES; 1 mM sodium pyruvate; 6 mM L-glutamine; 0.1 mM non-essential amino acids (NEAA); and 25 mM glucose. For experiments, they were seeded on day -7 in 96-well plates and grown until confluent, using DMEM with 2% FCS; 0.075% sodium bicarbonate; 1 mM sodium pyruvate; 0.1 mM NEAA; and pen-strep (= 100 U/ml penicillin and 0.1 mg/ml streptomycin). Human bronchial epithelial 16HBE cells (kind gift from P. Hoet, KU Leuven) and BHK-21J cells (donated by P. J. Bredenbeek) were maintained in DMEM with 10% FCS; 1 mM sodium pyruvate; 0.075% sodium bicarbonate; 0.1 mM NEAA; 4 mM L-glutamine; and 25 mM glucose. For experiments, 16HBE cells were seeded on day -1 in 96-well plates (30,000 cells per well), using the same medium but with 2% FCS and pen-strep. A549 cells expressing ACE2 and TMPRSS2 (A549$^{ACE2/TMPRSS2}$) were purchased from InvivoGen (No. a549d-cov2r) and cultured in DMEM supplemented with 6 mM L-glutamine; 25 mM glucose; 10% FCS; 100 µg/ml hygromycin and 0.5 µg/ml puromycin. HEK-Blue IFN-α/β cells (kindly donated by X. Saelens, UGent) were grown in DMEM with 4 mM L-glutamine; 25 mM glucose; 10% heat-inactivated FCS; pen-strep; 30 µg/ml blasticidin; and 100 µg/ml zeocin.

Human macrophages were purified from healthy donor buffy coats, obtained from the Red Cross Belgium according to KU Leuven agreement RKOV_19006. After isolating the mononuclear cell fraction by density gradient centrifugation on Pancoll (PAN-Biotech), the monocytes were purified by adherence

for 1.5 h at 37 °C, followed by three PBS washes to remove non-adherent cells. The cells were seeded

in 96-well plates (90,000 cells per well) and allowed to differentiate into macrophages during six days

incubation in RPMI1640 medium with 2 mM glutamine and supplemented with 10% heat-inactivated

FCS; 2 mM L-alanyl-glutamine; 20 ng/ml macrophage-colony stimulating factor (M-CSF, PeproTech);

and pen-strep. Fresh medium was provided on day 3. For infection experiments, we used the same

medium but with 2% FCS.

All cells were incubated in a humidified 5% CO_2 incubator at 37 °C.

Viruses

HCoV-229E (ATCC No. VR-740) was expanded in HEL299 cells. Reverse-engineered HCoV-229E virus

carrying substitution $H250A_{nsp15}$ was kindly provided by V. Thiel (University of Bern). The SARS-CoV-2

viruses were created by reverse-engineering (see below). The stocks of wild-type and nsp15-mutant

viruses (see below) were titrated in HEL299 (for HCoV-229E) or $A549^{ACE2/TMPRSS2}$ cells (for SARS-CoV-

2) and the 50% cell culture infective dose ($CCID_{50}$) was calculated by the method of Reed and Muench.

All infection experiments were done at 35 °C for HCoV-229E and 37 °C for SARS-CoV-2.

Antiviral and cytotoxicity assays

Unless specified differently, the compounds' antiviral activity was determined in HEL299 cells infected

with HCoV-229E or in $A549^{ACE2/TMPRSS2}$ cells infected with SARS-CoV-2, both at a multiplicity of infection

(MOI) of 100 $CCID_{50}$. Serial dilutions of the compounds were added to the cells in 96-well plates,

immediately followed by the virus. A mock-infected plate was prepared in parallel, to assess compound

cytotoxicity. To study combinations, different concentrations of the two compounds were added in a

matrix of 8x8 wells. For the CPE reduction assay, the viral CPE was scored by microscopy at day 5 p.i.,

then quantified by the colorimetric MTS cell viability assay. After replacing the supernatant by CellTiter

$96AQ_{ueous}$ MTS Reagent (Promega) and incubating the plates for 24 h (HEL299) or 6 h

($A549^{ACE2/TMPRSS2}$) at 37 °C, the A_{490} values were measured in a Tecan Spark 10M plate reader. To

calculate the EC_{50} (50% antivirally effective concentration) and CC_{50} (50% cytotoxic concentration)

values, we used the formulas specified elsewhere [40].

For virus yield reduction assays, the supernatants were harvested at day 3 to 5 p.i. To quantify the viral

load by RT-qPCR, the samples were mixed with Cells-to-cDNA II lysis buffer (Invitrogen) and incubated

for 10 min at 75 °C. The number of viral RNA genomes (= viral load) was determined by one-step RT-

qPCR (iTaq Universal Probes One-Step kit, BioRad), employing N-gene specific primers and probe (see

sequences provided in S2 Appendix) and an N-gene plasmid standard to allow absolute quantification

[41]. The RT-qPCR program consisted of: 10 min at 50 °C; 2 min at 95 °C; and 40 cycles of 15 s at 95

°C and 60 s at 60 °C. To quantify the yield of infectious virus, we incubated serial dilutions of the

supernatants with HEL299 cells in 96-well plates, scored the CPE at day 5 p.i. and calculated the $CCID_{50}$

titers as described above. Antiviral activity in these virus yield assays was expressed as the EC_{90} value,

i.e. compound concentration producing $1\text{-}log_{10}$ reduction in the number of viral RNA genomes or in the

$CCID_{50}$ titer. Compound cytotoxicity was assessed in parallel in mock-infected plates by MTS cell

viability assay at day 3 or 5 p.i.

Time-of-compound addition assay

Confluent HEL299 cells were infected with HCoV-229E and the compounds were added to separate

wells at -0.5, 0.5, 2, 4, 6, 8, 10, 12 or 14 h p.i. At 16 h p.i., the cells were washed with ice-cold PBS, then

lysed by adding Cells-to-cDNA II lysis buffer and incubating for 10 min at 75 °C. The viral RNA copy

number in each sample was determined by one-step RT-qPCR as explained above. The starting number

of viral RNA copies (= input RNA) was determined in a cell lysate (from the no compound condition)

made at 2 h p.i.

Selection of compound-resistant HCoV-229E viruses

Instead of performing serial virus passaging, we used a faster resistance selection method (that was

earlier developed for Chikungunya virus by Delang et al. [42]. First, HEL299 cells were exposed to serial

dilutions of HCoV-229E and compound (EPB-113 or JMLv-061). At day 4 p.i., the optimal condition for

resistance selection was defined as the condition receiving the highest MOI and lowest compound

concentration achieving complete prevention of CPE. Subsequently, this virus/compound condition was

applied to all wells of a 96-well plate with HEL299 cells. At day 4 p.i., the CPE was scored by microscopy

and the wells showing virus breakthrough were selected to harvest supernatants and expand the

progeny virus. For virus sequencing, we prepared RNA extracts from the expanded virus stocks

(QIAamp Viral RNA Kit; Qiagen) and performed high-fidelity RT-PCR (SuperScript IV One-Step RT-PCR

System; Invitrogen), yielding five overlapping cDNA fragments that covered the entire viral genome.

After Sanger sequencing (Macrogen Europe) with a set of 39 primers, the mutations selected under

EPB-113 or JMLv-061 were identified by aligning the sequences of compound-selected and wild-type

(WT) forms of HCoV-229E, using CLC Main Workbench (Qiagen). To conduct antiviral experiments, the

mutant viruses were plaque-purified in HEL299 cells grown in 6-well plates in the presence of 20 µM

compound and overlaid with 0.4% UltraPure Low Melting Point Agarose (Invitrogen).

Reverse-engineering on SARS-CoV-2

Starting from seven reverse genetics plasmids which were generously donated by P.-Y. Shi (World

Reference Center for Emerging Viruses and Arboviruses, Texas) [43], we first replaced their sequences

encoding SARS-CoV-2 strain SARS-CoV-2/USA_WA1/2020 by the analogous sequences of Omicron

subvariant BF.7. Therefore, viral RNA was extracted (QIAamp viral RNA kit, Qiagen) from a clinical

isolate (pango lineage BF.7; hCoV-19/Belgium/rega-43668/2022; EPI_ISL_14623620), followed by

reverse transcription (Superscript IV RT First-Strand synthesis system, Invitrogen) and PCR (Platinum

SuperFi PCR, Invitrogen). The resulting fragments were used to replace the WA1-sequences in the

seven reverse genetics plasmids (NEBuilder HiFi DNA Assembly Cloning Kit, New England Biolabs).

For generating mutants, site-directed mutagenesis was performed with overlapping mutagenic primers

and Platinum SuperFi II DNA Polymerase (Invitrogen). The procedure for reverse-engineering SARS-

CoV-2 was adapted from Xie et al [43]. The plasmids were transformed into One Shot TOP10 Chemically

Competent *E. coli* (Invitrogen; for pUC57-based plasmids) or TransforMax EPI300 *E. coli* (LGC

Biosearch Technologies; for pCC1-based plasmids) and purified with the Qiagen Plasmid Plus Maxi Kit.

All plasmids were validated by Sanger sequencing (Macrogen Europe). Next, SARS-CoV-2 DNA

fragments were recovered through restriction enzyme digestion of the seven plasmids and the

subsequent ligation step yielded full-length viral DNA. N-gene cDNA was prepared from the plasmid

containing the last part of the viral DNA via PCR (Platinum SuperFI II, Invitrogen). Using T7 polymerase

(mMESSAGE mMACHINE T7 transcription kit; Invitrogen), the full-length and N-gene viral DNA were *in*

vitro transcribed to obtain genomic and N-coding SARS-CoV-2 RNA, followed by purification by

phenol:chloroform extraction and isopropanol precipitation. Twenty µg of each was subsequently

electroporated into 8 million BHK-21J cells resuspended in PBS, using the Gene Pulser Xcell

Electroporation System (Bio-Rad). The electroporated cells were then seeded in a 75-cm² flask with

A549[ACE2-TMPRSS2] cells. The supernatants were harvested after three days, and further expanded in

A549[ACE2-TMPRSS2] cells. The resulting virus stocks were first subjected to nsp15 sequencing, with the

primers described by Xie et al. [43], to ensure that the desired mutations were present.

HCoV-229E replication kinetics and expression of innate immune factors

HEL299 cells, 16HBE cells and human macrophages were infected with WT or nsp15-mutant HCoV-229E and 2 h later, the inoculum was replaced by fresh medium. At 2, 24, 48 and 72 h p.i., supernatant samples were collected and the cells were extracted for total RNA (ReliaPrep RNA miniprep system, Promega). The supernatants were analyzed for number of viral RNA genomes, using the one-step RT-qPCR method from above, as well as for bioactive IFN-α/β (see below). To measure the mRNA levels for a selection of innate immune factors, the cellular RNA extracts were submitted to RT-qPCR (High-Capacity cDNA Reverse Transcription Kit, Applied Biosystems), using random primers for cDNA synthesis and gene-specific primers (primer sequences are listed in S3 table) for qPCR (GoTaq qPCR kit, Promega). The transcript levels for innate immune factors were normalized to the geometric mean of those for the housekeeping genes ACTB and HPRT (ΔC_T) and the fold change was calculated as $2^{-\Delta\Delta CT}$ over mock (ΔC_T target - ΔC_T mock).

The levels of bioactive IFN-α/β in the supernatants were quantified in HEK-blue IFN-α/β reporter cells, following the manufacturer's instructions. In brief, 20 µl supernatant was added to the wells of a 96-well plate and the reporter cells were added at 50,000 cells in a volume of 180 µl. Recombinant human IFN-β was included as a positive control. After overnight incubation at 37 °C, 20-µl supernatant samples were collected from the reporter cells, then combined with 180 µl QUANTI-blue solution (InvivoGen) in a 96-well plate and incubated for 2 h at 37 °C. The A_{620} values were measured in a Tecan Spark 10M plate reader. For calculations, all absorbance values were first adjusted by subtracting the blank, i.e. QUANTI-blue in the corresponding medium per cell line, then expressed as the fold increase for virus- versus mock-infected cells.

Competitive fitness of nsp15 inhibitor-resistant viruses

HEL299 cells were co-infected with WT and nsp15-mutant HCoV-229E virus, mixed at a ratio of 9:1, 1:1 or 1:9. After three days incubation at 35 °C, the supernatants were harvested, titrated and applied to fresh HEL299 cells. This procedure was repeated for an extra three passages. The supernatants of each passage were collected for RNA extraction (QIAamp Viral RNA Kit, Qiagen) followed by RT-PCR (SuperScript IV One-Step RT-PCR, Invitrogen) with primers 229E-For17743 and 229E-Rev24484

(sequences provided in S3 Table), resulting in ~6.7 kb amplicons covering the nsp15 gene. All PCR

products were sequenced using Oxford Nanopore Technologies devices. For passages 1 and 2, library

preparation was performed using the SQK-RBK110.96 kit and sequencing was done on an r9.4.1 flow

cell connected to a GridION (MinKNOW v23.04.6; super-accurate basecalling). For passages 3 and 4,

library preparation was done using the SQK-NBD114.96 kit and the resulting library was sequenced on

an r10.4.1 PromethION flow cell using a P2 Solo device connected to a GridION (MinKNOW v23.07.12;

super-accurate basecalling). Filtlong (v0.2.1) was used to subset for each barcode the 10% highest

quality reads with a length between 6.5 and 7 kb. These subsets were mapped against the HCoV-229E

reference sequence (NC_002645.1) using the minimap2 (v2.24) map-ont preset [44], resulting in a

minimal coverage of 1200X at each site. Detailed coverage statistics per position were determined using

bam-readcount (v1.0.1), after which a custom bash script was used to calculate simplified percentages

of WT and nsp-15 mutant motif prevalence at each virus passage [45].

Production of recombinant nsp15 protein from HCoV-229E and SARS-CoV-2

The pET28a(+) plasmid for bacterial expression of SARS-CoV-2 nsp15 was a kind gift from B. Canard

(Aix-Marseille University). To engineer the corresponding plasmid for HCoV-229E nsp15, the sequence

(codon-optimized for expression in *E. coli*) was supplemented with an N-terminal octahistidine tag and

TEV cleavage site and ordered as a gBlock from IDT. This DNA was cloned in the pET28a(+) vector

using the NEBuilder HiFi DNA assembly cloning kit (NEB) and corresponding instructions. To produce

the two proteins, T7 Express Competent *E. coli* cells (NEB) were transformed with pET28a(+)-nsp15

plasmid, as instructed by the supplier. A starter culture was grown overnight in 100 ml Luria–Bertani (LB)

broth with 50 µg/ml kanamycin, using a shaking incubator (100 rpm) at 37 °C. From this, 10 ml was

added to 1 liter LB medium with 50 µg/ml kanamycin, and the bacteria were grown until an OD_{600} of ~0.6

was reached. Protein expression was induced at 500 µM isopropyl β-D-1-thiogalactopyranoside, and

the cultures were incubated overnight at 18 °C, after which the cells were pelleted by centrifugation (30

min, 5000 g, 4 °C) and stored at -20 °C. For protein purification, the pellets were resuspended in lysis

buffer (50 mM Tris pH 8; 300 mM NaCl; 20 mM imidazole; and 1 mM dithiothreitol (DTT), supplemented

with 1 mM phenylmethylsulfonyl fluoride, 1% Triton-X, 30 mg lysozyme, 20 mM $MgSO_4$, 100 U DNase

I). After cell disruption by high pressure using EmulsiFlex-C3 (Avestin), 4 times at 20,000 bar, cell debris

was removed by centrifugation (40 min, 5000 x g, 4 °C) and filtration through a 0.45 μm Whatman GD/XP

Syringe Filter (Cytiva). The filtered lysates were subjected to Ni-NTA chromatography (Qiagen)

equilibrated with the lysis buffer without supplement, washed and eluted with the increasing

concentration of the imidazole (20-300 mM Imidazole). The eluted protein is then applied to a HiLoad

Superdex 16/600 200 pg size-exclusion chromatography (SEC) column (GE Healthcare) and run in a

buffer containing 25 mM HEPES at pH 8, 300 mM NaCl, 10% glycerol, and 1 mM tris(2-

carboxyethyl)phosphine. Finally, the protein solutions were concentrated with a centricon (VivaSpin 15R

centricon, 10,000 MWCO; Sartorius AG). The protein products were analyzed by dynamic light

scattering (DLS; S1 Fig), which showed that the nsp15 proteins were ≥ 98% pure and present in

hexameric form, in line with other reports [46, 47].

Biophysical evaluation of the compound-nsp15 interaction

The thermal shift assay was executed in white PCR plates. The wells contained 6 μM nsp15 (diluted in

assay buffer: 20 mM HEPES pH 7.5; 50 mM NaCl; 5 mM $MnCl_2$; and 1 mM DTT), 5x SYPRO orange

dye and either 100 μM test compound or the corresponding percentage of DMSO. Using a QuantStudio

5 Real-Time PCR system (Thermo Fisher), fluorescence was continuously monitored while the plate

was heated from 4 to 95 °C at a ramp rate of 0.05 °C/s. The nsp15 melting temperature (T_m) was

calculated by determining the temperature at which the negative first derivative of the fluorescence

versus temperature (-dFluorescence/dT) reaches its minimum, using Protein Thermal Shift Software

v1.3 (Thermo Fisher).

To assess the interaction between EPB-113 and nsp15 from SARS-CoV-2 or HCoV-229E by surface

plasmon resonance (SPR), we used a Biacore 8K instrument (Cytiva). Nsp15 was covalently coupled

onto a CM4 sensor chip (Cytiva) at a density of around 3500 resonance units (RU) using standard amine

coupling chemistry in 10 mM acetate buffer pH 5.5. A reference flow was used as a control for non-

specific binding and refractive index changes. All binding experiments were performed at 25 °C in 50

mM Tris buffer supplemented with 500 mM NaCl, 5 mM $MnCl_2$, 0.05% Tween20 and 5% DMSO. Serial

dilutions of EPB-113 or EPB-102 were injected for 120 s at a flow rate of 30 μl/min. The dissociation was

monitored for 6 min. Buffer blanks were used for double referencing and a DMSO solvent correction

was included. Biacore Insight Software 5.0.18 was used for analysis.

Enzymatic EndoU assay

For the FRET-based EndoU cleavage assay, we employed a 5-nt single-stranded DNA-RNA hybrid substrate (5′-6-FAM-AArUAA-3′-IABkFQ; synthesized by IDT), that contained a single U↓A-cleavage site, 5′-carboxyfluorescein (FAM) fluorophore and 3′-Iowa Black quencher [27]. The assay was conducted in black half-area 96-well plates, using the same assay buffer as in the thermal shift assay. The wells (25 µl total volume) contained 63 nM (HCoV-229E) or 4 nM (SARS-CoV-2) nsp15, 0.5 µM substrate and 100 µM of compound or the corresponding percentage of DMSO. Using a Tecan Spark 10M plate reader set at 485 nm emission and 535 nm excitation wavelength, the increasing fluorescence was continuously measured during 1 h at 37 °C. The initial cleavage rate (V_0) was obtained by calculating the slope (RFU/s) of the 5-10 min time interval of the cleavage reaction.

In parallel, the samples after 1 h reaction time were analyzed by HPLC on a capillary chromatography instrument (CapLC, Waters) equipped with a C18 reversed-phase column (PepMap 0.5 x 15 mm, LC Packings). The elution buffer contained 0.05% v/v *N,N*-dimethylaminobutane (DMAB, Acros) as ion pairing reagent and 1% v/v 1,1,1,3,3,3-hexafluoro-2-propanol (hexafluoroisopropanol, HFiP, Acros) in water (pH 8.0). The organic phase consisted of 50% v/v acetonitrile (Fisher Scientific) in water. Oligonucleotides were eluted at a flow rate of 12 µl per min, applying a gradient starting at 5% organic phase and increasing by 20% per min during 4 min. The mass spectrometer consisted of a quadrupole time-of-flight instrument (Synapt G2 HDMS, Waters) equipped with a standard electrospray source. Every 2 s, spectra were recorded in negative ion mode with following settings: cone voltage 35 V, capillary voltage 2200 V, source temperature 80 °C and N_2 desolvation gas 600 L/h. The acquired spectra were mass corrected using leucine enkephalin and deconvoluted applying the MaxEnt algorithm (MassLynx 4.1, Waters).

Data analysis and statistics

All graphs were created with GraphPad Prism 10.3.1 software, which was also used for statistical analysis. Statistical significance is shown as: ****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$, ns: not significant.

Results

EPB-113 is a strong replication inhibitor of HCoV-229E that also suppresses SARS-CoV-2

HCoV-229E differs from other human CoVs in causing manifest cytopathic effect (CPE) in different cell lines, while its manipulation is allowed at biosafety level 2. Based on this, we have been using since several years a CPE reduction assay in HCoV-229E-infected HEL299 cells, to conduct convenient phenotypic screening for novel CoV inhibitors. To assess the compounds' antiviral effect, the viral CPE is quantified by MTS cell viability assay in combination with microscopic examination. These two methods are also used to determine compound cytotoxicity in mock-infected cells.

Our screening campaign identified compound EPB-113 as a robust lead compound, exhibiting an EC_{50} value for HCoV-229E of 0.48 µM and CC_{50} value for HEL299 cells of 92 µM, thus yielding a favorable selectivity index (SI; ratio of CC_{50} to EC_{50}) of 193 (Figs 1B and 1D). Antiviral activity was very similar whether determined by MTS assay or by microscopic scoring (Fig 1D). The activity was confirmed when we quantified the virus in the supernatant at day 3 p.i. Here, EPB-113 showed an EC_{90} value of 0.65 µM based on RT-qPCR for viral genome copies and of 0.40 µM based on titration of infectious virus (Figs 1B and 1D). In HCoV-229E-infected HEL299 cells, EPB-113 outperformed GS-441524, the nucleoside form of remdesivir that was included as reference compound [EC_{50} value of 3.5 µM (CPE assay) and EC_{90} value of 2.5 and 4.6 µM (by infectious virus titration and RT-qPCR, respectively)].

Fig 1. Inhibitory activity of EPB-113 on the replication of HCoV-229E and SARS-CoV-2. (A) Chemical structures of EPB-113 and its structural analogues EPB-102 and JMLv-061. (B) Inhibition of HCoV-229E replication in HEL299 cells was determined by MTS-based CPE reduction assay; infectious virus titration on the supernatant; and RT-qPCR-based viral load assay. Inhibition of SARS-CoV-2 replication in A549[ACE2/TMPRSS2] cells was quantified by viral load assay. Cytotoxicity was determined by MTS cell viability assay in mock-infected cells and is shown in dashed lines (open circles). Data points are the mean ± SEM (N=3-12). (C) Anti-HCoV-229E activity and cytotoxicity in HEL299 cells for EPB-102 and JMLv-061, two structural analogues of EPB-113. Data points are the mean ± SEM (N=3). (D) Table summarizing the values for antiviral activity and cytotoxicity of EPB-113, JMLv-061, EPB-102 and reference compound GS-441524.

D

Compound	Virus	Antiviral activity				Cytotoxicity
		EC_{50} (CPE)	EC_{50} (MTS)	EC_{90} (infectious titer)	EC_{90} (viral load)	CC_{50}
EPB-113	HCoV-229E	1.0	0.48	0.40	0.65	92
	SARS-CoV-2	12	-	-	12	99
JMLv-061	HCoV-229E	0.33	0.25	-	-	> 100
EPB-102	HCoV-229E	12	7.0	-	-	48
GS-441524	HCoV-229E	3.5	2.6	2.5	4.6	> 100
	SARS-CoV-2	1.3	-	-	1.7	> 25

Chemically, EPB-113 is a diphenylurea compound featuring a *m*-guanidine function in an aromatic ring

and a *m*-pentafluorosulfanyl group in the other aromatic ring (Fig 1A). To determine the structure-activity

relationship for HCoV-229E, we synthesized a series of more than 20 structural analogues, the outcome

of which will be published in a separate report (S. Vazquez, manuscript in preparation). Of note, as the

very lipophilic *m*-pentafluorosulfanyl group is uncommon in medicinal chemistry, we also evaluated other

analogs lacking this group, such as JMLv-061 (Fig 1A), with an *o*-trifluoromethoxy group instead of the

m-pentafluorosulfanyl unit, but keeping the *m*-guanidine group on the other aromatic ring. While this

change had no impact on the antiviral activity (EC$_{50}$ for JMLv-061: 0.33 μM in HCoV-229E infected

HEL299 cells; Figs 1C and 1D), the analog lacking the guanidine function (= EPB-102; Fig 1A) was

shown to have much weaker activity (EC$_{50}$ of 12 μM in HCoV-229E infected HEL299 cells, reaching only

60% inhibition at 40 μM; Figs 1C and 1D), thus highlighting the relevance of the guanidine group in the

antiviral activity.

EPB-113 proved also a robust inhibitor of SARS-CoV-2 replication in A549$^{ACE2-TMPRSS2}$ cells. The

molecule was shown to have an EC$_{90}$ value (by RT-qPCR for viral load) of 12 μM and to produce more

than 5-log$_{10}$ reduction in viral load at 50 μM of compound (Fig 1B). The cell viability of mock-infected

A549$^{ACE2-TMPRSS2}$ cells was 100% at 50 μM EPB-113 and 45% at 100 μM compound (Fig 1B), which is

similar to what was seen in HEL299 cells.

Viral resistance to EPB-113 maps to the catalytic core of nsp15 EndoU

To identify the target of EPB-113, we performed selection of resistant virus. For this and all subsequent

mechanistic experiments, we chose to use HCoV-229E instead of SARS-CoV-2, since this allowed to

apply the compound at a concentration that was well above the EC$_{50}$ value. In the first step (Fig 2A), we

determined the optimal condition, being the highest MOI of HCoV-229E and lowest compound

concentration for which viral CPE was completely inhibited. Next, we applied this condition (i.e. 20 μM

of compound) to all wells of a 96-well plate with HEL299 cells. To maximize the chances of obtaining

resistance, we added EPB-113 to one plate and its analogue JMLv-061 to a second plate. At day 4 p.i.,

the few wells showing virus breakthrough were selected for virus expansion and subsequent full-genome

sequencing.

Fig 2. Resistance to EPB-113 maps to the catalytic core of nsp15 EndoU. (A) Setup of the resistance selection

experiment. (B) Alignment of partial nsp15 sequences from 11 coronaviruses, with EPB-113 and 5h resistance sites

marked with triangles. Both EPB-113 resistance sites, G248 and H250, are located at the catalytic core of nsp15

EndoU and strictly conserved in all CoVs, with H250 being part of the catalytic triad. Sequence similarities from

Clustal Omega-aligned sequences were rendered using ESPript 3.0. (C) Structure of the catalytic core of SARS-

CoV-2 EndoU bound to dsRNA (PDB 7TJ2). The residues associated with HCoV-229E resistance to EPB-113 or 5h

[39] are marked in blue and green, respectively. (D) Evaluation of EPB-113 and reference compound GS-441524

against nsp15-mutant CoVs, based on RT-qPCR-based viral load assay. Left panels: HEL299 cells infected with

HCoV-229E WT; the G248V$_{nsp15}$ or H250Y$_{nsp15}$ mutant (obtained in this study); or the K60R$_{nsp15}$ or T66$_{nsp15}$ mutant

(previously selected under compound 5h) [39]. Middle panels: 16HBE cells infected with H250A$_{nsp15}$ mutant HCoV-

229E, created by reverse genetics [11]. Right panels: A549$^{ACE2/TMPRSS2}$ cells infected with K61R$_{nsp15}$ or V67I$_{nsp15}$

mutant SARS-CoV-2, created by reverse genetics. Data points are the mean ± SEM (N=2-3). The panel legends

show the EC_{90} values and statistical significance of the difference between mutant and WT, based on (left panels)

ordinary one-way ANOVA with Dunnett's corrections for multiple comparisons and (middle panels) two-tailed

unpaired t-test.

The breakthrough viruses were found to carry mutations G248V and H250Y in the EndoU domain of

nsp15, after selection under EPB-113 and JMLv-061, respectively. After obtaining plaque-purified mutant

viruses, we confirmed their resistance by RT-qPCR-based viral load assay in which the increase in EC_{90}

value between mutant and WT was 20- to 25-fold (Fig 2D). Also another variant at the His250 site, i.e.

the $H250A_{nsp15}$ mutant form of HCoV-229E which was created by the V. Thiel team using reverse-

engineering [11], proved to be highly resistant to EPB-113 (Fig 2D). In this case the assay was done in

16HBE cells, since the $H250A_{nsp15}$ virus showed significantly impaired replication in HEL299 cells (see

below). The reference compound GS-441524 was equally active against the nsp15-mutant viruses and

WT.

Residues Gly248 and His250 are strictly conserved among CoVs and located in the catalytic center of

EndoU (Figs 2B and 2C). While His250 is part of the catalytic triad, the nearby Gly248 residue was

reported to engage in hydrogen bonding with the 3′-phosphate of the cleavage product [33].

As mentioned in the introduction, our group previously [39] identified betulonic acid derivative 5h as a

strong inhibitor of HCoV-229E which selects for resistance mutations $K60R_{nsp15}$ and $T66I_{nsp15}$ near the

EndoU catalytic core (Fig 2C). Considering that EPB-113 and 5h have totally different chemical

structures, we investigated whether the two compounds are synergistic when combined. Data analysis

with the zero-interaction potency (ZIP) model in SynergyFinder+ (S2 Fig) [48] indicated that the

combination of EPB-113 and 5h is synergistic (mean ZIP synergy score: 12), suggesting that these two

inhibitors possess non-overlapping mechanisms that potentiate each other. The combinations of EPB-

113 plus GS-441524 or 5h plus GS-441524 proved additive (mean ZIP synergy score below 10 in both

cases).

We next investigated potential cross-resistance between the two inhibitor classes. For this experiment

we used mutant viruses $T66I_{nsp15}$ and $K60R_{nsp15}$, which arose after virus passaging under 5h [39]. Fig

2D shows that the activity of EPB-113 was strongly affected by $T66I_{nsp15}$, the shift in EC_{90} value versus

WT virus being almost equally pronounced (12-fold) as for $G248V_{nsp15}$ and $H250Y_{nsp15}$. There was also

a 2-fold, yet non-significant, shift for the $K60R_{nsp15}$ mutant.

To assess whether the above resistance profile of HCoV-229E also applies to SARS-CoV-2, reverse

engineering was performed to introduce the above mutations in SARS-CoV-2. We were unable to rescue

SARS-CoV-2 virus carrying substitution $G248V_{nsp15}$, $H250Y_{nsp15}$ or $H250A_{nsp15}$. Although this observation

underscores the importance of a functional EndoU during SARS-CoV-2 replication, we did not expect it,

as all three mutations proved clearly viable in the context of HCoV-229E. On the other hand, we

successfully generated the $K61R_{nsp15}$ and $V67I_{nsp15}$ mutants of SARS-CoV-2 (corresponding to $K60R_{nsp15}$

and $V66I_{nsp15}$ in HCoV-229E), and both viruses proved to be less sensitive to EPB-113 than WT virus

(Fig 2D, right panels).

In short, resistance selection with HCoV-229E, combined with reverse engineering on SARS-CoV-2,

showed that the antiviral effect of EPB-113 is tightly linked with nsp15, and specifically its EndoU domain.

EPB-113-resistant mutants of HCoV-229E have a replication deficit

Attenuation in human macrophages

Since the nsp15 mutations causing reduced sensitivity to EPB-113 are located at (i.e. $G248V_{nsp15}$ and

$H250Y_{nsp15}$) or close to (i.e. $K60R_{nsp15}$ and $T66I_{nsp15}$) the EndoU catalytic core, we presumed that they

may impair the replication fitness of HCoV-229E. To thoroughly profile these mutants, we compared

them to WT virus as well as to the reverse-engineered $H250A_{nsp15}$ virus. The latter was previously

reported to be severely attenuated in human macrophages, due to early induction of IFN-I [11]. Hence,

we compared viral growth in human blood-derived macrophages, HEL299 human fibroblast cells (in

which our mutants arose under selective pressure of EPB-113 or 5h) and 16HBE human bronchial

epithelial cells.

In macrophages, no viral replication was detected for the $G248V_{nsp15}$, $H250Y_{nsp15}$ and $H250A_{nsp15}$

mutants (Fig 3A, left panels). The $K60R_{nsp15}$ and $T66I_{nsp15}$ mutants showed some growth impairment at

48 h and 72 h (3- to 17-fold reduction vs WT) but this was not significant. In 16HBE cells (Fig 3A, right

panels, most viruses showed very similar growth kinetics as WT virus, with only $G248V_{nsp15}$, $H250Y_{nsp15}$

and $H250A_{nsp15}$ showing a slight (3- to 8-fold) reduction at 48-72 h, which was only significant for

$H250Y_{nsp15}$ and $H250A_{nsp15}$ ($P \le 0.05$ at 48 h). An intermediate picture was seen in HEL cells (Fig 3A,

middle panels). Here, the $H250Y_{nsp15}$ and $H250A_{nsp15}$ mutants were significantly impaired (7- to 56-fold;

$P \le 0.01$) at 24 to 72 h p.i. The $G248V_{nsp15}$ and $K60R_{nsp15}$ mutants showed a modest (4- to 7-fold)

reduction, while the $T66I_{nsp15}$ virus was indistinguishable from WT virus.

Fig 3. HCoV-229E viruses bearing EPB-113 resistance mutations have impaired viral fitness. (A) Replication

kinetics in human macrophages, HEL299 cells and 16HBE cells, based on RT-qPCR for viral load. Data points are

the mean ± SEM (N=3). At each time point, statistical significance is shown for the difference between mutant and

490   WT (multiple unpaired t-tests, with Holm-Šídák's correction for multiple comparisons). (B) Competitive growth assay.

491   At the start of the experiment, each mutant was mixed with WT virus at a ratio of 9:1, 1:1 or 1:9, then applied to

492   HEL299 cells. After the virus was passaged three more times, Nanopore sequencing was performed on each

493   passage, to determine the percentage of viral genomes that contained the mutant or WT form of nsp15.

The mutant viruses are outcompeted by WT virus

To estimate the competitive fitness of the mutants vis-à-vis WT, we co-infected HEL299 cells with three

different WT:mutant ratios and then passaged the progeny virus three times; all this virus culturing was

done in the absence of compound. Each passage was submitted to Nanopore sequencing to assess

the percentage of mutant versus WT sequence in the virus population. This revealed that all nsp15

mutant forms of HCoV-229E were outcompeted by WT virus (Fig 3B). The two catalytic site mutants

($G248V_{nsp15}$ and $H250Y_{nsp15}$) showed the highest fitness cost. Their virus populations almost exclusively

consisted of WT after four passages, even if the mutant was 9-fold in excess at the start of the

experiment. Also the $K60R_{nsp15}$ and particularly the $T66I_{nsp15}$ virus were readily overtaken by WT virus.

Hence, in line with their shifted growth curves in HEL299 cells (see above), the catalytic site mutant

viruses proved unable to compete with WT virus. The other two mutations, $K60R_{nsp15}$ and $T66I_{nsp15}$, had

less dramatic effect but still rendered the virus less fit in comparison to WT virus.

Elevated expression of interferon and related players in innate immunity

So far, the only validated function of nsp15 is to avoid early induction of IFN-I by reducing the levels of

viral dsRNA replication intermediates [10-12]. Parallel to the viral growth experiments, we thus assessed

whether the mutants differed from WT virus in terms of induction of IFN-I and other innate immune

factors. The levels of bioactive IFN-α/β protein were determined in human macrophages, HEL299 and

16HBE cells, while all other factors were measured at transcript level and in HEL299 cells only. The data

are visualized as heatmaps in Fig 4, while the numeric data and levels of statistical significance are

provided in S3 Fig.

For WT virus (see top row in Fig 4B), the induction of IFN-I was evident at 48 and 72 h but not yet at 24 h p.i. The increases in bioactive IFN-I, compared to mock-infected cells, were up to 10-fold in macrophages, up to 2-fold in 16HBE cells and not seen (i.e. maximum 1.2-fold) in HEL299 cells (Fig 4B). The impact of the nsp15 mutations differed according to time and cell type and, overall, the viruses carrying EndoU catalytic site mutations had a different profile than the $K60R_{nsp15}$ and $T66I_{nsp15}$ mutants. When analyzed at 48 or 72 h p.i., macrophages infected with the $K60R_{nsp15}$ or $T66I_{nsp15}$ virus showed slightly higher levels of IFN-I than cells infected with WT virus. The $G248V_{nsp15}$, $H250Y_{nsp15}$ and $H250A_{nsp15}$ viruses not only showed the opposite profile (i.e. less IFN-I than WT at 48 and 72 h p.i.), but also differed from WT virus in showing an increase in IFN-I at 24 h p.i. Apparently, the levels of IFN-I induced by these mutant viruses within the first 24 h are sufficient to fully suppress their replication (as

evident from the growth curves above) which, in turn, prevents a further strong rise in IFN-I at later time points. The early IFN-I induction was not seen in the other two cell types. In 16HBE cells analyzed at 48 h or 72 p.i., the nsp15 mutants showed modest increases in IFN-I versus WT, although the difference reached significance ($P \leq 0.001$) for the G248V$_{nsp15}$ and H250Y$_{nsp15}$ viruses. In HEL cells, all nsp15 mutants generated markedly higher IFN-I levels than WT virus; the differences were more significant for the three catalytic site mutants (9-fold; $P \leq 0.0001$ at 72 h p.i.) than for K60R$_{nsp15}$ (4-fold; ns) and T66I$_{nsp15}$ (7-fold, $P = 0.002$).

The panel of innate immune factors, determined at transcript level in HEL299 cells, consisted of (Fig 4A): the pathogen recognition receptors (PRRs) retinoic acid-inducible gene I (RIG-I) and melanoma differentiation-associated protein 5 (MDA5); the signaling factors mitochondrial antiviral-signaling protein (MAVS), interferon regulatory factor 3 (IRF3) and 7 (IRF7); IFN-β; and antiviral effectors 2′-5′-oligoadenylate synthetases (OAS1-3), RNase L and protein kinase R (PKR). All nsp15 mutants generated significantly higher mRNA levels than WT virus for RIG-I, MDA5, IRF7, OAS1-3 and PKR. The increases were only evident at 48 and 72 h p.i., in line with the results obtained for the IFN-I bioassay in HEL299 cells. Particularly abundant expression was seen for OAS-1 and OAS-2: at 48 h p.i., their mRNA levels were 2000- to 25,000-fold increased versus mock, compared to 60- to 160-fold for WT virus, giving high significance ($P \leq 0.001$) for each comparison between mutant and WT [see the numeric values ($\log_{10}$ scale) and significance levels in S3 Fig). The catalytic site mutants showed higher expression of the innate immune factors than the other two mutants. For instance, for the G248V$_{nsp15}$, H250Y$_{nsp15}$ and H250A$_{nsp15}$ mutants, RIG-I mRNA was up to ~800-fold increased versus mock, while the K60R$_{nsp15}$ and T66I$_{nsp15}$ mutants showed a maximum increase of ~300-fold. For MAVS and RNase L, we observed no or only weak induction, regardless of whether the cells were infected with WT or mutant. Finally, we observed that HCoV-229E infection slightly reduces the transcription of IRF3 (3-fold), and that EndoU catalytic site mutations significantly ($P \leq 0.01$) revert this effect. This is in line with the finding that PEDV EndoU was reported to degrade the mRNA of IRF3 [49]. We note, however, that our analysis of innate immune factors looked only at their transcript levels, and not at their protein level, status of activation (e.g. from phosphorylation) or downstream activity.

The mutants are more sensitive to treatment with IFN-β

Considering that the EndoU-mutants induce higher levels of IFN-I effectors than WT virus, they may also be more responsive to exogenous interferon. This was indeed demonstrated by Kindler et al. [11], for H250A$_{nsp15}$-mutant HCoV-229E evaluated in human MRC5 cells. Hence, we determined the antiviral activity of recombinant human IFN-β in HEL299 cells (Fig 4C), which are the same cell type as MRC5 (i.e. both are human fibroblast cells isolated from normal fetal lung tissue). RT-qPCR for viral load was performed on the supernatants collected at day 3 p.i. Compared to WT virus, all nsp15-mutants exhibited markedly lower EC$_{90}$ values for IFN-β. The G248V$_{nsp15}$ and H250Y$_{nsp15}$ mutants showed a very strong shift of ~20-fold. The T66I$_{nsp15}$ and K60R$_{nsp15}$ mutants were less affected (= ~7- and ~3-fold lower EC$_{90}$ value than WT, respectively), but still showed significantly ($P \leq 0.0001$) higher IFN-β sensitivity than WT.

Collectively, the following conclusions can be made from this phenotypic profiling of the nsp15-mutant HCoV-229E viruses which arose under EPB-113 or 5h. Compared to WT virus, the mutants show an altered IFN-I response, but the kinetics, intensity and consequences of this effect depend on the cell type. Due to early induction of IFN-I, the EndoU catalytic site mutants are incapable of replicating in macrophages. This aligns with two previous studies on EndoU-mutant forms of HCoV-229E and MHV [10, 11]. In 16HBE cells, the mutants behave similarly to WT virus in terms of IFN-I induction and growth kinetics. An intermediate profile is seen in HEL299 cells. Here, the nsp15 mutants are prominent inducers of IFN-I and readily outcompeted by WT virus.

EPB-113 combines a direct antiviral effect with a subtle boost in type I interferon

Considering that EPB-113 selects mutations in the EndoU catalytic core, we hypothesized that it might act by altering the expression of IFN-I in CoV-infected cells. To test this hypothesis, we applied two different variations to our antiviral assay. First, we determined the inhibitory activity of EPB-113 in macrophages, HEL299 and 16HBE cells. Although EPB-113 proved nicely active in all three cell types, the EC$_{90}$ values were not identical. Compared to HEL299 cells, EPB-113 was 2-fold less potent in 16HBE cells and 6-fold more potent in macrophages (Fig 5). We also observed a difference for GS-441524, in

line with a report that the metabolic activation of this nucleoside analogue is cell type-dependent [50].

For EPB-113, a parallel can be noticed with the findings in the previous section. Namely, the nsp15

disturbance by EPB-113 proved strongest in macrophages, in line with the prominent growth impairment

of EndoU-mutant viruses in this cell type. The activity of EPB-113 was slightly higher in HEL299 than in

16HBE cells, which agrees with the competitive growth defect of the nsp15-mutant viruses seen in the

former cell line. Still, EPB-113 exerted clear activity in 16HBE cells, in which the EndoU-mutant viruses

were hardly distinguishable from WT virus in terms of growth kinetics and IFN-I induction. This was a

first indication that the compound may have a second effect that does not involve the EndoU/IFN-I axis.

Fig 5. The inhibitory potency of EPB-113 is cell-type dependent. EC_{90} values of EPB-113 (A) and GS-441524

(B) for inhibition of HCoV-229E WT virus in human macrophages, HEL299 cells and 16HBE cells, based on viral

load assay. The graph shows the individual values and mean from three independent experiments. The differences

in EC_{90} values among the three cell lines were evaluated with a one-way ANOVA with Tukey multiple comparisons

post-hoc correction.

Secondly, we assessed whether EPB-113 changes the expression of IFN-β in HCoV-229E-infected

HEL299 cells. When the compound was added together with virus (= at 0 h p.i.), the levels of IFN-β

mRNA (assessed at 48 h p.i.) were similar in cells receiving no compound, EPB-113, GS-441524 or

nirmatrelvir (S4 Fig). Strikingly, when compound addition was postponed until 24 h p.i., which gave the

virus the chance to start replicating, we observed 48 h later a significant spike in IFN-β in the condition

receiving 1 µM of EPB-113 (Fig 6B). The spike was evident from both IFN-β mRNA quantification and

IFN-I bioassay. For this delayed treatment schedule, the dose-response curve showed an extra plateau

at exactly this concentration of EPB-113 (Fig 6A). The biphasic shape of the curve indicates a dual

antiviral mechanism, in which the compound's direct antiviral effect combines with an indirect effect

through boosting of IFN-I. While GS-441524 had no such effect when added at 24 h p.i., the M^{pro} inhibitor

nirmatrelvir effectuated a strong and dose-dependent increase in the levels of IFN-β mRNA and IFN-I

protein. The IFN-I increase in the presence of nirmatrelvir implies that, besides its role in polyprotein

cleavage, the M^{pro} of HCoV-229E has immune-evading activity, as already demonstrated for the M^{pro}

enzymes of MERS-CoV [51] and SARS-CoV-2 [52]. M^{pro} was shown to exhibit cleavage activity towards

different cellular proteins [53], including RIG-I [52] and other components of the innate immune response

[54]. For both EPB-113 and nirmatrelvir, delaying treatment until 24 h p.i. was essential to ensure the

accumulation of sufficient viral RNA to function as a pathogen-associated molecular pattern (PAMP).

Subsequently, inhibiting the immune evasion activities of either nsp15 or M^{Pro} led to a significant increase

in interferon levels in response to the pre-existing viral RNA.

Fig 6. The anti-HCoV-229E activity of EPB-113 involves induction of IFN-I. Compound addition to HEL299 cells

infected with HCoV-229E WT was delayed until 24 h p.i. and 48 h later, samples were collected for assessment of

(A) viral load and (B) levels of IFN-I (left Y-axis: IFN-β mRNA; right Y-axis: bioactive IFN-I). GS-441524 and

nirmatrelvir were included as reference compounds. Data points are the mean ± SEM (N=4). Statistical analysis

was performed using a one-way ANOVA with Dunnett's correction for multiple comparisons, to compare each

compound concentration to untreated virus control (0 µM).

EPB-113 also acts at a post-entry and intermediate stage in first-cycle HCoV-229E replication

Since the above findings provided strong support for a direct effect of EPB-113 on viral replication, we wanted to gain basic insight into which stage of the replication cycle could be affected. To address this, we conducted one-cycle time-of-addition experiments in HCoV-229E-infected HEL299 cells. Besides EPB-113, we included four reference compounds: E64d (an inhibitor of cathepsin-mediated entry of the HCoV-229E ATCC strain) [55]; nirmatrelvir (an M^{pro} inhibitor); GS-441524 (an RdRp inhibitor that terminates viral RNA synthesis); and K22 (which prevents nsp6-mediated formation of the double-membrane vesicle structures where viral RNA synthesis takes place) [56]. The compounds were added to the cells at either 30 min before or several time points after virus infection. At 16 h p.i. (previously determined to mark the end of one HCoV-229E replication cycle), the cells were collected to quantify the level of intracellular viral RNA by RT-qPCR. We also analyzed a condition at 2 h p.i. (= RNA from input virus) to express the data as the fold increase in viral RNA at 16 h p.i. versus input. In the absence of compound, this fold increase was ~10,000-fold (= factor of 4 $\log_{10}$ in Fig 7).

A

B

Fig 7. Action point of EPB-113 and reference compounds in a one-cycle time-of-addition assay. (A) Experimental setup. (B) Timewise, EPB-113 has a similar action point as K22, located at an intermediate stage in HCoV-229E replication. The fold increase in intracellular viral RNA is expressed relative to the amount of input RNA at 2 h p.i. Compound concentrations: 15 µM E64d; 15 µM GS-441524; 15 µM K22 [56]; 3 µM nirmatrelvir; and 15 µM EPB-113. Data points are the mean ± SEM (N=3).

Whereas the entry inhibitor E64d lost all of its activity when added at 2 or more hours p.i., EPB-113 and K22 still had profound effect on the level of viral RNA (= ~300 times less than in the virus control) when added at 8 h p.i. (Fig 7). Nirmatrelvir already started to lose activity at 8 h p.i. (only 40-fold reduction versus virus control), consistent with inhibition of protease activity. GS-441524 was already less active when added at 4 h p.i., indicating that this nucleoside analogue requires a few hours to attain a sufficient level of its active 5′-triphosphate metabolite in HEL299 cells. The experiment further showed that neither of the tested compounds, including EPB-113, act at the stages of virus maturation or budding. Inhibitors of the latter processes would show no inhibition in our assay, since we measured the viral RNA levels

inside the cells and not the progeny virus released in the supernatant. Besides, the data corroborate

that EPB-113 also interferes with viral replication in a direct manner that does not involve IFN-I. The

mRNA for IFN-I became only detectable at 48 h p.i. of HEL299 cells (see above). Since the level of viral

RNA, produced within one replication cycle of ~16 h, is too low to induce a measurable level of IFN-I,

the strong activity of EPB-113 in the time-of-addition assay must be independent of this cytokine.

Taken together, the above results demonstrate that EPB-113 exhibits a dual effect on HCoV-229E

replication. The first effect is IFN-independent and evident from the nice activity in 16HBE cells and

strong suppression of first-cycle viral RNA synthesis. The second and indirect effect of EPB-113 occurs

via subtle boosting of IFN-I by disturbing EndoU activity. This effect provides an explanation for the

compound's superior activity in macrophages and biphasic dose-response curve in HEL299 cells.

679 The binding of EPB-113 to nsp15 destabilizes the protein
680 and stimulates its EndoU activity

In the final part, we used different biophysical and biochemical methods to study the potential binding

interaction between nsp15 protein and EPB-113, and how this binding could change the EndoU activity

of nsp15. The nsp15 proteins of HCoV-229E and SARS-CoV-2 were expressed in *E. coli* as described

by others [7, 32] and the hexameric proteins were purified by size-exclusion chromatography.

Additionally, dynamic light scattering (DLS) analysis confirmed that the two proteins were ≥98%

hexameric (S1 Fig).

Using a thermal shift assay, we found that EPB-113 causes quite pronounced reduction in the

thermostability of nsp15, producing a shift in the melting temperature (T_m) of 4°C (Fig 8A). The shift was

comparable for the proteins of HCoV-229E and SARS-CoV-2. It required a fairly high concentration of

100 μM EPB-113, which represents a 17-fold molar excess compared to the concentration of 6 μM for

nsp15 protein. The thermal shift was not at all observed with amine EPB-102, the analogue lacking the

guanidine function that exhibited much weaker anti-HCoV-229E activity in HEL299 cells (15-fold higher

EC_{50}, see first part of the results). The binding of EPB-113 to nsp15 was confirmed by surface plasmon

resonance, in which we immobilized nsp15 to the chip as ligand, then used EPB-113 and EPB-102 as

analytes. The sensorgrams (Fig 8B) show dose-dependent binding of EPB-113 to nsp15, with a similar

profile for the HCoV-229E and SARS-CoV-2 proteins. The binding was markedly less with EPB-102 (5-

to 8-fold less RU at 100 µM).

Fig 8. EPB-113 binds to hexameric nsp15 protein and enhances the EndoU cleavage reaction. (A) Thermal shift assay showing the decrease in T_m when 6 μM of nsp15 protein was incubated with 100 μM EPB-113. The controls contained 100 μM EPB-102 or the equivalent amount (= 5%) of DMSO. The top panel shows the curve of one representative experiment. The graph (bottom panel) shows the melting temperatures (individual and mean values) in three independent experiments. Statistical significance was determined with an ordinary one-way ANOVA with Dunnett's correction. (B) Dose-dependent binding between EPB-113 (top panels) or EPB-102 (bottom panels) and recombinant nsp15 from HCoV-229E (left panels) or SARS-CoV-2 (right panels), assessed by surface-plasmon resonance. RU: resonance units. (C) Scheme of the two-step EndoU cleavage reaction. (D) FRET assay in the presence of 100 μM EPB-113, 100 μM EPB-102 or the equivalent amount (= 5%) of DMSO. RFU: relative fluorescence units. An example of the real-time fluorescence curves is shown in the top panel, while the graph in the bottom panel shows the initial reaction velocity (V_0) derived from the 5-10 min interval. Individual and mean data from N=3. (E) HPLC-MS analysis on samples collected after 1 h reaction time. In the chromatograms, the colors indicate the FRET substrate; 2′,3′-cyclic intermediate; 5′-hydroxylated product; and 3′-phosphorylated product. The Y-axis shows the % normalized intensity.

To next investigate the impact of EPB-113 on EndoU cleavage, we conducted a convenient fluorescence resonance energy transfer (FRET) assay that allows to follow the reaction (Fig 8C) in real-time. Quite intriguingly, EPB-113 was found to enhance the EndoU cleavage reaction (Fig 8D). At 100 μM EPB-113, the initial reaction velocity (V_0) was increased by about 2- (SARS-CoV-2) or 4-fold (HCoV-229E) compared to the DMSO control condition (P < 0.0001). Also this effect was not observed with EPB-102. The activating effect of EPB-113 was also evident when we conducted HPLC-MS analysis on the mixtures after 1 h reaction time. This method enabled to discriminate between the starting substrate; 5′-hydroxylated reaction product; 2′,3′-cyclic phosphate reaction intermediate; and 3′-phosphorylated reaction product (see reaction scheme in Fig 8C). The latter two cannot be distinguished in the FRET assay, since they both carry the fluorophore. Although care should be taken when interpreting HPLC-MS peak areas in a quantitative manner, the chromatograms (Fig 8E) show that, in the DMSO control and EPB-102 samples, the 5′-hydroxylated product and 2′,3′-cyclic phosphate intermediate attained only low levels after 1 h reaction time. The minute amount of the 3′-phosphorylated product confirms a previous report that the 2′,3′-cyclic intermediate undergoes slow hydrolysis to this 3′-end product [32]. In the EPB-113 sample, both 5′- and 3′-end products as well as the 2′,3′-cyclic intermediate showed markedly higher peak areas, confirming that the compound stimulates the reaction. Together, these data

730   corroborate that EPB-113 induces prominent activation of the cleavage reaction, while not affecting the

731   relative abundance of the cyclic intermediate and final 3′-phosphate product.

Discussion

In this study, a detailed mechanistic investigation on the CoV replication inhibitor EPB-113 led to the discovery of a novel antiviral concept targeting the viral nsp15 protein. Additionally, our research highlights the broader importance of nsp15 for viral replication, extending beyond its EndoU activity and its role in evading innate immunity.

Our findings challenge the conventional pharmacological view that disrupting the function of nsp15 in virus-infected cells is best accomplished through inhibition of the EndoU-mediated RNA cleavage reaction. Based on this concept, the search for nsp15 inhibitors has typically relied on an enzymatic FRET assay that is suitable for high-throughput screening. Several hit compounds identified with the nsp15 protein from SARS-CoV [25] or SARS-CoV-2 [26-30] demonstrated favorable IC_{50} values in a FRET or similar enzymatic EndoU assay. However, in all cases, the compounds showed disappointing inhibitory activity in CoV-infected cells. Since these compounds have diverse chemical structures, it is unlikely that poor cell permeability is the sole issue. Rather, it may be that, unless a compound achieves complete inhibition of EndoU cleavage, viral replication cannot be effectively suppressed. Alternatively, the issue may be methodological: enzymatic EndoU assays may not accurately capture the complex functions of nsp15 in viral replication, while antiviral assays in interferon-deficient cell lines, such as Vero cells, overlook potential IFN induction by an EndoU inhibitor [38].

Our study with EPB-113 introduces a third and previously unrecognized possibility: disrupting the functions of nsp15 in virus-infected cells may be better achieved with an nsp15 binder that alters the protein's conformation in a way that enhances EndoU cleavage. The concept of enzyme activation offers several advantages, including the potential to achieve strong efficacy within cells, even if the drug only moderately activates the enzyme [57]. Although the activation by EPB-113 may seem modest based on fluorographs from the FRET assay, this effect was confirmed by HPLC-MS analysis which showed pronounced increase of the 5′-hydroxylated product, the 3′-phosphorylated product and the 2′-3′-cyclic intermediate. EPB-113 could induce these effects by causing a conformational change in the nsp15 hexameric structure, as suggested by the thermal shift assay. The reduction of T_m caused by EPB-113 is significant and contrasts with the effect reported for tipiracil [31], a UMP analogue that causes modest inhibition of the EndoU reaction and a slight increase (0.5 °C) in the thermostability of SARS-CoV-2

nsp15. In cryo-EM studies, nsp15 apoprotein is described as a highly flexible protein with 'wobbling'

EndoU domains that are stabilized upon RNA binding [32]. Although the specific structural changes

induced by EPB-113 remain speculative, we hypothesize that the compound might alter conformational

flexibility, inducing an RNA-bound state that is catalytically favorable, similar to the mechanism proposed

for the EndoU-stimulating effect of Mn^{2+} [58, 59]. Since the positively charged guanidine function (which

is fully protonated at physiological pH) is necessary for both the EndoU stimulation and strong antiviral

activity, EPB-113 may enhance the interaction between the RNA ligand and the negatively charged

surface of EndoU [33]. The idea that EPB-113 acts as a catalytic activator of EndoU aligns with its

resistance profile, as the virus escaped the compound by losing this enzymatic activity. A recent analysis

of the physiologic substrates of MHV EndoU suggested that its cleavage of specific motifs in viral

positive-sense RNA occurs in a regulated manner, to partially inhibit negative-strand RNA synthesis and

prevent accumulation of viral dsRNA [9]. It is possible that EPB-113 disrupts this finely balanced EndoU

activity. In summary, harnessing nsp15 for antiviral purposes may require rethinking the pharmacological

mechanisms best suited to achieve this objective.

As the first molecule reported to strongly suppress CoV replication in cells by binding to nsp15, EPB-

113 represents a unique tool for unravelling the functions of nsp15 beyond its role as an EndoU. An *in*

silico analysis, pending experimental validation, predicted a crucial role for nsp15 as the central scaffold

of the RTC structure [60]. Given that EPB-113 potently suppresses the replication of HCoV-229E in

16HBE cells, where the virus does not require a functional EndoU for replication, it is plausible that the

compound disrupts a structural function of nsp15. However, in HEL299 cells, EPB-113 exerts a second,

EndoU-dependent effect. The dual mechanism is evident from the biphasic dose-response curve and

subtle increase in IFN-I observed when the compound was added at 24 h post-infection. At high

concentrations, the direct effect of EPB-113 on nsp15 is dominant, whereas at intermediate

concentrations (~1 μM), the induced IFN-I helps to halt viral replication. The compound's direct effect is

further demonstrated by its strong activity in the one-cycle time-of-addition assay. Overall, the inhibitory

profile of EPB-113 strongly supports the idea that nsp15 combines roles in immune evasion and RNA

synthesis regulation.

Our phenotypic profiling of EndoU-mutant viruses confirms that, at least for HCoV-229E, having a

functional EndoU offers a clear advantage to the virus. At first sight, this seems contradictory to reports

of efficient replication of EndoU-deficient CoVs in various cell models [23, 61]. However, these studies

typically use immortalized cell lines, many of which have defects in the IFN-I pathway. Consistent with

a previous study [11], we confirm that EndoU-deficient HCoV-229E viruses (bearing nsp15 mutations

G248V, H250Y or H250A) trigger an early IFN-I response in human macrophages, rendering them

unable to replicate in these cells. While the mutants replicate efficiently in human lung fibroblasts, virus

competition experiments reveal that they are easily outcompeted by WT virus. The vulnerability of the

mutants stems from their ability to excessively stimulate the innate immune response, activating several

layers of defense, including IFN-I, PRRs, signaling factors and antiviral effectors. These responses

make the mutants more sensitive to IFN-I, amplifying their elimination when WT virus is also present.

Compared to HCoV-229E, SARS-CoV-2 may be even more dependent on EndoU for replication.

Mutations $G248V_{nsp15}$, $H250Y_{nsp15}$ and $H250A_{nsp15}$, which are viable for HCoV-229E, are not tolerated in

SARS-CoV-2. However, we successfully rescued two reverse-engineered SARS-CoV-2 mutants with

substitution $K61R_{nsp15}$ or $V67I_{nsp15}$ adjacent to the catalytic site. Using these mutants, we confirmed that

nsp15 is also the target for EPB-113 in SARS-CoV-2. Future development of EPB-113 will require

structural optimization, to achieve sufficient anti-SARS-CoV-2 activity.

Evidence suggests that the immune-evading function of nsp15 may not solely depend on EndoU

cleavage of specific motifs in viral RNA [9, 12], but could also involve cytoprotective effects or

interactions between nsp15 and a cellular protein involved in IFN production or cell cycle regulation [13,

14, 17]. While some of these functions have been proposed based on nsp15 transfection studies, they

await confirmation in authentic virus models. We cannot rule out the possibility that EPB-113 acts on

one of these alternative nsp15 functions. Considering recent links between nsp15 and stress granules

[18-20], one might hypothesize that EPB-113 induces a state of cell stress. However, this is highly

unlikely, as the compound has low cytotoxicity, even at concentrations that are well above its antiviral

range.

In conclusion, while the many unknowns regarding the function of nsp15 in CoV replication present a

challenge for antiviral drug development, our study demonstrates that nsp15 is a valid target for inhibiting

CoV replication. We provide evidence that strong antiviral activity can be achieved with a compound,

like EPB-113, which alters the conformation of nsp15 and enhances its EndoU activity. This mechanism

disrupts viral RNA synthesis while subtly boosting IFN-I production, offering a unique combination of

direct antiviral and indirect immunomodulatory effects that make the concept of nsp15 binders highly

attractive for antiviral therapy.

Acknowledgements

We wish to thank L. Delarbre, B. Ramos, E. Bernaerts and S. Wouters for excellent technical assistance and V. Thiel, B. Canard, P. Hoet, P. J. Bredenbeek, D. Jochmans, X. Saelens and P.-Y. Shi for the kind gift of viruses, plasmids or cells. Images were produced using Graphpad Prism 10.3.1 (GraphPad Software, Boston, Massachusetts USA), the UCSF Chimera package [62] from the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco (supported by NIH P41-GM103311); ESPript [63], SynergyFinder+ [48] and Marvin (https://www.chemaxon.com).

References

1. World Health Organization. WHO COVID-19 Dashboard. [23/08/2024]. Available from: https://data.who.int/dashboards/covid19/cases.

2. Infectious Diseases Society of America. IDSA guidelines on the treatment and management of patients with COVID-19. [23/08/2024]. Available from: https://www.idsociety.org/COVID19guidelines.

3. V'Kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. Coronavirus biology and replication: implications for SARS-CoV-2. Nat Rev Microbiol. 2021;19(3):155-70. Epub 2020/10/30. doi: 10.1038/s41579-020-00468-6. PubMed PMID: 33116300; PubMed Central PMCID: PMCPMC7592455.

4. Lou Z, Rao Z. The life of SARS-CoV-2 inside cells: replication-transcription complex assembly and function. Annu Rev Biochem. 2022;91:381-401. doi: 10.1146/annurev-biochem-052521-115653. PubMed PMID: 35729072.

5. Li G, Hilgenfeld R, Whitley R, De Clercq E. Therapeutic strategies for COVID-19: progress and lessons learned. Nat Rev Drug Discov. 2023;22(6):449-75. Epub 20230419. doi: 10.1038/s41573-023-00672-y. PubMed PMID: 37076602; PubMed Central PMCID: PMCPMC10113999.

6. Minkoff JM, tenOever B. Innate immune evasion strategies of SARS-CoV-2. Nat Rev Microbiol. 2023;21(3):178-94. Epub 20230111. doi: 10.1038/s41579-022-00839-1. PubMed PMID: 36631691; PubMed Central PMCID: PMCPMC9838430.

7. Ivanov KA, Hertzig T, Rozanov M, Bayer S, Thiel V, Gorbalenya AE, et al. Major genetic marker of nidoviruses encodes a replicative endoribonuclease. Proc Natl Acad Sci U S A. 2004;101(34):12694-9. Epub 2004/08/12. doi: 10.1073/pnas.0403127101. PubMed PMID: 15304651; PubMed Central PMCID: PMCPMC514660.

8. Deng X, Baker SC. An "old" protein with a new story: coronavirus endoribonuclease is important for evading host antiviral defenses. Virology. 2018;517:157-63. Epub 2018/01/09. doi: 10.1016/j.virol.2017.12.024. PubMed PMID: 29307596; PubMed Central PMCID: PMCPMC5869138.

9. Ancar R, Li Y, Kindler E, Cooper DA, Ransom M, Thiel V, et al. Physiologic RNA targets and refined sequence specificity of coronavirus EndoU. RNA. 2020;26(12):1976-99. Epub 2020/09/30. doi: 10.1261/rna.076604.120. PubMed PMID: 32989044; PubMed Central PMCID: PMCPMC7668261.

10. Deng X, Hackbart M, Mettelman RC, O'Brien A, Mielech AM, Yi G, et al. Coronavirus nonstructural protein 15 mediates evasion of dsRNA sensors and limits apoptosis in macrophages. Proc Natl Acad Sci U S A. 2017;114(21):E4251-E60. doi: 10.1073/pnas.1618310114. PubMed PMID: 28484023.

11. Kindler E, Gil-Cruz C, Spanier J, Li Y, Wilhelm J, Rabouw HH, et al. Early endonuclease-mediated evasion of RNA sensing ensures efficient coronavirus replication. PLoS Pathog. 2017;13(2):e1006195. Epub 2017/02/06. doi: 10.1371/journal.ppat.1006195. PubMed PMID: 28158275; PubMed Central PMCID: PMCPMC5310923.

12.   Hackbart M, Deng X, Baker SC. Coronavirus endoribonuclease targets viral polyuridine sequences to

evade activating host sensors. Proc Natl Acad Sci USA. 2020;117(14):8094-103. Epub 2020/03/22. doi:

10.1073/pnas.1921485117. PubMed PMID: 32198201; PubMed Central PMCID: PMCPMC7149396.

13.   Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, et al. A SARS-CoV-2 protein interaction

map reveals targets for drug repurposing. Nature. 2020;583(7816):459-68. Epub 2020/05/01. doi: 10.1038/s41586-

020-2286-9. PubMed PMID: 32353859; PubMed Central PMCID: PMCPMC7431030.

14.   Zhang D, Ji L, Chen X, He Y, Sun Y, Ji L, et al. SARS-CoV-2 Nsp15 suppresses type I interferon production

by inhibiting IRF3 phosphorylation and nuclear translocation. iScience. 2023;26(9):107705. Epub 20230823. doi:

10.1016/j.isci.2023.107705. PubMed PMID: 37680466; PubMed Central PMCID: PMCPMC10480782.

15.   Lei Y, Moore CB, Liesman RM, O'Connor BP, Bergstralh DT, Chen ZJ, et al. MAVS-mediated apoptosis

and its inhibition by viral proteins. PLoS One. 2009;4(5):e5466. Epub 2009/05/01. doi:

10.1371/journal.pone.0005466. PubMed PMID: 19404494; PubMed Central PMCID: PMCPMC2674933.

16.   Hayn M, Hirschenberger M, Koepke L, Nchioua R, Straub JH, Klute S, et al. Systematic functional analysis

of SARS-CoV-2 proteins uncovers viral innate immune antagonists and remaining vulnerabilities. Cell Rep.

2021:109126. Epub 2021/05/12. doi: 10.1016/j.celrep.2021.109126. PubMed PMID: 33974846; PubMed Central

PMCID: PMCPMC8078906.

17.   Bhardwaj K, Liu P, Leibowitz JL, Kao CC. The coronavirus endoribonuclease nsp15 interacts with

retinoblastoma tumor suppressor protein. J Virol. 2012;86(8):4294-304. Epub 2012/02/04. doi: 10.1128/JVI.07012-

11. PubMed PMID: 22301153; PubMed Central PMCID: PMCPMC3318636.

18.   Gao B, Gong X, Fang S, Weng W, Wang H, Chu H, et al. Inhibition of anti-viral stress granule formation by

coronavirus endoribonuclease nsp15 ensures efficient virus replication. PLoS Pathog. 2021;17(2):e1008690. Epub

2021/02/27. doi: 10.1371/journal.ppat.1008690. PubMed PMID: 33635931; PubMed Central PMCID:

PMCPMC7946191.

19.   Zhao J, Feng D, Zhao Y, Huang M, Zhang X, Zhang G. Role of stress granules in suppressing viral

replication by the infectious bronchitis virus endoribonuclease. J Virol. 2022;96(12):e0068622. Epub 20220531. doi:

10.1128/jvi.00686-22. PubMed PMID: 35638780; PubMed Central PMCID: PMCPMC9215229.

20.   Dolliver SM, Kleer M, Bui-Marinos MP, Ying S, Corcoran JA, Khaperskyy DA. Nsp1 proteins of human

coronaviruses HCoV-OC43 and SARS-CoV2 inhibit stress granule formation. PLoS Pathog. 2022;18(12):e1011041.

Epub 20221219. doi: 10.1371/journal.ppat.1011041. PubMed PMID: 36534661; PubMed Central PMCID:

PMCPMC9810206.

21.   Deng X, van Geelen A, Buckley AC, O'Brien A, Pillatzki A, Lager KM, et al. Coronavirus endoribonuclease

activity in porcine epidemic diarrhea virus suppresses type I and type III interferon responses. J Virol.

2019;93(8):e02000-18. Epub 2019/02/08. doi: 10.1128/JVI.02000-18. PubMed PMID: 30728254; PubMed Central

PMCID: PMCPMC6450110.

22. Comar CE, Otter CJ, Pfannenstiel J, Doerger E, Renner DM, Tan LH, et al. MERS-CoV endoribonuclease

and accessory proteins jointly evade host innate immunity during infection of lung and nasal epithelial cells. Proc

Natl Acad Sci U S A. 2022;119(21):e2123208119. Epub 20220520. doi: 10.1073/pnas.2123208119. PubMed PMID:

35594398; PubMed Central PMCID: PMCPMC9173776.

23. Otter CJ, Bracci N, Parenti NA, Ye C, Asthana A, Blomqvist EK, et al. SARS-CoV-2 nsp15

endoribonuclease antagonizes dsRNA-induced antiviral signaling. Proc Natl Acad Sci U S A.

2024;121(15):e2320194121. Epub 20240403. doi: 10.1073/pnas.2320194121. PubMed PMID: 38568967; PubMed

Central PMCID: PMCPMC11009620.

24. Xie Y, Chen C, Zhang D, Jiao Z, Chen Y, Wang G, et al. Diversity for endoribonuclease nsp15-mediated

regulation of alpha-coronavirus propagation and virulence. Microbiol Spectr. 2023;11(6):e0220923. Epub 20231108.

doi: 10.1128/spectrum.02209-23. PubMed PMID: 37938022; PubMed Central PMCID: PMCPMC10715224.

25. Ortiz-Alcantara JM, Bhardwaj K, Palaninathan S, Frieman M, Baric RS, Kao CC. Small molecule inhibitors

of the SARS-CoV nsp15 endoribonuclease. Virus Adapt Treat. 2010;2:125-33. doi: 10.2147/vaat.s12733.

26. Canal B, Fujisawa R, McClure AW, Deegan TD, Wu M, Ulferts R, et al. Identifying SARS-CoV-2 antiviral

compounds by screening for small molecule inhibitors of nsp15 endoribonuclease. Biochem J. 2021;478(13):2465-

79. Epub 2021/07/02. doi: 10.1042/BCJ20210199. PubMed PMID: 34198324.

27. Choi R, Zhou M, Shek R, Wilson JW, Tillery L, Craig JK, et al. High-throughput screening of the ReFRAME,

Pandemic Box, and COVID Box drug repurposing libraries against SARS-CoV-2 nsp15 endoribonuclease to identify

small-molecule inhibitors of viral activity. PLoS One. 2021;16(4):e0250019. Epub 2021/04/23. doi:

10.1371/journal.pone.0250019. PubMed PMID: 33886614; PubMed Central PMCID: PMCPMC8062000.

28. Hong S, Seo SH, Woo SJ, Kwon Y, Song M, Ha NC. Epigallocatechin gallate inhibits the uridylate-specific

endoribonuclease nsp15 and efficiently neutralizes the SARS-CoV-2 strain. J Agric Food Chem. 2021;69(21):5948-

54. Epub 2021/05/22. doi: 10.1021/acs.jafc.1c02050. PubMed PMID: 34015930; PubMed Central PMCID:

PMCPMC8146138.

29. Chen J, Farraj RA, Limonta D, Tabatabaei Dakhili SA, Kerek EM, Bhattacharya A, et al. Reversible and

irreversible inhibitors of coronavirus Nsp15 endoribonuclease. J Biol Chem. 2023;299(11):105341. Epub 20231011.

doi: 10.1016/j.jbc.2023.105341. PubMed PMID: 37832873; PubMed Central PMCID: PMCPMC10656235.

30. Ma Y, Luo M, Deng Y, Yang X, Wang X, Chen G, et al. Antibiotic-induced primary biles inhibit SARS-CoV-

2 endoribonuclease nsp15 activity in mouse gut. Front Cell Infect Microbiol. 2022;12:896504. Epub 2022/08/16.

doi: 10.3389/fcimb.2022.896504. PubMed PMID: 35967852; PubMed Central PMCID: PMCPMC9366059.

31. Kim Y, Wower J, Maltseva N, Chang C, Jedrzejczak R, Wilamowski M, et al. Tipiracil binds to uridine site

and inhibits nsp15 endoribonuclease NendoU from SARS-CoV-2. Commun Biol. 2021;4(1):193. Epub 2021/02/11.

doi: 10.1038/s42003-021-01735-9. PubMed PMID: 33564093.

32.   Pillon MC, Frazier MN, Dillard LB, Williams JG, Kocaman S, Krahn JM, et al. Cryo-EM structures of the

SARS-CoV-2 endoribonuclease nsp15 reveal insight into nuclease specificity and dynamics. Nat Commun.

2021;12(1):636. Epub 2021/01/29. doi: 10.1038/s41467-020-20608-z. PubMed PMID: 33504779; PubMed Central

PMCID: PMCPMC7840905.

33.   Frazier MN, Dillard LB, Krahn JM, Perera L, Williams JG, Wilson IM, et al. Characterization of SARS2

nsp15 nuclease activity reveals it's mad about U. Nucleic Acids Res. 2021;49(17):10136-49. Epub 2021/08/18. doi:

10.1093/nar/gkab719. PubMed PMID: 34403466; PubMed Central PMCID: PMCPMC8385992.

34.   Frazier MN, Wilson IM, Krahn JM, Butay KJ, Dillard LB, Borgnia MJ, et al. Flipped over U: structural basis

for dsRNA cleavage by the SARS-CoV-2 endoribonuclease. Nucleic Acids Res. 2022;50(14):8290-301. Epub

2022/07/09. doi: 10.1093/nar/gkac589. PubMed PMID: 35801916; PubMed Central PMCID: PMCPMC9371922.

35.   Ito F, Yang H, Zhou ZH, Chen XS. Structural basis for polyuridine tract recognition by SARS-CoV-2 Nsp15.

Protein Cell. 2024;15(7):547-52. Epub 20240413. doi: 10.1093/procel/pwae009. PubMed PMID: 38613382;

PubMed Central PMCID: PMCPMC11214832.

36.   Ricagno S, Egloff MP, Ulferts R, Coutard B, Nurizzo D, Campanacci V, et al. Crystal structure and

mechanistic determinants of SARS coronavirus nonstructural protein 15 define an endoribonuclease family. Proc

Natl Acad Sci USA. 2006;103(32):11892-7. doi: 10.1073/pnas.0601708103. PubMed PMID: 16882730; PubMed

Central PMCID: PMCPMC2131687.

37.   Joseph JS, Saikatendu KS, Subramanian V, Neuman BW, Buchmeier MJ, Stevens RC, et al. Crystal

structure of a monomeric form of severe acute respiratory syndrome coronavirus endonuclease nsp15 suggests a

role for hexamerization as an allosteric switch. J Virol. 2007;81(12):6700-8. Epub 2007/04/06. doi:

10.1128/JVI.02817-06. PubMed PMID: 17409150; PubMed Central PMCID: PMCPMC1900129.

38.   Van Loy B, Stevaert A, Naesens L. The coronavirus nsp15 endoribonuclease: A puzzling protein and

pertinent antiviral drug target. Antiviral Res. 2024;228:105921. Epub 20240531. doi:

10.1016/j.antiviral.2024.105921. PubMed PMID: 38825019.

39.   Stevaert A, Krasniqi B, Van Loy B, Nguyen T, Thomas J, Vandeput J, et al. Betulonic acid derivatives

interfering with human coronavirus 229E replication via the nsp15 endoribonuclease. J Med Chem.

2021;64(9):5632-44. Epub 2021/04/21. doi: 10.1021/acs.jmedchem.0c02124. PubMed PMID: 33877845; PubMed

Central PMCID: PMCPMC8084268.

40.   Vrijens P, Noppen S, Boogaerts T, Vanstreels E, Ronca R, Chiodelli P, et al. Influenza virus entry via the

GM3 ganglioside-mediated platelet-derived growth factor receptor beta signalling pathway. J Gen Virol.

2019;100(4):583-601. Epub 2019/02/15. doi: 10.1099/jgv.0.001235. PubMed PMID: 30762518.

41.   Szűcs Z, Naesens L, Stevaert A, Ostorházi E, Batta G, Herczegh P, et al. Reprogramming of the

antibacterial drug vancomycin results in potent antiviral agents devoid of antibacterial activity. Pharmaceuticals

(Basel). 2020;13(7):139. Epub 2020/07/03. doi: 10.3390/ph13070139. PubMed PMID: 32610683; PubMed Central

PMCID: PMCPMC7407158.

42.   Delang L, Segura Guerrero N, Tas A, Querat G, Pastorino B, Froeyen M, et al. Mutations in the

chikungunya virus non-structural proteins cause resistance to favipiravir (T-705), a broad-spectrum antiviral. J

Antimicrob Chemother. 2014;69(10):2770-84. doi: 10.1093/jac/dku209. PubMed PMID: 24951535.

43.   Xie X, Lokugamage KG, Zhang X, Vu MN, Muruato AE, Menachery VD, et al. Engineering SARS-CoV-2

using a reverse genetic system. Nat Protoc. 2021;16(3):1761-84. Epub 2021/01/31. doi: 10.1038/s41596-021-

00491-8. PubMed PMID: 33514944; PubMed Central PMCID: PMCPMC8168523.

44.   Li H. New strategies to improve minimap2 alignment accuracy. Bioinformatics. 2021;37(23):4572-4. doi:

10.1093/bioinformatics/btab705. PubMed PMID: 34623391; PubMed Central PMCID: PMCPMC8652018.

45.   Khanna A, Larson D, Srivatsan S, Mosior M, Abbott T, Kiwala S, et al. Bam-readcount - rapid generation

of basepair-resolution sequence metrics. J Open Source Softw. 2022;7(69). doi: 10.21105/joss.03722.

46.   Xu X, Zhai Y, Sun F, Lou Z, Su D, Xu Y, et al. New antiviral target revealed by the hexameric structure of

mouse hepatitis virus nonstructural protein nsp15. J Virol. 2006;80(16):7909-17. Epub 2006/07/29. doi:

10.1128/JVI.00525-06. PubMed PMID: 16873248; PubMed Central PMCID: PMCPMC1563835.

47.   Guarino LA, Bhardwaj K, Dong W, Sun J, Holzenburg A, Kao C. Mutational analysis of the SARS virus

nsp15 endoribonuclease: identification of residues affecting hexamer formation. J Mol Biol. 2005;353(5):1106-17.

Epub 2005/10/12. doi: 10.1016/j.jmb.2005.09.007. PubMed PMID: 16216269.

48.   Zheng S, Wang W, Aldahdooh J, Malyutina A, Shadbahr T, Tanoli Z, et al. SynergyFinder Plus: Toward

better interpretation and annotation of drug combination screening datasets. Genomics Proteomics Bioinformatics.

2022;20(3):587-96. Epub 20220125. doi: 10.1016/j.gpb.2022.01.004. PubMed PMID: 35085776; PubMed Central

PMCID: PMCPMC9801064.

49.   Wu Y, Zhang H, Shi Z, Chen J, Li M, Shi H, et al. Porcine epidemic diarrhea virus nsp15 antagonizes

interferon signaling by RNA degradation of TBK1 and IRF3. Viruses. 2020;12(6):599. Epub 20200531. doi:

10.3390/v12060599. PubMed PMID: 32486349; PubMed Central PMCID: PMCPMC7354440.

50.   Tao S, Zandi K, Bassit L, Ong YT, Verma K, Liu P, et al. Comparison of anti-SARS-CoV-2 activity and

intracellular metabolism of remdesivir and its parent nucleoside. Curr Res Pharmacol Drug Discov. 2021;2:100045.

Epub 20210812. doi: 10.1016/j.crphar.2021.100045. PubMed PMID: 34870151; PubMed Central PMCID:

PMCPMC8357487.

51.   van Huizen M, Vendrell XM, de Gruyter HLM, Boomaars-van der Zanden AL, van der Meer Y, Snijder EJ,

et al. The main protease of Middle East respiratory syndrome coronavirus induces cleavage of mitochondrial

antiviral signaling protein to antagonize the innate immune response. Viruses. 2024;16(2):256. Epub 20240205.

doi: 10.3390/v16020256. PubMed PMID: 38400032; PubMed Central PMCID: PMCPMC10892576.

52.   Liu Y, Qin C, Rao Y, Ngo C, Feng JJ, Zhao J, et al. SARS-CoV-2 Nsp5 demonstrates two distinct

mechanisms targeting RIG-I and MAVS to evade the innate immune response. mBio. 2021;12(5):e0233521. Epub

20210921. doi: 10.1128/mBio.02335-21. PubMed PMID: 34544279; PubMed Central PMCID: PMCPMC8546575.

53.   Melano I, Lo YC, Su WC. Characterization of host substrates of SARS-CoV-2 main protease. Front

Microbiol. 2023;14:1251705. Epub 20230821. doi: 10.3389/fmicb.2023.1251705. PubMed PMID: 37670988;

PubMed Central PMCID: PMCPMC10475589.

54.   Song L, Wang D, Abbas G, Li M, Cui M, Wang J, et al. The main protease of SARS-CoV-2 cleaves histone

deacetylases and DCP1A, attenuating the immune defense of the interferon-stimulated genes. J Biol Chem.

2023;299(3):102990. Epub 20230208. doi: 10.1016/j.jbc.2023.102990. PubMed PMID: 36758802; PubMed Central

PMCID: PMCPMC9907797.

55.   Shirato K, Kanou K, Kawase M, Matsuyama S. Clinical isolates of human coronavirus 229E bypass the

endosome for cell entry. J Virol. 2017;91(1):e01387-16. Epub 2016/11/01. doi: 10.1128/JVI.01387-16. PubMed

PMID: 27733646; PubMed Central PMCID: PMCPMC5165181.

56.   Lundin A, Dijkman R, Bergstrom T, Kann N, Adamiak B, Hannoun C, et al. Targeting membrane-bound

viral RNA synthesis reveals potent inhibition of diverse coronaviruses including the Middle East respiratory

syndrome virus. PLoS Pathog. 2014;10(5):e1004166. doi: 10.1371/journal.ppat.1004166. PubMed PMID:

24874215; PubMed Central PMCID: PMCPMC4038610.

57.   Zorn JA, Wells JA. Turning enzymes ON with small molecules. Nat Chem Biol. 2010;6(3):179-88. doi:

10.1038/nchembio.318. PubMed PMID: 20154666.

58.   Huang T, Snell KC, Kalia N, Gardezi S, Guo L, Harris ME. Kinetic analysis of RNA cleavage by coronavirus

nsp15 endonuclease: evidence for acid-base catalysis and substrate-dependent metal ion activation. J Biol Chem.

2023;299(6):104787. Epub 20230504. doi: 10.1016/j.jbc.2023.104787. PubMed PMID: 37149147; PubMed Central

PMCID: PMCPMC10158045.

59.   Bhardwaj K, Guarino L, Kao CC. The severe acute respiratory syndrome coronavirus Nsp15 protein is an

endoribonuclease that prefers manganese as a cofactor. J Virol. 2004;78(22):12218-24. Epub 2004/10/28. doi:

10.1128/JVI.78.22.12218-12224.2004. PubMed PMID: 15507608; PubMed Central PMCID: PMCPMC525082.

60.   Perry JK, Appleby TC, Bilello JP, Feng JY, Schmitz U, Campbell EA. An atomistic model of the coronavirus

replication-transcription complex as a hexamer assembled around nsp15. J Biol Chem. 2021;297(4):101218. Epub

20210923. doi: 10.1016/j.jbc.2021.101218. PubMed PMID: 34562452; PubMed Central PMCID:

PMCPMC8494237.

61.   Kang H, Bhardwaj K, Li Y, Palaninathan S, Sacchettini J, Guarino L, et al. Biochemical and genetic

analyses of murine hepatitis virus nsp15 endoribonuclease. J Virol. 2007;81(24):13587-97. Epub 2007/09/28. doi:

10.1128/JVI.00547-07. PubMed PMID: 17898055; PubMed Central PMCID: PMCPMC2168834.

62.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera - a

visualization system for exploratory research and analysis. J Comput Chem. 2004;25(13):1605-12. doi:

10.1002/jcc.20084. PubMed PMID: 15264254.

63.   Robert X, Gouet P. Deciphering key features in protein structures with the new ENDscript server. Nucleic

Acids Res. 2014;42(Web Server issue):W320-W4. Epub 2014/04/23. doi: 10.1093/nar/gku316. PubMed PMID:

24753421; PubMed Central PMCID: PMCPMC4086106.

1033

Supporting information

S1 Appendix. Chemical synthesis and analysis of EPB-113 and its two structural analogues EPB-102 and JMLv-061.

S1 Appendix: Chemistry

Table of contents

Chemical synthesis. General methods.

Commercially available reagents and solvents were used without further purification unless stated otherwise. Preparative normal phase chromatography was performed on a CombiFlash Rf 150 (Teledyne Isco) with pre-packed RediSep Rf silica gel cartridges. Thin-layer chromatography was performed with aluminum-backed sheets with silica gel 60 F_{254} (Merck, ref 1.05554), and spots were visualized with UV light and 1% aqueous solution of $KMnO_4$. Melting points were determined in open capillary tubes with a MFB 595010M Gallenkamp. 400 MHz ^{1}H, 100.6 MHz ^{13}C and 376.5 MHz ^{19}F NMR spectra were recorded on a Varian Mercury 400 or on a Bruker 400 Avance III spectrometers. The chemical shifts are reported in ppm (δ scale) relative to internal tetramethylsilane, and coupling constants are reported in Hertz (Hz). Assignments given for the NMR spectra of the new compounds have been carried out on the basis of COSY ^{1}H/^{13}C (gHSQC sequence) experiments. IR spectra were run on a Perkin-Elmer Spectrum RX I spectrophotometer. Absorption values are expressed as wave-numbers (cm^{-1}); only significant absorption bands are given. High-resolution mass spectrometry (HRMS) analyses were performed with an LC/MSD TOF Agilent Technologies spectrometer. The elemental analyses were carried out in a Flash 1112 series Thermofinnigan elemental microanalyzator (A5) to determine C, H, N and S. The structure of all new compounds was confirmed by elemental analysis and/or accurate mass measurement, IR, ^{1}H NMR, ^{13}C NMR and ^{19}F NMR. The analytical samples of all the new compounds, which were subjected to antiviral evaluation, possessed purity ≥ 95% as evidenced by their elemental analyses and/or their HPLC/MS. HPLC/MS for EPB-113 was determined with a HPLC Agilent 1260 Infinity II LC/MSD coupled to a photodiode array and mass spectrometer. 5 µL of sample 0.5 mg/mL in methanol:acetonitrile were injected, using a Agilent Poroshell 120 EC-C18, 2.7 µm, 50 mm x 4.6 mm column at 40 °C. The mobile phase was a mixture of A = water with 0.05% formic acid and B = acetonitrile with 0.05% formic acid, with the method described as follows: flow 0.6 mL/min, from 95% A - 5% B to 100% B in 3 min, 100% B 3 min, from 100% B to 95% A - 5% B in 1 min, 95% A – 5% B 3 min. Purity is given as % of absorbance at 254 nm. Data from mass spectra were analyzed by electrospray ionization in positive and negative between 100 and 1000 Da.

Synthesis of 1-(3-aminophenyl)-3-(3-(pentafluoro-λ⁶-sulfanyl)phenyl)urea, EPB-102

Step 1: Synthesis of 1-(3-nitrophenyl)-3-(3-(pentafluoro-λ⁶-sulfanyl)phenyl)urea

3-Nitrophenylisocyanate (500 mg, 3.05 mmol) was dissolved in THF (25 mL) and 3-pentafluorosulfanyl aniline (668 mg, 3.05 mmol) was added. The reaction mixture was stirred at 45 °C for 2 hours. The solvent was removed under reduced pressure to afford a yellow solid (1.10 g) that was crystallized in EtOAc to give pure 1-(3-nitrophenyl)-3-(3-(pentafluoro-λ⁶-sulfanyl)phenyl)urea (1.10 g, 94% yield) as a yellow crystalline solid.

Step 2: Synthesis of EPB-102

A suspension of 1-(3-nitrophenyl)-3-(3-(pentafluoro-λ⁶-sulfanyl)phenyl)urea (600 mg, 1.56 mmol) and PtO$_2$ hydrate (53 mg) in abs. ethanol (45 mL) was hydrogenated (1 atm) at room temperature for 6 hours. The suspension was then filtered, and the solvent was evaporated under vacuum to afford **EPB-102** (533 mg, 97% yield) as a white solid. The analytical sample was obtained by crystallization from EtOAc, mp (EtOAc) 193–194 °C. IR (ATR) ν: 643, 651, 682, 697, 789, 824, 845, 877, 913, 1167, 1229, 1313, 1420, 1465, 1485, 1566, 1590, 1651, 3092, 3294, 3389, 3489 cm⁻¹. ¹H-NMR (400 MHz, DMSO-d_6) δ: 5.05 (s, 2 H, NH$_2$), 6.21 (m, 1 H, 6-H or 4-H), 6.52 (m, 1 H, 4-H or 6-H), 6.84 (t, J = 2.0 Hz, 1 H, 2-H), 6.90 (t, J = 8.0 Hz, 1 H, 5-H), 7.40-7.55 (complex signal, 3 H, 4'-H, 5'-H, 6'-H), 8.28 (t, J = 2.0 Hz, 1 H, 2'-H), 8.51 (s, 1 H) and 9.03 (s, 1 H) (2 NH). ¹³C-NMR (100.6 MHz, DMSO-d_6) δ: 104.1 (CH, C2), 106.3 (CH, C4 or C6), 108.4 (CH, C6 or C4), 114.6 (quint, $^3J_{CF}$ = 5.0 Hz, CH, C2'), 118.5 (quint, $^3J_{CF}$ = 3.5 Hz, CH, C4'), 121.5 (CH, C6'), 129.0 (CH, C5), 129.6 (CH, C5'), 139.8 (C, C1), 140.7 (C, C1'), 149.2 (C, C3), 152.3 (C, CO), 153.3 (quint, $^2J_{CF}$ = 15.1 Hz, C, C3'). ¹⁹F-NMR (376.5 MHz, DMSO-d_6) δ: 63.6 (d, J = 150.6 Hz, 4 F, S<u>F</u>$_4$F), 87.8 (quint, J = 150.6 Hz, 1 F, SF$_4$<u>F</u>). HRMS-ESI⁻ *m/z* [M-H]⁻ calculated for [C$_{13}$H$_{12}$F$_5$N$_3$OS-H]⁻: 352.0548. Found: 352.0554. Elemental analysis calculated for C$_{13}$H$_{12}$F$_5$N$_3$OS: C 44.19%, H 3.42%, N 11.89%, S 9.07%. Found: C 44.24%, H 3.44%, N 11.63%, S 8.78%.

Synthesis of 1-(3-guanidinophenyl)-3-(3-(pentafluoro-λ^6-sulfanyl)-phenyl)urea hydrochloride, EPB-113

Step 1: Synthesis of 2-(3-nitrophenyl)-1,3-di-Boc-guanidine

To a stirred solution of 3-nitroaniline (1 g, 7.24 mmol) in DMF (10 mL) was added bis-Boc-pyrazolocarboxamidine (2.25 g, 7.25 mmol) and the reaction mixture was stirred at 40 °C for 24 hours. Water (15 mL) and EtOAc (15 mL) were added, and the phases were separated. The organic layer was further washed with water, dried over anhydrous Na_2SO_4 and filtered. Evaporation of the organics provided a yellowish gum (2.46 g), which was purified by column chromatography in silica gel (hexane/EtOAc mixtures). Fractions containing the desired product were collected and concentrated under vacuum to afford 2-(3-nitrophenyl)-1,3-di-Boc-guanidine (1.0 g, 36% yield) as a yellowish solid. The spectroscopic data matched with those previously described in the bibliography.[1] HRMS-ESI$^+$ *m/z* [M+H]$^+$ calculated for [C$_{17}$H$_{24}$N$_4$O$_6$+H]$^+$: 381.1769. Found: 381.1769.

Step 2: Obtention of 2-(3-aminophenyl)-1,3-di-Boc-guanidine

A suspension of 2-(3-nitrophenyl)-1,3-di-Boc-guanidine (1 g, 2.63 mmol) and PtO$_2$ hydrate (100 mg) in abs. ethanol (100 mL) was hydrogenated (1 atm) at room temperature for 3 hours. The suspension was then filtered, and the solvent was evaporated under vacuum to afford 2-(3-aminophenyl)-1,3-di-Boc-guanidine (0.91 g, quantitative yield) as a white foamy solid. The spectroscopic data matched with those previously described in the bibliography.[2]

Step 3: Synthesis of 1-(1,3-di-Boc-3-guanidinophenyl)-3-(3-(pentafluoro-λ^6-sulfanyl)phenyl)urea

i) 3-pentafluorosulfanyl aniline (80 mg, 0.36 mmol) was added to a stirring biphasic mixture of DCM (5 mL) and saturated aqueous NaHCO$_3$ solution (5 mL). Triphosgene (40 mg, 0.14 mmol) was then slowly added and the reaction mixture was stirred at room temperature for 30 minutes. The phases were separated and the organic layer was washed with brine, dried over anhydrous Na$_2$SO$_4$, filtered and half concentrated at room temperature to give 3-(pentafluoro-λ^6-sulfanyl)phenylisocyanate in DCM solution that was used in the next step without further purification.

ii) 2-(3-aminophenyl)-1,3-di-Boc-guanidine (139 mg, 0.40 mmol) dissolved in DCM (2 mL) was added to the previously obtained isocyanate in DCM (ca. 6 mL). The reaction mixture was stirred at room temperature overnight. Evaporation under vacuum of the solvent provided an orange solid (350 mg) which was purified by column chromatography in silica gel (hexane/EtOAc mixtures). Fractions containing the desired product were collected and concentrated under vacuum to give 1-(1,3-di-Boc-3-guanidinophenyl)-3-(3-(pentafluoro-λ^6-sulfanyl)phenyl)urea (60 mg, 29% overall yield) as a yellowish solid. The analytical sample was obtained by washing with Et$_2$O. ^{1}H-NMR (400 MHz, DMSO-d_6) δ: 1.41 (s, 9 H, Boc), 1.51 (s, 9 H, Boc), 7.15-7.35 (complex signal, 3 H, 2-H, 4-H, 6-H), 7.41-7.59 (complex signal, 3 H, 4'-H, 5-H, 6'-H), 7.64 (s, 1 H, 5'-H), 8.27 (m, 1 H, 2'-H), 8.93 (s, 1 H) and 9.17 (s, 1 H) (2 NH urea), 9.98 (s, 1 H) and 11.35 (s, 1 H) (2 NH guanidine).

Step 4: Synthesis of EPB-113 (as its hydrochloride salt)

To a stirred suspension of 1-(1,3-di-Boc-3-guanidinophenyl)-3-(3-(pentafluoro-λ^6-sulfanyl)phenyl)urea (32 mg, 0.05 mmol) in DCM (1 mL) was added a 4 M solution of HCl in 1,4-dioxane (4 mL, 16 mmol) and the mixture was stirred at room temperature overnight. Evaporation of the solvents provided a beige solid, which was washed with Et$_2$O and pentane to give **EPB-113** as its hydrochloride salt (19 mg, 88% yield), mp 219–220 °C. IR (ATR) ν: 754, 782, 812, 835, 857, 916, 1098, 1239, 1296, 1406, 1428, 1484, 1496, 1542, 1587, 1598, 1673, 2966, 3184, 3235, 3416 cm^{-1}. ^{1}H-NMR (400 MHz, DMSO-d_6) δ: 6.86 (m, 1 H, 4-H), 7.27 (m, 1 H, 6-H), 7.35 (t, J = 8.0 Hz, 1 H, 5-H), 7.38 (broad s, 4 H, 2 NH$_2$ guanidine), 7.46-7.55 (complex signal, 4 H, 2-H, 4'-H, 5'-H, 6'-H), 8.27 (m, 1 H, 2'-H), 9.31 (s, 1 H) and 9.53 (s, 1 H) (2 NH urea), 9.70 (s, 1 H, 3-NH). ^{13}C-NMR (100.6 MHz, CD$_3$OD) δ: 117.2 (CH, C2), 117.4 (quint, $^3J_{CF}$ = 5.0 Hz, CH, C2'), 119.4 (CH, C6), 120.5 (CH, C4), 120.9 (quint, $^3J_{CF}$ = 5.0 Hz, CH, C4'), 123.1 (CH, C6'), 130.3 (CH, C5'), 131.4 (CH, C5), 136.6 (C, C1), 141.3 (C, C1'), 142.0 (C, C3), 154.8 (C, CO), 155.4 (quint, $^2J_{CF}$ = 17.1 Hz, C, C3'), 158.0 (C, CN guanidine). ^{19}F-NMR (376.5 MHz, DMSO-d_6) δ: 63.6 (d, J = 150.6 Hz, 4 F, S$\underline{F}_4$F), 87.7 (quint, J = 150.6 Hz, 1 F, SF$_4$$\underline{F}$). HRMS-ESI$^+$ *m/z* [M+H]$^+$ calculated for [C$_{14}$H$_{14}$F$_5$N$_5$OS+H]$^+$: 396.0912. Found: 396.0916. HPLC (254 nm): t$_R$ = 3.75 min (98.4%).

Elemental analysis calculated for $C_{14}H_{14}F_5N_5OS \cdot HCl \cdot 0.4H_2O$: C 38.30%, H 3.63%, N 15.95%. Found: C 38.51%, H 3.88%, N 16.21%.

Synthesis of 1-(3-guanidinophenyl)-3-(2-(trifluoromethoxy)phenyl)urea dihydrochloride, JMLV-061

Step 1: Synthesis of 2-(3-nitrophenyl)-1,3-di-Boc-guanidine

To a solution of 3-nitroaniline (1.50 g, 10.86 mmol) and triethylamine (4.54 mL, 32.58 mmol) in anhydrous DCM (40 mL) was added *N,N'*-bis(tert-butoxycarbonyl)-S-methylisothiourea (3.31 g, 11.40 mmol) and HgCl$_2$ (3.24 g, 11.95 mmol) and the mixture was stirred at RT for 16 hours. The mixture was filtered through a pad of Celite® using DCM as eluting agent. Then, the solution was washed with water (2 x 80 mL). The organic layer was dried over anhydrous Na$_2$SO$_4$ and filtered. Solvents were concentrated under vacuum and the resulting crude was purified by column chromatography in silica gel (hexane/EtOAc mixtures). Fractions containing the desired product were collected and concentrated under vacuum to afford 2-(3-nitrophenyl)-1,3-di-Boc-guanidine as a pale-yellow solid (3.88 g, 94% yield). The spectroscopic data matched with those previously described in the bibliography.[1] HRMS-ESI$^+$ *m/z* [M+H]$^+$ calculated for [C$_{17}$H$_{24}$N$_4$O$_6$+H]$^+$: 381.1769. Found: 381.1769.

Step 2: Obtention of 2-(3-aminophenyl)-1,3-di-Boc-guanidine

A suspension of 2-(3-nitrophenyl)-1,3-di-Boc-guanidine (2.28 g, 5.99 mmol) and PtO$_2$ hydrate (250 mg) in abs. ethanol (50 mL) was hydrogenated at room temperature and atmospheric pressure for 3 hours. Anhydrous Na$_2$SO$_4$ and Celite® were added and the mixture was stirred for 2 minutes. Then, the suspension was filtered and the solvent was evaporated under vacuum. The resulting crude was purified by column chromatography in silica gel (hexane/EtOAc mixtures). Fractions containing the desired product were collected and concentrated under vacuum to afford 2-(3-aminophenyl)-1,3-di-Boc-guanidine (1.75 g, 83% yield) as a white solid. The spectroscopic data matched with those previously described in the bibliography.[2]

Step 3: Synthesis of JMLV-061 (as its hydrochloride)

i) To a stirring biphasic mixture of DCM (3 mL), saturated aqueous NaHCO$_3$ solution (3 mL) and triphosgene (101 mg, 0.34 mmol) was added 2-(trifluoromethoxy)aniline (93 µL, 120 mg, 0.68 mmol) portionwise. The mixture was then stirred at room temperature for 1 hour. The phases were separated and the organic layer was washed with brine, dried over anhydrous Na$_2$SO$_4$ and filtered. Solvent was concentrated until 1 mL of DCM was left to give 2-(trifluoromethoxy)phenylisocyanate in solution that was used in the next step without further purification.

ii) 2-(3-aminophenyl)-1,3-di-Boc-guanidine (80 mg, 0.23 mmol) and triethylamine (128 µL, 93 mg, 0.92 mmol) dissolved in DCM (2 mL) was added to the previous isocyanate solution and the mixture was stirred at room temperature for 16 hours. Saturated aqueous NaHCO$_3$ solution (15 mL) followed by DCM (10 mL) were added to the mixture and layers were separated. The organic layer was washed again with saturated aqueous NaHCO$_3$ solution (15 mL). Then, the organic layer was separated, dried over anhydrous Na$_2$SO$_4$, filtered and solvents were concentrated under vacuum to afford 1-(1,3-di-Boc-3-guanidinophenyl)-3-(2-(trifluoromethoxy)phenyl)urea that was used in the next step without further purification.

iii) A 4 M solution of HCl/1,4-dioxane (2 mL) and MeOH (0.5 mL) were added to the previously obtained crude and the mixture was stirred at room temperature for 48 hours. The solvents were concentrated under vacuum and the resulting crude was purified by column chromatography in silica gel (DCM/MeOH mixtures). Fractions containing the desired product were collected and concentrated under vacuum to afford **JMLv-061** in its dihydrochloride salt as a beige solid (39 mg, 48% yield), mp 218–219 °C. IR (ATR) ν: 624, 691, 758, 871, 1006, 1059, 1180, 1215, 1314, 1406, 1454, 1490, 1540, 1598, 1674, 2421, 3200, 3330 cm^{-1}. ^{1}H-NMR (400 MHz, CD$_3$OD) δ: 6.95 (ddd, J = 8.0 Hz, J' = 2.1 Hz, J'' = 1.0 Hz, 1 H, 4-H), 7.11 (ddd, J = 8.1 Hz, J' = 7.5 Hz, J'' = 1.6 Hz, 1 H, 4'-H), 7.29 (m, 1 H, 6-H), 7.30-7.34 (complex signal, 2 H, 3'-H, 5'-H), 7.40 (t, J = 8.0 Hz, 1 H, 5-H), 7.64 (t, J = 2.1 Hz, 1 H, 2-H), 8.20 (dd, J = 8.8 Hz, J' = 1.6 Hz, 1 H, 6'-H). ^{13}C-NMR (100.6 MHz, CD$_3$OD) δ: 116.8 (CH, C2), 119.0 (CH, C6), 120.4 (CH, C4), 122.11 (q, $^1J_{CF}$ = 256.5 Hz, C, OCF$_3$), 122.13 (CH, C3'), 123.1 (CH, C6'), 124.4 (CH, C4'), 128.7 (CH, C5'), 131.5 (CH, C5), 133.1 (C, C1'), 136.7 (C, C1), 139.9 (C, C2'), 142.1 (C, C3), 154.6 (C, CO), 158.0 (C, CN guanidine). HRMS-ESI$^+$ m/z [M+H]$^+$ calculated for [C$_{15}$H$_{14}$F$_3$N$_5$O$_2$+H]$^+$: 354.1172. Found: 354.1174. Elemental analysis calculated for C$_{15}$H$_{14}$F$_3$N$_5$O$_2$·HCl·0.6DCM: C 42.51%, H 3.70%, N 15.89%. Found: C 42.38%, H 3.90%, N 15.70%.

EPB-102

EPB-113

JMLV-061

References

1.   Guo Z-X, Cammidge AN, Horwell DC. A convenient and versatile method for the synthesis of protected guanidines. Synt. Commun. 2000;30(16):2933-2943.

2.   Kaloun EB, Schmitt P, Kruczynski, A. Isoquinolinone derivatives useful in the treatment of cancer. US9944641B2 (Patent) 2014.

S2 Appendix. Primer and probe sequences. Oligonucleotide sequences used for sequencing or (RT-)qPCR.

S2 Appendix: Primer and probe sequences

One-Step RT-PCR for sequencing of HCoV-229E: five primer pairs

Fragment 1	229E-For84	CTATGGCCGGCATCTTTGA	and 229E-Rev6270	CAAGCAGCCACCACTTAGAT
Fragment 2	229E-For5967	CAGTTGTTATGGTTGGTGGTTATG	and 229E-Rev12295	TATGTGTCCTGCGTAGTGTTG
Fragment 3	229E-For11960	GAACCACCTTGCAGATTTGTTATAG	and 229E-Rev18300	GACCCACCATTTACACCTTCA
Fragment 4	229E-For17743	GTTGCTTCTGGTGATGCTATTATG	and 229E-Rev24484	ACGTACAAGTCGTTAGTTGAGAG
Fragment 5	229E-For23936	TCAGTCGTGCTCATCTTTGTG	and 229E-Rev27281	TTTGTGTATCCATATCGAAACCGT

Sequencing of PCR fragments HCoV-229E

Fragment 1	229E-For84	CTATGGCCGGCATCTTTGA
	229E-Rev454	AACGATGCAATCCTGCAAATC
	229E-For854	CTACGCAATGGTTCTGTTCTTG
	229E-For1679	GACGTGTTTGTCACAGCTATTC
	229E-For2527	CACACAACAGGGTTTGCTTG
	229E-For3251	TCTGAATGGCCTCTTTCTGTTC
	229E-For4100	GAGAAGTTGAACGCCTTCCT
	229E-For4910	TGTGTTGTGCCATCCGAAA
	229E-For5709	ATGGTGTGCAAGTTGGTTATTG
Fragment 2	229E-For5967	CAGTTGTTATGGTTGGTGGTTATG
	229E-For6682	GATGAACTGCTTGTCACTGTTATT
	229E-For6757	CCCAGATTGTATTGCGTGTTC
	229E-For7617	CTGCTGAAGGTCGCAAGTAT
	229E-For8353	GCGTACTATTGCCACCAAATAC
	229E-For9213	CATCTGGCTTTGTGGAGAAATG
	229E-For9947	ACTGGTGTCTGTGTGGAAAG
	229E-For10793	TGTGTTAGTCCAGCCGAATTTA
	229E-For11627	GCTGGTGTTGTTTGGACATTAC
Fragment 3	229E-For11960	GAACCACCTTGCAGATTTGTTATAG
	229E-For12789	GCATGAGGGCAGAACCATTTA
	229E-For13635	GCCTGCCTTGGTGGATAAA
	229E-For14371	CGTATGTTGTCGGCTATGATCT
	229E-For15277	GCTGCTGGTCTTTGTGTAGTA
	229E-For15995	CAATGGCAAACCAGGAGAAATA
	229E-For16796	GGAGCAAGGCTGTGTTTATCT
	229E-For17590	GCTGGCTCATCTGATGTACTG
Fragment 4	229E-For17992	CGCACTGTGCAGAGTCATATT
	229E-For18811	GGTTTAACACCACCTCTCAGTAT
	229E-Rev19020	TCACAGCAGTAGCAGAGAATAAC
	229E-For19607	GGAGATGGATGCTATGGTGTAAAG
	229E-For20359	TGGCGTAACTCCACTGTTATG
	229E-For21258	GTACTGTTGCGTTAGCTTCTTATG
	229E-For22032	AGGGCACCATCTACTCTATCA
	229E-For22846	AGGTCTAACATCAGCCGTTTC
	229E-For23498	CGACAACCTAATCTTGCTCTTTAC
Fragment 5	229E-For23936	TCAGTCGTGCTCATCTTTGTG
	229E-For24698	CACCCAGCTGTTGGAATAGT
	229E-For25507	CGTGCCGAGCACTACTATAA
	229E-For26311	GTACTCCTAAGCCTTCTCGTAATC
	229E-Rev27281	TTTGTGTATCCATATCGAAACCGT

RT-qPCR for HCoV-229E

For	TTAGAGAGCGTGTTGAAGGTG
Rev	GTTCTGAATTCTTGCGCCTAAC
Probe	FAM-TCTGGGTTG/Zen/CTGTTGATGGTGCTA-3IABkFQ

RT-qPCR for SARS-CoV-2[1]

For	TTACAAACATTGGCCGCAAA
Rev	GCGCGACATTCCGAAGAA
Probe	FAM-ACAATTTGC/Zen/CCCCAGCGCTTCAG-3IABkFQ

[1]Lu X, Wang L, Sakthivel SK, Whitaker B, Murray J, Kamili S, et al. US CDC real-time reverse transcription PCR panel for detection of Severe Acute Respiratory Syndrome Coronavirus 2. Emerg Infect Dis. 2020;26(8):1654-65. doi: 10.3201/eid2608.201246.

qPCR for innate immune factors

hACTB_for	CACCATTGGCAATGAGCGGTTC
hACTB_rev	AGGTCTTTGCGGATGTCCACGT
hHPRT_for	CATTATGCTGAGGATTTGGAAAGG
hHPRT_rev	CTTGAGCACACAGAGGGCTACA
hRIGI_for	CACCTCAGTTGCTGATGAAGGC
hRIGI_rev	GTCAGAAGGAAGCACTTGCTACC
hMDA5_for	GCTGAAGTAGGAGTCAAAGCCC
hMDA5_rev	CCACTGTGGTAGCGATAAGCAG
hMAVS_for	ATGGTGCTCACCAAGGTGTCTG
hMAVS_rev	TCTCAGAGCTGCTGTCTAGCCA
hIRF3_for	TCTGCCCTCAACCGCAAAGAAG
hIRF3_rev	TACTGCCTCCACCATTGGTGTC
hIRF7_for	CCACGCTATACCATCTACCTGG
hIRF7_rev	GCTGCTATCCAGGGAAGACACA
hIFNB_for	CTTGGATTCCTACAAAGAAGCAGC
hIFNB_rev	TCCTCCTTCTGGAACTGCTGCA
hOAS1_for	AGGAAAGGTGCTTCCGAGGTAG
hOAS1_rev	GGACTGAGGAAGACAACCAGGT
hOAS2_for	GCTTCCGACAATCAACAGCCAAG
hOAS2_rev	CTTGACGATTTTGTGCCGCTCG
hOAS3_for	CCTGATTCTGCTGGTGAAGCAC
hOAS3_rev	TCCCAGGCAAAGATGGTGAGGA
hRNaseL_for	AAGGCTGTTCAAGAACTACACTTG
hRNaseL_rev	TGGATCTCCAGCCCACTTGATG
hPKR_for	GAAGTGGACCTCTACGCTTTGG
hPKR_rev	TGATGCCATCCCGTAGGTCTGT

S1 Fig. Particle size distribution of SEC-purified nsp15 samples

Particle size distribution of HCoV-229E and SARS-CoV-2 nsp15 as determined by dynamic light scattering (DLS). Cuvette-based DLS was performed on SEC-purified 0.5 mg/ml nsp15 solutions in storage buffer (25 mM HEPES at pH 8, 300 mM NaCl, 10% glycerol, and 1 mM tris(2-carboxyethyl)phosphine), using a DynaPro Nanostar and Dynamics software v7.10.1.21 (Wyatt Technologies). 15 µl of sample was mixed 5 µl water. The sample chamber was kept at 20 °C. The sample contained ≥ 98% particles with a diameter of ~13 nm, consistent with the size of hexameric nsp15, for both HCoV-229E (polydispersity index: 9%) and SARS-CoV-2 (polydispersity index: 11.7%).

S2 Fig. Synergism analysis of EPB-113, 5h and GS-441524.

To study drug combinations, we used an alternative CPE reduction assay with HCoV-229E in HEL299 cells. An 8x8-well matrix was created, each well containing different concentrations of two compounds. Data analysis was performed with the zero-interaction potency (ZIP) model in SynergyFinder+. Mean ZIP-scores are shown on top of each graph. A ZIP-score lower than -10 indicates antagonism, between -10 and 10 corresponds with additive effects, and higher than 10 indicates synergism. The combination of EPB-113 with 5h is synergistic (A), while the combinations of EPB-113 with GS-441524 (B) and 5h with GS-441524 (C) are additive.

A

mRNA
(log_{10} increase vs mock)

	RIG-I				MDA5				MAVS				IRF3				IRF7				IFN-β				OAS1				OAS2				OAS3				RNase L				PKR			
WT	0.2	-0.2	1.4	1.2	-0.1	0.1	1.0	0.7	-0.1	0.2	0.2	0.2	0.0	0.0	-0.2	-0.5	0.0	0.1	0.7	0.6	0.2	0.1	0.7	0.9	0.1	0.2	2.2	1.6	-0.1	0.2	1.8	1.3	0.0	0.3	0.9	1.1	0.0	0.1	0.5	0.3	0.0	0.1	0.6	0.3
G248V	0.3	0.1	2.9	2.7	0.0	0.2	2.5	1.9	0.1	0.1	0.4	0.0	0.1	0.1	0.2	-0.1	0.1	0.1	2.2	1.8	-0.2	0.6	3.4	3.7	0.1	1.1	4.3	3.6	0.1	0.6	3.9	3.2	0.1	0.1	2.6	2.5	0.0	0.1	0.7	0.3	0.0	0.1	1.4	0.8
			****	****			****	****								****			****	****			****	****			****	****			****	**			****	****			*				****	****
H250Y	0.5	0.6	2.9	2.6	0.1	0.5	2.6	2.1	0.2	0.2	0.3	-0.2	0.0	0.1	0.2	-0.1	0.3	0.4	2.4	2.0	0.0	1.2	3.9	3.9	0.9	2.2	4.4	3.6	0.5	1.6	4.0	3.3	0.2	0.4	2.8	2.8	0.0	0.1	0.5	0.1	0.1	0.2	1.4	0.9
			****	**			*	****								****			****	****			****	****			****	****			**	***			*	****							****	****
H250A	0.4	0.1	2.3	2.9	0.1	0.1	1.9	2.3	0.0	0.0	0.0	0.0	0.1	0.0	0.1	-0.1	0.3	0.1	1.5	1.7	0.2	0.5	2.6	4.1	0.7	1.1	3.7	3.7	0.5	0.7	3.3	3.3	0.1	0.1	1.9	2.4	0.0	0.4	0.0	0.0	0.0	0.2	0.9	0.9
			*	****			****	****							**	****			****	****			**	****			****	****			***	***			****	****			*				***	****
K60R	0.0	0.3	2.5	2.2	0.0	0.0	2.2	1.7	-0.1	0.0	0.4	0.0	0.0	0.0	-0.1	-0.4	0.0	0.1	1.8	1.6	0.0	0.5	1.9	2.3	0.1	0.8	3.8	3.2	0.2	0.4	3.4	2.8	0.1	0.0	2.2	2.2	0.0	0.1	0.5	0.3	0.0	0.0	1.1	0.8
			**	*			****	****											****	****			*	*			****	****			***	**			****	****							****	****
T66I	0.1	0.0	2.3	2.4	0.0	0.0	2.1	1.8	0.0	0.0	0.4	0.0	0.0	0.0	-0.1	-0.3	0.0	0.1	1.8	1.7	0.4	0.2	2.4	3.2	0.2	0.8	3.7	3.4	-0.1	-0.3	3.4	2.9	0.1	0.1	2.2	2.3	0.0	0.1	0.6	0.1	0.0	0.1	1.1	0.9
			*	**			****	****								*			****	****			***	****			****	****			***	***			****	****							****	****
	2	24	48	72	2	24	48	72	2	24	48	72	2	24	48	72	2	24	48	72	2	24	48	72	2	24	48	72	2	24	48	72	2	24	48	72	2	24	48	72	2	24	48	72

Time p.i. (h)

B

IFNα/β protein
(fold increase in A_{620} vs mock)

	16HBE					HEL					Macrophages			
WT	0.9	1.0	2.1	1.4		0.9	0.9	0.9	1.2		0.8	1.0	10	8.2
G248V	0.9	1.1	3.1	2.0		0.8	1.0	12	11		0.9	2.5	3.9	2.7
			***					****	****				****	****
H250Y	1.0	1.2	3.4	1.9		0.8	1.0	12	11		1.0	5.6	7.7	6.8
			****					****	****			**		
H250A	1.0	1.0	2.3	1.9		0.9	1.0	7.0	11		1.0	1.8	2.2	1.5
								*	****				****	****
K60R	1.1	1.0	2.7	1.7		0.8	1.0	3.5	5.3		0.9	1.4	13	12
														**
T66I	1.0	1.0	2.5	2.0		0.8	1.0	4.8	8.3		0.8	1.3	15	12
									**				**	**
	2	24	48	72		2	24	48	72		2	24	48	72

Time p.i. (h)

S3 Fig. Numeric values used to create the heatmap in Fig. 4, and statistical analysis of the data.

S4 Fig. EPB-113, nirmatrelvir and GS-441524 do not increase IFN-β mRNA levels in HEL299 cells when compound and HCoV-229E virus are added simultaneously.