The heat is on: behavioural, physiological and reproductive evidence of heat
stress in breeding king penguins.

Aude Noiret[1,2*], Agnès Lewden[3], Camille Lemonnier[1], Céline Bocquet[1], Marine Montblanc[1],
Fabrice Bertile[1], Marine Hoareau[1], Elsa Marçon[1], Jean-Patrice Robin[1], Pierre Bize[4,5], Vincent
A Viblanc[1] & Antoine Stier[1,2,6*]

[1] Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

[2]Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés, CNRS UMR 5023 Université
Claude Bernard Lyon 1, Université de Lyon, Lyon, France.

[3] Université de Brest - UMR 6539 CNRS/UBO/IRD/Ifremer, Laboratoire des sciences de
l'environnement marin - IUEM - Rue Dumont D'Urville - 29280 - Plouzané

[4] Swiss Ornithological Institute, CH-6204 Sempach, Switzerland

[5] Department of Ecology and Evolution, University of Lausanne, CH-1015 Lausanne, Switzerland

[6] Department of Biology, University of Turku, Turku, Finland

* corresponding authors: aude.noiret@gmail.com & amstie@utu.fi

Running Title: Heat stress in king penguins breeding on land

Key words: thermoregulation, temperature, seabird, climate change, behaviour

Author's contribution

Study design: AS, VAV, AL. Funding acquisition: AS, JPR, PB, VAV. Data collection in the field:
CL, CB, MM, MH, FB, EM, VAV, AS. Data analysis: AN. Writing original draft: AN and AS.
Writing review and editing: AL, CL, CB, MM, MH, FB, EM, JPR, PB, VAV.

Data availability

The datasets and script are available on *FigShare* (Noiret et al. 2024). HotKing dataset:
heat   stress   in   breeding   king   penguins.   figshare.   Dataset.
https://doi.org/10.6084/m9.figshare.26968723).

A   preprint   is   available   on   BioRxiv   at
https://biorxiv.org/cgi/content/short/2024.09.09.611977v1.

Number of words: 8391

Number of references: 73

Number of figures: 4; number of tables: 1; SEM: 9

© IPEV – A. Stier

© IPEV – A. Noiret

Abstract

Polar and sub-polar animals evolved to thrive in cold climates and may thus be
particularly vulnerable to rising temperatures associated with climate change. Penguins may
be especially vulnerable due to their dual habitat, alternating between foraging in cold waters
and breeding/moulting on an increasingly warm land. Here, we characterized heat stress
occurrence in breeding king penguins through behavioural observations (*e.g.* panting
occurrence) and body temperature measurements. We observed that behavioural signs of
heat stress are frequent in king penguins breeding in the sub-Antarctic region (> 20% of
observations at mid-day), and that subcutaneous temperatures increase under high heat load,
especially in penguins showing behavioural signs of heat stress. Subcutaneous and core body
temperatures were moderately correlated and both increased with heat load. Yet, their
responses were not parallel since core body temperature is markedly less sensitive to heat
load than subcutaneous temperature. Air temperature alone was a poor predictor of heat stress
occurrence, whereas the combination of high solar radiation, low wind speed and high air
temperatures provided the strongest predictive power. Finally, reproductive failures were more
likely to occur on warmer days, suggesting that heat stress may have significant sublethal
effects that could ultimately affect population dynamics.

Introduction

As climate is warming, it is expected to exert direct thermoregulatory pressure on most living organisms (Piatt *et al.* 2024; Stillman 2019). Although heat waves - which are known to cause mass mortality events amongst wild animals (Conradie *et al.* 2020) – are defined on different criteria between studies (Gimeno *et al.* 2025; Robinson 2001; Ton *et al.* 2021), climate can be defined as "hot" as soon as it triggers responses of the organism to maintain or restore body temperature and homeostasis, an energetically costly mechanism involving physiological and behavioural changes, here defined as heat stress (Hansen 2009). Indeed in endotherms, when ambient temperatures reach the upper limits of their thermoneutral zone (TNZ) (*i.e.* the range of ambient temperatures in which metabolic rate is minimal) (Romanovsky 2007, 2018), the costs of active thermoregulation or the failure to properly thermoregulate can impair other bodily functions of the organism (Boyles *et al.* 2011; Danner *et al.* 2021; Pipoly *et al.* 2013), eventually leading to sublethal to lethal effects in extreme situations. However, as climatic projections describing global warming are based primarily on air temperatures (T_{air}), the heat load faced by wild animals may be underestimated, as it generally depends on multiple climatic parameters, including T_{air}, but also solar radiation (SR), wind speed (WS) and humidity (Mitchell 2024). When available, such data can be used to compute various thermal indexes (Ioannou *et al.* 2022), which may help to refine the determination of the threshold beyond which an animal could be impacted by "hot" conditions, especially in a complex natural environments.

Studies on heat stress occurrence in wild animals are mostly conducted in species from arid, tropical or temperate regions, because of obvious extreme ambient temperatures (Bryant 2008; Conradie *et al.* 2020). Yet, cold-adapted species may be at greater risks of struggling under hot conditions because of a narrower, cold-shifted TNZ and thermoregulatory mechanisms specialized for cold environments (Beaman *et al.* 2024; Choy *et al.* 2021; Grémillet *et al.* 2012; O'Connor *et al.* 2021). Even if maximum temperatures in polar regions are seen as less extreme than in other parts of the globe, polar regions are warming at a faster rate than any other place on earth (Clem *et al.* 2020; Stammerjohn & Scambos 2020; You *et al.* 2021). Hence, animals adapted to cold environments may reach a breakpoint in thermoregulation efficacy more rapidly than other species in response to climate anomalies (Beaman *et al.* 2024; Choy *et al.* 2021; O'Connor *et al.* 2021).

Seabirds from polar and subpolar regions are expected to be particularly sensitive to heat stress, because they have to maintain a thick body insulation in order to forage in cold waters (Davis & Darby 2012), but are increasingly exposed to high temperatures on land (Gimeno *et al.* 2025). It has been suggested that since marine birds have an "easy access" to water to cool down, they are unlikely to suffer much from hot conditions on land (Beaman *et*

al. 2024). Yet, this assumption does not consider the high constraints that most seabirds face

during reproduction, which is also the timeframe where the heat load is expected to be the

highest. For instance, thermoregulatory behaviours such as bathing would directly impact nest

attendance, thereby increasing the egg or chick exposure to heat and predators (Olin *et al.*

2023). Direct and indirect (*e.g.* longer foraging times, Bost *et al.* 2015) effects of heat may

contribute to decrease reproductive success, leading to a trade-off between thermoregulation

and reproduction (or even survival) in breeding adults.

Direct observation of thermoregulatory behaviours is an efficient and non-invasive way

to study heat stress in free-living animals (Olin *et al.* 2023). Panting is considered a sign of

heat stress as it is an energy-expensive evaporative cooling mechanism (Tan & Knight 2018).

Panting behaviour during reproduction has been observed on days when air temperatures

were high, for example in nesting guillemot (Olin *et al.* 2023), great skuas (Oswald *et al.* 2008)

and African penguins (Welman & Pichegru 2022). The use of more passive thermoregulatory

behaviours (*i.e.* less energetically costly than panting) has also been described in breeding

seabirds, in order to increase convective and/or conductive heat loss for instance through

spreading wings (Olin *et al.* 2023), bathing (Oswald *et al.* 2008), crouching or standing (Cook

et al. 2020). To evaluate if thermoregulatory behaviours are sufficient to maintain

normothermia, various methods exist for measuring body temperature in wild animals (for a

review, see McCafferty *et al.* 2015). For animals being identified using sub-cutaneous pit-tags,

the replacement of traditional pit-tag with thermosensitive ones offers the opportunity to collect

subcutaneous temperature (T_{sc}) while minimizing invasiveness (Andreasson *et al.* 2023;

McCafferty *et al.* 2015). However, T_{sc} is measured close to the surface of the body and is

known to be more variable than core body temperature (T_b). Furthermore, the relationship

between T_{sc} and T_b is also known to vary between species, and with the location of the sensor

and environmental conditions (Lewden *et al.* 2017; Romanovsky 2018). Considering that

measuring T_{sc} is usually less invasive and expensive than T_b, it enables working with larger

sample sizes, but requires prior validation in any given species and environmental context.

Here, we hypothesized that king penguins *(Aptenodytes patagonicus)*, the second

largest penguin species, might be particularly sensitive to heat due to adaptation to cold

waters, with effective body insulation for deep-diving (Davis & Darby 2012). Modifying this

insulation during breeding is impossible, as penguins must alternate between

incubating/brooding on land during spring/summer at fluctuating ambient temperatures (see

ESM S1, numerous days with maximum temperature > 15°C at our study site), and foraging

at sea in *ca.* 4°C water (Stonehouse 1960). The risks of heat stress are further accentuated as

breeders reproduce on volcanic beaches with no access to shade, and must remain on land

while fasting for prolonged periods, around 2-3 weeks during egg incubation and around 1

week during chick rearing (Weimerskirch *et al.* 1992). During their stay on land, king penguins

breed in a dense colony, establishing a small territory against aggressive conspecifics (Côté

2000; Lemonnier *et al.* 2024) from which they cannot move or leave their egg/chick unattended

without very high risks of reproductive failure due to predation (Descamps *et al.* 2005). This

further accentuates the risk of heat stress by limiting access to cooling substrate (*i.e.* water)

and triggering frequent antagonistic interactions(Viblanc *et al.* 2012). Although behavioural

signs of heat stress have been observed in polar and sub-polar penguin species (Stier &

Lewden, *pers. obs.*), this phenomenon remains to be investigated in details.

In this study, we investigated the occurrence of heat stress in free-living king penguins

and, by extension, assessed whether this species can serve as an early sentinel of the risk

that sub-polar and polar seabirds face of experiencing land-based heat stress during

reproduction in the context of climate change. To this end, we examined: (1) the occurrence

of heat stress by studying thermoregulatory behaviours and the biological (*i.e.* breeding stage,

sex and fasting duration) and climatic (*i.e.* ambient temperature, solar radiation, humidity and

wind speed) factors triggering heat stress; (2) the relationship between subcutaneous

temperature and core body temperature under increasing heat load (3) the relationship

between thermoregulatory behaviours and subcutaneous temperature under hot climatic

conditions; (4) the performance of different thermal indexes in predicting heat stress

occurrence; and (5) the climatic conditions associated with reproductive failure.

Materials and Methods

Study model and site.

We studied king penguins breeding in the "Baie du Marin" colony on Possession Island in the Crozet Archipelago (46°25′S; 51°52′E). Data were collected over two breeding seasons, 2021-2022 and 2022-2023. During breeding, male and female partners alternate between periods of foraging at sea with periods on land caring for the egg or chick (Figure S1). After laying a single egg, females return to forage at sea while males take care of the first incubation shift. Alternated guard continues throughout incubation (*ca.* 53 days) and brooding until the chicks reach emancipation (*ca.* 35 days) (Davis & Darby 2012). In this study, we monitored 72 breeding pairs (20 during 2021-2022 and 52 during 2022-2023). For each breeding pair, male and female parents were individually marked during courtship using hair-dye (Schwarzkopf Palette Deep Black) and monitored daily throughout incubation and brooding. The number of days on land (D_{land}), from the return of the birds on land until the day of observation, was also considered in the analysis. The first incubation shift of males was excluded from the analysis as they fast 2-3 weeks longer than females, which influences greatly their physiology (Ancel *et al.* 2013).

Climatic parameters

Climatic data were provided by Météo France through a weather station located *ca.* 1km from the penguin colony (Station 98404004, 46°25'S; 51°51'E). The station provided the air temperature under shelter (T_{air}, °C), the hourly global solar radiation (SR, J/cm²), wind speed (WS, m/s), and relative humidity (RH, %). It also provided the dew-point temperature (Tw, °C). Although water vapor pressure is better than relative humidity to estimate the heat load and thermoregulatory constraints on an animal (Mitchell et al., 2024), it is highly colinear with temperature, and both cannot be included in the same model, contrary to RH. As our aim is to include the different meteorological variables within the same model, we thus used RH.

We used 4 different heat indexes to compare their predictive power to models including raw meteorological variables. These indexes are either practical metric of dry heat load to which the biological responses of wild animals should be assessed (T_g, Mitchell et al. 2024), the two most used indexes to predict heat stress occurrence in humans (THI, WBGT, Budd 2008), or two alternative indexes thought to better predict heat stress in humans and farm animals (HLI and WBGT; Carter *et al.* 2020; Liljegren *et al.* 2008; Silva *et al.* 2007). First, we estimated the globe temperature (T_g) from climatic parameters provided by the weather station using an equation derived from field measurement of T_g in our penguin colony (T_g, equation 1, see Figure S2 for details). The 3 other heat indexes we used were the Temperature-Humidity Index (THI, equation 2, Silva *et al.* 2007; Thom 1959), the Heat Load Index (HLI, equation 3,

178 Gaughan 2002; Silva *et al.* 2007) and the Wet-Bulb Globe Temperature (WBGT, equation 4,
179 Hajizadeh *et al.* 2017; Liljegren *et al.* 2008).

(1) $Tg = 2.89 + 0.83 \times Tair - 0.035 \times WS + 0.036 \times SR$ (*see ESM S2*)

(2) $THI = Tair + 0.36\,Tw + 41.5$ *where T_w is the dew-point temperature*

(3) $HLI = 33.2 + 0.2\,RH + 1.2\,Tg - (0.82\,WS)\,0.1 - \log(0.4\,WS^2 + 0.0001)$

(4) $WBGT = 0.7\,Tnwb \times 0.2\,Tg \times 0.1\,Tair$ *where T_{nwb} is the natural wet bulb temperature (for full equation, see Carter et al., 2020)*

Thermoregulatory behaviours

Penguins' thermoregulatory behaviours were recorded using scan sampling every 3
days over a period of 40 minutes with an observation recorded every 2 minutes (maximum 21
observations per scan). The scans were performed during midday (mean= 14:35 ± 1 hour and
35 minutes), when the heat load would be the highest, and to limit potential effects linked to
circadian variation in behaviour. We gathered a total number of 1175 scans, scattered over 60
days (from 28[th] November 2021 to 8[th] March 2022 and 4[th] December 2022 to 26[th] February
2023). The total number of scans per individual (mean of 9 ± 3 scans) varied with the duration
of their shift on land ('D_{land}') and the survival of their offspring. We focused on 3 behaviours
considered to have a thermoregulatory function, *i.e.* panting, wing spreading and exposure of
the brood pouch (*i.e.* a highly vascularized and featherless area that may serve as a thermal
window to dissipate heat; Handrich *et al.* 1997), and analysed their occurrence in terms of
presence/absence during the scan (1/0 respectively). Observations were carried out by 11
observers during 2 field seasons.

Body temperature measurements

We recorded sub-cutaneous temperature (T_{sc}, °C) using thermosensitive transponders
or "Pit-tags" (BioTherm13 tag, resolution ± 0.1°C, accuracy ± 0.5°C, between 33°C and 43°C,
Biomark, USA) used for individual identification before the beginning of the study and placed
in the upper back of all individuals (Oswald *et al.* 2018). T_{sc} was recorded every 3 days during
shifts on land (n = 856), using the Biomark HPR Plus Reader just after behavioural scans and
without handling of the animals. In our study species, T_{sc} changes have been shown to mirror
changes in core body temperature relatively well while on land (Eichhorn *et al.* 2011; Lewden
et al. 2017). Yet, to determine the extent to which T_{sc} can be informative of core body
temperature (T_b) in king penguins under high heat load, and thus of their actual
thermoregulatory capacity, we used another set of data where T_{sc} was measured alongside T_b
on 14 adults (7 males and 7 females, n = 37 observations) that are not part of the behavioural
study. T_b was measured as stomach temperature using Anipill loggers (Bodycap®, Anipill,

France, resolution ± 0.01°C, accuracy ± 0.2°C between 25 and 45°C) that were fed to the animals (Zuluaga *et al.* 2025). We estimated the raw correlation between T_b and T_{sc} simultaneously, then calculated their difference (Diff = T_b - T_{sc}, °C), and performed a Bland-Altman analysis. To further investigate the implication of the different climatic variables on T_b and T_{sc} variations, we paralleled the data with the climatic variables corresponding to the same time and date (T_{air}, SR, WS and RH). We tested the effect of climatic variables on T_{sc}, T_b and on their difference using linear mixed models (with individual identity as random intercept).

Reproductive failure

Survival of the egg or chick was monitored daily throughout the entire breeding season, and the day of egg abandonment or chick disappearance was recorded. The exact cause of reproductive failure (*e.g.* predation, parental abandonment) remained unknown in the vast majority of cases.

Statistical analysis.

Our behavioural observations and T_{sc} of breeding penguins were merged with climatic variables by the closest time of the day, with a maximum delay of 30 minutes. The statistical analyses were performed in R version 4.3.2 (R Core Team 2023). The effects of T_{air}, SR, WS and RH on the probability of displaying the 3 thermoregulatory behaviours during the scans were analysed using generalised linear mixed models (R package 'lme4', version 1.1-35.1; Bates *et al.* 2015), assuming a binomial distribution (link=logit) and after scaling (*i.e.* z-transformation) of the climatic parameters. Linear mixed models with a normal distribution were used to assess the effects of climatic parameters on T_{sc}. As individuals were measured several times over the breeding season, bird identity was included as a random intercept, as well as the identity of the observers (except for T_{sc}, recorded by a pit-tag reader and therefore not subject to inter-observer variation). Year (2021-22 vs. 2022-23), sex (Male vs. Female), reproductive stage (Incubation vs. Brooding) and day of the on-land shift ('D_{land}') were included as fixed effects in the models. T_{sc} was further analysed depending on the presence or absence of panting behaviour for each individual during the preceding behavioural scan. Breakpoint analysis of T_{sc} against T_{air}, SR, T_g and HLI, was performed using the *segmented* R package (version 2.0-2; Muggeo 2008). Segmented models were fitted from linear regressions without accounting for inter-individual variation of T_{sc}, as not all individuals had enough data points to enable a mixed model breakpoint analysis. A Davies test was used to test for the significance of a change in slope in each linear model (Muggeo 2008).

Models including all variables but no interaction terms (*i.e.* No-interaction models) were compared to two-ways interaction models between climatic variables (*e.g.* T_{air}*SR or WS*RH), and to a null model, using corrected Aikaike's information criterion (AICc). The models with

248 interaction terms presented in the results are the models with the lower AICc. To assess model

249 fit and statistical assumptions, 'DHARMa' package was used (version 0.4.6; (Hartig 2016).

250 Variance inflation factors from the No-interaction models were all under 3, suggesting no

251 substantial bias from collinearity between predictor variables (namely climatic variables)

252 occurred (see Zuur *et al.* 2010). Conditional and marginal R^2 were used to compare the

253 predictive power of each model (*i.e.* No-interaction model, Interaction model, heat indexes).

254 We tested for a difference in climatic conditions during days in which egg or chick

255 mortality was observed, compared to days where no reproductive failure was recorded. For

256 this, we used generalized models of the binomial family with the proportion of dead offspring

257 each day depending on maximum T_{air}, maximum SR, T_g, and HLI. Maximum daily values were

258 used since the exact time of day of reproductive failure was unknown, and maximum heat load

259 was predicted to be more relevant than daily average. We also used non-parametric Wilcoxon

260 tests to check for a difference in climatic condition between days with or without reproductive

261 failure being observed.

Results

Behavioural thermoregulation

Figure 1. Predicted probabilities of thermoregulatory behaviours (panting (brown), wing spreading (blue) and exposure of the brood pouch (red)) occurrence according to (A) solar radiation, (B) wind speed, (C) air temperature and (D) relative humidity. Predicted probabilities were calculated from generalized linear mixed models; N = 144 individuals, n = 1175 observations.

Independently of climatic conditions, panting was the most frequently observed thermoregulatory behaviour (23.4% of behavioural scans included panting behaviour), followed by wing spreading (9.7%) and exposure of the brood pouch (7.7%). Over the 60 days with behavioural scans, there were 27 days where at least one individual was observed panting (26 days for wing spreading and 16 days for brood pouch exposure).

The probability of observing panting behaviour was mainly explained by solar radiation (SR) and wind speed (WS), while air temperature (T_{air}) and relative humidity (RH) did not significantly impact panting as main effects (Table 1A, Fig 1). Four two-ways interactions significantly explained panting probability, with the most notable being the interaction between T_{air} and SR, revealing that panting only increased with increasing T_{air} when SR were moderate

to high (Fig 2A; see ESM S3 for details on other interactions). Males tended to display panting behaviour more frequently than females (p = 0.067), but the probability to observe panting behaviour did not significantly vary with reproductive stage or between the two breeding seasons we studied (Table 1A). Panting probability significantly decreased with the time spent on land (D_{land}), meaning that the birds' susceptibility to suffer from heat stress gradually decreased after returning from the sea (Table 1A, ESM S4 A).

Figure 2. Predicted probability of (A) Panting depending on air temperature and solar radiation; solar radiation is represented as maximum (380), median (264) and minimum (24) registered during behavioural scans; (**B) Wing spreading according to air temperature and wind speed** (2nd quartile (4.65), median (9.18), 3rd quartile (13.71))**; (C) Brood pouch exposure according to air temperature and humidity** (2nd quartile (67), median (73), 3rd quartile (84)). Predicted probabilities were calculated from generalized linear mixed models; N = 143 individuals, n = 1123 observations.

The probability of observing wing spreading increased under high solar radiation, high T_{air} and low wind speed, while relative humidity had no significant impact on this behaviour (Fig 1, Table 1B). Three two-ways interactions also significantly explained wing spreading probability, with the most notable being the interaction between T_{air} and WS (Table 1B), revealing that wing spreading increased with increasing temperature when wind speed was low to moderate (Fig 2B; see ESM S3 for details on other interactions). Males significantly spread their wings more often than females (p = 0.015) and brooding birds more than incubating ones (p < 0.001), but the probability to spread wings did not significantly change along the time spent on land nor between the two breeding seasons we studied (Table 1B).

The probability of observing the brood pouch lifted increased with SR, T_{air}, RH and decreased with WS (Fig 1, Table 1C). The interaction between T_{air} and RH was also significant, revealing that brood pouch exposure only increased with increasing T_{air} when RH was low (Table 1C, Fig 2C). Sex, breeding season and the time spent on land did not significantly influence the probability of exposing the brood pouch, but brooding individuals were more likely to exhibit this behaviour than incubating ones (p < 0.001; Table 1C).

	(A) Panting			(B) Wing spreading			(C) Brood pouch exposure			(D) Tsc		
	OR	CI	p	OR	CI	p	OR	CI	p	Est.	CI	p
(Intercept)	0.15	0.07 – 0.32	**<0.001**	0.07	0.02 – 0.20	**<0.001**	0.00	0.00 – 0.03	**<0.001**	37.71	37.50 – 37.92	**<0.001**
T_{air}	1.26	0.82 – 1.94	0.283	1.96	1.53 – 2.50	**<0.001**	1.55	1.03 – 2.33	**0.037**	0.08	0.00 – 0.15	**0.037**
RH	1.42	0.99 – 2.03	0.059	0.89	0.53 – 1.49	0.666	2.53	1.63 – 3.93	**<0.001**	0.09	0.02 – 0.16	**0.017**
SR	4.32	2.84 – 6.55	**<0.001**	4.15	2.75 – 6.28	**<0.001**	7.98	4.25 – 14.96	**<0.001**	0.45	0.39 – 0.52	**<0.001**
WS	0.27	0.18 – 0.40	**<0.001**	0.54	0.37 – 0.81	**0.003**	0.24	0.14 – 0.41	**<0.001**	-0.32	-0.39 – -0.26	**<0.001**
Season [2]	1.57	0.77 – 3.21	0.216	0.46	0.20 – 1.09	0.079	5.14	0.97 – 27.34	0.055	-0.05	-0.23 – 0.13	0.595
Sex [M]	1.57	0.97 – 2.54	0.067	1.81	1.12 – 2.92	**0.015**	0.71	0.39 – 1.29	0.259	0.09	-0.06 – 0.24	0.241
D_{land}	0.84	0.79 – 0.89	**<0.001**	1.00	0.94 – 1.06	0.996	0.98	0.91 – 1.06	0.635	-0.04	-0.05 – -0.02	**<0.001**
Repro. St [Incub.]	0.96	0.61 – 1.49	0.845	0.32	0.19 – 0.55	**<0.001**	0.12	0.06 – 0.25	**<0.001**	-0.20	-0.33 – -0.07	**0.002**
$T_{air} \times RH$	0.34	0.22 – 0.52	**<0.001**				0.37	0.23 – 0.58	**<0.001**			
$T_{air} \times SR$	1.57	1.04 – 2.38	**0.033**							0.25	0.18 – 0.31	**<0.001**
$T_{air} \times WS$				1.69	1.12 – 2.57	**0.013**				-0.22	-0.31 – -0.14	**<0.001**
SR × WS	0.67	0.44 – 1.03	0.068									
SR x RH	2.06	1.34 – 3.15	**0.001**	1.98	1.14 – 3.45	**0.016**						
WS x RH	1.82	1.21 – 2.73	**0.004**	1.80	1.09 – 2.94	**0.020**						
Random effects												
σ^2	3.29			3.29			3.29			0.68		
$T_{00\ ID\ bird}$	0.46			0.00			0.05			0.07		
$T_{00\ ID\ observer}$	0.09			1.18			7.31					
ICC	0.14						0.69			0.10		
$N_{ID\ bird}$	143			143			143			140		
$N_{ID\ observer}$	11			11			11					
N Obs.	1123			1123			1123			856		
R^2m / R^2c	0.684 / 0.729			0.646 / NA			0.479 / 0.839			0.369 / 0.430		

Relationships between sub-cutaneous temperature, core body temperature and climatic conditions

In a subsample of king penguins (N = 14, n = 37) where both T_{sc} and T_b were simultaneously measured, T_{sc} was significantly correlated with T_b (r = 0.62, p <0.001, Figure S5A). The Bland–Altman analysis indicated that the two measures presented a nearly significant mean bias of 0.24°C (95% CI: –0.03 to 0.5, t = 1.81; p = 0.079), indicating that T_{sc} slightly underestimated T_b, although not significantly (ESM S5B). However, the difference between T_b and T_{sc} significantly decreased with increasing body temperature (estimate of Difference ~ Mean = -0.88, p <0.001; ESM S5B). The difference between T_b and T_{sc} significantly decreased with increasing T_{air} and SR (Table S6A and S6B, ESM S5C and D). Importantly, both T_{sc} and T_b showed the same pattern of increasing significantly with increasing T_{air} and SR (Table S6A and S6B, ESM S5C and S5D). Wind speed and relative humidity did not affect T_{sc}, T_b, nor their difference (Table S6C and S6D, ESM S5E and S5F).

Relationships between subcutaneous temperature, heat load and thermoregulatory behaviour

Figure 3. Breeding king penguins' sub-cutaneous temperatures (T_{sc}) according to (A) the interaction between air temperature (T_{air}) and solar radiation (SR), and (B) air temperature and wind speed (WS). Values are predicted from the linear mixed model presented in Table 1D; *N = 140, n = 856.*

Subcutaneous temperature (T_{sc}) increased under high T_{air}, SR and RH, and decreased in response to high WS (Table 1D). The best predictive model included significant interactions between T_{air} and SR and between T_{air} and WS (Figure 3, Table 1D), revealing that king penguin's T_{sc} was especially elevated when T_{air} and SR increased concomitantly, along with a decrease in WS. Males and females did not show significant differences in their T_{sc}, but brooding individuals had a higher T_{sc} than incubating ones (p = 0.002; Table 1D). T_{sc} also significantly decreased with days on land (p < 0.001, Table 1D, ESM S4B).

Breakpoint analyses showed that T_{sc} significantly increased with T_{air} when a threshold
of 16.0 ± 1.1 °C was reached (Davies test: p = 0.005; ESM S7A), when SR reached 125.3 ±
11.7 J.cm-² (Davies test: p < 0.001; ESM S7B), or when the heat load index (HLI) was at 65.5
± 1.4 (Davies test: p = 0.002; ESM S7D). However, the breakpoint of T_{sc} with increasing T_g was
not significant (12.4 ± 0.6 °C; Davies test: p = 0.13; ESM S7C).

T_{sc} was significantly higher in panting than non-panting individuals (37.8°C vs. 37.2°C,
χ^2= 48.1, p < 0.001), and increased with T_{air} and SR in panting individuals while this was not
the case or less pronounced for non-panting animals (χ^2 Panting*T_{air} = 7.5; p = 0.006; Figure
4A; χ^2 Panting*SR = 14.3; p < 0.001; Figure 4B). Moreover, the increase of WS was not
sufficient to decrease T_{sc} in panting individuals compared to non-panting birds (χ^2 Panting*T_{air}
= 10.2; p = 0.001; Figure 4C). Relative humidity had no effect on T_{sc} as a main effect, but the
T_{sc} of panting individuals slightly decreased with increasing RH (χ^2 Panting*RH = 4.3; p = 0.038
Figure 4D).

Figure 4. Relationship between air temperature (A), solar radiation (B), wind speed (C) or relative humidity
(D) and king penguin's sub-cutaneous temperature according to the use of panting as a thermoregulatory
behaviour. *N = 140, n = 856.*

Evaluating the relevance of T_{air} and heat indexes in predicting heat stress occurrence

Models only including T_{air} poorly explained the occurrence of thermoregulatory

behaviours or variation in T_{sc}, with R² ranging from 0.07 to 0.23 (Table S8). Solar radiation

always performed better than T_{air} in predicting heat stress occurrence, with $R^2 > 0.24$ (Table

S8). Models containing all 4 parameters (T_{air}, RH, SR and WS) as main effects (R² ranging

from 0.30 to 0.67) and models including their two-ways interactions (R² ranging from 0.37 to

0.68) were the best predictive models for all explained variables (Table S8). Models including

heat indexes (Globe temperature (T_g), Temperature-Humidity (THI), Heat Load Index (HLI) and

Wet-Bulb Globe Temperature (WBGT) explained less variance in thermoregulatory behaviours

and T_{sc} than models (with or without interactions) including T_{air}, RH, SR and WS as explanatory

variables (Table S8). Among heat indexes, T_g (average R²m = 32.7%) and HLI (average R²m

32.0%) exhibited overall the best performance, while WBGT (average R²m 21.7%) exhibited

intermediate and THI (average R²m 13.5%) the lowest performance (Table S8).

Climatic conditions and reproductive failure

Over the two breeding seasons, 72 eggs were laid, 65 hatched and 53 chicks survived

until emancipation, resulting in a 26.4% rate of reproductive failure (19 eggs or chicks that did

not survive). Maximum T_{air} was significantly associated with a higher risk of reproductive failure

(odds-ratio = 1.16; 95% CI = 1.02-1.32; p = 0.025; Table S9A), while maximum SR, Tg and HLI

did not (Table S9B and D). The days where reproductive failure occurred (16 dates where

parents abandoned, or the chick, sometimes several, were found dead) were significantly

hotter than the days where no death was observed, but only regarding maximum T_{air} ($T_{air} = 10.8$

$\pm$ 3.3 °C, n = 170 for days without reproductive failure vs. 12.6 $\pm$ 3.4 °C, n = 16 for days with

reproductive failure, W = 955.5, p = 0.025). Indeed, maximum SR did not differ between days

with or without reproductive failure (SR = 283.8 $\pm$ 95.2 J.cm-² vs. 292.6 $\pm$ 73.6 J.cm-², W =

1582.5, p = 0.54), nor did maximum HLI (70.8 $\pm$ 4.6 vs. 71.8 $\pm$ 4.2, W = 1184, p = 0.20), or

maximum T_g ($T_g = 20.7 \pm 4.3$ °C vs. 21.9 $\pm$ 3.8 °C, W = 1183, p = 0.20).

Discussion

Sub-Antarctic islands are expected to warm by 2.3 to 2.9°C over the next 100 years (Nel *et al.* 2023), and although impactful, this increase in mean temperature still buffers the additional impacts of climate anomalies such as heat waves. Heat waves will likely become more frequent and intense in the coming decades (Gimeno *et al.* 2025; Piatt *et al.* 2024; Stillman 2019), especially during the austral summer, *i.e.* the breeding season of many endemic polar and sub-polar species (Chambers *et al.* 2014). Our study unambiguously shows that king penguins do experience heat stress while breeding on land, at a frequency being far from anecdotal, with >20% of the behavioural scans including signs of heat stress.

While many studies on heat stress focus on air temperature alone, the heat load experienced by wild animals depends markedly on other climatic variables (Mitchell et al. 2024). Here we show that the use of thermoregulatory behaviours in breeding king penguins was primarily influenced by solar radiation, followed by wind speed, air temperature and humidity in decreasing order of importance. Additionally, several interactions between climatic factors were significant in explaining the occurrence of heat stress, even if models without interactions were only slightly less predictive of heat stress occurrence than models including interactions (Table S8). Panting behaviour was the most frequently observed thermoregulatory behaviour, and occurred more often under high solar radiation and low wind speed, as well as at high air temperatures when solar radiation was moderate to high. Wing spreading was the second most frequently observed thermoregulatory behaviour, and it was more likely to occur under high solar radiation, high air temperature and low wind speed. Exposing the brood pouch was the least frequently observed behaviour, which can be explained by a trade-off between heat dissipation and the functions of the pouch: protecting the egg/chick from predators and keeping them warm. Yet, this behaviour was more frequent under high heat load (*i.e.* high solar radiation, high humidity, low wind speed and high air temperature), which supports our hypothesis regarding its potential use as a thermal window. High T_{air} has been reported to trigger heat stress in several seabirds and/or high latitude species (Cook *et al.* 2020; Holt & Boersma 2022; O'Connor *et al.* 2021; Olin *et al.* 2023; Welman & Pichegru 2022). While an increase in T_{air} as a main effect increased the probability to observe wing spreading or brood pouch exposure in breeding king penguins, it had no such effect on panting. Yet, air temperature in interaction with solar radiation explained panting behaviour frequency, to a point that virtually no bird would show panting around 20°C when solar radiation was low, while *ca.* 20% of the studied population was already observed panting below 10°C if sun exposure was maximal (Figure 3A). These results echo recent findings in guillemots, which showed an increased probability of panting with T_{air} in sun-exposed individuals around 15°C, compared with individuals in shaded areas in which panting started around 25°C (Olin *et al.* 2023).

Behavioural thermoregulation may enable animals to maintain normal body
temperature under hot conditions, or only to limit hyperthermia (Talbot *et al.* 2017). Here, we
first show in a sub-sample of 14 king penguins, that although sub-cutaneous temperature is
more variable than core body temperature, they are moderately correlated, and both increase
in response to high temperature and solar radiation. Yet, sub-cutaneous temperature over-
estimates climatic effects on core body temperature, which is not surprising as peripheral
tissues warm up at faster rates than core tissues due heat gain from solar radiation and
increased peripheral perfusion under high heat load to dissipate body heat through convection.
While acknowledging these limits, subcutaneous temperature may still serve as a useful proxy
of core body temperature when sampling a large number of individuals is required. Here, we
showed that subcutaneous temperatures rose under increasing heat load (high solar radiation,
air temperature and humidity, as well as low wind speed). Additionally, panting birds had a
higher T_{sc} than non-panting ones, and T_{sc} increased with air temperature only in panting king
penguins. This result may illustrate classic physiological thermoregulation: the panting birds
that would be the most subjected to heat stress, underwent peripheral vasodilatation in order
to evacuate body heat through convection, hence increasing T_{sc}. The particularly thick
feathered insulation coat of king penguins and limited thermal windows may reduce their
capacity to lose heat through their body surface (Dawson *et al.* 1999; McCafferty *et al.* 2013).
Additionally, breakpoint analysis allowed the detection of several threshold above which T_{sc}
would increase even further depending on T_{air}, SR and HLI, which could illustrate a point at
which core body temperatures increase. Although globe temperature has been suggested as
an integrative metric of the environmental heat load (Mitchell et al., 2024), our analysis did not
reveal a similar significant threshold for T_g. Yet, to ensure the robustness of our conclusions,
we would ideally need to measure core body temperature on a larger number of individuals to
properly assess the penguins' ability to maintain (or not) their core body temperature under
high heat load. Our results contrast with studies in dovekies (*Alle alle*) suggesting that the core
body temperature of this Arctic species is not challenged while attending the breeding colony
on land (Grunst *et al.* 2023; maximum T_{air} = 13.5°C over 11 days) or in a captive set up to 25°C
(*i.e.* substantially above the natural maximal temperature; Beaman *et al.* 2024). Yet, solar
radiation was not investigated in the aforementioned studies, and empirical evidence, including
in our king penguins, suggests that solar radiation is a key factor conditioning the total heat
load faced by wild animals (Mitchell et al. 2024).

Taken together, the 4 climatic variables (T_{air}, SR, WS and RH) we used predicted heat
stress occurrence more accurately than T_{air} alone, or the four classical thermal indexes we
used (T_g, THI, HLI, WBGT). These indexes have been established for humans and farm
animals (Gaughan 2002; Ioannou *et al.* 2023), which likely explain why they seem less suited

for wild seabirds that present morphological and physiological adaptations to the cold, and
breed in unsheltered conditions. Developing heat indexes specifically for wild endotherms
would enable to better predict the response of natural populations to heat waves. Considering
the tremendous impact of solar radiation and wind speed on animal heat load, modelling future
changes in climatic parameters beyond the sole air temperature seems of critical importance
to predict the coping capacity of wild species in the context of climate change. Yet, for many
species, general measures from meteorological stations may only reflect poorly the micro-
climatic conditions experienced by animals, and thus on-site measurement of micro-climate is
needed (Mitchell *et al.* 2024).

Beyond climatic parameters, biological factors were also important in predicting the
risks of king penguins to suffer from heat stress. First, sex did seem to play a role (albeit small)
in explaining variation in heat stress occurrence, since males spread their wings more often
and had a tendency to pant more than females. This may, at least partly, be linked to the higher
size of beak and flipper relative to body mass found in females (Kriesell *et al.* 2018), which
likely provides higher heat dissipation capacity through those thermal windows (Lewden *et al.*
2020b). Second, brooding parents generally suffered more from hot conditions than incubating
ones, as they spread their wings and exposed their brood pouch more often, as well as
exhibited higher subcutaneous temperature than incubating individuals. Yet, brooding adults
were not panting more than incubating ones. Considering that panting is extremely expensive
in terms of water (Adélie penguins can lose water up to 0.3% of their body mass per hour while
panting; Chappell & Souza 1988), our results may suggest that brooding individuals could limit
the use of panting to avoid severe dehydration (Smit *et al.* 2016), especially because breeding
king penguins do not appear to drink while fasting on land (Cherel *et al.* 1988). Brooding adults
are also known to exhibit higher aggression rates towards conspecifics than incubating ones
(Côté 2000), which could result in elevated body temperatures due to both physical activity
and stress response (Knoch *et al.* 2022). Finally, king penguins seemed especially sensitive to
heat stress when freshly returned from the sea to incubate or brood on land. Indeed, they
showed a higher probability to pant and a higher subcutaneous temperature during their first
few days on land. This may be explained by the gradual transition to the hypometabolic state
associated with fasting (2-5 days are needed to reach a stable hypometabolic state; Cherel *et*
al. 1988), but also to their thermal acclimation to on-land conditions. The mechanisms involved
in such a potential transient thermal acclimation on land remain to be investigated.

The impact of heat stress on breeding king penguins in our study was sublethal,
contrary to what has been reported in Magellanic (44°C; Holt & Boersma 2022) and king
penguins (two days at 24 and 30°C; Arriagada 2023) breeding in Argentina. Yet, sublethal
effects may have drastic consequences on population dynamics (Conradie *et al.* 2019),

especially if reproduction is affected (Morosinotto *et al.* 2025). Reproductive failure may be
related or accentuated by hot environmental conditions since parents may abandon
reproduction because of heat stress, or be more susceptible to egg/chick predation when
undergoing heat stress. Our results show that climatic conditions were hotter during days
where reproductive failure occurred, which suggests that heat stress may be an important
constraint on reproductive success for king penguins, as this has been demonstrated
previously in common Guillemot (Olin *et al.* 2023). Better understanding the sublethal
consequences of heat stress in king penguins is thus needed to predict more accurately the
changes in population dynamics expected due to the complex environmental challenges
brought by climate change.

Ethical note

Acknowledgments

We are grateful to the French Polar Institute (IPEV) and the Terres Australes et
Antarctiques Françaises for providing financial and logistical support for this study through the
polar project #119 (ECONERGY). We are grateful to Mathilde Lejeune, Thomas Faulman, Nina
Cossin-Sevrin, Robin Dardel, Laetitia Vuibert and the entire Alfred Faure field station for their
help in the field. This study is part of the long-term Studies in Ecology and Evolution (SEE-Life)
program of the CNRS. We wish to thank the Zone Atelier Antarctique et Terres Australes (ZATA)
from the CNRS for financial support, and the members of the Alfred Faure field station for their
help and support in the field. AL was supported by ISblue project, Interdisciplinary graduate
school for the blue planet (ANR-17-EURE-0015) and co-funded by a grant from the French
government under the program "Investissements d'Avenir" embedded in France 2030. AS was
supported by a 'Turku Collegium for Science and Medicine' Fellowship, a Marie Sklodowska-
Curie Postdoctoral Fellowships (#894963) and AN was funded by an IdEx grant from the
Université de Strasbourg (*HotPenguin*) granted to AS. CB, MM, MH and EM were funded by
the French Polar Institute.

References

Ancel, A., Beaulieu, M. & Gilbert, C. (2013). The different breeding strategies of penguins: A review. *Comptes Rendus Biologies*, 336, 1–12.

Andreasson, F., Rostedt, E. & Nord, A. (2023). Measuring body temperature in birds – the effects of sensor type and placement on estimated temperature and metabolic rate. *Journal of Experimental Biology*, 226, jeb246321.

Arriagada, M. (2023). King penguin mortality related to heat wave events in 2019 and 2020 at Bahía Inútil, Tierra del Fuego. In: *11th International Penguin Congress*.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using **lme4**. *J. Stat. Soft.*, 67.

Beaman, J.E., White, C.R., Clairbaux, M., Perret, S., Fort, J. & Grémillet, D. (2024). Cold adaptation does not handicap warm tolerance in the most abundant Arctic seabird. *Proc. R. Soc. B.*, 291, 20231887.

Bost, C.A., Cotté, C., Terray, P., Barbraud, C., Bon, C., Delord, K., *et al.* (2015). Large-scale climatic anomalies affect marine predator foraging behaviour and demography. *Nat Commun*, 6, 8220.

Boyles, J.G., Seebacher, F., Smit, B. & McKechnie, A.E. (2011). Adaptive Thermoregulation in Endotherms May Alter Responses to Climate Change. *Integrative and Comparative Biology*, 51, 676–690.

Bryant, D.M. (2008). Heat stress in tropical birds: behavioural thermoregulation during flight. *Ibis*, 125, 313–323.

Budd, G.M. (2008). Wet-bulb globe temperature (WBGT)—its history and its limitations. *Journal of Science and Medicine in Sport*, 11, 20–32.

Carter, A.W., Zaitchik, B.F., Gohlke, J.M., Wang, S. & Richardson, M.B. (2020). Methods for Estimating Wet Bulb Globe Temperature From Remote and Low-Cost Data: A Comparative Study in Central Alabama. *GeoHealth*, 4, e2019GH000231.

Chambers, L.E., Dann, P., Cannell, B. & Woehler, E.J. (2014). Climate as a driver of phenological change in southern seabirds. *Int J Biometeorol*, 58, 603–612.

Chappell, M.A. & Souza, S.L. (1988). Thermoregulation, gas exchange, and ventilation in Adelie penguins (Pygoscelis adeliae). *J Comp Physiol B*, 157, 783–790.

Cherel, Y., Robin, J.P., Walch, O., Karmann, H., Netchitailo, P. & Le Maho, Y. (1988). Fasting in king penguin. I. Hormonal and metabolic changes during breeding. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 254, R170–R177.

Choy, E.S., O'Connor, R.S., Gilchrist, H.G., Hargreaves, A.L., Love, O.P., Vézina, F., *et al.* (2021). Limited heat tolerance in a cold-adapted seabird: implications of a warming Arctic. *Journal of Experimental Biology*, 224, jeb242168.

Clem, K.R., Fogt, R.L., Turner, J., Lintner, B.R., Marshall, G.J., Miller, J.R., *et al.* (2020). Record warming at the South Pole during the past three decades. *Nat. Clim. Chang.*, 10, 762–770.

Conradie, S.R., Woodborne, S.M., Cunningham, S.J. & McKechnie, A.E. (2019). Chronic, sublethal effects of high temperatures will cause severe declines in southern African arid-zone birds during the 21st century. *Proc. Natl. Acad. Sci. U.S.A.*, 116, 14065–14070.

Conradie, S.R., Woodborne, S.M., Wolf, B.O., Pessato, A., Mariette, M.M. & McKechnie, A.E. (2020). Avian mortality risk during heat waves will increase greatly in arid Australia during the 21st century. *Conservation Physiology*, 8, coaa048.

Cook, T.R., Martin, R., Roberts, J., Häkkinen, H., Botha, P., Meyer, C., *et al.* (2020). Parenting in a warming world: thermoregulatory responses to heat stress in an endangered seabird. *Conservation Physiology*, 8, coz109.

Côté, S.D. (2000). Aggressiveness in king penguins in relation to reproductive status and territory location. *Animal Behaviour*, 59, 813–821.

Danner, R.M., Coomes, C.M. & Derryberry, E.P. (2021). Simulated heat waves reduce cognitive and motor performance of an endotherm. *Ecol. Evol.*, 11, 2261–2272.

Davis, L.S. & Darby, J.T. (2012). *Penguin Biology*. Elsevier.

Dawson, C., Vincent, J.F.V., Jeronimidis, G., Rice, G. & Forshaw, P. (1999). Heat Transfer through Penguin Feathers. *Journal of Theoretical Biology*, 199, 291–295.

Descamps, S., Gauthier-Clerc, M., Le Bohec, C., Gendner, J.-P. & Le Maho, Y. (2005). Impact of predation on king penguin Aptenodytes patagonicus in Crozet Archipelago. *Polar Biol*, 28, 303–310.

Eichhorn, G., Groscolas, R., Le Glaunec, G., Parisel, C., Arnold, L., Medina, P., *et al.* (2011). Heterothermy in growing king penguins. *Nat Commun*, 2, 435.

Gaughan, J. (2002). *Validation of the Heat Load Index for use in the feedlot industry* (No. 9781741918069).

Gauthier-Clerc, M., Le Maho, Y., Gendner, J.-P., Durant, J. & Handrich, Y. (2001). State-dependent decisions in long-term fasting king penguins, Aptenodytes patagonicus, during courtship and incubation. *Animal Behaviour*, 62, 661–669.

Gimeno, M., Ramírez, F., Coll, M., Chiaradia, A. & Artana, C. (2025). Cumulative Extreme Events Threaten Penguin Habitats Across the Southern Hemisphere. *Global Change Biology*, 31, e70562.

Grémillet, D., Meslin, L. & Lescroël, A. (2012). Heat dissipation limit theory and the evolution of avian functional traits in a warming world. *Funct Ecol*, 26, 1001–1006.

Grunst, M.L., Grunst, A.S., Grémillet, D., Kato, A., Gentès, S. & Fort, J. (2023). Keystone seabird may face thermoregulatory challenges in a warming Arctic. *Sci Rep*, 13, 16733.

Hajizadeh, R., Farhang Dehghan, S., Golbabaei, F., Jafari, S.M. & Karajizadeh, M. (2017). Offering a model for estimating black globe temperature according to meteorological measurements. *Meteorological Applications*, 24, 303–307.

Handrich, Y., Bevan, R.M., Charrassin, J.-B., Butler, P.J., Ptz, K., Woakes, A.J., *et al.* (1997). Hypothermia in foraging king penguins. *Nature*, 388, 64–67.

Hansen, P.J. (2009). Effects of heat stress on mammalian reproduction. *Phil. Trans. R. Soc. B*, 364, 3341–3350.

Hartig, F. (2016). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models.

Holt, K.A. & Boersma, P.D. (2022). Unprecedented heat mortality of Magellanic Penguins. *Ornithological Applications*, 124, duab052.

Ioannou, L.G., Ciuha, U., Fisher, J.T., Tsoutsoubi, L., Tobita, K., Bonell, A., *et al.* (2023). Novel Technological Advances to Protect People Who Exercise or Work in Thermally Stressful Conditions: A Transition to More Personalized Guidelines. *Applied Sciences*, 13, 8561.

Ioannou, L.G., Mantzios, K., Tsoutsoubi, L., Notley, S.R., Dinas, P.C., Brearley, M., *et al.* (2022). Indicators to assess physiological heat strain – Part 1: Systematic review. *Temperature*, 9, 227–262.

Knoch, S., Whiteside, M.A., Madden, J.R., Rose, P.E. & Fawcett, T.W. (2022). Hot-headed peckers: thermographic changes during aggression among juvenile pheasants (*Phasianus colchicus*). *Phil. Trans. R. Soc. B*, 377, 20200442.

Kriesell, H.J., Aubin, T., Planas-Bielsa, V., Benoiste, M., Bonadonna, F., Gachot-Neveu, H., *et al.* (2018). Sex identification in King Penguins *Aptenodytes patagonicus* through morphological and acoustic cues. *Ibis*, 160, 755–768.

Lemonnier, C., Schull, Q., Stier, A., Boonstra, R., Delehanty, B., Lefol, E., *et al.* (2024). Social, not genetic, programming of development and stress physiology of a colonial seabird. *Proceedings of the Royal Society B: Biological Sciences*, 291, 20240853.

Lewden, A., Bonnet, B. & Nord, A. (2020a). The metabolic cost of subcutaneous and abdominal rewarming in king penguins after long-term immersion in cold water. *Journal of Thermal Biology*, 91, 102638.

Lewden, A., Enstipp, M.R., Picard, B., Van Walsum, T. & Handrich, Y. (2017). High peripheral temperatures in king penguins while resting at sea: thermoregulation versus fat deposition. *Journal of Experimental Biology*, jeb.158980.

Lewden, A., Nord, A., Bonnet, B., Chauvet, F., Ancel, A. & McCafferty, D.J. (2020b). Body surface rewarming in fully and partially hypothermic king penguins. *J Comp Physiol B*, 190, 597–609.

Liljegren, J.C., Carhart, R.A., Lawday, P., Tschopp, S. & Sharp, R. (2008). Modeling the Wet
Bulb Globe Temperature Using Standard Meteorological Measurements. *Journal of*
Occupational and Environmental Hygiene, 5, 645–655.
McCafferty, D.J., Gallon, S. & Nord, A. (2015). Challenges of measuring body temperatures of
free-ranging birds and mammals. *Anim Biotelemetry*, 3, 33.
McCafferty, D.J., Gilbert, C., Thierry, A.-M., Currie, J., Le Maho, Y. & Ancel, A. (2013). Emperor
penguin body surfaces cool below air temperature. *Biol. Lett.*, 9, 20121192.
Mitchell, D., Maloney, S.K., Snelling, E.P., Carvalho Fonsêca, V.D.F. & Fuller, A. (2024).
Measurement of microclimates in a warming world: problems and solutions. *Journal of*
Experimental Biology, 227, jeb246481.
Morosinotto, C., Gasparini, C., Pilastro, A., Rosenthal, G.G. & Breedveld, M.C. (2025). Cryptic
Reproductive Costs of Heatwaves for Animal Populations. *Global Change Biology*, 31,
e70558.
Muggeo, V. (2008). segmented: an R Package to Fit Regression Models with Broken-Line
Relationships.
Nel, W., Hedding, D.W. & Rudolph, E.M. (2023). The sub-Antarctic islands are increasingly
warming in the 21st century. *Antarctic Science*, 35, 124–126.
O'Connor, R.S., Le Pogam, A., Young, K.G., Robitaille, F., Choy, E.S., Love, O.P., *et al.* (2021).
Limited heat tolerance in an Arctic passerine: Thermoregulatory implications for cold-
specialized birds in a rapidly warming world. *Ecol. Evol.*, 11, 1609–1619.
Olin, A., Dück, L., Berglund, P., Karlsson, E., Bohm, M., Olsson, O., *et al.* (2023). Breeding
failures and reduced nest attendance in response to heat stress in a high-latitude
seabird. *Mar. Ecol. Prog. Ser.*, HEAT.
Oswald, K.N., Evlambiou, A.A., Ribeiro, Â.M. & Smit, B. (2018). Tag location and risk
assessment for passive integrated transponder-tagging passerines. *Ibis*, 160, 453–
457.
Oswald, S.A., Bearhop, S., Furness, R.W., Huntley, B. & Hamer, K.C. (2008). Heat stress in a
high-latitude seabird: effects of temperature and food supply on bathing and nest
attendance of great skuas *Catharacta skua*. *Journal of Avian Biology*, 39, 163–169.
Piatt, J., Arimitsu, M., Thompson, S., Suryan, R., Wilson, R., Elliott, K., *et al.* (2024).
Mechanisms by which marine heatwaves affect seabirds. *Mar. Ecol. Prog. Ser.*, 737,
1–8.
Pipoly, I., Bókony, V., Seress, G., Szabó, K. & Liker, A. (2013). Effects of Extreme Weather on
Reproductive Success in a Temperate-Breeding Songbird. *PLoS ONE*, 8, e80033.
Robinson, P.J. (2001). On the Definition of a Heat Wave. *J. Appl. Meteor.*, 40, 762–775.
Romanovsky, A.A. (2007). Thermoregulation: some concepts have changed. Functional
architecture of the thermoregulatory system. *American Journal of Physiology-*
Regulatory, Integrative and Comparative Physiology, 292, R37–R46.
Romanovsky, A.A. (2018). The thermoregulation system and how it works. In: *Handbook of*
Clinical Neurology. Elsevier, pp. 3–43.
Silva, R.G. da, Morais, D.A.E.F. & Guilhermino, M.M. (2007). Evaluation of thermal stress
indexes for dairy cows in tropical regions. *R. Bras. Zootec.*, 36, 1192–1198.
Smit, B., Zietsman, G., Martin, R.O., Cunningham, S.J., McKechnie, A.E. & Hockey, P.A.R.
(2016). Behavioural responses to heat in desert birds: implications for predicting
vulnerability to climate warming. *Clim Chang Responses*, 3, 9.
Stammerjohn, S.E. & Scambos, T.A. (2020). Warming reaches the South Pole. *Nat. Clim.*
Chang., 10, 710–711.
Stillman, J.H. (2019). Heat Waves, the New Normal: Summertime Temperature Extremes Will
Impact Animals, Ecosystems, and Human Communities. *Physiology*, 34, 86–100.
Stonehouse, B. (1960). Breeding behaviour and development. In: *The King Penguin*
(Aptenodites patagonica) of South Georgia.
Talbot, W.A., McWhorter, T.J., Gerson, A.R., McKechnie, A.E. & Wolf, B.O. (2017). Avian
thermoregulation in the heat: evaporative cooling capacity of arid-zone caprimulgiforms
from two continents. *Journal of Experimental Biology*, jeb.161653.

Tan, C.L. & Knight, Z.A. (2018). Regulation of Body Temperature by the Nervous System.
Neuron, 98, 31–48.
Thom, E.C. (1959). The Discomfort Index. *Weatherwise*, 12, 57–61.
Ton, R., Stier, A., Cooper, C.E. & Griffith, S.C. (2021). Effects of Heat Waves During Post-natal
Development on Mitochondrial and Whole Body Physiology: An Experimental Study in
Zebra Finches. *Front. Physiol.*, 12, 661670.
Viblanc, V.A., Valette, V., Kauffmann, M., Malosse, N. & Groscolas, R. (2012). Coping with
social stress: heart rate responses to agonistic interactions in king penguins.
Behavioral Ecology, 23, 1178–1185.
Weimerskirch, H., Stahl, J.C. & Jouventin, P. (1992). The breeding biology and population
dynamics of King Penguins *Aptenodytes patagonica* on the Crozet Islands. *Ibis*, 134,
107–117.
Welman, S. & Pichegru, L. (2022). Nest microclimate and heat stress in African Penguins
Spheniscus demersus breeding on Bird Island, South Africa. *Bird Conservation*
International, 1–9.
You, Q., Cai, Z., Pepin, N., Chen, D., Ahrens, B., Jiang, Z., *et al.* (2021). Warming amplification
over the Arctic Pole and Third Pole: Trends, mechanisms and consequences. *Earth-*
Science Reviews, 217, 103625.
Zuluaga, J.D., Pretti, E., Leynaert, A., Marçon, E., Stier, A. & Lewden, A. (2025). Warming up
to a new coat: molting king penguins exhibit hyperthermia and increased peripheral
heat loss. *American Journal of Physiology-Regulatory, Integrative and Comparative*
Physiology, 329, R742–R752.
Zuur, A.F., Ieno, E.N. & Elphick, C.S. (2010). A protocol for data exploration to avoid common
statistical problems: *Data exploration. Methods in Ecology and Evolution*, 1, 3–14.
714

715 **Electronic supplementary material**

717 **ESM S1**

719 **Figure. S1 Climatic conditions (daily maximum air temperature (in red) and maximum solar**
720 **radiation (in orange) in 2022 in Crozet Archipelago along with a description of the breeding cycle of king**
721 **penguins.**

ESM S2: Predicting black globe temperature (T_g) based on meteorological parameters

Considering that T_g was not directly measured at our study site during the timeframe of this study, we relied on meteorological parameters to predict T_g, based on actual T_g measurements conducted during the 2023-24 field season with a Kestrel 5400 Heat Stress Tracker. We evaluated how measured T_g was best predicted by our four meteorological parameters (T_{air}, SR, WS and RH) by selecting the best linear model using the R function dredge (Package MuMIn, version 1.48.11), which excluded RH from the model (adjusted R^2 of the selected model = 0.76).

A

```
                Global model call: lm(formula = Tg ~ Tair + RU + WS + SR, data = data,
na.action = na.fail)
---
Model selection table
    (Intrc)      WS      SR   Tair        RH df     logLik   AICc  delta weight
8     2.887 -0.03470 0.03561 0.8265          5  -901.641 1813.4   0.00  0.411
7     2.598          0.03514 0.8211          4  -902.936 1814.0   0.54  0.314
16    2.483 -0.03583 0.03593 0.8204  0.005240 6  -901.551 1815.3   1.88  0.160
15    2.382          0.03530 0.8178  0.002734 5  -902.911 1816.0   2.54  0.115
11    1.031          0.04426        0.101100 4 -1053.295 2114.7 301.26  0.000
12    1.077 -0.01686 0.04457        0.102400 5 -1053.151 2116.5 303.02  0.000
14   12.280  0.07715         1.0350 -0.123600 5 -1062.429 2135.0 321.58  0.000
13   12.900                  1.0500 -0.123000 4 -1065.450 2139.0 325.57  0.000
3    10.120          0.03882          3 -1071.381 2148.8 335.39  0.000
4    10.000  0.01200 0.03865          4 -1071.313 2150.7 337.29  0.000
6     2.711  0.07382         0.9092          4 -1089.818 2187.7 374.30  0.000
5     3.348                  0.9237          3 -1092.241 2190.5 377.11  0.000
10   13.560  0.13930                -0.034460 4 -1188.072 2384.2 570.81  0.000
2    10.590  0.13590                  3 -1189.417 2384.9 571.46  0.000
9    14.730                        -0.031150 3 -1193.421 2392.9 579.47  0.000
1    12.010                          2 -1194.494 2393.0 579.59  0.000
Models ranked by AICc(x)
```

B

Predictors	Estimates	CI	p
	Tg		
(Intercept)	12.01	11.80 – 12.23	**<0.001**
T_{air}	2.52	2.29 – 2.76	**<0.001**
WS	-0.19	-0.41 – 0.04	0.101
SR	2.82	2.57 – 3.07	**<0.001**
RH	0.06	-0.20 – 0.31	0.673
Observations	408		
R2 / R2 adjusted	0.762 / 0.760		

C

Predicted LM vs. Actual Values

2.88 + 0.83 x Tair - 0.035 x WS + 0.036 x SR
adjusted R^2 = 0.76

Actual Values / Predicted Values

Figure S2. Prediction of the globe temperature (T_g) measured inside the penguin's colony depending on meteorological parameters from the nearby Meteo France station (air temperature, solar radiation, wind speed and relative humidity), using linear modelization. (A) dredge selection of lm(Tg~Tair+SR+WS+RH) depending on the lowest corrected Aikake's Coefficient; **(B)** relative weight of the different climatic variables on globe temperature after scaling (z-transformation) and **(C)** Predicted vs. Actual values of T_g.

Figure S3. **Predicted probability of (A) Panting depending on air temperature and relative humidity (B) panting depending on solar radiation and relative humidity (C) panting depending on wind speed and relative humidity (D) wing spreading depending on solar radiation and relative humidity and (E) wing spreading depending on wind speed and relative humidity;** probabilities were calculated from generalized linear mixed models; N = 143, n = 1123 relative humidity (%) is represented as second quartile (67), median (73) and 3rd quartile (84) registered during behavioural scans.

758 **ESM S4**

761 **Figure S4. Evolution of (A) panting probability and (B) subcutaneous temperature along the duration of the**
762 **shift on land.** Values are predicted from linear mixed models presented in Table 1. N = 143, n = 1123 for panting;
763 N = 140, n = 856 for T_{sc}.

765
766 **Figure S5.** Relationships between **sub-cutaneous temperature (T_sc, in °C) and core body temperature (T_b, in**
767 **°C). (A) Correlation (*r* = 0.62, p < 0.001), (B) Bland-Altman plot** representing T_b and T_{sc} difference ($Ts_c - T_b$) of
768 the same individuals against their mean, **(C-F) relationships between Tb (red), Tsc (blue) and their difference**
769 **(T_b – T_sc: black dotted line)** depending on climatic parameters (the significance of each relationship is marked with
770 an asterisk): air temperature (°C), solar radiation (J.cm-1), wind speed (m.s-1) and relative humidity (%).

ESM S6

A) Tair

Predictors	Tsc Est.	Tsc CI	Tsc p	Tb Est.	Tb CI	Tb p	Diff Est.	Diff CI	Diff p
(Intercept)	36.71	35.88 – 37.55	**<0.001**	37.59	37.23 – 37.96	**<0.001**	0.91	0.22 – 1.60	**0.012**
Tair	0.09	0.02 – 0.16	**0.010**	0.03	0.01 – 0.06	**0.014**	-0.06	-0.11 – -0.00	**0.041**

Random Effects

	Tsc	Tb	Diff
σ^2	0.77	0.12	0.50
$\tau_{00\ ID}$	0.12	0.08	0.11
ICC	0.14	0.40	0.18
N $_{ID}$	14	14	14
Observations	37	37	37
Marginal R^2 / Cond. R^2	0.161 / 0.275	0.114 / 0.469	0.100 / 0.263

B) SR

Predictors	Tsc Est.	Tsc CI	Tsc p	Tb Est.	Tb CI	Tb p	Diff Est.	Diff CI	Diff p
(Intercept)	35.67	34.19 – 37.15	**<0.001**	37.35	36.69 – 38.01	**<0.001**	1.79	0.60 – 2.99	**0.004**
SR	0.01	0.00 – 0.01	**0.006**	0.00	0.00 – 0.00	**0.047**	-0.00	-0.01 – -0.00	**0.011**

Random Effects

	Tsc	Tb	Diff
σ^2	0.69	0.13	0.42
$\tau_{00\ ID}$	0.25	0.07	0.23
ICC	0.27	0.36	0.35
N $_{ID}$	14	14	14
Observations	37	37	37
Marginal R^2 / Cond. R^2	0.178 / 0.400	0.090 / 0.416	0.147 / 0.443

C) WS

Predictors	Tsc Est.	Tsc CI	Tsc p	Tb Est.	Tb CI	Tb p	Diff Est.	Diff CI	Diff p
(Intercept)	37.44	36.59 – 38.29	**<0.001**	37.95	37.56 – 38.35	**<0.001**	0.54	-0.14 – 1.22	0.117
WS	0.04	-0.06 – 0.14	0.425	0.00	-0.04 – 0.05	0.826	-0.04	-0.12 – 0.04	0.336

Random Effects

	Tsc	Tb	Diff
σ^2	1.02	0.15	0.64
$\tau_{00\ ID}$	0.00	0.07	0.01
ICC	14	0.34	0.01
N $_{ID}$	37	14	14
Observations	0.018 / NA	37	37
Marginal R^2 / Cond. R^2	1.02	0.001 / 0.336	0.026 / 0.035

D) RH

Predictors	Tsc Est.	Tsc CI	Tsc p	Tb Est.	Tb CI	Tb p	Diff Est.	Diff CI	Diff p
(Intercept)	38.64	36.30 – 40.98	**<0.001**	38.53	37.57 – 39.50	**<0.001**	-0.27	-2.13 – 1.59	0.769
RH	-0.01	-0.05 – 0.02	0.438	-0.01	-0.02 – 0.01	0.250	0.01	-0.02 – 0.03	0.568

Random Effects

	Tsc	Tb	Diff
σ^2	0.98	0.14	0.59
$\tau_{00\ ID}$	0.05	0.08	0.07
ICC	0.05	0.35	0.11
N $_{ID}$	14	14	14
Observations	37	37	37
Marginal R^2 / Cond. R^2	0.017 / 0.066	0.030 / 0.373	0.009 / 0.120

Table S6. Summaries of the linear mixed models predicting sub-cutaneous temperature (T_{sc}), core body temperature (T_b) and their difference (Diff), depending on (A) ambient temperatures (T_{air}), (B) solar radiation (SR), (C) wind speed (WS) and (D) relative humidity (RH).

Figure S7. Breakpoint analyses between king penguins' sub-cutaneous temperatures (T$_{sc}$) and (A) air temperature; (B) solar radiation; (C) Globe temperature and (D) heat load index (HLI) computed from air temperature, solar radiation, wind speed and relative humidity. Segmented models are shown in red, and estimated breakpoints with confidence intervals are shown in blue. *N = 140, n = 856.*

ESM S8

Table S8. Model comparison for the prediction of panting, wing spreading, brood pouch exposure and Tsc based on Akaike's information criterion (AICc), marginal R² (R²m) and conditional R² (R²c). The best predictive model with interactions ('Inter.') between all 4 climatic parameters (Tair, SR, WS and RH), is compared to a model without interactions ('Total'), a model with only Tair or SR and 4 models including Heat indexes: globe

	Panting			Wing spreading			Brooding pouch exposure			Tsc		
	AICc	R²m	R²c	AICc	R²m	R²c	AICc	R²m	R²c	AICc	R²m	R²c
Tair	1067.6	0.23	0.30	659.82	0.14	0.29	517.72	0.15	0.55	2514.05	0.071	0.10
SR	928.96	0.49	0.52	601.93	0.33	0.45	462.88	0.42	0.69	2343.39	0.24	0.28
Total	740.06	0.67	0.71	554.86	0.44	0.59	401.65	0.50	0.80	2277.08	0.30	0.35
Inter.	695.47	0.68	0.73	545.37	0.57	0.69	382.09	0.48	0.84	2181.49	0.37	0.43
Tg		0.48	0.55		0.30	0.48		0.31	0.68		0.22	0.26
THI		0.21	0.28		0.13	0.27		0.14	0.58		0.062	0.092
WBGT		0.33	0.41		0.20	0.37		0.21	0.61		0.13	0.16
HLI		0.47	0.54		0.30	0.47		0.32	0.70		0.19	0.23

ESM S9

Table S9. Summaries of the generalized linear models (binomial) predicting offspring death depending on: **(A)** maximum ambient temperatures (Tmax), **(B)** maximum solar radiation (SRmax), (**C**) maximum Globe temperature (Tgmax) and **(D)** maximum Heat Load Index (HLImax).

Death prop.	(A) Tmax			(B) SRmax			(C) Tgmax			(D) HLImax		
Predictors	Odds Ratios	CI	p	Odds Ratios	CI	p	Odds Ratios	CI	p	Odds Ratios	CI	p
(Intercept)	0.00025	0.00 – 0.002	**<0.001**	0.00025	0.00- 0.003	**<0.001**	0.00025	0.00- 0.003	**<0.001**	0.00	0.00 – 0.02	**0.002**
Climatic predictor	1.16	1.02 – 1.32	**0.025**	1.00	1.00 – 1.01	0.58	1.09	0.97 – 1.21	0.14	1.07	0.97 – 1.17	0.20