1
14 **Functional separation of long-range inputs by intrinsic dynamics of dorsal raphe 5-HT**
15 **neurons**
16
17
18

19 Michael Lynn[1], Richard Naud[1], Leonard Maler[1] & Jean-Claude Béïque[1]
20
21
22
23 1: Department of Cellular & Molecular Medicine, University of Ottawa

Abstract:

The serotonergic dorsal raphe nucleus (DRN) receives diverse long-range synaptic inputs, yet the processing rules governing their contribution to DRN output spiking patterns are largely unknown. Here, we use electrophysiological, optogenetic and computational approaches to compare electrophysiological features of two major excitatory inputs onto DRN 5-HT neurons - the lateral habenula (LHb) and medial prefrontal cortex (mPFC). Dual-color opsin strategies revealed that a population of 5-HT neurons receive inputs from both mPFC and LHb. Subthreshold excitatory postsynaptic potentials triggered by both inputs were largely indistinguishable, yet suprathreshold spiking behavior exhibited input-specific latencies and dispersion statistics. A support vector machine classifier demonstrated that input identity can be accurately decoded from spike timing, but not subthreshold events, of under ten 5-HT neurons. Upon examining the intrinsic cellular mechanisms in 5-HT neurons that couple EPSPs to spiking dynamics, we uncovered two likely candidate mechanisms: a low-threshold calcium conductance that selectively boosts slow excitatory inputs, and a subthreshold, voltage-dependent membrane noise that generates variation of spike latency and jitter. Stochastic simulations suggest that these two intrinsic properties of 5-HT neurons are sufficient to transform LHb and mPFC inputs into distinct output spiking patterns. These results reveal that hub-like networks like the DRN can segregate distinct informational streams by a cell-intrinsic mechanism. The resulting emergent population spike synchrony code provides a means for the DRN to widely broadcast these streams as a multiplexed signals.

Introduction

The phylogenetically old monoaminergic nuclei share common core organizational features. They are anatomically compact, give rise to widespread and distributed axonal projections, and are controlled by expansive and convergent sets of synaptic inputs carrying distinct informational content. A corollary computational question is how these multiple, temporally overlapping, competing inputs are differentially processed by these hub networks. In principle, populations of neurons in these networks could either collapse independent streams of information into a low-dimensional rate code, or alternatively could independently encode features of the inputs in a way that can be decoded by downstream regions, a neural strategy termed 'multiplexing' which includes a diversity of rate or temporal code implementations (Naud & Sprekeler, 2018; Payeur et al., 2021; Oswald et al., 2004; Clarke & Maler, 2017; Lankarany et al, 2019; Ratte et al., 2013; Friedrich et al, 2004; Stringer et al., 2019; Nogueira et al., 2023). Physiological mechanisms for multiplexing can involve cell-intrinsic properties of individual neurons (Doiron et al., 2002; Laing et al., 2003), or alternatively can depend on large-scale network interactions, generating multiplexed neural codes at the population level (e,g, Stringer et al., 2019). It is currently unknown whether monoaminergic nuclei functionally discriminate competing excitatory inputs, an obligatory prerequisite of multiplexing, as well as what potential neural coding strategies could implement such a discrimination.

The midbrain dorsal raphe nucleus (DRN) is a monoaminergic nucleus that has received sustained attention notably because of its role in mood regulation and decision-making processes (Warden et al., 2012; Ren et al,. 2018; Li et al, 2016; Paquelet et al., 2022; Cohen et al., 2015). The anatomical and functional organization of DRN is complex, with DRN 5-HT neurons innervating a vast array of cortical and subcortical structures and receiving broad inputs, including from structures encoding cognitive and valence information such as the medial prefrontal cortex (mPFC) and lateral habenula (LHb) (Dorocic et al., 2014; Weissbourd et al., 2014; Geddes et al., 2016; Zhou et al., 2017). These two inputs provide excitatory synaptic inputs to partially overlapping anatomical divisions (Zhou et al., 2017) including the ventral section of DRN, yet are functionally distinct, with mPFC representing persistent cognitive states (Miller & Cohen, 2001; Bari et al., 2019) and LHb cells responding to transient negative reward prediction errors with high temporal precision spiking responses (Matsumoto & Hikosaka, 2007; Bromberg-Martin & Hikosaka, 2011).

Because of their fundamentally distinct informational content, we reasoned that the inputs from the PFC and the LHb present an ideal testbed to explore discriminative neural strategies in the DRN. We thus compared the LHb and PFC inputs to DRN 5-HT neurons using cellular electrophysiology and optogenetic approaches. While individual 5-HT neurons received broadly similar excitatory synaptic inputs from both sources, those from the PFC evoked spikes in 5-HT neurons with distinct statistics (*i.e.,* latency and jitter) than those triggered by LHb input activation. Machine learning approaches demonstrated that these timing differences were theoretically sufficient for downstream circuits receiving projections from DRN populations to functionally discriminate these input streams, suggesting a multiplexed population code for LHb and PFC input information. Physiological investigations and stochastic simulations defined a cell-intrinsic mechanism involving a low-threshold calcium channel-mediated conductance in 5-HT neurons that is responsible for generating these spike timing differences. Taken together, these data suggest that the phylogenetically old, expansively projecting DRN 5-HT system can broadcast multiple streams of information that are reflecting the contribution of distinct source inputs.

148
Results
Single 5-HT neurons integrate multiple long-range inputs

We first sought to determine whether LHb and mPFC inputs synapse onto distinct, non-overlapping populations of 5-HT DRN neurons or, alternatively, whether some subpopulations receive common input. To probe the integration of mPFC and LHb inputs at single 5-HT neurons, we employed a dual-color opsin approach (Klapoetke et al., 2014) wherein we bi-laterally injected adeno-associated viruses (AAVs) expressing Chronos (excited by blue light) in mPFC and Chrimson (excited by red light) in LHb (Fig. 1A). Using an *in vitro* slice physiology preparation, we recorded in whole-cell configuration from genetically identified 5-HT neurons (SERT-CRE-tdTomato mouse line; see Methods) primarily in the ventromedial and dorsomedial regions of the DRN. In voltage clamp mode (Vm=-70 mV), brief (1 ms) pulses of blue (473 nm) or red (550 nm) light were delivered to the slice through an optic fiber. In agreement with previous work (Geddes et al., 2016; Zhou et al., 2017), we observed reliable inward photocurrents (excitatory post-synaptic potentials, EPSCs) which were time-locked to optical stimulation (Fig. 1B), indicating the presence of long-range glutamatergic synaptic inputs from LHb or mPFC. Single 5-HT neurons showed a diversity of responses, responding to neither, one, or both wavelengths of light. Overall, 43.8% of 5-HT neurons responded to both wavelengths of light, 18.8% of neurons responded to neither, and the remainder responded to only one wavelength (Fig. 1C).

To rule out overlap between the excitation spectra of these two opsins, we delivered a set of patterned dual-color stimuli talking advantage of opsin desensitization, similar to Christoffel et al (2021) (see Methods). Briefly, if the same population of synapses expressing a single opsin are activated twice by paired-pulse stimulation, this should lead to release probability changes and measurable short-term dynamics, either facilitation due to presynaptic calcium accumulation, or depression due to receptor desensitization or vesicle depletion (Fig. S1A, top; Dobrunz & Stevens, 1997). However, if sequential pulses each activate distinct synaptic populations (*e.g.* populations expressing two spectrally distinct opsins), then no short-term dynamics should be possible. Consistent with this idea, we delivered paired-pulse stimulation at 20 Hz either with two red pulses (activation of the same synaptic population expressing Chrimson), or with a sequential red and blue pulse (activation of two synaptic populations expressing either Chrimson or Chronos). The red-red paired-pulse stimuli triggered prominent short-term depression (Fig. S1A, top), while the red-blue paired-pulse stimuli triggered minimal short-term dynamics, as assessed through comparison with the event amplitude of a solitary blue photostimulation (Fig. S1A, bottom). Paired comparison across the population indicated that the same neurons exhibited minimal dynamics with red-blue photostimulation (paired-pulse ratio close to 1) compared to red-red photostimulation (paired-pulse ratio < 1; Fig. S1B) This indicates an effective separation of the excitation spectra of both opsins in this subset of dual-responder neurons. Single-opsin controls (Chrimson in LHb alone; Fig. S1C-D) revealed that while blue light could occasionally activate the red opsin, our patterned dual-color stimulation accurately discriminated this condition, with similar depression dynamics observed in red-red paired-pulse and red-blue paired-pulse stimulation as expected (compare Fig. S1D with Fig. S1B). Together, these results indicate that potential misidentification of input specificity due to spectral overlap of complementary opsin pairs can be effectively screened with appropriately designed stimuli. While this overall strategy does not lend itself with ease to precise quantitative estimates, it nonetheless shows that at least a subpopulation of DRN 5-HT neurons receive joint long-range excitatory input from both mPFC and LHb.
196

Figure 1: Subthreshold and spike timing characteristics of long-range PFC and LHb input to 5-HT neurons. **A**: Experimental setup showing dual-color opsin strategy. **B**: Example voltage-clamp traces of individual 5-HT neurons responding only to red (top) and blue (bottom) photostimulation. **C**: Quantification of fraction of neurons responding to each light type (n=32). **D**: Example current-clamp traces of EPSPs of individual 5-HT neurons in response to photostimulation of PFC (red) or LHb (blue) inputs. **E-F**: Amplitude and decay (mean +- SE) of EPSPs in individual neurons in response to PFC (red, n=9 neurons) or LHb (blue, n=9 neurons) input (paired T-test). **G**: Example current-clamp traces of spiking responses of individual 5-HT neurons in response to LHb (blue) or PFC (red) photostimulation. **H**: Spike timing of individual neurons (rows) in response to LHb (blue, n=9 neurons) or PFC (red, n=9 neurons) photostimulation. **I-J**: Spike latency and jitter (mean +- SE) for individual neurons in response to LHb (blue, n=9 neurons) or PFC (red, n=9 neurons) photostimulation (paired T-test). **K**: Probability distribution of spike times in response to PFC photostimulation across all neurons (left) and plotted as a log-log relationship with line of best for power-law scaling (right). **L**: Probability distribution of spike times in response to LHb photostimulation across all neurons (left) and plotted as a log-log relationship with line of best fit for power-law scaling (right).

Subthreshold responses of DRN 5-HT neurons to mPFC and LHb inputs are similar
In principle, the information content encoded by distinct mPFC and LHb inputs to
individual neurons may be collapsed into a single low-dimensional code or, alternatively,
information from each input may be preserved. To address this question, we first investigated
synaptic features of PFC and LHb inputs to 5-HT neurons. We injected an AAV expressing
channelrhodopsin (ChR2) bilaterally into either LHb or mPFC in two groups of mice, and
conducted whole-cell recordings from identified 5-HT neurons (see Methods). Brief pulses of
blue light triggered fast inward currents, in keeping with previous reports showing
monosynaptic, glutamatergic connections between PFC and LHb axons and DRN 5-HT
neurons (Fig. S2A-B; Vm=-70mV). In current-clamp mode, photostimulation triggered time-
locked excitatory postsynaptic potentials (EPSPs) that were broadly similar between LHb and
mPFC inputs, although those induced from PFC inputs often displayed longer decay (Fig. 1D).
Indeed, while the mean amplitude and decay of EPSPs across cells were not statistically
different between inputs (Fig. 1D-F), the distribution of EPSP amplitudes and decays across
trials was slightly but significantly different between PFC and LHb evoked events (Fig. S2C-D).
Together, these results indicate that 5-HT neurons receive long-range inputs from LHb and
mPFC that are approximately matched in amplitude and timecourse at subthreshold
membrane potentials.
215
A spike-timing code discriminates input identity from mPFC and LHb in single 5-HT neurons
We next examined how the EPSPs from the PFC or the LHb are transformed into
spiking output in individual 5-HT neurons. We optogenetically stimulated long-range inputs
from either mPFC or LHb while recording from 5-HT neurons in current clamp, and examined in
details the resulting sparse spiking responses elicited by each input separately. By fine-tuning
direct current injection, we ensured: 1) comparable resting membrane potential (Fig. S2E), 2)
comparable fraction of trials eliciting a spike (Fig. S2F) between neurons and; 3) that
spontaneous spikes (*i.e.,* occurring before the optical stimulation) were not present. LHb
stimulation triggered low-latency spiking with very little variability (jitter) between trials, largely
consistent with theories of spike times encoding the derivative of inputs - in this case, EPSPs
with sharp rise-times (Mainen & Sejnowski, 1995). mPFC stimulation, in contrast, triggered
spikes with much longer latencies with high jitter (Fig. 1G-I). The latencies of the mPFC
triggered spikes, often lasting hundreds of milliseconds, were qualitatively similar across all 5-
HT neurons measured (Fig. 1H). In contrast, LHb-triggered spikes exhibited stereotyped, sharp
voltage trajectories to spike threshold that were highly reliable across trials, while in contrast
PFC-triggered spikes exhibited longer voltage trajectories that demonstrated less trial-to-trial
reliability (p=0.014; Fig. S2G-J). mPFC inputs yielded spike-times that were distributed with a
heavy right tail that could be well fit with a power-law decay (Fig. 1K), while LHb inputs
triggered spike-times that could also be fit with a power-law decay, but with a larger negative
exponent reflecting steeper decay (Fig. 1L). However, since spike-times spanned less than one
order of magnitude we could not formally distinguish this power-law decay from an exponential
decay (Fig. S2K-N). This is consistent with theories of fluctuation-driven activity regimes, where
spike timing depends on membrane potential and noise statistics (Gerstein & Mandelbrot,
1964; van Rossum et al., 2003).
Could differential stimulus-driven spiking for each input reflect an effective segregation
of each informational stream within populations of 5-HT neurons? One compatible theory is
that of synchrony-division multiplexing (Lankarany et al., 2019; Ratté et al., 2013; Friedrich et
al, 2004) suggesting that synchronous firing of neurons in a population can carry information
distinct from the time-averaged firing rate of individual neurons. Our data suggests that the
precisely time-locked response of many 5-HT DRN neurons to LHb inputs would trigger

synchronous spiking across the population. In contrast, mPFC input is expected to generate
asynchronous spiking of 5-HT neurons. These two firing patterns could, in principle, be
decoded by circuits downstream of 5-HT neurons. To explore this possibility, we first applied
linear classifiers to our spike timing data from each input, with the aim of determining whether
the identity of the input (LHb or mPFC) could be theoretically determined solely from the
spiking output of a population of 5-HT neurons. We generated synthetic training and test
datasets (5-fold cross-validation, see Methods) based on an unbiased sampling of
experimentally-derived spike-times from single trials for each input (Fig. 2A). By training a
support vector machine classifier and assessing performance on held-out data, we quantified
how spike timing-based decoding performance varied as a function of the number of spikes.
Surprisingly, we found that spike timing information provided by as few as 6 neurons allowed
good (>90%) decoding of the identity of the synaptic input driving these spikes (i.e., mPFC or
LHb) (Fig. 2B-C), indicating that synchrony/asynchrony-based decoding could be biologically
plausible even by areas receiving sparse innervation by 5-HT axons. We next determined
whether the EPSPs alone (i.e., subthreshold voltage amplitudes or decays) from both inputs
could be discriminated, and found poor decoding for both inputs, not exceeding 90% correct
classification with >30 observations (Fig. 2D-G). The poor discriminability between the EPSPs
triggered by PFC and LHb inputs onto 5-HT neurons stands in stark contrast to the robust
discriminability of 5-HT neuron's output (based on spike timing) induced by these same PFC
and LHb inputs. Thus, a latent cellular feature operant in 5-HT neurons is likely responsible for
transforming broadly indistinguishable synaptic inputs into spiking outputs with substantial
timing differences.
268
A Calcium channel-mediated conductance expressed in 5-HT neurons support differential spike
timing
We next sought to determine candidate cellular mechanisms in 5-HT neurons that could
generate reliable differences in spike timing between inputs. It has previously been shown that
low-threshold (T-type) calcium channels are highly expressed in several key subgroups of DRN
5-HT neurons (Templin et al., 2012; Okaty et al., 2020) and trigger rebound spiking in a subset
of putative 5-HT neurons (Burlhis & Aghajanian, 1987). We first examined their expression in
genetically identified 5-HT neurons and found that voltage steps to Vm = -40 mV triggered
delayed inward currents lasting several hundred milliseconds, which were blocked by
application of NiCl, a low-voltage calcium channel blocker (Fig. 3A; Lee et al., 1999, see
Methods). The inward current was voltage-dependent, with a protracted rise-time and decay
consistent with reported values for a low-threshold calcium channel (Fig. 3A-B; Perez-Reyes,
2003). Together, these data indicate that low-threshold calcium channels are expressed in 5-
HT neurons and are activated at depolarized voltages.
Next, we sought to examine how these low-threshold calcium channels modify
incoming synaptic inputs of variable kinetics. To parametrize this effect, we injected in
genetically identified 5-HT neurons synthetic EPSC waveforms which resembled incoming
EPSCs with varying decay kinetics (Fig. 3C, left). The injection of synthetic EPSCs with slow
decay kinetics led to variable voltage responses, both in terms of amplitude and decay, which
we hypothesized may be due to stochastic and transient calcium channel activation (Fig. 3C,
middle). Consistent with this hypothesis, bath application of low concentrations of NiCl
selectively blocked the slow and high amplitude subset of voltage responses triggered by the
slow synthetic EPSCs (Fig. 3C, right; also see Supp. Fig. S3). In contrast, charge-normalized
synthetic EPSCs with faster decays trigered low variability voltage responses that lacked
prolonged, high-amplitude events and that were not affected by application of NiCl (Fig. 3C;
also see Supp. Fig. S3).

Figure 2: Decoding long-range inputs by spike-timing and subthreshold features in 5-HT populations. **A**: Schematic showing approach for generating synthetic datasets and applying a support vector machine linear classifier to data (5-fold cross-validation). **B**: Example synthetic datasets of spike times from PFC (red) or LHb (blue). Each vertical line represents one example dataset of n=10 spike times. **C**: Decoding accuracy of PFC (red) and LHb (blue) synthetic datasets from linear classifier as a function of dataset size (neurons). Dotted line indicates chance decoding performance (50%). **D**: Example synthetic datasets of EPSP amplitude from PFC (red) or LHb (blue). Each vertical line represents one synthetic dataset of n=10 EPSP amplitudes. **E**: Decoding accuracy of PFC (red) and LHb (blue) synthetic datasets from linear classifier as a function of dataset size (neurons). Dotted line indicates chance decoding performance (50%). **F**: Example synthetic datasets of EPSP decay from PFC (red) or LHb (blue). Each vertical line represents one synthetic dataset of n=10 EPSP amplitudes. **G**: Decoding accuracy of PFC (red) and LHb (blue) synthetic datasets from linear classifier as a function of dataset size (neurons). Dotted line indicates chance decoding performance (50%).

Since the activation of calcium channels and other ionic conductances can affect the
timing of spikes (Cain & Snutch, 2013; van Rossum et al., 2003), we hypothesized that these
channels might play a similar role in regulating spike timing in 5-HT neurons. Blocking ionic
conductances should, in principle, increase the neuron's membrane resistance and make it
easier for a fixed magnitude of current input to depolarize the neuron sufficiently to trigger
spikes. However, we found, surprisingly, that bath application of NiCl could reduce or even
eliminate spiking response for a fixed perithreshold current input (Fig. 3D). This may indicate
complex interactions between voltage events and activation/inactivation kinetics of this
calcium conductance, as well as spatial profiles of channel expression throughout the dendrite
and soma. We used synthetic EPSCs to explore how the production of spikes is tuned by
calcium channels across the parameter space of event kinetics. The decay kinetics and
amplitudes of the injected synthetic EPSCs were systematically varied while we recorded the
voltage response of 5-HT neurons. We found, consistent with our earlier findings (Fig. 3D), that
NiCl application increased the amplitude of the synthetic EPSC required to elicit spiking
behavior, especially at long decay tau values (Fig. 3E) and reduced spiking probability across
all neurons at the same fixed EPSC amplitude (Fig. 3F). These results highlight an important
role of this calcium conductance in regulating the spiking output of 5-HT neurons.
We next sought a mechanistic explanation for how the activation of this peri-threshold
conductance could generate variation in spike timing, thus plausibly modulating network
synchrony. To this end, we adopted a dynamical approach and determined how voltage noise
statistics – a determinant of spike timing- are modulated by these channels. We conducted
experiments where finely spaced, long (20s) current steps were injected into identified 5-HT
neurons before and after application of NiCl (Fig. 3G). Throughout these experiments, a
cocktail of synaptic blockers was applied (see Methods). To focus on sub- and perithreshold
regimes, spikes were identified and removed from the recordings using a fractional first-
derivative method (Fig. 3G; see Methods; Trinh et al., 2019; Azouz and Gray, 2000). At
perithreshold membrane potentials (defined by the first depolarizing current step eliciting any
action potentials), we observed prominent voltage noise as well as depolarizing, non-spike-
generating, events lasting ~1s (Fig. 3I-J, top). We suspected that these slow depolarizing
events, as well as fast voltage noise, were partially generated by stochastic activation and
inactivation of these low-threshold calcium channels. Consistent with this, application of NiCl
modulated the amplitude of both the slow events and fast voltage noise, and decreased the tail
size of the membrane voltage distribution (Fig. 3I-J, bottom) without altering spike threshold
per se (Fig. 3H).
To quantify these observations across the population of recorded 5-HT neurons, we
measured the skewness (third central moment, indicating asymmetry) and kurtosis (fourth
central moment, indicating tailedness) of the membrane potential distribution, before and after
application of NiCl. Before NiCl application, the membrane voltage distribution had skewness
values above 0, indicating a right tail, that was not affected by NiCl application (Fig. 3K). The
distribution's kurtosis was, however, significantly reduced by NiCl application, consistent with
this calcium conductance being causally implicated in the generation of depolarized voltage
excursions (Fig. 3L). To determine whether these effects were localized to perithreshold voltage
regime, we considered the voltage dynamics during the first subthreshold current step (i.e.,
without spikes) and found no effect of NiCl bath application on kurtosis or skewness (Supp.
Fig. S4). This indicates that the effects of low-threshold calcium channel activation on
membrane dynamics is preponderant at perithreshold membrane potentials. Finally, we
performed a fast Fourier transform of the time-varying perithreshold membrane potential (again
with spikes removed) to determine how calcium channel blockade affected the frequency
content of the noise. This analysis revealed a NiCl-dependent boost in frequencies in a 0.1-

Figure 3: An inward calcium conductance in 5-HT neurons shapes EPSPs and contributes to membrane noise. **A**: Example traces in voltage-clamp showing an inward conductance at -40mV (right) but not -60mV (left) that is blocked by bath application of NiCl (blue trace). **B**: Voltage-dependence of the putative inward calcium conductance (n=3 neurons). **C**: Synthetic EPSCs injected in current-clamp into 5-HT neurons (left) produce transient voltage excursions (middle) blocked by bath application of NiCl (right). **D**: Example trace showing modulation of synthetic EPSC-triggered spiking by bath application of NiCl. **E**: Relationship between synthetic EPSC amplitude required to elicit a spike, and synthetic EPSC tau, in baseline conditions (black) and after bath application of NiCl (blue) (n=5 neurons, bars represent S.D.). **F**: Relationship between probability of obtaining an action potential across neurons, and synthetic EPSC tau, in baseline conditions (black) and after bath application of NiCl (blue) (n=5 neurons). **G**: Experimental setup for membrane noise experiments presented in **H-M** (left), and

example of spike threshold estimation using fractional first-derivative method (right). **H**: Spike threshold is not modulated by bath application of NiCl (n=8 neurons, paired T-test). **I**: Example traces of a single neuron recorded in current-clamp during injection of constant perithreshold depolarizing current, before (top) and after (bottom) application of NiCl. Red arrows depict large voltage excursions ('blips'). **J**: Probability distribution of membrane potentials for an example neuron before (top) and after (bottom) application of NiCl. **K**: Membrane potential distribution skewness is not significantly affected by bath application of NiCl (n=7, paired T-test). **L**: Membrane potential distribution kurtosis is significantly reduced by bath application of NiCl (n=8, paired T-test). **M**: Relative spectral power of calcium conductance, calculated by subtracting baseline membrane potential spectral power from post-NiCl washon membrane potential spectral power (n=8 neurons, mean shown.)

10Hz window (Fig. 3M). Collectively, these results indicate that low-threshold calcium channels
robustly modulate membrane noise around spike threshold in 5-HT neurons.
We next naturally hypothesized that this perithreshold calcium conductance is causally
involved in generating mPFC-evoked delayed spiking in 5-HT neurons. We thus repeated the
photostimulation experiments of mPFC axons in the DRN and found that it triggered high-
latency spikes (Fig. S5A, left), replicating our previous results (Fig. 1G). We then applied in
these same recordings voltage steps to -45mV that triggered inward currents that were
abolished by bath application of NiCl (Fig. S5B). We then resumed optogenetic stimulation of
mPFC axons and found that the resultant spikes displayed noticeably shorter latency and less
jitter (Fig. S5A, right; Fig. S5C). The effect of bath application of NiCl on reducing spike latency
was statistically significant across the population of 5-HT neurons recorded (Fig. S5D-E,
p=0.045, Kolmogorov-Smirnov test on spike times compiled across the population). Taken
together, these results establish a causal relationship between the calcium channel
conductance expressed in 5-HT neurons, and the delayed spike timing behavior evoked by
long-range mPFC input.
359
Dynamical regimes for synchronous and asynchronous spike generation
We next employed stochastic, biophysically grounded simulations to obtain a
mechanistic understanding of how calcium channels could regulate the initiation time of action
potentials, and thus modulate spike synchrony at the population level. We first considered a
simplified scenario where Ornstein-Uhlenbeck noise (simulating barrages of synaptic input
and/or intrinsic noise, see Methods; Destexhe et al., 2001) was injected to a toy neuron model
with subthreshold membrane potential. This produced a voltage signal that evolved over time
in a pseudo-random fashion (Fig. 4A). The time at which the voltage signal first crossed a
depolarized spike threshold (called the first-passage crossing time; yellow dot in Fig. 4A) was
recorded over 20,000 stochastic, independent simulations, generating a probability distribution
of spike times (Fig. 4B). These simulations effectively correspond to a population of 5-HT
neurons receiving a fixed synaptic input, and responding stochastically due to cell-intrinsic
mechanisms. This approach allowed us to ask how the spike timing distributions we observed
could be related to simple interactions between a stochastic noise process with defined
characteristics, and a fixed spike threshold.
Guided by our empirical findings showing that calcium channels altered the temporal
structure of membrane noise, we first used *in silico* simulations to ask how the rate of noise
fluctuations affected spike timing distributions. Ornstein-Uhlenbeck noise with a fast time
constant (tau=0.01) generated rapidly fluctuating voltage signals that to reach spike initiation
very quickly, with the spike times exponentially distributed and typically in the range of tens of
milliseconds (Fig. 4A-B). Ornstein Uhlenbeck noise with a slow time constant (tau=0.5), and
with a fixed membrane potential distribution compared to fast noise (Fig. 4C), generated time-
varying signals that evolved more slowly and tended to show longer delays before reaching
spike threshold. At the population level, these spike times were in the range of hundreds of
milliseconds to seconds, and were power-law distributed (Fig. 4E). The spectral characteristics
of these signals, revealed through a fast Fourier transform (Fig. 4F), showed that simulations
with a slower noise time constant were specifically enriched in very low frequencies (0.5-2Hz).
Together, these results provide a framework to understand how the distribution of spike
initiation times depends on intracellular noise features. In particular, at subthreshold membrane
potentials, noise with fast fluctuations tends to produce short spike latencies which are
exponentially distributed over time, while noise with slower fluctuations produces more
protracted spike latencies which are power-law distributed.

Figure 4: The statistics of membrane noise shapes stochastic spike timing in 5-HT neurons. **A**: Example simulation showing fast membrane potential noise represented by a fast (tau=0.01s) Ornstein-Uhlenbeck process. The dotted line represents spike threshold and the yellow dot represents the time of first-passage-crossing. **B**: Distribution of first-spike latencies across 20,000 independent initializations of OU noise. Black line represents an exponential fit. **C**: Distribution of simulated membrane voltages for fast and slow Ornstein-Uhlenbeck noise shown in **A** and **D**. **D**: Example simulation showing slow membrane potential noise represented by a slow (tau=0.5s) Ornstein-Uhlenbeck process. The dotted line represents spike threshold and the yellow dot represents the time of first-passage-crossing. **E**: Distribution of first-spike latencies across 20,000 independent initializations of OU noise. Black line represents a power-law fit. **F**: Spectral power of simulated membrane potential for fast and slow noise processes. **G**: Example simulations showing fast membrane potential noise (tau=0.01s Ornstein Uhlenbeck process, black) and fast membrane potential noise with calcium noise spectral power added in the frequency domain (pink). **H**: Distribution of first-spike latencies across 20,000 independent initializations of OU noise (black), and with increasing calcium noise spectral power added (pink). **I**: Spectral power of simulated membrane potential for fast noise process alone and with increasing calcium noise spectral power added.

To provide a more direct comparison to the membrane noise observed in 5-HT
neurons, we asked whether the observed spectral characteristics of this noise (Fig. 3M) could
modulate spike latency distributions in a reliable manner. We thus first added the spectral
components of the experimentally-observed calcium channel-dependent noise to the fast-
fluctuating Ornstein-Uhlenbeck noise process presented in Fig. 4A. Consistent with its
prominent low-frequency component, the addition of the spectral components of the calcium
conductance-dependent noise generated slow fluctuations in the simulated signal (Fig. 4G, I).
We determined how adding scaled components of the noise ranging from 0.5x to 10x (*i.e.,*
weak to strong activation of calcium channels), affected spike initiation times. Adding the noise
dramatically increased the average spike latency time, and prolonged the straight-line
component of the spike latency distribution (in a log-log plot) to span two orders of magnitude,
consistent with a power-law distribution of latencies (Fig. 4H). These results thus demonstrate
that distinct spike latency distributions at the population level can be readily explained by
stochastic noise processes occurring in single neurons, allowing these individual neurons to
toggle between distinct latency 'modes' through activation or inactivation of ionic
conductances. Ultimately, this suggests a mechanism by which populations of 5-HT neurons
receiving overlapping synaptic inputs can effectively segregate these inputs through dynamic
generation of synchronous (short-latency) or asynchronous (long-latency) spiking, utilizing the
simple building blocks of stochastic noise processes and ionic conductances with defined
spectral characteristics.
412
Discussion
In this work, we demonstrate that a subpopulation of individual 5-HT neurons in DRN
receives strong excitatory drive from both mPFC and LHb, and information from these two
inputs can be multiplexed at the population level by cell-intrinsic regulation of spike timing.
While synaptic inputs from LHb produce time-locked, low-latency spikes leading to population
synchrony, mPFC inputs produce highly jittered, asynchronous spikes. We show that these
two informational sources can be theoretically decoded by downstream circuits receiving
sparse 5-HT afferents. Finally, we demonstrate that the delayed spiking pattern from mPFC
inputs arise from a set of simple ionic mechanism: perithreshold calcium channels increase
membrane noise and trigger stochastic, delayed spike threshold crossings. Together, these
findings indicate that long-range inputs from mPFC and LHb can be multiplexed in local
serotonergic circuits by cell-intrinsic regulation of spike synchrony.
425
DRN 5-HT neurons multiplex information from multiple sources
Our results add to a body of work showing signals may be represented by precisely
timed activity patterns in 5-HT neurons, including during complex behaviors. Paquelet et al.
(2022) undertook miniscope calcium imaging of populations of 5-HT neurons during
emotionally salient behaviors, finding highly correlated (synchronous) ensembles of neurons
responding to sucrose or footshocks alongside non-selective ensembles. Li et al. (2016)
demonstrated that 5-HT neurons encode reward signals, including a subpopulation that
exhibits increased firing rate with millisecond precision. Cohen et al. (2015) reported that 5-HT
neurons exhibited tonic firing rates that tracked block reward size, alongside phasic responses
to direct rewards and punishments. These findings, taken alongside the results reported here,
suggest a model where information about ongoing environmental stimuli, such as rewards or
punishments, is encoded as synchronous population activity in DRN microcircuits, while
higher-order task features or environmental statistics may be represented as uncorrelated firing
of 5-HT neurons. Our dynamical model provides a putative explanation for both the
uncorrelated spiking and synchrony shown to be present in populations of 5-HT neurons *in*

vivo (Cohen et al., 2015; Paquelet et al., 2022). We expect that future work will help address
the question of how synchronous and asynchronous activity evolves in larger populations of 5-
HT neurons, and whether effective parsing of spiking in this manner provides meaningful
information about task variables or internal variables.
445
Calcium channel-dependent noise may provide a generalizable mechanism for separating
inputs through differential spike timing
Our results demonstrate that 5-HT neurons exhibit calcium conductances that are
activated in the perithreshold regime and add a broadband low-frequency component to
membrane noise. How do synaptic inputs from mPFC preferentially activate these calcium
conductances, while LHb inputs do not? Here we entertain two possibilities. First, a dendritic
gradient of calcium channel expression, combined with localized targeting of calcium channel-
rich dendritic segments by mPFC inputs, might produce mPFC-specific activation of calcium
conductances. Indeed, differential targeting of dendritic segments by long-range inputs can
profoundly shape spike output and integration in other brain areas (MacAskill et al., 2012),
indicating they are a plausible substrate for the effects reported here. A second possibility is
that mPFC synaptic inputs to 5-HT neuron dendrites could be more spatially clustered than
LHb synaptic inputs, allowing mPFC inputs to exclusively trigger to dendritic calcium events
when activated synchronously. Previous work mainly in cortex and hippocampus has shown
that clustered synaptic activation generates several dendritic non-linearities, both in terms of
calcium signaling and membrane depolarization (Lee et al., 2016; Higley & Sabatini, 2008; Mel,
1993; Losonczy & Magee, 2006). Future studies will be needed to disambiguate these and
other possibilities.
The findings reported here add to previous studies that link the actions of specific ionic
conductances to cellular and population spiking dynamics. For instance, electrosensory
neurons in the weakly electrical fish exhibit Na^+ channel noise, including "blips" that trigger
action potentials (Marcoux et al., 2016). This noise was shown to boost slow excitatory
synaptic inputs and the noise-triggered spikes enacted a stochastic process that prevented
spike synchronization at a population level. The blips, somewhat analogous to the membrane
noise observed here in DRN 5-HT neurons, were likely generated by clustered dendritic Na^+
channels (Turner et al., 1994). We thus propose that intrinsic neuron noise driven by
spontaneous opening of voltage-dependent Na^+ or Ca^{2+} channels may be a common
biophysical mechanism for decorrelating spiking outputs, even when the noisy cells are
embedded in neuronal networks that perform different computations.
475
Decoding of DRN input streams through differential 5-HT release in downstream brain areas
We used linear decoders to demonstrate that the observed spike timing patterns can
theoretically be decoded by downstream circuits innervated by 5-HT axons. There are multiple
biologically plausible mechanisms that could provide such a decoding capability. First, the
short-term facilitation of 5-HT release (Lynn et al., 2022) could functionally segregate high-
frequency and low-frequency spike trains, albeit from the same neuron, by differential 5-HT
release dynamics. Such a short-term facilitation decoding mechanism has been shown to be
theoretically viable in the electrosensory system of weakly electric fish (Middleton et al., 2011).
Here, the viability of this decoding scheme would rely on LHb or PFC inputs changing the
spiking statistics of single neurons at fine timescales corresponding to changes in input
patterns. Recent work, showing that 5-HT neurons can emit action potentials transiently at high
rates of ~20-60Hz, appears to provide a plausible substrate for such a decoding mechanism (Li
et al., 2016). A second biologically feasible mechanism involves a 5-HT affinity-based decoding
scheme, determined by the differential 5-HT receptor subtype expression across cortical

excitatory versus inhibitory neurons. In cortex, inhibitory 5-HT receptors of the 5-HT1A subtype
are broadly expressed in pyramidal neurons, while excitatory 5-HT2A receptors are strongly
expressed in apical dendrites of at least a subset of pyramidal neurons (Beique et al., 2007;
Jakab et al., 1999), while some neurons express both subtypes (Araneda & Andrade, 1991;
Avesar & Gulledge, 2012). Moreover, fast-spiking and somatostatin interneurons can either
express excitatory 5-HT2A or inhibitory 5-HT1A receptors, but typically not both (for review,
see Puig et al., 2011). The differential affinity of 5-HT1A and 5-HT2A receptors for 5-HT
(Peroutka et al., 1979; Nichols & Nichols, 2008), suggest a decoding scheme where 5-HT1AR-
expressing neurons may be inhibited by both synchronous and asynchronous 5-HT axon
release events, while 5-HT2AR-expressing interneurons and pyramidal neurons expressing 5-
HT2ARs may be depolarized by only highly synchronous 5-HT release events due to lower
receptor affinity. Such receptor binding rules in cortical microcircuitry could in principle sustain
demultiplexing of information carried by 5-HT axon synchrony. Further work is needed to
discriminate between these two hypothesized decoding schemes, and to establish the precise
network mechanisms by which such decoding schemes would function.
Taken together, our results demonstrate that populations of 5-HT neurons receiving
multiple sources of long-range synaptic input can effectively multiplex information, through
predominantly cell-intrinsic mechanisms that regulate spike synchrony at the population level.
These cell-intrinsic mechanisms involve stochastic membrane noise processes and differential
calcium channel activation. More broadly, these results suggest that a high-dimensional
population coding scheme such as synchrony-division multiplexing can plausibly arise from the
autonomous, stochastic noise processes occurring in individual neurons (assuming synaptic
volleys well separated in time), even in the absence of strong network-level interactions that
are conventionally thought to shape information representations in cortex.
The effects of synchronous versus asynchronous activation of DRN 5-HT cells on
individual pyramidal cells and interneurons will be amplified by local recurrent cortical circuitry.
Further, DRN has widespread projections to cortex and the extensive reentrant cortical
pathways will extend the effects of such inputs to cortical areas not directly associated with
persistent cognitive states (PFC) or transient negative reward prediction errors (LHb).
Extracting the signatures of the differential DRN 5-HT modulation of cortical activity from the
baseline spiking activity generated by local and long-range cortical circuitry will be a major
challenge for future studies.
522
523 **References**

524 G.R. Poe, et al., Locus coeruleus : A new look at the blue spot. *Nat Rev Neurosci* **21**, 644-659
525 (2020).
526
527 M.R. Gielow & L. Zaborszky, The input-output relationship of the cholinergic basal forebrain.
528 *Cell Rep* **18**, 1817-1830 (2017).
529
530 L. Faget et al., Afferent inputs to neurotransmitter-defined cell types in the ventral tegmental
531 area. *Cell Rep* **15**, 2796-2808 (2016).
532
533 D.E. Rumelhart, G.E. Hinton, R.J. Williams, Learning representations by back-propagating
534 errors. *Nature* **323**, 533-536 (1986).
535
536 R. Naud, H. Sprekeler, Sparse bursts optimize information transmission in a multiplexed neural
537 code. *Proc. Natl. Acad. Sci. U.S.A.* **115,** E6329-E6338 (2018).
538

539 A. Payeur, J. Guerguiev, F. Zenke, B.A. Richards, R. Naud, Burst-dependent synaptic plasticity
540 can coordinate learning in hierarchical circuits. *Nat Neurosci* **24**, 1010–1019 (2021).
541
542 A.-M Oswald, M.J. Chacron, B. Doiron, J. Bastian, L. Maler. Parallel processing of sensory
543 input by bursts and isolated spikes. *J Neurosci* **24**, 4351– 4362. (2004).
544
545 S.E. Clarke, L. Maler. Feedback synthesizes neural codes for motion. *Curr Biol* **27**, 1356–1361.
546 (2017).
547
548 M. Lankarany, D. Al-Basha, S. Ratté, S.A. Prescott, Differentially synchronized spiking enables
549 multiplexed neural coding. *Proc. Natl. Acad. Sci. U.S.A. 116,* 10097–10102 (2019).
550
551 S. Ratté, S. Hong, E. De Schutter, S.A. Prescott, Impact of neuronal properties on network
552 coding: roles of spike initiation dynamics and robust synchrony transfer. *Neuron* **78**, 758-772
553 (2013)
554
555 R.W. Friedrich, C.J. Habermann, G. Laurent, Multiplexing using synchrony in the zebrafish
556 olfactory bulb. *Nat Neurosci* **7**, 862-871 (2004).
557
558 C. Stringer et al., Spontaneous behaviors drive multidimensional, brainwide activity. *Science*
559 **364**, eaav7893 (2019).
560
561 R. Nogueira, C.C. Rogers, R.M. Bruno, S. Fusi, The geometry of cortical representations of
562 touch in rodents. *Nat Neurosci* **26**, 239-250 (2023).
563
564 B. Doiron, C. Laing, A. Longtin, L. Maler. Ghostbursting : A novel neuronal burst mechanism. *J*
565 *Comput Neurosci* **12**, 5-25. (2002).
566
567 C. Laing, B. Doiron, A. Longtin, L. Noonan, R.W. Turner, L. Maler. Type I burst excitability. *J*
568 *Comput Neurosci* **14**, 329-342. (2003).
569
570 Z.F. Mainen, T.J. Sejnowski, Reliability of spike timing in neocortical neurons. *Science* **268**,
571 1503-1506 (1995)
572
573 G.B. Ermentrout, R.F. Galan, N.N. Urban, Reliability, synchrony and noise. *Trends in Neurosci*
574 **31**, 428-434 (2008).
575
576 S. Schreiber, J.-M. Fellous, P. Tiesinga, T.J. Sejnowski, Influence of Ionic Conductances on
577 Spike Timing Reliability of Cortical Neurons for Suprathreshold Rhythmic Inputs. *J*
578 *Neurophysiol* **91**, 194-205 (2004).
579
580 C.M. Marcoux, S.E. Clarke, W.H. Nesse, A. Longtin, L. Maler. Balanced ionotropic receptor
581 dynamics support signal estimation via voltage- dependent membrane noise. *J Neurophysiol*
582 **115**, 530-545 (2016).
583
584 M.A. Kamaleddin et al., Physiological noise facilitates multiplexed coding of vibrotactile-like
585 signals in somatosensory cortex. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2118163119 (2022).
586

587 J. Ren *et al*., Anatomically Defined and Functionally Distinct Dorsal Raphe Serotonin Sub-
588 systems. *Cell* **175** 472–487 (2018).
589
590 M.R. Warden *et al*, A prefrontal cortex–brainstem neuronal projection that controls response to
591 behavioural challenge. *Nature* **492**, 428-432 (2012).
592
593 B. Weissbourd *et al.,* Presynaptic Partners of Dorsal Raphe Serotonergic and GABAergic
594 Neurons. *Neuron* **83** 645–662 (2014).
595
596 I.P. Dorocic *et al*., A Whole-Brain Atlas of Inputs to Serotonergic Neurons of the Dorsal and
597 Median Raphe Nuclei. *Neuron* **83**, 663–678 (2014).
598
599 S.D. Geddes *et al*., Target-specific modulation of the descending prefrontal cortex inputs to the
600 dorsal raphe nucleus by cannabinoids. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 5429-5434 (2016).
601
602 L. Zhou *et al*., Organization of Functional Long-Range Circuits Controlling the Activity of
603 Serotonergic Neurons in the Dorsal Raphe Nucleus. *Cell Reports* **18**, 3018–3032 (2017).
604
605 E.K. Miller, J.D. Cohen., An integrative theory of prefrontal cortex function. *Annu. Rev. Neurosci*
606 **24**, 167–202 (2001).
607
608 B. Bari *et al*., Stable Representations of Decision Variables for Flexible Behavior. *Neuron* **103**,
609 922–933 (2019).
610
611 M. Matsumoto, O. Hikosaka, Lateral habenula as a source of negative reward signals in
612 dopamine neurons. *Nature* **447**, 1111-1115 (2007).
613
614 E. Bromberg-Martin, O. Hikosaka, Lateral habenula neurons signal errors in the prediction of
615 reward information. *Nat Neurosci* **14**, 1209-1216 (2011).
616
617 N.C. Klapoetke *et al*., Independent optical excitation of distinct neural populations. *Nat*
618 *Methods* **11**, 338-346 (2014).
619
620 D.J. Christoffel *et al.,* Selective filtering of excitatory inputs to nucleus accumbens by
621 dopamine and serotonin. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2106648118 (2021).
622
623 L.E. Dobrunz & C.F. Stevens. Heterogeneity of release probability, facilitation, and depletion at
624 central synapses. *Neuron* **18**, 995–1008 (1997).
625
626 V.C. Caruso *et al.,* Single neurons may encode simultaneous stimuli by switching between
627 activity patterns. *Nat Comms* **9**, DOI: 10.1038/s41467-018-05121-8 (2018).
628
629 A. Kuhn, A. Aertsen, S. Rotter, Neuronal integration of synaptic input in the fluctuation-driven
630 regime. *J Neurosci* **24**, 2345–2356 (2004).
631
632 G.L. Gerstein, B. Mandelbrot, Random walk models for the spike activity of a single neuron.
633 *Biophys J* **4**, 42-68 (1964).
634

635 M.C.W. van Rossum, B.J. O'Brien, R.G. Smith, Effects of noise on the spike timing precision of
636 retinal ganglion cells. *J Neurophysiol* **89**, 2406-2419 (2003).
637
638 E.F. Harkin *et al*., Temporal derivative computation in the dorsal raphe network revealed by an
639 experimentally-driven augmented integrate- and-fire modeling framework. *eLife*,
640 DOI:https://doi.org/10.7554/eLife.72951 (2023).
641
642 T.M. Burlhis, G.K. Aghajanian, Pacemaker potentials of serotonergic dorsal raphe neurons:
643 Contribution of a low-threshold Ca2+ conductance. *Synapse* **1**, 582-588 (1987).
644
645 J.S. Templin, S.J. Bang, M. Soiza-Reilly, C.B. Berde, K.G. Commons, Patterned expression of
646 ion channel genes in mouse dorsal raphe nucleus determined with the Allen Mouse Brain Atlas.
647 *Brain Res* **1457**, 1-12 (2012).
648
649 B.W. Okaty *et al.,* A single-cell transcriptomic and anatomic atlas of mouse dorsal raphe Pet1
650 neurons. *eLife* **9**, e55523 (2020).
651
652 J.-H. Lee, J.C. Gomora, L.L. Cribbs, E. Perez-Reyes, Nickel block of three cloned T-type
653 calcium channels: low concentrations selectively block 1H. *Biophys J* **77**, 3034–3042 (1999).
654
655 E. Perez-Reyes, Molecular physiology of low-voltage-activated T-type calcium channels.
656 *Physiol Rev* **83**, 117-161 (2003).
657
658 S.M. Cain, T.P. Snutch, T-type calcium channels in burst-firing, network synchrony, and
659 epilepsy. *Biochimica et Biophysica Acta* **1828,**1572–1578 (2013).
660
661 A.-T. Trinh, S.E. Clarke, E. Harvey-Girard, L. Maler, Cellular and network mechanisms may
662 generate sparse coding of sequential object encounters in hippocampal-like circuits. *eNeuro* **6**,
663 ENEURO.0108-19.2019 (2019).
664
665 R. Azouz, C.M. Gray, Dynamic spike threshold reveals a mechanism for synaptic coincidence
666 detection in cortical neurons in vivo. *Proc. Natl. Acad. Sci. U.S.A.* **97,** 8110–8115 (2000).
667
668 A. Destexhe, M. Rudolph, J.-M. Fellous, T.J. Sejnowski, Fluctuating synaptic conductances
669 recreate in vivo-like activity in neocortical neurons. *Neuroscience* **107**, 13-24 (2001).
670
671 G.E. Paquelet *et al*., Single-cell activity and network properties of dorsal raphe nucleus
672 serotonin neurons during emotionally salient behaviors. *Neuron* **110**, 1-16 (2022).
673
674 Y. Li *et al*., Serotonin neurons in the dorsal raphe nucleus encode reward signals. *Nat Comms*
675 **7**, 10503 (2016).
676
677 J.Y. Cohen, M.W. Amoroso, N. Uchida, Serotonergic neurons signal reward and punishment on
678 multiple timescales. *eLife* **4**, e06346 (2015).
679
680 Y.-P. Zhang, T.G. Oertner, Optical induction of synaptic plasticity using a light-sensitive
681 channel. *Nature Methods* **4**, 139-141 (2007).
682

683   A.F. MacAskill, J.P. Little, J.M. Cassel, A.G. Carter, Subcellular connectivity underlies pathway-
684   specific signaling in the nucleus accumbens. *Nat Neurosci* **15**, 1624-1626 (2012).
685
686   K.F.H. Lee, C. Soares, J.-P. Thivierge, J.-C. Béïque. Correlated Synaptic Inputs Drive Dendritic
687   Calcium Amplification and Cooperative Plasticity during Clustered Synapse Development.
688   *Neuron* **89**, 784–799 (2016).
689
690   M.J. Higley, B.L. Sabatini. Calcium Signaling in Dendrites and Spines: Practical and Functional
691   Considerations. *Neuron* **59**, 902-913 (2008).
692
693   B.W. Mel. Synaptic integration in an excitable dendritic tree. *J Neurophysiol* **70**, 1086-1101
694   (1993).
695
696   A. Losonczy, J.C. Magee, Integrative properties of radial oblique dendrites in hippocampal CA1
697   pyramidal neurons. *Neuron* **50**, 291-307 (2006).
698
699   R.W. Turner, L. Maler, T. Deerinck, S.R. Levinson, M.H. Ellisman. TTX-Sensitive Dendritic
700   Sodium Channels Underlie Oscillatory Discharge in a Vertebrate Sensory Neuron. *J Neurosci*
701   **14**, 6453-6471 (1994).
702
703   M.B. Lynn, S. Geddes, M. Chahrour, S. Maillé, E. Harkin, E. Harvey-Girard, S. Haj-Dahmane, R.
704   Naud & J.-C. Beique. A slow 5-HT1AR-mediated recurrent inhibitory network in raphe
705   computes contextual value through synaptic facilitation. *Biorxiv*
706   doi:https://doi.org/10.1101/2022.08.31.506056 (2022).
707
708   J.W. Middleton, N. Yu, A. Longtin & L. Maler. Routing the flow of sensory signals using plastic
709   responses to bursts and isolated spikes: Experiment and theory. *J Neurosci* **31**, 2461-2473
710   (2011).
711
712   J.-C. Béïque, M. Imad, L. Mladenovic, J.A. Gingrich, R. Andrade, Mechanism of the 5-
713   hydroxytryptamine 2A receptor-mediated facilitation of synaptic activity in prefrontal cortex.
714   *Proc. Natl. Acad. Sci. U.S.A.* **104**, 9870–9875 (2007).
715
716   R.L. Jakab, P.S. Goldman-Rakic, 5-Hydroxytryptamine2A serotonin receptors in the primate
717   cerebral cortex: Possible site of action of hallucinogenic and antipsychotic drugs in pyramidal
718   cell apical dendrites. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 735–740 (1999).
719
720   R. Araneda, R. Andrade. 5-Hydroxytryptamine2 and 5-Hydroxytryptamine1A receptors mediate
721   opposing responses on membrane excitability in rat association cortex. *Neurosci* **44**, 399-412
722   (1991).
723
724   D. Avesar, A.T. Gulledge. Selective serotonergic excitation of callosal projection neurons.
725   *Front. Neur. Circuits* **6**, 12 (2012).
726
727   M.V. Puig, A.T. Gulledge, Serotonin and Prefrontal Cortex Function: Neurons, Networks, and
728   Circuits. *Mol Neurobiol* **44**, 449–464 (2011).
729

S.J. Peroutka, S.H. Snyder, Multiple Serotonin Receptors: Differential Binding of [3H]5-
Hydroxytryptamine, [3H]Lysergic Acid Diethylamide and [3H]Spiroperidol. *Mol Pharmacol* **16**,
687-699 (1979).

D.E. Nichols, C.D. Nichols, Serotonin receptors. *Chem Rev* **108**, 1614–1641 (2008).

Ting, J.T., Lee, B.R., Chong, P., Soler-Llavina, G., Cobbs, C., Koch, C., Zeng, H. & Lein, E.
(2018). Preparation of Acute Brain Slices Using an Optimized N-Methyl-D-glucamine Protective
Recovery Method. J. Vis. Exp. (132), e53825, doi:10.3791/53825.

Gillespie, D.T. (1996). Exact numerical simulation of the Ornstein-Uhlenbeck process and its
integral. *Phys Rev E* 54(2):2084-2091.

Methods

Animal use

SERTcre: B6.Cg-Tg(Slc6a4- cre)ET33Gsat/Mmcd

R26R-TdT: Rosa26R-TdTomato fluorescent protein (TdT) reporter mice

All experiments and procedures were performed in accordance with approved procedures and guidelines set forth by the University of Ottawa Animal Care and Veterinary Services. For this study, SERTcre mice were crossed with the R26R line to generate a SERTcre-TdT mouse line. Mice were kept on a 12:12 hour light/dark cycle, with access to food and water ad libitum unless otherwise noted. A mix of male and female SERTcre-TdT mice were used.

Slice physiology

Stereotaxic Injections

All experiments and procedures were performed in accordance with approved procedures and guidelines set forth by the University of Ottawa Animal Care and Veterinary Services. For stereotaxic injections, mice were injected with 0.05 mg/kg buprenorphine and anesthetized by inhalation of isoflurane. Injections were performed using either a 10 μl Hamilton syringe with a 33 gauge needle, or a Nanoject machine with a glass micropipette. For all injections, 0.2 - 0.3 μl of virus was injected per coordinate, at a rate of 0.1 – 0.2 ul per minute. Stereotaxic co-ordinates are as follows: LHb (from Bregma skull landmark, in mm: AP -1.7, ML +/- 0.4, DV -2.7 to -3.0) and mPFC (from Bregma skull landmark, in mm: AP +2.5, ML +0.4, DV -1.5 to -1.8 from the surface of the brain).

For ChR2 (H134R) expression, 0.2-0.3 μl of AAV9.hSyn.hChR2(H134R)- mCherry.WPRE.SV40 (titre ~1012 GC/mL) was injected bilaterally into either LHb or mPFC and animals were left to recover for 3 to 4 weeks prior to ex-vivo electrophysiological recordings.

For the dual opsin experiments (Chrimson/Chronos), 0.2-0.3 ul of AAV-Syn-ChrimsonR-TdTomato (Serotype:2/9) was injected bilaterally into LHb, and 0.2-0.3 ul of AAV-Syn-Chronos-GFP (Serotype:1) was injected bilaterally into mPFC. In some control experiments (marked in text and figures), only AAV-Syn-ChrimsonR-TdTomato (Serotype:2/9) was injected bilaterally into LHb to rule out spectral overlap between opsins. Animals were left to recover for 3 to 4 weeks prior to ex-vivo electrophysiological recordings.

Slice preparation

LED Photostimulation experiments (Fig. 1). DRN containing brainstem slices were prepared from mice aged P40-P90. Acute slices were prepared similarly to previously described methods (Geddes et al., 2016). In brief, mice were anesthetized by isoflurane inhalation (Baxter Corporation, Canada) and sacrificed by decapitation. The brain was removed and brain slices were sectioned while immersed in ice-cold choline chloride-based cutting solution of the following composition (in mM): 119 choline-Cl, 2.5 KCl, 1 $CaCl_2$, 4.3 $MgSO_4$-7H_2O, 1 NaH_2PO_4, 1.3 sodium L-ascorbate, 26.2 $NaHCO_3$, and 11 glucose, and equilibrated with 95% O_2, 5% CO_2. Slices were recovered in a chamber containing standard Ringer's solution of the following composition (in mM): 119 NaCl, 2.5 $CaCl_2$, 1.3 $MgSO_4$-7H_2O, 1 NaH_2PO_4, 26.2 $NaHCO_3$, and 11 glucose, at a temperature of 37°C, continuously bubbled with a mixture of 95% O_2, 5% CO_2. Slices were left to recover for 1 hour in a recovery chamber and equilibrate to a temperature of approximately 21°C until recordings were performed.

Calcium conductance and membrane noise experiments (Fig. 3): DRN containing brainstem
slices were prepared from mice aged P21-P40. To optimize the health of serotonin neurons for
these sensitive experiments, acute slices were prepared using a modified NMDG two-bath
cutting procedure, described in Ting et al. (2018). Briefly, mice were anesthetized by isoflurane
inhalation (Baxter Corporation, Canada) and were transcardially perfused with ice-cold NMDG-
based cutting solution with the following composition (in mM): 92 NMDG, 2.5 KCl, 1.2
NaH_2PO_4H, 30 $NaHCO_4$, 20 HEPES, 10 $MgSO_4$, 25 Glucose, 0.5 $CaCl_2$, 5 sodium ascorbate,
2 thiourea, 3 sodium pyruvate, with pH adjusted to 7.3-7.4 with HCl and osmolality adjusted to
300-310 mOsm/kg with sucrose, and equilibrated with 95% O_2, 5% CO_2. Following perfusion,
the brain was removed and 300um brain slices were sectioned in the same NMDG cutting
solution. Brain slices were incubated for 0.5h in NMDG cutting solution at 37C, with NaCl
concentration gradually increasing to 92 mM over the incubation time (details in Ting et al.,
2018), and then transferred to a HEPES holding ACSF bath, at room temperature, with the
following composition (in mM): 92 NaCl, 2.5 KCl, 1.2 NaH_2PO_4H, 30 $NaHCO_3$, 2
$MgSO_4.7H_2O$, 2 $CaCl_2.2H_2O$, 20 HEPES, 25 Glucose, 5 Sodium ascorbate, 2 Thiourea, 3
Sodium pyruvate, equilibrated with 95% O_2, 5% CO_2. Slices were left to recover for 1 hour in a
recovery chamber and equilibrate to a temperature of approximately 21°C until recordings
were performed.
810
<u>Whole-Cell Electrophysiology</u>
DRN neurons were visualized using an upright microscope: (1) Examiner D1 equipped with
Dodt-contrast or differential-interference contrast (DIC) (Zeiss, Oberkochen, Germany; 40x/
0.75NA objective) or, (2) BX51W1 equipped with DIC optics (Olympus, Center Valley, PA; 40x/
0.80NA objective). 5-HT neurons were visually identified by TdTomato signal driven by the
SERT promoter under EPI-fluorescence illumination (xCite Series 120pc). Whole-cell
recordings were carried out using an Axon Multiclamp 700B amplifier, sampled at 10 kHz,
digitized with an Axon Digidata 1440A (or 1550) digitizer and filtered at 2 kHz. Whole-cell
recordings were performed using borosilicate glass patch electrodes (3-6 MΩ; World Precision
Instruments, Florida) pulled on a Narishige PC-10 pipette puller (Narishige, Japan).
821
822   All LED photostimulation experiments (Fig. 1) were performed at room temperature (20C).
823
824   All calcium conductance and membrane dynamics experiments (Fig. 3) were conducted at a
825   measured 30-32°C through the use of an in-line bath heater and stage heater. The purpose of
826   these alterations were to replicate, as closely as possible, the membrane dynamics and
827   channel kinetics that would be observed *in vivo*.
828
Slices received constant bath application of Ringer containing (in mM): 119 NaCl, 2.5 KCl, 1.3
$MgSO_4$, 2.5 $CaCl_2$, 1.0 NaH_2PO_4, 11 glucose, 26.2 $NaHCO_3$, and 0.1 L-Tryptophan saturated
with 95% O_2 and 5% CO_2 (pH = 7.3; 295-310 mOsm/L). An intracellular solution of the
following composition was utilized: 115 mM potassium gluconate, 20 mM KCl, 10 mM sodium
phosphocreatine, 10 mM HEPES, 4 mM ATP(Mg2+), and 0.5 mM GTP, pH 7.25 (adjusted with
KOH; osmolarity, 280–290 mOsmol/L). For all voltage clamp recordings, access resistance was
continuously monitored using a 125 ms, 2 mV hyperpolarizing pulse, at least 245 ms prior to
stimulation. Recordings were discarded when the access resistance changed by >30%. Liquid
junction potential was not compensated for. For current clamp experiments, during LED
photostimulation, direct current injection was used to maintain membrane voltage at
approximately ~-50mV to allow a direct comparison between neurons in each stimulation
condition (described in Fig. S2).

LED Photostimulation
LED photostimulation was delivered using a PlexBright LED module and controller (465 nm; Plexon, Texas). Light was delivered through a 200 μm patch fiber cable with a bare fiber tip with ~ 1 cm of glass exposed. For EPSC/EPSP stimulation, unless otherwise stated, optically evoked-PSCs were stimulated with single light pulses of 1-5 ms in length (individual trials 12-15 seconds apart) and an irradiance of 792 mW/mm2.

Drugs and Chemicals
Picrotoxin, CNQX and nickel (ii) chloride were purchased from Abcam (Cambridge, MA), Tocris Bioscience (Bristol, UK) and Sigma-Aldrich (St. Louis, MO), respectively. The concentration of drugs employed was as follows: Picrotoxin 100uM, CNQX 10uM, NiCl 250uM.

Data Analysis
All electrophysiological recordings were analyzed offline in the Python computing environment using the Numpy and Scipy computational packages. All data are presented as means +/- SEM. n refers to the number of cells recorded unless otherwise stated. P values of less than 0.05 were considered statistically significant and are indicated with an asterisk(*).

Applying linear classifiers to sub-threshold and spiking statistics
The approach to apply linear classifiers to the LHb and mPFC event data was as follows. We first describe methods to generate unbiased synthetic datasets of varying sample sizes from our physiological data, and second describe application of a linear classifier to this data.

Synthetic datasets. For a given number of samples and an event statistic (*e.g.* spike latency or EPSP amplitude), we generated 1000 synthetic datasets for both LHb and mPFC inputs. These synthetic datasets were generated by iteratively performing the following: 1) randomly sampling a neuron, then 2) randomly sampling a trial from that neuron and storing the event statistic from that trial. This was iteratively performed until the desired number of samples was reached. This sampling procedure ensured that each neuron was equally represented in the dataset. Next, the 1000 synthetic datasets were partitioned into either *training* or *test* datasets, with 5-fold cross-validation.

Linear classifier. We first performed standardization of the data using a min-max scaler, and then trained a support vector machine linear classifier on the *training* dataset using stochastic gradient descent learning (employing the implementation in scikit-learn). The classifier was then used to predict labels (LHb or mPFC) in the test dataset, with classification accuracy registered as the fraction of correct classifications across the entire test dataset. This linear classifier training and evaluation was then repeated for a total of 5 times (5-fold cross-validation) while rotating the held-out data in the test set. The classification accuracy was reported as the average accuracy across all 5 rotations of test and training data.

Generation and injection of synthetic EPSCs (Fig. 3)
Synthetic EPSCs were generated as a waveform with instantaneous rise-time, and a monoexponential decay with τ_{decay} that varied from 30ms to 500ms. These synthetic EPSCs were injected into the neuron in current clamp to elicit an evoked EPSP response. In Fig. 3C and Fig. S3, as various values of τ_{decay} were presented to the same neuron, the total charge

889   transfer to the neuron was held constant by scaling the amplitude of each EPSC such that the
890   integral of current over time was calculated to be identical.
891
<u>Membrane noise analysis and spike threshold calculation (Fig. 3)</u>
For the analysis of membrane potential dynamics, spikes were first identified using a fractional
first-derivative method (Trinh et al., 2019; Azouz and Gray, 2000). Briefly, the maximum
derivative of voltage over time, $max(\frac{dV}{dt})$, was calculated for each spike, and the spike
threshold was identified as the first point in the depolarization where the voltage derivative
reached $0.01 \cdot max(\frac{dV}{dt})$. A spike comprised all points from spike threshold to 0.4s after the
threshold; all such points were masked and not included in analysis of membrane voltage
statistics or spectral characteristics. To calculate the spectral characteristics of membrane
voltage, a 1-dimensional discrete fast Fourier transform of the voltage time series was
computed, and the resulting spectrum was smoothed using a Savitzky-Golay filter with
polynomial order 3.
903
<u>Stochastic membrane noise simulations (Fig. 4)</u>
To investigate how distinct spike timing patterns could arise from stochastic fluctuations in
membrane voltage, we simulated membrane voltage as a dynamical process and solved the
level-crossing problem to infer how spike timing distributions depended on membrane
dynamics. We considered a simplified scenario where membrane voltage ($v_0 = -50mV$) was
initialized close to spike threshold ($v_{thresh} = -45mV$). Membrane voltage evolved
stochastically (described below) until spike threshold was reached. By repeating this over
20,000 simulations, we generated a distribution of level-crossing times (representing spike
timing distributions), allowing us to assess how membrane potential dynamics were related to
distinct regimes of spike timing.
914
915   We simulated the dynamics of membrane voltage as an Ornstein-Uhlenbeck noise with the
916   following equation:
917
918     $\frac{dv(t)}{dt} = -\frac{1}{\tau}(v(t) - v_0) + \sqrt{D} \cdot \eta(t)$ (1)
919
where τ is a constant denoting the correlation timescale of the noise (indicated in text, ranging
from 0.01 to 0.5), $v_0 = -50 \; mV$ represents the resting membrane potential, and $\eta(t)$ is a
white noise term. $\sqrt{D}$ is a noise diffusion coefficient (scaling factor) that was varied across
simulations with τ such that the quantity $\sigma^2 = D\tau/2$ was constant ($\sigma^2 = 100$); this produced
fixed membrane potential distributions that were independent of τ as it was varied (Fig. 4B).
The simulations proceeded until v reached the simulated spike threshold $v_{thresh} = -45 \; mV$,
at which point the time from simulation onset to spike threshold was recorded as a simulated
spike latency.
928
929   Numerically, the Ornstein-Uhlenbeck noise process was computed using the following
930   constant definitions and update rule, following the well-known algorithm derived by Gillespie
931   (1996) for exact numerical simulation of the Ornstein-Uhlenbeck Process:
932

$$A = \sqrt{\frac{D\tau}{2}\left(1 - e^{\frac{-2\Delta t}{\tau}}\right)} \qquad (2)$$

$$v(t + \Delta t) = v_0 + (v(t) - v_0) \cdot e^{-\Delta t/\tau} + A \cdot \mathcal{N}(0,1) \qquad (3)$$

With a timestep of $\Delta t = 0.001\ s$.

For simulations in which the spectral features of the noise were altered, we conducted
simulations as described above, then performed a fast Fourier transform of the simulated
voltage, altered the signal in the frequency domain, then transformed back into the time
domain.

Supplemental Figure 1: Dual-color opsin dissection of long-range inputs to 5-HT neurons. **A**: For 5-HT neurons that respond to both blue and red light, red paired-pulse stimuli produce depression (20Hz, top) while a red prepulse does not significantly affect the amplitude of a blue light-evoked current (bottom). **B**: Quantification of paired pulse ratio for red paired-pulse stimuli (red) and blue paired-pulse stimuli with red prepulse (blue). **C**: Single-opsin control with Chrimson injected in LHb. Red paired-pulse stimuli produce depression (20Hz, top) and a red prepulse significantly affect the amplitude of a blue light-evoked current (bottom). **D**: Quantification of paired pulse ratio for red paired-pulse stimuli (red) and blue paired-pulse stimuli with red prepulse (blue).

Supplemental Figure 2: Quantifying features of EPSPs and spikes elicited by LHb and PFC inputs to 5-HT neurons. **A-B**: Example traces in voltage clamp of light-evoked EPSCs in 5-HT neurons. **C**: Cumulative distribution of EPSP amplitudes of all LHb-evoked events (blue) and PFC-evoked events (red) (K-S test, n=631 events from 9 neurons, LHb; n=619 events from 9 neurons, PFC). **D**: Cumulative distribution of EPSP decay of all LHb-evoked events (blue) and PFC-evoked events (red) (K-S test, n=631 events from 9 neurons, LHb; n=619 events from 9 neurons, PFC). **E**: Mean baseline membrane potential before evoked spikes (n=9 neurons for LHb and PFC groups, bars indicate S.E.M; p>0.05, t-test). **F**: Mean probability for photostimulation to trigger a spike (n=9 neurons for LHb and PFC groups; bars indicate S.E.M; p>0.05, t-test). **G-H:** Trial-by-trial voltage trajectories to spike threshold in two example neurons receiving suprathreshold long-range LHb (**G**) or PFC (**H**) input. **I**: Population average of voltage trajectories to spike threshold (n=194 spikes from 9 neurons, LHb; n=203 spikes from 9 neurons, PFC; shaded lines denote S.E.M). **J:** Reliability of voltage trajectory to spike threshold across trials, for LHb and PFC inputs (Reliability is computed as pairwise pearson r coefficient of voltage trajectory across all trials within a neuron; n=9 neurons for PFC and LHb, p=0.014, t-test). **K-N**: Probability distribution of spike times triggered by PFC (red) and LHb (blue), shown with regular scaling (**K-M**) and log-log scaling (**L-N**). Black lines indicate exponential fit.

Supplemental Figure 3: Low-threshold calcium channels modulate the kinetics of EPSPs in 5-HT neurons. **A**: Example traces in current-clamp showing that injection of fast (tau=50ms) synthetic EPSCs produce EPSP responses that are modulated by bath application of NiCl. **B**: Example traces in current-clamp showing that injection of fast (tau=100ms) synthetic EPSCs produce EPSP responses that are modulated by bath application of NiCl. **C**: Example traces in current-clamp showing that injection of fast (tau=500ms) synthetic EPSCs produce EPSP responses that are modulated by bath application of NiCl.

Supplemental Figure 4: Subthreshold noise properties are not affected by calcium channels in 5-HT neurons. **A**: Example traces of a single neuron recorded in current-clamp during injection of constant subthreshold depolarizing current, before (top) and after (bottom) application of NiCl. 'Subthreshold' is defined as traces recorded during a current step just below spike threshold. **B**: Probability distribution of membrane potentials for an example neuron before (top) and after (bottom) application of NiCl. **C**: Membrane potential distribution skewness is not significantly affected by bath application of NiCl (n=7 neurons, paired T-test). **L**: Membrane potential distribution kurtosis not is significantly affected by bath application of NiCl (n=8 neurons, paired T-test).

Supplemental Figure 5: Low-threshold calcium channels contribute to spike timing properties of long-range PFC input to 5-HT neurons. Panels **A-C** depict results from a single representative 5-HT neuron. **A**: Representative traces from 5-HT neuron in current clamp depicting high-latency spiking in response to optogenetic PFC axon stimulation (left), and lower latency spiking after bath application of NiCl (right). **B**: Representative averaged trace in voltage clamp depicting inward current in response to a -45mV depolarizing step (black), which is abolished by bath application of NiCl (green). **C**: Cumulative distribution of spike latency before (black) and after (green) NiCl application from a single 5-HT neuron. **D**: Timing of spikes triggered by photostimulation of long-range PFC axons before (black) and after (green) bath application of NiCl. Each dot is a spike and each row depicts one neuron (n=4 neurons). **E**: Cumulative distribution of spike times triggered by photostimulation of PFC axons before (black) and after (green) bath application of NiCl (Kolmogorov-Smirnov test, p=0.045, n=96 spikes (base) and 62 spikes (Ni) from 4 neurons).