Rho-ROCK liberates sequestered claudin for rapid *de novo* tight junction formation

Yuma Cho[1], Akari Taniguchi[1], Akiharu Kubo[2] and Junichi Ikenouchi[1] *

1. Department of Biochemistry, Kyushu University Graduate School of Medical Sciences, Fukuoka, Japan

2. Division of Dermatology, Department of Internal Related, Graduate School of Medicine, Kobe University, Kobe, Japan

*Address correspondence to: Dr. Junichi Ikenouchi

Tel.: +81-92-642-6096; Fax: +81-92-642-6096; E-mail: ikenouchi.junichi.033@m.kyushu-u.ac.jp

Key words: Tight junction, Claudin, EpCAM, TROP2, epithelial barrier

Running title: Roles of the EpCAM-Claudin complex in the homeostatic maintenance of tight junctions

1   **Summary**
2
The epithelial cell sheet maintains its integrity as a barrier while undergoing turnover of constituent cells.
To sustain the barrier continuously, it's essential to preserve the 'old' tight junctions (TJs) between cells
being excluded from the sheet and their neighbors while simultaneously forming *de novo* TJs between
newly adjacent cells. However, the molecular mechanisms involved in the formation of *de novo* TJs
remain largely unknown. This study investigates two scenarios: the formation of *de novo* TJs during the
removal of apoptotic cells from monolayer epithelial sheets and during the differentiation of the granular
layer in stratified epidermis. We revealed that rapid claudin assembly is achieved by actively regulating
the dissociation of the EpCAM/TROP2-claudin complex in both situations. Furthermore, we found that the
Rho-ROCK pathway initiates the activation of matriptase, which cleaves EpCAM/TROP2, resulting in the
supply of polymerizable claudin from the stockpiled EpCAM/TROP2-claudin complex at the plasma
membrane to induce rapid *de novo* TJ formation.

14
15
16
17   **Introduction**
18
The surfaces of our body and its organs are covered by epithelial cell sheets, which serve to prevent the
entry of antigens and pathogens from the outside and the leakage of substances such as water and
glucose from the body [1,2]. This barrier function is achieved through the close attachment of the plasma
membranes of neighboring epithelial cells by intercellular adhesion structures called tight junctions (TJs).
The major component of TJs is claudin, a four trans membrane protein of around 25 kDa. Claudin binds in
cis on the plasma membrane and in *trans* between neighboring cells, forming a membrane structure
called TJ strands [3].
In both simple epithelia, such as the gastrointestinal tract, and stratified epithelia, such as skin,
the barrier is continuously maintained by various mechanisms that regulate the homeostasis of the
epithelial cell sheet, including the wound healing response [4]. For example, unnecessary inflammatory
responses are suppressed by mechanisms that rapidly eliminate apoptotic cells generated by constant
cell turnover or exposure to external stresses in the epithelial cell sheet [5]. During the process of apoptotic
cell extrusion, the apoptotic cells are physically pushed towards the apical side by an increase in
actomyosin contractility of cells adjacent to the apoptotic cells [6,7]. The barrier function of the epithelial
sheet is maintained during this process of apical extrusion [7]. In parallel with rapid apoptotic cell
elimination, it is necessary to maintain integrity of TJs between apoptotic cells and neighboring cells,
while rapidly forming *de novo* TJs between newly adjacent cells that result from apoptotic cell elimination.
However, the mechanism for the formation of such rapid *de novo* TJs is largely unknown.
In the epidermis, TJs form only between keratinocytes in the granular layer and act as a barrier
separating the inside and outside of the body. Claudin-1 knockout mice die within a day of birth due to

abnormal stratum corneum differentiation and disrupted granular layer barrier function, resulting in severe dehydration [8,9]. In the epidermis, stem cells in the basal layer divide to give rise to new keratinocytes, which differentiate towards the body surface into keratinocytes constituting the spinous layer, granular layer, and stratum corneum, forming a multilayered epithelium [10]. TJs are only formed in the second layer of the granular layer (SG2), even though claudin is also expressed in cells of the basal and spinous layers [11–13]. In the turnover of keratinocytes in the SG2 layer, the epidermal barrier is continuously maintained by the simultaneous formation of 'new' TJs between newly differentiating SG2 keratinocytes and pre-existing SG2 keratinocytes, before the 'old' TJs between SG2 keratinocytes that are differentiating into keratinocytes of the stratum corneum and neighboring SG2 keratinocytes are lost [13]. It has recently been reported that an E-cadherin-dependent increase of intercellular tension at adherens junctions (AJs) and inhibition of EGFR activity are required for induction of TJ formation in the SG2 layer [14]. However, how these E-cadherin-mediated changes regulate TJ formation in the SG2 layer and why TJs are never formed in keratinocytes other than the SG2 layer that express claudin remain open questions.

The formation of *de novo* tight junctions, which is common in these two phenomena, is rapid, suggesting that the formation of functional TJs is not regulated through *de novo* transcription or translation of claudin, but rather by controlling the polymerization state of claudin. In heterogeneous cells, such as L cells and COS-7 cells, forced expression of claudin alone is known to be sufficient to induce TJ formation [15,16]. On the other hand, in the lateral membranes of simple epithelial cells and keratinocytes, excess claudin is present in an unpolymerized. The molecular mechanisms that strictly regulate restrict the polymerization of claudin at the presumptive TJ region are largely unknown.

Recently, a single transmembrane protein, EpCAM, was identified as an interacting protein of claudin [17]. EpCAM binds to claudin in the lateral membrane of epithelial cells through the interactions between transmembrane regions [18,19]. By binding to EpCAM, claudin is stabilized at the plasma membrane, preventing it from being endocytosed and degraded [20]. Knockout of EpCAM in cultured cells does not affect claudin accumulation at TJs but selectively reduced claudin in the lateral membrane [20,21].

Another single transmembrane protein, TROP2, which shows 67% similarity in amino acid sequence to EpCAM, also binds to claudin [22]. In the small intestine, only EpCAM is expressed, while in the epidermis, both EpCAM and TROP2 are expressed [23,24]. In humans, loss-of-function mutations in the EpCAM gene cause a disruption in the barrier function of gastrointestinal epithelial cells, resulting in congenital tufted enteropathy (CTE), characterized by chronic diarrhea and growth delay [25,26]. EpCAM knockout mice also show similar phenotypes and die within two weeks after birth [27,28]. Similarly, TROP2 is the causative gene for human gelatinous drop-like corneal dystrophy (GDLD), which causes photophobia and vision loss due to amyloid deposition under the corneal epithelium [29,30]. This is because the barrier function of the corneal epithelium is impaired due to TROP2 dysfunction, allowing lactoferrin and

apolipoproteins in the tear fluid to penetrate and form amyloid deposits. It was also recently shown that the epidermal barrier function is impaired in TROP2 KO mice [24].

EpCAM and TROP2 are cleaved in their extracellular regions by matriptase, a transmembrane serine protease [31,32]. Cleavage of EpCAM disrupts the EpCAM-claudin complex, releasing making available unpolymerized claudin for TJ formation [21,32]. Higashi et al. proposed the physiological significance of this molecular mechanism, suggesting that when TJs are locally disrupted, serine proteases on the apical side shift their localization to the basolateral side and cleave EpCAM at the basolateral membrane, releasing unpolymerized claudin to repair TJs. In other words, local disruption of TJs allows passive translocation of proteases due to the partial loss of the fence function of TJs, leading to the breakdown of the EpCAM-claudin complex. In this study, we examined in detail the regulation of the EpCAM-claudin complex in the mechanisms to induce *de novo* TJs during the elimination of apoptotic cells and turnover of the SG2 layer of the epidermis.

Results

Rapid *de novo* TJ formation occurs during apoptotic cell elimination in a Rho-ROCK pathway-dependent manner

The pioneering work by Rosenblatt and colleagues established that the barrier function of the epithelial cell sheet is preserved even in the face of a mass cell death event triggered by acute UV irradiation [7]. This observation suggests that the epithelium is capable of removing apoptotic cells while maintaining the continuity of TJs uninterrupted. However, the complex logistics of simultaneously breaking down the TJ between apoptotic and neighbor cells and assembling them anew between formally non-interacting cells is not sufficiently known. We first examined claudin dynamics among relevant cells during apoptotic cell elimination in order to understand how these incongruous processes are carried out. Apoptosis was induced in a confluent monolayer of a representative epithelial cell line EpH4 stably expressing GFP-claudin-3 by pulsed laser irradiation of the nucleus in a targeted cell. Apical extrusion of the apoptotic cell and bicellular accumulation of claudin between newly-formed contacts were typically observed over the course of an hour **(Figure 1A, 1B and Movie 1)**. Throughout, GFP-claudin-3 remained enriched to some degree between the apoptotic and neighbor cells **(Figure 1B, purple arrowheads)**. Additionally, linear accumulation of GFP-claudin-3 could be observed more basally by about 40 min post-irradiation and steadily grew more pronounced over time **(Figure 1B, green arrowheads)**. Immunofluorescence staining showed that other TJ proteins, such as ZO-1 and occludin, colocalize with the basally-enriched claudin, indicating the *de novo* formation of functional TJs between the newly neighboring cells **(Figure 1C, green arrowheads)** before the complete loss of the apoptotic cell-neighbor cell TJs **(Figure 1C, purple arrowheads)**. Together, these data demonstrate that old and new TJs exist concurrently among the cells involved in apoptotic cell removal to ensure that there is no period of TJ absence that could compromise the epithelial barrier.

Previous studies have established that RhoA is activated in the neighbor cells at the interface

with the apoptotic cell to enhance the contractility of the actomyosin purse string that extrudes the

apoptotic cell [33]. RhoA is also required for the maturation of epithelial cell adhesions through its effect on

the underlying circumferential actomyosin and recent reports implicate it in the repair of TJ microinjuries [34].

We therefore hypothesized that RhoA could play a role in the rapid *de novo* TJ formation such as we see

during apoptotic cell removal. Observation of active RhoA in neighbor cells using an effector domain

biosensor (GFP-AnillinC) showed sustained activation at the apoptotic cell-neighbor cell interface as

previously reported **(Figure 1D and Movie 2, white arrowheads)**. Importantly, RhoA activity persisted at

newly formed cell contacts after full expulsion of the apoptotic cell **(Figure 1D, yellow arrowheads)**.

Furthermore, immunostaining revealed phosphorylation of the myosin regulatory light chain (ppMLC), a

substrate of the RhoA effector ROCK, at such contacts that are clearly located beneath the extruded cell

(Figure 1E). These data collectively point to the presence of an active pool of RhoA at the new TJs.

Apoptotic cells rapidly contract, causing the neighbor cells to extend lamellipodia, which acts to

seal the basal surface. Since contact between protrusions from separated cells is necessary to initiate

adhesion formation, we wondered if these lamellipodia alone, previously shown to be independent of both

the Rho-ROCK pathway and myosin contractility, could instigate claudin enrichment at newly formed

contacts [35]. By imaging the plasma membrane (mScarlet-PLCδ PH) together with GFP-claudin-3 under

ROCK inhibition, we found that claudin no longer accumulated at bicellular contacts once the old

apoptotic cell-neighbor cell TJs broke down, despite the completion of basal closure by neighbor cell

lamellipodia **(Figure 1F and Movie 3)**. This observation suggests that *de novo* TJ formation is impaired

by ROCK inhibition. Accordingly, inducing mass apoptosis in the epithelial monolayer by treatment with

the anticancer drug doxorubicin caused a significant increase in the leakage of 70 kDa FITC-Dextran

when ROCK was simultaneously inhibited, compared to the control condition **(Figure 1G)**. Signaling

through Rho-ROCK is clearly required for apical extrusion, since the apoptotic cell typically remained

loosely embedded in the cell sheet. However, these results collectively hint that its importance in epithelial

barrier homeostasis associated with apoptotic cell removal is not limited to this process: it is also critically

important for claudin enrichment and/or engagement at new cell-cell contacts, the failure of which

precludes the *de novo* formation of TJs **(Figure 1H)**.

De novo TJ formation in epidermis requires activation of the Rho-ROCK pathway

Given the technical difficulty in independently assessing the contribution of the Rho-ROCK

pathway in the two intertwined events arising in cells neighboring an apoptotic cell—increased

contractility to effect apical extrusion and *de novo* TJ formation to maintain the epithelial barrier—we

chose to investigate its involvement in the context of *de novo* TJ formation in keratinocyte differentiation.

The human keratinocyte cell line HaCaT differentiates into granular-like cells capable of forming TJs

when cultured for 24 hours in a medium supplemented with high concentration of calcium ion (9.8 mM)

and a JNK inhibitor (Ca + JNK inh medium) **(Figure 2A)** [36,37]. In undifferentiated HaCaT cells, both

claudin-1 and the adherens junction (AJ) receptor E-cadherin were broadly distributed throughout the

plasma membrane and their respective scaffolding proteins ZO-1 and α-catenin appeared cytoplasmic,

indicating the absence of functional epithelial adhesions **(Figure 2B, control)**. Following differentiation,

claudin-1 and ZO-1 were sharply enriched at apical intercellular adhesion sites and the AJ components

co-localized at the basolateral membrane **(Figure 2B, Ca + JNKinh)**.

Accordingly, in differentiated HaCaT cells, the actin cytoskeleton was re-organized to form the

circumferential actin ring associated with apical junctions (**Figure 2B,** F-actin). The morphological

changes observed here align well with those seen in differentiated keratinocytes of the SG2 layer *in vivo*.

Of note, western blotting revealed that there was no significant difference in claudin-1 expression level

before and after differentiation **(Figure 2C)**. Moreover, in the mouse ear epidermis, claudin-1 was present

across the entire plasma membrane in the spinous layers before differentiation into the SG2 layer **(Figure**

2D). Altogether, these observations suggest the existence of a regulatory mechanism that specifically

instigates *de novo* TJ formation in keratinocytes of the SG2 layer, independent of transcriptional and/or

translational control.

We then hypothesized the involvement of the Rho-ROCK pathway in *de novo* TJ formation

associated with keratinocyte differentiation, similar to its role in *de novo* TJ formation during apoptotic cell

elimination in monolayer epithelial cells. First, we investigated whether RhoA is activated during

keratinocyte differentiation. We detected a significant increase in RhoA activation upon differentiation

(Figures 3A and 3B). In differentiated HaCaT cells, phosphorylated MLCs were prominently observed at

cell-cell contacts **(Figures 3C and 3D)**. Similarly, in the mouse epidermis, ppMLC was notably enhanced

only at apical junctions of keratinocytes in the SG2 layer **(Figure 3E)**. These findings indicate an

augmented Rho-ROCK pathway in keratinocytes of the granular layer during differentiation, consistent

with a previous report that junctional tension is necessary for TJ formation in the granular layer [14].

We next examined whether activation of the Rho-ROCK pathway during differentiation is

necessary for the formation of *de novo* TJs upon keratinocyte differentiation. When ROCK activity was

inhibited by treatment with Y27632, claudin remained uniformly distributed across the entire plasma

membrane, indicating impairment of TJ formation **(Figures 3F and 3G)**. In contrast, E-cadherin signals at

apical junctions was not affected by ROCK inhibition in differentiated HaCaT cells **(Figure 3F)**. As we saw

in monolayer layer epithelia, these results suggest that Rho-ROCK signaling specifically regulates *de*

novo TJ formation in differentiating keratinocytes of the stratified epithelia by enabling claudin enrichment

at intercellular adhesions, i.e. shifting claudin from an unpolymerized state in the basolateral membrane

to a polymerized state in TJ strands **(Figure 3H)**.

Activation of ROCK alone can induce TJ formation in undifferentiated keratinocytes

Then how does the Rho-ROCK pathway regulate the polymerization state of claudin? Since

HaCaT cells already abundantly express claudins prior to differentiation, we wondered if activation of

RhoA or ROCK alone could induce *de novo* TJ formation in the absence of differentiation factors.

Overexpression of a constitutively active mutant of RhoA (Q61L, RhoA CA) led to the accumulation of

claudin-3 at cell-cell contacts but only when both cells expressed RhoA CA **(Figure 4A)**. By contrast, claudin-3 enrichment was not observed in cells expressing the other main Rho family GTPases, Rac1 and Cdc42 **(Figure S2A)**. Importantly, additional TJ components, such as ZO-1 and the transmembrane protein occludin, were recruited to claudin-enriched cell-cell contacts and actin reorganization was also observed, indicating that these structures were highly TJ-like **(Figures 4A and 4B)**. Moreover, activation of endogenous RhoA by treatment with the bacterial toxin cytotoxic necrotizing factor (Rho activator; **Figures 4C and 4G**), overexpression of a constitutively active mutant of ROCK (ROCKdelC; **Figure S2B**) and treatment with the ROCK activator Narciclasine also induced strong accumulation of claudin-3 in undifferentiated HaCaT cells **(Figures 4E and 4G)** to further support the proposition that signaling through RhoA-ROCK enables claudin polymerization. In some instances, the span of enriched claudin was significantly expanded laterally, indicating that the normally unpolymerized pool of claudin was being polymerized in situ, in the absence of positional cues **(Figures 4D and 4F)**.

We previously showed that polymerized claudin is preferentially partitioned into detergent resistant membranes (DRMs) enriched in cholesterol and very long-chain fatty acid-containing sphingomyelin [38]. Since both Rho activator and Narciclasine treatments can efficiently enrich claudin at cell-cell contacts, we next evaluated the polymerization state of claudin in treated cells biochemically. Rho activator and Narciclasine treatments markedly increased the amount of claudin-1 partitioned into the DRM fraction compared to the control condition. In contrast, in undifferentiated HaCaT cells, most of claudin-1 was solubilized by detergents, reflecting that claudin is present in an unpolymerized state in undifferentiated HaCaT cells **(Figures 4H and I)**. These findings altogether indicate that activation of the Rho-ROCK pathway induces the transition of claudin from an unpolymerized state to a polymerized state.

Matriptase activation by Rho-ROCK produces a pool of polymerizable claudin via TROP2 cleavage

Having established that signaling through Rho-ROCK triggers claudin polymerization, we next asked what the mechanism is. The single transmembrane proteins EpCAM and TROP2 bind to claudin and are abundantly expressed in epithelial cells and keratinocytes. They are thought to protect unpolymerized claudin from degradation to maintain a stable reserve apart from the TJ strand [20]. This prompted us to consider whether they could play a greater role, not only in protecting unpolymerized claudin from degradation but also in inhibiting unregulated claudin polymerization by interfering with the interaction between claudins. Matriptase, a transmembrane serine protease, cleaves the extracellular region of EpCAM and TROP2 to sever their interaction with claudin [31,32]. It was recently reported that, of the two, TROP2 plays a greater role in the epidermis [32]. Therefore, we reasoned that Rho-ROCK signaling releases TROP2-bound claudin by activating matriptase.

Matriptase is normally present at the plasma membrane in an inactive state but it is activated by autocleavage in response to external signals. After cleavage of the substrate, the Kunitz-type protease inhibitor family proteins Hai1 and Hai2 immediately bind to matriptase to block its activity [39–41]. The extracellular region of the matriptase-Hai complex is then cleaved and released from the cell **(Figure 5A, illustration)** [42]. To assess the activation state of Matriptase, we employed two mouse monoclonal

antibodies: M24, which detects total Matriptase protein levels, and M69, which specifically recognizes the

activated form of matriptase. These antibodies have been widely used in previous studies on

matriptase[43–47]. One such study showed that matriptase is activated by simply placing cells in an acidic

medium at pH 6.0 for 20 minutes [47]. Addition of acid medium to undifferentiated HaCaT cells decreased

the total amount of matriptase and increased the amount of cleaved TROP2 compared to the control,

indicating that matriptase is activated and cleaves TROP2 **(Figure 5A)**. This was accompanied by claudin

accumulation at cell-cell contacts and an increase in insoluble, i.e. polymerized, claudin **(Figures 5C-5F)**.

In an intriguing contrast to the results of Rho-ROCK activation by pharmacological means **(Figures 4C**

and 4E) where we saw development of junction-associated F-actin and AJ maturation simultaneous with

claudin polymerization, neither was observed here **(Figure 5C)**. This observation suggests that bypassing

Rho-ROCK and activating matriptase directly is sufficient to induce claudin strand assembly. However,

treatment with Rho-ROCK activators increased matriptase activity and enhanced TROP2 cleavage,

indicating that matriptase is a bona fide effector of Rho-ROCK signaling **(Figures 5A and 5B)**.

Furthermore, inhibition of matriptase activity with Camostat treatment suppressed claudin accumulation at

cell-cell contacts induced by Rho-ROCK activation, which further support the notion that activation of

matriptase occurs downstream of Rho-ROCK pathway **(Figures 5G and 5H)**. Collectively, these results

demonstrate that activation of matriptase by Rho-ROCK signaling liberates claudin from sequestration by

TROP2 and acts as the rate limiting process in *de novo* TJ formation; indeed, TROP2 was uniformly

present throughout the plasma membrane in undifferentiated cells but it was specifically excluded from

the region where claudin was polymerized between cells expressing Rho CA cells **(Figure 5I)**.

It was previously reported that the amount of claudin-1 protein is significantly reduced

TROP2-depleted keratinocytes, which we confirmed **(Figure 6A and 6B)** [32]. More interestingly, however,

quantification of claudin-1 surface expression revealed that the amount was almost halved in TROP2 KO

cells, even accounting for decreased overall expression **(Figures 6C and 6D)**. Moreover, claudin-1 was

most prominently found at the Golgi apparatus of TROP2 KO cells, where it was distributed throughout

the plasma membrane in undifferentiated wild-type cells, only appearing intermittently there as strongly

enriched dots **(Figure 6E)**. These punctate claudin-1 accumulations contained ZO-1 and occludin,

indicating that claudin-1 is incorporated in a quasi-TJ state **(Figure S3)**. In other words, claudin-1 cannot

stably exist in an unpolymerized state at the plasma membrane in the absence of TROP2. As to be

expected by the lack of available surface claudin, neither Rho-ROCK activation nor matriptase activation

by acid medium treatment could induce claudin polymerization in TROP2 KO cells **(Figures 6F-6H)**.

Taken together, these findings indicate that the regulation of claudin by TROP2 is not limited to stabilizing

unpolymerized claudin at the plasma membrane: it begins at the Golgi apparatus where—acting as a

chaperone—it mediates proper sorting and trafficking **(Figure 6I)**.

The results so far strongly suggest that the Rho-ROCK pathway enables claudin polymerization,

the initial step of *de novo* TJ formation, by activating matriptase, which frees the claudin pool sequestered

by TROP2 at the lateral membrane; this is independent of its long-established effect on junctional

maturation through actomyosin. We next performed a series of experiments to verify this hypothesis. First,

we found that matriptase was rapidly activated in HaCaT cells introduced into the Ca+ JNKinh medium to

induce differentiation; degradation of TROP2 was observed in parallel **(Figures 7A and 7B)**.

Immunostaining with the M24 antibody that detects total protein revealed that matriptase is

constitutively distributed along the lateral membrane, below TJs, regardless of HaCaT cell

differentiation state **(Figure 7C)**. By contrast, staining with the M69 antibody, which specifically

recognizes the activated form, revealed no signal in undifferentiated cells but following differentiation,

activated matriptase could be widely detected at the lateral membrane **(Figures 7D and 7E)**. These

observations indicate that while the subcellular localization of matriptase remains unchanged during

differentiation, it undergoes activation at the lateral membrane in differentiated granular-layer-like

cells.

We then examined HaCaT differentiation in the presence of the matriptase inhibitor Camostat.

Consistent with the result of our experiment with Rho CA overexpression, claudin failed to accumulate at

cell-cell junctions in Camostat-treated cells **(Figures 7F and 7H)**. A previous study demonstrated that

increase in intercellular tension at AJs is indispensable for the formation of TJs in the SG2 layer[14]. To

assess AJ tension, we stained with the α-18 monoclonal antibody, sensitive to the tension-dependent

conformational change of α-catenin, a core AJ component. In control cells, differentiation into

granular-layer-like cells led to a marked increase in α-18 signal at cell–cell adhesions. Importantly,

when HaCaT cells were treated with Camostat to inhibit matriptase before induction, we observed an

equivalent increase in α-18 signal at AJs **(Figure 7F)**. However, we did not detect claudin enrichment

at cell-cell contacts under these conditions **(Figures 7F and 7H)**. These results suggest that

matriptase inhibition specifically disrupts TJ formation during granular-layer differentiation without

impeding AJ maturation.

Similarly, in HaCaT cells overexpressing Hai1, an endogenous inhibitor of matriptase, TJ

formation was severely suppressed **(Figures 7G and 7H)**. Lastly, we sought to extend our findings to the

granular layer differentiation of the epidermis. As in undifferentiated HaCaT cells, TROP2 was distributed

throughout the plasma membrane of the spinous layer but was progressively lost within the granular

layer—particularly at the SG2 layer—as TJs appeared and matured **(Figure 7I)**. These results suggest

that ROCK-dependent activation of matriptase and cleavage of TROP2 by matriptase are crucial steps in

the formation of *de novo* TJs associated with granule cell differentiation in the epidermis **(Figure 7J)**.

Rho-ROCK liberates EpCAM-sequestered claudin through matriptase to enable rapid *de novo* TJ

formation during apoptotic cell elimination

We next investigated whether the same molecular mechanism of *de novo* TJ formation in

stratified epithelium plays a role in the homeostasis of the epithelial barrier in monolayer epithelial cells.

It has been established that EpCAM, rather than TROP2, regulates claudin in simple epithelium

[20,21,28,31]. Therefore, we established EpCAM KO EpH4 cells. In EpCAM KO cells, the amount of claudin-1

was significantly reduced, consistent with previous studies and TROP2 depletion in HaCaT cells **(Figure**

8A and 8B) [20]. Immunofluorescence with the standard Triton X-100 membrane permeabilization revealed

a sharp, linear localization of claudin-4 at apical junctions in EpCAM KO cells that was indistinguishable

from wild-type cells, giving the appearance that TJs are formed normally under steady state condition

(Figure 8C). However, when the detergent was changed to digitonin, significant claudin-7 staining was

observed in the lateral membranes in wild-type cells **(Figure 8D)**. By contrast, the lateral pool of claudin-7

was severely reduced in EpCAM KO cells; as in HaCaT cells depleted of TROP2, claudin was seen to

accumulate at the Golgi apparatus instead **(Figures 8D and 8E)**. These results indicate that claudin

transport is similarly regulated in simple epithelium and stratified epithelium by EpCAM and TROP2,

respectively.

We then wondered whether cleavage of EpCAM by forcibly activating matriptase through

treatment with acid medium would also induce *de novo* TJ formation here. When wild-type EpH4 cells

cultured on permeable supports were treated with the acid medium on the basolateral side, claudin

enrichment could be observed throughout the lateral membrane in contrast to EpCAM KO cells that

showed no such ectopic TJ expansion **(Figure 8F)**. These findings suggest that, similar to stratified

epithelium, cleavage of EpCAM by matriptase supplies polymerizable claudins for *de novo* TJ formation in

monolayer epithelium.

Finally, we investigated whether ROCK-dependent activation of matriptase and cleavage of

EpCAM by matriptase is important for rapid *de novo* TJ formation during apoptotic cell elimination. A

previous study showed that EpCAM cleavage mediates TJ repair at small leakages induced by acute

tension elevation [21]. Apical extrusion of the apoptotic cell induced by pulsed laser irradiation proceeded

without fail in EpCAM KO cells as in wild-type cells but the formation of *de novo* TJs between newly

adjacent cells was never observed **(Figure 8G and Movie4)**. Furthermore, *de novo* TJ formation after

apoptosis induction was likewise suppressed in EpH4 cells treated with Camostat or overexpressing Hai1

(Figures 8H, 8I and Movies 5, 6). These findings support the notion that *de novo* TJ formation occurring

in parallel with apical extrusion of apoptotic cells requires the degradation of EpCAM by matriptase.

The rapid restoration of TJ continuity upon the elimination of apoptotic cells is crucial for

epithelial barrier homeostasis. Even when apoptosis was induced by treatment with 4 µM doxorubicin,

leakage of 70 kDa FITC-Dextran was not detected in control EpH4 cells, indicating the preservation of

epithelial barrier integrity. In clear contrast, significant leakage of 70 kDa FITC-Dextran was detected in

EpCAM KO cells and epithelial cells in which matriptase function was inhibited **(Figure 8J)**. Altogether,

we conclude that Rho activation triggers matriptase-mediated EpCAM cleavage, which quickly introduces

a supply of polymerizable claudin to enable the preservation of barrier integrity in damaged epithelial cell

sheets.

Discussion

In this paper, we demonstrate that the rapid formation of *de novo* TJs observed during the

elimination of apoptotic cells in simple epithelium and the differentiation of granular cells in stratified

epithelium begins with the activation of the Rho-ROCK pathway and follows a common series of

reactions: (1) activation of matriptase, (2) destruction of the EpCAM/TROP2-claudin complex via

cleavage of EpCAM/TROP2 by activated matriptase, and (3) liberated supply of polymerizable claudin

(Figure 9).

4   The contribution of the Rho-ROCK pathway to the maintenance of epithelial barrier homeostasis

has been explicated in various experimental contexts. It is known to increase the contractile force of cells

adjacent to apoptotic cells to enable efficient apoptotic cell elimination [7,33,48], and recent research using

Xenopus embryos showed that local activation of Rho (referred to as Rho Flare) upregulates the

contractility of TJ-associated actomyosin at the site of TJ rupture, minimizing TJ disruption and thus

barrier breach [34]. Here, we utilized monolayer and stratified epithelial cells, which enabled us to

distinguish a novel role of signaling through Rho-ROCK, distinct from its effect on the cytoskeleton: to

release the stockpiled claudin—complexed with EpCAM/TROP2—from the lateral membrane to promote

rapid *de novo* TJ assembly. Several previous studies support our proposal. For one, overexpression of

constitutively activated RhoA induces ectopic TJs at the lateral membrane [49,50]. For a more

physiologically relevant example, hyperosmotic stress induces rapid expansion of TJs at the lateral

membrane [51,52]. Considering that Rho-ROCK pathway is known to be hyper-activated by the

hyperosmotic stress [53], the mechanism of supplying polymerizable claudin via the Rho-ROCK pathway is

likely to contribute to the adaptive expansion of TJs to hyperosmotic stress in epithelial cells.

While our results make clear that the cascade of reactions leading to claudin release is triggered

by Rho-ROCK-dependent activation of matriptase, the precise mechanism of this step is unclear. A

previous study showed that matriptase binds the actin cross-linking protein filamin, the depletion of which

severely impairs its activation [54]. Therefore, it is conceivable that actin reorganization associated with

Rho-ROCK activation allows filamin to engage matriptase and activate it. Alternatively, Rho-ROCK may

alter signaling pathways that regulate matriptase activity. Matriptase is intrinsically inhibited by Hai1; Hai1

is cleaved by the membrane type 1 matrix metalloproteases MMP14 and MMP7, which can stably

activate matriptase [55,56]. Notably, MMP14 is targeted to invadopodia in invasive cancer cells via

RhoA-dependent exocytosis [57]. The Rho-ROCK pathway may similarly facilitate the delivery of MMP14

and/or MMP7 to the plasma membrane to activate matriptase through Hai1 cleavage. Still another

possibility involves the type II transmembrane serine protease Tmprss2, the loss of which results in a

significant decrease in matriptase expression and disruption of the TJ barrier but the relationship between

Rho-ROCK and Tmprss2 is unknown [58]. Clearly, further studies are needed to elucidate how the

Rho-ROCK pathway activates matriptase.

Our analysis focused on matriptase as the protease that cleaves EpCAM/TROP2, but other

proteases that process EpCAM/TROP2 have been reported. Recently, Higashi et al. showed that

membrane-anchored serine proteinases (MASPs) cleave EpCAM [21]; MASPs are present at the apical

surface in polarized epithelial cells. Rupture of TJs results in a partial loss of the TJ fence function,

leading to a change in the localization of MASPs from the apical membrane to the lateral membrane. This

allows MASPs to cleave the EpCAM-claudin complex at the lateral membrane. By contrast, we found that

matriptase is always present in the lateral membrane and actively regulates *de novo* TJ formation under

the control of Rho-ROCK signaling. Whether these two pathways work independently or in parallel, or

whether each of these pathway functions only in specific tissues or contexts, needs to be investigated in

the future. Additionally, EpCAM is reportedly cleaved by A disintegrin and metalloprotease (ADAM) 10

and ADAM17 but it is unknown whether this contributes to *de novo* TJ formation [59]. In the present study,

overexpression of Hai1, which specifically inhibits matriptase activity, inhibited the formation of *de novo*

TJs in both apoptotic cell elimination and granular layer differentiation, suggesting that the primary means

of freeing sequestered claudin from EpCAM/TROP2 in these contexts is through matriptase.

In this paper, we show that Rho-ROCK-dependent matriptase activation enables *de novo* TJ

formation in the SG2 layer of the epidermis. Prior studies have shown that applying the ROCK activator

Narciclasine to mice with psoriatic dermatitis reduces inflammation [60]. Topical application of

oleoyl-l-α-lysophosphatidic acid (LPA), a potent Rho activator, was reported to increase the skin barrier in

mice [61]. However, the molecular basis for the activation of Rho-ROCK pathways to improve the epidermal

barrier was not elucidated. Our findings provide a rationale for the validity of these treatments.

Finally, we revealed that EpCAM/TROP2, besides its role in maintaining a stable pool of

unpolymerized claudin at the plasma membrane, functions as a chaperone to facilitate intracellular

transport of claudin from the Golgi apparatus to the plasma membrane. The molecular mechanism of

claudin transport to the plasma membrane remains largely unknown. In the future, it will be necessary to

further investigate how the interaction of EpCAM/TROP2 and claudin at the trans-Golgi networks helps

the intracellular transport of claudin.

Figure legends

Figure 1

Rho-ROCK controls *de novo* tight junction assembly between newly adjacent cells during apical

extrusion

(A)

Schematic showing the elimination of an apoptotic cell by apical extrusion and *de novo* tight junction (TJ)

formation. Tight junctions are established between newly adjacent cells (shown in green) concurrent with

extrusion of the apoptotic cell. The TJs between the apoptotic cell and the cells that effect extrusion

(shown in purple) gradually diminish as the new TJs mature, ensuring that the epithelial barrier is never

disrupted.

(B)

Live cell images of EpH4 cells expressing GFP-claudin-3 in cells neighboring laser-wounded cell. Cell

marked by an asterisk was wounded at time zero. The schematics represent the cells before laser

irradiation and after apical extrusion. Purple arrowheads indicate the old TJ with the apoptotic cell and

green arrowheads indicate the newly assembled TJ. The upper images (time = 60 and 80 min) show

higher magnification corresponding to each time point. The right panels are orthogonal views. Scale bar =

20 μm. (See also **Movie1**)

(C)

Immunofluorescence images showing apical extrusion at 60 min post-laser irradiation. Cells were stained with anti-claudin-3 pAb (green) and either anti-ZO-1 mAb (magenta, **upper**) or anti-occludin mAb (magenta, **lower**). Purple arrowheads indicate old TJs with the apoptotic cell and green arrowheads indicate newly assembled TJs. The bottom panels represent orthogonal views. Scale bar = 10 μm.

(D)

Live cell images of cells surrounding a laser-wounded cell expressing GFP-AnillinC as a probe for active RhoA. Cell marked by an asterisk was wounded at 0 min. The arrowheads indicate accumulation of active RhoA at the junction between an apoptotic cell and a neighboring cell (white) or between newly adjacent cells (yellow). Scale bar = 20 μm. (See also **Movie2**)

(E)

Immunofluorescence images showing apical extrusion at 60 min post-laser irradiation. Cells were stained with phosphorylated myosin light chain (ppMLC) mAb (magenta) and anti-NMIIB mAb (green). The asterisk in the X-Z image indicates the apoptotic cell. Scale bar = 10 μm.

(F)

Live cell images of cells surrounding a laser-wounded cell co-expressing GFP-claudin-3 and mScarlet-PLCδPH (membrane marker). Cells were pre-treated with H1152 and the cell marked by an asterisk was wounded at 0 min. The filled purple arrowhead indicates a TJ with an apoptotic cell and the unfilled purple arrowheads indicate the gradual dissolution of said TJ. Filled purple arrowheads indicate TJ between an apoptotic cell and neighboring cells, while unfilled purple arrowheads indicate the disappearance of TJ. Images of scarlet-PLCδ PH are projections of basal confocal slices and the extending lamellipodia are indicated by the white arrowheads. The inset enlarged at right—and illustrated below—show the absence of new TJs among the now adjacent cells A-E. Scale bar = 20 μm. (See also **Movie3**).

(G)

Bar graph showing the paracellular flux of 70kDa FITC-dextran tracer molecule at 9 h post-doxorubicin treatment (4 μM) (control: N=3, H1152: N=5; error bar: ±SD; Mann–Whitney U test).

(H)

Schematic illustrating the formation of new TJs between neighbor cells concurrent with extrusion of the apoptotic cell. The formation of these TJ requires the activation of RhoA.

Figure 2
Claudins in epidermal cells are present at the plasma membrane in an unpolymerized state before differentiation in the granular layer
(A)
Schematic showing the differentiation of human keratinocyte (HaCaT cells) into granular-like cells in high-calcium (9.8 mM) medium supplemented with a JNK inhibitor (40 μM) (Ca+JNK inh medium) for 24 h.
(B)

HaCaT cells were cultured in normal medium (control) or Ca+JNK inh medium for 24 h, fixed, and then

stained with anti-claudin-1 pAb (green), anti-E-cadherin mAb (magenta) and phalloidin (grayscale; **left**) or

anti-ZO-1 mAb (green), anti-alpha-catenin (magenta) and phalloidin (grayscale; **right**). X-Z images are

shown below. Yellow arrowheads indicate TJs. Scale bar = 20 µm.

(C)

Whole-cell lysates of cells cultured in normal medium or Ca+JNK inh medium for 24h were immunoblotted

with the indicated antibodies. Molecular weight measurements are in kDa.

(D)

Mouse ear whole-mount immunofluorescence analysis for claudin-1, ZO-1 and E-cadherin. The asterisk

indicates the nucleus of cells belonging to that layer. The panels in the bottom row represent orthogonal

views. Scale bar = 20 µm.

Figure 3

De novo TJ formation in the SG2 layer requires activation of Rho-ROCK

(A and B)

RhoA activities in undifferentiated and differentiated HaCaT cells. Cells were lysed and subjected to a

GST-Rhotekin pulldown assay. Total cell lysates and precipitates were analyzed by immunoblotting with

anti-RhoA mAb. Molecular weight measurements are in kDa. **B** is a graph illustrating the rate of RhoA

activities. RhoA activity was calculated by dividing the intensity of activated RhoA (pulldown signal)

by the intensity of input signal (N=6; error bar: ±SD; Student's t test).

(C and D)

HaCaT cells were cultured in normal medium or Ca+JNK inh medium for 24 h, fixed, and then stained

with anti-ppMLC mAb (green) and anti–E-cadherin mAb (magenta). Scale bar: 20 µm. Graph showing the

quantification of junctional enrichment of ppMLC signals **(D)**. The quantification methods are illustrated in

Figure S1 and details are described in **Methods** (control: N=5, Ca+JNK inh: N=8; error bar: ±SD;

Student's t test).

(E)

Mouse ear whole-mount immunofluorescence analysis for ppMLC, ZO-1 and E-cadherin. The asterisk

indicates the nucleus of cells belonging to the indicated layer.

(F and G)

HaCaT cells expressing GFP-claudin-3 were cultured in normal medium or Ca+JNK inh medium

supplemented with DMSO (control) or Y27632 (ROCK inhibitor) for 24 h, fixed, and then stained with

anti–E-cadherin mAb (magenta). Yellow arrowheads indicate TJs. Scale bar: 20 µm. **G** is a bar graph

illustrating the junctional enrichment of claudin-3 after differentiation (control: N=5, Ca+JNK inh: N=6;

error bar: ±SD; Student's t test).

(H)

Schematic illustrating TJ formation in the epidermal granular layer. In the epidermis, all keratinocytes of

basal, spinous and granular layer express claudin-1. However, functional TJs are only formed in the SG2

layer. The activation of the Rho-ROCK pathway is crucial for the formation of TJs in the SG2 layer.

Figure 4

Rho-ROCK activation induces ectopic TJ formation in undifferentiated keratinocytes

(A)

Undifferentiated HaCaT cells expressing constitutive active RhoA (RhoA CA) were stained with

anti-claudin-3 pAb (green) and phalloidin (grayscale). The arrowheads indicate ectopic tight junction.

Scale bar: 20 µm.

(B)

Undifferentiated HaCaT cells expressing RhoA CA were stained with anti-claudin-3 pAb (magenta) and

either anti-ZO-1 mAb (green; **upper**) or anti-occludin mAb (green; **lower**). The arrowheads indicate

ectopic TJs. Scale bar: 20 µm.

(C and D)

Confluent undifferentiated HaCaT cells were cultured in serum-free medium for 24h, treated with a

recombinant bacterial cytotoxic necrotizing factor (CNF) toxin (CN-03), also known as Rho activator II (1

µg/ml) for 2 h, fixed, and then stained with anti-claudin-3 pAb (green), anti-E-cadherin mAb (magenta)

and phalloidin (grayscale). Scale bar: 20 µm. **(D)** shows an alternate field of view corresponding to the

experiment in **(C)**.

(E and F)

Confluent undifferentiated HaCaT cells were treated with DMSO (control) or Narciclasine (100 nM; ROCK

activator) for 4 h, fixed, and then stained with anti-claudin-3 pAb (green), anti-E-cadherin mAb (magenta)

and phalloidin (grayscale). Scale bar: 20 µm. **(F)** shows an alternate field of view of the experiment in (**E**).

(G)

Junctional enrichment of claudin-3 was quantified based on the method described in **Figure S1** (N=6;

error bar: ±SD; Tukey-Kramer One-way Anova).

(H and I)

Western blot of Triton X-100–soluble fractions and -insoluble fractions from HaCaT cells treated with Rho

activator (CN-03) or Narciclasine. The proportion of insoluble claudin-1 was quantified in **(I)** (N=5; error

bar: ±SD; Kruskal-Wallis test followed by Steel-Dwass post hoc test; *: $p < 0.05$).

Figure 5

Rho-ROCK pathway acts upstream of matriptase activation in *de novo* TJ formation

(A and B)

Whole-cell lysates of WT treated with Rho activator (CN-03), Narciclasine or acid medium (pH 6.0) were

immunoblotted with the indicated antibodies. Molecular weight measurements are in kDa. Schematic

showing the sequential activation of matriptase. M24 mAb detects total Matriptase and M69 mAb the

activated form.

(B) Quantification of the cleaved TROP2 fragment relative to total TROP2. (N=4; error bar: ±SD;

Kruskal-Wallis test followed by Steel-Dwass post hoc test; * : $p < 0.05$).

(C and D)

Confluent undifferentiated HaCaT cells were cultured in normal (control) or acid medium (pH 6.0) for 20

min, fixed, and then stained with anti-claudin-3 pAb (green), anti-E-cadherin mAb (magenta) and

phalloidin (grayscale). Scale bar = 20 μm. **D** is a bar graph illustrating the junctional enrichment of

claudin-3 (N=6; error bar: ±SD; Student's t test).

(E and F)

Western blot of Triton X-100–soluble and -insoluble fractions from confluent undifferentiated HaCaT cells

cultured in acid medium. The proportion of insoluble claudin-1 was quantified in **(F)** (N=5; error bar: ±SD;

Student's t test).

(G)

Undifferentiated HaCaT cells treated with DMSO (control) or Camostat (serine protease inhibitor)

expressing constitutive active RhoA (Rho CA) were fixed and stained with anti-claudin-3 pAb (green) and

phalloidin (grayscale). Scale bar: 20 μm.

(H)

Bar graph illustrating the extent to which the matriptase inhibitor Camostat negated the

Narciclasine-induced claudin enrichment at cell-cell contacts. After inhibition of serine protease with

Camostat, confluent undifferentiated HaCaT cells were treated with Narciclasine (100 nM; ROCK

activator) for 4 h and stained to quantify the junctional enrichment of claudin-3 (N=6; error bar: ±SD;

Student's t test).

(I)

Undifferentiated HaCaT cells expressing constitutive active RhoA (Rho CA) were stained with

anti-TROP2 mAb (green) and claudin-4 mAb (magenta). Insets are enlarged images. Scale bar: 20 μm.

Figure 6

TROP2 is required for intracellular transport of claudin to the plasma membrane in keratinocytes

(A and B)

Whole-cell lysates of WT and TROP2 KO HaCaT cells were immunoblotted with the indicated antibodies.

Molecular weight measurements are in kDa. The expression level of claudin-1 normalized to α-tubulin

was quantified in **(B)** (N=4; error bar: ±SD; Student's t test).

(C and D)

Representative western blots of surface and total amount of claudin-1. Molecular weight measurements

are in kDa. The graph of **(D)** illustrates the quantification of claudin-1 present on the cell surface relative to

the total amount (N=3; error bar: ±SD; Student's t test).

(E)

Undifferentiated TROP2 KO HaCaT cells were stained with anti-claudin-1 pAb (green) and anti-GM130

mAb (magenta; Golgi marker). Insets are enlarged images. Scale bar: 20 μm.

(F)

Undifferentiated WT or TROP2 KO HaCaT cells expressing constitutive active RhoA (Rho CA) were

stained with anti-claudin-3 pAb (green) and phalloidin (grayscale). Scale bar: 20 μm.

(G)

Undifferentiated WT and TROP2 KO HaCaT cells were co-cultured, incubated in acid buffer (pH 6.0) for

20min, fixed, and then stained with anti-claudin-4 mAb (green) and anti-TROP2 mAb (magenta). Dotted

line overlays the border between WT and TROP2 KO cells. Scale bar: 20μm.

(H)

Bar graph illustrating the impairment of either Narciclasine- or acid medium-induced claudin-3

accumulation at cell-cell contacts by TROP2 depletion. Confluent undifferentiated WT or TROP2 KO

HaCaT cells were treated with Narciclasine (ROCK activator) or acid medium (pH 6.0) and stained to

quantify junctional enrichment of claudin-3 (N=5; error bar: ±SD; Tukey-Kramer One-way Anova).

(I)

Schematic illustrating the mechanism of ectopic TJ formation through RhoA activation. In wild-type

keratinocytes, claudin and TROP2 form a complex in the Golgi apparatus and are subsequently

transported to the plasma membrane. In contrast, in cells lacking TROP2, claudin, which cannot form a

complex with TROP2, accumulates in the Golgi apparatus, leading to a significant decrease in the amount

of claudin at the plasma membrane. In wild-type cells, TROP2 is cleaved by matriptase activated via the

Rho-ROCK pathway. This cleavage results in the breakdown of the TROP2-claudin complex, allowing

released claudins to form *de novo* TJs.

Figure 7

Matriptase cleaves TROP2 In the SG2 layer of epidermis.

(A)

WT HaCaT cells were cultured in Ca+JNK inh medium for indicated time points and time-course change

of indicated proteins were examined by western blotting. (N=5; error bar: ±SD; Kruskal-Wallis test

followed by Steel-Dwass post hoc test; * : $p < 0.05$).

(B) Quantification of the cleaved TROP2 fragment relative to total TROP2. (N=5; error bar: ±SD;

Kruskal-Wallis test followed by Steel-Dwass post hoc test; * : $p < 0.05$).

(C, D and E)

HaCaT cells were cultured in normal medium or Ca+JNK inh medium for 24h, fixed and then stained with

the m24 mAb (total matriptase: magenta) and anti-claudin-3 pAb (green ; **C**) or with the M69 mAb

(activated matriptase: magenta) and anti-claudin-3 pAb (green ; **D**). Scale bar = 20 μm. **E** is a bar graph

illustrating the junctional enrichment of activated matriptase (control: N=3, Ca+JNK inh: N=4; error bar:

±SD; Student's t test)**.**

(F)

HaCaT cells were cultured in normal medium or Ca+JNK inh medium supplemented with DMSO (control)

or Camostat (serine protease inhibitor) for 24 h, fixed, and then stained with anti-claudin-3 pAb (green),

anti-α-catenin mAb (α-18; magenta). Scale bar: 20 μm.

(G)

WT HaCaT cells and cells over-expressing Hai1-GFP were co-cultured in Ca+JNK inh medium for 24 h,

fixed, and then stained with anti-claudin-1 pAb. Scale bar: 20 μm.

(H)

Bar graph illustrating the effects of either Camostat treatment or over-expression of Hai1 on accumulation

of claudin-3 at cell-cell contacts in HaCaT cells cultured in Ca+JNK inh medium (N=6; error bar: ±SD;

Tukey-Kramer One-way Anova).

(I)

Mouse ear whole-mount immunofluorescence analysis for TROP2 and ZO-1. The asterisk indicates the

nucleus of cells belonging to the indicated layer.

(J)

Schematic illustrating the mechanism of TJ formation during differentiation into the granular layer in the

epidermis. In undifferentiated cells, claudin is expressed and bound to TROP2 on the plasma membrane.

As cells differentiate into the granular layer, RhoA becomes activated, followed by the activation of

matriptase, leading to the cleavage of TROP2. The cleavage of TROP2 results in the dissociation of

claudin, and the dissociated claudins then polymerized into *de novo* TJs.

Figure 8

Signaling through RhoA-Matriptase-EpCAM is critical for the formation of *de novo* TJ following

apical extrusion

(A and B)

Whole-cell lysates of WT and EpCAM KO EpH4 cells were immunoblotted with the indicated antibodies.

Molecular weight measurements are in kDa. The expression level of claudin-1 normalized to α-tubulin

was quantified in **(B)** (N=3; error bar: ±SD; Student's t test).

(C)

WT and EpCAM KO EpH4 cells were co-cultured, fixed, and then stained with anti-EpCAM pAb (green)

and anti-claudin-4 mAb (magenta). Dotted line overlays the border between WT and EpCAM KO cells.

Scale bar: 20 μm.

(D)

WT or EpCAM KO EpH4 cells were cultured, fixed, permeabilized with digitonin, and then stained with

anti-claudin-7 pAb (green) and anti-ZO-1 mAb (magenta). Yellow arrowheads indicate claudin-7 at the

lateral membrane, while white arrowheads indicate the intercellular accumulation of claudin-7. Scale bar:

20 μm.

(E)

EpCAM KO EpH4 cells permeabilized with digitonin were stained with anti-claudin-7 pAb (green) and anti-GM130 mAb (magenta; Golgi marker). Scale bar: 20 μm.

(F)

WT and EpH4 KO EpH4 cells were co-cultured, incubated in acid buffer (pH 6.0) for 20min, fixed, and then stained with anti-claudin-4 mAb (green) and anti-EpCAM pAb (magenta). Yellow arrowheads indicate the formation of ectopic TJ strands on the basolateral membrane. Scale bar: 20 μm.

(G, H and I)

Live cell imagings of EpCAM KO EpH4 cells expressing GFP-claudin-3 **(G)**, EpH4 cells expressing GFP-claudin-3 treated with Camostat **(H)**, or EpH4 expressing both Hai1-mScarlet and GFP-claudin-3 **(I)** after laser irradiation. Cell marked by an asterisk was wounded at time zero. The right diagram depicts the condition of TJs 60 min after laser irradiation. The navy color represents pre-existing TJs, and the dashed lines indicate the absence of newly formed TJs between neighboring cells. Scale bar = 20 μm. (See also **Movies 4-6**).

(J)

Bar graph showing the paracellular flux of 70kDa FITC-dextran tracer molecule at 9 h post-doxorubicin treatment (4 μM) (control: N=7, EpCAM KO: N=6, Camostat: N=4, Hai1 over-expression (OE): N=3; error bar: ±SD; Dunnett's test).

Figure 9

Activation of Rho-ROCK pathway triggers *de novo* TJ formation via disruption of the EpCAM/TROP2-claudin complex

De novo TJ formation associated with apoptotic cell elimination in simple epithelia and with keratinocyte differentiation in stratified epithelia, is regulated by a common molecular mechanism. In both types of epithelia, EpCAM or TROP2 is necessary for the intracellular transport of claudin from the Golgi apparatus to the plasma membrane. Claudin, when in complex with EpCAM or TROP2, cannot spontaneously polymerize into TJs at the plasma membrane. However, upon activation of the Rho-ROCK pathway, which activates matriptase, EpCAM or TROP2 is cleaved by matriptase. This cleavage breaks down the EpCAM/TROP-2-claudin complex, enabling claudin to polymerize into TJs. Consequently, rapid TJ formation occurs without *de novo* transcription or translation.

Supplemental figure legends

Figure S1

Details of the image analysis for quantification of junctional enrichment

Workflow for quantification of junctional enrichment of indicated molecules. Maximum intensity projections of E-cadherin immunostaining were processed to binarize and reduce noise by applying the Gaussian filter. After segmentation using cellpose2, junction areas were extracted by edge extraction and dilation of segmentation images, and junction intensities were obtained by overlaying the junction area. Junctional

enrichment of the indicated molecule (claudin-3 in this example) was quantified as the signal intensity of

the indicated molecule divided by the junction area.

Figure S2 (Related to Figure 4)

Claudin accumulation is induced by constitutive active ROCK but not by either constitutive active

Rac or Cdc42

Undifferentiated HaCaT cells expressing constitutive active Rac (**A; upper**), constitutive active Cdc42 (**A;**

lower) or ROCK delC (**B**) were stained with anti-claudin-3 pAb and phalloidin. The arrowheads indicate

ectopic TJs. Scale bar: 20 µm.

Figure S3 (Related to Figure 6)

Other TJ components are co-localized with claudin puncta on the plasma membrane of

undifferentiated TROP2 KO HaCaT cells

Undifferentiated TROP2 KO HaCaT cells were stained with anti-claudin-1 pAb (green) and ether

anti-ZO-1 mAb (magenta; **upper**) or anti-occludin mAb (magenta; **lower**). Scale bar: 20 µm.

Movie 1.

EpH4 cells expressing GFP-claudin-3 in cells neighboring laser-wounded cell were imaged. Frames were

taken every 1 min. (Scale bar, 20 µm.)

Movie 2.

EpH4 cells expressing GFP-AnillinC in cells neighboring laser-wounded cell were imaged. Frames were

taken every 1 min. (Scale bar, 20 µm.)

Movie 3.

EpH4 cells co-expressing GFP-claudin-3 and mScarlet-PLCδPH (membrane marker) in cells neighboring

laser-wounded were imaged. Cells were pre-treated with H1152 (ROCK inhibitor). Frames were taken

every 10 min. (Scale bar, 20 µm.)

Movie 4.

EpCAM KO EpH4 cells expressing GFP-claudin-3 in cells neighboring laser-wounded cell were imaged.

Frames were taken every 30 sec. (Scale bar, 20 µm.)

Movie 5.

EpH4 cells expressing GFP-claudin-3 in cells neighboring laser-wounded were imaged. Cells were

pre-treated with Camostat (matriptase inhibitor). Frames were taken every 5 min. (Scale bar, 20 µm.)

Movie 6.

EpH4 cells co-expressing GFP-claudin-3 and Hai1-mScarlet in cells neighboring laser-wounded were

imaged. Frames were taken every 5 min. (Scale bar, 20 μm.)

Materials and Methods

Antibodies and other reagents

The following primary antibodies were used for immunofluorescence microscopy and

immunoblotting: rabbit anti-claudin-3 pAb (34-1700; Thermo Fisher Scientific); mouse anti-claudin-4 mAb

(32-9400; Thermo Fisher Scientific); and rabbit anti-claudin-7 pAb (34-9100; Thermo Fisher Scientific);

rabbit anti-α-catenin pAb (C2081; Sigma Aldrich); rabbit anti-claudin-1 pAb (SAB4200534; Sigma

Aldrich); rat anti-E-cadherin mAb (ECCD2; Takara Bio); rabbit anti-RhoA mAb (2117; Cell Signaling

Technology); rabbit anti-phosoho-Myosin Light Chain 2 (Thr18/Thr19) mAb (95777; Cell Signaling

Technology); rabbit anti-TROP2 mAb (ab214488; abcam); rabbit anti-EpCAM pAb (ab71916; abcam);

mouse anti-GM130 mAb (610822; BD). Mouse anti–α-tubulin (12G10) mAb, rat anti-occludin (Moc37)

mAb, mouse anti–ZO-1 (T8-754) mAb and mouse myosin IIB (CMII-23) were produced in-house. Mouse

matriptase mAbs, M24 and M69, were generous gifts from Dr. Chen-yong Lin, Georgetown University

Medical Center, Washington DC. Rat anti–α-catenin mAb (α18) was a generous gift from Dr. A.

Nagafuchi, Nara Medical University, Nara, Japan.

Secondary antibodies were as follows: Cy2-conjugated donkey anti-rat IgG antibody

(712-225-150), anti-mouse IgG antibody (715-225-151), and anti-rabbit-IgG antibody (711-225-152);

Cy3-conjugated donkey anti-rat IgG antibody (712-165-153), anti-mouse IgG antibody (715-165-150),

and anti-rabbit IgG antibody (711-165-152);Cy5-conjugated donkey anti-rabbit IgG antibody

(711-175-152; Jackson ImmunoResearch) and HRP-conjugated anti-rat IgG antibody (HAF005; R&D

Systems), anti-mouse-IgG antibody (A90-516P; Bethyl Laboratories), and anti-rabbit IgG antibody

(4030-05; Southern Biotech).

F-actin was visualized with Alexa Fluor 647–phalloidin (A22287; Thermo Fisher Scientific).

Nucleus was visualized with DAPI (049-18801; Wako Pure Chemical Industries).

Chemicals and recombinant proteins were obtained as follows: H-1152 (555550; Sigma Aldrich);

Y27632 (08945-84; Nacalai Tesque); Doxorubicin Hydrochloride (040-21521; Wako); SP600125

(197-16591; Wako); Rho activator II (CN03-A; Cytoskeleton Inc); Camostat Mesylate (C2977; Tokyo

Chemical Industry Co., Ltd.).

Cell culture

EpH4 cells, HaCaT cells, and HEK293 cells were grown in DMEM supplemented with 10%

(vol/vol) FCS. All cell cultures were maintained in 5% CO_2 atmosphere at 37˚C.

For live imaging, cells were cultured on 35 mm glass-base dish for 48 h before observation. For

immunostaining, cells were cultured on 15mm coverslip. For mix-culture experiments, two types of cells

were co-cultured at a ration of 1:1.

For differentiation of HaCaT cells into granular-like cells, cells were cultured in high-calcium (9.8

mM) medium supplemented with a JNK inhibitor (40 µM) for 24 h. For inhibition of ROCK, cells were

treated H-1152 (final 2 µM) or Y27632 (final 20 µM) for 30min. For inhibition of serine protease, cells were

treated Camostat (final 10 µM) for 6 h. For activation of Rho, after serum starved for 24 h, cells were

treated Rho activator (final 1 µg/ml) for 2 h. For activation of ROCK, cells were treated Narciclasine (final

100 nM) for 4 h. For activation of matriptase, cells were cultured in pH 6.0 citric acid-phosphate buffer for

20 min.

Plasmids

We established KO cells by using the CRISPR-Cas9 system. Oligonucleotides were

phosphorylated, annealed, and cloned into the BsmBI site of pLenti-CRISPR v2 vector according to the

Zhang laboratory protocols (F. Zhang, MIT, Cambridge, MA). The target sequences were as follows;

Mouse EpCAM: 5′- GGGCGATCCAGAACAACGAT -3′

Human TROP2: 5′- GGCGGCGGTGACCGGCCACA -3′

Other expressing vectors were constructed on the pLenti-CMV backbone.

Transfection and generation of knockout cell lines

Transfections were performed by using the PEI-max (24765-1; Polysciences Inc.). Lentiviruses

were produced in HEK293 cells by transfecting the expression vector with the packaging and envelop

vectors (psPAX2 and pMD2.G) and harvested by centrifugation. Infections wre carried out in low calcium

medium with polybrene. Cells remained in infection media for 48 h, followed by antibiotic selection (400

µg/ml neomycin; 4.5 µg/ml puromycin; 200 µg/ml hygromycin).

To establish knockout cell lines, cells were seeded into 96-well plates after drug selection, and at

least 10 clones were collected and examined by western blotting to check for loss of expression of the

protein of interest. HaCaT cells expressing GFP-claudin-3 or Hai1-GFP and EpH4 cells expressing

GFP-claudin3 and/or Hai1-scarlet were obtained using a cell sorter (SH800; SONY)

Live imaging

Fluorescence imaging was performed using a confocal microscope (LSM900; Carl Zeiss

MicroImaging) equipped with a 63x/1.40 NA, oil immersion Plan-APO objective with a heated stage set to

37˚C and cells were cultured in Leibovitz's L-15 Medium (1415064; Thermo Fisher Scientific)

supplemented with 10 % (vol/vol) FCS. Images were acquired using Zen 2012 (Carl Zeiss MicroImaging).

For induced apoptosis, nuclei of any selected cells were injured by MicroPoint Laser (wavelength =

435nm, duration < 4 nsec, laser energy < 40 µJ, single pulse, OXFORD Instruments).

Immunofluorescence microscopy

Cells cultured on coverslips were fixed with 3% formalin prepared in PBS for 15 min at room

temperature (RT), permeabilized with 0.4% Triton X-100/PBS for 5 min and blocked with 1% BSA

prepared in PBS for 1h at RT. In **Figures 8D and 8E**, cells were fixed with 4% paraformaldehyde (PFA)

/PBS for 15 min and permeabilized with 100 µM digitonin/PBS for 10min at RT. When using

anti-matriptase mAbs, cells were fixed with 4% PFA/PBS and permeabilized with 0.4% Triton

X-100/PBS. Antibodies were diluted in the blocking solution. Cells were incubated with primary

antibodies for 1 h at RT and with secondary antibodies for 30 min at RT. Samples were observed at RT

with the confocal microscope.

7   To quantify junctional enrichment of indicated molecule, cells were segmented based on

maximum projection of E-cadherin immunostaining using the cellpose2 according to the schematic in

Figure S1[62]. Junction areas were obtained by the "find edge" and "dilute" command in ImageJ. Junctional

signal intensities were obtained by the "Image calculator" command in ImageJ. Finally, junctional

enrichment of the indicated molecule was quantified as the signal intensity of the indicated molecule

divided by the junction area.

Mouse ear whole-mount immunofluorescence microscopy

15   The detachment of the epidermis from the mouse ear and fluorescence immunostaining were

performed as previously described [63]. In brief, after euthanasia of ICR mouse, the ears were immediately

cut off at its root and floated in calcium-containing PBS. The dorsal skin was removed, followed by the

removal of cartilage from the ventral side skin. The epidermis and dermis were separated after dispase

treatment. The separated epidermis was fixed with 4% PFA for 10 min and permeabilized with 1% Triton

X-100 for 10 min. After washing, samples were blocked with 1% BSA/PBS, and incubated with primary

antibodies overnight at 4°C and with secondary antibodies for 2 h at RT. The samples were mounted in

Mowiol. Fluorescence imaging was performed using a confocal microscope equipped with a 40x/1.40 NA,

oil immersion Plan-APO objective.

Immunoblotting

26   Cells were cultured in 30 mm dish or 6-well plate and lysed in SDS sample buffer and boiled for

5 min. Lysates were prepared in SDS sample buffer without 2-Mercaptoethanol and were not boiled.

Samples were resolved by SDS-PAGE and transferred to nitrocellulose membrane. The membrane was

blocked with 5% skim milk/0.1% Tween 20/TBS(TBST) for 1 h at RT. Indicated proteins were probed by

sequential incubation with the primary antibody and HRP-conjugated secondary antibody prepared in 5%

BSA/TBST for 1 h at RT each. After each antibody reaction, nitrocellulose membranes were washed three

times with TBST for 10 minutes. Signals were detected by mixing A (100 mM Tris-HCl [pH 8.5], 0.4 mM

p-coumaric acid, 5 mM luminol) and B (100 mM Tris-HCl [pH 8.5], 0.04% H_2O_2). Chemiluminescence was

captured using the LAS-3000 Imaging system (Fujifilm).

Cell surface protein biotinylation

37   Cell surface protein biptonylation was performed according to a previously reported protocol [64].

The biotinylation of surface proteins was performed using EZ-Link sulfo-NHS-SS-biotin (21217; Thermo

Fisher Scientific). Biotinylated proteins were isolated using High-Capacity Streptavidin Agarose Resin

(20361; Thermo Fisher Scientific).

Paracellular tracer flux measurement

10^5 EpH4 cells were seeded on a 12-mm-diameter cell culture insert (83.3931.040; SARSTEDT)

and cultured with daily medium changes. After 6 days, cells were basally treated with 4 µM Doxorubicin

Hydrochloride for 9 h, and 70kDa FITC-dextran (46945; Sigma Aldrich) was added to the medium in the

apical compartment at a concentration of 1 mg/ml. Medium was collected from the basal compartment

after 1 h, and the FITC signal was measured with a fluorometer.

Active RhoA pulldown assay

RhoA activation was assessed using GST-tagged rhotekin RBD domain (GST-RBD), which

binds to the GTP-bound form of RhoA. Undifferentiated or differentiated HaCaT cells were lysed

using lysis buffer (20 mM HEPES (pH 7.5), 150 mM NaCl, 5 mM MgCl2, 1% Triton X, 1 mM DTT, 100

µM APMSF, 10 µg/ml aprotinin, 10 µg/ml leupeptin), and after centrifugation, the supernatant was

aliquoted as input and the remainder incubated with GST-RBD bound to glutathione Sepharose

beads at 4°C for 2 h. After washing with lysis buffer, pulldown samples were eluted with SDS sample

buffer. Immunoblotting was performed using an anti-RhoA mAb, and the band intensities of RhoA in

both input and pulldown samples were quantified using Fiji/ImageJ. RhoA activity was calculated by

dividing the intensity of activated RhoA (pulldown) by the intensity of input, with undifferentiated cells

set to 1.00.

Quantification and Statistical analysis

Error bars indicate 95% confidence intervals. The number and statistical details of

independent experiments is indicated in figure legend. Each point on the graph represents an

individual biological replicate. Unless otherwise specified, microcopy images are representatives of

at least three independent experiments. The microscope images of the mouse epidermis are

representative ones sourced from a single mouse. Statistical analysis was performed in Python 3.11.3

using scipy 1.10.1, statsmodels 0.14.0 and scikit_posthocs 0.9.0 packages. Graphs were created using

matplotlib 3.7.1 and seaborn 0.12.2. The number and statistical details of independent experiments is

indicated in figure legend. Error bar indicates SD.

Acknowledgments

We thank all members of the Ikenouchi laboratory (Department of Biology, Faculty of Sciences, Kyushu

University) and Prof. Kyoko Shirakabe (Ritsumeikan Univ) for helpful discussions. This work was

supported by JSPS KAKENHI (JP22H02618, JP23K18141, JP25H01325 and JP25H00994 [J.I.],

Author contributions

References

1. Piontek, J., Krug, S.M., Protze, J., Krause, G., and Fromm, M. (2020). Molecular architecture and

assembly of the tight junction backbone. Biochim. Biophys. Acta Biomembr. *1862*, 183279.

2. Horowitz, A., Chanez-Paredes, S.D., Haest, X., and Turner, J.R. (2023). Paracellular permeability

and tight junction regulation in gut health and disease. Nat. Rev. Gastroenterol. Hepatol. *20*,

417–432.

3. Tsukita, S., Furuse, M., and Itoh, M. (2001). Multifunctional strands in tight junctions. Preprint,

https://doi.org/10.1038/35067088 https://doi.org/10.1038/35067088.

4. Gu, Y., and Rosenblatt, J. (2012). New emerging roles for epithelial cell extrusion. Curr. Opin. Cell

Biol. *24*, 865–870.

5. Duszyc, K., von Pein, J.B., Ramnath, D., Currin-Ross, D., Verma, S., Lim, F., Sweet, M.J., Schroder,

K., and Yap, A.S. (2023). Apical extrusion prevents apoptosis from activating an acute inflammatory

program in epithelia. Dev. Cell *58*, 2235-2248.e6.

6. Takeuchi, Y., Narumi, R., Akiyama, R., Vitiello, E., Shirai, T., Tanimura, N., Kuromiya, K., Ishikawa,

S., Kajita, M., Tada, M., et al. (2020). Calcium Wave Promotes Cell Extrusion. Curr. Biol. *30*,

670-681.e6.

7. Rosenblatt, J., Raff, M.C., and Cramer, L.P. (2001). An epithelial cell destined for apoptosis signals

its neighbors to extrude it by an actin- and myosin-dependent mechanism. Curr. Biol. *11*, 1847–1857.

8. Furuse, M., Hata, M., Furuse, K., Yoshida, Y., Haratake, A., Sugitani, Y., Noda, T., Kubo, A., and

Tsukita, S. (2002). Claudin-based tight junctions are crucial for the mammalian epidermal barrier: A

lesson from claudin-1-deficient mice. J. Cell Biol. *156*, 1099–1111.

9. Sugawara, T., Iwamoto, N., Akashi, M., Kojima, T., Hisatsune, J., Sugai, M., and Furuse, M. (2013).

Tight junction dysfunction in the stratum granulosum leads to aberrant stratum corneum barrier

function in claudin-1-deficient mice. J. Dermatol. Sci. *70*, 12–18.

10. Kulukian, A., and Fuchs, E. (2013). Spindle orientation and epidermal morphogenesis. Philos. Trans.

R. Soc. Lond. B Biol. Sci. *368*, 20130016.

11. Brandner, J.M., Kief, S., Grund, C., Rendl, M., Houdek, P., Kuhn, C., Tschachler, E., Franke, W.W., and Moll, I. (2002). Organization and formation of the tight junction system in human epidermis and cultured keratinocytes. Eur. J. Cell Biol. *81*, 253–263.

12. Yoshida, K., Yokouchi, M., Nagao, K., Ishii, K., Amagai, M., and Kubo, A. (2013). Functional tight junction barrier localizes in the second layer of the stratum granulosum of human epidermis. J. Dermatol. Sci. *71*, 89–99.

13. Yokouchi, M., Atsugi, T., van Logtestijn, M., Tanaka, R.J., Kajimura, M., Suematsu, M., Furuse, M., Amagai, M., and Kubo, A. (2016). Epidermal cell turnover across tight junctions based on Kelvin's tetrakaidecahedron cell shape. Elife *5*. https://doi.org/10.7554/elife.19593.

14. Rübsam, M., Mertz, A.F., Kubo, A., Marg, S., Jüngst, C., Goranci-Buzhala, G., Schauss, A.C., Horsley, V., Dufresne, E.R., Moser, M., et al. (2017). E-cadherin integrates mechanotransduction and EGFR signaling to control junctional tissue polarization and tight junction positioning. Nat. Commun. *8*, 1250.

15. Furuse, M., Sasaki, H., Fujimoto, K., and Tsukita, S. (1998). A single gene product, claudin-1 or -2, reconstitutes tight junction strands and recruits occludin in fibroblasts. J. Cell Biol. *143*, 391–401.

16. Gonschior, H., Schmied, C., Van der Veen, R.E., Eichhorst, J., Himmerkus, N., Piontek, J., Günzel, D., Bleich, M., Furuse, M., Haucke, V., et al. (2022). Nanoscale segregation of channel and barrier claudins enables paracellular ion flux. Nat. Commun. *13*, 4985.

17. Ladwein, M., Pape, U.-F., Schmidt, D.-S., Schnölzer, M., Fiedler, S., Langbein, L., Franke, W.W., Moldenhauer, G., and Zöller, M. (2005). The cell-cell adhesion molecule EpCAM interacts directly with the tight junction protein claudin-7. Exp. Cell Res. *309*, 345–357.

18. Nübel, T., Preobraschenski, J., Tuncay, H., Weiss, T., Kuhn, S., Ladwein, M., Langbein, L., and Zöller, M. (2009). Claudin-7 regulates EpCAM-Mediated functions in tumor progression. Mol. Cancer Res. *7*, 285–299.

19. Barth, A.I.M., Kim, H., and Riedel-Kruse, I.H. (2018). Regulation of epithelial migration by epithelial cell adhesion molecule requires its Claudin-7 interaction domain. PLoS One *13*, e0204957.

20. Wu, C.-J., Mannan, P., Lu, M., and Udey, M.C. (2013). Epithelial Cell Adhesion Molecule (EpCAM) Regulates Claudin Dynamics and Tight Junctions. J. Biol. Chem. *288*, 12253–12268.

21. Higashi, T., Saito, A.C., Fukazawa, Y., Furuse, M., Higashi, A.Y., Ono, M., and Chiba, H. (2023). EpCAM proteolysis and release of complexed claudin-7 repair and maintain the tight junction barrier. J. Cell Biol. *222*. https://doi.org/10.1083/jcb.202204079.

22. Lenárt, S., Lenárt, P., Šmarda, J., Remšík, J., Souček, K., and Beneš, P. (2020). Trop2: Jack of All Trades, Master of None. Cancers *12*. https://doi.org/10.3390/cancers12113328.

23. Nakato, G., Morimura, S., Lu, M., Feng, X., Wu, C., and Udey, M.C. (2020). Amelioration of Congenital Tufting Enteropathy in EpCAM (TROP1)-Deficient Mice via Heterotopic Expression of TROP2 in Intestinal Epithelial Cells. Cells *9*. https://doi.org/10.3390/cells9081847.

24. Szabo, R., Ward, J.M., Artunc, F., and Bugge, T.H. (2022). EPCAM and TROP2 share a role in claudin stabilization and development of intestinal and extraintestinal epithelia in mice. Biol. Open *11*. https://doi.org/10.1242/bio.059403.

25. Sivagnanam, M., Mueller, J.L., Lee, H., Chen, Z., Nelson, S.F., Turner, D., Zlotkin, S.H., Pencharz, P.B., Ngan, B.-Y., Libiger, O., et al. (2008). Identification of EpCAM as the gene for congenital tufting enteropathy. Gastroenterology *135*, 429–437.

26. Kozan, P.A., McGeough, M.D., Peña, C.A., Mueller, J.L., Barrett, K.E., Marchelletta, R.R., and Sivagnanam, M. (2015). Mutation of EpCAM leads to intestinal barrier and ion transport dysfunction. J. Mol. Med. *93*, 535–545.

27. Guerra, E., Lattanzio, R., La Sorda, R., Dini, F., Tiboni, G.M., Piantelli, M., and Alberti, S. (2012). mTrop1/Epcam knockout mice develop congenital tufting enteropathy through dysregulation of intestinal E-cadherin/β-catenin. PLoS One *7*, e49302.

28. Lei, Z., Maeda, T., Tamura, A., Nakamura, T., Yamazaki, Y., Shiratori, H., Yashiro, K., Tsukita, S., and Hamada, H. (2012). EpCAM contributes to formation of functional tight junction in the intestinal epithelium by recruiting claudin proteins. Dev. Biol. *371*, 136–145.

29. Tsujikawa, M., Kurahashi, H., Tanaka, T., Nishida, K., Shimomura, Y., Tano, Y., and Nakamura, Y. (1999). Identification of the gene responsible for gelatinous drop-like corneal dystrophy. Nat. Genet. *21*, 420–423.

30. Takaoka, M., Nakamura, T., Ban, Y., and Kinoshita, S. (2007). Phenotypic investigation of cell junction-related proteins in gelatinous drop-like corneal dystrophy. Invest. Ophthalmol. Vis. Sci. *48*, 1095–1101.

31. Wu, C.-J., Morimura, S., and Udey, M.C. (2017). Matriptase-mediated cleavage of EpCAM destabilizes claudins and dysregulates intestinal epithelial homeostasis The Journal of Clinical Investigation. J. Clin. Invest. *127*. https://doi.org/10.1172/JCI88428.

32. Wu, C.-J., Lu, M., Feng, X., Nakato, G., and Udey, M.C. (2020). Matriptase Cleaves EpCAM and TROP2 in Keratinocytes, Destabilizing Both Proteins and Associated Claudins. Cells *9*. https://doi.org/10.3390/cells9041027.

33. Duszyc, K., Gomez, G.A., Lagendijk, A.K., Yau, M.K., Nanavati, B.N., Gliddon, B.L., Hall, T.E., Verma, S., Hogan, B.M., Pitson, S.M., et al. (2021). Mechanotransduction activates RhoA in the neighbors of apoptotic epithelial cells to engage apical extrusion. Curr. Biol. *31*, 1326-1336.e5.

34. Stephenson, R.E., Higashi, T., Erofeev, I.S., Arnold, T.R., Leda, M., Goryachev, A.B., and Miller, A.L. (2019). Rho Flares Repair Local Tight Junction Leaks. Dev. Cell *48*, 445-459.e5.

35. Bonfim-Melo, A., Noordstra, I., Gupta, S., Chan, A.H., Jones, M.J.K., Schroder, K., and Yap, A.S. (2022). Rapid lamellipodial responses by neighbor cells drive epithelial sealing in response to pyroptotic cell death. Cell Rep. *38*, 110316.

36. Aono, S., and Hirai, Y. (2008). Phosphorylation of claudin-4 is required for tight junction formation in a human keratinocyte cell line. Exp. Cell Res. *314*, 3326–3339.

37. Kitagawa, N., Inai, Y., Higuchi, Y., Iida, H., and Inai, T. (2014). Inhibition of JNK in HaCaT cells induced tight junction formation with decreased expression of cytokeratin 5, cytokeratin 17 and desmoglein 3. Histochem. Cell Biol. *142*, 389–399.

38. Shigetomi, K., Ono, Y., Inai, T., and Ikenouchi, J. (2018). Adherens junctions influence tight junction formation via changes in membrane lipid composition. J. Cell Biol. *217*, 2373–2381.

39. Friis, S., Sales, K.U., Schafer, J.M., Vogel, L.K., Kataoka, H., and Bugge, T.H. (2014). The Protease Inhibitor HAI-2, but Not HAI-1, Regulates Matriptase Activation and Shedding through Prostasin *. J. Biol. Chem. *289*, 22319–22332.

40. Chiu, Y.-L., Wu, Y.-Y., Barndt, R.B., Lin, Y.-W., Sytwo, H.-P., Cheng, A., Yang, K., Chan, K.-S., Wang, J.-K., Johnson, M.D., et al. (2022). Differential subcellular distribution renders HAI-2 a less effective protease inhibitor than HAI-1 in the control of extracellular matriptase proteolytic activity. Genes Dis. *9*, 1049–1061.

41. Larsen, B.R., Steffensen, S.D.R., Nielsen, N.V.L., Friis, S., Godiksen, S., Bornholdt, J., Soendergaard, C., Nonboe, A.W., Andersen, M.N., Poulsen, S.S., et al. (2013). Hepatocyte growth factor activator inhibitor-2 prevents shedding of matriptase. Exp. Cell Res. *319*, 918–929.

42. List, K., Bugge, T.H., and Szabo, R. (2006). Matriptase: potent proteolysis on the cell surface. Mol. Med. *12*, 1–7.

43. Benaud, C., Dickson, R.B., and Lin, C.Y. (2001). Regulation of the activity of matriptase on epithelial cell surfaces by a blood-derived factor. Eur. J. Biochem. *268*, 1439–1447.

44. Benaud, C., Oberst, M., Hobson, J.P., Spiegel, S., Dickson, R.B., and Lin, C.-Y. (2002). Sphingosine 1-Phosphate, Present in Serum-derived Lipoproteins, Activates Matriptase*. J. Biol. Chem. *277*, 10539–10546.

45. Chen, Y.-W., Wang, J.-K., Chou, F.-P., Chen, C.-Y., Rorke, E.A., Chen, L.-M., Chai, K.X., Eckert, R.L., Johnson, M.D., and Lin, C.-Y. (2010). Regulation of the Matriptase-Prostasin Cell Surface Proteolytic Cascade by Hepatocyte Growth Factor Activator Inhibitor-1 during Epidermal Differentiation. J. Biol. Chem. *285*, 31755–31762.

46. Gaymon, D.O., Barndt, R., Stires, H., Riggins, R.B., and Johnson, M.D. (2023). ROS is a master regulator of in vitro matriptase activation. PLoS One *18*, e0267492.

47. Tseng, I.-C., Xu, H., Chou, F.-P., Li, G., Vazzano, A.P., Kao, J.P.Y., Johnson, M.D., and Lin, C.-Y. (2010). Matriptase Activation, an Early Cellular Response to Acidosis. J. Biol. Chem. *285*, 3261–3270.

48. Slattum, G., McGee, K.M., and Rosenblatt, J. (2009). P115 RhoGEF and microtubules decide the direction apoptotic cells extrude from an epithelium. J. Cell Biol. *186*, 693–702.

49. Jou, T.S., Schneeberger, E.E., and Nelson, W.J. (1998). Structural and functional regulation of tight junctions by RhoA and Rac1 small GTPases. J. Cell Biol. *142*, 101–115.

50. Takaishi, K., Sasaki, T., Kotani, H., Nishioka, H., and Takai, Y. (1997). Regulation of cell-cell adhesion by rac and rho small G proteins in MDCK cells. J. Cell Biol. *139*, 1047–1059.

51. Madara, J.L. (1983). Increases in guinea pig small intestinal transepithelial resistance induced by osmotic loads are accompanied by rapid alterations in absorptive-cell tight-junction structure. J. Cell Biol. *97*, 125–136.

52. Shiomi, R., Shigetomi, K., Inai, T., Sakai, M., and Ikenouchi, J. (2015). CaMKII regulates the strength of the epithelial barrier. Sci. Rep. *5*, 13262.

53. Thirone, A.C.P., Speight, P., Zulys, M., Rotstein, O.D., Szászi, K., Pedersen, S.F., and Kapus, A. (2009). Hyperosmotic stress induces Rho/Rho kinase/LIM kinase-mediated cofilin phosphorylation in tubular cells: key role in the osmotically triggered F-actin response. Am. J. Physiol. Cell Physiol. *296*, C463-75.

54. Kim, C., Cho, Y., Kang, C.-H., Kim, M.G., Lee, H., Cho, E.-G., and Park, D. (2005). Filamin is essential for shedding of the transmembrane serine protease, epithin. EMBO Rep. *6*, 1045–1051.

55. Domoto, T., Takino, T., Guo, L., and Sato, H. (2012). Cleavage of hepatocyte growth factor activator inhibitor-1 by membrane-type MMP-1 activates matriptase. Cancer Sci. *103*, 448–454.

56. Ishikawa, T., Kimura, Y., Hirano, H., and Higashi, S. (2017). Matrix metalloproteinase-7 induces homotypic tumor cell aggregation via proteolytic cleavage of the membrane-bound Kunitz-type inhibitor HAI-1. J. Biol. Chem. *292*, 20769–20784.

57. Sakurai-Yageta, M., Recchi, C., Le Dez, G., Sibarita, J.-B., Daviet, L., Camonis, J., D'Souza-Schorey,
C., and Chavrier, P. (2008). The interaction of IQGAP1 with the exocyst complex is required for
tumor cell invasion downstream of Cdc42 and RhoA. J. Cell Biol. *181*, 985–998.

58. Rickman, O.J., Guignard, E., Chabanon, T., Bertoldi, G., Auberson, M., and Hummler, E. (2024).
Tmprss2 maintains epithelial barrier integrity and transepithelial sodium transport. Life Sci. Alliance *7*.
https://doi.org/10.26508/lsa.202302304.

59. Maetzel, D., Denzel, S., Mack, B., Canis, M., Went, P., Benk, M., Kieu, C., Papior, P., Baeuerle, P.A.,
Munz, M., et al. (2009). Nuclear signalling by tumour-associated antigen EpCAM. Nat. Cell Biol. *11*,
162–171.

60. Kong, Y., Jiang, J., Huang, Y., Liu, X., Jin, Z., Li, L., Wei, F., Liu, X., Yin, J., Zhang, Y., et al. (2022).
Narciclasine inhibits phospholipase A2 and regulates phospholipid metabolism to ameliorate
psoriasis-like dermatitis. Front. Immunol. *13*, 1094375.

61. Sumitomo, A., Siriwach, R., Thumkeo, D., Ito, K., Nakagawa, R., Tanaka, N., Tanabe, K., Watanabe,
A., Kishibe, M., Ishida-Yamamoto, A., et al. (2019). LPA Induces Keratinocyte Differentiation and
Promotes Skin Barrier Function through the LPAR1/LPAR5-RHO-ROCK-SRF Axis. J. Invest.
Dermatol. *139*, 1010–1022.

62. Pachitariu, M., and Stringer, C. (2022). Cellpose 2.0: how to train your own model. Nat. Methods *19*,
1634–1641.

63. Kubo, A., Aoki, S., and Fujita, H. (2022). Whole-mount preparation and microscopic analysis of
epidermis. Curr. Protoc. *2*, e464.

64. Tarradas, A., Selga, E., Riuró, H., Scornik, F., Brugada, R., and Vergés, M. (2013). Cell surface
protein biotinylation and analysis. Bio Protoc. *3*. https://doi.org/10.21769/bioprotoc.857.

A

TJ · old TJ · *de novo* TJ

B *WT EpH4 expressing GFP-claudin-3*

0min · 20min · 40min · 60min · 80min · 60min · 80min

X-Z · 0min · 20min · 40min · 60min · 80min

de novo TJ

C

ZO-1 · claudin-3 · Merge · X-Z

Occludin · claudin-3 · Merge · X-Z

D *WT EpH4 expressing GFP-AnillinC (active RhoA)*

0min · 5min · 10min · 15min · 20min · 25min

E

ppMLC · NMIIB · Merge · X-Z

F *WT EpH4 expressing GFP-claudin-3 and mScarlet PLCδ-PH + H1152 treatment*

GFP-claudin-3 · 0min · 60min · 80min · 100min · 200min · 300min · 400min

mScarlet PLCδPH · basal

A · B · E · C · D

G p=0.0357

Permeability (ratio to Doxorubicin(-))

control · H1152

H

Rho activation ? · *de novo* TJ

A

claudin-3

RhoA CA

F-actin

B

ZO-1   claudin-3   Merge   Scarlet-RhoA CA

occludin   claudin-3   Merge   Scarlet-RhoA CA

C

claudin-3   E-cadherin   Merge   F-actin

PBS

Rho activator

D

claudin-3 zoom

E

claudin-3   E-cadherin   Merge   F-actin

DMSO

Narciclasine

F

claudin-3 zoom

G

claudin-3

Junction enrichment
(ratio to control)

p = 0.0001
p = 0.0004

control   Rho activator   Narciclasine

H

control   Rho activator   Narciclasine

insoluble   20

soluble   20

IB: claudin-1

I

insoluble / soluble
(ratio to control)

*

control   Rho activator   Narciclasine

skin (stratified squamous epithelium)

Stratum corneum (SC)

SG1

SG2

SG3

RhoA activation

TJ

Spinous layer

claudin (+) cells

Basal layer

SG3 → SG2

Matriptase

TROP2

de novo TJ

PM

sphingomyelin / cholesterol rich domain

claudin

RhoA activation

WT

TROP2 KO

Golgi

simple epithelium

TJ

old TJ

RhoA activation

de novo TJ

WT

EpCAM KO

Golgi

Matriptase

EpCAM

de novo TJ

PM

sphingomyelin / cholesterol rich domain

claudin

RhoA activation