Redistribution of ancestral functions underlies the evolution of venom production in marine predatory snails

Giulia Zancolli[1,2*], Maria Vittoria Modica[3], Nicolas Puillandre[4], Yuri Kantor[5], Agneesh Barua[1,2], Giulia Campli[1,2], Marc Robinson-Rechavi[1,2]

[1]Department of Ecology and Evolution, University of Lausanne, 1015, Lausanne, Switzerland

[2]Swiss Institute of Bioinformatics, Lausanne, Switzerland

[3]Department of Biology and Evolution of Marine Organisms, Stazione Zoologica Anton Dohrn, 1 Via Gregorio Allegri, 00198, Roma, Italy

[4]Institut Systématique Evolution Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, 57 rue Cuvier, CP51, 75005, Paris, France

[5]Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, 33 Leninsky prospect, 119034, Moscow, Russian Federation

*Corresponding author: giulia.zancolli@gmail.com

Keywords: Neogastropoda, transcriptome evolution, ancestral reconstruction, toxin, venom, UPR, ER stress, secretion

Abstract

Venom-secreting glands are highly specialised organs evolved throughout the entire animal kingdom to synthetise and secrete toxins for predation and defence. Venom is extensively studied for its toxin components and application potential; yet, how animals become venomous remains poorly understood. Venom systems therefore offer a unique opportunity to understand the molecular mechanisms underlying functional innovation. Here, we conducted a multi-species multi-tissue comparative transcriptomics analysis of 12 marine predatory gastropods, including species with venom glands and species with homologous non-venom producing glands, to examine how specialised functions evolve through gene expression changes. We found that while the venom gland specialised for the mass production of toxins, its homologous glands retained the ancestral digestive functions. The functional divergence and specialisation of the venom gland was achieved through a redistribution of its ancestral digestive functions to other organs, specifically the oesophagus. This entailed concerted expression changes and accelerated transcriptome evolution across the entire digestive system. The increase in venom gland secretory capacity was achieved through the modulation of an ancient secretory machinery, particularly genes involved in endoplasmic reticulum stress and unfolded protein response. This study shifts the focus from the well-explored evolution of toxins to the lesser-known evolution of the organ and mechanisms responsible for venom production. As such, it contributes to elucidating the molecular mechanisms underlying organ evolution at a fine evolutionary scale, highlighting the specific events that lead to functional divergence.

Introduction

Across diverse branches of the animal kingdom, organisms have independently evolved the ability to produce and deliver venom, a cocktail of bioactive toxin molecules. In many venomous animals, these toxins are synthesised in specialised exocrine organs known as venom glands (Casewell et al. 2013; Schendel et al. 2019). While significant research has focused on the molecular evolution of toxins and venom composition, the molecular mechanisms underlying the evolution of venom-producing organs and their specialised function remains poorly understood (Zancolli and Casewell 2020). Recent genomics studies in snakes, for instance, have highlighted how regulatory networks are co-opted to drive high toxin gene expression in venom glands, notably through the involvement of trans-regulatory factors from the extracellular signal-regulated kinase (ERK) and unfolded protein response (UPR) pathways (Perry et al. 2020; Perry et al. 2022; Westfall et al. 2023). Moreover, the upregulation of UPR and endoplasmic reticulum (ER) stress pathways in the venom glands

of several distinct venomous taxa suggests that similar molecular solutions may have convergently evolved across venomous lineages to support the high secretory demands of toxin production (Perry et al. 2020; Barua and Mikheyev 2021; Zancolli et al. 2022). However, a broader understanding of how venom glands become highly specialised and optimised for the efficient mass production of toxins is still lacking, particularly outside of snake models.

Investigating homologous structures with divergent functions offers unique insights into the genetic basis of functional innovation, as demonstrated in large-scale evolutionary studies on vertebrate limbs (Zuniga 2015) and feathers (Benton et al. 2019). Among non-vertebrates, the mid-oesophageal glands of marine predatory snails in the subclass Caenogastropoda (Fig. 1) provide an excellent system to examine, at a finer evolutionary scale, the specific events that led to the emergence of new physiological functions. In some caenogastropods, like those in the family Naticidae, the oesophageal gland consist of a dilated section of the oesophageal wall involved in secreting digestive enzymes and mucus (Reid and Friesen 1980; Lobo-da-Cunha 2019). By contrast, in many caenogastropods of the order Neogastropoda, this glandular section of the oesophagus has evolved into a distinct organ known as the gland of Leiblein, which connects to the oesophagus via a duct (Modica and Holford 2010; Lobo-da-Cunha 2019). Ultrastructural observations in species from the families Muricidae and a Nassariidae suggest that the gland of Leiblein plays a role in food processing, particularly nutrient absorption and storage (Andrews and Thorogood 2005). In the superfamily Conoidea, which comprises 18 families (Abdelkrim et al. 2018) including the well-known cone snails (family Conidae) (Nguyen et al. 2023), the oesophageal gland has undergone further modifications into a long, convoluted duct that secretes a mixture of hundreds of primarily neurotoxic peptides, mainly known as 'conotoxins' (Modica and Holford 2010; Abdelkrim et al. 2018; Morales Duque et al. 2019). The venom gland is attached to a large muscular bulb that contracts to push venom through the duct and into the buccal cavity, where a modified radula injects venom into prey or predators (Modica and Holford 2010; Safavi-Hemami et al. 2010; Salisbury et al. 2010). Interestingly, some neogastropods, such as those in the family Mitridae (Harasewych 2009) and Terebridae (Holford et al. 2009), have entirely lost the mid-oesophageal gland.

The evolution of venom production from digestive-related functions in these snails represents a remarkable example of functional specialisation, shifting the gland's secretion targets from internal digestive roles (e.g., lysosomal activity) (Andrews and Thorogood 2005) to external roles that affect other organisms (receptor binding in prey) (Morales Duque et al. 2019). This transition likely provided significant adaptive advantages, enabling cone snails to diversify their diet to include fast-moving organisms, such as fish, while also offering a

defence against powerful predators (Olivera et al. 2012). However, the processes through
which an organ originally dedicated to digestive functions transformed into a specialised

toxin-producing factory remain unclear.

Figure 1: Species and tissues investigated. Overview of the anatomy of a marine snail
with the color-coded sampled tissues and the phylogeny of the species used in this study
based on Fedosov et al. (2024). The abbreviations used for the tissues and species used in
other figures are shown. The anatomy drawing is modified after (Modica and Holford 2010),
while the schematic of the foregut apparatus of caenogastropods with their mid-oesophageal
glands was modified after (Page 2011).

Here, we analyse gene expression data from the mid-oesophageal glands and other tissues
of 12 marine caenogastropod species to investigate the link between transcriptome evolution
and functional divergence. By studying the genetic underpinnings of venom gland evolution,
we aim to uncover how specific genes, pathways, and regulatory networks drive organ
specialisation to meet the distinct physiological demands of venom production. Our study
addresses the following key questions: (i) Do mid-oesophageal glands across species share
similar gene expression profiles, given their common origin? (ii) Considering the venom
gland's unique function, does its transcriptome evolve more rapidly compared to the other
mid-oesophageal glands? (iii) Which gene expression changes led to the evolution of toxin
production in the mid-oesophageal gland?
To answer these questions, we first conducted species-level analyses to characterise sets of
overexpressed genes and to delineate the functional specialisation of each gland type. We
then performed between-species comparisons to explore the gene expression dynamics that

led to the evolution of venom production. Our findings reveal that the venom gland has a
markedly distinct gene expression profile compared to its homologous organs, which are
more similar to each other. Genes encoding secreted proteins in venom glands are
expressed at exceptionally high levels, far exceeding those of other tissues. This
specialisation for toxin secretion was achieved through modulation of a conserved secretory
machinery, while ancestral digestive functions were redistributed to other organs. This shift
involved high evolutionary rates not only in the venom gland itself but across the entire
digestive system, suggesting concerted changes that underscore the adaptive flexibility of
these organisms.

129   Results

Summary of species and tissues investigated

We sampled ten species of Neogastropoda and two outgroup species within the same
subclass Caenogastropoda (Fig. 1). The two outgroup species, the Cymatidae *Monoplex*
pilearis (Linnaeus, 1758) and the Naticidae *Polinices mammilla* (Linnaeus, 1758), possess a
simple oesophageal gland (OEG) attached to the oesophagus, representing the ancestral
state. Among the Neogastropoda, two species from the Muricidae family, *Nassa serta*
(Bruguière, 1789) and *Murex tenuirostrum* (Lamarck, 1822), have a gland of Leiblein (LEG),
with *Murex tenuirostrum* also possessing dorsal glandular folds on the mid-oesophagus,
often referred to as glande framboisée (Lobo Cunha 2019). Additionally, we sampled the
gland of Leiblein in the Olividae *Oliva amethystina* (Rőding, 1798), the Vasidae *Vasum*
turbinellus (Linnaeus, 1758), and the Nassaridae *Phos senticosus* (Linnaeus, 1758). The
venomous species, possessing a venom gland (VG), belonged to the family Conidae,
including *Conus imperialis* (Linnaeus, 1758), *C. marmoreus* (Linnaeus, 1758), *C. virgo*
(Linnaeus, 1758), and *C. quercinus* (Lightfoot, 1786). We also collected tissue samples from
the Mitridae *Mitra mitra* (Linnaeus, 1758), which lacks a mid-oesophageal gland
(Harasewych 2009). Besides the glands, we sampled either the foot or the columellar
muscle, oesophagus, salivary glands, dorsal glandular folds in *M. tenuirostrum*, and
muscular venom bulb in cone snails.

Sequencing and de novo assembly statistics

A total of 150 libraries were sequenced, yielding an average of 45 million reads per library
across 12 species, with an average of three biological replicates per species (supplementary
dataset S1). We generated *de novo* assemblies by pooling all libraries within each species
and processing them through our quality-filtering pipeline (see Methods). After filtering, the
assemblies contained between 26,215 and 59,431 annotated, non-redundant protein-coding

genes, with an average of 40,192 sequences per assembly (supplementary dataset S2). The
completeness of these assemblies ranged from 82% to 93%, with an average of 89%.
Prior to data analysis, we evaluated the quality of read alignment. Samples with low mapping
rates or those that did not cluster with other samples of the same tissue type in a Principal
Component Analysis (PCA) plot were excluded (see Methods). After this quality filtering
step, we retained a total of 140 samples (supplementary fig. S1, Supplementary Material
online).
162

Within-species analysis
Characterisation of tissue-specific gene sets
To better understand the functional specialisation of the mid-oesophageal glands, we
identified sets of genes that were overexpressed in each organ. Genes were classified as
tissue-specific if their expression in a tissue was at least twice that of the second most highly
expressed tissue. On average, 14% (range: 10-19%) of genes were tissue-specific, with
43% (range: 38-51%) of these genes exclusively expressed in one tissue. Tissue specificity
was consistent across organs and species, although we observed a higher number of tissue-
specific genes in the venom glands compared to the glands of Leiblein and oesophageal
glands (Fig. 2a). The smallest set observed was the oesophagus of *M. tenurostrum* (N =
318). Notably, this species possesses dorsal glandular folds on the oesophagus (Fig. 1). The
tissue-specificity of the gene sets was further validated by differential expression analysis
using a likelihood ratio test across all tissues (supplementary text 1, Supplementary Material
online).
Gene Ontology (GO) enrichment analysis of the tissue-specific gene sets revealed marked
differences between the mid-oesophageal glands (Fig. 2b), as well as lineage-specific
patterns (supplementary text 2 and figs. S2-S10, Supplementary Material online). In OEG-
specific gene sets, we observed enrichment for terms related to intracellular trafficking and
communication (e.g., 'AP-5 adaptor complex') and transmembrane transport. Additionally,
terms related to detoxification, homeostasis, and response to external stimuli (e.g.,
'response to inorganic substance', 'superoxide dismutase activity') were enriched
(supplementary figs. S2-S4, Supplementary Material online). In contrast, gene sets specific
to the gland of Leiblein across multiple species were enriched for terms related to
intracellular digestion, particularly protein digestion and absorption, with several lysosome-
related terms (e.g., 'endopeptidase activity', 'WASH complex'), aligning with previous
ultrastructural studies (Andrews and Thorogood 2005). Terms related to iron homeostasis
and metabolism were also enriched. However, *O. amethystina* showed some unique
enrichments, including extracellular rather than intracellular compartment terms, reflecting
possible lineage-specific adaptations linked to diet (supplementary figs. S5-S7,

Supplementary Material online). In venom glands, GO terms associated with toxin and
neurotoxic activity (e.g., 'host cell postsynaptic membrane') were over-represented in all
species, as expected (Fig. 2b). Additionally, we found terms related to protein synthesis
(e.g., 'signal peptide processing', 'Golgi organisation') and to post-translational modifications
(e.g., 'hydroxylation', 'disulfide oxidoreductase activity'). Notably, terms related to ER stress
and UPR were also enriched (e.g., 'response to unfolded proteins', 'response to ER stress')
(supplementary figs. S8-S10, Supplementary Material online).
In summary, while the non-venomous mid-oesophageal glands are primarily involved in
digestion, homeostasis, absorption, and storage, the homologous venom gland has
specialised into a factory for toxin synthesis and secretion. Despite these differences, all
three gland types shared enriched terms related to iron homeostasis and metabolism (e.g.,
'iron binding') and methylation ('methionine adenosyltransferase activity', 'betaine-
homocysteine S-methyltransferase activity'). Additionally, we found terms related to hormone
response across all three gland types, including 'thyroid hormone generation' in OEG- and
LEG-specific gene sets, and 'response to thyroglobulin triiodothyronine' in VG-specific sets.

Figure 2: Overview of tissue-specific gene sets and gland secretomes. a) Number of genes within each tissue-specific set across all organs (x-axis) and species (y-axis). Species abbreviations as in Fig. 1. The venom glands have a significant higher number of tissue-specific genes (mean N = 1,424) compared to the gland of Leiblein (mean N = 771; t = 3.6, df = 6, p = 0.005) and oesophageal gland (mean N = 699; t = 5.6, df =3, p = 0.005). Missing tissue samples are marked with a "-". b) Gene Ontology (GO) enrichment results of the OEG-, LEG-, and VG-specific gene sets. c) Mean expression levels as Transcript per Million (TPM) of genes possessing a signal peptide, therefore comprising the secretomes, expressed in the organs (x-axis) of each species (y-axis). Species abbreviations as in Fig. 1. Missing tissue samples are marked with a "-". d) GO enrichment results of the oesophageal gland, gland of Leiblein, and venom gland secretomes.

Characterisation of the glands' secretomes

Given the fundamental role of secretion in the evolution of the venom gland, we analysed the 'secretome' by assessing the diversity and expression levels of genes predicted to have a signal peptide with SignalP (Teufel et al. 2022).

The number of expressed genes encoding secreted proteins was relatively consistent across
tissue types, although it was higher in the venom gland (mean N = 565.25) compared to the
gland of Leiblein (mean N = 354.25) and the oesophageal gland (mean N = 379.5)
(supplementary fig. S11a, Supplementary Material online). When focussing on tissue-
specific genes, the venom gland secretome was also more diverse than the other glands
(supplementary fig. S11b, Supplementary Material online). Interestingly, the salivary glands
did not show particularly high diversity despite being exocrine glands like the venom gland
(supplementary fig. S11, Supplementary Material online).

In terms of expression levels (mean TPM), the differences between tissues were more
pronounced (Fig. 2c). Gene expression was generally higher in venom glands, although it
varied among cone snail species (Fig. 2c). This variation could reflect genuine lineage-
specific differences, technical factors, or differences in the venom replenishment circle at the
time of sampling. However, the latter is unlikely, as individuals were sampled randomly and
thus would not all be at the same point in their replenishment cycle. Additionally, all
specimens were kept for 1-3 days in captivity prior dissection to minimise environmental
variation. We also did not observe ingested prey in any stomachs, suggesting that the last
feeding – and thus venom expulsion - did not occur close to the time of dissection for any
individual. In non-venomous species, the highest expression levels were observed in the
salivary glands. Interestingly, in the oesophagus, lower expression levels were observed in
non-venomous species (mean = 21 TPM) compared to venomous species (mean = 260; t = -
3, df = 3, p = 0.02), suggesting higher secretory activity in the latter.

As anticipated, the secretomes of the oesophageal gland and gland of Leiblein were
enriched in hydrolases and peptidases, enzymes essential for digestive processes (Fig. 2d).
Additionally, *O. amethystina* showed enrichment in 'toxin activity' (see next section), while *M.*
pilearis was enriched in terms related to communication and transport (e.g., 'metal ion
transport'). Although the salivary glands also expressed hydrolases and peptidases, these
enzymes were extracellular, whereas those in the gland of Leiblein were primarily
intracellular, consistent with enrichment in cellular compartments like 'lysosome' and
'organelle lumen". In contrast, the venom gland secretomes were dominated by toxins
released in the extracellular space and of genes involved in the ER function (Fig. 2d).

Identification and characterisation of conotoxins

Cone snail venom is composed primarily of small, disulfide-rich peptides known as
conotoxins. The number of putative conotoxin transcripts predicted in the venomous species'
assemblies was consistent with previous findings for cone snail venom gland *de novo*
transcriptomes (Abalde et al. 2018; Dutt et al. 2019), with 150-250 toxins predicted per
species. Notably, conotoxin-like sequences were also identified in other species, with

numbers ranging from 33 in *M. mitra* to 88 in *O. amethystina*. However, when restricting the

analysis to sequences with both a signal peptide and a predicted conotoxin domain, the

numbers were greatly reduced, ranging from 16 to 108 in venomous species, and 7 to 28 in

non-venomous species.

As expected, conotoxins were predominantly expressed in the venom gland (supplementary

fig. S12, Supplementary Material online). A few were also highly and specifically expressed

in the salivary glands, consistent with previous reports in other cone snail species (Fedosov

et al. 2023). In non-venomous species, predicted conotoxins were expressed at much lower

levels and across multiple tissues, although a trend towards expression in salivary glands

was observed (supplementary figs. S13-S14, Supplementary Material online).

It is important to note that the prediction tool used, ConoPrec (Kaas et al. 2012), is design to

predict only conoidean toxins. Consequently, toxins found outside Conoidea, like echotoxin

(Shiomi et al. 2002), were not predicted, leading to an underestimation of the toxic potential

of gastropod salivary glands (Ponte and Modica 2017). Our focus on conotoxins stems from

their critical role as the primary weapon of cone snails, whose massive gene expansion and

diversification likely drove the evolution of the venom gland.

Between-species analysis

Orthogroup assignment

We assigned 330,491 genes (68% of the total) to 41,720 orthogroups (OGs), with 2,588

OGs shared across all species. For comparative transcriptomics, we created a multi-species

expression matrix using the 2,588 OGs. Since many OGs contained multiple genes per

species, we selected a representative gene for each OG using two methods: 1) randomly

selecting a single transcript's TPM value, and 2) calculating the mean TPM across all the

transcripts within an OG. Both approaches produced similar outcomes, therefore the results

presented here are based on the first method. Results from the second approach are

provided in the Supplementary Material online.

Transcriptome similarity and shared tissue specificity between organs and species

To determine whether homologous glands share similar global gene expression profiles or if

their functional specialisations align them more closely with non-homologous organs, we

analysed whole transcriptome similarity patterns by means of correlation matrix and PCA.

Overall, samples primarily clustered by tissue type (Fig. 3a, supplementary figs. S15-S16,

Supplementary Material online). The oesophageal glands and glands of Leiblein grouped

together, while the venom glands clustered with the salivary glands rather than with the

homologous counterparts. Interesting patterns were observed when comparing tissue

specificity across species. We found that the oesophageal glands on average shared more

tissue-specific OGs with the glands of Leiblein (mean N = 61, sd = 17.7) and the
oesophagus of the glandless *M. mitra* (mean N. = 56, sd = 10.6) rather than with each other
(mean N. = 41, sd = 0). This suggests substantial variation between the two OEG-species,
as corroborated by the GO enrichment analysis (supplementary figs. S2-S4, Supplementary
Material online). The glands of Leiblein, in contrast, shared more tissue-specific OGs among
themselves (mean N. 103, sd = 45.9), and with the oesophagus of *M. mitra* (mean N. = 100,
sd = 38.2). In contrast, venom glands shared more tissue-specific OGs among themselves
(mean N = 111, sd = 20.4) and with the salivary glands of *M. mitra* (mean N = 71, 11.5),
while very few with the oesophageal glands (mean N = 11, sd = 3.6) and glands of Leiblein
(mean N = 16, 7.7). In summary, while the oesophageal gland and gland of Leiblein share
more similar transcriptomes, the venom gland markedly diverged from its homologous
organs and converged towards the other exocrine organ, the salivary glands
For subsequent analyses, we defined OGs as tissue-specific if they were upregulated in the
same tissue of at least two species, resulting in 41 OEG-specific OGs, 343 LEG-specific
OGs, and 405 VG-specific OGs.

313
Figure 3. Transcriptome similarity and shared tissue specificity. a) Heatmap of Pearson
correlation coefficients between tissues and species. The expression tree was made using
neighbour-joining based on the correlation matrix. The samples are colored based on the
tissue and the gland type that the species possess. b) Average number of OEG-, LEG-, and
VG-specific OGs shared with other tissue-specific OG sets, where species have been
grouped by their gland type (OEG-, LEG-, VG- species and glandless).
320
Rates of gene expression evolution

Given the marked functional divergence of the venom gland, we hypothesised that its
transcriptome evolves faster than that of its homologous glands. We tested this hypothesis
using CAGEE (Bertram et al. 2023), which employs a bounded Brownian motion model to
estimate the most likely value of the evolutionary rate parameter (σ^2) consistent with an
ultrametric species tree and observed expression values at the tip of the tree. We ran
CAGEE for the mid-oesophageal glands, salivary glands, and oesophagus, as these tissues
were sampled across most species representing all three gland types.

We evaluated four different evolutionary models (supplementary fig. S17, Supplementary
Material online). The first model estimated a single rate σ^2. The second model estimated two
distinct rates, one for the venomous clade and one for all other species. In the third model,
species were grouped by gland type and the rates were estimated for each group
separately. The final model also calculated three rates but assigned them randomly across
the phylogeny. Overall, the third model had the best fit (Table 1, supplementary table S1,
Supplementary Material online). Notably, the model with the poorest fit was the first one,
indicating that some degree of variation in evolutionary rates, even if random, is more
consistent with the data than assuming a uniform rate across all lineages. When comparing
across lineages, the venom gland showed the highest σ^2 value, supporting our hypothesis of
accelerated evolution in the venom gland relative to the other homologous glands. A similar
trend was observed in other organs, with higher evolutionary rates in venomous species
than non-venomous ones. However, when comparing across organs, none of the glands had
the highest σ^2 value, suggesting that other organs also underwent accelerated evolution,
even more so than the mid-oesophageal glands, aligning with the hypothesis of concerted
evolution across the entire digestive system.

Table 1. Evolutionary rates for the tested models.

tissue	model	N. of evolutionary rates ($\sigma2$)	likelihood (-ln L)	OEG-species $\sigma2$	LEG-species $\sigma2$	glandless-species $\sigma2$	VG-species $\sigma2$
gland	random	3	17109.7				
	1	1	17649.4	1.02	1.02	-	1.02
	2	2	16525.1	0.59	0.59	-	1.79
	3	**3**	**16018.7**	**0.23**	**0.72**	**-**	***1.80***
salivary glands	random	3	17480				
	1	1	17953.2	0.93	0.93	0.93	0.93
	2	2	15434.8	0.51	0.51	0.51	2.93
	3	**3**	**15314.9**	**0.35**	**0.58**	**0.35**	***2.86***
oesophagus	random	3	22018.2				

1	1	22728.1	1.15	1.15	1.15	1.15
2	2	21316.9	0.70	0.70	0.70	2.11
3	**3**	**21089.5**	**0.39**	**0.81**	**0.39**	*2.10*

347 For each organ is reported: the number of evolutionary rates estimated (σ^2), the likelihood,
348 and the σ^2 estimates for each group.
349

Expression changes of tissue-specific orthogroups along the phylogeny

Given the better fit of the third evolutionary model, we used this one for ancestral state
reconstruction to assess the number and direction of expression changes at each node of
the phylogenetic tree (Fig. 4b). We observed substantial changes at node 18, which leads to
the glandless *M. mitra* and the venomous clade (Fig. 4a, supplementary fig. S18,
Supplementary Material online). At this node, many OEG- and LEG-specific OGs showed a
marked decrease in expression in the ancestral gland, coupled with an increase in
expression in the oesophagus and salivary glands. Conversely, VG-specific OGs underwent
upregulation in the gland as well as across other tissues (Fig. 4a).

Among the OEG- and LEG-specific OGs, the strongest downregulation in the gland was a
galectin, which simultaneously showed the highest upregulation in the oesophagus (Fig. 4d),
alongside with a member of the ependymin family. The top three VG-specific OGs that were
upregulated in the gland included an integral membrane protein of the DAD family, a
disulfide-isomerase, and a selenoprotein F (Fig. 4d), all involved in the ER.

We then utilised the ancestral state reconstruction from CAGEE to calculate tissue specificity
at each node, akin to our approach for extant species. The goal was to understand whether
gland-specific OGs were tissue-specific in the ancestral lineages. Our findings reveal that
some OEG- and LEG-specific OGs were gland-specific in ancestral lineages until node 18,
where they shifted mainly to the oesophagus (Fig. 4c, supplementary fig. S18,
Supplementary Material online). The galectin and ependymin mentioned earlier were among
the genes that transitioned to oesophagus specificity, alongside other genes involved in
gluconeogenesis. In contrast, VG-specific OGs were primarily salivary gland-specific in early
nodes but shifted to VG-specific at node 18 (Fig. 4c). These shifts suggest that the organs of
the digestive system had to adapt to the new function, or loss of, the mid-oesophageal
gland.

Figure 4. Gene expression dynamics across the phylogeny. a) Number of OEG-, LEG, and VG-specific OGs decreasing and increasing their expression levels in the ancestral salivary glands, oesophagus, or mid-oesophageal gland at each internal node of the gastropod phylogeny. The expression changes were calculated based on gene expression reconstruction at each node of the phylogeny. b) Species phylogeny with the number of the internal nodes. c) Number of OEG-, LEG-, and VG-specific OGs which were tissue-specific in the ancestral mid-oesophageal gland, salivary glands, and oesophagus at each node of the phylogeny. d) Ancestral reconstruction of gene expression of a galectin (OG2631) and selenoprotein F (OG465) in the mid-oesophageal gland and oesophagus. The missing tissues are marked with a *.

Discussion

Understanding how organs evolve is central to studying functional innovation. Our
investigation into marine gastropod mid-oesophageal glands provides a unique perspective
on the specific events driving the emergence of specialised functions like venom production.
Venom glands in animals are highly specialised exocrine organs dedicated primarily to toxin
synthesis and secretion (Schendel et al. 2019). Consistent with this, our functional
enrichment analyses reveals that venom glands in cone snails are distinguished by
exceptionally highly upregulation of protein secretion pathways, mirroring findings in other

lineages (Perry et al. 2020; Barua and Mikheyev 2021; Zancolli et al. 2022). In contrast,
genes specifically expressed in the gland of Leiblein, and, to a lesser extent, the
oesophageal gland, are associated with intracellular digestion, particularly of proteins, as
well as uptake and storage of lipids and carbohydrates (Andrews and Thorogood 2005). The
oesophageal gland, in particular, also appears to play roles in homeostasis, transport of
fluids and nutrients, and secretion of digestive enzymes (Reid and Friesen 1980).
The phylogenetic distribution of these mid-oesophageal glands, combined with the functions
inferred from their transcriptomes suggest a scenario of concerted functional and
morphological evolution closely linked to their trophic ecology. Initially, the role of these
glands in nutrient absorption and intracellular digestion would have required prey to be either
pre-digested by salivary enzymes or naturally soft-tissues. For example, species in the
Muricidae family that drill holes into prey and consume soft tissues have a well-developed
gland of Leiblein (Andrews and Thorogood 2005), and their salivary glands open near the tip
of the proboscis, enabling immediate interaction of salivary products with prey (Kantor 1996).
As gastropods diversified their diets and some species adopted macrophagous feeding
strategies, the glands evolved accordingly, trending towards reduction or even loss. For
instance, Buccinoidea species ingest prey whole or in large chunks (Kantor 2003) and
possess a simple gland structure (Andrews and Thorogood 2005). In cases where the gland
no longer functioned in absorption, it either disappeared, as in the Mitridae, or specialised for
new functions, as seen in conoideans like cone snails. While it is often proposed that venom
evolved to enable animals to capture and consume larger prey, our findings suggest an
intriguing reversal in marine gastropods: an initial shift towards macrophagus feeding may
have paved the way for venom evolution rather than vice versa.
The functional transformation of the mid-oesophageal gland in Conoidea was facilitated by
coordinated adaptations in other digestive organs, which shifted their functions accordingly.
Our ancestral state reconstruction reveals extensive changes in gene expression across the
phylogeny, reflecting the dynamic nature of this system and the adaptation of these snails to
diverse ecological niches and feeding strategies. Notably, the venomous clade experienced
more substantial expression changes and higher evolutionary rates than non-venomous
species, consistent with the drastic functional divergence of the venom gland from its
homologous counterparts. However, the venom gland itself did not exhibit higher
evolutionary rates than the homologous glands, nor higher rates compared to other organs.
This suggests concerted evolution across the digestive systems, with other organs co-
evolving to support the gland's new function. In the ancestor of the venomous species and
the glandless *M. mitra* (node 18), several genes became downregulated in the gland while
upregulated in the oesophagus (Fig. 4a, c). Among these were genes involved in
gluconeogenesis, such as galectin and PEPCK, indicating that the ancestral gland's role in

producing energy from non-carbohydrate substrates was reassigned to other digestive
tissues. This concerted adaptation is particularly evident in the glandless species, where
many OEG- and LEG-specific genes are now oesophagus-specific (Fig. 3b), especially
those related to lysosomal functions. Mitridae snails feed on soft-bodied Sipuncula worms
which they may ingest whole (Taylor 1989; Taylor 1993), or by pumping the worm's viscera
(West 1990), including coelomic fluids and eggs (West 1991), into the buccal cavity. In such
cases, nutrients may be pre-digested by salivary enzymes, allowing the oesophagus to take
over functions previously performed by the gland of Leiblein.

During conoidean evolution, the mid-oesophageal gland lost its original digestive functions
but gained enhanced secretory capacity, primarily through the modulation of genes involved
in pre-existing secretory pathways. Our analysis identified key upregulated genes in the
ancestral venom gland (node 18) that encode proteins active in the ER, including DAD1,
which is critical for N-glycosylation and protein translocation (Zhang et al. 2016), disulfide
isomerases, and a selenoprotein F likely involved in ER protein folding quality control
(Takeda et al. 2014). Additionally, genes associated with the UPR and ER stress response
were enriched in cone snail venom glands, echoing patterns seen across other venomous
lineages (Perry et al. 2020; Zancolli et al. 2022). This finding is significant for two reasons:
first, gastropod venom glands reaffirm the trend of convergent transcriptomic evolution in
venom glands across Metazoa (Zancolli et al. 2022). Second, the upregulation of these
pathways appears unique to venom glands, as neither the extant homologous glands not the
ancestral organ show this pattern based on transcriptome reconstruction. The strong
upregulation of the UPR pathway in venom glands is notable. Even when compared to other
exocrine organs with high secretory demands, such as salivary glands (this study) or the
pancreas (Perry et al. 2020), venom glands show much higher expression levels. This
indicates that heightened UPR expression is a distinct adaptation that evolved specifically
within venom glands to support their unique physiological demands and directly controlling
toxin expression. Studies in snakes have led to a model where venom production activates
the UPR, creating a positive feedback loop that enhances venom production through up-
regulation and binding of UPR transcription factors targeting toxin genes (Perry et al. 2020;
Perry et al. 2022; Westfall et al. 2023).

In earlier nodes of the tree, OEG- and LEG-specific genes were expressed in the ancestral
gland, while VG-specific genes were found in salivary glands - a pattern still evident in
modern species (Fig. 3, Fig. 4c). This convergence between venom and salivary glands is
logical, as both are exocrine organs secreting products into the extracellular environment.
Notably, the salivary glands of some neogastropods secrete toxins to immobilise prey
(Ponte and Modica 2017), thereby expanding their function. In these cases, salivary

secretions serve both endogenous roles (e.g., pre-digestion enzymes) and exogenous
functions (e.g., toxins), indicating a functional convergence between these two non-
homologous organs. Interestingly, the salivary glands of the glandless species *M. mitra*
share several tissue-specific genes with the venom glands. Unlike other snails, *Mitra* has
salivary ducts that open at the tip of the epiproboscis, an extendible muscular rod within the
proboscis (Harasewych 2009), likely facilitating the direct delivery of secretions to prey (West
1990). We observed overexpression of cysteine-rich secreted proteins, peptidases, and
serine proteases in *Mitra*'s salivary glands, which suggests roles in tissue degradation and
potential toxin activity. Despite these functional similarities, salivary glands retained their
original digestive role and only secondarily adopted a toxin-secreting function, while the
venom gland became fully specialised solely for the latter. A plausible explanation for why
the salivary glands did not evolve into specialised venom glands lies, again, in the diet –
Mitridae snails primarily feed on Sipuncula worms, a relatively inactive and scarcely targeted
prey group. This reduced competition may have lessen the selective pressure to develop a
specialised venom apparatus, unlike cone snails that faced greater competition and adapted
to prey on fast-moving organisms that required paralysing toxins for successful capture.
486

Conclusions
This study provides new insights into the genetic basis of functional innovation by examining
the mid-oesophageal glands of marine gastropods, with a particular focus on the evolution of
venom production. Our results indicate that while mid-oesophageal glands share a common
origin, they have diverged in gene expression profiles and functions, shaped by adaptations
to different feeding strategies. In cone snails, ancestral digestive functions of the mid-
oesophageal gland were relocated to other digestive tissues in a process of concerted
evolution, enabling the venom gland to specialise exclusively in toxin production through
modulation of pathways related to secretion and cellular stress management. Overall, this
study underscores the link between transcriptome evolution and functional divergence and
identify the specific events leading to the emergence of new physiological functions.
498

Materials and Methods
Sample collection and sequencing
Between two and five individuals from Neogastropoda, and two outgroup species, were
collected in Koumac, New Caledonia, under permit N°609011- 55 /2019/DEPART/JJC.
Individuals were kept for a minimum of one day and a maximum of three days in aquaria
with fresh sea water before dissection. Salivary glands, foot, columellar muscle,
oesophagus, oesophageal gland, gland of Leiblein, dorsal glandular folds (only in *M.*

tenuirostrum), venom gland, and muscular bulb (only in cone snails) were dissected and

preserved in RNA*later* (Invitrogen) (supplementary dataset S1).

Tissue samples were homogenised with Trizol (Invitrogen) and total RNA purified with the

PureLink™ RNA Mini kit (Invitrogen) with an additional DNase I treatment following

manufacturer's protocol. cDNA libraries were constructed using the NEBnext Ultra II

Directional RNA Library Preparation Kit with polyA selection and dUTP method (New

England BioLabs) and sequenced on an Illumina NovaSeq 2x150bp. Raw paired-end reads

were checked with FastQC 0.11.9 (Andrews 2010), quality-filtered and trimmed with FastP

0.20.1 (Chen et al. 2018).

De novo assembly and annotation

All the reads from a species were pooled to generate *de novo* transcriptome assemblies

using rnaSpades 3.15.2 (Bushmanova et al. 2019). To reduce assembly size and

redundancy, and remove spurious transcripts, we adopted a series of filtering steps. First,

we translated the transcripts to amino acid sequences using Borf 1.2.1 (Signal and Kahlke

2021) and kept only those with a complete open read frame. Second, we compared the

translated sequences with BlastP (Altschul et al. 1997) against a suite of databases

(downloaded on 30.07.2021) including UniprotKB/Swiss-Prot (The UniProt Consortium

2023), a set of 11 Gastropoda genomes (supplementary table S2, Supplementary Material

online), Conoserver (Kaas et al. 2012) and Tox-Prot (Jungo et al. 2012). Protein domains

were identified with PSIblast (Altschul et al. 1997) against the Pfam (Mistry et al. 2021) and

Cdd (Wang et al. 2022) databases. Only hits with evalue < 1e-05 were retained. Additionally,

we annotated signal peptides with signalP 6.0 (Teufel et al. 2022). Putative conotoxins were

predicted with the ConoPrec tool available on ConoServer (Kaas et al. 2012). All sequences

with a UniprotKB/Swiss-Prot hit to a non-Metazoa organism were removed. We then

trimmed the retained transcripts to their coding region, reduced redundancy by clustering

sequences with 98% or more identical nucleotide sequences with CD-HIT-EST 4.6 (Li and

Godzik 2006), and kept only transcripts with more than one read count in at least one library.

Finally, we evaluated assembly completeness using Omark on the webserver

omark.omabrowser.org (Nevers et al. 2024).

For each species we performed GO annotation by combining the annotation from Pannzer2

(Törönen et al. 2018) and DeepGOPlus 1.0.2 (Kulmanov and Hoehndorf 2020). Based on

the score values distribution, we used 0.3 as a score threshold for both methods.

Within-species analysis

Expression levels

We mapped all libraries to the respective species assembly with Kallisto 0.48.0 (Bray et al.

2016) with 100 bootstrap and quantified gene expression using the package Sleuth 0.30.1

(Pimentel et al. 2017) using R 4.2.2 (R Core Team 2019). We employed a suite of quality
control steps to identify and remove outliers. First, we analysed read count distribution with
vioplot 0.4.0 and removed samples with particularly different distributions, generally with
read counts lower than the average. Then, we utilise dimensionality reduction techniques,
including PCA (*dudi_PCA*) and multidimensional scaling (*plotMDS*) to verify that samples
were clustering by tissue type. If a sample had unusual clustering, i.e., it was far from the
others of the same tissue, we excluded it. Normalised estimated counts and TPM
abundances were then re-calculated after outlier libraries' removal. As in some species
samples still tended to cluster by individual rather than tissue type, we account for the
specimen effect using an empirical Bayes method implemented via the *ComBat_seq*
function in the package sva 3.46.0 (Leek et al. 2012) which specifically targets RNA-Seq
data. We build a full model as ~ specimen + tissue_type, and a reduced model as ~
specimen.
556
Tissue-specific gene sets
For each species, we identified tissue-specific genes based on their fold change (FC)
calculated as the ratio between the TPM value of the first most-highly expressed tissue and
the TPM value of the second-most highly expressed tissue. A gene with TPM $\geq$ 2 and FC $\geq$ 2
was classified as specific of the top tissue. We validated our FC method by confirming that
the tissue-specific genes identified were also significantly differentially expressed when
analysed using the likelihood ratio test in Sleuth. Differences in the number of tissue-specific
genes between glands were tested by means of one-tailed t-tests. Functional enrichment of
tissue-specific gene sets was performed with TopGO 2.50.0 (Alexa and Rahnenfuhrer 2019)
using the *elim* algorithm and Fisher test. The foreground was the list of tissue-specific genes
while the background included all the genes expressed in that species.
568
Between-species analysis
Orthogroup expression matrix
Amino acid sequences were assigned to orthogroups (OGs) with the OrthoDB standalone
pipeline OrthoLoger 3.0.2 (Kuznetsov et al. 2023) using default parameters. Since most OGs
included more than one gene per species (i.e., one-to-many or many-to-many orthologs), we
randomly selected one representative sequence for each OG in each species and used the
TPM value estimated for that gene as the orthogroup expression value. This method was
shown to be robust (Zancolli et al. 2022). Alternatively, we calculated the mean TPM values
across all genes belonging to the same OG. All samples were then merged into a multi-
species multi-tissue matrix and the expression levels corrected for the species effect using
ComBat_seq. All the downstream analyses reported in the main text are based on the

random matrix, while the results from the mean-based matrix are reported in Supplementary

Material online.

To have an overview of global transcriptome similarity across tissues and species, we

calculated pairwise distances as 1- Spearman correlation and used it to reconstruct a gene

expression tree using the neighbour-joining method. Additionally, we calculated the

proportion of shared OGs among tissue-specific genes. For downstream analyses, OGs

were classified tissue-specific if at least one gene within that OG was specific in a tissue,

and if it was found specific in at least two species.

Gene expression evolution analysis

We analysed changes in gene expression with the program CAGEE 1.1.1 (Bertram et al.

2023) which uses Brownian motion to model gene expression across a phylogenetic tree.

The tree was derived from the phylogeny of Fedosov et al. (2024) by excluding the families

not encompassed in this study and retaining only the branches most closely related to the

species that we examined. The rates of expression changes (σ^2) were calculated for the

mid-oesophageal glands, salivary glands, and oesophagus separately. We fit a series of

nested models in which σ^2 varies across branches of the species tree to test for different

hypotheses as outlined in the results. Ancestral transcriptomes at inner nodes reconstructed

in the best fit model were used to calculate ancestral tissue specificity for each OG as we did

for extant species.

Evolution of novel genes

To test whether new genes evolved along with the venom gland, we examined the number of OGs that were found exclusively in the venomous species, as well as whether venom gland-specific OGs had, on average, a higher number of genes in venomous species compared to the other species (i.e., mean OG size in VG-species > mean OG size of all the other species groups).

Acknowledgments

This work has received funding from the European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant agreement 845674 to G.Z, and grant agreement 865101 to N.P. We are grateful to Philippe Buchet, the team of the expeditions Koumac 2.1 and 2.3 of the Muséum National d'Histoire Naturelle, Paris, and the Province Nord and Government of New Caledonia for providing the logistic support and expertise, David Massemin and Patrick Marti for their skills in finding the species needed. We thank Alexander Fedesov for providing the phylogenetic tree, Consolée Aletti for support with the RNA extraction, the personnel of the Genomic Technologies Facility and Genewiz for the RNA-seq experiment, and members of the Robinson Rechavi's lab for bioinformatics support and scientific discussions. The first version of the manuscript was greatly improved thanks to the feedback of two anonymous reviewers and additional reviewers through Qeios.

Author contributions

G.Z., M.V.M., N.P., Y.K., and M.R.-R. conceived the study design; Y.K., and M.V.M perform the dissections; G.Z. generated the RNA-seq data, perfomed data analysis, and wrote the manuscript; A.B. performed the GO annotation and G.C. the orthogroup assignment. All the authors contributed to the interpretation and discussion of the results, and to the final version of the manuscript.

Competing interests

The authors declaire no competing interests.

Data availability

The RNA-seq data generated in this study have been deposited in the NCBI SRA archive with the accession number PRJNA1158673. Additional data generated in this study are available at 10.5281/zenodo.13685166. Additional information are provided as Supplementary files.

References

Abalde S, Tenorio MJ, Afonso CML, Zardoya R. 2018. Conotoxin diversity in *Chelyconus ermineus* (Born, 1778) and the convergent origin of piscivory in the Atlantic and Indo-Pacific cones. *Genome Biology and Evolution* 10:2643–2662.

Abdelkrim J, Aznar-Cormano L, Fedosov AE, Kantor YI, Lozouet P, Phuong MA, Zaharias P, Puillandre N. 2018. Exon-capture-based phylogeny and diversification of the venomous gastropods (Neogastropoda, Conoidea). *Molecular Biology and Evolution* 35:2355–2374.

Alexa A, Rahnenfuhrer J. 2019. topGO: Enrichment analysis for gene ontology. R package version 2.38.1.

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25:3389–3402.

Andrews EB, Thorogood KE. 2005. An ultrastructural study of the gland of Leiblein of muricid and nassariid neogastropods in relation to function, with a discussion on its homologies in other caenogastropods. *Journal of Molluscan Studies* 71:269–300.

Andrews S. 2010. FastQC: A quality control tool for high throughput sequence data. Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Barua A, Mikheyev AS. 2021. An ancient, conserved gene regulatory network led to the rise of oral venom systems. *Proceedings of the National Academy of Sciences* 118:e2021311118.

Benton MJ, Dhouailly D, Jiang B, McNamara M. 2019. The early origin of feathers. *Trends in Ecology & Evolution* 34:856–869.

Bertram J, Fulton B, Tourigny JP, Peña-Garcia Y, Moyle LC, Hahn MW. 2023. CAGEE: Computational analysis of gene expression evolution. *Molecular Biology and Evolution* 40:msad106.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* 34:525–527.

Bushmanova E, Antipov D, Lapidus A, Prjibelski AD. 2019. rnaSPAdes: a *de novo*
transcriptome assembler and its application to RNA-Seq data. *GigaScience* 8:giz100.

Casewell NR, Wüster W, Vonk FJ, Harrison RA, Fry BG. 2013. Complex cocktails: the
evolutionary novelty of venoms. *Trends in Ecology & Evolution* 28:219–229.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34:i884–i890.

Dutt M, Dutertre S, Jin A-H, Lavergne V, Alewood PF, Lewis RJ. 2019. Venomics reveals
venom complexity of the piscivorous cone snail, *Conus tulipa*. *Marine Drugs* 17:71.

Fedosov A, Tucci CF, Kantor Y, Farhat S, Puillandre N. 2023. Collaborative expression:
Transcriptomics of *Conus virgo* suggests contribution of multiple secretory glands to
venom production. *J Mol Evol* 91:837–853.

Fedosov AE, Zaharias P, Lemarcis T, Modica MV, Holford M, Oliverio M, Kantor YI,
Puillandre N. 2024. Phylogenomics of Neogastropoda: The backbone hidden in the
bush. *Systematic Biology*:syae010.

Harasewych MG. 2009. Anatomy and biology of *Mitra cornea* Lamarck, 1811 (Mollusca,
Caenogastropoda, Mitridae) from the Azores. *Açoreana* 6:121–135.

Holford M, Puillandre N, Terryn Y, Cruaud C, Olivera B, Bouchet P. 2009. Evolution of the
Toxoglossa venom apparatus as inferred by molecular phylogeny of the Terebridae.
Molecular Biology and Evolution 26:15–25.

Jungo F, Bougueleret L, Xenarios I, Poux S. 2012. The UniProtKB/Swiss-Prot Tox-Prot
program: A central hub of integrated venom protein data. *Toxicon* 60:551–557.

Kaas Q, Yu R, Jin A-H, Dutertre S, Craik DJ. 2012. ConoServer: updated content,
knowledge, and discovery tools in the conopeptide database. *Nucleic Acids*
Research 40:D325–D330.

Kantor Y. 1996. Phylogeny and relationships of Neogastropoda. In: Taylor J, editor. Origin
and evolutionary radiation of the Mollusca. Oxford: Oxford University Press. p. 221–
230.

Kantor YI. 2003. Comparative anatomy of the stomach of Buccinoidea (Neogastropoda). *J*
Molluscan Stud 69:203–220.

Kulmanov M, Hoehndorf R. 2020. DeepGOPlus: improved protein function prediction from
sequence. *Bioinformatics* 36:422–429.

Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, Zdobnov EM.
2023. OrthoDB v11: annotation of orthologs in the widest sampling of organismal
diversity. *Nucleic Acids Research* 51:D445–D451.

Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. 2012. The sva package for removing
batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics 28:882–883.

Li W, Godzik A. 2006. CD-HIT: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* 22:1658–1659.

Lobo-da-Cunha A. 2019. Structure and function of the digestive system in molluscs. *Cell
Tissue Res* 377:475–503.

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto
SCE, Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families
database in 2021. *Nucleic Acids Research* 49:D412–D419.

Modica MV, Holford M. 2010. The Neogastropoda: Evolutionary innovations of predatory
marine snails with remarkable pharmacological potential. In: Pontarotti P, editor.
Evolutionary Biology – Concepts, Molecular and Morphological Evolution: 13th
Meeting 2009. Berlin, Heidelberg: Springer. p. 249–270. Available from:
https://doi.org/10.1007/978-3-642-12340-5_15

Morales Duque H, Campos Dias S, Franco OL. 2019. Structural and functional analyses of
cone snail toxins. *Marine Drugs* 17:370.

Nevers Y, Warwick Vesztrocy A, Rossier V, Train C-M, Altenhoff A, Dessimoz C, Glover NM.
2024. Quality assessment of gene repertoire annotations with OMArk. *Nat
Biotechnol*:1–10.

Nguyen LTT, Craik DJ, Kaas Q. 2023. Bibliometric review of the literature on cone snail
peptide toxins from 2000 to 2022. *Marine Drugs* 21:154.

Olivera BM, Watkins M, Bandyopadhyay P, Imperial JS, de la Cotera EPH, Aguilar MB, Vera
EL, Concepcion GP, Lluisma A. 2012. Adaptive radiation of venomous marine snail

725   lineages and the accelerated evolution of venom peptide genes. *Annals of the New*
726   *York Academy of Sciences* 1267:61–70.

727   Page LR. 2011. Metamorphic remodeling of a planktotrophic larva to produce the predatory
728   feeding system of a cone snail (Mollusca, Neogastropoda). *The Biological Bulletin*
729   221:176–188.

730   Perry BW, Gopalan SS, Pasquesi GIM, Schield DR, Westfall AK, Smith CF, Koludarov I,
731   Chippindale PT, Pellegrino MW, Chuong EB, et al. 2022. Snake venom gene
732   expression is coordinated by novel regulatory architecture and the integration of
733   multiple co-opted vertebrate pathways. *Genome Res.* 32:1–16.

734   Perry BW, Schield DR, Westfall AK, Mackessy SP, Castoe TA. 2020. Physiological demands
735   and signaling associated with snake venom production and storage illustrated by
736   transcriptional analyses of venom glands. *Scientific Reports* 10:18083.

737   Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. 2017. Differential analysis of RNA-seq
738   incorporating quantification uncertainty. *Nat Methods* 14:687–690.

739   Ponte G, Modica MV. 2017. Salivary glands in predatory mollusks: Evolutionary
740   considerations. *Frontiers in Physiology* 8:fphys.2017.00580.

741   R Core Team. 2019. R: A language and environment for statistical computing. R Foundation
742   for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.

743   Reid R, Friesen J. 1980. The digestive system of the moon snail *Polinices lewisii* (Gould,
744   1847) with emphasis on the role of the oesophageal gland. *Veliger* 23:25–34.

745   Safavi-Hemami H, Young ND, Williamson NA, Purcell AW. 2010. Proteomic interrogation of
746   venom delivery in marine cone snails: novel insights into the role of the venom bulb.
747   *J Proteome Res* 9:5610–5619.

748   Salisbury SM, Martin GG, Kier WM, Schulz JR. 2010. Venom kinematics during prey capture
749   in *Conus*: the biomechanics of a rapid injection system. *Journal of Experimental*
750   *Biology* 213:673–682.

751   Schendel V, Rash LD, Jenner RA, Undheim EAB. 2019. The diversity of venom: The
752   importance of behavior and venom system morphology in understanding its ecology
753   and evolution. *Toxins* 11:666.

Shiomi K, Kawashima Y, Mizukami M, Nagashima Y. 2002. Properties of proteinaceous
toxins in the salivary gland of the marine gastropod (*Monoplex echo*). *Toxicon*
40:563–571.

Signal B, Kahlke T. 2021. Borf: Improved ORF prediction in de-novo assembled
transcriptome annotation. :2021.04.12.439551.

Takeda Y, Seko A, Hachisu M, Daikoku S, Izumi M, Koizumi A, Fujikawa K, Kajihara Y, Ito
Y. 2014. Both isoforms of human UDP-glucose:glycoprotein glucosyltransferase are
enzymatically active. *Glycobiology* 24:344–350.

Taylor JD. 1989. The diet of coral-reef Mitridae (Gastropoda) from Guam; with a review of
other species of the family. *Journal of Natural History* 23:261–278.

Taylor JD. 1993. Dietary and anatomical specialization of mitrid gastropods (Mitridae) at
Rottnest island, Western Australia. In: Wells FE, Walker DI, Kirkman H, Lethdridge R,
editors. Proceedings of the fifth international marine biological workshop: the marine
flora and fauna of Rottnest Island, Western Australia. Perth: Western Australian
Museum. p. 583–599.

Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, Winther
O, Brunak S, von Heijne G, Nielsen H. 2022. SignalP 6.0 predicts all five types of
signal peptides using protein language models. *Nat Biotechnol* 40:1023–1025.

The UniProt Consortium. 2023. UniProt: the Universal Protein Knowledgebase in 2023.
Nucleic Acids Research 51:D523–D531.

Törönen P, Medlar A, Holm L. 2018. PANNZER2: a rapid functional annotation web server.
Nucleic Acids Res 46:W84–W88.

Wang J, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu S, Marchler GH, Song JS,
Thanki N, Yamashita RA, et al. 2022. The conserved domain database in 2023.
Nucleic Acids Res 51:D384–D388.

West TL. 1990. Feeding behavior and functional morphology of the epiproboscis of *Mitra*
idae (Mollusca: Gastropoda; Mitridae). *Bulletin of Marine Science* 46:761–779.

West TL. 1991. Functional morphology of the proboscis of *Mitra catalinae* Dall 1920
(Mollusca: Gastropoda: Mitridae), and the evolution of the mitrid epiproboscis.
Bulletin of Marine Science 48:702–718.

Westfall AK, Gopalan SS, Perry BW, Adams RH, Saviola AJ, Mackessy SP, Castoe TA.
2023. Single-cell heterogeneity in snake venom expression is hardwired by co-option
of regulators from progressively activated pathways. *Genome Biology and Evolution*
15:evad109.

Zancolli G, Casewell NR. 2020. Venom systems as models for studying the origin and
regulation of evolutionary novelties. *Molecular Biology and Evolution* 37:2777–2790.

Zancolli G, Reijnders M, Waterhouse RM, Robinson-Rechavi M. 2022. Convergent evolution
of venom gland transcriptomes across Metazoa. *PNAS* 119:e2111392119.

Zhang Y, Cui C, Lai Z-C. 2016. The defender against apoptotic cell death 1 gene is required
for tissue growth and efficient N-glycosylation in *Drosophila melanogaster*.
Developmental Biology 420:186–195.

Zuniga A. 2015. Next generation limb development and evolution: old questions, new
perspectives. *Development* 142:3810–3820.

a

Pearson correlation

1
0.5
0

tissue
species group

tissue:

■ oesophageal gland
■ gland of Leiblein
■ venom gland

■ salivary glands
■ oesophagus
■ foot
■ columellar muscle

■ dorsal glandular folds
■ muscular bulb

b

N. shared tissue-specific OGs

100
50
0

OEG-OGs

LEG-OGs

VG-OGs

OEG-species LEG-species glandless-species VG-species

a

N. OGs with -/+ changes

b

tissue:
- salivary gland
- oesophagus
} gland

OEG-OGs   LEG-OGs   VG-OGs

c

N. tissue-specific OGs

d

galectin

selenoprotein F

log2(TPM)
2   8