Additive and substitutable prey responses to feral and native predator combinations.

Justin R. Saint Juliana[1], S.S. Bleicher[2]*, S. Mukherjee[3,4], V. Sundararaj[5], J.S. Brown[5], B.P Kotler[6]

1. Cornell University, Dept. of Ecology and Evolutionary Biology, Ithaca, NY, 14853, USA
2. Loyola University New Orleans, Biology Department, 6363 St. Charles Ave, New Orleans, LA, 70118, USA.
3. Biological and Life Sciences Division, School of Arts and Sciences, Ahmedabad University, Ahmedabad, Gujarat, India
4. Ecology Environment and Climate Change Research Cluster, Ahmedabad University, Ahmedabad, Gujarat, India
5. Texas A&M University, Biological Sciences
6. Moffit Cancer Research Center, Department of Integrated Mathematical Oncology, Tampa, Florida, 33612, USA
7. Ben Gurion University of the Negev, Blaustein Institute for Desert Research, Sde Boker, Israel

*Corresponding author: Sonny S. Bleicher, sbleiche@loyno.edu , +1 (415) 670-0595

Abstract

In captive experiments of Negev Desert granivores, we investigated the ways in which combinations of
feral mesocarnivores and native predators alter wild prey behavior. We hypothesized that feral
mesocarnivores would have a greater impact on prey energy acquisition, reflected in foraging dynamics,
than native predators. Allenby's gerbils (*Gerbillus andersoni allenbyi*) and greater Egyptian gerbils
(*Gerbillus pyramidum*) were used as prey species, with feral dogs (*Canis lupus familiaris*), feral cats
(*Felis catus*), barn owls (*Tyto alba*), and horned vipers (*Cerastes gasperetti*) as predators. Gerbil
perceived risk was measured using optimal patch-use theory, with exposures to tethered predators
occurring hourly throughout the night. Some nights, two predators were alternated every other hour. We
found that human-commensal predators, particularly feral cats, induced stronger foraging than native
predators, such as barn owls. Combined predators caused gerbils to decrease foraging only when a higher-
risk predator was introduced, as indicated by higher giving-up densities (GUDs) for the dog and cat
combination compared to the dog alone, and a nonsignificant increase compared to the cat alone. The
impact of feral cats especially appears to outweigh that of native predators. This highlights the
conservation challenges to arid environments where feral cats have become ubiquitous.

Key Words: Ecology of Fear; Optimal Foraging; Predator-prey Dynamics; Global Change Ecology;

<u>Highlights</u>

• Feral predators, particularly cats and dogs, elicit stronger anti-predator responses in Negev Desert
gerbils than native predators, significantly altering foraging behavior and habitat use.
• The presence of feral predators in the Negev Desert leads to cumulative impacts on prey species,
resulting in increased vigilance and reduced resource acquisition, which suggests a disruption of
native predator-prey dynamics.
• The study's findings emphasize the unique conservation challenges posed by human-commensal
species in arid ecosystems, where limited refugia increases the predation risk from invasive and
feral species.
• This research demonstrates how predator-prey interactions in the arid communities tested are
impacted by both additive predator effects between native predators, and between feral predators,
but when combining feral predators with native ones, the effect is substitutable with feral
predators disproportionately influencing prey behavior.

Graphical Abstract

Introduction

Predators are known to impact the richness of biological communities through top-down trophic interactions [1–3]. Predator-prey interactions enrich biological communities by allowing adaptive strategies of varying prey populations to coexist in space and time [4,5]. While the original perception of predator influence on communities was predominantly consumptive [6], the ecological research community has come to accept the non-consumptive impacts of predators to be equal or greater to those of direct predation [7–9]. Overall, increases in risk, and increases in the diversity of predators promote increased diversity of anti-predator adaptations among prey, which in turn help facilitate prey coexistence [10–13]. In a period of great biodiversity loss, especially predator loss [14], the impacts of changes in predator communities are expected to reverberate down through the prey communities. While the composition of predator assemblages has impacts globally, a disproportionate impact to predation risk has always been attributed in arid ecosystems, namely due to scarcity of refuge, despite a rich biodiversity of vulnerable prey species [15–17]. In addition, the studies evaluating the structure of predator assemblages have scarcely incorporated human-commensal, feral and invasive predators as part of the same analysis.

Despite the agreement on the impacts of risk on prey diversity, measuring risk intensity in empirical experiments has been proven complicated, and quantifying prey response to said variation is harder still. The current standard for testing impact of predation risk on prey, involves captive experiments, managing titration of food and safety through the addition of predators [18], or regulating available resources in the environment [19]. Current evidence that the effects of predator abundance on prey behavior tends to be cumulative, but non-additive [20–22]. For example, gerbils exposed to one owl decreased foraging by 18%, however the addition of a second owl only decreased foraging by 5% further and a third owl reduced foraging only by 1% [23]. This may be due to a "buy one get one half off" type of situation where the value of vigilance is greater when risk is higher. In other words, prey being vigilant for one owl, will also detect a 2^{nd} or 3^{rd} one. This type of benefit may apply to similar predators, but not predators that are very different. In contrast, with some predators, for example foxes, desert rodents tend to only "tune in" when the level reaches a certain threshold[24]..

While titration of risk is a clear influencer of risk accumulation, rarely is it ecologically relevant to observe a single predator species as the only source of risk in an environment. Multi-predator impacts have been predominantly studies in consumptive studies in aquatic and arachnid model communities, and have shown to take two major forms: facilitation or interference between the two predators [25,26].

Behaviorally, the impact of predators on each other depends on how prey perceive risk from these predators, thus impacting their habitat choices and the amount of time they spend foraging. Predator facilitation benefits predators in two major ways: 1. One predator drives prey towards the habitat preferred by another predator [25,27,28] in a mutualistic benefit. 2. Presence of one predator overpowers the attention given to another predator thus making the prey more vulnerable to the other predator that is less focused on, *i.e.*, a commensal relationship between the predators [29]. For many prey species, the risk from one predator requires much greater vigilance, or other types of defensive behavior, and thus marginalizes the attention (vigilance behavior) and energy expenditure (as decerned through time foraging) that can be allocated to avoidance of the lower risk predator [30,31]. It is important to be careful when interpreting the latter case when researching predation based on prey behavior. The predator that has a lower impact on the prey's fitness may seem to cause no risk, because it is masked by the predator that the prey perceives to be riskier.

Predator facilitation should narrow the availability of perceived safe habitat and narrow the realized niche that the prey can occupy. This logically changes both community interactions, and the

populations dynamics of both predators and prey. The lower the abundance and distribution of prey, the
less fit both predators become in the long run. For example, the desert pocket mouse (*Cheatodipus*
penicillatus), from the Mojave Desert, favors a strategy of torpor and reduction of foraging to a minimal
sustainable level in response to combinations of owls and snakes in their environment [32]. When this
occurs, neither predator is able to hunt effectively, despite the scenario suggesting facilitation. On the flip
side, when predators interfere with each other, the prey should effectively be able to use more resources
as a result of the conflict between the predators, reducing the overall risk [33,34]. This interaction is
typically observed when the two predators use the same habitat, and drive the prey to use habitat that is
less optimal for both predators. For example, when a population of gerbils is exposed to both owls and
foxes, they prefer to forage under the safety of bushes where neither predator has easy access to them
[35].

Different prey species will respond to the interaction of predators in different ways. This is a
consequence of evolutionary changes driven both by predator avoidance [36,37] and by intraguild
competition strategies pushing prey populations to evolve strategies that balance both competition and
predation costs [30–32,38–41]. Specifically, the foraging strategy of competing prey species will alter
how susceptible they may be to predation risk [42,43]. Examples include locomotion type, with saltatory
motion conveying advantages for predator avoidance [44,45], sensory prowess as in the example of inner
ear bullae allowing for extra sensitive hearing and early escape [46], and even unique special adaptations
such as torpor as in the example of the abovementioned pocket mice [32]. In this experiment, we test the
responses of two competing gerbil species from the Negev Desert in Israel, *Gerbillus andersoni allenbyi*
(GA) and *Gerbillus pyramidum* (GP). We expect that the same experiments would result in different
outcomes from species applying different anti-predator strategies. Specifically, the two gerbil species
differ in their size with GA being significantly smaller than GP [47]. The size difference between those
two well-studied model species has given rise to significant behavioral differences both in intra specific
competition strategies and management of predation risk. The variation between the species, interspecific
competitive dynamics between them, and intraspecific competition within the populations combined with
the richness of the environment should also alter the impact of combined predators. We expect the
addition of predators to have a greater impact when the competition for resources is lower (*cf.* Makin and
Kotler 2020).

This study aims to compare the response of two desert gerbil species to titrations of predation risk
through the exposure to four different predators: two human-commensal species (a feral cat and a dog), a
barn owl, and an olfactory cue of a horned viper in the form of musk imbued sand. We intend to test each
predator alone and in combination with each of the other predators to assess their cumulative and additive
impacts on prey behavior, or whether their impacts are substitutable.

Given excessive impacts of human-commensal predators on small mammal communities
worldwide [49,50] we predict a potentially disproportionate assessment of the risk these predators pose
compared with native predators. The directionality of disproportionality may increase perceived risk, if
the prey species understand the risks they pose, but may just as likely be significantly lower if the prey are
naïve to the risks they pose [51]. Despite expected differences between the prey species' behavior, we
predict that the general response to predator stressors will follow similar patterns, as their anti-predator
evasion strategies are similar enough. We predicted that the combination of any two predators would
result in additive responses of the behavior of the gerbils. Additionally, we predicted that the closer the
hunting strategy of any two predators, the lower the added impact will be on gerbil apprehension and
foraging. *i.e.* adjustment in behavior applied towards one predator would automatically apply to the
similar predator. For example, the response of gerbils to a fox would be to move the majority of their

foraging under the safety of bushes. The logical response to a jackal in the environment would be very
similar, therefore the response would not vary much when both are present. However, when avoiding
snakes that ambush under bush cover, foraging in the open is the logical safe response [33,48]. Therefore,
combing a fox and a snake would increase the risk throughout the environment, thus registering as a much
greater threat than two ambush predators, or two active searching predators. (Table 1). Where the risk
from one predator type outweighs the risk from another, the effect may register as overshadowing of the
impact of the lower-risk predator [30].

Methods

Species

For this experiment we used two common gerbils from the Negev Desert of Israel: the greater
Egyptian gerbil *Gerbillus pyramidum* (GP), 40 g, and Allenby's gerbil *Gerbillus andersoni allenbyi*
(GA), 24 g [52]. Both these gerbils are nocturnal, are commonly found on sandy substrates such as sand
dunes, and are considered granivores, despite foraging on available insects, and green vegetation when
available [53]. Both gerbils have adaptations to reduce the risk of predation, including saltatorial
locomotion for enhanced escape abilities and auditory adaptations to increase hearing acuity [31,54,55].
However, the species are known to vary in their response to risk, with the larger species being more
responsive to changes in both patch quality and risk [41,56]. Similarly, the foraging patterns differ
between the species, the larger applies a cream-skimming pattern, only collecting easily found grain in
risky conditions, and the smaller applies a crumb-picking strategy in which they dig and search for seeds
more thoroughly [42].

The gerbils were collected in the Mashabim Dunes (N 31_0'14.531'', E 34_44'47.31''), some
30km from the experimental enclosure. The barn owls were donated to the lab from a rescue center. For
this experiment we used *Cerastes gasperetti* snakes that were trapped in the vicinity of the experimental
enclosure, and housed in the zoological collection at Midreshet Ben-Gurion, Israel. All the animals were
held in climate-controlled animal-rooms at the Blaustein Institutes for Desert Research within 300 m of
the experimental vivarium. Both the feral dog (*Canis lupus familiaris*) and the feral cat (*Felis catus*) were
trapped (adopted) in the Sde Boker environment and were loaned for the experiment by their owners.

Experimental Set-Up

Animals:

The measurements were collected for each species in the absence of direct interspecific
competition. This allowed us to make a comparative study of the effects of predation risk in the exact
same setup in the absence of competition stressors. To best allow for comparison across gerbil species, we
populated the vivarium with roughly the same biomass of rodents corrected for metabolic rate, leading to:
24 Allenby's gerbils, 16 Greater Egyptian gerbils. In each of the experiments we used a 1:1 mix of adult
males and females to avoid sex-based, and age-based biases and a stage structured predator prey dynamic
[57]. We tracked depredation using subcutaneous RFID PIT tags which were identified by scanning the
owl pellets with a reader. All other predators were muzzled to avoid physical harm to the gerbil.
Depredated individuals were replaced on the following night.

Dates:

Each experiment was conducted for 100 days starting April 20[th] 2004 with GA being on one side
of the vivarium and GP on the other.

The Enclosure

We conducted our experiments in a semi-natural, outdoor enclosure (vivarium 17 X 34 m)
located on the Sede Boker Campus of Ben-Gurion University, Blaustein Institutes for Desert Research,
Midreshet Ben-Gurion, Israel (N 30 51' 25.978'', E 34 46' 51.284''). The vivarium was divided into two
halves. It is covered with a wire net 4.5 m high such that aerial predators can be flown within. Our
vivarium mimics a dune habitat with a layer of 10 cm of sand covering a loess-clay subflooring with
escape buried to 1 m below ground. To replicate the sparse vegetation cover in semi-stabilized desert
dunes we created heterogeneity by adding 16 trellises in a randomized array with at least 2 m separation
from the next trellis. Each trellis (80 X 50 X 15 cm) at each station was topped with a pile of cut brush to
create a spherical ''bush'' approximately 1m in diameter. The environment beneath each trellis provides a
sheltered environment mimicking a bush microhabitat, while the space between trellises replicates the
open space occurring between vegetation clumps in the gerbils' natural environment.

Data Collection

We measured gerbil foraging using a randomized grid of 32 patches sieved nightly to obtain a
GUD measurement. Each patch (38 X 28 X 8 cm) held 3 liters of sand and was stocked each evening with
3 g of millet. At sunrise, each patch was sieved and remaining seeds weighed to 1/100 g and logged for
analysis. Daily, patches that were set up as covered patches (bush) were set outside the trellis for the
following collection night as open, and vice versa. The data sampling in this experiment combined two
related experiments, which used the same predators. This manuscript is a reflection of a meaningful
subset of data collected. However, this resulted in an uneven number of repetitions for each predator
treatment we used.

For each of the gerbil species, in accordance with each of the predator treatments and
combinations of predators, two nights were sampled for each moonphase. This resulted for 8 nights of
sampling for every predator alone and 4 nights for every predator combination. Additionally, 8 control
nights were sampled for each rodent species in accordance with the same moonphase regimen. Because of
the experimental set-up, randomization, and removal of additional data we deemed more effective to
analyze separately, three of the predator combinations (cat-owl, cat-snake, and owl-snake) were run for an
additional 2 nights. We randomized the predator sequence so that no two treatments would follow each
other on two consequent moonphases. This means that the control nights were interspersed between other
treatment nights. On nights when a single predator was used, the predator would be introduced to the
vivarium once every 60 minutes from sunset to sunrise. For the snake treatment, sand imbued with the
natural musk of the snakes from their enclosures was spread around the foraging stations on the same
schedule. Based on the lunar calendar we marked the moonphase associated with the night of collection.
Of the 64 patches, 32 were set as "bottom" trays and 32 as "full" trays. For bottom trays, we mixed the
seeds into half of the sand, spread that sand over the tray, and then placed the remaining sand without
seeds on top. This created a more complex tray in which the seeds were in higher concentrations, but
harder to access. For full trays, we simply mixed all the seeds into all of the sand and spread that across
the tray. Such setups allow us to gauge apprehension by comparing changes in selectivity [58,59].
Moonlight ambient illumination and the temperature were recorded using digital meters.

Data Availability

The data used for this experiment available at Mendeley Data Repository [60]

Data analyses

We separated the cumulative data collected in the experiment into two data sets: single predator
data (including the control) and multi-predator data (using only nights with predator encounters every 60
minutes). We then ran analyses on the multi-species data set both as a single unit, and separately for each
of the gerbil species. Figures were generated using the open source Jamovi project [61], running on R ,
while the statistical analyses were conducted in Systat13.

<u>Single predator</u>

We ran a series of t-tests for both gerbil species combined, and for each apart. These tests
compared whether the gerbils responded to the risk associated with each predator species as unique and
different from the same environment in absence of predators.

<u>Multi-predator</u>

To address differences between predators, we ran a General Linear Model (GLM) analysis for the
data collected from the experiment using both gerbil species. We increased the explanatory value of the
models, by adding interactions shown significant in a multi-factorial ANOVA that had very low
explanatory power, and made sure to include all biologically meaningful interactions. The factors we used
were gerbil species (GS), predator treatment (PT- type of predator present on the night of), moonphase,
resource distribution (RD; clumped or dispersed resources – at the bottom of a tray or mixed throughout),
and microhabitat (open and bush). For all variables and interactions found statistically significant in the
model (p-values below 0.05) we ran post-hoc Tukey Tests of Honest Significance Difference (HSD).

To ease in the interpretation of post-hoc analyses, we re-ran multi-factorial ANOVAs for each of
the gerbil species separately. Both these models used predator treatment, moonphase, Resource
distribution and microhabitat as independent variables and added covariates of ambient temperature and
moonlight intensity. Post-hoc Tukey HSD pairwise comparisons were calculated for all significant factors
and interactions.

<u>Interpretation of data</u>

To measure the cumulative responses of various predators, we must interpret the effects of the
predators on each other, through the behavior of their prey as our experiments minimize the actual
depredation our predators were allowed to perform (through restricting the predators). We use quitting
harvest rates of the gerbils as gauged by giving-up densities (GUDs) in artificial patches to measure
perceived risk [62,63]. We attributed patches with greater foraging activity (lower GUDs) as areas
perceived as safer, and patches with higher GUDs as areas perceived as risky in accordance with previous
publications[63]. The comparison of the mean GUDs for a single predator with the combination including
another predator will allow us to assess the impacts of the predators on each other. By calculating a
change in foraging dynamics dGUD (GUD of single – GUD of combined predators) we can expect a few
patterns (Table 2). Because of predator microhabitat preference, our prediction is that facilitation will
occur between the snake treatment and any of the other predators. This will result in a negative dGUD as
the overall perceived risk in the environment will be higher when the two predators are present.
Additionally, because we used musk in lieu of free-ranging snakes, it is likely that the response to the risk
from active predators will swamp the risk from the snakes entirely. Last, if the predators facilitate each
other, then the overall dGUD should be zero or positive, as the foraging in the entire experimental arena
should not decline as it simply equalizes between the microhabitats [33].

<u>Predator housing and control:</u>

Efforts were made to reduce predator strikes on gerbils: owls were flown tethered to a rope and
were fed prior to the experiment, the muzzled dog ran free in the enclosure, the cat was on a harness and
tether but allowed to roam freely, the tether was used to prevent the cat from making fatal strikes. As
snakes were hard to control, sand imbued with their musk from their terraria was used as an olfactory cue
instead.

Results:

In a simple t-test, all predators were perceived as giving rise to greater risks than the control, with
exception to risk posed by owls. Allenby's gerbils (GA) categorically associated the risk from all
treatments as greater than the control, whereas the Greater Egyptian Sand Gerbils (GP) did not
differentiate the snake treatment from the control and associated the owl treatment as lower risk than the
control (Fig 1, Table 3). None of the treatments were perceived as a very high risk, as even the highest
giving-up densities (means) were at 1.71g±0.03g, meaning that more than one third of resources were
consumed in patches on a nightly average. While low, the variation seen here does not stray from
published observations, and is in line with natural occurring intra-species variation.

The GLM comparing the data collected for both gerbil species resulted in three significant single-
factors and seven two-way interactions (Table 5). The two more multi-factorial ANOVAs for GA and GP
separately found predator treatment, moonphase and microhabitat, and both covariates significant as well
as the interactions of PT X moonphase and RD and moonphase. The three-way interaction yielded
significant but hard to interpret results (SI 1,2). GA decreased foraging with increased temperature, while
GP increased foraging as temperatures rose. No clear discernible patterns emerge for the moon
illumination (SI 3). Regression analyses were non-significant and relatively low correlations were
observed with Pearson Correlation Indices for moonlight illumination were -0.1 and 0.0 for GP and GA
respectively. For temperature the correlation indices were 0.034 and 0.08 for GP and GA respectively.

GP foraged to lower GUDs than GA, with means being 1.694±0.02 and 1.489±0.02 g
respectively. Both gerbils perceived the bush microhabitat as safer (lower GUDs). The means were
1.465±0.016 and 1.718±0.017 g for bush and open microhabitats respectively. Gerbil response to
moonphase was not uniform for both species. GA foraged less on the full and waxing periods of the
month, and had lower GUDs on the waning and new moons. The mean GUDs were 1.709±0.033,
1.751±0.030, 1.735±0.033, 1.521±0.033 g for the new, waxing, full, and waning moonphases
respectively. GP foraged the least on the new and waning moonphases with GUDs being 1.521±0.033,
1.485±0.030, 1.418±0.033, 1.553±0.033 g for the new, waxing, full, and waning moonphases
respectively.

Predator type (PT) yielded significant patterns in the two gerbils combined dataset, and a less
impactful relationship for each separately (Fig 2). Tukey HSD tests revealed that GA response to predator
treatments was a lot less definitive than with GP (Figs 2B, and 2C respectively, SI 4,5). Neither the RD,
nor the interaction of moonphase with PT yielded few clear patterns of biological significance.

Overall, the impact of the snake treatment was negligible for both species suggesting that the
gerbils were likely ignoring the olfactory cues in presence of all live predators except for the owl (Fig 3).
For GA, the exposure to the dog resulted in a cumulative trend that was not statistically significant. The
owl treatment was always replaced by the risk of the dog, leading to higher GUDs when the dog was
present. The risk from the dog and cat treatments overall caused the largest decrease in foraging from all
the predators, but were comparable to each other. The dog treatments resulted in mean GUDs of
1.61±0.037, 1.681±0.052, 1.539±0.052 g for both gerbils, GA and GP respectively. The cat, resulted in
GUDs of 1.685±0.037, 1.720±0.052 and 1.650±0.052 g respectively. To better understand the predator

interactions however, we examined the interaction of predator treatment and microhabitat, but only for
both species combined, as the three-way interaction did not prove to be significant (Table 4, Fig 3).

The two ground foraging predators seemed to have elicited the largest "fear" response in the
gerbils, corresponding with highest mean GUDs. Between the two predators, the cat was perceived as a
much greater threat than the dog (Fig 3 A&B). The change in foraging in the open microhabitat decreased
by 10.8% of the total available food from the dog treatment to the dog and cat combined. The decrease
was only 4.6% from the cat to the cat + dog combination. Neither the snake treatment alone, nor owl
treatment were substantially different from the control in this experiment, in neither the bush nor the open
microhabitat (Fig 3 C&D). For the owl, the open microhabitat became significantly riskier only in
combination with the dog and the snake, but not with the cat (an anomalous result). Last, the snake
treatment resulted in very little differences from the control. However, when combined with the dog, both
microhabitats were equally risky pointing to facilitation between the two predators.

Discussion

This experiment highlights and strengthens a few previous finds, namely that not all predators
exude the same intensity of predation risk on foragers [29,64,65], and that different forager species- based
on their unique adaptive strategies- do not respond equally to the same risks [66,67]. Meanwhile, our data
also supports observation made by numerous authors that the suggest that intensity of a predator cue
affects the response by the prey [68], and that the presence of various predators simultaneously have
cumulative impacts and not substitutive effects. However, the perceived risk that a predator exudes on the
prey will dictate whether the effect of the additional risk factor is negligible or cumulative.

Foraging strategy matters

Stronger interference competitors that employ a cream-skimming foraging strategy appear to be
more attuned to the specifics of risk in their environment. GPs were much more specific in their
management of light (moonphase) and predator activity than the GA were. This suggests that when
foragers have the ability to exclude their competition, they will use resources in a more discerning manner
than a lesser competitor [39]. A number of hypotheses have been generated over time that explain why
the larger competitor tends to "give-up" sooner than the smaller one. The leading ones include that the
larger prey have greater value to the predator and thus draw greater   focus from the predator [69];
another is that larger foragers have higher energetic foraging costs due to the larger metabolic costs
associated with greater body mass [20]. Regardless of explanation, the evidence collected herein supports
this observation. The consequence of the sensitivity of the larger competitor is that it "opens up" time,
space, and resources for the lesser competitor who is less discerning [70,71].

The observations relating to the smaller competitor, GA, suggest a different pattern of behavior
than its larger counterpart. In congruence with former observations, the GA accept what they must as the
weaker competitor. They take more risks [20], utilize suboptimal times [67] and suboptimal substrates
[53,71]. Last, they forage in patches that have much lower energetic value (Kotler et al. 1993b, Ben-
Natan et al. 2004). The two major observations supporting this dynamic in our data were a hyper focus on
the greatest risk predators in the experiment, where they foraged more, and took risks with all predators
except the highest risk (dog or cat). Additionally, the support of the suboptimal times, the GA preferred
the "in-between" times when the moon was waning and waxing especially at times that the risk was
greatest [72].

Predator treatments

The patterns of facilitation, interference and cumulative risks derived from the combinations of
predators significantly deviated from our expectations. The response to olfactory cues were smaller than
the responses to live predators, and even a live predator did not always differ significantly from the
control treatment (as seen in the example of the owl). Below we aim to break down the possible drivers of
the interactions we observed.

Predation risk administration

It is not surprising that a predator cue in absence of the actual predator, was perceived as a lower
risk than live predators. Many studies have tested the use of olfactory cues of predators as deterrents to
control agricultural damage and found conflicting outcomes. This scenario, where the gerbils
underestimate the risk from the snake musk, also aligns with a significant body of literature that suggests
that olfactory cues result in less impactful results from prey. A couple of examples include Tasmanian
wallabies having little response to dingo urine [73] and squirrels in Illinois not responding to red fox urine
[74,75]. Moreover, the debate about using olfactory cues to simulate predation has been ongoing for
decades. Banks et al. [76,77] have shown that the response to olfactory cues can vary significantly,
especially when the source of the cue is not fresh. They also argue that these types of cues are used in a
subjective manner by the foragers, for example bush rats in their experiments had less response to old
urine when exposed to fresh urine on the same night. Olfactory cues have also been shown to be
interpreted by prey differently based on ambient temperature and air humidity [78]. Lastly, foragers have
even been found to use olfactory cues, of both predators and competitors, as camouflage from live
predators[68].

We mention all these challenges to raise the caveat that we interpret our results to suggest that the
gerbil responses to the olfactory cues for the presence of snakes, based on musk-imbued sand, do not
support the view that gerbils would interpret the scent as a sign of predation risk. However, the gerbils
did show an interaction between these cues and the presence of other predators, namely owl and dog. In
addition, the responses were much greater for the GPs than for the GA, evidence supporting the above
claims on variation between the species and their competitive foraging strategies.

Hunting strategy matters and impacts prey's risk perception.

Foraging has been shown to decrease based on various predator related characteristics, namely
the lethality of the predator, but potentially more importantly, based on perceived risk which at times may
veer from the actual lethality of the predator [79,80]. Thus, as a generalization, prey often overestimate
the risk posed by a predator, as the cost of under-estimating the risk is significantly greater than the
opposite [81–84]. Additionally, the information a prey species has of their predator's behavior, innate or
learned, will change the behavioral tactics they will employ to avoid that risk. While the most commonly
reported is a shift in microhabitat use, the entire spatial and temporal distribution may change based on
the type of predator the foragers interact with. For example, desert rodents from multiple continents have
been shown to reassess risk distribution, sometimes referred to as a landscape of fear, and refocused their
anti-predator strategies to the predator they perceived as more lethal. In that example the greater risk was
provided by an owl that overshadowed live snakes [65].

Our observations in this experiment suggest a general agreement between the two foragers. Both
the gerbils foraged the least in the presence of the active ground hunting predators, followed by the owl
and the snake musk having a significantly lower impact on their behavior. Similarly, the cat was
perceived as a greater risk than the dog as evident by the significant decrease in foraging when the
combination of the cat and dog are compared to the isolated dog treatment. However, when comparing
the combined predators with an isolated cat, the pattern is cumulative but not significant.

<u>Predator facilitation, interference, and cumulative effects</u>.

The interactions between two predator treatments overwhelmingly showed cumulative effects.
Where the interaction was not significant, this can be interpreted as additive interactions. Where the
interaction was less than additive (significant positive dGUD) we can interpret the interaction as
interference. When the interaction was more than additive (significant negative dGUD) we interpret the
result to be facilitation. The gerbils of both species foraged less in combinations of predators over the
control and over each predator alone. However, the specific interactions do shed light on a couple of
patterns. Where interference between predators was observed, as measured through increased foraging in
the combined treatment, it was never statistically significant, and thus the interpretation results in a
conclusion that these are all additive interactions. Additive patterns were more likely in the GP population
than in GA and were more likely in predators of lower lethality (owl and snake) and especially when
these predators hunt using different strategies as in the snake and the cat (present in both species). A
plausible explanation for why these appear to exist is that different parts of the enclosure are associated
with the risk of specific predators. For example, both Merriam's kangaroo rats (*Dipodomys merriami*) and
GAs have been shown to identify areas of risk of predators they prioritize [30] and change their
geographic distribution within experimental enclosures in response. The kangaroo rats had identified the
locations where snakes were present and depleted the available forage in other locations, meanwhile GAs
avoided a transact of land corresponding to the flight path between two perches favored by the barn owls
but were shown to be willing to risk the physical space where snakes were observed. Having the
combination of multiple predators here, drives the gerbils to search for resources in environment they did
not previously forage, and thus increasing their harvest intake to compensate for increased movement
activity [48].

The interplay of predator hunting strategies was expected to yield more significant habitat
selections in the gerbils than we observed. The clearest of textbook examples for predator facilitation is
that of an active-searching predator, like an owl, pushing the foragers towards ambush predators awaiting
them under bushes. The presence of ambush predators alternatively encourages the foragers to avoid the
refuge for the exposed environment where the owls can pick them off [33]. As mentioned above, our
ambush predator treatment, in the form of snake musk, was not as effective as we had planned and
resulted in the bush habitat being safer in every interaction where an owl was present. The dog treatment
overall interplayed similarly to that owl, as a larger mesocarnivores that would predominantly impose risk
in the open spaces and could not fit under the trellises. The cat on the other hand equalized the risk in both
microhabitats, especially when interacting with the other active searching predators. This may also
explain the overall greater impact that are perceived by feral cats, and their known impact to small
vertebrate biodiversity in arid zones [85].

Despite our expectation, the theory of predator facilitation did hold up when we look at the
microhabitat use in this experiment. This perhaps is a result of the lack of significance of the three-way
interaction of gerbil species, predator treatment and microhabitat. Therefore, we were only observing
interactions of the combined data set for both species. Here, the only example of facilitation was not
between the cat and snake as we had initially expected, but between the dog and the snake. In that
treatment combination the gerbils equalized the resource use between the microhabitats. To our best
interpretation, this comes from the general behavioral pattern both gerbil species were observed to follow
where they shift their focus to the greatest risk in the environment [30]. While all the other interactions
appear to be facilitative and cumulative with the foraging decreasing in the open microhabitat only, it
would be very difficult to tease out which of the two each interaction represents. However, the same
pattern of shift of attention leads us to assume that most of the interactions are cumulative.

The anthropogenic human-commensal predator, conservation implications

Our use of feral, human commensal species of predators allows us to comment on an unexpected
aspect of conservation importance. Significant numbers of papers focus on the impact that feral cats
specifically [85–88] and human commensal predators in general [89–91] have on biological communities
globally. Our observations were both encouraging and support the overall fear of the exploding
populations of feral cats, a politically explosive topic in arid zones, such as Australia [92], the Middle
East and North Africa [93] and on arid islands [94]. The encouragement we observe comes from the fact
that the gerbils clearly understood the cat to be the greatest threat to them. In addition, the gerbils
interpreted the risk from the cats and dogs correctly and responded by shifting their microhabitat
preference. Thus the overall impact these feral predators may have on their populations is curbed in
comparison to the impacts they have on naïve prey on islands and the Australian continent [95,96].

In addition, the impact the dog had on the foraging of the gerbils surprised us. Predominantly,
aerial predators, and specifically barn owls, tend to garner the most anti-predator focus in gerbils [35]. It
is unusual to find ground dwelling, active searching predator, as the main risk factor for desert granivores.
The fact that the dog, despite being perceived as a lower risk than the cat, still resulted in greater
apprehension by the gerbils, suggests that human commensal predators have a significant potential to
impact community-wide interactions around the globe.

Community-wide impacts of feral cats and dogs have predominantly focused on consumptive and
injury-based effects, with strong emphasis on the naiveté of prey [49,51,94,95,97]. Studies on feral cat
impacts far outnumber those on feral dogs[98]. The study of community-wide impacts of feral dogs has
been mostly limited to the role of dingoes replacing apex predators in Australian arid zones. Two major
factors differentiate this case from the present studies: first, dingoes act as apex predators in Australia,
while in our study, feral dogs functioned as meso-carnivores within Negev and Saharan Desert
ecosystems; second, management efforts for dingoes—and therefore research on them—far exceed those
for feral dogs in other locations. Meanwhile, reviews of prey responses to dingoes, albeit focused on
larger prey species, indicate that responses vary based on cue type and recent exposure to the predator. In
other feral dog populations, most research has focused solely on consumptive and injury-based impacts.

However, recent evidence challenges the assumption that native prey are naïve to introduced
predators[98,99].While previously attributed biodiversity demise in arid zones was attributed to naïveté of
the prey, more recent studies show that small mammals in arid regions are rapidly adapting to recognize
the presence of feral and invasive predators. The exception lies in feral cats [98]. Our observations here
stand at odds with these observations from the Australian studies. The gerbils clearly acknowledged the
risk posed by the cats in their behavioral choices. We offer two interpretive reasons for this difference.
First domestic cats are believed to have been domesticated from an ancestral species *Felis libyaca* in this
region of the world[100], suggesting a much longer period of habituation with both the feral dogs and
cats. Second, we expect a certain evolved recognition given the diversity of native felid and canine
mesocarnivores in the habitat where the gerbils were collected, namely jackals, sand foxes, red foxes,
caracals and sand cats.

While not groundbreaking, our observations have significant implications for conservation and
wildlife management. Predominantly, we can acknowledge the need for strategies beyond predator
removal, which is economically and ecology costly and perceived as a controversial practice by the
general public [101]. However, this raises the need to manage wildlife by enhancing natural antipredator
behaviors, potentially through controlled exposure to predator cues, which should reduce the long-term
impacts of the feral predators. Additionally, habitat modifications that increase cover and refuge

availability can mitigate predation risk, and reduce consumptive impacts. This would support small mammal populations in arid environments where exposure to predators increases in movement between shelters [15]. By integrating these approaches with traditional predator control measures, conservation efforts can better safeguard biodiversity in landscapes where human-commensal predators exert disproportionate influence.

Conclusion

The experiment was designed to tease out the interactions of various predator types on the foraging behavior of small, wild, desert mammals. Through the behavior of the gerbils, the facilitation and interference between the predators was tested. The interactions between predators proved to be predominantly cumulative. Facilitation was found to be indisputable between lower-lethality snake olfactory cues and the dog. The impacts of ground foraging, human-commensal species, was shown to draw the attention of the gerbils over predators that naturally occur in their environment, such as a barn owl and a horned viper. This also supports a tendency of habituation to these predators as integral parts of the predator community, and not a novelty risk to which prey species are naïve. The tendency of the gerbils to focus their apprehension on the greatest risk in the environment supports the use of single predator type titrations favored in behavioral common-garden experiments, but potentially only true to these specific species. Therefore, these findings might or might not generalize to other arid-zone small mammals or other introduced carnivores.

Acknowledgements:

The research was funded by a United States–Israel Binational Science Foundation grant #1999–109 to BPK and JSB. We thank the Midreshet Ben Gurion Zoological Collection Reptile Collection for providing snake musk imbued bedding used in these experiments. Graphical abstract illustrated by Reza Dalvand.

Declaration of generative AI and AI-assisted technologies in the writing process:

During the preparation of this work the authors used Chat GPT in order to generate base illustrations based on personal photographs for the graphical abstract. No AI was used in writing or editing this work.

Funding

The research was funded by a United States–Israel Binational Science Foundation grant #1999–109 to Burt P. Kotler and Joel S. Brown.

Author Contributions

Conceptualization: Justin St Juliana and Burt Kotler, Methodology: Justin St. Juliana, Joel Brown and Burt Kotler; Formal analysis and investigation: Sonny Bleicher, Justin St Juliana, Shomen Mukherjee and Vijayan Sundararaj; Writing – original draft preparation: Sonny Bleicher; Writing – review and editing: Justin St Juliana, Burt Kotler and Sonny Bleicher, Funding acquisition: Burt Kotler and Joel Brown; Resources: Burt Kotler and Joel Brown; Supervision: Burt Kotler

Ethical Statement

Conflicts of interest:

All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript.

Research involving Human Participants and/or Animals:

The research was conducted in compliance with the provisions of the Israel Nature and Parks Authority, in lieu of an ethics committee established at Ben Gurion University at the time of this experiment. The animals collected in the sand dunes were collected under permit from the Israel Nature and Parks Authority Permit number 01/2003 to BPK. The procedures used in this study adhere to the tenets of the Declaration of Helsinki.

References

1. Ale, S.B.; Whelan, C.J. Reappraisal of the Role of Big, Fierce Predators! *Biodivers. Conserv.* **2008**, *17*, 685–690, doi:10.1007/s10531-008-9324-5.

2. Laundré, J.W.; Hernandez, L.; Ripple, W.J. The Landscape of Fear: Ecological Implications of Being Afraid. *Open Ecol. J.* **2010**, *3*, 1–7, doi:10.2174/1874213001003030001.

3. Rosenzweig, M.L.; MacArthur, R.H. Graphical Representation and Stability Conditions of Predator-Prey Interactions. *Am. Nat.* **1963**, *97*, 209–223.

4. Stephan, J.G.; Stenberg, J.A.; Björkman, C. Consumptive and Non-Consumptive Effect Ratios Depend on Interaction between Plant Quality and Hunting Behaviour of Omnivorous Predators Running Head: Plant Quality and Hunting Behaviour Affecting Prey. *Ecol. Evol.* **2017**, *7*, 2327–2339, doi:10.1002/ece3.2828.

5. Peckarsky, B.L.; Abrams, P.A.; Bolnick, D.I.; Dill, L.M.; Grabowski, J.H.; Luttbeg, B.; Orrock, J.L.; Peacor, S.D.; Preisser, E.L.; Schmitz, O.J.; et al. Revisiting the Classics: Considering Nonconsumptive Effects in Textbook Examples of Predator Prey Interactions. *Ecology* **2008**, *89*, 2416–2425, doi:10.1890/07-1131.1.

6. Paine, R.T. Food Web Complexity and Species Diversity. *Am. Nat.* **1963**, *100*, 65–75, doi:10.2307/2678832.

7. Lima, S.L. Nonlethal Effects in the Ecology of Predator-Prey Interactions. *Bioscience* **1998**, *48*, 25–34, doi:10.2307/1313225.

8. Orrock, J.L.; Grabowski, J.H.; Pantel, J.H.; Peacor, S.D.; Peckarsky, L.; Sih, A.; Werner, E.E.; Werner, E. Consumptive and Nonconsumptive Effects of Predators on Metacommunities of Competing Prey. *Ecology* **2008**, *89*, 2426–2435.

9. Sih, A.; Bolnick, D.I.; Luttbeg, B.; Orrock, J.L.; Peacor, S.D.; Pintor, L.M.; Preisser, E.; Rehage, J.S.; Vonesh, J.R. Predator-Prey Naïveté, Antipredator Behavior, and the Ecology of Predator Invasions. *Oikos* **2010**, *119*, 610–621, doi:10.1111/j.1600-0706.2009.18039.x.

10. Schmitz, O.J.; Beckerman, A.P.; Brien, K.M.O. Behaviorally Mediated Trophic Cascades: Effects of Predation Risk on Food Web Interactions. *Ecology* **1997**, *78*, 1388–1399.

11. Letnic, M.; Ritchie, E.G.; Dickman, C.R. Top Predators as Biodiversity Regulators: The Dingo Canis Lupus Dingo as a Case Study. *Biol. Rev.* **2012**, *87*, 390–413, doi:10.1111/j.1469-185X.2011.00203.x.

12. Beschta, R.L.; Ripple, W.J. Large Predators and Trophic Cascades in Terrestrial Ecosystems of the

563   Western United States. *Biol. Conserv.* **2009**, *142*, 2401–2414, doi:10.1016/j.biocon.2009.06.015.

564   13.   Kotler, B.P.; Holt, R.D. Predation and Competition: The Interaction of Two Types of Species.
565   *Oikos* **1989**, *54*, 256–260.

566   14.   Manlick, P.J.; Newsome, S.D. Adaptive Foraging in the Anthropocene: Can Individual Diet
567   Specialization Compensate for Biotic Homogenization? *Front. Ecol. Environ.* **2021**, *19*, 510–518,
568   doi:10.1002/fee.2380.

569   15.   Bleicher, S.S.; Dickman, C.R. On the Landscape of Fear: Shelters Affect Foraging by Dunnarts
570   (Marsupialia, Sminthopsis Spp.) in a Sandridge Desert Environment. *J. Mammal.* **2020**, *101*, 281–
571   290, doi:10.1093/jmammal/gyz195.

572   16.   Pavey, C.R.; Eldridge, S.R.; Heywood, M. Population Dynamics and Prey Selection of Native and
573   Introduced Predators during a Rodent Outbreak in Arid Australia. *J. Mammal.* **2008**, *89*, 674–683,
574   doi:10.1644/07-MAMM-A-168R.1.

575   17.   Dickman, C.R.; Greenville, A.C.; Tamayo, B.; Wardle, G.M. Spatial Dynamics of Small
576   Mammals in Central Australian Desert Habitats: The Role of Drought Refugia. *J. Mammal.* **2011**,
577   *92*, 1193–1209, doi:10.1644/10-MAMM-S-329.1.

578   18.   Abramsky, Z.; Rosenzweig, M.L.; Subach, A. The Costs of Apprehensive Foraging. *Ecology*
579   **2002**, *83*, 1330–1340.

580   19.   Kotler, B.P.; Blaustein, L. Titrating Food and Safety in a Heterogeneous Environment: When
581   Are the Risky and Safe Patches of Equal Value? *Oikos* **1995**, *74*, 251–258.

582   20.   Brown, J.S.; Kotler, B.P. Hazardous Duty Pay and the Foraging Cost of Predation. *Ecol. Lett.*
583   **2004**, *7*, 999–1014, doi:10.1111/j.1461-0248.2004.00661.x.

584   21.   Relyea, R.A. How Prey Respond to Combined Predators: A Review and an Empirical Test.
585   *Ecology* **2003**, *84*, 1827–1839.

586   22.   Arditi, R.; Ginzburg, L.R. Coupling in Predator-Prey Dynamics: Ratio Dependence. *J. Theor. Biol.*
587   **1989**, *139*, 311–326.

588   23.   St. Juliana, J.; Kotler, B.P.; Brown, J.S.; Mukherjee, S.; Bouskila, A. The Foraging Response of
589   Gerbils to a Gradient of Owl Numbers. *Evol. Ecol. Res.* **2011**, *13*, 869–878.

590   24.   Mukherjee, S. Understanding the Interplay Between Time Allocation, Vigilance and State
591   Department Foraging Behavior. (Doctoral Dissertation, Ben Gurion University of the Negev).,
592   Ben Gurion University, 2010.

593   25.   Berec, L. Impacts of Foraging Facilitation among Predators on Predator-Prey Dynamics. *Bull.*
594   *Math. Biol.* **2010**, *72*, 94–121, doi:10.1007/s11538-009-9439-1.

595   26.   Soluk, D.A.; Collins, N.C. Between Fish and Stoneflies: Facilitation Interactions Synergistic and
596   Interference among Stream Predators. *Oikos* **1988**, *52*, 94–100.

597   27.   Embar, K.; Kotler, B.P.; Bleicher, S.S.; Brown, J.S. Pit Fights: Predators in Evolutionarily
598   Independent Communities. *J. Mammal.* **2018**, *99*, 1183–1188, doi:10.1093/jmammal/gyy085.

599   28.   Kotler, B.P.; Blaustein, L.; Brown, J.S. Predator Facilitation: The Combined Effect of Snakes and
600   Owls on the Foraging Behavior of Gerbils. *Ann. Zool. Fenn.* **1992**, *29*, 199–206.

601   29.   Bleicher, S.S.; Brown, J.S.; Embar, K.; Kotler, B.P. Novel Predator Recognition by Allenby's
602   Gerbil (Gerbillus Andersoni Allenbyi): Do Gerbils Learn to Respond to a Snake That Can "See" in

the Dark? *Isr. J. Ecol. Evol.* **2016**, *62*, 178–185, doi:10.1080/15659801.2016.1176614.

30.   Bleicher, S.S.; Kotler, B.P.; Brown, J.S. Comparing Plasticity of Response to Perceived Risk in
the Textbook Example of Convergent Evolution of Desert Rodents and Their Predators; a
Manipulative Study Employing the Landscape of Fear. *Front. Behav. Neurosci.* **2019**, *13*, 1–12,
doi:10.3389/fnbeh.2019.00058.

31.   Kotler, B.P.; Brown, J.S.; Bleicher, S.S.; Embar, K. Intercontinental-Wide Consequences of
Compromise-Breaking Adaptations: The Case of Desert Rodents. *Isr. J. Ecol. Evol.* **2016**, *62*,
186–195, doi:10.1080/15659801.2015.1125832.

32.   Bleicher, S.S.; Kotler, B.P.; Downs, C.J.; Brown, J.S. Playing to Their Evolutionary Strengths;
Heteromyid Rodents Provide Opposite Snake Evasion Strategies in the Face of Known and Novel
Snakes. *J. Arid Environ.* **2020**, *173*, 104025, doi:10.1016/j.jaridenv.2019.104025.

33.   Embar, K.; Raveh, A.; Hoffmann, I.; Kotler, B.P. Predator Facilitation or Interference: A Game of
Vipers and Owls. *Oecologia* **2014**, *174*, 1301–1309, doi:10.1007/s00442-013-2760-2.

34.   Leo, V.; Reading, R.P.; Letnic, M. Interference Competition: Odours of an Apex Predator and
Conspecifics Influence Resource Acquisition by Red Foxes. *Oecologia* **2015**, *179*,
doi:10.1007/s00442-015-3423-2.

35.   Kotler, B.P.; Brown, J.S.; Mukherjee, S.; Berger-Tal, O.; Bouskila, A.; Kotier, B.P.; Brown, J.S.;
Mukherjee, S.; Berger-Tal, O.; Bouskila, A. Moonlight Avoidance in Gerbils Reveals a
Sophisticated Interplay among Time Allocation , Vigilance and State-Dependent Foraging. *Proc.*
R. Soc. B **2010**, *277*, 1469–1474, doi:10.1098/rspb.2009.2036.

36.   Wasko, D.K.; Bonilla, F.; Sasa, M. Behavioral Responses to Snake Cues by Three Species of
Neotropical Rodents. *J. Zool.* **2014**, *292*, doi:10.1111/jzo.12086.

37.   Orizaola, G.; Dahl, E.; Laurila, A. Reversibility of Predator-Induced Plasticity and Its Effect at a
Life-History Switch Point. *Oikos* **2012**, *121*, 44–52, doi:10.1111/j.1600-0706.2011.19050.x.

38.   Bleicher, S.S.; Kotler, B.P.; Downs, C.J.; Brown, J.S. Intercontinental Test of
Constraint☐Breaking Adaptations; Testing Behavioural Plasticity in the Face of a Predator with
Novel Hunting Strategies. *J. Anim. Ecol.* **2020**, *89*, 1837–1850, doi:10.1111/1365-2656.13234.

39.   Abramsky, Z.; Strauss, E.; Subach, A.; Riechman, A.; Kotler, B.P. The Effect of Barn Owls (Tyto
Alba) on the Activity and Microhabitat Selection of Gerbillus Allenbyi and G. Pyramidum.
Oecologia **1996**, *105*, 313–319, doi:10.1007/BF00328733.

40.   Rosenzweig, M.L. Habitat Selection as a Source of Biological Diversity. *Evol. Ecol.* **1987**, *1*, 315–
330, doi:10.1007/BF02071556.

41.   Ben-Natan, G.; Abramsky, Z.; Kotler, B.P.; Brown, J.S. Seeds Redistribution in Sand Dunes☐: A
Basis for Coexistence of Two Rodent Species. *Oikos* **2004**, *105*, 325–335, doi:10.1111/j.0030-
1299.2004.12948.x.

42.   Brown, J.S.; Kotler, B.P.; Mitchell, W.A. Competition between Birds and Mammals☐: A
Comparison of Giving-up Densities between Crested Larks and Gerbils. *Evol. Ecol.* **1997**, *11*,
757–771, doi:10.1023/A:1018442503955.

43.   Kotler, B.P.; Brown, J.S.; Dall, S.R.X.; Gresser, S.; Ganey, D.; Bouskila, A. Foraging Games
between Gerbils and Their Predators☐: Temporal Dynamics of Resource Depletion and
Apprehension in Gerbils. *Evol. Ecol. Res.* **2002**, *4*, 495–518.

44.   Biewener, A.; Blickhan, R. Kangaroo Rat Locomotion: Design for Elastic Energy Storage or
645   Acceleration? *J. Exp. Biol.* **1988**, *140*, 243–255.

646   45.   Longland, W.S.; Price, M. V Direct Observations of Owls and Heteromyid Rodents : Can
647   Predation Risk Explain Microhabitat Use ? *Ecology* **1991**, *72*, 2261–2273.

648   46.   Webster, D.B.; Strother, W.F. Middle-Ear Mor- Phology and Auditory Sensitivity of Heteromyid
649   Rodents. *Am. Zool.* **1972**, *12*, 727.

650   47.   Abramsky, Z.; Rosenzweig, M.L.; Subach, A. The Cost of Interspecific Competition in Two
651   Gerbil Species. *J. Anim. Ecol.* **2001**, *70*, 561–567.

652   48.   Makin, D.F.; Kotler, B.P. How Do Allenby ' s Gerbils Titrate Risk and Reward in Response to
653   Different Predators ? *Behav. Ecol. Sociobiol.* **2020**, *74*, 1–10, doi:https://doi.org/10.1007/s00265-
654   019-2785-6 (2020).

655   49.   Letnic, M.; Koch, F. Are Dingoes a Trophic Regulator in Arid Australia? A Comparison of
656   Mammal Communities on Either Side of the Dingo Fence. *Austral Ecol.* **2010**, *35*, 167–175,
657   doi:10.1111/j.1442-9993.2009.02022.x.

658   50.   Berger, S.; Wikelski, M.; Romero, L.M.; Kalko, E.K. V; Rödl, T. Behavioral and Physiological
659   Adjustments to New Predators in an Endemic Island Species, the Galápagos Marine Iguana.
660   *Horm. Behav.* **2007**, *52*, 653–663, doi:10.1016/j.yhbeh.2007.08.004.

661   51.   Carthey, A.J.R.; Blumstein, D.T. Predicting Predator Recognition in a Changing World. *Trends
662   Ecol. Evol.* **2018**, *33*, 106–115, doi:10.1016/j.tree.2017.10.009.

663   52.   Goodfriend, W.; Ward, D.; Subach, A. Standard Operative Temperatures of Two Desert Rodents,
664   Gerbillus Allenbyi and Gerbillus Pyramidum: The Effects of Morphology, Microhabitat and
665   Environmental Factors. *J. Therm. Biol.* **1991**, *16*, 157–166, doi:10.1016/0306-4565(91)90038-4.

666   53.   Kotler, B.P.; Brown, J.S.; Oldfield, A.; Thorson, J.; Cohen, D. Foraging Substrate and Escape
667   Substrate: Patch Use by Three Species of Gerbils. *Ecology* **2001**, *82*, 1781–1790,
668   doi:10.1890/0012-9658(2001)082[1781:FSAESP]2.0.CO;2.

669   54.   Ellard, C.G.; Eller, M.C. Spatial Cognition in the Gerbil: Computing Optimal Escape Routes from
670   Visual Threats. *Anim. Cogn.* **2009**, *12*, 333–345, doi:10.1007/s10071-008-0193-9.

671   55.   Lay, D.M. The Anatomy, Physiology, Functional Significance and Evolution of Specialized
672   Hearing Organs of Gerbilline Rodents. *J. Morphol.* **1972**, *138*, 41–120,
673   doi:10.1002/jmor.1051380103.

674   56.   Emerson, S.E.; Kotler, B.P.; Sargunaraj, F. Foraging Efficiency in the Face of Predation Risk: A
675   Comparative Study of Desert Rodents. *Evol. Ecol. Res.* **2018**, *19*, 61–70.

676   57.   Panday, P.; Pal, N.; Samanta, S.; Tryjanowski, P.; Chattopadhyay, J. Dynamics of a Stage-
677   Structured Predator-Prey Model : Cost and Benefit of Fear-Induced Group Defense. *J. Theor.
678   Biol.* **2021**, *528*, 110846, doi:10.1016/j.jtbi.2021.110846.

679   58.   Kotler, B.P.; Brown, J.S.; Bouskila, A.; Mukherjee, S.; Goldberg, T. Foraging Games between
680   Gerbils and Their Predators: Seasonal Changes in Schedules of Activity and Apprehention. *Isr. J.
681   Ecol. Evol.* **2004**, *50*, 255–271.

682   59.   Dall, S.R.X.; Kotler, B.P.; Bouskila, a Attention, "apprehension" and Gerbils Searching in
683   Patches. *Ann. Zool. Fennici* **2001**, *38*, 15–23.

60.   Bleicher, S.S. Multi-Predator Common Garden Experiment with Two Gerbil Species.

61.   Jamovi The Jamovi Project 2023, Version 2.4.

62.   Brown, J.S. Vigilance , Patch Use and Habitat Selection: Foraging under Predation Risk. *Evol.*
Ecol. Res. **1999**, *1*, 49–71.

63.   Brown, J.S. Patch Use as an Indicator of Habitat Preference, Predation Risk, and Competition.
Behav. Ecol. Sociobiol. **1988**, *22*, 37–47, doi:10.1007/BF00395696.

64.   Bouskila, A. A Habitat Selection Game of Interactions between Rodents and Their Predators. *Ann.*
Zool. Fenn. **2001**, *38*, 55–70.

65.   Bleicher, S.S.; Kotler, B.P.; Shalev, O.; Austin, D.; Embar, K.; Brown, J.S.; Dixon, A. Divergent
Behavior amid Convergent Evolution: A Case of Four Desert Rodents Learning to Respond to
Known and Novel Vipers. *PLoS One* **2018**, *13*, 1–17, doi:10.1101/362202.

66.   Kotler, B.P.; Brown, J.S.; Mitchell, W.A. Environmental Factors Affecting Patch Use in Two
Species of Gerbelline Rodents. *J. Mammal.* **1993**, *74*, 614–620, doi:10.2307/1382281.

67.   Kotler, B.P.; Brown, J.S.; Subach, A. Temporal Foragers: Of Optimal Coexistence of Species
Mechanisms of Sand Dune Gerbils by Two Species Partitioning. *Oikos* **1993**, 548–556.

68.   Bleicher, S.S.; Ylönen, H.; Käpylä, T.; Haapakoski, M. Olfactory Cues and the Value of
Information: Voles Interpret Cues Based on Recent Predator Encounters. *Behav. Ecol. Sociobiol.*
2018, *72*, 187–199, doi:10.1007/s00265-018-2600-9.

69.   Porter, W.P.; Budaraju, S.; Stewart, W.E.; Ramankutty, N. Calculating Climate Effects on Birds
and Mammals: Impacts on Biodiversity, Conservation, Population Parameters, and Global
Community Structure. *Am. Zool.* **2000**, *40*, 597–630, doi:10.1093/icb/40.4.597.

70.   Kotler, B.P.; Brown, J.S. Mechanisms of Coexistence of Optimal Foragers as Determinants of
Local Abundances and Distributions of Desert Granivores. *J. Mammal.* **1999**, *80*, 361–374.

71.   Ziv, Y.; Abramsky, Z.; Kotler, B.P.; Subach, A. Interference Competition and Temporal and
Habitat Partitioning in Two Gerbil Species, 1993, Vol. 66.

72.   Dixon, A.K. Tradeoffs of Food and Safety in Contrasting Environments: From the Deserts of the
Mojave and the Negev to the Coral Reefs of Eilat. Ph.D. Thesis., Ben Gurion University of the
Negev, 2017.

73.   Blumstein, D.T.; Mari, M.; Daniel, J.C.; Ardron, J.G.; Griffin, A.S.; Evans, C.S. Olfactory
Predator Recognition: Wallabies May Have to Learn to Be Wary. *Anim. Conserv.* **2002**, *5*, 87–93,
doi:10.1017/S1367943002002123.

74.   Thorson, J.M.; Morgan, R.A.; Brown, J.S.; Norman, J.E. Direct and Indirect Cues of Predatory
Risk and Patch Use by Fox Squirrels and Thirteen-Lined Ground Squirrels. *Behav. Ecol.* **1998**, *9*,
151–157.

75.   Morgan, R.A.; Brown, J.S.; Thorson, J.M. The Effect of Spatial Scale on the Functional Response
of Fox Squirrels. *Ecology* **2013**, *78*, 1087–1097, doi:10.1890/0012-
9658(1997)078[1087:TEOSSO]2.0.CO;2.

76.   Bytheway, J.P.; Carthey, A.J.R.R.; Banks, P.B. Risk vs. Reward: How Predators and Prey
Respond to Aging Olfactory Cues. *Behav. Ecol. Sociobiol.* **2013**, *67*, 715–725,
doi:10.1007/s00265-013-1494-9.

77.   Banks, P.B.; Bytheway, J.P.; Carthey, A.J.R.; Hughes, N.K.; Price, C.J. Olfaction and Predaotr-
Prey Interactions Amongst Mammals in Australia. In *Predators in Australia*; Glen, A.S.,

Dickman, C.R., Eds.; CSIRO PUBLISHING: Collingwood, 2014; p. Chapter 17.

78.   Bleicher, S.S. Heat and Humidity Alter Predation Cues in Gerbillus Andersoni Allebyi. MS
Thesis. Ben Gurion University of the Negev, Ben Gurion University of the Negev, 2012.

79.   Laundré, J.W.; Calderas, J.M.M.; Hernández, L. Foraging in the Landscape of Fear , the
Predator's Dilemma: Where Should I Hunt□? *Open Ecol. J.* **2009**, *2*, 1–6.

80.   Vance-Chalcraft, H.D.; Soluk, D.A.; Ozburn, N. Is Prey Predation Risk Influenced More by
Increasing Predator Density or Predator Species Richness in Stream Enclosures? *Oecologia* **2004**,
139, 117–122, doi:10.1007/s00442-003-1484-0.

81.   Martín, J.; López, P.; Polo, V. Temporal Patterns of Predation Risk Affect Antipredator Behaviour
Allocationby Iberian Rock Lizards. *Anim. Behav.* **2009**, *77*, 1261–1266,
doi:10.1016/j.anbehav.2009.02.004.

82.   Abrams, P.A. Should Prey Overestimate the Risk of Predation? *Am. Nat.* **1994**, *144*, 317–328.

83.   Bouskila, A.; Blumstein, D.T.; Mangel, M. Prey under Stochastic Condition Should Probably
Overstimate Predation Risk: A Reply to Abrams. *Am. Nat.* **1995**, *145*, 1015–1019.

84.   Bouskila, A.; Blumstein, D.T. Rules of Thumb for Predation Hazard Assessment□: Predictions
from a Dynamic Model. *Am. Nat.* **1992**, *139*, 161–176.

85.   Dickman, C.R. Overview of the Impacts of Feral Cats On. *Nature* **1996**, 92.

86.   Doherty, T.S.; Dickman, C.R.; Johnson, C.N.; Legge, S.M.; Ritchie, E.G.; Woinarski, J.C.Z.
Impacts and Management of Feral Cats Felis Catus in Australia. *Mamm. Rev.* **2017**, *47*, 83–97,
doi:10.1111/mam.12080.

87.   Hardman, B.; Moro, D.; Calver, M. Direct Evidence Implicates Feral Cat Predation as the Primary
Cause of Failure of a Mammal Reintroduction Programme. *Ecol. Manag. Restor.* **2016**, *17*, 152–
158, doi:10.1111/emr.12210.

88.   Moseby, K.E.; Peacock, D.E.; Read, J.L. Catastrophic Cat Predation: A Call for Predator Profiling
in Wildlife Protection Programs. *Biol. Conserv.* **2015**, *191*, 331–340,
doi:10.1016/j.biocon.2015.07.026.

89.   Labra, A.; Leonard, R. Intraspecific Variation in Antipredator Responses of Three Species of
Lizards (Liolaemus): Possible Effects of Human Presence. *J. Herpetol.* **1999**, *33*, 441–448.

90.   Avilés-Rodríguez, K.J.; Kolbe, J.J. Escape in the City: Urbanization Alters the Escape Behavior of
Anolis Lizards. *Urban Ecosyst.* **2019**, *22*, 733–742, doi:10.1007/s11252-019-00845-x.

91.   Zuñiga-Palacios, J.; Zuria, I.; Castellanos, I.; Lara, C.; Sánchez-Rojas, G. What Do We Know (and
Need to Know) about the Role of Urban Habitats as Ecological Traps? Systematic Review and
Meta-Analysis. *Sci. Total Environ.* **2021**, *780*, 1–11, doi:10.1016/j.scitotenv.2021.146559.

92.   Glen, A.S.; Dickman, C.R. Complex Interactions among Mammalian Carnivores in Australia, and
Their Implications for Wildlife Management. *Biol. Rev. Camb. Philos. Soc.* **2005**, *80*, 387–401,
doi:10.1017/s1464793105006718.

93.   Brito, J.C.; Barrio, G. Del; Stellmes, M.; Pleguezuelos, J.M.; Saarinen, J.; Barrio, G. Del;
Stellmes, M.; Pleguezuelos, J.M.; Saarinen, J. Drivers of Change and Conservation Needs for
Vertebrates in Drylands□: An Assessment from Global Scale to Sahara-Sahel Wetlands an
Assessment from Global Scale to Sahara-Sahel Wetlands. *Eur. Zool. J.* **2021**, *88*, 1103–1129,
doi:10.1080/24750263.2021.1991496.

94. Medina, F.M.; Nogales, M. A Review on the Impacts of Feral Cats (Felis Silvestris Catus) in the
Canary Islands□: Implications for the Conservation of Its Endangered Fauna. *Biodivers. Conserv.*
2009, *18*, 829–846, doi:10.1007/s10531-008-9503-4.

95. Banks, P.B.; Dickman, C.R. Alien Predation and the Effects on Multiple Levels of Prey Naivite.
Trends Ecol. Evol. **2007**, *22*, 229–230, doi:10.1016/j.tree.2007.02.003.

96. Carthey, A.J.R.; Banks, P.B. Naïveté in Novel Ecological Interactions: Lessons from Theory and
Experimental Evidence. *Biol. Rev. Camb. Philos. Soc.* **2014**, *89*, 932–949, doi:10.1111/brv.12087.

97. Rödl, T.; Berger, S.; Romero, L.M.; Wikelski, M. Tameness and Stress Physiology in a Predator-
Naive Island Species Confronted with Novel Predation Threat. *Proc. Biol. Sci.* **2007**, *274*, 577–
582, doi:10.1098/rspb.2006.3755.

98. Banks, P.B.; Carthey, A.J.R.; Bytheway, J.P.; Banks, P.B.; Carthey, A.J.R.; Bytheway, J.P.
Australian Native Mammals Recognize and Respond to Alien Predators□: A Meta- Analysis.
2018, *285*, 1–8.

99. Ramp, D.; Benítez-lópez, A.; Carroll, S.; Carthey, A.J.R.; Rogers, E.I.E.; Zawada, K.J.A.;
Middleton, O.; Lundgren, E.; Svenning, J.; Avidor, E. Savviness of Prey to Introduced Predators.
2023, 1–11, doi:10.1111/cobi.14012.

100. Serpell, J.A. Domestication and History of the Cat. In *The domestic cat: The biology of Iits*
behavior; Turne, D.C., Bateson, P., Eds.; Cambridge Univ. Press.: Cambridge, UK, 2000; pp.
180–192 ISBN 9781139177177.

101. Banks, P.B.; Dickman, C.R.; Newsome, A.E. Ecological Costs of Feral Predator Control: Foxes
and Rabbits. *J. Wildl. Manage.* **1998**, *62*, 766–772, doi:10.2307/3802353.

Figure and Table Legends

Table 1. Predictive patterns of gerbil behavior in relation to four predators used in the experiment.

Table 2. Inference table associating gerbil foraging with likely predator relationship. Replacement is only
likely with low lethality predators, or in this experiment where an olfactory-cue was used in lieu of using
live snakes.

Table 3. T-test comparisons between the control and every single predator treatment, for the combined
data set, and each of the species separately.

Table 4: ANOVA table for GLM using data from both species (N= 2752, multiple R= 0.51, and multiple
R^2 = 0.26)

Table 5. Three-way interactions post hoc Tukey Tests of Honestly Significant Difference for A. Gerbillus
andersoni allebyi, and B. Gerbillus pyramidum. Only significant interactions are listed, and * delineates
one tailed distribution significance

Fig 1. Foraging activity of each of the gerbil species and both combined in isolated, single predator
treatments. α dictates a significant pairwise comparison (between control and each predator) for the
combined gerbil species, β for the smaller gerbil (GA), and γ for the larger species (GP) corresponding to
Table 3. The patterns shown here suggest a greater selectivity and "evaluation" of risk performed by GP
than GA.

Fig 2. Gerbil foraging adjustment in response to perceived risk from combinations of predators. The bars
represent the mean change in foraging from a treatment with a single predator and a combination of that
same predator with others. (A) combined species dataset, (B) GA, and (C) GP. If the adjustment is
positive (*i.e.,* combined predator GUD was lower than the single predator) the trend is suggestive of
predator interference. When the adjustment is negative, the gerbils foraged less with the combination of
predators than the single predator, thus suggesting a pattern of facilitation or accumulation of risk. Bold
bars delineate post hoc significant pairwise comparisons (Tukey HSD) between the single predator and
the combination with another. The brackets delineate significant difference between predator
combinations. Asterisks represent one tailed significance (P-values between 0.05-0.1). The combined data
suggests opposite patterns than observed for each species independently.

Fig 3. Combined Gerbil foraging (GUD±SE) as influenced by predator type and microhabitat (bush/cover
in gray and open in white). Each panel focuses on a different predator and the combination with other
predators where (A) focuses on the Dog, (B) the Cat, (C) the Owl, and (D) Snake musk. Greek letters
represent significant Tukey Pairwise Comparisons, read from left to right for each panel.

Tables

Table 1. Predictive patterns of gerbil behavior in relation to four predators used in the experiment.

Predator	Safer microhabitat for gerbils	Predator hunting activity type	Potential interference	Expected facilitation	Potential replacement to
Cat	Bush	Ground foraging	Dog, Owl	Snake	Snake
Dog	Bush	Ground foraging	Cat, Owl	Snake	Snake
Snake	Open	Ambush		Cat, Dog, Owl	
Owl	Bush	Aerial	Cat, Dog	Snake	Snake

826 **Table 2.** Inference table associating gerbil foraging with likely predator relationship. Replacement is only
827 likely with low lethality predators, or in this experiment where an olfactory-cue was used in lieu of using
828 live snakes.

Main Predator	Predator Combination	Predicted Likely Outcomes		
		dGUD if Interference	**dGUD if Facilitation/ Cumulative Effects**	**dGUD if Replacement (Likely focus predator)**
Snake	Cat		= between microhabitats/- ↓	-↓ (Cat)
	Dog		= between microhabitats /- ↓	-↓ (Dog)
	Owl		= between microhabitats /- ↓	-↓ (Owl)
Owl	Snake		= between microhabitats /- ↓	+↑ (Owl)
	Dog	+ ↑		
	Cat	+ ↑		
Cat	Dog	+ ↑		
	Owl	+ ↑		
	Snake		= between microhabitats /- ↓	=/+↑ (Cat)
Dog	Cat	+ ↑		
	Owl	+ ↑		
	Snake		= between microhabitats / - ↓	=/+↑ (Dog)

831   Table 3. T-test comparisons between the control and every single predator treatment, for the combined
832   data set, and each of the species separately.

		Both Gerbils			GA			GP		
Group 1	Group 2	t-critical	df	P-value	t-critical	df	P-value	t-critical	df	P-value
Control	Snake	-4.03	1342	<0.001	-4.21	670	<0.001	-1.55	670	0.122
Control	Owl	-1.44	1342	0.162	-4.32	670	<0.001	+2.62	670	0.009
Control	Cat	-8.33	1278	<0.001	-6.33	638	<0.001	-5.74	638	<0.001
Control	Dog	-11.11	1278	<0.001	-6.74	638	<0.001	-9.16	638	<0.001

836   Table 4: ANOVA table for GLM using data from both species (N= 2752, multiple R= 0.51, and multiple
837   R^2 = 0.26)

Source	Type III SS	df	Mean Squares	F-Ratio	p-Value
Gerbil Species (GS)	27.483	1	27.483	79.288	0.000
Predator Treatment (PT)	50.070	9	5.563	16.050	0.000
Moonphase	1.139	3	0.380	1.095	0.350
Resource Distribution (Clumped/ Evenly Mixed) (RD)	0.031	1	0.031	0.090	0.764
Microhabitat	41.602	1	41.602	120.021	0.000
GS X PT	9.019	9	1.002	2.891	0.002
Moonphase X GS	6.922	3	2.307	6.656	0.000
Microhabitat X GS	22.670	1	22.670	65.401	0.000
Moonphase X PT	72.831	27	2.697	7.782	0.000
Microhabitat X PT	6.701	9	0.745	2.148	0.023
Microhabitat X RD	3.834	1	3.834	11.060	0.001
RD X GS	0.256	1	0.256	0.739	0.390
GS X PT X Moonphase	59.957	27	2.221	6.407	0.000
PT X Moonphase X RD	18.991	27	0.703	2.029	0.001
Error	911.965	2,631	0.347		

838
839

Table 5. Three-way interactions post hoc Tukey Tests of Honestly Significant Difference for A. Gerbillus andersoni allebyi, and B. Gerbillus pyramidum. Only significant interactions are listed, and * delineates one tailed distribution significance.

GA							
#	Fixed Variable	Category	Dynamic Variable	Relationship			P-value
1	Moonphase	Full	Predators	Dog	>	Snake	0.022
2	Moonphase	New	Predators	Dog+ Snake	>	Snake	<0.001
3	Moonphase	New	Predators	Owl	>	Snake	<0.001
4	Moonphase	Waning	Predators	Cat + Dog	>	Cat + Snake	<0.001
5	Moonphase	Waning	Predators	Dog + Owl	>	Owl	0.009
6	Moonphase	Waxing	Predators	Dog + Cat	>	Dog + Snake	0.009
7	Moonphase	Waxing	Predators	Dog + Cat	>	Dog	<0.001
8	Moonphase	Waxing	Predators	Dog + Cat	>	Dog + Snake	0.001
9	Predators	Cat + Snake	Moonphase	Full	>	Waning	<0.001
10	Predators	Dog	Moonphase	New	>	Full	0.023
11	Predators	Dog	Moonphase	Full	>	Waxing	<0.001
12	Predators	Dog+ Snake	Moonphase	New	>	Waxing	0.013
13	Predators	Owl	Moonphase	Waxing	>	New	0.059*
14	Predators	Cat + Owl	Moonphase	Waxing	>	Waning	0.031
GP							
#	Fixed Variable	Category	Dynamic Variable	Relationship			P-value
1	Moonphase	Full	Predators	Cat	>	Dog	0.058*
2	Moonphase	Full	Predators	Cat + Owl	>	Dog	0.044
3	Moonphase	Full	Predators	Cat + Dog	>	Cat + Snake	<0.001
4	Moonphase	Full	Predators	Cat + Dog	>	Dog	<0.001
5	Moonphase	Full	Predators	Owl	>	Dog	<0.001
6	Moonphase	New	Predators	Cat	>	Owl	<0.001
7	Moonphase	New	Predators	Cat + Owl	>	Owl	<0.001
8	Moonphase	New	Predators	Dog	>	Dog + Snake	0.048
9	Moonphase	New	Predators	Dog	>	Owl	<0.001

10	Moonphase	New	Predators	Dog + Cat	>	Dog + Snake	<0.001
11	Moonphase	New	Predators	Dog + Owl	>	Owl	<0.001
12	Moonphase	Waning	Predators	Cat	>	Snake	<0.001
13	Moonphase	Waning	Predators	Cat + Owl	>	Dog + Owl	0.008
14	Moonphase	Waning	Predators	Cat + Snake	>	Snake	<0.001
15	Moonphase	Waxing	Predators	Dog + Owl	>	Dog + Cat	0.022
16	Moonphase	Waxing	Predators	Dog	>	Owl	<0.001
17	Moonphase	Waxing	Predators	Dog	>	Snake	<0.001
18	Predators	Cat + Snake	Moonphase	Waning	>	Full	<0.001
19	Predators	Dog	Moonphase	New	>	Full	<0.001
20	Predators	Dog	Moonphase	Waxing	>	Full	<0.001
21	Predators	Dog+ Snake	Moonphase	Waxing	>	New	0.046

845   **Fig 1.**

 Fig. 2

 Fig 3.

List of Appendices:

SI 1. Multi-way ANOVA for GA, including post-hoc Tukey tests for all significant variables and
interactions.

SI 2. Multi-way ANOVA for GP, including post-hoc Tukey tests for all significant variables and
interactions.

SI 3. Pearson Correlation matrices for the covariates against GUD for each of the gerbil species
separately.

SI 4. Response of combined gerbil species to predator encounters.

SI 5. Response of gerbils to predator treatments as affected by moonphase.

SI 6. Raw-data file used for analyses.