Additive, substitutable, and antagonistic prey responses to feral and native
predator combinations.

Justin R. Saint Juliana[1], S.S. Bleicher[2*], S. Mukherjee[3,4], V. Sundararaj[5], J.S. Brown[6], B.P Kotler[7]

1. Cornell University, Dept. of Ecology and Evolutionary Biology, Ithaca, NY, 14853, USA

2. Loyola University New Orleans, Biology Department, 6363 St. Charles Ave, New Orleans, LA, 70118, USA

3. Biological and Life Sciences Division, School of Arts and Sciences, Ahmedabad University, Ahmedabad,
Gujarat, 6 India 7

4. Ecology Environment and Climate Change Research Cluster, Ahmedabad University, Ahmedabad,
Gujarat, India

5. Texas A&M International University, Loredo, TX, USA.

6. Moffit Cancer Research Center, Department of Integrated Mathematical Oncology, Tampa, Florida,
33612, USA

7. Ben Gurion University of the Negev, Blaustein Institute for Desert Research, Sde Boker, Israel

*Corresponding author: Sonny S. Bleicher, sbleiche@loyno.edu

Abstract

Question: Do combinations of human commensal mesocarnivores with native predators alter wild prey
behavior in additive or substitutable responses?

Hypothesis: Feral mesocarnivores will have a greater impact on prey energy acquisition than native
predators.

Organisms: Allenby's gerbils (*Gerbillus andersoni allebyi*) and Greater Egyptian Gerbils (*Gerbillus*
pyramidum) as prey and feral dogs (*Canis lupus familliaris*), feral cats (*Felis catus*), barn owls (*Tyto*
alba) horned vipers (*Cerastes gasperetti*) as predators.

Field Site: A 17x34 meter semi natural arena mimicking sand dune environments in Sde Boker, Israel.

Methods: Gerbil perceived risk was measured using optimal patch-use, predator exposures were
conducted with guided (leashed) predators on an hourly basis throughout the night. Viper musk was used
in lieu of live vipers. On nights with combined predators, two predators were exposed, rotating between
the predators every hour.

Conclusions: Human-commensal predators induced a stronger foraging aversion from the gerbils than the
owl. When combining predators, the gerbils significantly decrease their foraging only when a predator
perceived to be of greater risk is introduced. This was exemplified by a significantly higher GUD for dog
and cat combination over dog alone, but insignificant increase, compared to the cat alone. The impact of
human commensals, especially feral cats, appears to outweigh the ecological impacts of native predators.

Key Words: Predator-prey dynamics; Common-garden experiments; Feral mesocarnivores; Foraging
ecology; Ecology of Fear.

Introduction

Predators are known to regulate the richness of biological communities through top-down trophic interactions (Rosenzweig and MacArthur, 1963; Ale and Whelan, 2008; Laundré *et al.*, 2010). Predator-prey interactions enrich biological communities by allowing adaptive strategies of varying prey populations to coexist in space and time (Peckarsky *et al.*, 2008; Stephan *et al.*, 2017). While the original perception of predator influence on communities was predominantly consumptive (Paine, 1963), the ecological research community has come to accept the non-consumptive impacts of predators to be equal or greater to those of direct predation (Lima, 1998; Orrock *et al.*, 2008; Sih *et al.*, 2010). Overall, increases in risk, and increases in the diversity of predators promote increased diversity of anti-predator adaptations among prey, which in turn help facilitate prey coexistence (Kotler and Holt, 1989; Schmitz *et al.*, 1997; Beschta and Ripple, 2009; Letnic *et al.*, 2012). In a period of great biodiversity loss, especially predator loss (Manlick and Newsome, 2021), the impacts of changes in predator communities are expected to reverberate down through the prey communities.

Despite the agreement on the impacts of risk on prey diversity, measuring risk intensity in empirical experiments has been proven complicated, and quantifying prey response to said variation is harder still. The current standard for testing impact of predation risk on prey, involves captive experiments, managing titration of food and safety through the addition of predators (*c.f.* Abramsky et al. 2002), or regulating available resources in the environment (Kotler and Blaustein, 1995). Current evidence that the effects of predator abundance on prey behavior tends to be cumulative, but non-additive (Arditi and Ginzburg, 1989; Relyea, 2003; Brown and Kotler, 2004). For example when gerbils in an aviary were exposed to one owl, gerbil foraging decreased by 18%, however the addition of a second owl only decreased foraging by 5% further and a third owl reduced foraging only by 1% (St. Juliana *et al.*, 2011). This may be due to a "buy one get one half off" type of situation where the value of vigilance is greater when risk is higher. In other words, a gerbil being vigilant for one owl, will also detect a 2^{nd} or 3^{rd} owl. This type of benefit may apply to similar predators, but not predators that are very different. In contrast, with some predators, for example foxes, the gerbils tend to only "tune in" when the level reaches a certain threshold (unpublished work). In our experience the threshold was two foxes.

While titration of risk is a clear influencer of risk accumulation, rarely is it ecologically relevant to observe a single predator species as the only source of risk in an environment. Multi-predator impacts have been predominantly studies in consumptive studies in aquatic and arachnid model communities, and have shown to take two major forms: facilitation or interference between the two predators (Soluk and Collins, 1988; Berec, 2010).

Behaviorally, the impact of predators on each other depends on how prey perceive risk from these predators, thus impacting their habitat choices and the amount of time they spend foraging. Predator facilitation benefits predators in two major ways: 1. One predator drives prey towards the habitat preferred by another predator (Kotler *et al.*, 1992; Berec, 2010; Embar *et al.*, 2018) in a mutualistic benefit. 2. Presence of one predator overpowers the attention given to another predator thus making the prey more vulnerable to the other predator that is less focused on, *i.e.*, a commensal relationship between the predators (Bleicher et al. 2016). For many prey species, the risk from one predator requires much greater vigilance, or other types of defensive behavior, and thus marginalizes the attention (vigilance behavior) and energy expenditure (as decerned through time foraging) that can be allocated to avoidance of the lower risk predator (Kotler *et al.*, 2016; Bleicher *et al.*, 2019). It is important to be careful when interpreting the latter case when researching predation based on prey behavior. The predator that has a lower impact on the prey's fitness may seem to cause no risk, because it is masked by the predator that the prey perceives to be riskier.

Predator facilitation should narrow the availability of perceived safe habitat and narrow the
realized niche that the prey can occupy. This logically changes both community interactions, and the
populations dynamics of both predators and prey. The lower the abundance and distribution of prey, the
less fit both predators become in the long run. For example, the desert pocket mouse (*Cheatodipus*
penicillatus), from the Mojave Desert, favors a strategy of torpor and reduction of foraging to a minimal
sustainable level in response to combinations of owls and snakes in their environment (Sonny S. Bleicher
et al., 2020). When this occurs, neither predator is able to hunt effectively, despite the scenario suggesting
facilitation. On the flip side, when predators interfere with each other, the prey should effectively be able
to use more resources as a result of the conflict between the predators, reducing the overall risk (Embar *et*
al., 2014; Leo *et al.*, 2015). This interaction is typically observed when the two predators use the same
habitat, and drive the prey to use habitat that is less optimal for both predators. For example, when a
population of gerbils is exposed to both owls and foxes, they prefer to forage under the safety of bushes
where neither predator has easy access to them (Kotler *et al.*, 2010).

Different prey species will respond to the interaction of predators in different ways. This is a
consequence of evolutionary changes driven both by predator avoidance (Orizaola *et al.*, 2012; Wasko *et*
al., 2014) and by intraguild competition strategies pushing prey populations to evolve strategies that
balance both competition and predation costs (Rosenzweig, 1987; Abramsky *et al.*, 1996; Ben-Natan *et*
al., 2004; Kotler *et al.*, 2016; Bleicher *et al.*, 2019; Sonny S. Bleicher *et al.*, 2020; Sonny Shlomo
Bleicher *et al.*, 2020). Specifically, the foraging strategy of competing prey species will alter how
susceptible they may be to predation risk (Brown *et al.*, 1997; Kotler *et al.*, 2002). Examples include
locomotion type, with saltatory motion conveying advantages for predator avoidance (Biewener and
Blickhan, 1988; Longland and Price, 1991), sensory prowess as in the example of inner ear bullae
allowing for extra sensitive hearing and early escape (Webster and Strother, 1972), and even unique
special adaptations such as torpor as in the example of the abovementioned pocket mice (Sonny S.
Bleicher *et al.*, 2020). In this experiment, we test the responses of two competing gerbil species from the
Negev Desert in Israel, *Gerbillus andersoni allenbyi* (GA) and *Gerbillus pyramidum* (GP). We expect that
the same experiments would result in different outcomes from species applying different anti-predator
strategies. Specifically, the two gerbil species differ in their size with GA being significantly smaller than
GP (Abramsky *et al.*, 2001). The size difference between those two well-studied model species has given
rise to significant behavioral differences both in intra specific competition strategies and management of
predation risk. The variation between the species, interspecific competitive dynamics between them, and
intraspecific competition within the populations combine with the richness of the environment should also
alter the impact of combined predators. We expect the addition of predators to have a greater impact when
the competition for resources is lower (*cf.* Makin and Kotler 2020).

This study aims to compare the response of two gerbil species to titrations of predation risk
through the exposure to four different predators: two human-commensal species (a feral cat and a dog), a
barn owl, and an olfactory cue of a horned viper in the form of musk imbued sand. We intend to test each
predator alone and in combination with each of the other predators. Despite the differences between the
species' behavior, we predict that the general response to predator stressors will follow similar patterns,
as their anti-predator evasion strategies are similar enough. We predicted that the combination of any two
predators would result in additive responses of the behavior of the gerbils. Additionally, we predicted that
the closer the hunting strategy of two predators, the lower the added impact will be on gerbil
apprehension and foraging. *i.e.* adjustment in behavior applied towards one predator would automatically
apply to the similar predator. For example, the response of gerbils to a fox would be to move the majority
of their foraging under the safety of bushes. The logical response to a jackal in the environment would be
very similar, therefore the response would not vary much when both are present. However, when avoiding

snakes that ambush under bush cover, foraging in the open is the logical safe response (Embar *et al.*,
2014; Makin and Kotler, 2020). Therefore, combing a fox and a snake would increase the risk throughout
the environment, thus registering as a much greater threat than two ambush predators, or two active
searching predators. (Table 1). Where the risk from one predator type outweighs the risk from another,
the effect the effect may register as overshadowing of the impact of the lower-risk predator (Bleicher *et*
al., 2019).

Methods

Species

For this experiment we used two common gerbils from the Negev Desert of Israel: the greater
Egyptian gerbil *Gerbillus pyramidum* (GP), 40 g, and Allenby's gerbil *Gerbillus andersoni allenbyi*
(GA), 24 g (Goodfriend *et al.*, 1991). Both these gerbils are nocturnal, are commonly found on sandy
substrates such as sand dunes, and are considered granivores, despite foraging on available insects, and
green vegetation when available (Kotler *et al.*, 2001). Both gerbils have adaptations to reduce the risk of
predation, including saltatorial locomotion for enhanced escape abilities and auditory adaptations to
increase hearing acuity (Lay, 1972; Ellard and Eller, 2009; Kotler *et al.*, 2016). However, the species are
known to vary in their response to risk, with the larger species being more responsive to changes in both
patch quality and risk (Ben-Natan *et al.*, 2004; Emerson *et al.*, 2018). Similarly, the foraging patterns
differ between the species, the larger applies a cream-skimming pattern, only collecting easily found
grain in risky conditions, and the smaller applies a crumb-picking strategy in which they dig and search
for seeds more thoroughly (Brown *et al.*, 1997).

The gerbils were collected in the Mashabim Dunes (N 31_0'14.531'', E 34_44'47.31''), some
30km from the experimental enclosure. The barn owls were donated to the lab from a rescue center. For
this experiment we used *Cerastes gasperetti* snakes that were trapped in the vicinity of the experimental
enclosure, and housed in the zoological collection at Midreshet Ben-Gurion, Israel. All the animals were
held in climate-controlled animal-rooms at the Blaustein Institutes for Desert Research within 300 m of
the experimental vivarium. Both the feral dog (*Canis lupus familiaris*) and the feral cat (*Felis catus*) were
trapped (adopted) in the Sde Boker environment and were loaned for the experiment by their owners.

Experimental Set-Up

Dates and Animals:

The measurements were collected for each species in the absence of direct interspecific
competition. This allowed us to make a comparative study of the effects of predation risk in the exact
same setup in the absence of competition stressors. To best allow for comparison across gerbil species, we
populated the vivarium with roughly the same biomass of rodents corrected for metabolic rate, leading to:
24 Allenby's gerbils, 16 Greater Egyptian gerbils. Each experiment was conducted for 100 days starting
April 20[th] 2004 with GA being on one side of the vivarium and GP on the other. Based on subcutaneous
RFID PIT tags we tracked depredation.

The Enclosure

We conducted our experiments in a semi-natural, outdoor enclosure (vivarium 17 X 34 m)
located on the Sde Boker Campus of Ben-Gurion University, Blaustein Institutes for Desert Research,
Midreshet Ben-Gurion, Israel (N 30 51' 25.978'', E 34 46' 51.284''). The vivarium was divided into two
halves. It is covered with a wire net 4.5 m high such that aerial predators can be flown within. Our
vivarium mimics a dune habitat with a layer of 10 cm of sand covering a loess-clay subflooring with

escape buried to 1 m below ground. To replicate the sparse vegetation cover in semi-stabilized desert
dunes we created heterogeneity by adding 16 trellises in a randomized array with at least 2 m separation
from the next trellis. Each trellis (80 X 50 X 15 cm) at each station was topped with a pile of cut brush to
create a spherical ''bush'' approximately 1m in diameter. The environment beneath each trellis provides a
sheltered environment mimicking a bush microhabitat, while the space between trellises replicates the
open space occurring between vegetation clumps in the gerbils' natural environment.

Data Collection

We measured gerbil foraging using a randomized grid of 32 patches sieved nightly to obtain a
GUD measurement. Each patch (38 X 28 X 8 cm) held 3 liters of sand and was stocked each evening with
3 g of millet. At sunrise, each patch was sieved and remaining seeds weighed to 1/100 g and logged for
analysis. Daily, patches that were set up as covered patches (bush) were set outside the trellis for the
following collection night as open, and vice versa. The data sampling in this experiment combined two
related experiments, which used the same predators. This manuscript is a reflection of a meaningful
subset of data collected. This resulted in an uneven number of repetitions for each predator treatment we
used.

On nights when a single predator was used, the predator would be introduced to the vivarium
once every 60 minutes from sunset to sunrise. For the snake treatment, sand imbued with the natural musk
of the snakes from their enclosures was spread around the foraging stations on the same schedule. Based
on the lunar calendar we marked the moonphase associated with the night of collection. Of the 64
patches, 32 were set as "bottom" trays and 32 as "full" trays. For bottom trays, we mixed the seeds into
half of the sand, spread that sand over the tray, and then placed the remaining sand without seeds on top.
This created a more complex tray in which the seeds were in higher concentrations, but harder to access.
For full trays, we simply mixed all the seeds into all of the sand and spread that across the tray. Such
setups allow us to gauge apprehension by comparing changes in selectivity (Dall *et al.*, 2001; Kotler *et*
al., 2004). Moonlight ambient illumination and the temperature were recorded using digital meters.

Data analyses

197　　　We separated the cumulative data collected in the experiment into two data sets: single predator
198　data (including the control) and multi-predator data (using only nights with predator encounters every 60
199　minutes). We then ran analyses on the multi-species data set both as a single unit, and separately for each
200　of the gerbil species.

Single predator

202　　　We ran a series of t-tests for both gerbil species combined, and for each apart. These tests
203　compared whether the gerbils responded to the risk associated with each predator species as unique and
204　different from the same environment in absence of predators.

Multi-predator

To address differences between predators, we ran a General Linear Model (GLM) analysis for the
data collected from the experiment using both gerbil species. We increased the explanatory value of the
models, by adding interactions shown significant in a multi-factorial ANOVA that had very low
explanatory power, and made sure to include all biologically meaningful interactions. The factors we used
were gerbil species (GS), predator treatment (PT- type of predator present on the night of), moonphase,

resource distribution (RD; clumped or dispersed resources – at the bottom of a tray or mixed throughout),
and microhabitat (open and bush). For all variables and interactions found statistically significant in the
model (p-values below 0.05) we ran post-hoc Tukey Tests of Honest Significance Difference (HSD).

To ease in the interpretation of post-hoc analyses, we re-ran multi-factorial ANOVAs for each of
the gerbil species separately. Both these models used predator treatment, moonphase, Resource
distribution and microhabitat as independent variables and added covariates of ambient temperature and
moonlight intensity. Post-hoc Tukey HSD pairwise comparisons were calculated for all significant factors
and interactions.

Interpretation of data

To measure the cumulative responses of various predators, we have to interpret the effects of the
predators on each other, through the behavior of their prey as our experiments minimize the actual
depredation our predators were allowed to perform (through restricting the predators). We use quitting
harvest rates of the gerbils as gauged by giving-up densities (GUDs) in artificial patches to measure
perceived risk (Brown, 1988, 1999). We are attributed patches with greater foraging activity (lower
GUDs) as areas perceived as safer, and patches with higher GUDs as areas perceived as risky in
accordance with previous publications. The comparison of the mean GUDs for a single predator with the
combination including another predator will allow us to assess the impacts of the predators on each other.
By calculating a change in foraging dynamics dGUD (GUD of single – GUD of combined predators) we
can expect a few patterns (Table 2). Because of predator microhabitat preference, our prediction is that
facilitation will occur between the snake treatment and any of the other predators. This will result in a
negative dGUD as the overall perceived risk in the environment will be higher when the two predators are
present. Additionally, because we used musk in lieu of free-ranging snakes, it is likely that the response to
the risk from active predators will swamp the risk from the snakes entirely. Last, if the predators facilitate
each other, then the overall dGUD should be zero or positive, as the foraging in the entire experimental
arena should not decline as it simply equalizes between the microhabitats (Embar *et al.*, 2014).

Predator housing and control:

Efforts were made to reduce predator strikes on gerbils: owls were flown tethered to a rope and
were fed prior to the experiment, the muzzled dog ran free in the enclosure, the cat was on a harness and
tether but allowed to roam freely, the tether was used to prevent the cat from making fatal strikes. As
snakes were hard to control, sand imbued with their musk from their terraria was used as an olfactory cue
instead.

Results:

In a simple t-test, all predators were perceived as giving rise to greater risks than the control, with
exception to risk posed by owls. Allenby's gerbils (GA) categorically associated the risk from all
treatments as greater than the control, whereas the Greater Egyptian Sand Gerbils (GP) did not
differentiate the snake treatment from the control, and associated the owl treatment as lower risk than the
control (Fig 1, Table 3). None of the treatments were perceived as a very high risk, as even the highest
giving-up densities (means) were at $1.71g\pm0.03g$, meaning that more than one third of resources were
consumed in patches on a nightly average. While low, the variation seen here does not stray from
observations collected in previous experiments, and is in line with natural occurring intra-species
variation.

The GLM comparing the data collected for both gerbil species resulted in three significant single-
factors and seven two-way interactions (Table 5). The two more multi-factorial ANOVAs for GA and GP
separately found predator treatment, moonphase and microhabitat, and both covariates significant as well
as the interactions of PT X moonphase and RD and moonphase. The three-way interaction yielded
significant but hard to interpret results (SI 1,2). GA decreased foraging with increased temperature, while
GP increased foraging as temperatures rose. No clear discernible patterns emerge for the moon
illumination (SI 3). Regression analyses were non-significant and relatively low correlations were
observed with Pearson Correlation Indices for moonlight illumination were -0.1 and 0.0 for GP and GA
respectively. For temperature the correlation indices were 0.034 and 0.08 for GP and GA respectively.

GP foraged to lower GUDs than GA, with means being 1.694 ± 0.02 and 1.489 ± 0.02 g
respectively. Both gerbils perceived the bush microhabitat as safer (lower GUDs). The means were
1.465 ± 0.016 and 1.718 ± 0.017 g for bush and open microhabitats respectively. Gerbil response to
moonphase was not uniform for both species. GA foraged less on the full and waxing periods of the
month, and had lower GUDs on the waning and new moons. The mean GUDs were 1.709 ± 0.033,
1.751 ± 0.030, 1.735 ± 0.033, 1.521 ± 0.033 g for the new, waxing, full, and waning moonphases
respectively. GP foraged the least on the new and waning moonphases with GUDs being 1.521 ± 0.033,
1.485 ± 0.030, 1.418 ± 0.033, 1.553 ± 0.033 g for the new, waxing, full, and waning moonphases
respectively.

Predator type (PT) yielded significant patterns in the two gerbils combined dataset, and a less
impactful relationship for each separately (Fig 2). Tukey HSD tests revealed that GA response to predator
treatments was a lot less definitive than with GP (Figs 2B, and 2C respectively, SI 4,5). Neither the RD,
nor the interaction of moonphase with PT yielded few clear patterns of biological significance.

Overall, the impact of the snake treatment was negligible for both species suggesting that the
gerbils were likely ignoring the olfactory cues in presence of all live predators except for the owl (Fig 3).
For GA, the exposure to the dog resulted in a cumulative trend that was not statistically significant. The
owl treatment was always replaced by the risk of the dog, leading to higher GUDs when the dog was
present. The risk from the dog and cat treatments overall caused the largest decrease in foraging from all
the predators, but were comparable to each other. The dog treatments resulted in mean GUDs of
1.61 ± 0.037, 1.681 ± 0.052, 1.539 ± 0.052 g for both gerbils, GA and GP respectively. The cat, resulted in
GUDs of 1.685 ± 0.037, 1.720 ± 0.052 and 1.650 ± 0.052 g respectively. To better understand the predator
interactions however, we examined the interaction of predator treatment and microhabitat, but only for
both species combined, as the three-way interaction did not prove to be significant (Table 4, Fig 3).

The two ground foraging predators seemed to have elicited the largest "fear" response in the
gerbils, corresponding with highest mean GUDs. Between the two predators, the cat was perceived as a
much greater threat than the dog (Fig 3 A&B). The change in foraging in the open microhabitat decreased
by 10.8% of the total available food from the dog treatment to the dog and cat combined. The decrease
was only 4.6% from the cat to the cat + dog combination. Neither the snake treatment alone, nor owl
treatment were substantially different from the control in this experiment, in neither the bush or the open
microhabitat (Fig 3 C&D). For the owl, the open microhabitat became significantly riskier only in
combination with the dog and the snake, but not with the cat (an anomalous result). Last, the snake
treatment resulted in very little differences from the control. However, when combined with the dog, both
microhabitats were equally risky pointing to facilitation between the two predators.

Discussion

This experiment highlights and strengthens a few previous finds, namely that not all predators
exude the same intensity of predation risk on foragers (Bouskila 2001; Bleicher et al. 2016, 2018a), and
that different forager species- based on their unique adaptive strategies- do not respond equally to the
same risks (Burt P Kotler *et al.*, 1993; Burt Phillip Kotler *et al.*, 1993). Meanwhile, our data also builds
on previous findings that suggests that the intensity of a predator cue affects the response by the prey
(Sonny S. Bleicher *et al.*, 2018), and that the presence of various predators simultaneously have
cumulative impacts and not substitutive effects. However, the perceived risk that a predator exudes on the
prey will dictate whether the effect of the additional risk factor is negligible or cumulative.

Foraging strategy matters

Stronger interference competitors that employ a cream-skimming foraging strategy appear to be
more attuned to the specifics of risk in their environment. GPs were much more specific in their
management of light (moonphase) and predator activity than the GA were. This suggests that when
foragers have the ability to exclude their competition, they will use resources in a more discerning manner
than a lesser competitor (Abramsky *et al.*, 1996). A number of hypotheses have been generated over time
that explain why the larger competitor tends to "give-up" sooner than the smaller one. The leading ones
include that the larger prey have greater value to the predator and thus draw greater   focus from the
predator (Porter *et al.*, 2000); another is that larger foragers have higher energetic foraging costs due to
the larger metabolic costs associated with greater body mass (Brown and Kotler, 2004). Regardless of
explanation, the evidence collected herein supports this observation. The consequence of the sensitivity of
the larger competitor is that it "opens up" time, space, and resources for the lesser competitor who is less
discerning (Ziv *et al.*, 1993; Kotler and Brown, 1999).

The observations relating to the smaller competitor, GA, suggest a different pattern of behavior
than its larger counterpart. In congruence with former observations, the GA accept what they must as the
weaker competitor. They take more risks (Brown and Kotler, 2004), utilize suboptimal times (Burt P
Kotler *et al.*, 1993) and suboptimal substrates (Ziv *et al.*, 1993; Kotler *et al.*, 2001). Last, they forage in
patches that have much lower energetic value (Kotler et al. 1993b, Ben-Natan et al. 2004). The two major
observations supporting this dynamic in our data were a hyper focus on the greatest risk predators in the
experiment, where they foraged more, and took risks with all predators except the highest risk (dog or
cat). Additionally, the support of the suboptimal times, the GA preferred the "in-between" times when the
moon was waning and waxing especially at times that the risk was greatest (Dixon, 2017).

Predator treatments

The patterns of facilitation, interference and cumulative risks derived from the combinations of
predators significantly deviated from our expectations. The response to olfactory cues were smaller than
the responses to live predators, and even a live predator did not always differ significantly from the
control treatment (as seen in the example of the owl). Below we aim to break down the possible drivers of
the interactions we observed.

Predation risk administration

It is not surprising that a predator cue in absence of the actual predator, was perceived as a lower
risk than live predators. Many studies have tested the use of olfactory cues of predators as deterrents to
control agricultural damage and found conflicting outcomes. An example of gray kangaroos exposed to
dingo feces showed that the prey changed their behavior in the presence of predator feces compared to the
control and sham (sheep feces) treatments, however vigilance rates remained low (Mella *et al.*, 2014).
Wallabies did not respond to predator urine (Blumstein *et al.*, 2002). Additionally, some studies even

suggest that some prey will be attracted to predator olfactory cues, as a mechanism to take advantage of
resources with lower competition (Parsons *et al.*, 2018). Moreover, the information and prior experiences
the foragers have greatly alters the meaning they draw from an olfactory cue. In an experiment with bank
voles and weasels, voles interpreted a sham control (rabbit bedding) and weasel bedding as equally
suspicious, however after exposure to a caged weasel the voles interpreted the weasel odor to be a greater
threat (Bleicher *et al.*, 2018).

The gerbil responses to the olfactory cues for the presence of snakes based on musk-imbued sand
do not support the view that gerbils would interpret the scent as a sign of predation risk. However, the
gerbils did show an interaction between these cues and the presence of other predators, namely owl and
dog. In addition, the responses were much greater for the GPs than for the GA, evidence supporting the
above claims on variation between the species and their competitive foraging strategies.

<u>Hunting strategy matters and impacts prey's risk perception</u>.

Foraging has been shown to decrease based on various predator related characteristics, namely
the lethality of the predator, but potentially more importantly, based on perceived risk which at times may
veer from the actual lethality of the predator (Vance-Chalcraft *et al.*, 2004; Laundré *et al.*, 2009). Thus,
as a generalization, prey often overestimate the risk posed by a predator, as the cost of under-estimating
the risk is significantly greater than the opposite (Bouskila and Blumstein, 1992; Abrams, 1994; Bouskila
et al., 1995; Martín *et al.*, 2009). Additionally, the information a prey species has of their predator's
behavior, innate or learned, will change the behavioral tactics they will employ to avoid that risk. While
the most commonly reported is a shift in microhabitat use, the entire spatial and temporal distribution may
change based on the type of predator the foragers interact with. In an experiment with Saharan Horned
Vipers (*Cerastes cerastes*), Sidewinder rattlesnakes (*Crotalus cerastes*) & barn owls, conducted in the
same vivarium in 2010 and 2012 for GA and GP respectively, both of these gerbils species reassessed the
risk distribution in the enclosure, and refocused their anti-predator space-use to the predator they
perceived as more lethal, in that example that was the owl (Bleicher *et al.*, 2018).

Our observations in this experiment suggest a general agreement between the two foragers. Both
the gerbils foraged the least in the presence of the active ground hunting predators, followed by the owl
and the snake musk having a significantly lower impact on their behavior. Similarly, the cat was
perceived as a greater risk than the dog as evident by the significant decrease in foraging when the
combination of the cat and dog are compared to the isolated dog treatment. However, when comparing
the combined predators with an isolated cat, the pattern is cumulative but not significant.

<u>Predator facilitation, interference, and cumulative effects</u>.

The interactions between two predator treatments overwhelmingly showed cumulative effects.
Where the interaction was not significant, this can be interpreted as additive interactions. Where the
interaction was less than additive (significant positive dGUD) we can interpret the interaction as
interference. When the interaction was more than additive (significant negative dGUD) we interpret the
result to be facilitation. The gerbils of both species foraged less in combinations of predators over the
control and over each predator alone. However, the specific interactions do shed light on a couple of
patterns. Where interference between predators was observed, as measured through increased foraging in
the combined treatment, it was never statistically significant, and thus the interpretation results in a
conclusion that these are all additive interactions. Additive patterns were more likely in the GP population
than in GA and were more likely in predators of lower lethality (owl and snake) and especially when
these predators hunt using different strategies as in the snake and the cat (present in both species). A
plausible explanation for why these appear to exist is that different parts of the enclosure are associated

with the risk of specific predators. For example, in an experiment conducted in 2011-2013 in the same
enclosure, a populations of Merriam's kangaroo rats (*Dipodomys merriami*) and GAs identified areas of
risk of predators they prioritize (Bleicher *et al.*, 2019) and changed their geographic distribution within
the enclosure. The kangaroo rats identified the locations where snakes were present and depleted the
available forage in other locations, meanwhile GAs avoided a transact of land corresponding to the flight
path between two perches favored by the barn owls in the experiment, but were willing to risk the space
where snakes were present. Having the combination of multiple predators here, drives the gerbils to
search for resources in environment they did not previously forage, and thus increasing their harvest
intake to compensate for increased movement activity (Makin and Kotler, 2020).

This theory doesn't hold up when we look at the microhabitat use, perhaps because of the lack of
significance of the three-way interaction of gerbil species, predator treatment and microhabitat. Therefore,
we were only observing interactions of the combined data set for both species. Here, the only example of
facilitation was not between the cat and snake as we had initially expected, but between the dog and the
snake. In that treatment combination the gerbils equalized the resource use between the microhabitats. To
our best interpretation, this comes from the general behavioral pattern both gerbil species were observed
to follow where they shift their focus to the greatest risk in the environment (Bleicher *et al.*, 2019). While
all the other interactions appear to be facilitative and cumulative with the foraging decreasing in the open
microhabitat only, it would be very difficult to tease out which of the two each interaction represents.
However, the same pattern of shift of attention leads us to assume that most of the interactions are
cumulative.

<u>The anthropogenic human-commensal predator</u>

Our use of feral, human commensal species of predators allow us to comment on an unexpected
aspect of conservation importance. Significant numbers of papers focus on the impact that feral cats
specifically (Dickman, 1996; Moseby *et al.*, 2015; Hardman *et al.*, 2016; Doherty *et al.*, 2017) and human
commensal predators in general (Labra and Leonard, 1999; Avilés-Rodríguez and Kolbe, 2019; Zuñiga-
Palacios *et al.*, 2021) have on biological communities globally. Our observations were both encouraging
and support the overall fear of the exploding populations of feral cats. The encouragement we observe
comes from the fact that the gerbils clearly understood the cat to be the greatest threat to them. In
addition, the gerbils interpreted the risk from the cats and dogs correctly and responded by shifting their
microhabitat preference. Thus the overall impact these feral predators may have on their populations is
curbed in comparison to the impacts they have on naïve prey on islands and the Australian continent
(Banks and Dickman, 2007; Carthey and Banks, 2014).

In addition, the impact the dog had on the foraging of the gerbils surprised us. Previous
experiments suggested that the aerial predators, specifically the barn owl tends to be the focus of the anti-
predator behavior of the gerbils (Kotler *et al.*, 2010). The only case in a 30 year research program
studying the interactions of gerbils with their predators that the ground foraging predator (a fox) was
identified as a greater risk, was where a young pup was introduced (Bleicher et al. 2016). That fox was
highly energetic and dug up all the rodent borrows in the vivarium. The fact that the dog, despite being
perceived as a lower risk than the cat, still resulted in greater apprehension by the gerbils, suggests that
human commensal predators have a significant potential to impact community-wide interactions around
the globe.

<u>Conclusion</u>

The experiment was designed to tease out the interactions of various predator types on the
foraging behavior of small mammals. Through the behavior of the gerbils, the facilitation and

interreference between the predators was tested. The interactions between predators proved to be predominantly cumulative. Facilitation was found to be indisputable between lower-lethality snake olfactory cues and the dog. The impacts of ground foraging, human-commensal species was shown to draw the attention of the gerbils over predators that naturally occur in their environment, such as a barn owl and a horned viper. The tendency of the gerbils to focus their apprehension on the greatest risk in the environment supports the use of single predator type titrations favored in behavioral common-garden experiments.

Acknowledgements:

The research was funded by a United States–Israel Binational Science Foundation grant #1999–109 to BPK and JSB. We thank the Midreshet Ben Gurion Zoological Collection Reptile Collection for providing snake musk imbued bedding used in these experiments.

References

Abrams PA (1994) Should Prey Overestimate the Risk of Predation? Am Nat 144:317–328

Abramsky Z, Rosenzweig ML, Subach A (2002) The costs of apprehensive foraging. Ecology 83:1330–1340

Abramsky Z, Rosenzweig ML, Subach A (2001) The cost of interspecific competition in two gerbil species. J Anim Ecol 70:561–567

Abramsky Z, Strauss E, Subach A, et al (1996) The effect of barn owls (*Tyto alba*) on the activity and microhabitat selection of *Gerbillus allenbyi* and *G. pyramidum*. Oecologia 105:313–319. https://doi.org/10.1007/BF00328733

Ale SB, Whelan CJ (2008) Reappraisal of the role of big, fierce predators! Biodivers Conserv 17:685–690. https://doi.org/10.1007/s10531-008-9324-5

Arditi R, Ginzburg LR (1989) Coupling in predator-prey dynamics: Ratio Dependence. J Theor Biol 139:311–326

Avilés-Rodríguez KJ, Kolbe JJ (2019) Escape in the city: urbanization alters the escape behavior of Anolis lizards. Urban Ecosyst 22:733–742. https://doi.org/10.1007/s11252-019-00845-x

Banks PB, Dickman CR (2007) Alien predation and the effects on multiple levels of prey naivite. Trends Ecol Evol 22:229–230. https://doi.org/10.1016/j.tree.2007.02.003

Ben-Natan G, Abramsky Z, Kotler BP, Brown JS (2004) Seeds redistribution in sand dunes : a basis for coexistence of two rodent species. Oikos 105:325–335. https://doi.org/10.1111/j.0030-1299.2004.12948.x

Berec L (2010) Impacts of foraging facilitation among predators on predator-prey dynamics. Bull Math Biol 72:94–121. https://doi.org/10.1007/s11538-009-9439-1

Beschta RL, Ripple WJ (2009) Large predators and trophic cascades in terrestrial ecosystems of the western United States. Biol Conserv 142:2401–2414. https://doi.org/10.1016/j.biocon.2009.06.015

Biewener A, Blickhan R (1988) Kangaroo rat locomotion: design for elastic energy storage or acceleration? J Exp Biol 140:243–55

Bleicher SS (2023) Multi-predator Common Garden Experiment with Two Gerbil Species. In: Mendeley Data, V1

Bleicher SS, Brown JS, Embar K, Kotler BP (2016) Novel predator recognition by Allenby's gerbil (*Gerbillus andersoni allenbyi*): do gerbils learn to respond to a snake that can "see" in the dark? Isr J Ecol Evol 62:178–185. https://doi.org/10.1080/15659801.2016.1176614

Bleicher SS, Kotler BP, Brown JS (2019) Comparing Plasticity of Response to Perceived Risk in the Textbook Example of Convergent Evolution of Desert Rodents and Their Predators; a Manipulative Study Employing the Landscape of Fear. Front Behav Neurosci 13:1–12. https://doi.org/10.3389/fnbeh.2019.00058

Bleicher SS, Kotler BP, Downs CJ, Brown JS (2020a) Playing to their evolutionary strengths; heteromyid rodents provide opposite snake evasion strategies in the face of known and novel snakes. J Arid Environ 173:104025. https://doi.org/10.1016/j.jaridenv.2019.104025

Bleicher SS, Kotler BP, Downs CJ, Brown JS (2020b) Intercontinental Test of Constraint-Breaking Adaptations; Testing Behavioural Plasticity in the Face of a Predator with Novel Hunting Strategies. J Anim Ecol 89:1837–1850. https://doi.org/10.1111/1365-2656.13234

Bleicher SS, Kotler BP, Shalev O, et al (2018a) Divergent behavior amid convergent evolution : A case of four desert rodents learning to respond to known and novel vipers. PLoS One 13:1–17. https://doi.org/10.1101/362202

Bleicher SS, Ylönen H, Käpylä T, Haapakoski M (2018b) Olfactory cues and the value of information : voles interpret cues based on recent predator encounters. Behav Ecol Sociobiol 72:187–199. https://doi.org/10.1007/s00265-018-2600-9

Blumstein DT, Mari M, Daniel JC, et al (2002) Olfactory predator recognition: wallabies may have to learn to be wary. Anim Conserv 5:87–93. https://doi.org/10.1017/S1367943002002123

Bouskila A (2001) A habitat selection game of interactions between rodents and their predators. Ann Zool Fenn 38:55–70

Bouskila A, Blumstein DT (1992) Rules of Thumb for Predation Hazard Assessment : Predictions from a Dynamic Model. Am Nat 139:161–176

Bouskila A, Blumstein DT, Mangel M (1995) Prey under stochastic condition should probably overstimate predation risk: A reply to Abrams. Am Nat 145:1015–1019

Brown JS (1988) Patch use as an indicator of habitat preference, predation risk, and competition. Behav Ecol Sociobiol 22:37–47. https://doi.org/10.1007/BF00395696

Brown JS (1999) Vigilance , patch use and habitat selection : Foraging under predation risk. Evol Ecol Res 1:49–71

Brown JS, Kotler BP (2004) Hazardous duty pay and the foraging cost of predation. Ecol Lett 7:999–1014. https://doi.org/10.1111/j.1461-0248.2004.00661.x

Brown JS, Kotler BP, Mitchell WA (1997) Competition between birds and mammals : A comparison of giving-up densities between crested larks and gerbils. Evol Ecol 11:757–771. https://doi.org/10.1023/A:1018442503955

Carthey AJR, Banks PB (2014) Naïveté in novel ecological interactions: lessons from theory and experimental evidence. Biol Rev Camb Philos Soc 89:932–49. https://doi.org/10.1111/brv.12087

Dall SRX, Kotler BP, Bouskila a (2001) Attention, "apprehension" and gerbils searching in patches. Ann Zool Fennici 38:15–23

Dickman CR (1996) Overview of the Impacts of Feral Cats on. Nature 92

Dixon AK (2017) Tradeoffs of food and safety in contrasting environments: From the deserts of the Mojave and the Negev to the coral reefs of Eilat. Ph.D. Thesis. Ben Gurion University of the Negev

Doherty TS, Dickman CR, Johnson CN, et al (2017) Impacts and management of feral cats Felis catus in Australia. Mamm Rev 47:83–97. https://doi.org/10.1111/mam.12080

Ellard CG, Eller MC (2009) Spatial cognition in the gerbil: Computing optimal escape routes from visual threats. Anim Cogn 12:333–345. https://doi.org/10.1007/s10071-008-0193-9

Embar K, Kotler BP, Bleicher SS, Brown JS (2018) Pit fights: Predators in evolutionarily independent communities. J Mammal 99:1183–1188. https://doi.org/10.1093/jmammal/gyy085

Embar K, Raveh A, Hoffmann I, Kotler BP (2014) Predator facilitation or interference: a game of vipers and owls. Oecologia 174:1301–9. https://doi.org/10.1007/s00442-013-2760-2

Emerson SE, Kotler BP, Sargunaraj F (2018) Foraging efficiency in the face of predation risk: A comparative study of desert rodents. Evol Ecol Res 19:61–70

Goodfriend W, Ward D, Subach A (1991) Standard operative temperatures of two desert rodents, Gerbillus allenbyi and Gerbillus pyramidum: The effects of morphology, microhabitat and environmental factors. J Therm Biol 16:157–166. https://doi.org/10.1016/0306-4565(91)90038-4

Hardman B, Moro D, Calver M (2016) Direct evidence implicates feral cat predation as the primary cause of failure of a mammal reintroduction programme. Ecol Manag Restor 17:152–158. https://doi.org/10.1111/emr.12210

Kotler BP, Blaustein L (1995) Titrating Food and Safety in a Heterogeneous Environment : When Are the Risky and Safe Patches of Equal Value ? Oikos 74:251–258

Kotler BP, Blaustein L, Brown JS (1992) Predator Facilitation: the combined effect of snakes and owls on the foraging behavior of gerbils. Ann Zool Fenn 29:199–206

Kotler BP, Brown JS (1999) Mechanisms of coexistence of optimal foragers as determinants of local Abundances and distributions of desert granivores. J Mammal 80:361–374

Kotler BP, Brown JS, Bleicher SS, Embar K (2016) Intercontinental-wide consequences of compromise-breaking adaptations: the case of desert rodents. Isr J Ecol Evol 62:186–195. https://doi.org/10.1080/15659801.2015.1125832

Kotler BP, Brown JS, Bouskila A, et al (2004) Foraging games between gerbils and their predators: seasonal changes in schedules of activity and apprehention. Isr J Ecol Evol 50:255–271

Kotler BP, Brown JS, Dall SRX, et al (2002) Foraging games between gerbils and their predators : temporal dynamics of resource depletion and apprehension in gerbils. Evol Ecol Res 4:495–518

Kotler BP, Brown JS, Mitchell WA (1993a) Environmental factors affecting patch use in two species of gerbelline rodents. J Mammal 74:614–620. https://doi.org/10.2307/1382281

Kotler BP, Brown JS, Mukherjee S, et al (2010) Moonlight avoidance in gerbils reveals a sophisticated interplay among time allocation , vigilance and state-dependent foraging. Proc R Soc B 277:1469–1474. https://doi.org/10.1098/rspb.2009.2036

Kotler BP, Brown JS, Oldfield A, et al (2001) Foraging substrate and escape substrate: patch use by three species of gerbils. Ecology 82:1781–1790. https://doi.org/10.1890/0012-9658(2001)082[1781:FSAESP]2.0.CO;2

Kotler BP, Brown JS, Subach A (1993b) temporal foragers : of optimal coexistence of species

mechanisms of sand dune gerbils by two species partitioning. Oikos 548–556

Kotler BP, Holt RD (1989) Predation and Competition: The Interaction of Two Types of Species. Oikos
54:256–260

Labra A, Leonard R (1999) Intraspecific Variation in Antipredator Responses of Three Species of Lizards
(*Liolaemus*): Possible Effects of Human Presence. J Herpetol 33:441–448

Laundré JW, Calderas JMM, Hernández L (2009) Foraging in the landscape of fear , the predator's
dilemma: Where should I hunt ? Open Ecol J 2:1–6

Laundré JW, Hernandez L, Ripple WJ (2010) The landscape of fear: Ecological implications of being
afraid. Open Ecol J 3:1–7. https://doi.org/10.2174/1874213001003030001

Lay DM (1972) The anatomy, physiology, functional significance and evolution of specialized hearing
organs of gerbilline rodents. J Morphol 138:41–120. https://doi.org/10.1002/jmor.1051380103

Leo V, Reading RP, Letnic M (2015) Interference competition: odours of an apex predator and
conspecifics influence resource acquisition by red foxes. Oecologia 179:.
https://doi.org/10.1007/s00442-015-3423-2

Letnic M, Ritchie EG, Dickman CR (2012) Top predators as biodiversity regulators: The dingo Canis
lupus dingo as a case study. Biol Rev 87:390–413. https://doi.org/10.1111/j.1469-
185X.2011.00203.x

Lima SL (1998) Nonlethal effects in the ecology of predator-prey interactions. Bioscience 48:25–34.
https://doi.org/10.2307/1313225

Longland WS, Price M V (1991) Direct observations of owls and heteromyid rodents : Can predation risk
explain microhabitat use ? Ecology 72:2261–2273

Makin DF, Kotler BP (2020) How do Allenby's gerbils titrate risk and reward in response to different
predators ? Behav Ecol Sociobiol 74:1–10. https://doi.org/https://doi.org/10.1007/s00265-019-2785-
6 (2020)

Manlick PJ, Newsome SD (2021) Adaptive foraging in the Anthropocene: can individual diet
specialization compensate for biotic homogenization? Front Ecol Environ 19:510–518.
https://doi.org/10.1002/fee.2380

Martín J, López P, Polo V (2009) Temporal patterns of predation risk affect antipredator behaviour
allocationby Iberian rock lizards. Anim Behav 77:1261–1266.
https://doi.org/10.1016/j.anbehav.2009.02.004

Mella VSA, Cooper CE, Davies SJJF (2014) Behavioural responses of free-ranging western grey
kangaroos (*Macropus fuliginosus*) to olfactory cues of historical and recently introduced predators.
Austral Ecol Ecol 39:115–121. https://doi.org/10.1111/aec.12050

Moseby KE, Peacock DE, Read JL (2015) Catastrophic cat predation: A call for predator profiling in
wildlife protection programs. Biol Conserv 191:331–340.
https://doi.org/10.1016/j.biocon.2015.07.026

Orizaola G, Dahl E, Laurila A (2012) Reversibility of predator-induced plasticity and its effect at a life-
history switch point. Oikos 121:44–52. https://doi.org/10.1111/j.1600-0706.2011.19050.x

Orrock JL, Grabowski JH, Pantel JH, et al (2008) Consumptive and Nonconsumptive Effects of Predators
on Metacommunities of Competing Prey. Ecology 89:2426–2435

Paine RT (1963) Food web complexity and species diversity. Am Nat 100:65–75. https://doi.org/10.2307/2678832

Parsons MH, Apfelbach R, Banks PB, et al (2018) Biologically meaningful scents: a framework for understanding predator–prey research across disciplines. Biol Rev 93:98–114. https://doi.org/10.1111/brv.12334

Peckarsky BL, Abrams PA, Bolnick DI, et al (2008) Revisiting the classics: Considering nonconsumptive effects in textbook examples of predator prey Interactions. Ecology 89:2416–2425. https://doi.org/10.1890/07-1131.1

Porter WP, Budaraju S, Stewart WE, Ramankutty N (2000) Calculating climate effects on birds and mammals: Impacts on biodiversity, conservation, population parameters, and global community structure. Am Zool 40:597–630. https://doi.org/10.1093/icb/40.4.597

Relyea RA (2003) How Prey Respond to Combined Predators : A Review and an Empirical Test. Ecology 84:1827–1839

Rosenzweig ML (1987) Habitat selection as a source of biological diversity. Evol Ecol 1:315–330. https://doi.org/10.1007/BF02071556

Rosenzweig ML, MacArthur RH (1963) Graphical Representation and Stability Conditions of Predator-Prey Interactions. Am Nat 97:209–223

Schmitz OJ, Beckerman AP, Brien KMO (1997) Behaviorally Mediated Trophic Cascades : Effects of Predation Risk on Food Web Interactions. Ecology 78:1388–1399

Sih A, Bolnick DI, Luttbeg B, et al (2010) Predator-prey naïveté, antipredator behavior, and the ecology of predator invasions. Oikos 119:610–621. https://doi.org/10.1111/j.1600-0706.2009.18039.x

Soluk DA, Collins NC (1988) between fish and stoneflies : facilitation interactions Synergistic and interference among stream predators. Oikos 52:94–100

St. Juliana J, Kotler BP, Brown JS, et al (2011) The foraging response of gerbils to a gradient of owl numbers. Evol Ecol Res 13:869–878

Stephan JG, Stenberg JA, Björkman C (2017) Consumptive and non-consumptive effect ratios depend on interaction between plant quality and hunting behaviour of omnivorous predators Running head : Plant quality and hunting behaviour affecting prey. Ecol Evol 7:2327–2339. https://doi.org/10.1002/ece3.2828

Vance-Chalcraft HD, Soluk DA, Ozburn N (2004) Is prey predation risk influenced more by increasing predator density or predator species richness in stream enclosures? Oecologia 139:117–122. https://doi.org/10.1007/s00442-003-1484-0

Wasko DK, Bonilla F, Sasa M (2014) Behavioral responses to snake cues by three species of Neotropical rodents. J Zool 292:. https://doi.org/10.1111/jzo.12086

Webster DB, Strother WF (1972) Middle-ear mor- phology and auditory sensitivity of heteromyid rodents. Am Zool 12:727

Ziv Y, Abramsky Z, Kotler BP, Subach A (1993) Interference Competition and Temporal and Habitat Partitioning in Two Gerbil Species

Zuñiga-Palacios J, Zuria I, Castellanos I, et al (2021) What do we know (and need to know) about the role of urban habitats as ecological traps? Systematic review and meta-analysis. Sci Total Environ 780:1–11. https://doi.org/10.1016/j.scitotenv.2021.146559

Figure and Table Legends

Table 1. Predictive patterns of gerbil behavior in relation to four predators used in the experiment.

Table 2. Inference table associating gerbil foraging with likely predator relationship. Replacement is only likely with low lethality predators, or in this experiment where an olfactory-cue was used in lieu of using live snakes.

Table 3. T-test comparisons between the control and every single predator treatment, for the combined data set, and each of the species separately.

Table 4: ANOVA table for GLM using data from both species (N= 2752, multiple R= 0.51, and multiple R^2 = 0.26)

Table 5. Three-way interactions post hoc Tukey Tests of Honestly Significant Difference for A. Gerbillus andersoni allebyi, and B. Gerbillus pyramidum. Only significant interactions are listed, and * delineates one tailed distribution significance

Fig 1. Foraging activity of each of the gerbil species and both combined in isolated, single predator treatments. α dictates a significant pairwise comparison (between control and each predator) for the combined gerbil species, β for the smaller gerbil (GA), and γ for the larger species (GP) corresponding to Table 3. The patterns shown here suggest a greater selectivity and "evaluation" of risk performed by GP than GA.

Fig 2. Gerbil foraging adjustment in response to perceived risk from combinations of predators. The bars represent the mean change in foraging from a treatment with a single predator and a combination of that same predator with others. (A) combined species dataset, (B) GA, and (C) GP. If the adjustment is positive (*i.e.,* combined predator GUD was lower than the single predator) the trend is suggestive of predator interference. When the adjustment is negative, the gerbils foraged less with the combination of predators than the single predator, thus suggesting a pattern of facilitation or accumulation of risk. Bold bars delineate post hoc significant pairwise comparisons (Tukey HSD) between the single predator and the combination with another. The brackets delineate significant difference between predator combinations. Asterisks represent one tailed significance (P-values between 0.05-0.1). The combined data suggests opposite patterns than observed for each species independently.

Fig 3. Gerbil foraging (GUD±SE) as influenced by predator type and microhabitat (bush/cover in gray and open in white). Each panel focuses on a different predator and the combination with other predators where (A) focuses on the Dog (A), (B) the Cat, (C) the Owl, and (D) Snake musk. Brackets indicate significant Tukey Pairwise Comparisons.

660 **Tables**

661 Table 1. Predictive patterns of gerbil behavior in relation to four predators used in the experiment.

Predator	Safer microhabitat for gerbils	Predator hunting activity type	Potential interference	Expected facilitation	Potential replacement to
Cat	Bush	Ground foraging	Dog, Owl	Snake	Snake
Dog	Bush	Ground foraging	Cat, Owl	Snake	Snake
Snake	Open	Ambush		Cat, Dog, Owl	
Owl	Bush	Aerial	Cat, Dog	Snake	Snake

664 **Table 2.** Inference table associating gerbil foraging with likely predator relationship. Replacement is only
665 likely with low lethality predators, or in this experiment where an olfactory-cue was used in lieu of using
666 live snakes.

Main Predator	Predator Combination	Predicted Likely Outcomes		
		dGUD if Interference	dGUD if Facilitation/ Cumulative Effects	dGUD if Replacement (Likely focus predator)
Snake	Cat		= between microhabitats/- ↓	-↓ (Cat)
	Dog		= between microhabitats /- ↓	-↓ (Dog)
	Owl		= between microhabitats /- ↓	-↓ (Owl)
Owl	Snake		= between microhabitats /- ↓	+↑ (Owl)
	Dog	+ ↑		
	Cat	+ ↑		
Cat	Dog	+ ↑		
	Owl	+ ↑		
	Snake		= between microhabitats /- ↓	=/+↑ (Cat)
Dog	Cat	+ ↑		
	Owl	+ ↑		
	Snake		= between microhabitats / - ↓	=/+↑ (Dog)

669   Table 3. T-test comparisons between the control and every single predator treatment, for the combined
670   data set, and each of the species separately.

		Both Gerbils			GA			GP		
Group 1	Group 2	t-critical	df	P-value	t-critical	df	P-value	t-critical	df	P-value
Control	Snake	-4.03	1342	<0.001	-4.21	670	<0.001	-1.55	670	0.122
Control	Owl	-1.44	1342	0.162	-4.32	670	<0.001	+2.62	670	0.009
Control	Cat	-8.33	1278	<0.001	-6.33	638	<0.001	-5.74	638	<0.001
Control	Dog	-11.11	1278	<0.001	-6.74	638	<0.001	-9.16	638	<0.001

674   Table 4: ANOVA table for GLM using data from both species (N= 2752, multiple R= 0.51, and multiple
675      $R^2 = 0.26$)

Source	Type III SS	df	Mean Squares	F-Ratio	p-Value
Gerbil Species (GS)	27.483	1	27.483	79.288	0.000
Predator Treatment (PT)	50.070	9	5.563	16.050	0.000
Moonphase	1.139	3	0.380	1.095	0.350
Resource Distribution (Clumped/ Evenly Mixed) (RD)	0.031	1	0.031	0.090	0.764
Microhabitat	41.602	1	41.602	120.021	0.000
GS X PT	9.019	9	1.002	2.891	0.002
Moonphase X GS	6.922	3	2.307	6.656	0.000
Microhabitat X GS	22.670	1	22.670	65.401	0.000
Moonphase X PT	72.831	27	2.697	7.782	0.000
Microhabitat X PT	6.701	9	0.745	2.148	0.023
Microhabitat X RD	3.834	1	3.834	11.060	0.001
RD X GS	0.256	1	0.256	0.739	0.390
GS X PT X Moonphase	59.957	27	2.221	6.407	0.000
PT X Moonphase X RD	18.991	27	0.703	2.029	0.001
Error	911.965	2,631	0.347		

676
677

678 Table 5. Three-way interactions post hoc Tukey Tests of Honestly Significant Difference for A. Gerbillus
679 andersoni allebyi, and B. Gerbillus pyramidum. Only significant interactions are listed, and * delineates
680 one tailed distribution significance.

GA							
#	Fixed Variable	Category	Dynamic Variable	Relationship			P-value
1	Moonphase	Full	Predators	Dog	>	Snake	0.022
2	Moonphase	New	Predators	Dog+ Snake	>	Snake	<0.001
3	Moonphase	New	Predators	Owl	>	Snake	<0.001
4	Moonphase	Waning	Predators	Cat + Dog	>	Cat + Snake	<0.001
5	Moonphase	Waning	Predators	Dog + Owl	>	Owl	0.009
6	Moonphase	Waxing	Predators	Dog + Cat	>	Dog + Snake	0.009
7	Moonphase	Waxing	Predators	Dog + Cat	>	Dog	<0.001
8	Moonphase	Waxing	Predators	Dog + Cat	>	Dog + Snake	0.001
9	Predators	Cat + Snake	Moonphase	Full	>	Waning	<0.001
10	Predators	Dog	Moonphase	New	>	Full	0.023
11	Predators	Dog	Moonphase	Full	>	Waxing	<0.001
12	Predators	Dog+ Snake	Moonphase	New	>	Waxing	0.013
13	Predators	Owl	Moonphase	Waxing	>	New	0.059*
14	Predators	Cat + Owl	Moonphase	Waxing	>	Waning	0.031
GP							
#	Fixed Variable	Category	Dynamic Variable	Relationship			P-value
1	Moonphase	Full	Predators	Cat	>	Dog	0.058*
2	Moonphase	Full	Predators	Cat + Owl	>	Dog	0.044
3	Moonphase	Full	Predators	Cat + Dog	>	Cat + Snake	<0.001
4	Moonphase	Full	Predators	Cat + Dog	>	Dog	<0.001
5	Moonphase	Full	Predators	Owl	>	Dog	<0.001
6	Moonphase	New	Predators	Cat	>	Owl	<0.001
7	Moonphase	New	Predators	Cat + Owl	>	Owl	<0.001
8	Moonphase	New	Predators	Dog	>	Dog + Snake	0.048
9	Moonphase	New	Predators	Dog	>	Owl	<0.001

10	Moonphase	New	Predators	Dog + Cat	>	Dog + Snake	<0.001
11	Moonphase	New	Predators	Dog + Owl	>	Owl	<0.001
12	Moonphase	Waning	Predators	Cat	>	Snake	<0.001
13	Moonphase	Waning	Predators	Cat + Owl	>	Dog + Owl	0.008
14	Moonphase	Waning	Predators	Cat + Snake	>	Snake	<0.001
15	Moonphase	Waxing	Predators	Dog + Owl	>	Dog + Cat	0.022
16	Moonphase	Waxing	Predators	Dog	>	Owl	<0.001
17	Moonphase	Waxing	Predators	Dog	>	Snake	<0.001
18	Predators	Cat + Snake	Moonphase	Waning	>	Full	<0.001
19	Predators	Dog	Moonphase	New	>	Full	<0.001
20	Predators	Dog	Moonphase	Waxing	>	Full	<0.001
21	Predators	Dog+ Snake	Moonphase	Waxing	>	New	0.046

List of Appendices:

SI 1. Multi-way ANOVA for GA, including post-hoc Tukey tests for all significant variables and
interactions.

SI 2. Multi-way ANOVA for GP, including post-hoc Tukey tests for all significant variables and
interactions.

SI 3. Pearson Correlation matrices for the covariates against GUD for each of the gerbil species
separately.

SI 4. Response of combined gerbil species to predator encounters.

SI 5. Response of gerbils to predator treatments as affected by moonphase.

SI 6. Raw-data file used for analyses.