1
2
3
4   # Statistical learning dynamically shapes auditory perception

Sahil Luthra[1,a], Austin Luor[1,b], Adam T. Tierney[c], Frederic Dick[d], Lori L. Holt[b]

13
14
[a] Department of Psychology, Carnegie Mellon University, Pittsburgh, PA 15213
[b] Department of Psychology & Center for Perceptual Systems, The University of Texas at
Austin, Austin, TX 78712
[c] Department of Psychological Sciences, Birkbeck College, University of London, United
Kingdom WC1E 7HX
[d] Experimental Psychology, University College London, United Kingdom WC1E 6BT
21
22   [1] Equal contributions
23
24
25
Corresponding Author:
Professor Lori L. Holt
Department of Psychology & Center for Perceptual Systems
The University of Texas at Austin
Sarah M. & Charles E Seay Building
108 E Dean Keeton St, Austin, TX 78712
lori.holt@austin.utexas.edu
(512) 471-1157

34
35   **Preprint Server:**

37
38   **Keywords:**
39   Statistical learning, expectation, perception, auditory
40

Abstract

Humans and other animals use information about how likely it is for something to happen. The absolute and relative probability of an event influences a remarkable breadth of behaviors, from foraging for food to comprehending linguistic constructions -- even when these probabilities are learned implicitly. It is less clear how, and under what circumstances, statistical learning of simple probabilities might drive changes in perception and cognition. Here, across a series of 29 experiments, we probe listeners' sensitivity to task-irrelevant changes in the probability distribution of tones' acoustic frequency across tone-in-noise detection and tone duration decisions. We observe that the task-irrelevant frequency distribution influences the ability to detect a sound and the speed with which perceptual decisions about its duration are made. The shape of the probability distribution, its range, and a tone's relative position within that range impact observed patterns of suppression and enhancement of tone detection and decision making. Perceptual decisions are also modulated by a newly discovered perceptual bias, with lower frequencies in the distribution more often and more rapidly perceived as longer, and higher frequencies as shorter. Perception is sensitive to rapid distribution changes, but distributional learning from previous probability distributions also carries over. In fact, massed exposure to a single point along the dimension results in seemingly maladaptive loss of sensitivity - occurring entirely in the absence of feedback or reward – along a range of subsequently encountered frequencies. This points to a gain mechanism that suppresses sensitivity to regions along a perceptual dimension that are less likely to be encountered.

Significance Statement

Organisms as diverse as honeybees and humans pick up on probabilities in the world around them. People implicitly learn the likelihood of a color, price range, or even syntactic structure. How does statistical learning affect how we detect events and make decisions, especially when probabilities are completely irrelevant to the task at hand, and can change without warning? We find that people learn and track changes in perceptual probabilities irrelevant to a task and that this learning drives dynamic shifts in perception characterized by graded effects of enhancement – and primarily – suppression across acoustic frequency. This can result in a remarkably long-lived diminishment of perceptual sensitivity that seems maladaptive but may instead reflect use of likelihood to guide and sharpen perception.

Introduction

We implicitly pick up information about the probability of white versus red cars on the road, the spatial position of objects in a room, and how likely different sounds might be within a soundscape – for instance, hearing a cow moo in a barnyard versus a hospital. The detailed distributional structure of sensory input leads us to expect some events and to be surprised by others.

How does distributional learning influence perception? Many studies have focused on learning across probabilistic input, whereby organisms implicitly discover regularities across continuous input dimensions (e.g., Love, 2003; McMurray, Aslin, Toscano, 2009; Rosenthal, Fusi, & Hochstein, 2001). For example, unsupervised learning of clustering in speech sounds may scaffold infants' language acquisition (Werker, Yeung, & Yoshida, 2012; Cristià, 2011; Schatz et al., 2022). Other studies have focused on the outcome of distributional learning, manipulating stimulus probability to operationalize expectation and characterizing the influence of expectation

89   on perception and neural representation (Summerfield & de Lange, 2014; Summerfield & Egner,
90   2009).
91
Some theoretical accounts of these expectation-driven effects predict that perception will prioritize
high-probability expected input (consistent with Bayesian inference; de Lange et al., 2018).
Indeed, frequent, expected stimuli are better detected than rare stimuli (Pinto et al., 2015; Stein
& Peelen, 2015) and perceptual decisions about expected stimuli are speedier and more accurate
(Summerfield & de Lange, 2014; Summerfield & Egner, 2009). This enhanced perception might
be achieved via adjustments of weights on sensory channels that modulate gain, sharpening
representation of frequent relative to rare input. Alternately, perceptual enhancements might be
mediated by expectation-congruent memory representations (Summerfield & de Lange, 2014;
Kok et al., 2012). Neuroimaging studies have revealed that representation of expected stimuli is
enhanced via suppressed activity in voxels tuned away from expected stimuli (Kok et al., 2012;
Yon et al., 2018).
103
Other accounts conclude, instead, that distributional learning accentuates infrequent, unexpected
events (see Press et al., 2020). This prioritization is accomplished by suppressing expected input
(Blakemore et al., 1998; Kilteni & Ehrsson, 2017; Richeter et al., 2018; Meyer & Olson, 2011;
Kumar et al., 2017), leading to improved detection of rare stimuli (Milne et al., 2024). A third
account posits that expectation can lead to enhancement in some contexts and suppression in
others, with initial perceptual biases that tilt toward expected stimuli but can be cancelled out by
highly surprising input (Press et al., 2020). But complicating matters, probability distributions
experienced across a perceptual dimension may influence the bottom-up salience (Alink & Blank,
2021; Zivony & Eimer, 2024) or task relevance (Rungratsameetaweemana & Serences, 2019) of
a dimension, each with the potential to impact perception. In sum, there is no consensus about
how likelihood influences perception.
115
We propose that opposing theoretical perspectives may persist, at least in part, as a byproduct of
empirical focus on dichotomous frequent-versus-rare likelihoods that necessarily limit the
resolution with which the relationship between expectation and gain can be estimated. More
complex probability distributions sampled across a continuous perceptual dimension have the
potential to reveal granular, graded influences of expectation built from distributional learning
across probability.
122
Here, we shape expectation by sampling stimuli probabilistically across the primary
representational axis of the auditory system, acoustic frequency. Crucially, *acoustic frequency is*
task-irrelevant across our 29 studies. This decouples expectation from task utility, unveiling the
influence of distributional statistical learning across a task-irrelevant dimension on perception. We
test how this learning impacts perception across unimodal, bimodal, and equiprobable
distributions varying in statistical volatility and sampling density. Given evidence that distinct task
goals can influence the impact of short-term input regularities on perception (Fritsche, Mostert, &
de Lange, 2017), we use two tasks inspired by classic psychoacoustics literatures. One task
examines detection of near-threshold tones in continuous noise. In an influential study, Greenberg
and Larkin (1968) led listeners to expect a single constant-frequency tone to appear in noise but
tone frequency varied on a minority of trials. Detection accuracy was superior for the expected,
high-probability frequency with graded diminishment of detection accuracy as a function of
distance from the expected frequency. This graded sensitivity has been interpreted as a
frequency-selective attentional filter centered at the distribution mode (Scharf et al., 1987). Here,
detection accuracy of near-threshold tones in noise provides a graded metric of the perceptual

gain function arising from expectation built from distributional learning that supports directional
assessment of enhancement versus suppression.
140
Complementing this, the speed of tone duration decisions reveals the influence of distributional
learning across task-irrelevant acoustic frequency. Capitalizing on classic studies of Schröger and
Wolf (1998) developed as a model of auditory distraction, participants decide whether a sound is
"long" or "short" across tones with distinct and equiprobable durations. The tones' acoustic
frequency is task-irrelevant but carries a distributional regularity that impacts responses, with
slower duration decisions consistent with longer processing time for tones with low-probability
frequencies. Our parallel tests of tone-in-noise detection and tone duration decisions allow us to
examine the influence of distributional learning on perceptual tasks possessing distinct processing
demands.
150
To foreshadow, our distributional learning approach reveals influences of expectation that would
be unobservable across simple frequency-versus-rare likelihoods. We find that distributional
learning is not simply frequentist accumulation of events: equally likely events are differentially
perceived as a function of their position within a probability distribution and are influenced by filter
properties of both the sensory system and selective attention. We observe exquisite sensitivity to
distribution shifts and robust carryover of influence from previously experienced distributions at
multiple timescales. The influence of distributional learning on perception arises rapidly after a
shift in distributions, but our data also provide evidence for 'stubborn predictions' that linger (Yon,
de Lange, & Press, 2019). In fact, massed exposure to a single point along the frequency
dimension results in sustained diminishment of perceptual sensitivity of subsequently
encountered frequencies as a function of their distance from that single point. This would seem
maladaptive for perceptual precision but instead may reflect a mechanism that centers attentional
gain around the most likely region(s) along a perceptual dimension. Across studies, the impact of
distributional learning on perception is most consistent with a *sharpening* of representations
through *suppression* of processing units tuned to less-likely stimuli (e.g., Yon et al., 2018).
166
167 **Results**
168
169 Given the large number of experiments and results we report only test type, exact *p* values, and
170 power for each statistical test in the main text; all tests are corrected for multiple comparisons.
171 **Table S3** provides details on each reported analysis, including the relevant filename of the
172 subject-wise data and analysis files available at https://osf.io/xdgnw/.
173
174 **Distributional learning alters the detection of tones in noise**
175
176 We first ask whether distributional learning across a continuous sensory dimension affects the
177 most basic perceptual process: detection. Does the probability with which a sound occurs
178 influence the ability to hear it in noise?
179
180 For each detection study, individual detection thresholds are established immediately before the
181 experiment using three iterations of a standard staircase technique adapted for online testing,
182 Zhao et al., 2022; see **Materials and Methods**, **Fig 1a**, top). Thereafter listeners detect a tone
183 presented at threshold in continuous white noise within one of two intervals (**Fig 1a**, bottom).
184
Exp 1a establishes baseline detection accuracy when a single acoustic frequency (1000 Hz) is
100% probable. Exp 1b-f draw from a pool of five easily differentiable frequencies (800, 920,

1000, 1080, 1200 Hz) spaced ~13x the just-noticeable difference in frequency (Sek & Moore,
1995). In Exp 1b-d, one highly probable frequency comprises 75% of the 320 trials. The remaining
four tones each occur on just 6.25% of trials, creating a unimodal distribution across frequency.
Exp 1e has a bimodal probability distribution with 800 Hz and 1200 Hz frequencies each
presented on 40.6% of trials, with each other frequency presented on 6.25% of trials. Exp 1f is
identical to Exp 1e, except that the frequency for threshold estimation is 1080 Hz, rather than
1000 Hz as in Exp 1a-e. **Fig 1c** illustrates these distributions across the acoustic frequency
dimension.
195

Figure 1. Tasks and Distributional Regularities. (A) The tone-in-noise detection task involved two
phases: adaptive threshold estimation followed by the tone-in-noise detection task. Threshold estimation
trials began with continuous noise and a fixation cross (750 ms), after which a 1000-Hz tone was presented
with equal probability in one of three 250-ms detection windows (250 ms ISI), each indicated by a number
(1, 2, or 3) on the screen. A prompt 250-ms after the third detection window elicited participants' report of
the interval containing a tone (here, shown in the first interval). Tone intensity followed the 3-down, 1-up
procedure to estimate 79% accuracy (see **Methods and Materials**). The noise continued through the tone-
in-noise detection task, shown in the bottom of **(A)**. For each trial, 500 ms preceded a 250 ms fixation cross
and another 500 ms period. A 250-ms sinewave tone with intensity + 0.75 dB above the threshold estimated
in the adaptive thresholding task appeared in one of two 250-ms intervals (250 ms ISI), indicated by a "1"
and a "2" on the screen, respectively. Participants reported which interval contained the tone (here, shown
as interval 2). Tone frequency varied according to the distributions in **(C)**. **(B)** In the duration-decision task,
each trial involved a 1000-ms fixation followed by a 50 or 90 ms sine wave tone (equal probability) and
participants reported "long" or "short" with a button press. **(C)** Probability distributions for each experiment,
as a function of acoustic frequency. Blue distributions indicate tone-in-noise detection experiments. Orange
distributions indicate duration-decision experiments.
213
214
In Exp 1, stimulus probability strongly modulates tone detection in noise across Exp 1b-f with
better detection of high-probability frequencies at the distribution mode (**Fig 2a**; ANOVA, Freq x
Exp interaction, $p = 1.761 \times 10^{-31}$, $\eta^2 = 0.169$). Detection of only 1000 Hz (Exp 1a: 100% probability;
average accuracy 77.9%) does not differ from detection of the highest-probability frequency in
unimodal distributions (Exp 1b-d: 75% probability; average accuracy 75.3%; ANOVA, $p = 0.242$,
$\eta^2 = 0.012$). But detection of the modal frequencies in the bimodal distributions (40% probable) is
lower than when a single frequency is 80% or 100% probable (Exp 1e-f: 40.6% probability;
average accuracy 70.3%; ANOVA, $p = 0.006$, $\eta^2 = 0.053$ versus Exp 1b-d, ANOVA, $p = 0.003$, η^2
$= 0.101$ versus Exp 1a).
224

Figure 2. Distributional learning alters the detection of tones in noise. Each panel plots mean detection accuracy as a function of tones' acoustic frequency. The histograms to the left show distributional regularities for each experiment. Marker size scales with tone probability. Error bars are standard error of the mean. **(A)** Detection accuracy for a single-point distribution at 1000 Hz in Exp 1a approximates the expected detection accuracy estimated by the preceding threshold procedures and serves as a reference baseline for single frequency detection. For Exp 1b-d the distribution mode is detected best, with equivalently low-probability tones detected more poorly as a function of distance from the mode (see inset). **(B)** Bimodal distributions produce a 'dual spotlight' with detection accuracy best at the modes. Exp 1e-f differ only in the frequency used to estimate the threshold (1000 and 1080 Hz, respectively).

Proximity to the high probability tone also influences detection (**Fig 2a**). The low-probability
frequencies of Exp 1b-d share the same probability, yet those closer to a high-probability
frequency are better detected than those further away (ANOVA, p = 0.014, η^2 = 0.022). When the
high-probability frequency is centered in the range of frequencies defining the distribution, this
graded detection accuracy difference is symmetric (near > far to high-probability frequency,
ANOVA, p = 0.004, η^2 = 0.259). When the high-probability frequency is nearer to the distribution
edge (Exp 1b and Exp 1d), there is an asymmetric detection curve (ANOVA, p = 0.015, η^2 =
0.034): a sharp detection decrement toward the distribution edge is contrasted with a more
gradual 'ski slope' decrement toward the middle of the frequency range (see inset, **Fig 2a**). In
sum, equiprobable rare tones are detected more accurately if they are adjacent to the distribution
mode, but this advantage is modulated by the position of the high probability tone relative to the
range of the frequency distribution.

These results must be understood in the context of the operating characteristics of the auditory
system, for which there is a critical bandwidth of approximately 1/6 of an octave within which the
cochlea has limited ability to resolve stimuli (Fletcher, 1940). At 1000 Hz, the critical bandwidth is
~130 Hz. Thus, taking Exp 1c (1000 Hz mode) as an example, these sensory filter properties
likely contribute to the 'rescue' of detection of low-probability tones with frequencies situated close
to the distribution mode (920, 1080 Hz). Even so, the asymmetry of the shallow 'ski slope'
decrement toward the middle of the frequency range compared to the steep decrement toward
the edges of the range in Exp 1b and Exp 1d suggests that other mechanisms are also at play.
Where the high probability tone is situated within the range of experienced frequencies also has
an influence. If effects were driven solely by integration within an auditory critical band, we should
not observe such contextual influence.

More complex probability distributions also modulate detection (**Fig 2b**). Exp 1e shows that a
bimodal probability distribution with higher-probability (40.6%) frequencies at the edges of the
distribution (800 and 1200 Hz) induces a 'dual spotlight' across the frequency dimension.
Listeners detect the higher-probability tones more accurately than neighboring low-probability
tones (920 and 1080 Hz, linear contrast, Bonferroni-corrected p = 3.590 x 10^{-5}, Cohen's d = 0.757)
with a marginal difference compared to the middle 1000 Hz tone (linear contrast as above, p =
0.072, Cohen's d = 0.418).

Note that for Exp 1e, detection of 1000 Hz tones has a numerical (but not significant) detection
advantage compared to the other low-probability tones (**Fig 2b**). Two 'spotlights' centered at the
high-probability tone frequencies would yield a "V" rather than this observed "W" detection profile.
We speculated that the numerical detection advantage for 1000 Hz might arise from experience
with 1000 Hz in the 90-trial threshold-setting procedure that precedes Exp 1e. Exp 1f falsifies this
hypothesis. Changing the initial threshold-setting frequency to 1080 Hz elicits a similar "W" profile
and, importantly, replicates the overall 'dual spotlight' at 800 and 1200 Hz (linear contrast as
above, p = 4.62 x 10^{-8}, Cohen's d = 0.985, **Fig 2b**).

In summary, Exp 1 demonstrates that distributional learning modulates sound detection.
Replicating and extending classic studies in psychoacoustics (Greenberg & Larkin, 1968), tones
with higher-probability frequencies are better detected in noise than lower-probability frequencies.
The impact of distributional learning is graded across frequency, with better detection of low-
probability frequencies that lie closer to high-probability frequencies than equally improbable, but
more distant, frequencies. This effect is further influenced by the overall distributional context: the
protective effect of proximity to the high-probability tone depends on its position within the range

of encountered frequencies. Moreover, bimodal distributions with two higher-probability
frequencies at the edges of the frequency range elicit a 'dual spotlight'.
286
287 **Distributional learning across a task-irrelevant dimension impacts perceptual decisions**
288
289 Listeners track probabilities across acoustic frequency despite the irrelevance of frequency to the
290 Exp 1 detection task. Previous findings show that similar probability distribution manipulations
291 affect perceptual decision response times (Schröger & Wolff, 1998; Garrido, Dolan & Sahani,
292 2011). We next ask whether statistical learning over a probability distribution defined across task-
293 irrelevant *frequency* impacts the time course of decisions about a sound's *duration*.
294
In Exp 2a-c, participants report whether a tone is long or short, with 50 ms and 90 ms tones
presented equiprobably across 400 trials (**Fig 1b**; see **Materials and Methods**). Task-irrelevant
tone frequency varies across five frequencies (800-1200 Hz) in the manner of Exp 1 (**Fig 1c**).
There are four improbable tone frequencies (each 5% of trials), and a single probable frequency
(80% of 400 trials, Exp 2a: 920 Hz; Exp 2b, 1000 Hz; Exp 2c: 1080 Hz). In Exp 2d, 800 Hz and
1200 Hz are presented on 40.625% of trials with the other frequencies each presented on 6.25%
of trials to create a bimodal distribution (320 trials). In Exp 2e, the five tones are equiprobable
(20%) across the first half of the study and then switch to the bimodal distribution of Exp 2d (640
total trials).
304
Across Exp 2a-c, the probability of a tone's *frequency* significantly impacts the speed of *duration*
decisions (ANOVA, $p = 7.620 \times 10^{-7}$, $\eta^2 = 0.017$, **Fig 3a**). Response times (RTs) are slower for
tones with low, compared to high, probability frequencies (linear contrast, unpooled error, p
$=1.628 \times 10^{-17}$, Cohen's $d = 0.516$). Further, RTs for duration decisions to equiprobably rare
frequencies are graded as a function of their distance from the high-probability distribution mode.
Compared to RTs to the most probable frequency, those to the adjacent low-probability
frequencies are slower (Holm-corrected post hoc comparison, $p = 4.316 \times 10^{-12}$, Cohen's $d =$
0.387) and frequencies furthest away from the high-probability frequency are yet slower than the
frequencies nearer the high-probability one (Holm-corrected $p = 5.137 \times 10^{-6}$, Cohen's $d = 0.258$).
(These patterns hold true for each Exp 2a-c study, $p < .05$ Holm-corrected). This replicates and
extends classic observations from psychoacoustics (Schröger & Wolff, 1998) and mirrors the
graded influence on Exp 1 detection accuracy (**Fig 2a**).

Figure 3. Distributional learning across a task-irrelevant dimension impacts perceptual decisions.
Each panel plots mean response time as a function of tones' acoustic frequency. The histograms to the left
show distributional regularities for each experiment. Marker size scales with tone probability. Error bars are
standard error of the mean. **(A)** Response time to report tone duration is impacted by the probability of
tones' acoustic frequency across Exp 2a-c. The influence is graded, with faster decision times for
equivalently low-probability tones closer to the distribution mode (see inset). **(B)** Unlike the dual spotlight
for tone detection in Exp 1e-f, there is no significant response time difference for the two more probable
modes in Exp 2d, a consequence of a frequency-duration perceptual bias (see **Fig S1**). **(C)** Exp 2e
evaluated the frequency-duration bias across an equiprobable distribution in the first half of the study
(orange, dashed) with a switch to the bimodal distribution at study midpoint (yellow, solid). The bias is
largest at the edges of the distribution where it interacts with the bimodal distributional regularity (see **Fig
S1**).

However, unlike the dual spotlight for tone detection in Exp 1e-f, there is no significant RT
advantage for making duration decisions about the higher-probability 800 and 1200 Hz tones in
Exp 2d (**Fig 3b**; ANOVA effect of frequency, $p = 0.526$, $\eta^2 = 0.010$). To examine this more closely,
Exp 2e introduces a distribution change: five initially equiprobable (20%) frequencies (320 trials)
shift to mirror the Exp 2d bimodal distribution mid-study (320 trials; see **Fig 1c**). This allows us to
characterize potential frequency-duration interactions that may exist, independent of probability.
Indeed, and unexpectedly, when tone frequencies are equiprobable in the first half of trials,
decision RTs are *longer* for 800 Hz and 1200 Hz compared to other frequencies (ANOVA on
Frequency, $p = 0.031$, $\eta^2 = 0.026$; post-hoc linear contrast $p = 0.002$, Cohen's d = 0.205) (**Fig 3c**).

Investigating this reveals a *novel frequency-duration perceptual bias*: duration decisions for lower-
frequency tones (800, 920 Hz) are more accurate and faster for long (90 ms) compared to short
(50 ms) tones whereas those for the highest frequency tone (1200 Hz) are more accurate and
faster for short compared to long tones (**Fig S1**; ANOVA, Frequency x Duration interaction, RT:
$p = 0.003$, $\eta^2 = 0.029$; Accuracy (Acc): $p = 3.738 \times 10^{-5}$, $\eta^2 = 0.037$). This perceptual bias is
mirrored, but only qualitatively, in Exp 2d (**Fig S1**; $p > 0.05$, $\eta^2 = 0.010$, with lower frequencies
related to longer durations and higher frequencies with shorter durations). Notably, the bias is
largest at the edges of the frequency distribution (800 and 1200 Hz) where it interacts with the
bimodal distribution modes of Exp 2d-e, helping to explain why the dual spotlight observed in Exp
1e-f detection is not apparent in Exp 2d duration decisions. When we inspect the data from Exp
2a-c (**Fig S1**) we also observe the longer-lower/shorter-higher bias in the context of the unimodal
distributions (ANOVA, Frequency x Duration interaction, RT: $p = 3.968 \times 10^{-6}$, $\eta^2 = 0.020$; Acc: p
$= 0.003$, $\eta^2 = 0.020$). In other words, listeners found it easier to identify long durations when tones
were relatively lower in frequency; conversely, it was easier to identify short durations when the
sound was a relatively higher frequency tone. This impacted response time and interacted with
the probability manipulation.

In summary, distributional learning across a task-irrelevant dimension affects perceptual
decisions. The speed with which participants report the *duration* of a tone is impacted by the
probability of the tone's frequency. As with tone detection in noise in Exp 1a-f, learning across the
probability distribution produces a graded influence on perceptual decisions: decisions across
equivalently low-probability tones differ as a function of the tone's distance in frequency from a
high-probability tone. Moreover, Exp 2 demonstrates that seemingly intrinsic biases across
acoustic dimensions may influence and/or disguise the impact of short-term statistical input
regularities (for other examples see Roark & Holt, 2022; Bröker et al., 2024). These "intrinsic"
biases might arise from distributional learning across longer timescales or perhaps sensory

processing (see **Discussion**), and interact with short-term distributional regularities as shown in
Exp 2a-e.
368
369 **Perceptual sensitivity and decisions rapidly update in volatile statistical contexts**
370
Studies of statistical learning often investigate static distributions. But real-world environments
can be volatile: for example, listeners often encounter talkers speaking different accents with
different distributional regularities. The perceptual weight of different speech cues can rapidly alter
in response to shifts in distributional regularities (e.g., Hodson et al., 2023; Murphy et al., 2023).
It is not clear whether fundamental processes like detection and perceptual decisions are
modulated by statistical volatility across *task-irrelevant* sensory dimensions.
377
Here, across six studies, we examine distributions composed of two tones: one high probability
frequency and one low probability frequency (**Fig 1c**), akin to dichotomous probability distributions
often used in studies of expectation and attention (e.g., Zivony & Eimer, 2024). In Exp 3a-b
(detection) and Exp 4a-b (duration decision) we examine static two-frequency distributions to
assure that effects of distributional learning observed across 5-tone distributions in Exp 1 and Exp
2 hold even in the simplest 2-tone sensory context over 320 trials. Exp 3a and Exp 4a examine
detection and duration decisions, respectively, with 1000 Hz occurring across 75% of trials and
1155 Hz occurring over the remaining 25% of trials. Exp 3b and Exp 4b examine detection and
duration across the complementary probability distribution. In Exp 3c and Exp 4c, we model a
dynamic statistical context where these two-frequency distributions alternate every 160 trials.
Participants experience four 160-trial blocks, with 1000 Hz high-probability (75%) and 1155 Hz
low-probability (25%) in the first block, and probabilities alternating across frequencies in
subsequent blocks.
391

 Figure 4. Perceptual sensitivity and decisions rapidly update in volatile statistical contexts. For Exp

3a-c mean detection accuracy as a function of acoustic frequency is plotted in blue; for Exp 4a-c mean
response times in the duration-decision task are plotted in orange. Marker size scales with tone probability.
In **(A)** and **(B)** the insets show the probability distributions. In **(C)** and **(D)** color indicates the tone frequency
and marker size indicates its probability. Error bars are standard error of the mean. **(A)** Probability
distributions defined across just two acoustic frequencies impact tone detection, with more accurate
detection for high-probability tones in Exp 3a-b. **(B)** Two-tone distributions defined across task-irrelevant
acoustic frequency also impact the response time to make duration decisions, with slower decisions to low-
probability tones in Exp 4a-b. **(C)** As tone probability shifts every 160 trials across four blocks in Exp 3c,
detection is more accurate for the high-probability, compared to low-probability, tones. **(D)** Similarly, in Exp
4c, the speed of duration decisions mirrors volatile probability changes: lower probability tone frequencies
elicit slower decisions.

Across Exp 3a and Exp 3b, we find equal and opposite effects of frequency probability, with the
high probability tone detected on average ~6% more accurately than the low probability tone (**Fig
4a**; ANOVA, Freq x Prob interaction, $p = 3.361 \times 10^{-6}$, $\eta^2 = 0.073$). In Exp 4a and Exp 4b, RTs to
the high probability tone frequency are on average ~28 ms faster than those to the low-probability
frequency (**Fig 4b**, ANOVA, $p = 1.375 \times 10^{-6}$, $\eta^2 = 0.010$). We also observe the perceptual 'low-
frequency à long-duration / high frequency à short-duration' bias of Exp 2 even in this dichotomous
probability distribution, with faster RTs for long-low/short-high duration-to-frequency pairings
(ANOVA, Freq x Duration interaction, RT: $p = 9.34 \times 10^{-6}$, $\eta^2 = 0.013$; Acc: $p = 6.318 \times 10^{-5}$, $\eta^2 =$
0.023). In summary, a 2-tone frequency probability distribution affects tone in noise detection. It
also affects individuals' speed in making perceptual decisions across a different, task-relevant
input dimension, but this effect is modulated by pre-existing perceptual biases.
415
416   In the statistically volatile context established in Exp 3c, there is a detection advantage for the
417   more probable frequency, with significant 'flips' in detection accuracy due to short-term reversals
418   in tone probability for the first three blocks of Exp 3c (**Fig 4c**; ANOVA, Freq x Block interaction, p
419    $= 2.495 \times 10^{-5}$, $\eta^2 = 0.099$, each block $p < 0.05$). In the final block, there is no significant difference
420   in detection accuracy across frequencies.
421
Likewise, transient changes in probability distribution affect the efficiency of perceptual decisions
in Exp 4c (**Fig 4d**, ANOVA, Freq x Block interaction, $p = 5.253 \times 10^{-7}$, $\eta^2 = 0.040$). RTs are slowest
for the less probable frequency in all blocks (all $p < 0.05$ Bonferroni-corrected). Even in this
dynamic context we again observe the systematic frequency-duration perceptual bias discovered
in Exp 2 (ANOVA, Freq x Duration interaction, RT: $p = 0.019$, $\eta^2 = 0.039$; Acc: $p = 0.019$, $\eta^2 =$
0.077).
428
429   In summary, probability distributions defined across two acoustic frequencies elicit implicit
430   statistical learning that impacts perception. The influence is rapid: probability exerts its influence
431   across just 160 trials. As input statistics change, implicit statistical learning influences sound
432   detection and perceptual decision making.
433
434   **The influence of distributional learning is consistent with a gain mechanism exhibiting**
435   **hysteresis**
436
We observe strong influences of distributional learning across unimodal probability distributions
on detection accuracy and the speed of duration decisions (Exp 1 and Exp 2) that holds for
dichotomous probabilities and follows volatile statistics across an experiment (Exp 3 and Exp 4).
Here in Exp 5 (detection) and Exp 6 (duration decisions), we borrow from the distribution-switch

design established in Exp 2e (**Fig 1c**). This distribution manipulation enables us to investigate
how distributional learning influences detection and duration decisions across a changing
statistical context. Moreover, by establishing perception across equiprobable distributions as a
baseline, we reveal granular and graded changes in detection and decision making that emerge
as distributional learning builds expectations, including enhancement and suppression of
expected stimuli.

Figure 5. The influence of distributional learning is consistent with a gain mechanism exhibiting hysteresis. In Exp 5a-b mean detection accuracy as a function of acoustic frequency is plotted in blue; in Exp 6a-b mean response times in the duration-decision task are plotted in orange. The histograms to the left show distributional regularities for each experiment. Marker size scales with tone probability. In each panel, the darker color (dotted line) indicates behavior in the first half of the experiment; the lighter color (solid line) indicates behavior in the second half, when distributional regularities shift. Error bars are standard error of the mean. **(A)** Exp 5a establishes detection accuracy across a equiprobable distribution, then shifts to a unimodal distribution centered on 1000 Hz. Detection accuracy improves for the distribution mode with increased probability and decreases for frequencies with decreased probability. **(B)** Exp 5a switches from a unimodal distribution centered at 1000 Hz to an equiprobable distribution. Note the hysteresis at 1000 Hz, where detection remains elevated even into the second half of the study. **(C)** In Exp 6a, duration-decision times are flat with equiprobable frequencies in the first half. Introduction of a unimodal distribution centered at 1000 Hz leads to faster duration decisions at the mode. **(D)** In Exp 6b the unimodal distribution shifts to equiprobable at the study midpoint and response times shift substantially; note that this effect interacts with the frequency-duration bias identified in Exp 2.

With equiprobable frequencies in the first half of Exp 5a, detection accuracy is consistent across frequency (**Fig 5a**; overall ~65%, with unexpectedly better detection for 800 Hz, ANOVA, $p = 0.009$, $\eta^2 = 0.129$). In the second half of Exp 5a, probabilities shift to mirror Exp 1b (1000 Hz 75%; all others 6.25%). This shift drives changes in accuracy which differ across frequencies ($p = 8.511 \times 10^{-7}$, $\eta^2 = 0.142$). The 1000 Hz tones, which are now more probable, are better detected than they were in the first (equiprobable) half of Exp 5a ($p = 0.002$, Cohen's d = 0.648), whereas the frequencies nearest (marginal $p = 0.058$, Cohen's d = 0.554) and furthest ($p = 0.027$, Cohen's d = 0.727) from 1000 Hz, which are now less probable, are more poorly detected than they were in the first half of the study (all Bonferroni-corrected linear contrasts with pooled error).

In Exp 5b, we reverse distribution order. With a unimodal distribution centered on 1000 Hz in the first half of Exp 5b, detection generally resembles Exp 1c (**Fig 5b**), with better accuracy for high-probability 1000 Hz compared to low-probability frequencies (Bonferroni-corrected linear contrast, $p = 1.023 \times 10^{-7}$, Cohen's d = 1.255), but with only a numerical detection advantage for frequencies nearest (920 and 1080 Hz) versus furthest (800 and 1200 Hz) from the probable center frequency ($p = 0.262$, Cohen's d = 0.267, Bonferroni-corrected). When tone frequencies become equiprobable mid-study, again the probability shift drives differential changes in accuracy ($p = 1.815 \times 10^{-4}$, $\eta^2 = 0.056$). Here, the influence of the unimodal distribution carries over to confer a detection advantage to 1000 Hz, which was formerly highly probable, compared to other frequencies, which were formerly less probable (Bonferroni-corrected linear contrast, $p = 1.383 \times 10^{-4}$, Cohen's d = 0.577). Detection of 1000 Hz tones decreased in accuracy from the first to the second study half due to the probability shift ($p = 1.162 \times 10^{-5}$, Cohen's d = 0.496), but detection accuracy for the formerly low-probability tones did not change, despite a more than 3-fold probability increase ($p = 1$, Bonferroni corrected).

In sum, statistical learning across a unimodal distribution provokes a persistent effect on detection. For example, in Exp 5b, the initially highly probable 1000 Hz tone continued to be detected more accurately than other tones even after tone frequencies became equiprobable. Conversely, the tones adjacent 1000 Hz, which were initially relatively improbable, continued to be detected poorly even after the shift to the equiprobable distribution. Next, we use this distribution shift design to examine duration decisions.

Exp 6a begins with equiprobable frequencies and shifts mid-study to a unimodal distribution centered at 1000 Hz (80%, each other frequency 5%; **Fig 1c**). Exp 6b reverses this order. In the

first half of Exp 6a, RTs in the duration-decision task across equiprobable frequencies are similar
(**Fig 5c**, ANOVA, p = 0.163, η^2 = 0.018). When probabilities shift to a unimodal distribution
centered on 1000 Hz mid-study, RTs drop overall (ANOVA, p = 0.011, η^2 = 0.061). Although there
is a numerical 'V-shaped' RT advantage for the now-probable 1000 Hz compared to increasingly
more distant frequencies, this pattern does not differ significantly from the first half of the
experiment (ANOVA, p = 0.245, η^2 = 0.005).

In the first, unimodal probability half of Exp 6b, decision response times exhibit the "V" shape
around the high-probability 1000 Hz tone also observed in Exp 2b (ANOVA, effect of frequency,
p =6.847 x 10^{-8}, η^2 = 0.080, **Fig 5d**). Decisions about low-probability frequencies near to 1000 Hz
are only numerically slowed compared to 1000 Hz itself (this and following test Bonferroni-
corrected linear contrasts, p = 0.058, Cohen's d = 0.227) but faster than to those further away
from 1000 Hz (p = 0.014, Cohen's d = 0.299).

When all frequencies become equiprobable mid-study in Exp 6b, there is a change in the degree
to which frequency modulates decision RTs (ANOVA, p = 0.024, η^2 = 0.017), but the 1000 Hz
decision advantage persists in the second half (**Fig 5d**). Even though 1000 Hz is now 20%
probable, RTs are not significantly different than in the first experiment half when it was 80%
probable (linear contrast, p = 0.720, Cohen's d = 0.025). Like detection in Exp 5b, there is
carryover from experience with the unimodal distribution in the first half of the study, such that
decision RTs are still modulated by frequency (ANOVA, p = 8.306 x 10^{-5}, η^2 = 0.068). RTs to
report decisions for 1000 Hz continue to be significantly faster than for the now-equally-probable
far frequencies (p = 0.020, Cohen's d = 0.199), although not significantly faster than nearby
frequencies (p = 0.668, Cohen's d = 0.054). Finally, we again observe the duration-frequency bias
established in the prior duration-decision studies (ANOVA, Freq x Duration interaction, RT: p =
1.608 x 10^{-4}, η^2 = 0.013; Acc: p = 0.006, η^2 = 0.021).

In summary, the impact of distributional learning on both detection and perceptual decisions
emerges quickly and exhibits hysteresis, persisting even after the unimodal probability distribution
flattens so that tones are equiprobable.

The detailed shape of statistically-driven gain is modulated by range, distribution, and
sampling density

In Exp 7, we make a more in-depth exploration of how expectations built up from distributional
learning are impacted by statistical context, including frequency range and sampling density.
Across six tone-in-noise detection studies, Exp 7 provides detailed information about the shape
of the gain that emerges from distributional learning and how it evolves after an abrupt change in
distributional statistics. We use these within-experiment distributional changes to estimate the
emergence of enhancement and suppression of frequencies via distributional learning.

Exp 7a-f incorporate a mid-study change in distribution from equiprobable to unimodal or vice
versa. The studies vary the range and density of 7 tone frequencies that define the distributions
(**Fig 1c**) from *narrow* (Exp 7a,b; 5.5 semitone range), *intermediate* (Exp 7c,d; 9.47 semitones
range), to *wide* (Exp 7e,f; 11.36 semitone range). In each range, frequencies are symmetrically
arranged around 1000 Hz (like Exp 1c). As in prior studies, we group frequencies according to
their distance (near, middle, and far) from the center frequency, which changes from highly
probable to equiprobable or vice versa. In Exp 7a,c,e, the 7 frequencies are equiprobable (14.3%)

until the experiment mid-point when 1000 Hz tones comprise the majority (71.4%) of trials and
the other six tones are lower probability (4.8%). This order is reversed in Exp 7b,d,f. Below, we
first describe detection accuracy patterns separately for Exp 7a,c,e (equiprobable to unimodal)
and Exp 7b,d,f (unimodal to equiprobable), and then aggregate detection data across the
unimodal conditions from each experiment to maximize power to detect effects of statistical
context.
551
In Exp 7a,c,e, an equiprobable distribution precedes a switch to a unimodal distribution centered
on 1000 Hz (see **Fig 6a-c**). Across these three studies, detection accuracy in the equiprobable
first halves does not vary across frequency (ANOVA, p = 0.399, η^2 = 0.003), nor is it modulated
by the different frequency ranges across Exp 7a,c,e (ANOVA, p = 0.115, η^2 = 0.035), and there
is no interaction of frequency and range (p = 0.119, η^2 = 0.011). Average detection accuracy
across these equiprobable distributions is 64%, which does not differ significantly from that of the
5-frequency equiprobable distribution of Exp 5 (ANOVA, p = 0.219, η^2 = 0.039).
559
The introduction of the unimodal distribution differentially affects detection, depending on distance
of tones from 1000 Hz (ANOVA, p = 1.622 x 10^{-11}, η^2 = 0.030). When 1000 Hz shifts from
equiprobable (14.3%) to highly probable (71.4%), there is a small but reliable *increase* in detection
accuracy (ANOVA, p = 0.002, η^2 = 0.026). It is notable that this five-fold increase in probability
(and ~16-fold increase in relative probability compared to low-probability frequencies) only
confers an average 3.7% detection improvement. This mild enhancement is not significantly
influenced by the range of frequencies (ANOVA, p = 0.365, η^2 = 0.005). Examining the off-center
frequencies that drop in probability (14.3% to 4.8%) upon introduction of a unimodal distribution,
we observe a significant *decrease* in detection accuracy of 4.7% (ANOVA, p = 4.798 x 10^{-9}, η^2 =
0.051), the magnitude of which does not differ significantly across range (ANOVA, p = 0.337, η^2
= 0.003). In brief, when probabilities switch from equiprobable to unimodal we observe a modest
increase in detection accuracy for the center frequency that increased in probability and a
decrease in detection accuracy for the off-center frequencies that decreased in probability.
573
Turning next to Exp 7b,d,f (**Fig 6d-f**), what happens when initial experience with a unimodal
distribution shifts mid-study to equiprobable presentation? As now expected from prior results,
detection of the high-probability mode of a unimodal distribution is considerably more accurate
than detection of improbable frequencies (linear contrast, p = 5.198 x 10^{-30}, Cohen's d = 1.332,
Fig 7d-f). Detection of low-probability frequencies is impacted by proximity to the high-probability
center frequency (ANOVA, p = 0.010, η^2 = 0.016); accuracy is higher for frequencies nearest the
high-probability center frequency compared those at middle (p = 0.023, Cohen's d = 0.269) or far
frequencies (p = 0.023, Cohen's d = 0.274, both contrasts Holm-corrected). However, the
relatively preserved detection accuracy for tones near the high-probability frequency compared
to those is observed only in Exp 7b for the narrow range (near vs. middle, p = 0.014, Cohen's d =
0.512, near vs. far, p = 1.161 x 10^{-3}, Cohen's d = 0.697, all linear contrasts Bonferroni-corrected).
It is noteworthy that the tones sampling narrow distributions remain highly differentiable at ~8x
larger than typical just-noticeable frequency differences.
587

Figure 6. The detailed shape of statistically-driven gain is modulated by range, distribution, and
sampling density. See Fig 1c for histograms of distributional regularities. Marker size scales with tone
probability. In each panel, the darker color (dotted line) indicates behavior in the first half of the experiment;
the lighter color (solid line) indicates behavior in the second half, when distributional regularities shift. Each
panel plots mean detection accuracy as a function of acoustic frequency. Error bars indicate standard error
of the mean. The top row shows Exp 7a,c,e for which the equiprobable distribution preceded the unimodal
distribution. The bottom row shows Exp 7b,d,f for which a unimodal distribution preceded the switch to an
equiprobable distribution. Panels **(A)** and **(D)** plot the narrow distribution (5.5 semitone range), Panels **(B)**
and **(E)** plot the intermediate distribution (9.47 semitone range), and Panels **(C)** and **(F)** plot the wide
distribution (11.36 semitone range). In each panel, the insets show detection accuracy for the high-
probability tone (in the unimodal half of the experiment) and equiprobable low-probability tones near,
intermediate, and far from the high-probability 1000-Hz tone.

The effects on detection of proximity to the high-probability 1000 Hz mode are modulated by the
switch to an equiprobable distribution (ANOVA, $p = 3.279 \times 10^{-11}$, $\eta^2 = 0.023$). We observe a
continued, but smaller, detection advantage for the formerly-high-probability center frequency
compared to formerly improbable frequencies (linear contrast, $p = 1.194 \times 10^{-8}$, Cohen's d =
0.552). This change is driven by a *decrease* (difference of 7.1%, ANOVA, $p = 1.137 \times 10^{-12}$, $\eta^2 =$
0.085) in detection accuracy for the center frequency as it becomes 5 times less probable, as well
as a smaller (difference of ~2%, ANOVA, $p = 0.007$, $\eta^2 = 0.006$) *increase* in accuracy as off-center
frequencies become 3 times more probable; this is potentially compatible with a relative release
from suppression. This residual advantage does not vary significantly with distance from the
center frequency (ANOVA, $p = 0.213$, $\eta^2 = 0.002$) or interact with the range of frequencies
presented (ANOVA, $p = 0.202$, $\eta^2 = 0.004$). In sum, there is hysteresis from experience with the
unimodal distribution such that the formerly high-probability frequency remains better detected
than other frequencies.
614
Next, we ask if hysteresis is also observed in detection accuracy for 1000 Hz in a unimodal
distribution *after* prolonged initial exposure to an equiprobable distribution (second half of Exp
7a,c,e) compared to when the experiment begins with a unimodal distribution (first half of Exp
7b,d,f). We find that pre-exposure to 336 trials of the flat probability distribution diminishes
detection rates for the high-probability 1000 Hz tone in the subsequent unimodal distribution by
5.8% relative to when the identical unimodal distribution is encountered first (ANOVA, $p = 6.394$
$\times 10^{-4}$, $\eta^2 = 0.064$). The persistent damping effect of first encountering the equiprobable
distribution is not significantly affected by the range of frequencies encountered (ANOVA, $p =$
0.768, $\eta^2 = 0.003$).
624
Finally, we aggregate detection data for off-center frequencies across the unimodal conditions
from Exp 7a,c,e (when the unimodal distribution was preceded by equiprobable) and Exp 7b,d,f
(when it was first) to maximize the power to detect influences of frequency range and distance
from the higher-probability center frequency. Frequency range influences detection in unimodal
probability distributions (ANOVA, $p = 0.005$, $\eta^2 = 0.039$). Specifically, a wide frequency range
impairs overall off-center detection accuracy, compared to when the frequency range is narrow
(post hoc Holm-corrected for following three contrasts, $p = 0.006$, Cohen's d = 0.471). The middle
frequency range falls in-between and differs significantly from detection in wide ($p = 0.037$,
Cohen's d = 0.354) but not narrow ranges ($p = 0.429$, Cohen's d = 0.118). Moreover, the shape
of the drop-off in detection accuracy from the high-probability center frequency is significantly
graded only in the narrow frequency range, with a significant difference between the near and mid
frequency band conditions (linear contrast, $p = 0.015$, Cohen's d = 0.388), and a non-significant
decrease between the middle and far frequencies (linear contrast, $p = 0.316$, Cohen's d = 0.229).

638
To summarize Exp 7, we again observe that listeners' ability to detect a tone in noise is modulated
by dynamic changes in statistical distributions. Decreases in probability are met with diminished
detection and increases in probability improve detection. However, as we previously observed,
the degree of proximity to a more-probable center frequency in unimodal distributions partially
rescues detectability of low-probability frequencies. The impact of distributional learning on
detection reflects both the probability distribution and the range over which it is defined.
645
646   **Experience with a single-frequency point distribution results in suppressive reduction in**
647   **perceptual sensitivity to other frequencies**
648
The prior experiments leave open the possibility that perceptual interactions across adjacent trials
may account for the graded impact on detection, for example through spectrally contrastive
influences among tones with different frequencies (Holt, 2005). Exp 8 makes a critical test of
whether patterns of relative gain, characterized in the prior experiments, involves enhancement
of the high-probability frequency, suppression of low-probability frequencies, or a combination of
enhancement and suppression.
655
To do so, Exp 8 establishes a context in which participants detect *only* 1000 Hz tones in noise, or
an equiprobable distribution of 20 tones finely sampling frequency between 800-1200 Hz that
does not include 1000 Hz (**Fig 1c**). In Exp 8a, the first 320 trials involve 20 different equiprobable
(6.25%) tone frequencies (35-cent intervals from 800-1200 Hz, excluding 1000 Hz) and the
second 320 trials present exclusively 1000 Hz tones (100% probability). Exp 8b begins with 320
1000-Hz trials, then transitions to the 20-frequency equiprobable distribution (excluding 1000 Hz)
across 320 trials. Excluding 1000 Hz from the stimulus set provides a control for possible
perceptual interactions across adjacent trials that may have an influence and establishes a
baseline against which to evaluate evidence of enhancement and suppression.
665

666
667
Figure 7. Experience with a single-frequency point distribution results in suppressive reduction in
perceptual sensitivity to other frequencies. Exp 8 makes a critical test of whether the gain characterized
in the prior experiments involves enhancement of the high-probability frequency, suppression of low-
probability frequencies, or a combination of enhancement and suppression. The histograms to the left show
distributional regularities for Exp 8a and Exp 8b. Marker size scales with tone probability. Mean detection
accuracy is shown as a function of acoustic frequency, with standard error of the mean indicated by error
bars. In Exp 8a (dark blue, dashed line), detection trials included 20 equiprobable tones (800-1200 Hz,
excluding 1000 Hz) in the first half of the study. In the second half, tones were exclusively 1000 Hz. In Exp
8b (light blue, solid line) the first half of the study involved only 1000 Hz whereas the second half shifted to
20 equiprobable frequencies (800-1200 Hz, excluding 1000 Hz). The inset shows detection in the context
of equiprobable distributions for each experiment, as a function of distance from 1000 Hz. Note that
detection is somewhat 'rescued' around 1000 Hz and that detection of frequencies distant from 1000 Hz is
suppressed in Exp 8b relative to Exp 8a.

We first ask whether the consistent experience with 1000 Hz in the first half of Exp 8b yields
accumulating detection accuracy improvements (**Fig 7b**). It does not: accuracy in the first quarter
of trials (first half of the first half) is 78% (aligned with expectations from listener-specific
thresholding) then decreases slightly to plateau at 75% for the remaining trials in the first half of
the study (ANOVA, p = 0.015, η^2 = 0.068). Similarly, neither Exp 1a (ANOVA, p = 0.210, η^2 =
0.050) or Exp 8a (ANOVA, p = 0.451, η^2 = 0.050) exhibit improved detection across a block of
trials with only 1000 Hz tones. There is a similar initial detection decrement of ~5% across the
first quarter of the 20-equiprobable-frequency trials of Exp 8a with no further change (ANOVA, p
= 9.669 x 10^{-6}, η^2 = 0.049). This same pattern emerges in the initial equiprobable blocks of Exp
7a,c,e (ANOVA, p = 1.375 x 10^{-5}, η^2 = 0.058). Detection accuracy for equiprobable distributions
that are experienced in the first half of a study does not differ over experiments (ANOVA, Exp 5a,
7a,c,e, and 8a; p = 0.387, η^2 = 0.024).

Turning next to the nature of the gain, we first examine whether initial experience with the 20-tone
equiprobable distribution in Exp 8a (which does not include 1000 Hz) impacts subsequent
detection in the 1000-Hz-only block (**Fig 7a**). It does not: detection of 1000 Hz in the second half
of Exp 8a did not differ from either Exp 1a (Tukey-corrected post hoc contrast, p = 0.315, Cohen's
d = 0.333) or the first half of Exp 8b (Tukey-corrected p = 0.837, Cohen's d = 0.104), each of
which involved blocks of trials with only 1000 Hz at the beginning of the study.

In contrast, massed exposure to 1000 Hz in the first half of Exp 8b drives a dramatic, long-lasting,
and frequency-specific detection decrement for the subsequently encountered 20 equiprobable
frequencies, as compared to detection across equiprobable frequencies in Exp 8a (ANOVA,
interaction of Distance-from-1000-Hz x Exp, p = 2.618 x 10^{-4}, η^2 = 0.007). Specifically, as shown
in **Fig 7b**, detection of frequencies at far (2 to 3.9 semitones) and intermediate (1 to 2 semitones)
distances from 1000 Hz were detected much less accurately after massed experienced with 1000
Hz (Exp 8b, linear contrasts; far: p = 2.427 x 10^{-3}, Cohen's d = 0.330; intermediate: p = 9.784 x
10^{-4}, Cohen's d = 0.346), compared to equiprobable presentation at the beginning of the study
(Exp 8a). This suppressive effect was rescued by proximity to the now-absent 1000 Hz in the
second half of Exp 8b, with frequencies within about a semitone from 1000 Hz eliciting detection
accuracies roughly on par with those from Exp 8a (p = 0.332, Cohen's d = 0.311). Thus, a half-
hour of 1000-Hz exposure induces a lasting attentional filter that impacts the ability to detect
frequencies varying from 800-1200 Hz, even though 1000 Hz was never again encountered.

One might expect that any initial learning across the 1000-Hz-only distribution would be
overwhelmed by the mid-study shift to the high-uncertainty 20-frequency equiprobable
distribution. However, we see the opposite: across the second half of Exp 8b, there is no
significant change in overall detection accuracy (ANOVA p = 0.165, η^2 = 0.008), nor any change
across time in relative accuracy of detection across frequencies (ANOVA, p = 0.568, η^2 = 0.006).
The large advantage for detection of frequencies near 1000 Hz compared to intermediate and far
frequencies persists to the final 80 trials of Exp 8b (linear contrast, p = 0.002, Cohen's d = 0.257).
This effect is further evidenced by comparing the second half of Exp 8b with the first half of Exp
8a. Here, there is strong suppression of frequencies at far and intermediate distances from 1000
Hz in Exp 8b compared to detection of the same frequencies in the equiprobable half of Exp 8a.
As for the within-experiment comparison, this difference is observed through the entirety of the
second half of the study, again extending even to the last quarter of trials (linear contrast, p =
0.008, Cohen's d = 0.440). The absence of 1000 Hz over this period rules out the possibility that
trial-wise perceptual interactions or the experience of a relative probability difference for a

particular frequency were strong contributors to the hysteresis observed in Exp 5 and Exp 7. See
Fig S2.

Discussion

Is perception guided toward what we expect, or by what surprises us? Here, across 29
experiments, we examine two perceptual tasks for which distributional regularities accumulate
over a task-irrelevant dimension without instruction, directed attention, or feedback. We find that
distributional learning drives dynamic shifts in perception across tasks. It affects sound detection:
a faint tone of a particular frequency is better detected in noise if it occurs frequently than if it
occurs rarely. Distributional learning also affects duration decisions, which are faster for tones
that possess a frequency that has been more common in recent experience. Across tasks, our
results converge to indicate that effects of expectation are largely driven by suppression of less-
probable, unexpected stimuli that are, to a lesser degree, supported by modest enhancement of
highly probable stimuli.

Our examination of expectation built across distributions (rather than dichotomous probabilities)
affords a wider vantage point for understanding how perceptual gain is modulated by expectation.
Our results reveal an influence on perception that is graded as a function of the distribution mode,
the range of the distribution, and the position of a stimulus within the distribution. The detailed
shape of the distribution is important, as well, as shown by the bimodal profile of tone detection
evoked by a bimodal frequency distribution. Strikingly, equally probable rare events are perceived
differently as a function of their perceptual distance from the distribution mode(s). Decades ago,
Greenberg and Larkin (1968) examined tone detection in a similar paradigm (albeit with overt
instructions about tone probability instead of distributional learning) and interpreted the graded
gain to be indicative of a frequency-selective attentional filter situated at the high-probability mode
with increasingly suppressive sidebands with greater distance from the mode. Indeed, in the time
since there has been sustained interest (e.g., Summerfield & Egner, 2009; Zivony & Eimer, 2024)
in isolating the influence of *expectation* - operationalized by manipulating the probability of stimuli
– from *attention* – defined according to the utility or relevance of these stimuli to a task
(Summerfield & de Lange, 2014; Kok et al., 2012). Under these definitions, the present tasks are
attention-neutral and involve manipulations of *expectation* only. Yet, our results suggest that
expectation built across distributional learning establishes a selection filter that impacts how (and
whether) subsequent stimuli are perceived. Whether this is described as a dimension-selective
attentional filter (as proposed by Greenberg & Larkin, 1968) or more neutrally as an experience-
driven predictive filter, the present results are distinct from manipulations of task utility or
relevance that have been previously attributed to attention interacting with expectation (Zivony &
Eimer, 2024; Rungratsameetaweemana & Serences, 2019).

In the time domain, the influence of distributional learning on perception is persistent: effects of a
unimodal distribution provoke lasting influence with a continued advantage for tones that were
previously probable and a lasting disadvantage for the tones that were previously improbable,
even after exposure to a uniform distribution. Even so, there remains sensitivity to volatile
distribution changes with both detection and perceptual decisions dynamically adjusting when
dichotomous probabilities flip. Future work will be needed to resolve the interpretive tension
between the rapid adjustment we observe across changing dichotomous probabilities in Exp 3
and Exp 4 versus the lingering influence of bimodal (Exp 5,6,7) and point (Exp 8) distributions.
Candidate contributors include the magnitude of differences in stimulus probabilities,
dichotomous versus more fully sampled distributions, lower information conveyance by uniform

distributions, and the relative volatility experienced across a listening session. The present
paradigms provide a basis for further discovery, with implications for 'stubborn predictions'
examined in other literatures (Yon et al., 2023).

The impact of these distributional regularities on perception is evident for both detection and
perceptual decisions, emphasizing the breadth of influence of distributional learning across
distinct task demands (Fritsche, Mostert, & de Lange, 2017). Even so, detection provides a unique
window through which to observe effects of distributional learning and resulting expectations, as
it has a natural baseline set by individuals' thresholds. The detection results make it especially
clear that the net impact of distributional learning is to prioritize the high-probability distribution
mode - not by enhancing detectability of the expected stimulus, but instead by suppressing
detectability of rare, unexpected stimuli. We observe this repeatedly across experiments. Despite
considerable headroom for detection accuracy to improve in the context of a threshold set at
~79% accuracy we do not observe substantial enhancement of detection of the high-probability
tone. Indeed, in the original Greenberg and Larkin (1968) study, exposure to tens of thousands
of trials of a high-probability frequency did not enhance detection above the initially established
perceptual threshold. This lack of enhancement at the mode is somewhat surprising given the
literature on perceptual learning (Amitay, Zhang, Jones, & Moore, 2014; Watanabe & Sasaki,
2015; Wright, Wilson, & Sabin, 2010), where intensive practice with attentionally-demanding
perceptual paradigms can drive improved detection. But, in contrast to most perceptual learning
approaches, the influences we observe accrue across a task-irrelevant perceptual dimension,
without directed attention, reward, or feedback.

It would seem inefficient for a system to track distributional regularities irrelevant to the task at
hand. However, 'optimal' selectivity to a task-relevant dimension may not always be adaptive for
perception: in natural environments with shifting demands, it may be effective to 'keep an ear out'
by tracking evolving regularities with potential utility for future behavior (Heilbron & Chait, 2017).
Moreover, the reduction in sensitivity to subsequently encountered frequencies that we observe
following massed exposure to a single frequency would seem to be a maladaptive loss of
perceptual sensitivity. Instead, it may reflect gain mechanisms that suppress sensitivity to
region(s) along a perceptual dimension that are less likely to be encountered. In the sense that
one cannot be surprised by something if one is not sure it has occurred (Press et al., 2020), the
suppressive effects we observe for low-probability stimuli distant from a distribution mode are
substantial enough that these stimuli would seem to be less likely to contribute to subsequent
distributional learning. Distributional effects on perception thus may have the potential to
snowball, exaggerating regularities relative to the true distribution of events.

Our results emphasize that layered histories of experience with distributional regularities impact
behavior. For example, unimodal distributions have lingering effects even after a switch to
equiprobable stimulus presentation. At a longer timescale, we observe a consistent frequency-
duration bias in our perceptual decision experiments. The effect is persistent across decision
experiments (even when only two frequencies were present) and appears to be associated with
the ordinal position of frequencies in the distribution range rather than absolute frequency.
Although acoustic frequency and duration would seem to be good candidates for orthogonal
acoustic input dimensions – and indeed, older studies had suggested this (Allan & Kristofferson,
1974; Woods, Sorkin, & Boggs, 1979) – the ubiquity of interactions between acoustic dimensions
is seen clearly in auditory category learning studies in which rotating the sampling of acoustic
category exemplars in an ostensibly orthogonal acoustic space produces radically different

learning outcomes due to prior expectations about the relationship between the dimensions
(Roark & Holt, 2022; Bröker et al., 2024; Bröker, Plaut & Holt, 2024).

Life-long exposure to the distributional statistics of natural sound environments may drive at least
some of the ubiquitous bias to perceive relatively lower frequencies as longer, and relatively
higher frequencies as shorter (Fiser, Berkes, Orbán, & Lengyel, 2010; Berkes, Orbán, Lengyel, &
Fiser, 2011). Pinning down the etiology of this endogenous bias will be challenging, as multiple
environmental and acoustic factors may contribute. From different decay characteristics for struck
strings on the piano (undamped bass notes decay much more slowly than treble notes; Fletcher,
Blackham & Stratton, 1962) to the longer reverberance for lower versus higher frequencies
(Backus, 1977) there are complex, and likely consistent, regularities across acoustic frequency
and duration that individuals may learn about over a lifetime of listening. Just as important, there
may be sensory contributions such that the longer duration of cochlear filters at lower frequencies
may facilitate identification of lower-frequency tones as long duration, consistent with the bias we
observe across experiments (Patterson, 1987). For example, an auditory filter centered at 800 Hz
is expected to have a longer duration than a filter centered at 1200 Hz. Thus, longer-lasting
peripheral excitation at lower frequencies may facilitate identification of lower-frequency tones as
long duration.
837
Sensory factors may play a role in the subtle influence across frequencies observed for tone
detection, as well. As noted above, the cochlea's limited ability to resolve stimuli creates critical
bands within which it is difficult to distinguish frequencies (Fletcher, 1940). As described above,
the 'rescue' of detection accuracy for low probability tone frequencies that lie closer to the
distribution mode may partly reflect limitations of frequency resolution within a critical band.
Similarly, changes in the width of the critical band across our 800-1200 Hz frequency range may
account for the subtle, consistent tendency for detection to be less accurate at higher frequencies
(Fig 2, 5, 6). Yet, beyond sensory influences, our manipulation of distributions (instead of binary
probabilities) helps to reveal that other factors are at play. For example, both the distribution range
and the position of the high probability tone with the distribution range influence detection.
848
The present results are potentially informed by rich literatures studying neural response across
stimuli that vary in probability. Repeated exposure to a stimulus changes neural firing patterns in
visual (Schoups, Vogels, Qian, & Orban, 2001) and auditory (Khouri & Nelken, 2015) cortex. Two
neural phenomena - the mismatch negativity (MMN, Naatanen et al., 1978), and stimulus specific
adaptation (SSA, Ulanovsky et al., 2004) – are extensively studied in the auditory domain using
an 'oddball' paradigm in which common and rare stimuli are intermixed in a sequence. This
probability manipulation reveals exaggerated neural response to low-probability sounds, seeming
to run counter to the principally suppressive behavioral effects we observe for low-probability
tones. However, these neural phenomena can be observed in active freely moving states
(Polterovich et al., 2018) as well as under anesthesia (Yaron et al., 2012) and in disordered
consciousness (Bekinschtein et al., 2009); we do not yet have a strong understanding of how they
relate to auditory behavior. There is much more to understand in relating exaggerated neural
response to low-probability sounds with slower decisions and less accurate detection.
Nonetheless, these neurobiological literatures support the possibility that effects of distributional
learning can emerge robustly in primary auditory cortex, and possibly even auditory midbrain
(Malmierca et al., 2009; Duque et al., 2012; Cruces-Solís et al., 2018).
865
Schröger and Wolf (1998), who pioneered the duration-decision task we use here, argued from
human electroencephalography results that, at least for perceptual decisions, effects may arise

from a memory-based mechanism that detects deviance from expectations and orients attention
to the rare stimulus frequency. This leaves fewer resources for perceptual decision making,
slowing response times. However, in a case of convergent experimental design, Mondor and
Bregman (1994) used a very similar duration-decision paradigm to argue that the reaction time
advantage for probable or cued frequencies instead arises from allocation of selective attention
to the probable (not the improbable) frequency. This echoes the interpretational challenges of the
larger literature on expectation and attention effects (see Introduction) and is mirrored, as well, in
literatures attempting to resolve patterns of behavioral repetition priming and neural repetition
suppression (McMahon & Olson, 2007; Feuerriegel, Vogels, & Kovács, 2021).

Organisms as diverse as humans and honeybees are exquisitely sensitive to patterns that unfold
across sensory input. We find that distributional learning affects fundamental aspects of
perception: the very ability to detect whether a stimulus is present and to make a simple judgment
about it. Listeners rapidly and implicitly apprehend distributional regularities of how often stimuli
occur, even when the regularities emerge across sensory dimensions irrelevant to the task at
hand. This statistical learning across input distributions arises rapidly even in the context of
statistically dynamic contexts and has a substantial and lasting influence on perception driven by
collaborative influences of sensory processing, distributional learning, and selective attention in
sculpting a gain function involving modest enhancement and robust suppression.

Materials and Methods

Experiment materials, code, and analyses can be found at https://osf.io/xdgnw/.

Participants. Participants (ages 18-35 yrs) were recruited online and compensated via Prolific.co
(Damer & Bradley, 2014). All self-reported normal hearing. **Table S1** provides experiment-wise
demographic details. Based on power analyses of pilot data collected using the same tasks, we
targeted recruitment of 30 participants/experiment.

Stimuli. Sinewave tones and white noise were generated in the lossless FLAC format using the
Sound eXchange sound processing software (SoX, http://sox.sourceforge.net/) at 44.1kHz and
16-bit precision.

Procedure. All experiments were conducted online following best-practices described by Zhao
et al. (2022) using PsychoPy (2022.1.2, pavlovia.org) for tone-in-noise detection experiments and
Gorilla (Anwyl-Irvine et al., 2020) for duration-decision experiments. Online participants used the
Chrome browser on their own laptop or desktop computer (no smartphones or tablets) with a brief
listening test assuring headphone compliance (Milne et al., 2020). **Fig 1** illustrates the trial
structure for each task. Table S2 provides experiment-level details.

Tone-in-Noise Detection Task. Continuous white noise commenced +40 dB relative to the level
just detectable over participants' own computer and headphones, as determined by a brief
system-calibration procedure (Zhao et al., 2022). Adaptive thresholding commenced with the
onset of a 300-sec white noise (200-ms cosine amplitude onset/offset ramps) that looped
continuously through the end of the study. Adaptive thresholding entailed detecting a 250-ms (10-
ms cosine onset/offset ramps), 1000-Hz sinewave tone (1080-Hz in Exp 1f) in a three-interval
forced choice task (**Fig 1a**). The first 6 trials served as practice, with feedback and -13.75 dB
SNR. Thereafter, there was no feedback across three 40-trial adaptive thresholding runs. Each
run began at -13.75 dB SNR with tone intensity decreasing 1.5 dB after each correct detection

until the SNR reached -19.75 dB, or until an incorrect response. Subsequently, tone intensity
decreased -.75 dB after three correct responses and increased +.75 dB after each incorrect
response. Threshold tone-in-noise detection was computed as the mean of the mode tone
intensity across the three runs (Zhao et al. 2022) which estimates threshold at 79.4% correct
detection (Levitt, 1971).

Adaptive thresholding established a by-participant threshold tone intensity for the tone-in-noise
experiment. The first experiment block was practice, with -13.75 dB SNR, feedback, and tone
frequencies that matched the initial experiment distributional regularity (**Fig 1a**). After practice,
tone intensity was set to -.75 dB relative to the threshold estimate for the remainder of the
experiment. Participants reported which of two intervals contained the tone (**Fig 1a**). Participants
were not informed about the task-irrelevant distributional regularities across acoustic frequency
(**Fig 1c**). The entire protocol took about 30 minutes, except in experiments with double the trials
(see **Table S2**). We report mean detection accuracy.

Duration-Decision Task. Each trial involved a single sinewave tone presented in quiet at a
comfortable level. Tones were 50 or 90 ms, with equal probability and random presentation.
Participants reported whether the tone was "long" or "short" with a key press and were not
instructed about the task-irrelevant distributional regularities across acoustic frequency (**Fig 1b**).
Each experiment began with a practice block involving feedback and a distributional regularity
that mirrored the main experiment. There was no feedback for the remainder of the experiment.
Table S2 provides experiment-wise details. The entire protocol took about 30 minutes, except in
experiments with double the trials. Analyses focused on decision response time, measured from
tone offset to response. Trials for which response time was shorter than 300 ms or longer than
1500 ms (non-inclusive) were excluded from analyses (see **Table S1** for percent of trials
excluded).

Approach to Analysis. Data were preprocessed using JMP Pro 17.0.0, and statistical analyses
were conducted in JASP (JASP team, Amsterdam, Netherlands, 10/19/22, version 0.16.4). We
report Greenhouse-Geisser corrected degrees of freedom and *p* values for ANOVAs for which
the assumption of sphericity was violated, as determined by a Mauchly test. Multiple comparison
correction for linear contrasts was carried out using Bonferroni correction, and for post-hoc tests
using Holm correction. Study-wise analysis details are provided in **Table S3**.

Acknowledgements and Funding Sources

The work was supported by grants from National Institutes of Health (R01DC004674) and the
National Science Foundation (SBE/BCS 2414066, 2420979) to LLH and FD, an institutional
training grant from the National Institute of General Medical Sciences to LLH that supported AL
(T32GM081760), and an individual NRSA from the National Institute on Deafness and
Communication Disorders to SL (F32DC020625) under the mentorship of LLH and FD. The work
reported here comprised a chapter of AL's 2024 PhD dissertation (Carnegie Mellon University).
Christi Gomez, Erin Smith, and Gracee Swatsworth contributed to the study. Drs. Clare Press,
Daniel Yon, and Alain de Chevigné provided very helpful comments on a manuscript draft.

Competing Interests

The authors declare that they have no competing interests.

References

Allan, L. G., & Kristofferson, A. B. (1974). Psychophysical theories of duration discrimination. *Perception & Psychophysics*, *16*(1), 26-34.

Alink, A., & Blank, H. (2021). Can expectation suppression be explained by reduced attention to predictable stimuli?. *NeuroImage*, *231*, 117824. https://doi.org/10.1016/j.neuroimage.2021.117824

Amitay, S., Zhang, Y. X., Jones, P. R., & Moore, D. R. (2014). Perceptual learning: top to bottom. *Vision research*, *99*, 69–77. https://doi.org/10.1016/j.visres.2013.11.006

Backus, John, The Acoustical Foundations of Music, 2nd Ed, W W Norton, New York, 1977

Bekinschtein, T. A., Dehaene, S., Rohaut, B., Tadel, F., Cohen, L., & Naccache, L. (2009). Neural signature of the conscious processing of auditory regularities. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(5), 1672–1677. https://doi.org/10.1073/pnas.0809667106

Berkes, P., Orbán, G., Lengyel, M., & Fiser, J. (2011). Spontaneous Cortical Activity Reveals Hallmarks of an Optimal Internal Model of the Environment. Science, 331(6013), 83–87. https://doi.org/10.1126/science.1195870

Blakemore, S. J., Wolpert, D. M., & Frith, C. D. (1998). Central cancellation of self-produced tickle sensation. *Nature neuroscience*, *1*(7), 635–640. https://doi.org/10.1038/2870

Bröker, F., Holt, L. L., Roads, B. D., Dayan, P., & Love, B. C. (2024). Demystifying unsupervised learning: how it helps and hurts. *Trends in cognitive sciences*, S1364-6613(24)00227-4. Advance online publication. https://doi.org/10.1016/j.tics.2024.09.005

Bröker, F., Plaut, D., & Holt, L.L. (2024). Supervised learning outperforms unsupervised learning only for categories of intermediate difficulty. https://osf.io/preprints/psyarxiv/3rfjk

Cristià A. (2011). Fine-grained variation in caregivers' /s/ predicts their infants' /s/ category. *The Journal of the Acoustical Society of America*, *129*(5), 3271–3280. https://doi.org/10.1121/1.3562562

Cruces-Solís H, Jing Z, Babaev O, Rubin J, Gür B, Krueger-Burg D, et al. (2018) Auditory midbrain coding of statistical learning that results from discontinuous sensory stimulation. PLoS Biol 16(7): e2005114. https://doi.org/10.1371/journal.pbio.2005114

de Lange, F. P., Heilbron, M., & Kok, P. (2018). How Do Expectations Shape Perception?. Trends in cognitive sciences, 22(9), 764–779. https://doi.org/10.1016/j.tics.2018.06.002

Duque, D., Pérez-González, D., Ayala, Y. A., Palmer, A. R., & Malmierca, M. S. (2012). Topographic distribution, frequency, and intensity dependence of stimulus-specific adaptation in the inferior colliculus of the rat. Journal of Neuroscience, 32(49), 17762-17774.

Fiser, J., Berkes, P., Orbán, G., & Lengyel, M. (2010). Statistically optimal perception and learning: from behavior to neural representations. Trends in Cognitive Sciences, 14(3), 119–130. https://doi.org/10.1016/j.tics.2010.01.003

Fletcher, H. (1940). Auditory Patterns. Reviews of Modern Physics, 12, 47–65. doi.org/10.1103/RevModPhys.12.47.

Fletcher, H., Blackham, E. D., & Stratton, R. (1962). Quality of piano tones. The Journal of the Acoustical Society of America, 34(6), 749-761.

Feuerriegel, D., Vogels, R. & Kovács, G. (2021). Evaluating the evidence for expectation suppression in the visual system. Neurosci. Biobehav. Rev. 126, 368–381.

Friston, K. (2005). A theory of cortical responses. Philosophical Transactions of the Royal Society, B: Biological Sciences, 360(1456), 815–836.

Fritsche, M., Mostert, P., & Lange, F. P. de. (2017). Opposite Effects of Recent History on Perception and Decision. *Current Biology*, 27(4), 590–595. doi.org/10.1016/j.cub.2017.01.006

Garrido, M. I., Dolan, R. J., & Sahani, M. (2011). Surprise leads to noisier perceptual decisions. i-Perception, 2, 112-120. dx.doi.org/10.1068/i0411

Greenberg, G. Z., & Larkin, W. D. (1968). Frequency-response characteristic of auditory observers detecting signals of a single frequency in noise: the probe-signal method. *The Journal of the Acoustical Society of America*, 44(6), 1513–1523. https://doi.org/10.1121/1.1911290

Heilbron, M., & Chait, M. (2017). Great expectations: Is there evidence for predictive coding in auditory cortex? *Neuroscience*, 1–20. Retrieved from https://doi.org/10.1016/j.neuroscience.2017.07.061

Holt L. L. (2005). Temporally nonadjacent nonlinguistic sounds affect speech categorization. *Psychological science*, *16*(4), 305–312. https://doi.org/10.1111/j.0956-7976.2005.01532.x

Khouri, L., & Nelken, I. (2015). Detecting the unexpected. *Current opinion in neurobiology*, *35*, 142–147. https://doi.org/10.1016/j.conb.2015.08.003

Kilteni, K., & Ehrsson, H. H. (2017). Body ownership determines the attenuation of self-generated tactile sensations. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(31), 8426–8431. https://doi.org/10.1073/pnas.1703347114

Kok, P., Jehee, J. F., & de Lange, F. P. (2012). Less is more: expectation sharpens representations in the primary visual cortex. *Neuron*, *75*(2), 265–270. https://doi.org/10.1016/j.neuron.2012.04.034

Kumar, S., Kaposvari, P., & Vogels, R. (2017). Encoding of Predictable and Unpredictable Stimuli by Inferior Temporal Cortical Neurons. *Journal of cognitive neuroscience*, *29*(8), 1445–1454. https://doi.org/10.1162/jocn_a_01135

1048 Logothetis, N. What we can do and what we cannot do with fMRI. *Nature* **453**, 869–878 (2008).
1049 https://doi.org/10.1038/nature06976
1050
1051 Love B. C. (2003). The multifaceted nature of unsupervised category learning. *Psychonomic*
1052 *bulletin & review*, *10*(1), 190–197. https://doi.org/10.3758/bf03196484
1053
1054 Malmierca, M. S., Cristaudo, S., Pérez-González, D., & Covey, E. (2009). Stimulus-specific
1055 adaptation in the inferior colliculus of the anesthetized rat. Journal of Neuroscience, 29(17), 5483-
1056 5493.
1057
1058 McMahon, D. B. T., & Olson, C. R. (2007). Repetition Suppression in Monkey Inferotemporal
1059 Cortex: Relation to Behavioral Priming. Journal of Neurophysiology, 97(5), 3532–3543. Retrieved
1060 from https://doi.org/10.1152/jn.01042.2006
1061
1062 McMurray, B., Aslin, R. N., & Toscano, J. C. (2009). Statistical learning of phonetic categories:
1063 insights from a computational approach. *Developmental science*, *12*(3), 369–378.
1064 https://doi.org/10.1111/j.1467-7687.2009.00822.x
1065
1066 Meyer, T., & Olson, C. R. (2011). Statistical learning of visual transitions in monkey inferotemporal
1067 cortex. *Proceedings of the National Academy of Sciences of the United States of America*,
1068 *108*(48), 19401–19406. https://doi.org/10.1073/pnas.1112895108
1069
1070 Milne, A. E., Chait, M., & Conway, C. M. (2024). Probing sensitivity to statistical structure in rapid
1071 sound sequences using deviant detection tasks. bioRxiv 2024.04.19.590221; doi:
1072 https://doi.org/10.1101/2024.04.19.590221
1073
1074 Mondor, T. A., & Bregman, A. S. (1994). Allocating attention to frequency regions. *Perception &*
1075 *psychophysics*, *56*(3), 268–276. https://doi.org/10.3758/bf03209761
1076
1077 Näätänen, R., Gaillard, A. W., & Mäntysalo, S. (1978). Early selective-attention effect on evoked
1078 potential reinterpreted. *Acta psychologica*, *42*(4), 313–329. https://doi.org/10.1016/0001-
1079 6918(78)90006-9
1080
1081 Pinto, Y., van Gaal, S., de Lange, F. P., Lamme, V. A., & Seth, A. K. (2015). Expectations
1082 accelerate entry of visual stimuli into awareness. *Journal of vision*, *15*(8), 13.
1083 https://doi.org/10.1167/15.8.13
1084
1085 Polterovich A, Jankowski MM, Nelken I (2018) Deviance sensitivity in the auditory cortex of freely
1086 moving rats. PLoS ONE 13(6): e0197678. https://doi.org/10.1371/journal.pone.0197678
1087
1088 Press, C., Kok, P., & Yon, D. (2020). The Perceptual Prediction Paradox. *Trends in cognitive*
1089 *sciences*, *24*(1), 13–24. https://doi.org/10.1016/j.tics.2019.11.003
1090
1091 Richter, D., Ekman, M., & de Lange, F. P. (2018). Suppressed Sensory Response to Predictable
1092 Object Stimuli throughout the Ventral Visual Stream. *The Journal of Neuroscience*, *38*(34), 7452–
1093 7461. https://doi.org/10.1523/JNEUROSCI.3421-17.2018
1094

Roark, C. L., & Holt, L. L. (2022). Long-term priors constrain category learning in the context of
short-term statistical regularities. *Psychonomic bulletin & review*, *29*(5), 1925–1937.
https://doi.org/10.3758/s13423-022-02114-z

Rosenthal, O., Fusi, S., & Hochstein, S. (2001). Forming classes by stimulus frequency: behavior
and theory. *Proceedings of the National Academy of Sciences of the United States of America*,
98(7), 4265–4270. https://doi.org/10.1073/pnas.071525998

Rungratsameetaweemana, N., & Serences, J. T. (2019). Dissociating the impact of attention and
expectation on early sensory processing. *Current opinion in psychology*, *29*, 181–186.
https://doi.org/10.1016/j.copsyc.2019.03.014

Scharf, B., Quigley, S., Aoki, C., Peachey, N., & Reeves, A. (1987). Focused auditory attention
and frequency selectivity. *Perception & psychophysics*, *42*(3), 215–223.
https://doi.org/10.3758/bf03203073

Schatz, T., Feldman, N. H., Goldwater, S., Cao, X. N., & Dupoux, E. (2021). Early phonetic
learning without phonetic categories: Insights from large-scale simulations on realistic input.
Proceedings of the National Academy of Sciences of the United States of America, *118*(7),
e2001844118. https://doi.org/10.1073/pnas.2001844118

Schoups, A., Vogels, R., Qian, N., & Orban, G. (2001). Practising orientation identification
improves orientation coding in V1 neurons. *Nature*, *412*(6846), 549–553.
https://doi.org/10.1038/35087601

Schröger, E., & Wolff, C. (1998). Behavioral and electrophysiological effects of task-irrelevant
sound change: A new distraction paradigm. Cognitive Brain Research, 7, 71–87.
https://doi:10.1016/S0926-6410(98)00013-5

Sek, A., & Moore, B. C. (1995). Frequency discrimination as a function of frequency, measured
in several ways. *The Journal of the Acoustical Society of America*, *97*(4), 2479–2486.
https://doi.org/10.1121/1.411968

Stein, T., & Peelen, M. V. (2015). Content-specific expectations enhance stimulus detectability by
increasing perceptual sensitivity. *Journal of experimental psychology. General*, *144*(6), 1089–
1104. https://doi.org/10.1037/xge0000109

Summerfield, C., & Egner, T. (2009). Expectation (and attention) in visual cognition. *Trends in*
cognitive sciences, *13*(9), 403–409. https://doi.org/10.1016/j.tics.2009.06.003

Summerfield, C., & de Lange, F. P. (2014). Expectation in perceptual decision making: neural and
computational mechanisms. *Nature reviews. Neuroscience*, *15*(11), 745–756.
https://doi.org/10.1038/nrn3838

Ulanovsky, N., Las, L., Farkas, D., & Nelken, I. (2004). Multiple time scales of adaptation in
auditory cortex neurons. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience, *24*(46), 10440–10453. https://doi.org/10.1523/JNEUROSCI.1905-04.2004
1142

Watanabe, T., & Sasaki, Y. (2015). Perceptual learning: toward a comprehensive theory. *Annual review of psychology*, *66*, 197–221. https://doi.org/10.1146/annurev-psych-010814-015214

Werker, J. F., Yeung, H. H., & Yoshida, K. A. (2012). How do infants become experts at native-speech perception? *Current Directions in Psychological Science, 21*(4), 221–226. https://doi.org/10.1177/0963721412449459

Woods, D. D., Sorkin, R. D., & Boggs, G. J. (1979). Stimulus context and duration discrimination. *Perception & Psychophysics, 26(2)*, 127-132.

Wright, B. A., Wilson, R. M., & Sabin, A. T. (2010). Generalization Lags behind Learning on an Auditory Perceptual Task. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 30(35), 11635–11639. Retrieved from https://doi.org/10.1523/jneurosci.1441-10.2010

Yaron, A., Hershenhoren, I., & Nelken, I. (2012). Sensitivity to complex statistical regularities in rat auditory cortex. *Neuron*, *76*(3), 603–615. https://doi.org/10.1016/j.neuron.2012.08.025

Yon, D., Gilbert, S.J., de Lange, F.P., & Press, C. (2018). Action sharpens sensory representations of expected outcomes. *Nature Communications, 9*, 4288. https://doi.org/10.1038/s41467-018-06752-7

Zhao, S., Brown, C. A., Holt, L. L., & Dick, F. (2022). Robust and Efficient Online Auditory Psychophysics. *Trends in Hearing*, *26*, 23312165221118792. https://doi.org/10.1177/23312165221118792

Zivony, A., & Eimer, M. (2024). A dissociation between the effects of expectations and attention in selective visual processing. *Cognition*, *250*, 105864. Advance online publication. https://doi.org/10.1016/j.cognition.2024.105864