1

The holobiont's extracellular microbiome

modulates the adaptation of an intracellular endosymbiont

Madangchanok Imchen[1], Rupinder Kaur[1], Angelina McGarry[1], and Seth R. Bordenstein[1]*

[1]Departments of Biology and Entomology, One Heath Microbiome Center, Huck Institutes of the

Life Sciences, The Pennsylvania State University, USA

* Corresponding author

E-mail: srb6251@psu.edu

Keywords: cytoplasmic incompatibility, microbiome, intracellular bacteria, symbiosis,

Drosophila melanogaster.

Abstract

Animal holobionts contain two broad classes of microorganisms, intra- and extracellular microbes, that are largely studied in symbiotic siloes. Here we demonstrate the impact of the extracellular gut microbiome on an intracellular reproductive endosymbiont using the *Drosophila melanogaster* holobiont and its endosymbiont *Wolbachia* that adaptively cause a paternal-effect lethality of embryos termed cytoplasmic incompatibility (CI). We report the endosymbiont's adaptation is not only self-sufficient but stronger in a host without an extracellular microbiome. Indeed, host transcripts associated with the CI adaptation are enhanced, while cell biological hallmarks exhibit complex quantitative and qualitative differences in microbiome-free hosts. Decreases in the gut bacteria *Lactobacillus* specifically link with high penetrance of the CI adaptation. Notably, both the extracellular microbiome and endosymbiont harbor markedly higher cellular densities in a complete holobiont setting than when either constituent is reared alone. Altogether, this study uncovers a novel interaction whereby the extracellular microbiome modulates the phenotype of an intracellular endosymbiont, revealing intricate symbiotic networks within the holobiont.

Introduction

Animal and plant holobionts contain a consortium of microorganisms spanning intracellular endosymbionts and extracellular microbes with varying degrees of interdependencies [1,2]. The co-existence of these endosymbionts and extracellular microbes with different lifestyles and functions can impact each other in competitive or collaborative ways [3], potentially generating trade-offs and synergies across symbiotic types and host anatomical sites throughout a host's lifespan. The net outcome of such multipartite interactions may leave symbiotic ripple effects in a holobiont.

Challenges to assessing the specific impacts of the extracellular microbiome on endosymbioses are two-fold. First, microbiome elimination methods are often indiscriminate to the extracellular and intracellular microbiome, and second, most endosymbionts cannot be cultured outside of the host cell [4]. Here, we overcome these factors using *Drosophila melanogaster* that can be reared without its simple extracellular gut microbiome [5] while maintaining the vertically-inherited, intracellular endosymbiont. This is accomplished by dechorionation of the fertilized eggshell layer that removes the extracellular microbiome while maintaining the internal endosymbiont.

The α-proteobacterium *Wolbachia* is an archetypical intracellular reproductive endosymbiont of *D. melanogaster* and one of the most widespread, maternally-inherited, bacteria in multicellular eukaryotes [6]. *Wolbachia* are widely known for their diverse effects on manipulating host reproductive biology, in particular cytoplasmic incompatibility (CI), whereby crosses between symbiotic males and aposymbiotic females result in embryonic death [7–11]. Notably, the embryos are rescued if the females harbor *Wolbachia*. Thus, CI is a selfish

evolutionary adaptation that enhances the relative fitness of symbiont-transmitting females and

thus *Wolbachia*'s spread through host populations [8].

Here, we leveraged the *D. melanogaster* holobiont to disentangle the impact of the

extracellular microbiome on an endosymbiont's adaptation, densities, and fitness impacts on the

host. We address three central questions: Can an intracellular endosymbiont operate independently

of an animal's extracellular microbiome? If so, does the endosymbiont benefit from this altered

holobiont? What alterations in host and endosymbiont biology manifest in a strict, single-pair,

animal-microbe endosymbiosis relative to the complete holobiont setting?

Results

We generated *D. melanogaster* lines both with and without *Wolbachia* and the extracellular

microbiome (e.g., gut microbiome). *w*Mel *Wolbachia* is a member of the fly microbiome, yet for

simplicity and the purposes of the paper, we separately define the conventional microbiome as the

community of extracellular microbes and the endosymbiont *Wolbachia* as the intracellular

microbe. Thus, we adhere to the typical nomenclature system of conventional (CV) and germ-free

(GF) lines, both with *Wolbachia* (CV+ and GF+) and without it (CV- and GF-). We confirmed

Wolbachia infection status by PCR (Fig S1) and GF status of *Wolbachia*-infected and uninfected

flies by plating whole fly homogenates of GF reared flies (see methods) and demonstrating no

colony-forming units (CFUs) (Fig S2).

Absence of an extracellular microbiome promotes an endosymbiotic adaptation and host gene activation

To determine whether or not the CI adaptation is contingent on the extracellular fly

microbiome, we set up single-pair matings of <1-day-old, CI-inducing males carrying *Wolbachia*

(CV+ and GF+) and aposymbiotic females lacking *Wolbachia* (CV- and GF-). While CI levels by

CV+ males were moderate and typical for *w*Mel CI (median embryonic hatch rate = 36.17%), GF+

males notably induced 2.6-fold stronger CI (median = 13.66%) (Fig 1a). This strengthening of

embryonic lethality was fully rescued by CV+ females, demonstrating it was bona fide CI and not

an artifactual host sterility (Fig 1a).

The cell biological hallmarks of CI induction in male testes include abnormal histone

retention and decreased protamine deposition in developing sperm [7]. To evaluate if these gametic

modifications occur in GF+ males that express stronger CI, we first performed RNA-seq on late

male pupae when the histone-to-protamine transition commences in *D. melanogaster* [12,13]. The

core nucleosome structure is composed of H2A, H2B, H3, and H4 classes of histone proteins [1].

The expression of corresponding *His2Av*, *His2B*, *His3.3A*, *His3.3B*, and *His4r* genes were

upregulated in GF+ pupae compared to CV+ (Table S2). In *Drosophila* spermiogenesis, H3.3

variant along with the bulk histones are removed in the early canoe stage during the histone-to-

protamine transition [14]. Notably, expression of the *His3.3B* gene encoding the H3.3 class of

histone protein [15] was significantly upregulated in GF+ males relative to CV+ males (fold

change = 1.82, Log2FC= 0.87, p < 0.05) (Fig 1b). Moreover, expression of histone

methyltransferase 4-20 (*Hmt4-20*) that marks DNA for damage repair [16] was also significantly

upregulated (fold change = 1.82, Log2FC= 0.86), suggesting the host is attempting to keep up with

the elevated DNA damage in developing CI sperm [2]. Furthermore, comparative analysis indicated

that four of the five significant differentially-expressed genes involved in histone functions had an

enhanced relative gene expression ratio in GF+ vs. CV+ compared to GF- vs. CV-, suggesting that

Wolbachia can drive enhanced expression of CI-related cellular features in the GF+ state.

Further, there are four well-characterized protamine and protamine-like genes essential for

Drosophila mature sperm formation [17,18]. As expected for the CI-induced depletion of

protamine abundance in sperm, *ProtB*, *Prtl99C*, and *Mst77F* were significantly downregulated in

GF+ vs. CV+ male pupae (Fig 1b), and *ProtA* expression trended non-significantly lower (Table

S2). Consistent with the histone gene expression pattern above and protamine depletion as causal

to CI, three out of the four significant protamine genes had weakened transcription in GF+/CV+

relative to GF-/CV-.

Many male-specific transcripts involved in secretory proteins and spermatid maturation,

e.g., *Mst57Da*, *Mst57Db*, and *Mst57Dc*, were also significantly downregulated in GF+ compared

to CV+, collectively suggesting that CI-*Wolbachia* generally weaken gene expression of sperm-

associated proteins. Consistent with the aforementioned gene expression trends for CI-related gene

functions, 13 of the 15 significant *Mst* genes had a more pronounced fold change in GF+/CV+

relative to GF-/CV-.

Overall, the results specify that transcriptomic changes of key cell biological pathways in

CI links with augmented CI penetrance in flies that lack an extracellular microbiome. Interestingly,

GF- males without *Wolbachia* appear transcriptionally primed to express CI, but to a lesser degree

than when *Wolbachia* occur. This observation suggests moderate enhancements of transcriptome

changes coupled with expression of the CI genes in GF+ males may tip the cellular balance towards

enhanced CI expression.

Fig 1. CI penetrance is stronger in germ-free males and associates with gene expression variation in histone, protamine, and male-specific transcripts. (a) Germ-free males with *Wolbachia* (GF+) induce significantly higher CI or incompatibility (purple) than conventional (CV+) males. CI is a metric for embryonic viability that is measured by the percentage of embryos that hatch into larvae. Compatible crosses (grey bars) with normal embryonic hatching percentages are controls that express either no CI or rescue of CI. P-values were calculated based on a Kruskal-Wallis test followed by Benjamini, Krieger, and Yekutieli multiple correction test. The vertical bars represent the median value, and the error bars represent the 95% confidence interval. (b) Comparative transcriptomic analysis of GF+ vs. CV+ male pupae (bars filled with black dots) and GF- vs. CV- male pupae (bars without black dots) revealed many significant Differentially Expressed Genes, which were further screened to 24 genes that represented histone, protamine, and male-specific transcripts (MSTs). Significant differences are highlighted with a red colored asterisk (*). Notably, the histone-related genes genes (blue) were upregulated, while protamine (grey) and MST (yellow) genes were downregulated. Log2 fold change was calculated using DESeq2, and its default parameters; Walt test and Benjamini-Hochberg procedure were used to calculate p and adjusted p-values. The asterisk *, **, and *** represent adjusted p-values less than 0.05, 0.005, and 0.0005, respectively.

Cell biological modifications of host chromatin integrity occur in flies lacking an extracellular microbiome

To test if CI-related host nucleoprotein signatures occur in GF+ flies, we next compared histone abundance in spermatids and protamine deficiency in matured sperms of GF+ males.

Immunostaining <8-hour-old testes of virgin CI-inducing GF+ and CV+ males using a core histone

antibody showed significant histone retention in late canoe spermatid bundles, as expected

compared to GF- and CV-, respectively. The level of histone retention between the two groups

varied, with GF+ canoe spermatids exhibiting a median of 15.48% histone retention compared to

0% for GF- males (Fig 2a,b), and CV+ testes showing 57.14% histone retention compared to

12.14% for CV- males. As such, quantitative differences in histone retention are enhanced in CV+

males, while qualitative differences are larger in GF+ males. These differences are discussed

below.

Finally, cytochemical staining of mature sperm from <1-day-old virgin males with the

chromomycin A3 (CMA3) stain that binds to DNA regions devoid of protamines specified the

expected protamine deficiency is 1.45-fold elevated in mature sperm of GF+ compared to GF-

males (Fig 2b,d). These results are consistent with *Wolbachia* in a GF+ host modulating the

protamine deposition onto mature sperm in a slightly enhanced manner relative to the deficiency

of a 1.3 fold difference in CV+ vs. CV- males (Fig 2). Previous work demonstrated that enhanced

protamine deficiency is causal to increases in CI penetrance [3].

Fig 2. CI defects in the histone-to-protamine transition occur in germ-free and conventional

males. (a) Testes (N = ~10) from <8-hour-old virgin males of GF-, GF+, CV-, and CV+ lines were

dissected and immunostained to visualize and quantify histone abundance (magenta) in late-canoe

elongating spermatids. DAPI stain (grey) was used to label spermatid nuclei. (b) Total spermatid

bundles with DAPI signals and those with retained histones were manually counted and graphed.

Signal intensity in each sperm nucleus head was quantified in ImageJ and graphed. GF+ and CV+

males show a significantly higher number of spermatid bundles with retained histones compared

to their negative counterparts GF- and CV-, respectively. (c) Fully mature sperm were extracted

from <1 day-old virgin males of GF-, GF+, CV-, and CV+ lines and stained with fluorescent

CMA3 (green) for detection of protamine deficiency. (d) Quantification of the signal intensity

using ImageJ shows that GF+ and CV+ sperm have enhanced CMA3 intensity, consistent with

protamine deficiency levels compared to GF- and CV- negative controls, respectively. Vertical

bars represent the median, and error bars represent the standard deviation. P-values in the bracket

are calculated using the Mann-Whitney U test and the p-values outside of the bracket are calculated

using the Kruskal-Wallis test.

Endosymbiont density decreases in a microbiome-free host

Higher CI strength can depend on several, common factors including but not limited to an

increased density of *Wolbachia* [19], increased expression of *cif* genes [20], and/or reduced host

resistance to CI [21]. We measured relative *Wolbachia* density (*groEL)* compared to *D.*

melanogaster housekeeping gene *rp49*. Interestingly, *Wolbachia* density was reduced 4.58-fold in

strong CI-inducing GF+ males relative to CV+ males expressing moderate CI (Fig 3a). Moreover,

the relative expression of CI-associated genes *cifA* and *cifB* compared to *Wolbachia groEL* was

similar between CV+ and GF+ in adults (Fig 3b, c) and larvae (Fig S3), indicating that relative *cif*

gene transcription is independent of the *Wolbachia* density decline and loss of the extracellular

microbiome. Taken together, we conclude that GF+ males experience stronger CI not because they

experience an acute increase in symbiont number or gene expression (e.g., exemplary of a hyper-

pathogenic state), but instead because the hosts are physiologically more susceptible to the same

CI-induced modifications induced with lower *Wolbachia* densities in a microbiome-free or

conventional state.

Fig 3. *Wolbachia* density is markedly lower and relative *cif* gene expression is equal in a

germ-free host. (a) *Wolbachia* density was determined in CV+ and GF+ adult males by

quantifying single copy *Wolbachia* housekeeping gene *groEL* relative to *D. melanogaster*

housekeeping gene *rp49*. *Wolbachia* density was significantly lower in the GF+. (b) *cifA* and (c)

cifB gene expression were quantified by the ratio of cDNA from single copy *cifs* relative to the

Wolbachia housekeeping gene *groEL*. Gene expression was similar between GF+ and CV+,

indicating higher CI penetrance is not attributable to *cif* gene expression. The p-values are shown

for the non-parametric Kolmogorov-Smirnov test of equal distributions and Mann-Whitney U test

of population differences in the bracket.

Endosymbiont promotes extracellular microbiome growth

The microbiome of *D. melanogaster* is predominantly composed of 5 core culturable

species from the genera Lactobacillus and Acetobacter (Ludington & Ja, 2020). To assess if there

are microbiome abundance differences between conventionally reared flies with (CV+) and

without *Wolbachia* (CV-) that associated with CI penetrance variation, we plated whole adult male

homogenates onto *Lactobacillus* (MRS agar plates) and *Acetobacter* spp. (MYPL agar plates)

selective media (Aranda-Díaz et al., 2020) and counted the number of colony forming units

(CFUs) after 48 hours. CFUs of *Acetobacter* and *Lactobacillus* species in CV+ were notably 7.07-

and 9.78-fold higher compared to CV- (Fig 4a). Thus, an intact holobiont with the endosymbiont

contains nearly an order of magnitude more cells of the extracellular microbiome. Moreover, there

was a positive correlation between the overall CFUs of *Acetobacter* and *Lactobacillus* (Fig 4b)

that validated previous reports [22,23]. In parallel, we demonstrated no CFUs from negative-

control homogenates from GF+ and GF- males, as expected (Fig S2). Taken together with data in

Fig 3a, an intact holobiont contains markedly more cells of both the extracellular microbiome and

Wolbachia endosymbiont.

Based on this observation, we next tested if the abundance of cells from the extracellular

microbiome link with the penetrance of CI. Since GF+ males without a gut microbiome yield

significantly lower levels of embryos hatching and thus higher CI, we hypothesized that variation

in CI penetrance in CV+ males is associated with variation in the abundance of some gut microbes.

Indeed, there is a strong positive correlation between embryonic hatch rates from CI-causing CV+

males and the CFU count of *Lactobacillus* spp. (Fig 4d). Conversely, there is no association with

the CFU counts of *Acetobacter* spp. (Fig 4e), specifying then that higher *Lactobacillus* spp.

abundance links with reduced CI penetrance.

Fig 4. Holobionts with *Wolbachia* have higher colony forming units (CFUs) of gut bacteria,

of which *Lactobacillus* negatively associates with CI strength. (a) Homogenates of less than

one day old CV- (white circles) and CV+ males (orange circles) (N=5) in triplicates were spread

on MRS and MYPL plates. (b) *Wolbachia* infection (CV+) significantly increased the CFU load

of *Acetobacter spp*. and *Lactobacillus spp*., suggesting a positive interaction between the

endosymbiont and the microbiome. The p-values are calculated with a Mann-Whitney U test. (c)

The CFU of *Acetobacter* spp. and *Lactobacillus* spp. in CV- (data points filled in white) and CV+

(data points with filled in orange color) males have a significant positive correlation. (d) The CFU

of *Lactobacillus* spp. in CV+ males showed a significant correlation with the embryonic CI hatch

rate (HR) levels from the cross ♂CV+ x ♀wMel-. (e) Conversely, the CFU of *Acetobacter* spp. in

CV+ males did not exhibit a significant correlation with the hatch rate.

The extracellular microbiome and endosymbiont independently impact survival across development

Next, we examined the impact of the extracellular microbiome and endosymbiont on host

fitness and development across various life stages. We reared flies with (CV+ and GF+) and

without *Wolbachia* (CV- and GF-), and we measured the survival of the fly through metamorphosis

from embryo-to-larvae, larvae-to-pupae, and pupae-to-adults. GF+ flies with only *Wolbachia* had

a significantly higher embryonic hatch rate to larvae (71.43%) than GF- flies without *Wolbachia*

(66.95%, Fig 5a). CV+ flies similarly had higher survival (87%) than CV- flies lacking *Wolbachia*

(82%). Thus, the presence of *Wolbachia* generally enhanced egg-to-larvae survivability to a

modest degree. However, CV+ embryos exhibited higher embryo-to-larval hatching (87%) than

GF+ (71.43%), suggesting that a complete microbiome is beneficial for embryo-to-larvae

survivability. Thus, holobiont development with *Wolbachia* is optimized at this stage. From

larval-to-pupal (L-P) and pupal-to-adult (P-A) development stages, *Wolbachia* universally reduced

survival compared to their negative control counterparts, specifying a modest, negative fitness cost

at the larval stage onwards (4-13% reduction in survival, Fig 5b,c). Nonetheless, holobiont

survival was higher during the L-P stage (CV+ vs GF+, 11.67% increase, Fig 5b), indicating a

compensatory effect of the extracellular microbiome. Altogether, these findings specify that

Wolbachia's impacts on survival (beneficial to embryonic development, but harmful to subsequent

stages) are independent of those of the extracellular microbiome.

Fig 5. The fitness impacts of *Wolbachia* are independent of the extracellular microbiome and stage-dependent on metamorphosis. (a) The extracellular microbiome and *Wolbachia* exert independent and positive effects on embryo-to-larval survival, as calcuated by the number of surving larvae divided by the number of emryos laid. (b) The extracellular microbiome is beneficial to larval-to-pupal survival, while *Wolbachia* imposes a fitness cost in a microbiome-independent manner. (c) Similarly, *Wolbachia* lower pupal-to-adult survival percentage in a microbiome-independent manner. Each circle represents a family, and color denotes genotype status: orange bar (CV), blue bar (GF), open circle (no *Wolbachia*), and filled cirlce (*Wolbachia*). The p-values are shown were calculated with Kruskal-Wallis test followed by Benjamini, Krieger, and Yekutieli multiple correction test.

Discussion

How the extracellular microbiome influences intracellular symbiont biology is poorly known. Here we addressed this knowledge gap in a model system by examining the impacts of *D. melanogaster*'s extracellular microbiome on an intracellular endosymbiont and its CI adaptation, which manifests as a paternal-effect embryonic lethality when a *Wolbachia*-symbiotic male mates with an aposymbiotic female. We report five specific findings: (i) CI penetrance is stronger in GF+ *D. melanogaster* lacking an extracellular microbiome. (ii) CI strength in GF+ vs. CV+ males is independent of *Wolbachia* density and CI gene expression. (iii) transcriptomic reveal moderate upregulation of histone genes and downregulation of protamine and testes-specific genes that are consistent with known CI functions [7,24]. (iv) cell biological analyses in male reproduction

uncover complex quantitative and qualitative differences in histone retention and a slightly

increased protamine deficiency in germ-free males, as expected for elevated CI levels [3]. (v)

Notably, intact holobionts with both *Wolbachia* and gut bacteria have markedly higher microbial

densities, and penetrance of CI negatively correlates with a higher abundance of the gut bacteria

Lactobacillus. (vi) *Wolbachia* generally increase *D. melanogaster* fitness during embryo-to-larval

development but exert a fitness cost during larvae-to-pupae and pupa-to-adult development.

Influences of the gut microbiome on endosymbiont biology are generally rare because

experiments can be complex due to germ-free rearing, and binary interactions are simpler to study

than tripartite ones involving multi-species microbial communities. Notably, the gut bacterial

genus *Asaia* can impede the vertical transmission of *Wolbachia* in mosquitoes [25]. As such, the

nested symbiotic interactions between symbionts in different anatomical sites raise a fresh suite of

questions about how mechanistically extracellular and intracellular bacterial interactions impact

endosymbiotic functions. These nascent results imply ripple effects of an intricate network of

symbiotic interactions within holobionts that are likely to occur in other holobiont systems.

CI penetrance in various arthropods was previously known to vary with endosymbiont

density, *cif* gene expression, host age, and genetic background, among other traits. [26–30]. Here,

we add an unexpected dimension of the gut microbiome to the sources of variation, suggesting

heightened sensitivity of the germ-free or microbiome-depleted host to the CI molecular

mechanism. Specifically, the load of *Lactobacillus* spp. in males is inversely correlated to the CI

strength, the mechanism of which will be subject to future investigations.

The impact of the extracellular microbiome and endosymbiont on host fitness across

developmental stages revealed an intriguing cascade of intendent fitness impacts of the microbes

as well. *Wolbachia* promoted egg-to-larvae survivability, while hindering larval-to-pupal and

pupal-to-adult development stages. These effects were microbiome-independent, as evidenced by

similar outcomes in GF flies, suggesting that *Wolbachia*'s influences on fly fitness are conditional

on developmental stage in a gut microbiome-independent manner, likely due to the endosymbiont

residing outside the gut [31].

In conclusion, our study disentangled the intricate relationship of the gut microbiome's

impact on a key adaptation and replication of a common reproductive endosymbiont. This work

aligns with the growing recognition that sources of phenotypic variation, even in presumed simple

and binary symbiotic systems such as arthropods and their intracellular *Wolbachia*, can require a

broader view of the nested network of symbiotic interactions and functions in holobiont systems.

As biology is likely to observe increasing cases of biomolecular interactions of the intracellular

microbiome, extracellular microbiome, and the host, there will be new opportunities to understand

and apply knowledge that shapes evolutionary symbiotic outcomes and deployment of symbiotic

traits for positive outcomes in ecosystem health.

Materials and methods

Fly genotype, rearing condition, and food preparation

Drosophila melanogaster y^{1}w* (BDSC 1495) lines infected with *Wolbachia* strain *w*Mel were used for all the experiments. *w*Mel uninfected lines were previously established through tetracycline treatment [27]. *Wolbachia* infection status was confirmed through PCR using Wolb_F and Wolb_R3 primers (Table S1). Flies were reared in a 12:12 h light/dark cycle at 25°C and 60-70% relative humidity. The food for CV and GF flies has the same composition but differs in handling – food for GF flies was handled in a sterile condition, and the food for conventionally reared flies was handled in a non-sterile condition. The composition for 1L media consisted of 100gm yeast, 100g dextrose, 40g cornmeal, 7g agar, and 8.8 ml of 10% propionic acid dissolved in ethanol [32].

Rearing of flies to obtain mothers and fathers for hatch rate assay

Adult flies of both sexes were seeded and then cleared on the 4th day. The virgin female offspring (referred to as PGMs – paternal grandmothers) were collected on the 10th to 13th day. The PGM virgins were reared for 4-5 days to ensure virginity and to obtain moderate CI penetrance to test if germ-free treatment yields higher or lower CI penetrance [26]. Virgin PGMs were crossed with males to obtain embryos. In brief, ~30 individuals of 4-7 days old virgin PGMs were allowed to mate with 10 males overnight in a bottle without food secured with a grape plate. The next morning, the grape plate was replaced by a fresh one and incubated for 30 minutes. The grape plate was again replaced by a new one and incubated for 1 hour to obtain 1-hour embryos. In all

instances, grape plates were smeared with yeast. The embryos were processed for conventional

(CV) and germ-free (GF) treatment. On the 2nd and 9th days after the CV and GF treatments, larvae

and pupae survival were counted. On the 10th to 13th day, adult emergence was noticed. Males

(referred to as fathers) less than 24 hours old were collected to perform hatch rate assay.

Meanwhile, to obtain mothers, adult flies of both sexes were seeded for 4 days and cleared. The 1-

3 days old virgin female offspring (referred to as mothers) were collected on the 10th to 13th day.

Hatch rate (HR) assay

All crosses were performed with young <1-day-old males [26] and 3 days-old females. The

matings were performed in an 8 oz polypropylene bottle and secured with a grape plate. The flies

were reared overnight, and the grape plate was replaced by a fresh new grape plate. After 24 hours

of rearing, the flies were cleared, and the number of embryos on the grape plate was counted

(referred to as the 0-hour count). The embryos were incubated for an additional 36 hours, and the

unhatched embryos were counted (referred to as 36-hour count). Any plate with less than 25

embryos was discarded. The hatching percentage was determined by calculating the percentage of

embryos hatched on the 36-hour count compared to the 0-hour count.

Germ-free check and CFU quantification

To score the bacterial CFU in CV flies or to determine the germ-free status in GF flies, less than

24 hours old adult male flies (N=5) from each treatment were collected. However, single males

from each single-pair cross were collected to assess the bacterial CFU in males from the CI assay.

The flies were surface sterilized by washing twice in 70% Ethanol and once in 1x Phosphate-

buffered saline (PBS). The surface sterilized flies were crushed with a sterile micro pestle, diluted

in 300 uL 1xPBS, and 20 uL was used to spread onto MYPL (*Acetobacter* spp. selective media)

media (Aranda-Díaz et al., 2020) and *Lactobacillus* spp. selective media MRS (Sigma, #69964,

USA) plates. The MYPL media was composed of 1% D-mannitol (Sigma, #M4125, USA), 1%

Yeast extract (VWR, #J850, USA), 0.5% peptone (Thermo Fisher Scientific, # R451102, USA),

1% lactic acid (Sigma, #L6661, USA), 0.1% tween 80 (Thermo Fisher Scientific, #278632500,

USA), adjusted to pH 7 with NaOH pellets (Spectrum Chemical, #S1301, USA), and 1.5% agar

(Sigma, #A1296, USA) was added. The plates were incubated for 48 hours at 30°C and the CFU

per fly was quantified using the equation:

$$CFU\ per\ fly\ = \left. \frac{\substack{Total\ volume\ of \\ homogenate\ x\ CFU\ per\ plate}}{} \middle/ Total\ flies\ x\ Volume\ plated \right.$$

RNA/DNA isolation for RT/qPCR and transcriptome

Less than 6 hours old embryos (N=7), 2 days old larvae (N=7), late-stage pupae (N=1), and less

than 24 hours old adults (N=1) in triplicates from each treatment were used to extract RNA and

perform RT/qPCR and transcriptome from each treatment in triplicates. Embryos were crushed

using a needle, while larvae/pupae/adults were frozen in liquid nitrogen and crushed with a sterile

micro pestle. RNA was extracted using 1mL TRI Reagent (Sigma, #93289, USA) and 200 uL

chloroform (Sigma, #C2432, USA). Samples were incubated for 3 minutes in an ice block with

occasional mixing and centrifuged for 15minutes/12000rpm/4°C. The upper aqueous phase was

transferred to an RNase-free microcentrifuge tube containing 500 uL isopropanol & 1 uL

glycogen. Samples were incubated for 10 minutes in an ice block with occasional mixing, and

centrifuged for 10min/14000rpm/4°C. The supernatant was discarded, and the pellet was washed

in 500 uL 75% EtOH followed by centrifugation for 5min/7500rpm/4°C. The pellet was air-dried

inside the laminar flow hood at room temperature (RT) until it became translucent. The RNA pellet

was resuspended in 30 uL of nuclease-free water and incubated at 55°C for 15 minutes to dissolve

the pellet. Any contaminating DNA was eliminated by treating with DNA-free™ DNA Removal

Kit (Thermofisher Scientific, #AM9937, USA) as per the manufacturer's instructions. RNA was

confirmed DNA-free through negative PCR amplification of *rp49 Drosophila* gene. DNA-free

RNA samples were then purified using the ethanol precipitation method (see below) and stored at

-80°C until further use. DNA was isolated from the lower non-aqueous layers from the samples

used for RNA isolation using Qiagen PureGene Core Kit A kit (Qiagen, #158667, USA) following

manufacturer's instructions. However, during the DNA precipitation step we added 500 uL of

100% Isopropanol and 1 uL of glycogen to enhance the DNA recovery.

Ethanol precipitation

All the steps were handled in an ice block at all times. The sample volume was raised to 100 uL

using nuclease-free water. 10 uL of 3M Sodium Acetate (Thermofisher Scientific, # R1181, USA)

and 300 uL of 100% EtOH were added in the sample, mixed thoroughly & incubated at -80°C

overnight. The next morning, samples were centrifuged at full speed (14000 rpm) for 30 minutes

at 4°C and supernatant was discarded. 500 uL of freshly prepared 70% EtOH was added and

centrifuged at full speed for 20 mins at 4°C. The supernatant was discarded and pellet was air dried

for 30 minutes inside the biosafety cabinet until it became transparent. Pellet was resuspended in

20 ul nuclease-free water and incubate at 55°C for 15 minutes to fully dissolve. The purified RNA

sample was stored at -80°C until further use.

Sexing of pupae

410   Kl-5 primers were used to sex individual pupae collected at the late stage [33]. PCR conditions

411   used were 95°C for 2 mins and 30 cycles of 95°C for 30 sec, 56°C for 30 sec, and 72°C for 1 min,

412   final extension of 72°C for 5 mins.

Transcriptome

Samples were prepared by generating a uniquely indexed library from each of the RNA samples

isolated from male pupae using the Illumina Stranded Total RNA Prep with RiboZero Plus

Microbiome kit (Illumina, USA) and custom *D. melanogaster* probes for the rRNA depletion. The

libraries were pooled into a single pool containing an approximately equimolar amount of the

samples and outsourced to the Genomic Sciences Laboratory, North Carolina State University, for

sequencing using NovaSeq6000 S4 150 PE lane yielding ~25 million pairs of reads per sample on

average. The raw reads were subjected to quality control using Trimgalore v0.6.10 with a PHRED

score of 20 and minimum read length of 25 bp. The cleaned reads were mapped to *D. melanogaster*

reference genome (v.6.12) using STAR v2.7 to generate BAM files. The read counting was

performed using the BAM files with featureCounts v2.0.6. The raw reads were rarefied using

EcolUtils v0.1 with 100 permutations setting the lowest read (16M) as the threshold. The reads

were then processed for downstream differentially expressed genes (DEGs) using DeSeq2 v1.42.0

Histone staining and quantification

Less than 8 hours old males from the same population used for HR assay above were dissected to

collect testes in chilled 1x PBS solution and perform histone staining as described previously [7].

In brief, the testes were fixed for 30 minutes at RT in 4% formaldehyde prepared in 1x PBS with

subsequent washing with 1x PBST solution (1x PBS + 0.3% Triton X-100) and blocking in 1%

BSA in PBST at RT for an hour. Samples were incubated overnight with core histone antibody

(MilliporeSigma, Cat#MABE71, USA) (1:1,000) at 4°C in rotating shaker. After washing with 1x

PBST solution (1x PBS containing 0.1% Triton X-100), samples were incubated with goat anti-

mouse Alexa Fluor 488 secondary antibody (Thermo Fisher Scientific, Cat#A28175, 1:500 diluted

in 1x PBST) in the complete dark at RT for 4 hours. After washing in 1x PBST, samples were then

mounted in vectashield DAPI-containing mounting media (Vector Laboratories, Cat#H-1200-10,

USA) for imaging using Zeiss LSM 880 confocal microscope with constant exposure settings

across treatment groups. The total number of late canoe-stage sperm bundles and those that

retained histones were quantified in each testis, and the percentage of bundles with histone

retention was plotted in GraphPad Prism.

444   Protamine staining and quantification

Protamine staining and quantification was evaluated using chromomycin 3 (CMA3) stain as

described previously [7]. In brief, mature sperm were extracted from seminal vesicles of less than

8 hours old males in chilled 1x PBS. Samples were fixed in 3:1 vol/vol methanol:acetic acid for

20 mins at 4°C. Subsequently, the slide was air-dried and treated with 0.25 mg/mL of CMA3 in

McIlvain's buffer, pH 7.0, with 10 mM $MgCl_2$ for 20 minutes. The samples were washed with 1x

PBS and mounted in vectashield DAPI-containing mounting media. Imaging was performed using

Zeiss LSM 880 confocal microscope. Fluorescence quantification was performed by scoring

fluorescent pixels in arbitrary units (A.U.) within individual sperm heads using ImageJ. The flies

were reared at 21°C instead of 25°C for this assay and a hatch rate was also conducted in parallel

to compare protamine quantification with CI induction at 21°C.

Statistical analysis

A comparison of data between two groups was performed using Kolmogorov-Smirnov tests and

a Mann-Whitney U test. For more than two groups, the Kruskal Wallis test with Benjamini, Krieger

& Yekutieli p-value correction was performed. A correlation test was performed using the Pearson

correlation coefficient. To calculate tests and plot graphs, GraphPad Prism v10.0.0 was used.

Acknowledgments

The authors thank the Huck Institutes Genomics Core Facility (RRID:SCR_023645) at The

Pennsylvania State University for generating a uniquely indexed library from each of the RNA

samples and custom *D. melanogaster* probes for the rRNA depletion. The co-authors also thank

the Microscopy Core Facility (RRID:SCR_024457) for use of the Zeiss LSM 880 confocal

microscope. This work was supported by The Pennsylvania State University funds to the lab of

SRB.

References

472   1.   McFall-Ngai M, Hadfield MG, Bosch TCG, Carey H V., Domazet-Lošo T, Douglas AE,
473   et al. Animals in a bacterial world, a new imperative for the life sciences. Proceedings of
474   the National Academy of Sciences. 2013;110: 3229–3236. doi:10.1073/pnas.1218525110
475

476   2.   Bordenstein SR, Theis KR. Host Biology in Light of the Microbiome: Ten Principles of
477   Holobionts and Hologenomes. PLoS Biol. 2015;13: e1002226.
478   doi:10.1371/journal.pbio.1002226
479

480   3.   Brinker P, Fontaine MC, Beukeboom LW, Falcao Salles J. Host, Symbionts, and the
481   Microbiome: The Missing Tripartite Interaction. Trends Microbiol. 2019;27: 480–488.
482   doi:10.1016/j.tim.2019.02.002
483

484   4.   Masson F, Lemaitre B. Growing Ungrowable Bacteria: Overview and Perspectives on
485   Insect Symbiont Culturability. Microbiol Mol Biol Rev. 2020;84.
486   doi:10.1128/MMBR.00089-20
487

488   5.   Ludington WB, Ja WW. Drosophila as a model for the gut microbiome. PLoS Pathog.
489   2020;16: e1008398. doi:10.1371/journal.ppat.1008398
490

491   6.   Weinert LA, Araujo-Jnr E V., Ahmed MZ, Welch JJ. The incidence of bacterial
492   endosymbionts in terrestrial arthropods. Proceedings of the Royal Society B: Biological
493   Sciences. 2015;282: 20150249. doi:10.1098/rspb.2015.0249
494

495   7.   Kaur R, Leigh BA, Ritchie IT, Bordenstein SR. The Cif proteins from Wolbachia
496   prophage WO modify sperm genome integrity to establish cytoplasmic incompatibility.
497   PLoS Biol. 2022;20: e3001584. doi:10.1371/journal.pbio.3001584
498

499   8.   Shropshire JD, Leigh B, Bordenstein SR. Symbiont-mediated cytoplasmic
500   incompatibility: What have we learned in 50 years? Elife. 2020;9.
501   doi:10.7554/eLife.61989
502

503   9.   Hoffmann AA, Turelli M, Harshman LG. Factors affecting the distribution of cytoplasmic
504   incompatibility in Drosophila simulans. Genetics. 1990;126: 933–948.
505   doi:10.1093/genetics/126.4.933
506

507   10. Shropshire JD, Hamant E, Conner WR, Cooper BS. *cifB*- transcript levels largely explain
508   cytoplasmic incompatibility variation across divergent *Wolbachia*. PNAS Nexus. 2022;1.
509   doi:10.1093/pnasnexus/pgac099
510

511   11. Landmann F, Orsi GA, Loppin B, Sullivan W. Wolbachia-Mediated Cytoplasmic
512   Incompatibility Is Associated with Impaired Histone Deposition in the Male Pronucleus.
513   PLoS Pathog. 2009;5: e1000343. doi:10.1371/journal.ppat.1000343
514

515   12.   Ingham P. The development of Drosophila melanogaster. Trends in Genetics. 1994;10:
516   299. doi:10.1016/0168-9525(90)90017-Z
517
518   13.   Williamson A, Lehmann R. GERM CELL DEVELOPMENT IN *DROSOPHILA*. Annu
519   Rev Cell Dev Biol. 1996;12: 365–391. doi:10.1146/annurev.cellbio.12.1.365
520
521   14.   Rathke C, Baarends WM, Jayaramaiah-Raja S, Bartkuhn M, Renkawitz R, Renkawitz-
522   Pohl R. Transition from a nucleosome-based to a protamine-based chromatin
523   configuration during spermiogenesis in *Drosophila*. J Cell Sci. 2007;120: 1689–1700.
524   doi:10.1242/jcs.004663
525
526   15.   Akhmanova AS, Bindels PCT, Xu J, Miedema K, Kremer H, Hennig W, et al. Structure
527   and expression of histone H3.3 genes in *Drosophila melanogaster* and *Drosophila hydei*.
528   Genome. 1995;38: 586–600. doi:10.1139/g95-075
529
530   16.   Sakaguchi A, Karachentsev D, Seth-Pasricha M, Druzhinina M, Steward R. Functional
531   characterization of the Drosophila Hmt4-20/Suv4-20 histone methyltransferase. Genetics.
532   2008;179: 317–22. doi:10.1534/genetics.108.087650
533
534   17.   Jayaramaiah Raja S, Renkawitz-Pohl R. Replacement by *Drosophila melanogaster*
535   Protamines and Mst77F of Histones during Chromatin Condensation in Late Spermatids
536   and Role of Sesame in the Removal of These Proteins from the Male Pronucleus. Mol Cell
537   Biol. 2005;25: 6165–6177. doi:10.1128/MCB.25.14.6165-6177.2005
538
539   18.   Eren-Ghiani Z, Rathke C, Theofel I, Renkawitz-Pohl R. Prtl99C Acts Together with
540   Protamines and Safeguards Male Fertility in Drosophila. Cell Rep. 2015;13: 2327–2335.
541   doi:10.1016/j.celrep.2015.11.023
542
543   19.   Ikeda T, Ishikawa H, Sasaki T. Infection density of Wolbachia and level of cytoplasmic
544   incompatibility in the Mediterranean flour moth, Ephestia kuehniella. J Invertebr Pathol.
545   2003;84: 1–5. doi:10.1016/S0022-2011(03)00106-X
546
547   20.   Ritchie IT, Needles KT, Leigh BA, Kaur R, Bordenstein SR. Transgenic cytoplasmic
548   incompatibility persists across age and temperature variation in Drosophila melanogaster.
549   iScience. 2022;25: 105327. doi:10.1016/j.isci.2022.105327
550
551   21.   Bordenstein SR, Uy JJ, Werren JH. Host Genotype Determines Cytoplasmic
552   Incompatibility Type in the Haplodiploid Genus Nasonia. Genetics. 2003;164: 223–233.
553   doi:10.1093/genetics/164.1.223
554
555   22.   Newell PD, Douglas AE. Interspecies Interactions Determine the Impact of the Gut
556   Microbiota on Nutrient Allocation in Drosophila melanogaster. Appl Environ Microbiol.
557   2014;80: 788–796. doi:10.1128/AEM.02742-13
558

559   23.   Sommer AJ, Newell PD. Metabolic Basis for Mutualism between Gut Bacteria and Its
560   Impact on the *Drosophila melanogaster* Host. Appl Environ Microbiol. 2019;85.
561   doi:10.1128/AEM.01882-18
562

563   24.   Kaur R, McGarry A, Shropshire JD, Leigh BA, Bordenstein SR. Prophage proteins alter
564   long noncoding RNA and DNA of developing sperm to induce a paternal-effect lethality.
565   Science (1979). 2024;383: 1111–1117. doi:10.1126/science.adk9469
566

567   25.   Hughes GL, Dodson BL, Johnson RM, Murdock CC, Tsujimoto H, Suzuki Y, et al. Native
568   microbiome impedes vertical transmission of *Wolbachia* in *Anopheles* mosquitoes.
569   Proceedings of the National Academy of Sciences. 2014;111: 12498–12503.
570   doi:10.1073/pnas.1408888111
571

572   26.   Layton EM, On J, Perlmutter JI, Bordenstein SR, Shropshire JD. Paternal Grandmother
573   Age Affects the Strength of *Wolbachia* -Induced Cytoplasmic Incompatibility in
574   Drosophila melanogaster. mBio. 2019;10. doi:10.1128/mBio.01879-19
575

576   27.   LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, et al.
577   Prophage WO genes recapitulate and enhance Wolbachia-induced cytoplasmic
578   incompatibility. Nature. 2017;543: 243–247. doi:10.1038/nature21391
579

580   28.   Bourtzis K, Nirgianaki A, Markakis G, Savakis C. Wolbachia Infection and Cytoplasmic
581   Incompatibility in Drosophila Species. Genetics. 1996;144: 1063–1073.
582   doi:10.1093/genetics/144.3.1063
583

584   29.   Cooper BS, Ginsberg PS, Turelli M, Matute DR. Wolbachia in the *Drosophila yakuba*
585   Complex: Pervasive Frequency Variation and Weak Cytoplasmic Incompatibility, but No
586   Apparent Effect on Reproductive Isolation. Genetics. 2017;205: 333–351.
587   doi:10.1534/genetics.116.196238
588

589   30.   Shropshire JD, On J, Layton EM, Zhou H, Bordenstein SR. One prophage WO gene
590   rescues cytoplasmic incompatibility in *Drosophila melanogaster*. Proceedings of the
591   National Academy of Sciences. 2018;115: 4987–4991. doi:10.1073/pnas.1800650115
592

593   31.   Simhadri RK, Fast EM, Guo R, Schultz MJ, Vaisman N, Ortiz L, et al. The Gut
594   Commensal Microbiome of Drosophila melanogaster Is Modified by the Endosymbiont
595   *Wolbachia*. mSphere. 2017;2. doi:10.1128/mSphere.00287-17
596

597   32.   Suito T, Nagao K, Juni N, Hara Y, Sokabe T, Atomi H, et al. Regulation of
598   thermoregulatory behavior by commensal bacteria in *Drosophila*. Biosci Biotechnol
599   Biochem. 2022;86: 1060–1070. doi:10.1093/bbb/zbac087
600

601   33.   Lott SE, Villalta JE, Schroth GP, Luo S, Tonkin LA, Eisen MB. Noncanonical
602   Compensation of Zygotic X Transcription in Early Drosophila melanogaster Development
603   Revealed through Single-Embryo RNA-Seq. PLoS Biol. 2011;9: e1000590.
604   doi:10.1371/journal.pbio.1000590

Supporting information

S1 Fig. Molecular confirmation of *Wolbachia* infection status of the parental lines *w*Mel-

(CV-) and *w*Mel+ (CV+).

S2 Fig. Conformation of microbiome absence and presence by colony forming units (CFUs)

in GF and CV lines.

S3 Fig. The larvae-to-pupae fitness cost in a germ-free state is neither due to *Wolbachia*

density nor *cif* gene expression.

S1 Table: Primers used for relative quantification of Wolbachia infection check, sexing of

pupae, and quantification of cifA, cifB, and Wolbachia density.

S2 Table: Differentially-expressed histone, protamine, and MST genes in male pupae.

(a)

Embryos hatched (%)

- Incompatible cross
- Compatible cross

<0.0001

<0.0001

0.0371

<0.0001

Cross (male x female)

CV+ x CV−   GF+ x CV−   GF− x CV−   CV+ x CV+   GF+ x CV+

(b)

Relative gene expression changes (Log2Fold)

Comparison
- GF−/CV−
- GF+/CV+

Markers
- Histones
- MSTs
- Protamines

Differentially expressed genes (in male pupae)

Hers, His2Av, His3.3B, Hmt4-20, His4r, BigH1, Hinfp, Utx, His3.3A, Mst57Da, Mst57Db, Mst57Dc, Mst84Dc, Mst98Ca, Mst98Cb, Mst77Y-13, Mst87F, Mst84Db, Mst84Da, Mst84Dd, Mst84B, Mst89B, Mst36Fb, Mst36Fa, Mst33A, Mst77Y-7, MstProx, Mst77Y-16Psi, Mst27D, mst, Mst77Y-4, Mst77Y-8Psi, Mst85C, Prtl99C, ProtB, Mst77F, ProtA

(a)

	α-His	DAPI	Merged
GF-			
GF+	Elongating spermatid histone		

	α-His	DAPI	Merged
CV-			
CV+			

(b)

% of spermatid bundles with histone retention

0.0334

GF-   GF+

% of spermatid bundles with histone retention

0.0011

CV-   CV+

(c)

CMA3

(d)

Protamine-deficient CMA3 fluorescence per sperm

<0.0001

GF-   GF+

Protamine-deficient CMA3 fluorescence per sperm

0.0029

CV-   CV+

(a) *Wolbachia* **density in adults**

(b) *cifA/groEL* **in adults**

(c) *cifB/groEL* **in adults**