1

Gut-Germline Axis:

A Reproductive Endosymbiont's Adaptation

is Modulated by the Gut Microbiome

Madangchanok Imchen[1], Rupinder Kaur[1], Darlene Huilin Zhou Luo[1], Mia W. Bitman[1], Emilie Lefoulon[1], Angelina McGarry[1], Sarah R. Bordenstein, and Seth R. Bordenstein[1]*

[1]Departments of Biology and Entomology, One Heath Microbiome Center, Huck Institutes of the Life Sciences, The Pennsylvania State University, USA

*Corresponding author: srb6251@psu.edu

Keywords: cytoplasmic incompatibility, microbiome, intracellular bacteria, endosymbiosis, *Drosophila melanogaster*

Abstract

Holobionts contain extra- and intracellular microbes that are typically studied in symbiotic and anatomical siloes. Here we demonstrate a previously unknown impact of the extracellular gut microbiome on the adaptation of a reproductive endosymbiont. Specifically, a paternal-effect embryonic lethality caused by *Wolbachia*-induced cytoplasmic incompatibility (CI) is attenuated by the gut microbiome. Gut microbiome-mediated variation in the CI phenotype causes key gene expression changes in the symbiosis and defining cell biological defects in the evolutionary conserved histone-to-protamine transition that drives CI. Reintroducing the gut microbiome into germ-free flies recapitulates the CI attenuation. Notably, the integrated holobiont exhibits significantly elevated gut microbiome and endosymbiont densities relative to isolated constituent rearing, highlighting a synergistic relationship. Finally, impacts of the symbionts on survival are independent and inversely related throughout development, and fitness changes are recapitulated when the gut microbiome is restored. Altogether, this study uncovers a gut-germline axis in which bacteria shape an entangled network of symbiotic functions within a holobiont. Interorgan, symbiotic relationships have far-reaching implications for the study of trait variation.

Main

Animal and plant holobionts harbor consortia of microorganisms, encompassing endosymbionts and extracellular microbes with varying degrees of interdependencies[1–3]. The co-existence of these microbial classes with different lifestyles and functions can hypothetically exert influences in competitive, collaborative, or stochastic ways[4]. If such trade-offs and synergies occur across diverse symbiotic types and host anatomical sites, then they can drive cascading, ripple effects of symbiosis throughout the entire holobiont. However, such effects have not been a focal point of study owing to two primary factors. First, conventional microbiome elimination methods are indiscriminate, affecting both symbiont classes. Second, the obligatory intracellular nature of endosymbionts often precludes their cultivation[5]. Here, we overcome these limitations in a system that contains a vertically-inherited, intracellular endosymbiont and an environmentally-acquired gut microbiome.

In the *Drosophila melanogaster* holobiont, the α-proteobacterium *Wolbachia* is an archetypal reproductive endosymbiont of *D. melanogaster* and one of the most prevalent maternally-inherited bacteria in animals[6]. The endosymbiont is widely known for replication in the reproductive tract and diverse manipulations of arthropod reproductive biology. The most common adaptation that *Wolbachia* deploy is cytoplasmic incompatibility (CI), a phenomenon in which crosses between symbiotic males and aposymbiotic females result in embryonic lethality[7–11]. Embryos are rescued if the females harbor *Wolbachia*. Consequently, CI selfishly enhances the relative fitness of symbiont-transmitting females, thereby spreading *Wolbachia*[8]. In addition, a combined body of evidence specifies that *D. melanogaster* houses a simple and beneficial gut microbiome of a handful of core species that can be easily removed[12]. *Wolbachia* are absent from the *D. melanogaster* gut lumen[13], suggesting there is no obvious route for an interaction with its

simple gut microbiome. However, indirect associations occur wherein *Wolbachia* associate with

gut microbiome composition[13,14].

The last decade of work in host-microbial interactions has featured a large increase in

functional studies of binary symbioses or experimental studies of single anatomical sites of

symbiotic interactions. This wave of experiments has inspired a good deal of new conceptual and

theoretical work on the complexity of holobionts that house a rich aggregate of microbes across

the body. Here we address three fundamental questions to unravel symbiotic functions in a

holobiont context: (i) Can a reproductive endosymbiont (*Wolbachia*) function autonomously of

the gut microbiome? (ii) What effects does the gut microbiome have on densities, penetrance,

gene expression, and cell biological underpinnings of the endosymbiont? And (iii) How is host

fitness independently and synergistically shaped by the gut and reproductive microbiome? By

tackling these inquiries, this study reveals an overlooked, functional interplay between bacteria

in different host organs and their combined influence on holobiont biology.

Results

We generated *D. melanogaster* with and without *Wolbachia* and the gut microbiome by

leveraging dechorionation of the fertilized eggshell layer in *Drosophila melanogaster,* a process

that eliminates the extracellular gut microbiome[12] while preserving the vertically-inherited,

reproductive endosymbiont. For simplicity and the purposes of the paper, we separately define

the conventional microbiome as the community of gut microbes. Thus, we adhere to the typical

nomenclature of conventional (CV) and germ-free (GF) hosts, both with *Wolbachia* (CV+ and

GF+) and without it (CV- and GF-) (Fig 1A). We confirmed *Wolbachia* infection status by PCR

and GF status by plating whole fly homogenates (see methods) to detect the presence/absence of

colony-forming units (CFUs) (Fig S1). We also confirmed that the CV+/- lines have the same

genetic makeup (Table S1).

83
84
The adaptation and host regulation of a reproductive endosymbiont

are altered by the gut microbiome

If the *Wolbachia* reproductive endosymbionts in the reproductive tissue are modulated by

the presence of a gut microbiome, then we expect that CI will increase or decrease in the absence

of a gut microbiome. To test this prediction, we mated CI-inducing males with *Wolbachia* (CV+

and GF+) and aposymbiotic females without *Wolbachia* (CV-) in single, pairwise matings. All

males in these CI assays were <1-day-old since they cause a strong CI lethality[15], which is

measured as a percent reduction of embryonic hatching into larvae. In multiple experiments,

GF+ males induced 2.6-fold stronger CI (median embryonic hatch rate = 13.66%) relative to

CV+ males (median = 36.17%) (Fig 1B). This increase in embryonic lethality was fully rescued

by CV+ females, confirming the effect is bona fide CI rather than artifactual host sterility. These

results specify that *Wolbachia* can alone induce CI and the penetrance of the trait is attenuated by

the presence of the gut microbiome. To validate this interpretation, we conducted a microbiome

transplant experiment into GF+ males via fecal-oral transmission. Restoration of the CV+ native

microbiome likewise attenuated the enhanced CI penetrance (Fig 1C). Thus, the enhanced CI

penetrance in GF+ was reversible with a 4.72-fold decrease that matched that of CV+ CI (Fig

1C), demonstrating the gut microbiome's causal role in modulating CI-*Wolbachia*.

Given that *Wolbachia* initiate CI with nucleoprotein modifications in developing

sperm[7,16], we next reasoned that the fly transcriptome could be a sensitive measure of host

regulatory changes in GF+ vs. CV+ males. Indeed, expression analyses revealed changes to CI-

defining gametic modifications in the histone-to-protamine transition in late male pupae, which

is when this protein transition begins in *D. melanogaster*[17,18]. The core nucleosome structure is

composed of histone classes H2A, H2B, H3, and H4[19], and expression of the *His2Av*, *His2B*,

His3.3A, *His3.3B*, and *His4r* genes were upregulated in GF+ pupae that induce stronger CI

compared to CV+ (Fig 1D; Table S2). In total, expression of 8 out of 10 histone associated genes

were more elevated in GF+/CV+ compared to GF-/CV-. These results are consistent with the

increased histone retention of CI sperm[7]. In *Drosophila* spermiogenesis, the histone variant *H3.3*

along with the bulk histones are removed in the early canoe stage during the protein transition[20].

Expression of the *His3.3B* gene encoding the H3.3 histone protein[21] was significantly

upregulated in GF+ males relative to CV+ males (fold change = 1.84, Log2FC= 0.88 p < 0.05).

Moreover, expression of histone methyltransferase 4-20 (*Hmt4-20*) that marks DNA for damage

repair[22] was also significantly upregulated (fold change = 1.83, Log2FC= 0.87), which is

consistent with the fly quiescing elevated DNA damage also known to occur in developing CI

sperm[23]. Consequently, *Wolbachia* enhance gene expression of histone-related CI features in the

GF+ state.

There are four well-characterized protamine and protamine-like genes essential for

Drosophila mature sperm formation[24,25]. Consistent with CI's causal role in depleting protamine

abundance in sperm[7,16], *ProtB*, *Prtl99C*, and *Mst77F* were significantly downregulated in GF+

vs. CV+ male pupae (Fig 1C), and *ProtA* expression trended lower (Table S2). Three out of the

four significant protamine genes expressed lower transcription in GF+/CV+ relative to GF-/CV-.

Moreover, 13 of the 15 significant male-specific transcript (*Mst*) genes, involved in secretory

proteins and spermatid maturation, e.g., *Mst57Da*, *Mst57Db*, and *Mst57Dc*, had a more

pronounced change in GF+/CV+ relative to GF-/CV-. Thus, *Wolbachia* generally weaken gene

expression of these sperm-associated proteins. Collectively, results show that transcriptomic

alterations in key CI cellular pathway genes link to the augmented CI penetrance in gut

microbiome-deficient flies.

Finally, analysis of the top 25 differentially-expressed host transcripts between GF vs CV

flies revealed gene expression changes associated with male accessory gland proteins that are

transferred to females during mating (*Mst57Da*, CG9997, and CG9029) (Fig S2). There were

also transcripts involved in chitin-based cuticle development, cuticle and trachea development[26],

light-sensitive photoreceptor rhodopsins, and unannotated markers. Taken together, the results

imply that GF+ flies are transcriptionally altered in diverse ways, with mechanistic relevance to

Wolbachia pathology and the acute sperm development changes that underpin CI.

Endosymbiont-driven sperm chromatin remodeling is sensitive to the gut microbiome

Based on the sperm-associated regulatory changes above, we next leveraged cell

biological markers of chromatin integrity to test if CI-nucleoprotein signatures increase in GF+

flies during the critical sperm developmental stages when the histone-to-protamine transition

takes place (Fig 2A). We first examined the total histone levels in GF+ spermatids using a core

histone antibody and show histone retention in both GF+ and CV+ males in late canoe spermatid

bundles compared to GF- and CV-, respectively. The level of histone retention between the two

groups varied, with GF+ canoe spermatids exhibiting a median of 15.48% histone retention

compared to 0% for GF- males (Fig 2B), and CV+ testes showing 57.14% histone retention

compared to 12.14% for CV- males. As such, quantitative differences in histone retention are

enhanced in CV+ males, while qualitative differences are greater in GF+ males.

Chromomycin A3 (CMA3) stain binds DNA regions devoid of protamines and assays if

CI sperm experience protamine deficiency. Cytochemical staining of mature sperm from <1-day-

old virgin males showed that protamine deficiency is 1.45-fold elevated in mature sperm of GF+

compared to GF- males (Fig 2C), which is slightly larger than the 1.3-fold difference in CV+ vs.

CV- males (Fig 2C). Taken together, these results link CI variation and the transcriptome

changes to sperm integrity changes *in situ*, namely that *Wolbachia* in a GF+ host enhance several

central dogma underpinnings of CI. Our previous genetic work demonstrated that enhanced

protamine deficiency is causal to increases in CI penetrance[23].

The gut microbiome lowers densities and transcriptional activity of the endosymbiont

CI strength varies with densities of *Wolbachia*[27], expression of the *cifA* and *cifB* genes[28],

host genetic effects, among other facrtors[29]. To examine the impacts of the gut microbiome on

Wolbachia densities, we quantified relative *Wolbachia* counts (*groEL)* normalized to *D.*

melanogaster housekeeping gene *rp49*. Unexpectedly, *Wolbachia* density was reduced 2.07-fold

($p<0.05$) in strong CI-inducing GF+ males compared to CV+ males, but not in larvae (Fig 3A,B

and S3). Furthermore, the relative gene expression of *cifA* and *cifB* compared to *Wolbachia*

groEL was similar between CV+ and GF+ in adults (Fig 3C,D) and larvae (Fig 3E,F), indicating

that *cif* transcription is independent of the *Wolbachia* density decline and loss of the extracellular

microbiome. These findings indicate that an intact CV+ fly holobiont has higher *Wolbachia*

copies than a GF+ fly with *Wolbachia*, and the enhanced CI in GF+ males is surprisingly not

attributable to increased symbiont abundance or CI gene expression. Since there were no

expression changes to the CI genes, we investigated *Wolbachia* transcriptome variation by

analyzing *Wolbachia* reads in the gene expression dataset above across embryos, larvae, male

pupae, and male adults. During the larval stage when the testes develop in flies and both the

spermatogonia and spermatocytes take form and store RNA transcripts for remodeling sperm

development, *Wolbachia* in GF+ significantly overexpress ribosomal RNAs and mRNAs relative

to CV+. The highest number of annotated *Wolbachia* functions includes host-microbe

interactions, secretion system and virulence effectors, as well as ankyrin-containing proteins

widely involved in host-microbe relationships (Fig 3G, blue category). Changes in protein

synthesis transcripts and membrane transport further support that *Wolbachia* are transcriptionally

altered in GF+ larvae to hypermodify host biology.

Finally, the microbiome of *D. melanogaster* is predominantly composed of 5 core

culturable species from the genera *Lactobacillus* and *Acetobacter*[12]. To assess if *Wolbachia* also

have a reverse impact on gut bacterial abundance between CV+ and CV- flies, we plated whole

adult male homogenates onto *Lactobacillus* (MRS agar plates) and *Acetobacter* spp. (MYPL

agar plates) selective media[30] and counted the number of colony forming units (CFUs) after 48

hours (Fig 4A). CFUs of *Acetobacter* and *Lactobacillus* species in CV+ were notably 7.07- and

9.78-fold higher compared to CV- (Fig 4B). Thus, an intact holobiont with the endosymbiont

contains nearly an order of magnitude more cells of the extracellular microbiome. Moreover,

there was a positive correlation between the overall CFUs of *Acetobacter* and *Lactobacillus* (Fig

4C) that validated previous reports[31,32]. In parallel, we demonstrated no CFUs from negative-

control homogenates from GF+ and GF- males, as expected (Fig S1). Taken together with data in

Fig 3, the intact holobiont contains markedly more cells of both the extracellular gut microbiome

and *Wolbachia* endosymbiont.

The gut microbiome and endosymbiont independently impact survival across development

By disentangling the gut microbiome and endosymbiont into constituent parts relative to

the intact holobiont, we tested how the gut microbiome and *Wolbachia* differentially impact host

fitness (survival) across metamorphosis. GF+ flies had a significantly higher embryonic hatch

rate to larvae (71.43%) than GF- flies without *Wolbachia* (66.95%, Fig 5A). CV+ flies similarly

also had higher embryo-to-larvae survival (87%) than CV- flies lacking *Wolbachia* (82%). Thus,

the presence of *Wolbachia* modestly enhanced early life survivability. Moreover, CV+ embryos

with higher *Wolbachia* counts also exhibited higher embryo-to-larval hatching (87%) than GF+

(71.43%), indicating that the complete microbiome and/or its possible impacts on an increase in

Wolbachia density are beneficial for embryo-to-larvae survivability. From larval-to-pupal (L-P)

and pupal-to-adult (P-A) development stages, *Wolbachia* generally reduced CV and GF host

survival compared to their negative control counterparts, specifying a modest, negative fitness

cost at the larval stage onwards (4-13% reduction in survival, Fig 5B,C). Nonetheless, holobiont

survival was higher during the L-P stage (CV+ vs GF+, 11.67% increase, Fig 5B), indicating a

compensatory effect of the extracellular microbiome. Overall, GF+ embryo-to-adult survival

(38.56%) was noticeably lower than CV+ survival (51.56%) (Fig 5D), which was causally

compensated upon microbiome reintroduction into GF+ (GF$_M$+ egg-to-adult survival = 51.35%)

(Figure 5E). Altogether, these findings specify independent impacts of the symbiont classes on

survival across development, namely *Wolbachia* are beneficial to embryonic-to-larval

development and harmful to subsequent stages, while the gut microbiome increases embryo-to-

larval and larval-to-pupal survival during metamorphosis. Such effects can shape host fitness and

will depend on genetic variation of both the host[33] and the endosymbiont.

Discussion

Holobionts are units in biology's hierarchy of organization that encompass the host

organism and its associated microbial communities[3], including both intracellular

(endosymbionts) and extracellular microbes. The interwoven existence of these distinct

microbial groups spanning different anatomical locations, each with unique lifestyles and

functional roles, is recognized after decades of study but rarely studied in a functional context.

Here we addressed this knowledge gap by examining the adaptation, densities, transcriptomes,

host modifications, and fitness impacts of a reproductive endosymbiont in the presence and

absence of a gut microbiome. The accumulated lines of evidence add an unexpected dimension

of the causes of variation that affect intimate, host-microbial symbioses.

We report five main findings: (i) CI penetrance is heightened in a host without a gut

microbiome. (ii) Transplantation of the gut microbiome by fecal ingestion into a germ-free host

causes the reduced CI penetrance. (iii) Host gene expression and cell biological hallmarks in the

histone-to-protamine transition of sperm development underpin CI variation in germ-free vs.

conventional hosts[7,23]. (iv) *Wolbachia* gene expression shows host interactions are linked to the

heightened CI effect. (v) Intact holobionts have markedly higher microbial densities of both the

endosymbiont and gut microbiome relative to when one microbial class is missing, suggesting

synergies on their within-host fitness.

The study of within-host symbiotic dynamics, especially across organs, remains in its

infancy. Not surprisingly, much of the history of studying host-microbe interactions has

emphasized simple, binary, host-microbe interactions or single, anatomical sites in the case of

the host-associated microbial communities. Consequently, assumptions prevail that the causes of

functional variation in host-microbe interactions are due to genetic and environmental variation

of the interacting partners. Indeed, CI penetrance in various arthropods can vary with

endosymbiont density, *cif* gene expression, host age, and host genetic background[34–43]. This

broad assumption is now questionable in light of the complexity of holobiont systems with

varying numbers and types of microorganisms across body sites that can affect the cell biology

and physiology of the host. In mosquitoes, the gut bacterial genus *Asaia* can impede the vertical

transmission of *Wolbachia*[44]. As such, nested symbiotic interactions between symbionts in

different anatomical sites raise a fresh perspective and suite of questions on how one microbial

class impacts the functions of another across the biogeography of the body. If these effects are

widespread as we suspect, it might also aid explanations of the missing heritability problem in

symbioses in which host genetic variation insufficiently explains variation in microbial traits.

In conclusion, this work disentangles the intricate relationship of the gut microbiome's

impact on a key adaptation and replication of a common reproductive endosymbiont. It aligns

with the growing recognition that sources of phenotypic variation, even in presumed simple and

binary symbiotic systems, such as arthropods and their intracellular *Wolbachia*, can require a

broader view of the nested network of symbiotic interactions and functions in holobionts[3]. A gut-

germline axis for microbial symbionts can also, in principle, shape evolutionary outcomes and

symbiotic traits used in the biocontrol of arthropods.

Materials and methods

Fly genotype, rearing condition, and food preparation

Drosophila melanogaster y^{1}w* (BDSC 1495) lines infected with *Wolbachia* strain *w*Mel were used for all the experiments. *D. melanogaster* white and yellow genetic background is extensively used in cytoplasmic incompatibility (CI) studies[7,35,36,40,46,47]. Furthermore, the *Wolbachia* strain present in this line has been effectively utilized in vector control initiatives by the World Mosquito Program[46,48]. *w*Mel uninfected lines were previously established through tetracycline treatment[35]. *Wolbachia* infection status was confirmed through PCR using Wolb_F and Wolb_R3 primers (Table S3). Flies were reared in a 12:12 h light/dark cycle at 25°C and 60-70% relative humidity. The food for both CV and GF flies had the same composition and was autoclaved. However, the autoclaved food for GF flies was maintained under sterile conditions, while the food for conventionally reared flies was maintained under non-sterile conditions. The composition for 1L media consisted of 100g yeast, 100g dextrose, 40g cornmeal, 7g agar, and 8.8 mL of 10% propionic acid dissolved in ethanol[49].

Rearing of flies to obtain mothers and fathers for hatch rate assay

Adult flies of both sexes were kept for four days for oviposition. The virgin female offspring (referred to as PGMs – paternal grandmothers) were collected on the 10th to 13th day. The PGM virgins were kept in isolation for 4-5 days to ensure virginity and to obtain moderate CI penetrance to test if germ-free treatment yields higher or lower CI penetrance[34]. Virgin PGMs were crossed with males to obtain embryos. In brief, ~30 virgin PGMs, aged 4-7 days, were

allowed to mate overnight with 10 males in an empty bottle secured with a grape plate smeared

with yeast. The next morning, the grape plate was replaced by a fresh one and incubated for 30

min. The grape plate was again replaced by a new one and incubated for 1 hour to obtain 1-hour

embryos. In all instances, grape plates were smeared with yeast. The embryos were processed for

conventional (CV) and germ-free (GF) treatment[51]. The GF treatment was carried out as

previously described[50]. Briefly, embryos were dechorionated twice using 10% bleach, followed

by a single wash in 70% ethanol and three washes in PBS containing 0.1% Triton X (Fig S4). On

the 2^{nd} and 9^{th} days after the CV and GF treatments, larvae and pupae survival were counted,

respectively. On the 10^{th} to 13^{th} day, adult emergence was noticed. Males (referred to as fathers)

less than 24 hours old were collected to perform hatch rate assay. Meanwhile, parental flies of

both sexes were reared on food for 4 days to lay fertilized eggs, after which they were removed.

The F1 virgin female offspring (referred to as mothers) were collected on the 10^{th} to 13^{th} day.

CI assays

All crosses were performed using <1-day-old males[34] and 3-day-old females in a conventional

incubator under a 12:12 h light/dark cycle at 25°C with 60–70% relative humidity[16]. The matings

were performed in an 8 oz polypropylene bottle and secured with a grape plate. The flies were

kept overnight, and the grape plate was replaced by a fresh new grape plate. After 24 hours, the

flies were cleared, and the number of embryos on the grape plate was counted (referred to as the

0-hour count). The embryos were incubated for an additional 36 hours, and the unhatched

embryos were counted (referred to as 36-hour count). Any plate with less than 25 embryos was

discarded. The hatching percentage was determined by calculating the percentage of embryos

hatched on the 36-hour count compared to the 0-hour count. CI (incompatible) crosses were

performed between *Wolbachia*-infected males (CV+ and GF+) and *Wolbachia*-uninfected

females (CV-). Compatible rescue crosses were performed between *Wolbachia*-infected males

(CV+ and GF+) and *Wolbachia*-infected females (CV+). Using CV+ females for rescue

experiments serves as a persistent control that ensures consistency in rescue studies[7,23,34,35,38,40]

and avoids confounders by adding the extra variable of GF+ females.

Germm-free check and CFU quantification

To score the bacterial CFU in CV flies or to determine the germ-free status in GF flies, less than

24 hours old adult male flies (N=5) from each treatment were collected. However, single males

from each single-pair cross were collected to assess the bacterial CFU in males from the CI

assay. The flies were surface sterilized by washing twice in 70% Ethanol and once in 1x

Phosphate-buffered saline (PBS). The surface sterilized flies were crushed with a sterile micro

pestle, diluted in 300 uL 1xPBS, and 20 uL was used to spread onto MYPL (*Acetobacter* spp.

selective media) media (Aranda-Díaz et al., 2020) and *Lactobacillus* spp. selective media MRS

(Sigma, #69964, USA) plates. The MYPL media was composed of 1% D-mannitol (Sigma,

#M4125, USA), 1% yeast extract (VWR, #J850, USA), 0.5% peptone (Thermo Fisher Scientific,

R451102, USA), 1% lactic acid (Sigma, #L6661, USA), 0.1% tween 80 (Thermo Fisher

Scientific, #278632500, USA), adjusted to pH 7 with NaOH pellets (Spectrum Chemical,

#S1301, USA), and 1.5% agar (Sigma, #A1296, USA) was added. The plates were incubated for

48 hours at 30°C and the CFU per fly was quantified using the equation:

$$CFU\ per\ fly\ = \left. \frac{Total\ volume\ of}{homogenate\ x\ CFU\ per\ plate} \middle/ Total\ flies\ x\ Volume\ plated \right.$$

Microbiome reintroduction

A natural method of gut microbiome transfer based on fecal material ingestion followed a

previously published protocol[51]. First, 50 adult *w*Mel+ males of random age were left in a GF

vial for 7 min. Second, vials were flipped upside down for another 7 min so that the fecal

material was deposited onto the plug and food surface. Finally, the resulting vials were externally

surface sterilized by spraying them with ample 70% EtOH before adding the GF embryos to

grow and ingest the bacteria.

RNA/DNA isolation for RT/qPCR and transcriptome

RNA was extracted and used for RT-qPCR (N=6 to 12 replicates) and RNA-Seq (N=3

replicates) from samples collected at various developmental stages for each treatment: 2-day-old

larvae (N=7), late-stage pupae (N=1), and less than 24-hour-old adults (N=1). Whole body

larvae/pupae/adults were frozen in liquid nitrogen and crushed with a sterile micro pestle. For

Wolbachia density measurements, the adult abdomen was dissected and processed. RNA was

extracted using 1 mL TRI Reagent (Sigma, #93289, USA) and 200 uL chloroform (Sigma,

#C2432, USA). Samples were incubated for 3 min in an ice block with occasional mixing and

centrifuged for 15min/12000rpm/4°C. The upper aqueous phase was transferred to an RNase-

free microcentrifuge tube containing 500 uL isopropanol & 1 uL glycogen. Samples were

incubated for 10 min in an ice block with occasional mixing, and centrifuged for

10min/14000rpm/4°C. The supernatant was discarded, and the pellet was washed in 500 uL 75%

EtOH followed by centrifugation for 5min/7500rpm/4°C. The pellet was air-dried inside the

laminar flow hood at room temperature (RT) until it became translucent. The RNA pellet was

resuspended in 30 uL of nuclease-free water and incubated at 55°C for 15 min for dissolution.

Any contaminating DNA was eliminated by treating with DNA-free™ DNA Removal Kit

(Thermofisher Scientific, #AM9937, USA) as per the manufacturer's instructions. RNA was

confirmed DNA-free through negative PCR amplification of *rp49 Drosophila* gene. DNA-free

RNA samples were then purified using the ethanol precipitation method by initially raising to

100 uL using nuclease-free water. 10 uL of 3M Sodium Acetate (Thermofisher Scientific, #

R1181, USA) and 300 uL of 100% EtOH were added in the sample, mixed thoroughly &

incubated at -80°C overnight. The next morning, samples were centrifuged at full speed (14000

rpm) for 30 min at 4°C and supernatant was discarded. 500 uL of freshly prepared 70% EtOH

was added and centrifuged at full speed for 20 min at 4°C. The supernatant was discarded and

pellet was air dried for 30 min inside the biosafety cabinet until it became transparent. Pellet was

resuspended in 20 uL nuclease-free water and incubate at 55°C for 15 min to fully dissolve. The

purified RNA sample was stored at -80°C until further use. cDNA was synthesized from RNA

(100 ng for larvae and 1 ug for adults) with SuperScript VILO kit (Invitrogen, #11755250) and

diluted 1:10 using Ambion nuclease-free water.

DNA was isolated from the non-aqueous layers from the samples used for RNA isolation using

Qiagen PureGene Core Kit A kit (Qiagen, #158667, USA) following manufacturer's instructions.

However, during the DNA precipitation step we added 500 uL of 100% Isopropanol and 1 uL of

glycogen to enhance the DNA recovery. qPCR was performed in 2 technical replicates using 1

uL of the diluted cDNA or DNA (10ng) and PowerTrack SYBR Green Mastermix (Applied

Biosystems, #A46109) on a QuantStudio 6 Pro Real-Time PCR system using the following

conditions: 50°C 2 min, 95°C 10 min, 40 cycles (15 s at 95°C , and 30 s at 56°C and 60°C for

whole body and abdomen samples, respectively), 95°C 15 s, 60°C 1 min, 95°C 15 s. Gene

expression of *cifA* and *cifB* (Table S3) was normalized to *groEL* and *Wolbachia* density was

measured by $2^{-\Delta Ct}$ method using *groEL* relative to *rp49*. Ct values above 30 were omitted from

the analysis.

Sexing of pupae

380 Kl-5 primers were used to sex individual pupae collected at the late stage[52]. PCR conditions used

381 were 95°C for 2 min and 30 cycles of 95°C for 30 sec, 56°C for 30 sec, and 72°C for 1 min, final

382 extension of 72°C for 5 min.

Transcriptome

Samples were prepared by generating a uniquely indexed library from each of the RNA samples

isolated from male pupae using the Illumina Stranded Total RNA Prep with RiboZero Plus

Microbiome kit (Illumina, USA) and custom *D. melanogaster* probes for the rRNA depletion.

The libraries were pooled into a single pool containing an approximately equimolar amount of

the samples and outsourced to the Genomic Sciences Laboratory, North Carolina State

University, for sequencing using NovaSeq6000 S4 150 PE lane yielding ~25 million pairs of

reads per sample on average. The raw reads were subjected to quality control using Trimgalore

v0.6.10 with a PHRED score of 20 and minimum read length of 25 bp. The cleaned reads were

mapped to *D. melanogaster* reference genome (v.6.12) using STAR v2.7 to generate BAM files.

The read counting was performed using the BAM files with featureCounts v2.0.6. The raw reads

were rarefied using EcolUtils v0.1 with 100 permutations setting the lowest read (16M) as the

threshold. The reads were then processed for downstream differentially expressed genes (DEGs)

using DeSeq2 v1.42.0.

Histone staining and quantification

Less than 8 hours old males from the same population used for hatch rate assay above were

dissected to collect testes in chilled 1x PBS solution and perform histone staining as described

previously[7]. In brief, the testes were fixed for 30 min at RT in 4% formaldehyde prepared in 1x

PBS with subsequent washing with 1x PBST solution (1x PBS + 0.3% Triton X-100) and

blocking in 1% BSA for core histone at RT for an hour. Samples were incubated overnight with

core histone antibody (MilliporeSigma, Cat#MABE71, USA) (1:1,000 dilution in BSA-PBST) at

4°C in rotating shaker. After washing with 1x PBST solution (1x PBS containing 0.1% Triton X-

100), samples were incubated with goat anti-mouse Alexa Fluor 488 secondary antibody

(Thermo Fisher Scientific, Cat#A28175, 1:500 diluted in 1x PBST) for core histone in the

complete dark at RT for 4 hours. After washing in 1x PBST, samples were then mounted in

vectashield DAPI-containing mounting media (Vector Laboratories, Cat#H-1200-10, USA) for

imaging using Zeiss LSM 880 confocal microscope with constant exposure settings across

treatment groups. The signal intensity from late canoe-stage sperm bundles that retained core

histones was quantified as previously described[7], and the intensity from all stages of

spermatogenesis that retained the H4ac signal was quantified and plotted in GraphPad Prism.

Protamine staining and quantification

Protamine deficiency was evaluated using chromomycin 3 (CMA3) stain as described

previously[7]. In brief, mature sperm were extracted from seminal vesicles of less than 8 hours old

males in chilled 1x PBS. Samples were fixed in 3:1 vol/vol methanol:acetic acid for 20 min at

4°C. Subsequently, the slide was air-dried and treated with 0.25 mg/mL of CMA3 in McIlvain's

buffer, pH 7.0, with 10 mM $MgCl_2$ for 20 min. The samples were washed with 1x PBS and

mounted in vectashield DAPI-containing mounting media. Imaging was performed using Zeiss

LSM 880 confocal microscope. Fluorescence quantification was performed by scoring

fluorescent pixels in arbitrary units (A.U.) within individual sperm heads using ImageJ. The flies

were reared at 21°C instead of 25°C for this assay.

Statistical analysis

428   A comparison of data between two groups was performed using Kolmogorov-Smirnov tests and

429   a Mann-Whitney U test. For more than two groups, the Kruskal Wallis test with Benjamini,

430   Krieger & Yekutieli p-value correction was performed. A correlation test was performed using

431   the Pearson correlation coefficient. To calculate tests and plot graphs, GraphPad Prism v10.0.0

432   was used.

Resource availability

Data deposition: The data reported in this paper have been deposited in Sequence Read Archive (SRA; https://www.ncbi.nlm.nih.gov/sra) (accession no. PRJNA1209084).

Acknowledgments

The authors thank the Huck Institutes Genomics Core Facility (RRID:SCR_023645) at The Pennsylvania State University for generating a uniquely indexed library from each of the RNA samples and custom *D. melanogaster* probes for the rRNA depletion. The co-authors also thank the Microscopy Core Facility (RRID:SCR_024457) for use of the Zeiss LSM 880 confocal microscope. The authors also thank Will Ludington's lab, Department of Embryology, Carnegie Institution for Science, Baltimore, MD, USA, for their expert guidance in generating germ-free flies and proofreading of the manuscript. This work was supported by The Pennsylvania State University funds to the lab of SRB.

Author contributions

449   M.I. and S.R.B conceived the project. M.I. and S.R.B designed the research. M.I., R.K, D.H.Z.L,

450   M.W.B, E.L, and A.M. performed the research. M.I., R.K, E.L, and S.R.B analyzed the data.

451   M.I. and S.R.B wrote the manuscript.

452   **Declaration of interests**

453   The authors declare no conflict of interest.

References

1.   McFall-Ngai, M., Hadfield, M. G., Bosch, T. C. G., Carey, H. V., Domazet-Lošo, T.,
Douglas, A. E., Dubilier, N., Eberl, G., Fukami, T., Gilbert, S. F., Hentschel, U., King, N.,
Kjelleberg, S., Knoll, A. H., Kremer, N., Mazmanian, S. K., Metcalf, J. L., Nealson, K.,
Pierce, N. E., Rawls, J. F., Reid, A., Ruby, E. G., Rumpho, M., Sanders, J. G., Tautz, D. &
Wernegreen, J. J. Animals in a bacterial world, a new imperative for the life sciences.
Proceedings of the National Academy of Sciences **110,** 3229–3236 (2013).
2.   Bordenstein, S. R. & Theis, K. R. Host Biology in Light of the Microbiome: Ten
Principles of Holobionts and Hologenomes. *PLoS Biol* **13,** e1002226 (2015).
3.   Bordenstein, S. R., The Holobiont Biology Network, Gilbert, M. T. P., Ginnan, N.,
Malacrinò, A., Martino, M. E., Bahrndorff, S., Mundra, S., Martin, M. D., Theis, K. R.,
Hird, S. M., Caro-Quintero, A., Sharpton, T. J., Kohl, K. D., Barnes, C. J., Eisenhofer, R.,
Aizpurua, O., Andersen, S. B., Brealey, J. C., Noer, C. L., Medina, M., Limborg, M. T. &
Alberdi, A. The disciplinary matrix of holobiont biology. *Science (1979)* **386,** 731–732
(2024).
4.   Brinker, P., Fontaine, M. C., Beukeboom, L. W. & Falcao Salles, J. Host, Symbionts, and
the Microbiome: The Missing Tripartite Interaction. *Trends Microbiol* **27,** 480–488
(2019).
5.   Masson, F. & Lemaitre, B. Growing Ungrowable Bacteria: Overview and Perspectives on
Insect Symbiont Culturability. *Microbiol Mol Biol Rev* **84,** (2020).
6.   Weinert, L. A., Araujo-Jnr, E. V., Ahmed, M. Z. & Welch, J. J. The incidence of bacterial
endosymbionts in terrestrial arthropods. *Proceedings of the Royal Society B: Biological
Sciences* **282,** 20150249 (2015).
7.   Kaur, R., Leigh, B. A., Ritchie, I. T. & Bordenstein, S. R. The Cif proteins from
Wolbachia prophage WO modify sperm genome integrity to establish cytoplasmic
incompatibility. *PLoS Biol* **20,** e3001584 (2022).
8.   Shropshire, J. D., Leigh, B. & Bordenstein, S. R. Symbiont-mediated cytoplasmic
incompatibility: What have we learned in 50 years? *Elife* **9,** (2020).
9.   Hoffmann, A. A., Turelli, M. & Harshman, L. G. Factors affecting the distribution of
cytoplasmic incompatibility in Drosophila simulans. *Genetics* **126,** 933–948 (1990).
10.   Shropshire, J. D., Hamant, E., Conner, W. R. & Cooper, B. S. *cifB-* transcript levels
largely explain cytoplasmic incompatibility variation across divergent *Wolbachia. PNAS
Nexus* **1,** (2022).
11.   Landmann, F., Orsi, G. A., Loppin, B. & Sullivan, W. Wolbachia-Mediated Cytoplasmic
Incompatibility Is Associated with Impaired Histone Deposition in the Male Pronucleus.
PLoS Pathog **5,** e1000343 (2009).
12.   Ludington, W. B. & Ja, W. W. Drosophila as a model for the gut microbiome. *PLoS
Pathog* **16,** e1008398 (2020).
13.   Simhadri, R. K., Fast, E. M., Guo, R., Schultz, M. J., Vaisman, N., Ortiz, L., Bybee, J.,
Slatko, B. E. & Frydman, H. M. The Gut Commensal Microbiome of Drosophila
melanogaster Is Modified by the Endosymbiont *Wolbachia. mSphere* **2,** (2017).
14.   Ye, Y. H., Seleznev, A., Flores, H. A., Woolfit, M. & McGraw, E. A. Gut microbiota in
Drosophila melanogaster interacts with Wolbachia but does not contribute to Wolbachia -
mediated antiviral protection. *J Invertebr Pathol* **143,** 18–25 (2017).

15.   Shropshire, J. D., Hamant, E. & Cooper, B. S. Male Age and *Wolbachia* Dynamics:
Investigating How Fast and Why Bacterial Densities and Cytoplasmic Incompatibility
Strengths Vary. *mBio* **12,** (2021).

16.   Kaur, R., Meier, C. J., McGraw, E. A., Hillyer, J. F. & Bordenstein, S. R. The mechanism
of cytoplasmic incompatibility is conserved in Wolbachia-infected Aedes aegypti
mosquitoes deployed for arbovirus control. *PLoS Biol* **22,** e3002573 (2024).

17.   Ingham, P. The development of Drosophila melanogaster. *Trends in Genetics* **10,** 299
(1994).

18.   Williamson, A. & Lehmann, R. GERM CELL DEVELOPMENT IN *DROSOPHILA*.
Annu Rev Cell Dev Biol **12,** 365–391 (1996).

19.   Kornberg, R. D. & Lorch, Y. Twenty-five years of the nucleosome, fundamental particle
of the eukaryote chromosome. *Cell* **98,** Preprint at https://doi.org/10.1016/S0092-
8674(00)81958-3 (1999)

20.   Rathke, C., Baarends, W. M., Jayaramaiah-Raja, S., Bartkuhn, M., Renkawitz, R. &
Renkawitz-Pohl, R. Transition from a nucleosome-based to a protamine-based chromatin
configuration during spermiogenesis in *Drosophila*. *J Cell Sci* **120,** 1689–1700 (2007).

21.   Akhmanova, A. S., Bindels, P. C. T., Xu, J., Miedema, K., Kremer, H., Hennig, W., Xu, J.
& Hennig, W. Structure and expression of histone H3.3 genes in *Drosophila melanogaster*
and *Drosophila hydei*. *Genome* **38,** 586–600 (1995).

22.   Sakaguchi, A., Karachentsev, D., Seth-Pasricha, M., Druzhinina, M. & Steward, R.
Functional characterization of the Drosophila Hmt4-20/Suv4-20 histone
methyltransferase. *Genetics* **179,** 317–22 (2008).

23.   Kaur, R., McGarry, A., Shropshire, J. D., Leigh, B. A. & Bordenstein, S. R. Prophage
proteins alter long noncoding RNA and DNA of developing sperm to induce a paternal-
effect lethality. *Science (1979)* **383,** 1111–1117 (2024).

24.   Jayaramaiah Raja, S. & Renkawitz-Pohl, R. Replacement by *Drosophila melanogaster*
Protamines and Mst77F of Histones during Chromatin Condensation in Late Spermatids
and Role of Sesame in the Removal of These Proteins from the Male Pronucleus. *Mol Cell*
Biol **25,** 6165–6177 (2005).

25.   Eren-Ghiani, Z., Rathke, C., Theofel, I. & Renkawitz-Pohl, R. Prtl99C Acts Together with
Protamines and Safeguards Male Fertility in Drosophila. *Cell Rep* **13,** 2327–2335 (2015).

26.   Scholl, A., Ndoja, I., Dhakal, N., Morante, D., Ivan, A., Newman, D., Mossington, T.,
Clemans, C., Surapaneni, S., Powers, M. & Jiang, L. The Osiris family genes function as
novel regulators of the tube maturation process in the Drosophila trachea. *PLoS Genet* **19,**
e1010571 (2023).

27.   Ikeda, T., Ishikawa, H. & Sasaki, T. Infection density of Wolbachia and level of
cytoplasmic incompatibility in the Mediterranean flour moth, Ephestia kuehniella. *J*
Invertebr Pathol **84,** 1–5 (2003).

28.   Ritchie, I. T., Needles, K. T., Leigh, B. A., Kaur, R. & Bordenstein, S. R. Transgenic
cytoplasmic incompatibility persists across age and temperature variation in Drosophila
melanogaster. *iScience* **25,** 105327 (2022).

29.   Bordenstein, S. R., Uy, J. J. & Werren, J. H. Host Genotype Determines Cytoplasmic
Incompatibility Type in the Haplodiploid Genus Nasonia. *Genetics* **164,** 223–233 (2003).

30.   Aranda-Díaz, A., Obadia, B., Dodge, R., Thomsen, T., Hallberg, Z. F., Güvener, Z. T.,
Ludington, W. B. & Huang, K. C. Bacterial interspecies interactions modulate pH-
mediated antibiotic tolerance. *Elife* **9,** (2020).

31.   Newell, P. D. & Douglas, A. E. Interspecies Interactions Determine the Impact of the Gut
547   Microbiota on Nutrient Allocation in Drosophila melanogaster. *Appl Environ Microbiol*
548   **80,** 788–796 (2014).
549   32.   Sommer, A. J. & Newell, P. D. Metabolic Basis for Mutualism between Gut Bacteria and
550   Its Impact on the *Drosophila melanogaster* Host. *Appl Environ Microbiol* **85,** (2019).
551   33.   Chaston, J. M., Dobson, A. J., Newell, P. D. & Douglas, A. E. Host genetic control of the
552   microbiota mediates the Drosophila nutritional phenotype. *Appl Environ Microbiol* **82,**
553   (2016).
554   34.   Layton, E. M., On, J., Perlmutter, J. I., Bordenstein, S. R. & Shropshire, J. D. Paternal
555   Grandmother Age Affects the Strength of *Wolbachia* -Induced Cytoplasmic
556   Incompatibility in Drosophila melanogaster. *mBio* **10,** (2019).
557   35.   LePage, D. P., Metcalf, J. A., Bordenstein, S. R., On, J., Perlmutter, J. I., Shropshire, J. D.,
558   Layton, E. M., Funkhouser-Jones, L. J., Beckmann, J. F. & Bordenstein, S. R. Prophage
559   WO genes recapitulate and enhance Wolbachia-induced cytoplasmic incompatibility.
560   *Nature* **543,** 243–247 (2017).
561   36.   Bourtzis, K., Nirgianaki, A., Markakis, G. & Savakis, C. Wolbachia Infection and
562   Cytoplasmic Incompatibility in Drosophila Species. *Genetics* **144,** 1063–1073 (1996).
563   37.   Cooper, B. S., Ginsberg, P. S., Turelli, M. & Matute, D. R. Wolbachia in the *Drosophila*
564   *yakuba* Complex: Pervasive Frequency Variation and Weak Cytoplasmic Incompatibility,
565   but No Apparent Effect on Reproductive Isolation. *Genetics* **205,** 333–351 (2017).
566   38.   Shropshire, J. D., On, J., Layton, E. M., Zhou, H. & Bordenstein, S. R. One prophage WO
567   gene rescues cytoplasmic incompatibility in *Drosophila melanogaster*. *Proceedings of the*
568   *National Academy of Sciences* **115,** 4987–4991 (2018).
569   39.   Reynolds, K. T. & Hoffmann, A. A. Male age, host effects and the weak expression or
570   non-expression of cytoplasmic incompatibility in Drosophila strains infected by
571   maternally transmitted Wolbachia. *Genet Res* **80,** (2002).
572   40.   Yamada, R., Floate, K. D., Riegler, M. & O'Neill, S. L. Male Development Time
573   Influences the Strength of Wolbachia-Induced Cytoplasmic Incompatibility Expression in
574   *Drosophila melanogaster*. *Genetics* **177,** 801–808 (2007).
575   41.   Numajiri, Y., Kondo, N. I., Toquenaga, Y. & Kageyama, D. Intraspecies variation in
576   cytoplasmic incompatibility intensity in the bean beetle Callosobruchus analis. *Evol Ecol*
577   **38,** 861–870 (2024).
578   42.   Sasaki, T., Kubo, T. & Ishikawa, H. Interspecific Transfer of Wolbachia Between Two
579   Lepidopteran Insects Expressing Cytoplasmic Incompatibility: A Wolbachia Variant
580   Naturally Infecting *Cadra cautella* Causes Male Killing in *Ephestia kuehniella*. *Genetics*
581   **162,** 1313–1319 (2002).
582   43.   Suwanchaisri, K., Roddee, J. & Wangkeeree, J. Wolbachia Transinfection and Effect on
583   the Biological Traits of Matsumuratettix hiroglyphicus (Matsumura), the Leafhopper
584   Vector of Sugarcane White Leaf Disease. *Agriculture* **14,** 1236 (2024).
585   44.   Hughes, G. L., Dodson, B. L., Johnson, R. M., Murdock, C. C., Tsujimoto, H., Suzuki, Y.,
586   Patt, A. A., Cui, L., Nossa, C. W., Barry, R. M., Sakamoto, J. M., Hornett, E. A. &
587   Rasgon, J. L. Native microbiome impedes vertical transmission of *Wolbachia* in
588   *Anopheles* mosquitoes. *Proceedings of the National Academy of Sciences* **111,** 12498–
589   12503 (2014).

45.   Detcharoen, M., Jiggins, F. M., Schlick-Steiner, B. C. & Steiner, F. M. Wolbachia
endosymbiotic bacteria alter the gut microbiome in the fly Drosophila nigrosparsa. *J*
Invertebr Pathol **198,** 107915 (2023).
46.   Walker, T., Johnson, P. H., Moreira, L. A., Iturbe-Ormaetxe, I., Frentiu, F. D.,
McMeniman, C. J., Leong, Y. S., Dong, Y., Axford, J., Kriesner, P., Lloyd, A. L., Ritchie,
S. A., O'Neill, S. L. & Hoffmann, A. A. The wMel Wolbachia strain blocks dengue and
invades caged Aedes aegypti populations. *Nature* **476,** (2011).
47.   Blagrove, M. S. C., Arias-Goeta, C., Failloux, A. B. & Sinkins, S. P. Wolbachia strain
wMel induces cytoplasmic incompatibility and blocks dengue transmission in Aedes
albopictus. *Proc Natl Acad Sci U S A* **109,** (2012).
48.   O'Neill, S. L. in *Adv Exp Med Biol* **1062,** (2018).
49.   Suito, T., Nagao, K., Juni, N., Hara, Y., Sokabe, T., Atomi, H. & Umeda, M. Regulation
of thermoregulatory behavior by commensal bacteria in *Drosophila*. *Biosci Biotechnol*
Biochem **86,** 1060–1070 (2022).
50.   Koyle, M. L., Veloz, M., Judd, A. M., Wong, A. C.-N., Newell, P. D., Douglas, A. E. &
Chaston, J. M. Rearing the Fruit Fly Drosophila melanogaster
Under Axenic and Gnotobiotic Conditions. *Journal of Visualized Experiments* (2016).
doi:10.3791/54219
51.   Chandler, J. A., Innocent, L. V., Martinez, D. J., Huang, I. L., Yang, J. L., Eisen, M. B. &
Ludington, W. B. Microbiome-by-ethanol interactions impact Drosophila melanogaster
fitness, physiology, and behavior. *iScience* **25,** 104000 (2022).
52.   Lott, S. E., Villalta, J. E., Schroth, G. P., Luo, S., Tonkin, L. A. & Eisen, M. B.
Noncanonical Compensation of Zygotic X Transcription in Early Drosophila
melanogaster Development Revealed through Single-Embryo RNA-Seq. *PLoS Biol* **9,**
e1000590 (2011).
615
616
617

Fig 1. CI penetrance is stronger in germ-free males and links with host regulation of

histone, protamine, and male-specific transcripts. (A) Schematic representation of the fly

lines showing (from left to right): complete symbiont community, gut microbiome only,

endosymbiont (*Wolbachia*) only, and symbiont-free. (B) Germ-free males with *Wolbachia* (GF+)

induce significantly higher incompatibility (dark pink) than conventional (CV+) males. CI is a

metric for embryonic viability that is measured by the percentage of embryos that develop into

larvae. Compatible crosses (grey bars) with normal embryonic hatching percentages are controls

that express either no CI or rescue of CI. P-values were calculated based on a Kruskal-Wallis test

followed by Benjamini, Krieger, and Yekutieli multiple correction test. (C) GF+ males with a

native microbiome reintroduced (GF+$_M$) recapitulate the CI strength similar to CV+ males. (D)

Comparative transcriptomic analysis of GF+ vs. CV+ male pupae and GF- vs. CV- male pupae

revealed significant differentially expressed genes narrowed here to 37 genes expressing histone,

protamine, and male-specific transcripts (MSTs). Significant differences are highlighted with a

black colored asterisk (*). Notably, the histone-related genes genes (red) were upregulated, while

protamine (green) and MST (blue) genes were downregulated. Log2 fold change was calculated

using DESeq2, and its default parameters; Walt test and Benjamini-Hochberg procedure were

used to calculate p and adjusted p-values. The asterisk *, **, and *** represent adjusted p-values

less than 0.05, 0.005, and 0.0005, respectively.

Fig 2. CI defects in the histone-to-protamine transition occur in germ-free and

conventional males. (A) During *Drosophila melanogaster* sperm development, chromatin is

initially packaged with histones. In the late canoe stage, histones are replaced by protamines and

protamine-like proteins, facilitating tight chromatin condensation within the sperm head. We

tested if there is variation in the histone-protamine transition using testes (N = ~10) from <8-
hour-old virgin males of GF-, GF+, CV-, and CV+ lines were dissected and (B) immunostained
to visualize and quantify histone abundance (magenta) in late-canoe elongating spermatids.
DAPI stain (grey) was used to label spermatid nuclei, and (C) CMA3 stained for protamine
deficiency. Signal intensity from each stained spermatid with signal was quantified in ImageJ
and graphed. (B) GF+ and CV+ late canoe spermatids both significantly retain histones
compared to their negative counterparts GF- and CV-, respectively. (C) GF+ and CV+ mature
sperm have enhanced CMA3 intensity, consistent with protamine deficiency levels compared to
GF- and CV- negative controls, respectively. Vertical bars represent the median, and error bars
represent the standard deviation. P-values are calculated using the Mann-Whitney U test in
Prism.

Fig 3. *Wolbachia* density and transcriptional activity vary with host developmental stage
and microbiome status. (A) Relative to CV+ males, *Wolbachia* densities were significantly
lower in the GF+ adult males and (B) similar in GF+ larvae using sigle copy genes of the
endosymbiont (*groEL*) and host (*rp49*). (D-F) *cifA* and *cifB* gene expression in adults and larvae
were similar between CV+ and GF+ males, suggesting that the neither *Wolbachia* density nor *cif*
gene expression underpin exacerbated CI in GF+ flies. The p-values are shown for the non-
parametric Kolmogorov-Smirnov [and Mann-Whitney U] tests. (G) *Wolbachia* transcriptomes
across development reveal that *Wolbachia* are transcriptionally more active in GF+ compared to
CV+, particularly in the larval stage. Among the differentially-expressed genes, the most
common types of transcripts were hypothetical and microbial interactions, such as transcripts
with ankyrin repeat domains.

Fig 4. Holobionts with *Wolbachia* have higher colony forming units (CFUs) of gut bacteria. (A) Homogenates of less than one day old CV- (white circles) and CV+ males (orange circles) (N=5) in triplicates were spread on MRS and MYPL plates. (B) *Wolbachia* infection (CV+) significantly increased the CFU load of *Acetobacter spp.* and *Lactobacillus spp.*, suggesting a positive interaction between the endosymbiont and the microbiome. The p-values are calculated with a Mann-Whitney U test. (C) The CFU of *Acetobacter* spp. and *Lactobacillus* spp. in CV- (data points filled in white) and CV+ (data points with filled in orange color) males have a significant positive correlation. All flies were surface sterilized to avoid transient bacteria residing on the fly surface. A CFU count below 500 is typical in most flies, as young flies (<1 day old) have not yet developed a fully established gut microbiome.

Fig 5. The fitness impacts of *Wolbachia* and gut microbiome are independent and developmentally staged, and microbiome reintroduction into germ-free flies restores fly fitness. (A) The extracellular microbiome and *Wolbachia* exert independent and positive effects on embryo-to-larval survival, as calcuated by the number of surving larvae divided by the number of emryos laid for CV+ vs. CV- and GF+ vs. GF-. (B) The extracellular microbiome is beneficial to larval-to-pupal survival, while *Wolbachia* imposes a fitness cost in a microbiome-independent manner. (D) Similarly, *Wolbachia* lower pupal-to-adult survival percentage in a microbiome-independent manner. (D) The overall trend from embry-to-adult shows GF+ flies consistently performing weker comapred to CV+. The top green bar represents the positive fitness effect of the microbiome on the host, while the green and red lines at the bottom indicate the positive and negative effects, respectively, of *Wolbachia* on the host. (E) When the

conventional microbiome was re-introduced into GF+ flies through fecal-to-oral transmission,

the egg-to-adult survival of the recipient flies (GF$_M$+) significantly increased to the same level as

that of CV+ flies. Each circle represents a family, and color denotes genotype status: orange bar

(CV), blue bar (GF), open circle (no *Wolbachia*), and filled cirlce (*Wolbachia*). The p-values

were calculated with Kruskal-Wallis test followed by Benjamini, Krieger, and Yekutieli multiple

correction test.

693
694

 SUPPORTING INFORMATION

S1 Fig. Confirmation of gut microbiome absence and presence by colony forming units

(CFUs) in GF and CV lines. The germ-free status of the GF-/+ lines was confirmed by the

absence of colonies when 20uL of fly homogenate were plated on to MYPL and MRS plates.

The conventionally-reared positive control lines had ample colonies with colony number

significantly elevated in CV+ relative to CV- flies.

S2 Fig. The top misregulated transcripts link with reproductive, chitin, cuticle, and sensory

development. The diffrentially-expressed transcripts that were significant (adjusted p<0.05) are

shown for (A) GF+ vs CV+ male pupae and (B) GF- vs CV- male puape.

S3 Fig. Copy number of host *rp49* in CV+ and GF+ are comparable. (A) Standard curves for

the absolute quantification of *rp49* were generated using a log10 dilution series of known

amounts of PCR product amplified from a larger region of the same *rp49* gene (404 bp). PCR

amplification was performed using primers listed in Table S1 at a concentration of 0.4 µM. The

cycling conditions were as follows: 30 cycles of 95°C for 15 seconds, 60°C for 30 seconds, and

72°C for 30 seconds. The PCR products were purified using the Monarch DNA Purification Kit

(NEB) and their concentrations were measured using a NanoDrop One spectrophotometer. The

number of *rp49* copies was calculated using the following formula: Number of DNA copies =

(amount of DNA in ng * 6.022 x 10^23) / length of amplicon in bp x 10^9 x 660). Standard

curve was plotting using Design & Analysis Software v2.8.0 (B) Mann Whitney U test of the

CV+ and GF+ shows that the absolute count of CV+ and GF+ *rp49* gene in their abdomen are

similar.

S4 Fig. Dechorination of *Drosophila melanogaster* embryos. The germ-free treatment

followed the standard procedure of bleaching fertilized embryos. Briefly, CV- and CV+ females

were allowed to oviposit overnight on a grape plate smeared with yeast. The next morning, the

grape plates were replaced with fresh ones smeared with yeast, and the flies were allowed to

oviposit for 30 minutes. This was followed by another replacement with a fresh grape plate

smeared with yeast, where the flies oviposited for an additional hour to collect 1-hour embryos.

The collected embryos were transferred to a petri dish and approximately half was transferred

into a cell strainer. These embryos were bleached twice in 10% bleach for 2.5 minutes each,

followed by a single wash in 70% ethanol for 30 seconds and three quick washes in PBS

containing 0.1% Triton X for 10 seconds each. The dechorionated eggs were carefully counted

under a stereomicroscope and then placed on a germ-free grape plate or a germ-free food vial

before being incubated under sterile conditions.

S1 Table: Genetic comparison between CV- and CV+ lines

S2 Table: Differentially-expressed histone, protamine, and MST transcripts in male pupae.

S3 Table: Primers used for measuring *Wolbachia* density, sex of pupae, and *cifA* and *cifB*

expression.

A

CV+   CV−   GF+   GF−

🪱 Gut Microbiome   🦠 Reproductive Wolbachia

B

Embryos hatched (%)

<0.0001
<0.0001
0.0371
<0.0001

CV+ x CV− GF+ x CV− GF− x CV− CV+ x CV+ GF+ x CV+

C

Embryos hatched (%)

0.2530
<0.0001
<0.0001

CV+ x CV− GF+ x CV− GF+ₘ x CV−

Cross (male x female)
- 🟣 Incompatible cross
- ⚪ Compatible cross

D

Sperm associated transcripts

	GF+/CV+	GF−/CV−
Mst57Da (MSTs)	***	***
Mst57Db (MSTs)	***	
Mst57Dc (MSTs)	***	***
Mst84Da (MSTs)		
Mst87F (MSTs)	*	***
Mst84Db (MSTs)	*	***
Mst84Dc (MSTs)	*	***
Mst98Cb (MSTs)	**	***
Mst77Y−7 (MSTs)		*
Mst98Ca (MSTs)	*	**
Mst84Dd (MSTs)	*	*
Mst77Y−13 (MSTs)	*	
Mst89B (MSTs)	*	*
Mst36Fb (MSTs)	*	
Mst33A (MSTs)		
MstProx (MSTs)		
Mst36Fa (MSTs)		
Mst84B (MSTs)	*	
Mst77Y−16Psi (MSTs)		
Mst27D (MSTs)		
mst (MSTs)		
Mst77Y−8Psi (MSTs)		
Mst85C (MSTs)	*	**
Mst77Y−4 (MSTs)		
Prtl99C (Prot/PL)	*	*
ProtB (Prot/PL)	*	
Mst77F (Prot/PL)	*	*
ProtA (Prot/PL)		
His3.3A (Histones)		
Utx (Histones)		
Hinfp (Histones)		
BigH1 (Histones)		
His2B:CG17949 (Histones)		
His4r (Histones)		
Hmt4−20 (Histones)	*	*
His3.3B (Histones)	*	
His2Av (Histones)		
Hers (Histones)		***

Log2 Fold Change

−12 −10 −8 −6 −4 −2 0 2

A Sperm developmental stages

Germline stem cells

Spermatogonia

Spermatocytes

Round onion spermatids

Histone
Protamine
transition

Canoe spermatids

Needle Spermatids

Matured sperm

B

| DAPI | α-His | Merged |

CV+

GF+

Elongating spermatid histone

0.0011

% of spermatid bundles with histone retention

CV-   CV+

0.0334

% of spermatid bundles with histone retention

GF-   GF+

C

CMA3

CV-

CV+

GF-

GF+

0.0029

Protamine-deficient CMA3 fluorescence per sperm

CV-   CV+

<0.0001

Protamine-deficient CMA3 fluorescence per sperm

GF-   GF+

Microbiome — Increases host fitness

A

Embryo to larval survival (%)

<0.0001
<0.0001
0.0071   0.0201

CV- CV+ GF- GF+
Genotype

B

Larval to pupal survival (%)

<0.0001
0.0065
0.0324   0.0002

CV- CV+ GF- GF+
Genotype

C

Pupal to adult survival (%)

0.0758
0.0878
0.0027   0.0027

CV- CV+ GF- GF+
Genotype

D

Survival %

CV+
GF+

Eggs Larvae Pupae Adult
Developmental stages

E

Egg-to-adult survival (%)

0.1513
0.0402
0.0004

CV+ GF+ GF$_M$+
Genotype

Wolbachia — Increases host fitness

Reduces host fitness