Title: *FishPi*: a bioinformatic prediction tool to link piRNA and transposable elements

Alice M. Godden[1], Benjamin Rix[1], Simone Immler[1]

[1] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,

NR4 7TJ, United Kingdom

Corresponding author:

Alice M. Godden Email: alice.godden@uea.ac.uk

Abstract

Background

Piwi-interacting RNAs (piRNA)s are non-coding small RNAs that post-transcriptionally

affect gene expression and regulation. Through complementary seed region binding with

transposable elements (TEs), piRNAs protect the genome from transposition. A tool to

link piRNAs with complementary TE targets will improve our understanding of the role of

piRNAs in genome maintenance and gene regulation. Existing tools such as *TEsmall*

can process sRNA-seq datasets to produce differentially expressed piRNAs, and

piRScan developed for nematodes can link piRNAs and TEs but it requires knowledge

about the target region of interest and works backwards.

Results

We have therefore developed *FishPi* to predict the pairings between piRNA and TEs for

available genomes from zebrafish, medaka and tilapia, with full user customisation of

parameters including orientation of piRNA, mismatches in the piRNA seed binding to TE

and scored output lists of piRNA-TE matches. *FishPi* works with individual piRNAs or a

list of piRNA sequences in fasta format. The software focuses on the piRNA-TE seed

region and analyses reference TEs for piRNA complementarity. TE type is examined,

counted and stored to a dictionary, with genomic loci recorded. Any updates to piRNA-

TE binding rules can easily be incorporated by changing the seed-region options in the

graphic user-interface. *FishPi* provides a graphic interface using *tkinter* for the user to

input piRNA sequences to generate comprehensive reports on piRNA-TE interactions.

FishPi can easily be adapted to genomes from other species and taxa opening the

interpretation of piRNA functionality to a wide community.

Conclusions

Users will gain insight into genome mobility and *FishPi* will help further our

understanding of the biological role of piRNAs and their interaction with TEs in a similar

way that public databases have improved the access to and the understanding of the

role of small RNAs.

Keywords

PiRNAs, transposons, evolution, zebrafish

Background

Environmental changes and related biotic and abiotic stressors can disrupt the balance and

expression of small RNAs and transposable elements (TEs) in the genome [1]. TEs are mobile

DNA elements that can cause insertions in the genome, as well as deletions and genomic

rearrangements, initially identified by McClintock in maize but later found to be present

across taxa [2,3]. The mechanisms underlying TE transposition in zebrafish can be divided into

two categories: retrotransposons that synthesise an RNA intermediate to insert themselves

into the host genome [4] and DNA transposons that use a cut-and-paste mechanism to insert

themselves into the host genome [4-6]. Therefore, knowing the family of TE that is mobile can

improve our understanding of the mode of TE action and impact on the genome.

Piwi-interacting RNAs (piRNAs) have evolved to repress the action of TEs and constitute a

substantial fraction of small RNAs [7]. The main function of piRNAs is to silence TEs and

repetitive elements in animal germ cells to preserve the integrity of the germline genome [8].

In zebrafish *Danio rerio*, piRNAs are 100x more frequent than miRNAs in zebrafish ovaries

and testes [7], and small RNAs post-transcriptionally regulate gene expression and regulation

[9] and the loss of Ziwi leads to apoptosis of germ cells in embryonic development [7]. Ziwi and

Zili proteins provide a piRNA amplification system that is highly conserved. This is important

as the availability of complementary piRNAs is essential for survival of germ cells [7]. In the

nematode *Caenorhabditis elegans,* loss of TE silencing by piRNAs can lead to sterility [10].

TEs are not randomly distributed in the genome but they have preferred regions for insertion,

including distorted and bent DNA [5,11], or open chromatin regions [11,12]. Vast populations of

piRNAs are complementary to TE messenger RNAs (mRNA) [13]. In *C. elegans*, the 2nd to the

7th nucleotide at the 5' end of the piRNA [14] or the 2nd to the 8th nucleotide [15] have both been

suggested to be the piRNA-TE seed region. Similarly, in the house mouse *Mus musculus*,

the 2nd to 8th nucleotides have been identified as a seed region [16]. In contrast to mammals,

most zebrafish piRNAs map to TEs, and the LTR families in particular [7]. In zebrafish and

other teleost species, the first ten nucleotides at the 5' end of the piRNA are loaded into the

Argonaute machinery to produce Ziwi (zebrafish Piwi)-bound piRNAs complementary to TE

transcripts [7,13]. The same is true for many other organisms including *Drosophila*, as piRNAs

show complementarity to other antisense piRNAs in their first ten nucleotides, which

associate with PIWI proteins to generate the mature 5' end of the piRNA [7] forming part of the

ping-pong amplification cycle where piRNAs act and bind complementarily to transposons [17].

It is the PIWI-bound piRNA with the piRNA seed that recognises a complementary region on

the transposon and leads to silencing [18,8]. In *Drosophila*, piRNAs cleave their target mRNAs

(TEs) after the 10th nucleotide from 5' end of the piRNA[19]. Given the evolutionary

conservation of piRNA biogenesis and TE silencing mechanisms [7,20], using the first ten

nucleotides of piRNA as a putative piRNA seed region is what we use as a starting point.

However, *FishPi* allows users to propose custom piRNA seed regions to predict piRNA:TE

activity in the event of further work confirming or changing this prediction.

TEs are known to be activated through specific environmental stressors. In *C. elegans*

mutant strains lacking the ability to generate piRNAs, heat-treated worms displayed

transgenerational impairments for up to three subsequent generations with thermal stress

reducing the rate of piRNA biogenesis [21,22]. In *Drosophila,* the *mariner-Mos1* transposon

was enriched following heat-stress but not following ultraviolet irradiation [23]. TEs are

expressed in a tissue-specific manner and can affect transcription and gene expression [24] by

generating new genes and mRNAs through genomic rearrangements and *de novo*

mutations, and cause alteration of regulatory networks and even trigger immune responses

[5]. Understanding the interaction between piRNAs and their targets is therefore key to

understand the defence mechanisms in the germ line against the effects of environmental

conditions.

We developed the software *FishPi* to enable mapping zebrafish piRNAs against

complementary TEs and link them to understand their biological association [13]. *FishPi*

facilitates research into the function of piRNAs and accelerates analyses in the same way as

microRNA databases like *Targetscan* have supported small RNA research [25]. *FishPi*

provides a user-friendly graphic interface (Figure 1) and generates a comprehensive output

that can be exported as a report including complementary TE counts, family level annotation,

chromosomal and counts plots, with the option to export the full list of complementary TEs

and their sequences. The user can enter individual piRNAs or lists of piRNAs in fasta format.

For zebrafish and other model organisms, a list of piRNA sequences are available at

piRBase   (http://bigdata.ibp.ac.cn/piRBase/browse.php),   [26]   or   piRNAclusterDB

(https://www.smallrnagroup.uni-mainz.de/piRNAclusterDB/) [27]. *FishPi* can easily be used for

any species for which the relevant information such as a high-resolution reference genome

and sufficient piRNA sequences are available.

Implementation

FishPi is a Python-based programme and uses the following packages and versions: Python

v3.11, Tkinter (included in Python v3.11 [28]), Pillow 10.0.1 [29], and Matplotlib 3.8.0 [30]. The

code is platform independent (see Figure 2 for overview of processes) and follows the

installation guidance for Conda on Github (https://github.com/alicegodden/fishpi). The TE

families assigned to each class were checked using RepeatMasker and Dfam (Table 1) [31].

Input TE fasta files can be found on Zenodo: https://zenodo.org/records/13975588.

Input file preparation

To generate a fasta file of the TE sequences from the zebrafish reference genome GRCz11,

a TE.bed file with clade was created by using the UCSC table genome browser [32] as follows:

Vertebrate, group: Variation and Repeats, genome: zebrafish, assembly: May 2017

GRCz11, track: RepeatMasker, table: rmsk. Reference genome files for zebrafish, medaka

and tilapia were obtained from Ensembl. Genomes were decompressed with *gunzip* and

indexed with *samtools faidx refgenome.fa*. A gtf TE annotation file was generated with: *perl*

makeTEgtf.pl -c 6 -s 7 -e 8 -o 10 -t 11 fish_rmsk > fish_TE.gtf. This script was obtained from

the   Hammel   lab   (https://labshare.cshl.edu/shares/mhammelllab/www-

[data/TEtranscripts/TE_GTF/makeTEgtf.pl.gz](https://labshare.cshl.edu/shares/mhammelllab/www-data/TEtranscripts/TE_GTF/makeTEgtf.pl.gz)) [33]. Only necessary columns were retained

with following command in fish_TE.gtf: *cut -f1,2,3,4,5,7,8,9,10,11,12,13,14,15,16 fish_TE.gtf*

> fish_TE.use.gtf. The awk script to convert .gtf to .bed file can be found on the

README.md https://github.com/alicegodden/fishpi/blob/main/README.md , along with *sed*

commands to handle different chromosome naming conventions. Finally, the steps to extract

DNA sequences from the reference genome are: *bedtools getfasta -s -name -fi*

Danio_rerio.GRCz11.dna.primary_assembly.fa   -fo   GRCz11.teseqs.use.fasta   -bed

GRCz11.teannotation.bed.

To generate chromosome length files for chromosomal plots "species_chrom_end.txt" files,

the user needs to navigate to the UCSC table browser (https://genome.ucsc.edu/cgi-

bin/hgTables) and select the species and reference genomes as above. Under the "Group"

drop down menu, select "*All Tables*". Under the "*Table*" drop down menu select

"cytoBandIdeo". To download the file in .txt format, the first column needs to be chromosome

names/numbers and second column to be the length of each chromosome. The function that

represents the inclusion of custom species files is *select_custom_files().*

Users can choose their own piRNA sequence(s) through individual pasting of piRNA

sequences or upload a list of named piRNAs, *upload_pirna_file.* The piRNA seed region

sequence can be specified. For teleosts*,* we recommend 1-10 nucleotides (default setting).

"*Reverse complement*" can be checked to add the reverse of piRNA seed for

complementarity matching. Mismatches in the piRNA seed can be included, the number of

these can be specified, where zero is the default setting. Once executed with

analyse_sequence() or *analyse_uploaded_pirna(),* complementary TEs are identified in any

orientation with *is_complementary()* and with mismatches with *introduce_mismatches().*

Visualisations are provided by the function *create_results()*. The resulting bar chart includes

a binomial distribution test, with stars on the bar chart denoting $P \leq 0.05$. This is to show that

the likelihood of a specific TE family is complementary to the piRNA at a significantly higher

than expected ratio. The functions *classify_te_type()* group complementary TEs by family for

plotting and *plot_te_chromosomal_locations()* displays a chromosomal plot of

complementary TE loci in the genome. All results can be exported with

export_complementary_te_list(), a .csv output file of results including piRNA names (when

fasta file of piRNAs is uploaded), TE name and location, orientation of complementarity and

complementarity percentage of piRNA (not including seed region) to TE. The functions

export_bar_chart() and *export_chrom_plot()* allow users to save high resolution images for

presentation and publication.

Results & Discussion

FishPi is designed for users with minimal coding experience. A detailed manual can be

found in the README on the git repository:

https://github.com/alicegodden/fishpi/blob/main/README.md. The simple graphic user

interface offers a point and click tool to rapidly gain insight into many piRNA sequences.

Each time a new piRNA sequence is provided, the programme resets itself before providing

the new results. As more research into piRNA-TE seed sites evolve, users can change their

search criteria. Additionally, users can generate and use other reference TE genome files

and use *FishPi* for other model systems through the custom options. This makes *FishPi*

particularly powerful for users looking at non-model systems. The software has a graphic

user interface for *FishPi* (Figure 1A), and the outputs can be exported as high-resolution

figures presenting raw count data of the different target TE categories for presentation

(Figure 1B). Additionally, users can export a .csv file with all complementary TE mRNA

sequences, along with genomic co-ordinates, allowing the user to conduct further

downstream analyses and explore each class of TE individually. *FishPi*'s advantage is that

as more research into piRNA-TE seeds are validated, the user can customise their search

and input different seed regions for analysis. Additionally, users can use this for any species

and there is option for custom species input.

Tools for the study of piRNAs and tools for piRNA target analyses do exist but have limited

application bandwidths. *PiRNAQuest* V.2 for example is an online tool to annotate and

identify piRNA clusters and it can identify piRNA targets, but these are limited to mouse and

diseases related to cancers [34]. *piRPred* is piRNA identification tool based on an identification

algorithm with k- nearest neighbours' kernel to improve piRNA classification [35]. The tool

miRanda is primarily designed for miRNA gene target prediction to look at piRNA target

genes, rather than TEs [36,37]. *FishPi* fills a current gap and offers a user-friendly, adaptable

and customisable interface for identifying transposons targeted by piRNAs.

Conclusions

Currently available tools to link piRNA and TEs are limited and require knowledge of known

targets or are restricted to a pre-defined genome and species due to piRNA binding rules [14-

16,38]. *FishPi* is ideal for users with a list of candidate piRNAs. This is tool that can easily be

adapted to any organism in a user-friendly GUI format. *FishPi* is fully adaptable and

customisable so will continue to be user-friendly and useable as our understanding of piRNA

targeting improves. It generates detailed reports and files that can be used for further

analysis including statistical analysis of TE family targets in enrichment bar charts, loci

analysis in chromosomal plots, and exportable spreadsheets with piRNA:TE targets with

scoring via percentage complementarity for further filtering. *FishPi* software is freely

available and is distributed under GPL-3.0 licence.

Availability and requirements

Project name: FishPi

Project home page: https://github.com/alicegodden/fishpi

Operating system(s): Platform independent

Programming language: Python

Other requirements: Python v3.11

License: GPL-3.0

Any restrictions to use by non-academics: GPL-3.0 license needed

Declarations

Ethics

No ethics approvals were required for this research project.

Consent for publication

All authors consent to publication.

Availability of data and materials

FishPi is available on the Git repository here: https://github.com/alicegodden/fishpi

Competing interests

Authors declare no financial or competing interests.

Funding

This project was funded by a Consolidator Grant from the European Research Council to SI

(SELECTHAPLOID – 101).

Author's contributions

AMG conceived the idea for the project, AMG and BR wrote the software, BR tested the

software, AMG wrote the manuscript and SI contributed to the writing and provided guidance

throughout.

Acknowledgements

AMG would like to thank Immler lab colleagues for testing FishPi.

List of abbreviations

mRNA- Messenger RNA

PiRNA- Piwi-Interacting RNA

TE- Transposable element

References

1   Godden, A. M. & Immler, S. The potential role of the mobile and non-coding
genomes in adaptive response. *Trends Genet* (2022).
https://doi.org/10.1016/j.tig.2022.08.006
2   Doolittle, W. F. & Sapienza, C. Selfish genes, the phenotype paradigm and genome
evolution. *Nature* **284**, 601-603 (1980). https://doi.org/10.1038/284601a0
3   McClintock, B. The origin and behavior of mutable loci in maize. *Proc Natl Acad Sci*
U S A **36**, 344-355 (1950). https://doi.org/10.1073/pnas.36.6.344
4   Boeke, J. D., Garfinkel, D. J., Styles, C. A. & Fink, G. R. Ty elements transpose
through an RNA intermediate. *Cell* **40**, 491-500 (1985). https://doi.org/10.1016/0092-
8674(85)90197-7
5   Bourque, G. *et al.* Ten things you should know about transposable elements. *Genome*
Biol **19**, 199 (2018). https://doi.org/10.1186/s13059-018-1577-z
6   Chang, N. C., Rovira, Q., Wells, J., Feschotte, C. & Vaquerizas, J. M. Zebrafish
transposable elements show extensive diversification in age, genomic distribution,
and developmental expression. *Genome Res* **32**, 1408-1423 (2022).
https://doi.org/10.1101/gr.275655.121
7   Houwing, S. *et al.* A role for Piwi and piRNAs in germ cell maintenance and
transposon silencing in Zebrafish. *Cell* **129**, 69-82 (2007).
https://doi.org/10.1016/j.cell.2007.03.026

8   Brennecke, J. *et al.* Discrete small RNA-generating loci as master regulators of
transposon   activity   in   Drosophila.   *Cell*   **128**,   1089-1103   (2007).
https://doi.org/10.1016/j.cell.2007.01.043

9   Wei, C., Salichos, L., Wittgrove, C. M., Rokas, A. & Patton, J. G. Transcriptome-
wide analysis of small RNA expression in early zebrafish development. *RNA* **18**, 915-
929 (2012). https://doi.org/10.1261/rna.029090.111

10   Spichal, M. *et al.* Germ granule dysfunction is a hallmark and mirror of Piwi mutant
sterility. *Nat Commun* **12**, 1420 (2021). https://doi.org/10.1038/s41467-021-21635-0

11   Sultana, T., Zamborlini, A., Cristofari, G. & Lesage, P. Integration site selection by
retroviruses and transposable elements in eukaryotes. *Nat Rev Genet* **18**, 292-308
(2017). https://doi.org/10.1038/nrg.2017.7

12   Gebrie, A. Transposable elements as essential elements in the control of gene
expression. *Mob DNA* **14**, 9 (2023). https://doi.org/10.1186/s13100-023-00297-3

13   Houwing, S., Berezikov, E. & Ketting, R. F. Zili is required for germ cell
differentiation and meiosis in zebrafish. *EMBO J* **27**, 2702-2711 (2008).
https://doi.org/10.1038/emboj.2008.204

14   Zhang, D. *et al.* The piRNA targeting rules and the resistance to piRNA silencing in
endogenous   genes.   *Science*   **359**,   587-592   (2018).
https://doi.org/10.1126/science.aao2840

15   Shen, E. Z. *et al.* Identification of piRNA Binding Sites Reveals the Argonaute
Regulatory Landscape of the C. elegans Germline. *Cell* **172**, 937-951 e918 (2018).
https://doi.org/10.1016/j.cell.2018.02.002

16   Gou, L. T. *et al.* Pachytene piRNAs instruct massive mRNA elimination during late
spermiogenesis. *Cell Res* **24**, 680-700 (2014). https://doi.org/10.1038/cr.2014.41

17   Kaaij, L. J., Hoogstrate, S. W., Berezikov, E. & Ketting, R. F. piRNA dynamics in
divergent zebrafish strains reveal long-lasting maternal influence on zygotic piRNA
profiles. *RNA* **19**, 345-356 (2013). https://doi.org/10.1261/rna.036400.112

18   Gunawardane, L. S. *et al.* A slicer-mediated mechanism for repeat-associated siRNA
5' end formation in Drosophila. *Science* **315**, 1587-1590 (2007).
https://doi.org/10.1126/science.1140494

19   Wang, C. & Lin, H. Roles of piRNAs in transposon and pseudogene regulation of
germline   mRNAs   and   lncRNAs.   *Genome   Biol*   **22**,   27   (2021).
https://doi.org/10.1186/s13059-020-02221-x

20   Gainetdinov, I., Colpan, C., Arif, A., Cecchini, K. & Zamore, P. D. A Single
Mechanism of Biogenesis, Initiated and Directed by PIWI Proteins, Explains piRNA
Production   in   Most   Animals.   *Mol   Cell*   **71**,   775-790   e775   (2018).
https://doi.org/10.1016/j.molcel.2018.08.007

21   Belicard, T., Jareosettasin, P. & Sarkies, P. The piRNA pathway responds to
environmental signals to establish intergenerational adaptation to stress. *BMC Biol* **16**,
103 (2018). https://doi.org/10.1186/s12915-018-0571-y

22   Kurhanewicz, N. A., Dinwiddie, D., Bush, Z. D. & Libuda, D. E. Elevated
Temperatures   Cause   Transposon-Associated   DNA   Damage   in   C.   elegans
Spermatocytes.   *Curr   Biol*   **30**,   5007-5017   e5004   (2020).
https://doi.org/10.1016/j.cub.2020.09.050

23   Jardim, S. S., Schuch, A. P., Pereira, C. M. & Loreto, E. L. Effects of heat and UV
radiation on the mobilization of transposon mariner-Mos1. *Cell Stress Chaperones* **20**,
843-851 (2015). https://doi.org/10.1007/s12192-015-0611-2

24   He, J. *et al.* Identifying transposable element expression dynamics and heterogeneity
during development at the single-cell level with a processing pipeline scTE. *Nat*
Commun **12**, 1456 (2021). https://doi.org/10.1038/s41467-021-21808-x

25 Agarwal, V., Bell, G. W., Nam, J. W. & Bartel, D. P. Predicting effective microRNA
target sites in mammalian mRNAs. *Elife* **4** (2015).
https://doi.org/10.7554/eLife.05005
26 Wang, J. *et al.* piRBase: a comprehensive database of piRNA sequences. *Nucleic*
Acids Res **47**, D175-D180 (2019). https://doi.org/10.1093/nar/gky1043
27 Rosenkranz, D., Zischler, H. & Gebert, D. piRNAclusterDB 2.0: update and
expansion of the piRNA cluster database. *Nucleic Acids Res* **50**, D259-D264 (2022).
https://doi.org/10.1093/nar/gkab622
28 Foundation, P. S. Python Language reference, vesion 3.11.
https://python.org/downloads/release/python-3110/ (2022).
https://doi.org/https://python.org/downloads/release/python-3110/
29 Clark, A. https://buildmedia.readthedocs.org/media/pdf/pillow/latest/pillow.pdf
(readthedocs, 2015).
30 Hunter, J. D. Matplotlib: A 2D graphics environment. *Computing in Science and*
Engineering **9**, 90-95 (2007). https://doi.org/10.1109/MCSE.2007.55
31 Storer, J., Hubley, R., Rosen, J., Wheeler, T. J. & Smit, A. F. The Dfam community
resource of transposable element families, sequence models, and genome annotations.
Mob DNA **12**, 2 (2021). https://doi.org/10.1186/s13100-020-00230-y
32 Karolchik, D. *et al.* The UCSC Table Browser data retrieval tool. *Nucleic Acids Res*
32, D493-496 (2004). https://doi.org/10.1093/nar/gkh103
33 Jin, Y., Tam, O. H., Paniagua, E. & Hammell, M. TEtranscripts: a package for
including transposable elements in differential expression analysis of RNA-seq
datasets. *Bioinformatics* **31**, 3593-3599 (2015).
https://doi.org/10.1093/bioinformatics/btv422
34 Ghosh, B. *et al.* piRNAQuest V.2: an updated resource for searching through the
piRNAome of multiple species. *RNA Biol* **19**, 12-25 (2022).
https://doi.org/10.1080/15476286.2021.2010960
35 Brayet, J., Zehraoui, F., Jeanson-Leh, L., Israeli, D. & Tahi, F. Towards a piRNA
prediction using multiple kernel fusion and support vector machine. *Bioinformatics*
30, i364-370 (2014). https://doi.org/10.1093/bioinformatics/btu441
36 John, B. *et al.* Human MicroRNA targets. *PLoS Biol* **2**, e363 (2004).
https://doi.org/10.1371/journal.pbio.0020363
37 Wang, X. *et al.* Global expression and functional analysis of human piRNAs during
HSV-1 infection. *Virus Res* **328**, 199087 (2023).
https://doi.org/10.1016/j.virusres.2023.199087
38 Wu, W. S. *et al.* pirScan: a webserver to predict piRNA targeting sites and to avoid
transgene silencing in C. elegans. *Nucleic Acids Res* **46**, W43-W48 (2018).
https://doi.org/10.1093/nar/gky277
343

 Figures and Tables

 Table 1- Transposable element families assigned to transposable element class.

Class	Transposable element family
DNA	hAT, Tc1, Tc-Mar, Harbinger, Enspm, Kolobok, Merlin, Crypton, PiggyBac, Dada, Zatar, Ginger, TDR, Polinton, Maverick, Acrobat, Looper, TZF, Angel, Mariner
LTR	Gypsy, DIRS, Ngaro, ERV, Pao, Copia, BEL, HERV, Bhikari
LINE	L1, L2, L1-Tx1, Rex-Babar, RTE, Penelope, Keno, Rex
SINE	Alu, tRNA-V-RTE
RC	Helitron
Satellite	BRSATI, MOSAT

Figure 1- *The graphic user interface of FishPi with and pop-out window with exportable*
results for dre-piRNA 5'-TACACGAAGACTGTGGTGTGATTGGGCG-3'. The buttons
included allow users to export the plots at hi-resolution for presentation and publication.
Additionally, the user may export a .csv file list of all complementary TE mRNAs for further
downstream analysis.

357   ***Figure 2** – Flow chart covering the key steps and processes of FishPi.*