1 **Title: *FishPi*: a bioinformatic prediction tool to link piRNA and transposable elements**
2 **in zebrafish**
3

4 Alice M. Godden[1], Benjamin Rix[1], Simone Immler[1]
5
6 [1] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,
7 NR4 7TJ, United Kingdom
8
9 Corresponding author:
10 **Alice M. Godden** Email: alice.godden@uea.ac.uk

**Abstract**

**Background**

Piwi-interacting RNAs (piRNA)s are non-coding small RNAs that post-transcriptionally

affect gene expression and regulation. Through complementary seed region binding with

transposable elements (TEs), piRNAs protect the genome from transposition, and

therefore a tool to link piRNAs with complementary TE targets is needed. Tools like

*TEsmall* can process sRNA-seq datasets to produce differentially expressed piRNAs and

*piRScan* developed for nematodes can link piRNAs and TEs but it requires the user to

know the target region of interest and work backwards.

**Results**

We have therefore developed *FishPi* to predict the pairings between piRNA and TEs.

*FishPi* works with individual piRNAs or a list of piRNA sequences in fasta format. The

software focuses on the piRNA:TE seed region and analyses reference TEs for piRNA

complementarity. TE type is examined, counted and stored to a dictionary, with genomic

loci recorded.  Any updates to piRNA-TE binding rules, can easily be incorporated by

updating the code underlying *FishPi*. *FishPi* provides a graphic interface, using *tkinter*,

that requires the user to input piRNA sequences to generate comprehensive reports on

piRNA:TE dynamics. *FishPi* can easily be adapted to other genomes opening the

interpretation of piRNA functionality to a wide community.

**Conclusions**

Users will gain insight into genome age and *FishPi* will help further our understanding of

the biological role of piRNAs and their interaction with TEs in a similar way that public

databases have improved the access to and the understanding of the role of small

RNAs.

40
41  **Keywords**
42  PiRNAs, transposons, evolution, zebrafish

## Background

Environmental changes and related biotic and abiotic stressors can disrupt the balance and expression of small RNAs and transposable elements (TEs) in the genome [1]. TEs are parasitic DNA that can cause insertions in the genome, as well as deletions and genomic rearrangements [2]. To counter this activity, Piwi-interacting RNAs (piRNAs), have evolved to repress the action of these repetitive elements [3]. PiRNAs constitute a substantial fraction of small RNAs and are predominantly found in animal germ cells [4]. The main function of piRNAs is to silence TEs and repetitive elements in animal germ cells to preserve the integrity of the germline genome [5] and loss of TE silencing by piRNAs can lead to sterility for example in the nematode *Caenorhabditis elegans* [6]. TEs are parasitic and are not randomly distributed in the genome but have preferred regions for insertion in the genome, including distorted and bent DNA [7,8], or open chromatin regions [8,9]. Large populations of piRNAs are complementary to TE sequences and messenger RNAs (mRNA) [10]. In *C. elegans*, the 2[nd] to the 7[th] nucleotide at the 5' end of the piRNA [11] or the 2[nd] to the 8[th] nucleotide [12] have both been suggested to be the piRNA:TE seed region. Similarly, in *Mus musculus*, the 2[nd] to 8[th] nucleotide was a predicted seed region [13] and in *Drosophila melanogaster*, the first 10 nucleotides at the 5' end of the piRNA are predicted to be the piRNA:TE seed binding site [14]. Finally, in zebrafish and other teleost fish, the first 10 nucleotides at the 5' end of the piRNA are loaded into the Argonaute machinery to produce Ago2-bound piRNAs complementary to TEs [3,15].

TEs are known to be activated through environmental stressors. In *C. elegans* for example, thermal stress has been shown in mutant piRNA biogenesis strains that lack the ability to generate piRNAs. In this work nematodes were heat-treated and displayed transgenerational impairments for up to three subsequent generations: with thermal stress decreasing the rate of piRNA biogenesis. [16,17]. In *Drosophila,* the *mariner-Mos1* transposon is enriched following heat-stress but not following ultraviolet irradiation [18]. TEs are expressed in a tissue-specific manner and can affect transcription and gene expression [19], by generating new genes and

mRNAs through genomic rearrangements and *de novo* mutations, and cause alteration of

regulatory networks and even trigger immune responses [7]. Understanding the interaction

between piRNAs and their target activities is therefore key to understand the defence

mechanisms in the germ line against the effects of environmental conditions.

In the zebrafish (*Danio rerio*) genome, small RNAs post-transcriptionally regulate gene

expression and regulation [20]. Piwi proteins are predominantly expressed in the animal germ

line and associate with piRNAs. The loss of Ziwi (zebrafish Piwi) leads to apoptosis of germ

cells in embryonic development [3]. Ziwi and Zili zebrafish Piwi proteins provide a piRNA

amplification system that is highly conserved. TEs represent over 50% of the zebrafish

genome [21] and it is therefore key to improve our understanding of the mechanisms that

evolved to allow an organism to regulate their activity. TEs have over 2000 families in the

zebrafish and exploit the host's genome by dynamic transposition events. The mechanisms

underlying this transposition can be divided into two categories: retrotransposons, these TEs

synthesise an RNA intermediate that can insert itself into the host genome [22]. In contrast,

DNA transposons cut-and-paste themselves into the host genome [7,22,23].

We therefore developed the software FishPi to provide a means to map zebrafish piRNAs

against complementary TEs and link them to understand their biological association [10].

FishPi facilitates research into the function of piRNAs and accelerates analyses in the same

way as microRNA databases like Targetscan have supported small RNA research [24]. FishPi

provides a user-friendly graphic interface (Figure 1), which is based on the latest zebrafish

reference genome and TE sequences and generates a comprehensive output that can be

exported as a report including complementary TE counts, family level annotation,

chromosomal and counts plots, with the option to export the full list of TEs and their

sequences that are complementary to the piRNA sequence fed in by the user. The user can

enter individual piRNAs or lists of piRNAs in fasta format. For zebrafish and other model

organisms, a list of piRNA sequences are available at piRBase

([http://bigdata.ibp.ac.cn/piRBase/browse.php](http://bigdata.ibp.ac.cn/piRBase/browse.php)), [4] or piRNAclusterDB

([https://www.smallrnagroup.uni-mainz.de/piRNAclusterDB/](https://www.smallrnagroup.uni-mainz.de/piRNAclusterDB/)), [25].

**Implementation**

FishPi is a python-based programme and uses the following packages and versions: Python

v3.11, Tkinter (included in Python v3.11 [26]), Pillow 10.0.1 [27], and Matplotlib 3.8.0 [28]. The

code for the programme was written and tested using PyCharm Community Edition 2023.2.1

and is platform-independent, as it is designed to be executed from the Linux command line;

however, the software can be run and used in PyCharm IDE. The software initially defines

functions to assay piRNA TE complementarity, creating a dictionary to count the TEs by

class. The families assigned to each class were checked using dfam.org (Table 1), [29].

To generate a fasta file of the TE sequences to use in place of the zebrafish reference

genome GRCz11 TE sequences, a TE.bed file with clade was created by using the UCSC

table genome browser [30] as follows: Vertebrate, group: Variation and Repeats, genome:

zebrafish, assembly: May 2017 GRCz11, track: RepeatMasker, table: rmsk. To extract DNA

sequences from the reference genome based on the co-ordinates supplied in the TE.bed

file, the following command lines were entered:

bedtools getfasta -s -name -fi Danio_rerio.GRCz11.dna.primary_assembly.fa -fo
GRCz11.teseqs.use.fasta -bed GRCz11.teannotation.bed

The following command line serves to mask the sequences in the reference genome:

bedtools maskfasta -fi Danio_rerio.GRCz11.dna.primary_assembly.fa -fo
GRCz11.masked.fasta -bed GRCz11.teannotation.bed

To create the TE-merged reference genome [31]:

cat GRCz11.masked.fasta GRCz11.teseqs.1.fasta > GRCz11.teseqs.fishpi.fasta

This preparatory work was based on the pipeline described for PopoolationTE2 [32].  The

Ensembl reference genome GRCz11 v107 was used in this project and can be retrieved

here:                         https://ftp.ensembl.org/pub/release-107/fasta/danio_rerio/dna/

"Danio_rerio.GRCz11.dna.primary_assembly.fa.gz" [33]. These instructions can be adjusted

for other reference genomes and allow the user to run the analyses on any available

genome version.

**Results & Discussion**

FishPi successfully links piRNAs and TEs and is a helpful tool for anyone interested in

studying the role of TEs and sRNAs in shaping the germline genome. FishPi is designed to

be used by those with minimal coding experience. A detailed walk-through use and

installation    is    listed    in    the    readme    on    the    git    repository    here:

https://github.com/alicegodden/fishpi . The simple graphic user interface offers a point and

click tool to rapidly gain insight into many piRNA sequences. Each time a new piRNA

sequence is provided, the programme resets itself before providing the new results. As more

research into piRNA:TE seed sites evolve, FishPi can be adapted and updated to reflect

changes and updates to the reference genome. Alternatively, users can generate and use

other reference TE genome files and use FishPi for other model systems. This makes FishPi

particularly powerful for users looking at non-model systems. The software has a graphic

user interface for FishPi (Figure 1), and the outputs can be exported as hi-resolution bar-

charts and chromosomal plots presenting raw count data of the different target TE

categories for presentation (Figure 1, Supplementary Files 2 & 3). Additionally, users can

export a .csv file with all complementary TE mRNA sequences, along with genomic co-

ordinates, allowing the user to conduct further downstream analyses and explore each class

of TE individually (Supplementary File 1). Benchmarking data can be found in

Supplementary File 4 for optimal running of the software. FishPi's advantage over other tools

is that it can be easily updated and modified as more research into piRNA:TE seeds are

validated and will be fully customisable for future genome releases for other understudied

species including fish and other model systems.

**Conclusions**

By analysing piRNA complementarity to a TE, we can infer recent genomic expansion but

also adaptive evolution. For DNA class TEs, we can find footprints of evolutionary activity, as

DNA TEs are often associated with past transposition events. In the zebrafish genome, DNA

TEs comprise 40%, with retrotransposons comprising 10% zebrafish genome [23]. Currently

available tools to link piRNA and TEs are limited and require knowledge of known targets or

are restricted to a pre-defined genome and species due to piRNA binding rules [11-13,34].

Further analysis with RNA-seq data and tools such as RepeatMasker and RepeatLandscape

would complement this by Kimura distance analysis [35]. Additionally, use of the Retroseq

pipeline can highlight non-reference insertion mutations by TEs from high-coverage DNA-

seq data and help predict TE activity [36]. In the future, other adaptations of the software could

be made to look at other small RNAs. The software allows users to rapidly confirm

predictions of complementary TEs from specific piRNAs. FishPi software is freely available

and is distributed under GPL-3.0 licence.

**Availability and requirements**

**Project name:** FishPi

**Project home page:** https://github.com/alicegodden/fishpi/tree/fishpi

**Operating system(s):** Platform independent

**Programming language:** Python

**Other requirements:** Python v3.11

**License:** GPL-3.0

**Any restrictions to use by non-academics:** GPL-3.0 license needed

**Declarations**

**Ethics**

No ethics approvals were required for this research project.

**Consent for publication**

All authors consent to publication.

**Availability of data and materials**

FishPi is available on the Git repository here: https://github.com/alicegodden/fishpi/tree/fishpi

**Competing interests**

Authors declare no financial or competing interests.

**Funding**

This project was funded by a Consolidator Grant from the European Research Council to SI

(SELECTHAPLOID – 101).

**Author's contributions**

AMG conceived the idea for the project, AMG and BR wrote the software, BR tested the software, AMG wrote the manuscript and SI contributed to the writing and provided guidance throughout.

**Acknowledgements**

AMG would like to acknowledge Prof. Oliver Buckley from University of East Anglia for software publication and licensing advice.

**List of abbreviations**

mRNA- Messenger RNA

PiRNA- Piwi-Interacting RNA

TE- Transposable element

**References**

1    Godden, A. M. & Immler, S. The potential role of the mobile and non-coding genomes in adaptive response. *Trends Genet* (2022). https://doi.org/10.1016/j.tig.2022.08.006

2    Kofler, R. piRNA Clusters Need a Minimum Size to Control Transposable Element Invasions. *Genome Biol Evol* **12**, 736-749 (2020). https://doi.org/10.1093/gbe/evaa064

3    Houwing, S. *et al.* A role for Piwi and piRNAs in germ cell maintenance and transposon silencing in Zebrafish. *Cell* **129**, 69-82 (2007). https://doi.org/10.1016/j.cell.2007.03.026

4    Wang, J. *et al.* piRBase: a comprehensive database of piRNA sequences. *Nucleic*
*Acids Res* **47**, D175-D180 (2019). https://doi.org/10.1093/nar/gky1043
5    Brennecke, J. *et al.* Discrete small RNA-generating loci as master regulators of
transposon    activity    in    Drosophila.    *Cell*    **128**,    1089-1103    (2007).
https://doi.org/10.1016/j.cell.2007.01.043
6    Spichal, M. *et al.* Germ granule dysfunction is a hallmark and mirror of Piwi mutant
sterility. *Nat Commun* **12**, 1420 (2021). https://doi.org/10.1038/s41467-021-21635-0
7    Bourque, G. *et al.* Ten things you should know about transposable elements. *Genome*
*Biol* **19**, 199 (2018). https://doi.org/10.1186/s13059-018-1577-z
8    Sultana, T., Zamborlini, A., Cristofari, G. & Lesage, P. Integration site selection by
retroviruses and transposable elements in eukaryotes. *Nat Rev Genet* **18**, 292-308
(2017). https://doi.org/10.1038/nrg.2017.7
9    Gebrie, A. Transposable elements as essential elements in the control of gene
expression. *Mob DNA* **14**, 9 (2023). https://doi.org/10.1186/s13100-023-00297-3
10   Houwing, S., Berezikov, E. & Ketting, R. F. Zili is required for germ cell
differentiation    and    meiosis    in    zebrafish.    *EMBO J* **27**, 2702-2711 (2008).
https://doi.org/10.1038/emboj.2008.204
11   Zhang, D. *et al.* The piRNA targeting rules and the resistance to piRNA silencing in
endogenous    genes.    *Science*    **359**,    587-592    (2018).
https://doi.org/10.1126/science.aao2840
12   Shen, E. Z. *et al.* Identification of piRNA Binding Sites Reveals the Argonaute
Regulatory Landscape of the C. elegans Germline. *Cell* **172**, 937-951 e918 (2018).
https://doi.org/10.1016/j.cell.2018.02.002
13   Gou, L. T. *et al.* Pachytene piRNAs instruct massive mRNA elimination during late
spermiogenesis. *Cell Res* **24**, 680-700 (2014). https://doi.org/10.1038/cr.2014.41
14   Aravin, A. A., Hannon, G. J. & Brennecke, J. The Piwi-piRNA pathway provides an
adaptive defense in the transposon arms race. *Science* **318**, 761-764 (2007).
https://doi.org/10.1126/science.1146484
15   La, Y. *et al.* Identification and Characterization of Piwi-Interacting RNAs for Early
Testicular    Development    in    Yak.    *Int J Mol Sci* **23** (2022).
https://doi.org/10.3390/ijms232012320
16   Belicard, T., Jareosettasin, P. & Sarkies, P. The piRNA pathway responds to
environmental signals to establish intergenerational adaptation to stress. *BMC Biol* **16**,
103 (2018). https://doi.org/10.1186/s12915-018-0571-y
17   Kurhanewicz, N. A., Dinwiddie, D., Bush, Z. D. & Libuda, D. E. Elevated
Temperatures Cause Transposon-Associated DNA Damage in C. elegans
Spermatocytes.    *Curr    Biol*    **30**,    5007-5017    e5004    (2020).
https://doi.org/10.1016/j.cub.2020.09.050
18   Jardim, S. S., Schuch, A. P., Pereira, C. M. & Loreto, E. L. Effects of heat and UV
radiation on the mobilization of transposon mariner-Mos1. *Cell Stress Chaperones* **20**,
843-851 (2015). https://doi.org/10.1007/s12192-015-0611-2
19   He, J. *et al.* Identifying transposable element expression dynamics and heterogeneity
during development at the single-cell level with a processing pipeline scTE. *Nat*
*Commun* **12**, 1456 (2021). https://doi.org/10.1038/s41467-021-21808-x
20   Wei, C., Salichos, L., Wittgrove, C. M., Rokas, A. & Patton, J. G. Transcriptome-
wide analysis of small RNA expression in early zebrafish development. *RNA* **18**, 915-
929 (2012). https://doi.org/10.1261/rna.029090.111
21   Huang, C. R., Burns, K. H. & Boeke, J. D. Active transposition in genomes. *Annu Rev*
*Genet* **46**, 651-675 (2012). https://doi.org/10.1146/annurev-genet-110711-155616

22　Boeke, J. D., Garfinkel, D. J., Styles, C. A. & Fink, G. R. Ty elements transpose
through an RNA intermediate. *Cell* **40**, 491-500 (1985). https://doi.org/10.1016/0092-
8674(85)90197-7

23　Chang, N. C., Rovira, Q., Wells, J., Feschotte, C. & Vaquerizas, J. M. Zebrafish
transposable elements show extensive diversification in age, genomic distribution,
and developmental expression. *Genome Res* **32**, 1408-1423 (2022).
https://doi.org/10.1101/gr.275655.121

24　Agarwal, V., Bell, G. W., Nam, J. W. & Bartel, D. P. Predicting effective microRNA
target sites in mammalian mRNAs. *Elife* **4** (2015).
https://doi.org/10.7554/eLife.05005

25　Rosenkranz, D., Zischler, H. & Gebert, D. piRNAclusterDB 2.0: update and
expansion of the piRNA cluster database. *Nucleic Acids Res* **50**, D259-D264 (2022).
https://doi.org/10.1093/nar/gkab622

26　Foundation, P. S. Python Language reference, vesion 3.11.
https://python.org/downloads/release/python-3110/ (2022).
https://doi.org/https://python.org/downloads/release/python-3110/

27　Clark, A. https://buildmedia.readthedocs.org/media/pdf/pillow/latest/pillow.pdf
(readthedocs, 2015).

28　Hunter, J. D. Matplotlib: A 2D graphics environment. *Computing in Science and*
*Engineering* **9**, 90-95 (2007). https://doi.org/10.1109/MCSE.2007.55

29　Storer, J., Hubley, R., Rosen, J., Wheeler, T. J. & Smit, A. F. The Dfam community
resource of transposable element families, sequence models, and genome annotations.
*Mob DNA* **12**, 2 (2021). https://doi.org/10.1186/s13100-020-00230-y

30　Karolchik, D. *et al.* The UCSC Table Browser data retrieval tool. *Nucleic Acids Res*
**32**, D493-496 (2004). https://doi.org/10.1093/nar/gkh103

31　Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841-842 (2010).
https://doi.org/10.1093/bioinformatics/btq033

32　Kofler, R., Gomez-Sanchez, D. & Schlotterer, C. PoPoolationTE2: Comparative
Population Genomics of Transposable Elements Using Pool-Seq. *Mol Biol Evol* **33**,
2759-2764 (2016). https://doi.org/10.1093/molbev/msw137

33　Cunningham, F. *et al.* Ensembl 2022. *Nucleic Acids Res* **50**, D988-D995 (2022).
https://doi.org/10.1093/nar/gkab1049

34　Wu, W. S. *et al.* pirScan: a webserver to predict piRNA targeting sites and to avoid
transgene silencing in C. elegans. *Nucleic Acids Res* **46**, W43-W48 (2018).
https://doi.org/10.1093/nar/gky277

35　Smit, A. F., Hubley, R. & Green, P. RepeatMasker software (2013).
https://doi.org/http://www.repeatmasker.org

36　Keane, T. M., Wong, K. & Adams, D. J. RetroSeq: transposable element discovery
from next-generation sequencing data. *Bioinformatics* **29**, 389-390 (2013).
https://doi.org/10.1093/bioinformatics/bts697

 **Figures and Tables**

302 *Table 1- Transposable element families assigned to transposable element Class (verified*

303 *using dfam.org).*

| Class | Transposable element family |
|---|---|
| DNA | hAT, Tc1, Tc-Mar, Harbinger, Enspm, Kolobok, Merlin, Crypton, PiggyBac, Dada, Zatar, Ginger, TDR, Polinton, Maverick, Acrobat, Looper, TZF, Angel, Mariner |
| LTR | Gypsy, DIRS, Ngaro, ERV, Pao, Copia, BEL, HERV, Bhikari |
| LINE | L1, L2, L1-Tx1, Rex-Babar, RTE, Penelope, Keno, Rex |
| SINE | Alu, tRNA-V-RTE |
| RC | Helitron |
| Satellite | BRSATI, MOSAT |

306 *Figure 1- The graphic user interface of FishPi with and pop-out window with exportable*

307 *results for dre-piRNA-53095 5'- CGGATAAGCATATTGCCCTAGCAACATC - 3'. The buttons*

308 *included allow users to export the plots at hi-resolution for presentation and publication.*

309 *Additionally, the user may export a .csv file list of all complementary TE mRNAs for further*

310 *downstream analysis.*