1 **Sheltered load in fungal mating-type chromosomes revealed by fitness**
2 **experiments**
3

4 Lou Guyot[1,2], Elizabeth Chahine[1], Elsa De Filippo[1], Christophe Lalanne[3], Sylvain Brun[4],
5 Fanny E. Hartmann[1*],Tatiana Giraud[1*]
6
1 Ecologie, Systematique et Evolution, CNRS, Universite Paris-Saclay, AgroParisTech, Gif-
sur-Yvette, France
2 Master de Biologie, Ecole Normale Supérieure de Lyon, Universite Claude Bernard Lyon
1, Universite de Lyon, 69342 Lyon Cedex 07, France
3 Laboratoire Interdisciplinaire des Energies de Demain, Université Paris Cité, Paris, France
4 Université Paris Cité, Institut Jacques Monod UMR-7592, 15 rue Hélène Brion 73205 Paris
cedex 13.
14
15 *FEH and TG are joint senior authors on this work.
16
17
18 Keywords: heterozygous advantage, mutational load, self-incompatibility locus, sex
19 chromosome, fungi, mating-type locus
20
21 Corresponding author: tatiana.giraud@universite-paris-saclay.fr

22
23

Abstract

Sex chromosomes and mating-type chromosomes can carry large regions with suppressed

recombination. As a result of a lower efficacy of selection, recessive deleterious mutations

are expected to accumulate in these non-recombining regions. Multiple genomic analyses

have indirectly inferred the presence of deleterious mutations in sex and mating-type

chromosomes, but direct experimental evidence remains scarce. Here, we performed fitness

assays in fungi with megabase-large and young non-recombining regions around the mating-

type locus, using three Sordariales species, to test whether heterokaryons (diploid-like,

heterozygous at the mating-type locus) exhibited a fitness advantage over homokaryons

(haploid-like, with a single mating-type allele), in terms of spore germination dynamics or

mycelium growth speed, under different conditions of light and temperature. We found a faster growth of heterokaryons compared to one of the homokaryons for *Podospora anserina* at 18°C and for *Schizothecium tetrasporum* and *Schizothecium tritetrasporum* at 22°C under light. These findings suggest the presence of a sheltered load, i.e., recessive deleterious mutations at the heterozygous state in or near non-recombining regions, associated to a specific mating-type allele. Genomic analyses indeed suggested that the non-recombining regions around the mating-type locus likely carries heterozygous deleterious mutations, while the rest of the genome was mostly homozygous. We also showed that the difference in growth rates did not result from different numbers or densities of nuclei between homokaryons and heterokaryons. Leveraging the experimental assets of fungi, allowing cultivating separately haploid-like and diploid-like life stages, our experiments provided one of the rare direct experimental evidence of sheltered load around mating-compatibility loci, which is crucial for our understanding of sex-related chromosome evolution.

Social Media Abstract

Experimental evidence for sheltered load around the mating-type locus in filamentous fungi: slower growth of haploid versus diploid-like mycelia, revealing recessive deleterious mutations, in *Podospora anserina* and other Sordariales fungi.

Introduction

Even in sexual species, genomes can display large non-recombining regions (Wright et al., 2016, Schwander et al., 2014, Bergero & Charlesworth, 2009). Such non-recombining regions are particularly frequent around sex-determining loci and mating-type loci (Charlesworth, 2017, Ma & Veltsos, 2021, Hartmann et al., 2021b, Jay et al., 2024, Nicolas et al., 2004, Bachtrog, 2013). Deprived of recombination, these genomic regions experience a reduced efficacy of selection (Felsenstein, 1974, Muller, 1932, Hill & Robertson, 1966, Rice & Chippindale, 2001). As a consequence, it has been theoretically suggested that non-recombining regions are bound to degenerate (Smith & Haigh, 2009, Muller, 1932, Hill & Robertson, 1966, Fisher, 1930), i.e., to accumulate recessive deleterious mutations following recombination suppression. In addition, theoretical models have shown that recessive deleterious mutations can be maintained for some times in populations when tightly linked to self-incompatibility or mating-type loci, especially when weakly deleterious, generating mutational load associated to a specific self-incompatibility or mating-type allele (Tezenas et al., 2023, Antonovics & Abrams, 2004, Glémin et al., 2001, Uyenoyama, 2005). One therefore expects recessive deleterious mutations to be present in or near non-recombining regions around sex-determining or mating-type determining alleles. Such recessive deleterious mutations associated with a permanently heterozygous allele constitute a sheltered load, as they are also heterozygous and therefore not exposed to selection, being protected by the functional copy of the alternative allele in the heterozygote.

The presence of deleterious mutations in non-recombining mating-type and sex chromosomes has been indirectly inferred through genomic analyses, that have for example revealed gene disruptions, gene losses, non-synonymous substitutions, transposable element accumulation

and chromosomal rearrangements (Carpentier et al., 2022, Vittorelli et al., 2023, Duhamel et

al., 2023, Bachtrog, 2013, Hough et al., 2014, Whittle et al., 2011). However, there has been

little direct evidence for the existence of such sheltered load through experimental fitness

assays so far. To perform such experiments, one indeed needs to compare fitness between a

haploid phase or a homozygous state and the heterozygous diploid state, i.e., between

situations where mutations are exposed to selection to situations where deleterious mutations

are sheltered at a heterozygous state. While Y-like chromosomes are never homozygous, non-

recombining regions can be found homozygous in some autosomal supergenes, i.e. large

regions in autosomes governing adaptive and complex phenotypes and segregating as a single

unit (Schwander et al., 2014). Using such autosomal supergenes that can be homozygous, the

existence of sheltered load could be demonstrated in some cases (Jay et al., 2021). A few

balanced haplo-lethal systems even exist, in which recessive deleterious mutations are lethal

at the homozygous state in non-recombining supergenes (Wielstra, 2020, Hill et al., 2023).

However, in the majority of organisms with non-recombining regions on sex-related

chromosomes, haploid or homozygous states do not exist in the life cycle and cannot be

generated. The rare exceptions include self-incompatibility loci in plants that can be rendered

homozygous under certain conditions. Sheltered load was found associated with some self-

incompatibility alleles in the plants *Arabidopsis halleri* (Stone, 2004, Llaurens et al., 2009,

Le Veve et al., 2024) and *Solanum carolinense* (Mena-Alí et al., 2009).

Fungi can be highly tractable models for investigating the existence of a sheltered load in

sex-related chromosomes. Indeed, there can be recombination suppression in large genomic

regions around their mating-type loci in species with a long heterokaryotic (diploid-like)

phase, i.e., with nuclei of opposite mating types per cell, but unfused, which is functionally

similar to a diploid stage (Branco et al., 2018, Hartmann et al., 2021b, Grognet et al., 2014,

Menkis et al., 2008, Vittorelli et al., 2023). Genomic footprints of degeneration have been

found in these non-recombining regions, in the form of gene losses, transposable element

accumulation, non-synonymous substitutions and non-optimal codon usage and expression

(Carpentier et al., 2022, Vittorelli et al., 2023, Duhamel et al., 2023, Whittle et al., 2011).

Mating types in fungi do not prevent selfing, as mating compatibility is controlled at the

haploid stage (Billiard et al., 2012, Giraud et al., 2008). As a matter of fact, the fungal

lineages with recombination suppression around their mating-type loci most often undergo

regular automixis i.e., intra-tetrad selfing (Billiard et al., 2012, Giraud et al., 2008, Hood &

Antonovics, 2000, Zakharov, 2005, Zakharov, 2023, Grognet et al., 2014, Jay et al., 2024,

Vittorelli et al., 2023). Such life cycles in fungi are called pseudo-homothallic, with the

production of self-fertile heterokaryotic spores, encompassing two nuclei of opposite mating

types originating from a single tetrad after meiosis, thus germinating as a self-fertile

heterokaryon and undergoing mostly automixis. Self-sterile spores with a single mating type,

germinating as homokaryotic mycelia, can also generally be produced in laboratory

conditions, although their importance in nature is unknown.

With such life cycles, experiments can be set up for comparing fitness between homokaryons

(i.e., with a single type of nuclei per cell) and heterokaryons (i.e., with nuclei of opposite

mating types in each cell) in order to study the heterozygous advantage in the region around

the mating-type locus. Indeed, natural heterokaryotic strains are typically highly homozygous

genome-wide except around the mating-type locus, because of high selfing rates (Branco et

al., 2017, Grognet et al., 2014, Hartmann et al., 2021a, Hartmann et al., 2021b, Vittorelli et

al., 2023). Therefore, any variability in fitness between homokaryotic and heterokaryotic

stages should mainly be due to heterozygous recessive deleterious mutations present in the

non-recombining region around the mating-type locus and possibly in its flanking region due to linkage disequilibrium (Branco et al., 2017, Tezenas et al., 2023, Glémin et al., 2001) (Fig 1A). Recessive lethal alleles linked to the mating-type locus have been documented in anther-smut fungi of the *Microbotryum* genus and in the fungal-like oomycete *Phytophthora infestans* (Judelson et al., 1995, Hood & Antonovics, 2000, Thomas et al., 2003, Oudemans et al., 1998).

We studied here three species of two phylogenetically distant fungal species complex within the Sordariales order of Ascomycota: *Schizothecium tetrasporum* and *Schizothecium tritetrasporum,* belonging to the *Schizothecium tetrasporum* species complex (Vittorelli et al., 2023, De Filippo et al., 2025), and *Podospora anserina* (Grognet et al., 2014, Hartmann et al., 2021a, Boucher et al., 2017). These species are coprophilous, saprophyte or endophyte, and can be found in herbivore faeces and soil (Silar, 2020). These fungi are excellent biological models to investigate the presence of a sheltered load around their mating-type locus, displaying two mating-type alleles (*MAT1-1* and *MAT1-2*, also called *mat-* and *mat+*, respectively). Indeed, these fungi display recombination suppression around their mating-type locus, as shown by progeny segregation and nucleotide divergence analyses (Grognet et al., 2014, Hartmann et al., 2021a, Vittorelli et al., 2023). These species are pseudo-homothallic, i.e. with their main life cycle phase being heterokaryotic (Grognet et al., 2014, Vittorelli et al., 2023, De Filippo et al., 2025), so that sheltered load can be expected. Recombination suppression around the mating-type locus spans approximately 1 Mb in the *S. tetrasporum* and *P. anserina* species complexes and has been suggested to be quite recent, as indicated by the low divergence between mating types and the lack of trans-specific polymorphism (Vittorelli et al., 2023, Grognet et al., 2014, Hartmann et al., 2021a). Indirect evidence of sheltered load has been reported by genomic analyses in the non-recombining

regions of the reference strains, in particular a few gene losses or disruptions (Vittorelli et al.,

2023, Grognet et al., 2014). Furthermore, recombination suppression around the mating-type

locus represents convergent events between the *S. tetrasporum* and *P. anserina* species

complexes, that diverged over 150 million years ago (Hyde et al., 2017), which provides

independent events of recombination suppression to study. Even within each of these two

species complexes, recombination suppression has likely evolved independently in the

various species, as suggested by the lack of trans-specific polymorphism and sizes of the non-

recombining regions (Hartmann et al., 2021a). In addition, natural strains are highly

homozygous genome-wide, except around the mating-type locus, as shown in multiple strains

of *P. anserina* and in the reference strain of *S. tetrasporum* (Vittorelli et al., 2023, Grognet et

al., 2014, Hartmann et al., 2021a). Another asset of these models is that one can cultivate *in*

vitro homokaryotic and heterokaryotic mycelia (Fig 1B). In addition, while most ascospores

are heterokaryotic, homokaryotic spores are sometimes produced and are easily

distinguishable based on their smaller size (Vittorelli et al., 2023, Silar, 2020) (Fig 1C). In *S.*

tetrasporum and *P. anserina,* previous experiments on a single strain each did not detect any

heterokaryon advantage (Grognet et al., 2014, Vittorelli et al., 2023). Studying multiple

strains is however essential in evolutionary biology, as the effects of deleterious mutations

can depend on epistatic effects and/or be present in only some individuals in populations,

while still having important population effects. Studying various situations and diverse fitness

components is also important, as the sheltered load may only be detectable in some

conditions, for example if recessive deleterious mutations affect genes expressed only in

specific situations, or if stressful conditions exacerbate deleterious effects. Examples of

stressful conditions for these species can be temperature or darkness, as fungi do not tolerate

high temperatures and use light as an important regulator of diverse physiological processes,

as shown in particular in *P. anserina* (Shen et al., 2022, Silar, 2020).

Here, we thus conducted assays using different fitness proxies on a dozen strains of each of *S.*

tetrasporum, *S. tritetrasporum* and *P. anserina*, by comparing heterokaryons and

homokaryons, in terms of mycelium growth speed under light or dark, at different

temperatures, and in terms of spore germination dynamics, and we analysed genomes. We

expected that i) strains would be homozygous genome-wide except around the mating-type

locus, ii) heterozygous non-synonymous mutations would be present in the non-recombining

region around the mating-type locus of our strains, iii) at least one homokaryon would have a

lower fitness than the heterokaryon, at least under some conditions, due to recessive

deleterious alleles associated to a specific mating type. Our experiments revealed as expected

a fitness advantage of heterokaryons over one of the homokaryons (either *MAT1-1* or *MAT1-*

2), in at least one tested condition, for *S. tetrasporum*, *S. tritetrasporum* and *P. anserina*. This

indicates the likely existence of a sheltered load, associated to a specific mating type allele, in

the non-recombining region present around the mating-type locus and/or in its flanking

region. Genome analyses indeed showed that most of the strains used here in *S. tetrasporum*

and *S. tritetrasporum* were homozygous genome-wide, except around the mating-type locus,

as previously shown in multiple strains in *P. anserina* (Hartmann et al., 2021a) and in the

reference strain of *S. tetrasporum* (Vittorelli et al., 2023). Furthermore, we detected

heterozygous non-synonymous mutations in the non-recombining region, with in particular

gains of stop codons and loss of start codons, that are likely deleterious. Counts of nuclei in

the reference strain of *P. anserina* suggested that the difference in growth rates did not result

from different numbers or densities of nuclei between homokaryons and heterokaryons,

which had, surprisingly, not been investigated in this model organism yet.

Material and Methods

Strains

The information on the strains is given in Supplementary Table S1 (strain ID, species, collection, year, substrate, geographic region of collection, accession number and bioproject number on Genbank). We used 10 heterokaryotic strains of *S. tetrasporum* and 13 heterokaryotic strains of *S. tritetrasporum* stored at -80°C (De Filippo et al., 2025). In the strain ID, the letters A, B, C and D, when present, refer to the spore ID within the isolated tetrads, that were the same as in (De Filippo et al., 2025). We also used 17 heterokaryotic strains of *P. anserina,* from their *MAT1-1* and *MAT1-2* homokaryotic mycelium stored at -80°C (Silar et al., 2024). Most of these strains were isolated as heterokaryons in 2019-2022 from nature in different areas in France, including the Corsica island, but some were isolated from other countries, including New Zealand, Italy, the Netherlands and Canada. Strains were isolated from soil or from faeces of rabbit, cow, bull, sheep, hare, goat, wallaby or horse. These strains had been stored at -80°C upon isolation. In addition, we used two *S. tetrasporum* strains from the Westerdijk Fungal Biodiversity Institute Collection and the reference *P. anserina* S strain. These three strains may have experienced culture in laboratories, including as homokaryons.

For *S. tetrasporum* and *S. tritetrasporum*, we obtained homokaryotic mycelium of the *MAT1-1* and *MAT1-2* mating types for each heterokaryotic strain stored at -80°C as described previously (Vittorelli et al., 2023, Silar, 2020). Briefly, a plug of heterokaryotic mycelium (5 mm³) was grown for each strain on a V8 medium Petri dish for two to three weeks until perithecia formation. Then, agar Petri dishes were added on the top, upside down, to collect projected spores (Silar, 2020). Homokaryotic spores were then isolated, identified by their smaller size compared to heterokaryotic spores, and grown on Petri dishes with G medium

(Silar, 2020) (Fig 1C). These Petri dishes were subjected to a heat shock (30 minutes at 65°C)

to induce spore germination. After a few days of growth, mycelia were transferred on M2

medium for growth. The mating-type allele of homokaryotic mycelium grown from those

spores was identified by PCR. For this goal, DNA was extracted using Chelex and the

mating-type allele (*MAT1-1* or *MAT1-2*) was determined using the PCR conditions described

previously (Vittorelli et al., 2023) and the following primer pairs: *MAT1-2*

5''GAGGGTGCGAGTCGTGAT'3' and 3''CCAGCCACACGAAAATCC'5'; *MAT1-1*

5''GATCGGATTCGCTCCAGA'3' and 3''CGACCGTTGGAAATGACC'5'.

Genome sequencing and analysis

We used the genomes of 13 of our heterokaryotic strains (CBS815.71, PSN779, PSN881A

and PSN710A for *S. tetrasporum,* and PSN352A, PSN571A, PSN1070A, PSN667A,

PSN777A, PSN829C, PSN1040, PSN1236 and PSN1433A for *S. tritetrasporum*) available

from previous studies (Vittorelli et al., 2023, De Filippo et al., 2025). Genomes have been

downloaded at NCBI Genbank under Bioproject numbers PRJNA882797 and

PRJNA1190690 (accession numbers in Table S1). DNA had been extracted with the

Nucleospin soil kit from Macherey Nagel. Library preparation and genome sequencing had

been performed in six different batches by Genewiz, Genoscope or Novogene. The genomes

had been sequenced with the Illumina technology (2x150 bp paired-end Novaseq) and had an

average coverage of 89X. SNPs were called in (De Filippo et al., 2025). To summarise

briefly, we checked Illumina raw read quality using FastQC v0.11.1 v.0.11.5 (Andrews,

2010). Reads were trimmed with Trimmomatic v0.36 (Bolger et al., 2014) in paired-end

mode with TruSeq3 adapter and the following options: PE -threads 2 -phred33 TruSeq3-

PE.fa:2:30:10 LEADING:10 TRAILING:10 SLIDINGWINDOW:5:10 MINLEN:50.

Illumina trimmed reads were mapped against the CBS815.71-sp3 (MAT1-2) long-read

assembly available from the Genbank BioProject accession number PRJNA882797

(assembly accession number JAQKAE000000000 (Vittorelli et al., 2023)) with bowtie2

v2.3.4.1 (Langmead & Salzberg, 2012) with the options -very-sensitive-local -phred33 - X

1000. We used the MarkDuplicates tool from PicardCommandLine v2.8.1 to get rid of

duplicated sequences. We called SNPs against the CBS815.71-sp3 assembly with the

HaplotypeCaller tool of the Genome Analysis Toolkit (GATK) v4.1.2.0 (McKenna et al.,

2010) in the diploid mode. We used CombineGVCF, GenotypeGVCF and SelectVariants to

merge data from all strains and conjointly call SNPs. GATK Good Practice rules were

followed with the following options: QD=20.0; FS=60.0; MQ=20.0; MQRankSumNeg=-2.0;

MQRankSumPos=2.0; <SOR=3.0; QUAL=100. SelectsVariants and VCFtools v0.1.17 were

used to filter out indels, and to keep only good-quality SNPs with the 90% missingness

option. We predicted SNP effect with SNPeff (Cingolani et al., 2012), recording whether they

were synonymous, missense or inducing the gain/loss of stop or start codons, compared to the

CBS815.71-sp3 reference genome. We considered that autosomes carried heterozygous

regions when at least four consecutive genes carried at least one heterozygous SNP, with less

than 50kb between the midpoints of two consecutive genes.

The divergence time between haplotypes in the non-recombining region of *S. tetrasporum*,

i.e. a proxy for the age of recombination suppression, was estimated using computed per-gene

d_S values (synonymous divergence) from a previous study on the *S. tetrasporum* reference

genome CBS815.71 (Vittorelli et al., 2023). The number of generations T_g since

recombination suppression can be estimated by $T_g = d_S/2\mu$, μ being the synonymous

substitution rate per generation. Previous estimates of synonymous substitution rates in

fungal ascomycetes ranged from $0.9.10^{-9}$ to $16.7.10^{-9}$ substitutions per site per year (Kasuga

et al., 2002). We used these two values for our computations to obtain a range of dates, as the

substitution rate is unknown in *S. tetrasporum*. We considered that *S. tetrasporum* undergoes

five generations per year. Indeed, a maximum of 15 generations/year can be obtained in

optimal laboratory conditions, which are generally harsher in the wild, with often too low

temperature in autumn, winter and spring for growth. To obtain T_y, the time since

recombination suppression in years, we thus use the formula $T_y = T_g/5$.

Inoculum preparation for experiments

Prior to each experiment, *MAT1-1* and *MAT1-2* homokaryotic mycelia of the studied

heterokaryotic strains were grown on M2 medium (Silar, 2020) for about 7 to 15 days at

27°C, 50% humidity and light and then stored at 4°C for a few days until the beginning of the

experiments. On the day of inoculation, fresh inoculum was prepared. Mycelium square plugs

of the two mating types for each strain were crushed separately (to form homokaryotic

mycelium solutions) and also together (to generate heterokaryotic mycelium solutions; (Silar,

2020, Vittorelli et al., 2023)), with 200 µL of sterile water for *S. tritetrasporum* and *S.*

tetrasporum and 150 µL of sterile water for *P. anserina*, using a TissueLyser at 30 Hz for 20

to 30 min.

Growth speed

Spots of 20 µL of the obtained solutions for homokaryotic mycelium *MAT1-1*, homokaryotic

mycelium *MAT1-2* or heterokaryotic mycelium were deposited on Petri dishes with M2

medium (Silar, 2020), with three replicates per strain and condition. One central spot was

deposited per Petri dish, corresponding to a replicate. For *S. tritetrasporum*, two batches were

performed at different times. Plates were incubated at the conditions of temperature and light

exposure detailed in Table 1. Pictures of the plates were taken on two days (see details of the

days for each species in Table 1). As the mycelium limits of *P. anserina* were not always

clearly distinguishable on pictures, the contour of the mycelial colony was drawn with a

marker pen under the Petri dish before taking the picture. Diameters of mycelia were

measured using ImageJ 1.54g (Schneider et al., 2012), and the growth speed was computed as

the difference in diameter between the two days divided by the number of growth days. We

measured one diameter per Petri dish, at the same location for the different time points, using

as a reference the label that had been put under the Petri dishes before growth. Some Petri

dishes were excluded from the analysis when contamination, dryness or satellite colonies

occurred (see Table S2 with raw data).

Heterokaryotic state persistence and loss of nuclei of one or the other mating type

PCRs were performed on heterokaryotic mycelia 14 days post-inoculation for all *S.*

tritetrasporum strains used for measures in the two experiment batches to check whether they

remained heterokaryotic. Nine plugs (5 mm³) were sampled in each Petri dish from the inner

to the outer of the mycelium ring. DNA was extracted using Chelex and the mating-type

allele (*MAT1-1*, *MAT1-2* or both) was determined using the PCR conditions and the primer

pairs described above. When only a faint band appeared for one of the two mating types, the

PCR was re-run to check the presence of the second band. Some PCRs did not yield any band

at all, for neither mating type, and the corresponding plugs were excluded from the analyses

(see Table S2 with raw data).

Number of nuclei per cell in heterokaryons versus homokaryons

We measured the number and density of nuclei per cell in the S strain of *P. anserina* in

homokaryons and heterokaryons. Three replicates of each of the *MAT1-1* and the *MAT1-2*

homokaryons, as well as of the heterokaryon, were cultured on standard M2 medium for 6

days at 18°C under light. For microscopic analysis, a one centimeter square sample located at

the edge of the thallus was cut, fixed in paraformaldehyde (ThermoFischer) 4% in PBT (PBS

Phosphate buffered saline + triton X100 (Merck) 0.3%) for 1h, washed in PBT (3X 10 min),

stained with 5 µg.mL^{-1} DAPI (4′,6-diamidino-2-phenylindole) in PBT for 30 min then

washed in PBT. Samples were mounted inverted on a coverslip in PBT and observed on a

Zeiss Axio Observer Z.1 fully motorized inverted microscope; Cool LED pE 400 White;

DAPI filter (365 D395 445/50); objective 63x/1.4 Plan-Apo Oil, WD: 0.19mm, ∞/0.17. For

each sample, observations were made in the two first millimeters from the edge of the thallus

(growth zone). Images were treated with the Fiji software (Schindelin et al., 2012). A total of

30 articles (the hyphal cells delimited by two septa) were selected at random in each

replicate, to avoid any biased selection of articles depending on their number of nuclei, and

their length was measured in the transmitted light canal. Then, the number of nuclei per

article was counted and the density was computed as the number of nuclei divided by the

length of articles (see Table S3 with raw data).

Germination dynamics

For the 10 heterokaryotic strains of *S. tetrasporum*, three spots of 20 µL of heterokaryotic

mycelium solution were deposited on a single Petri dish with V8 medium (Vittorelli et al.,

2023) and incubated at 22°C until perithecia formation (about 15 days). PSN881A and

PSN779 strains did not produce any perithecia and were excluded from the analysis. Agar

Petri dishes were added on the top, upside down, to collect projected spores (Vittorelli et al.,

2023). We isolated 10 large spores and 10 small spores per heterokaryotic strain, which were

deposited on a Petri dish with G medium (Silar, 2020), allowing germination (two batches per

strain). Small spores are homokaryotic, while large spores are heterokaryotic at the mating-

type locus with a 90% probability, as estimated in (Vittorelli et al., 2023) based on

genotyping of the germinated mycelium, which is a minimum estimate, as the mycelium

could also have lost a nucleus during growth before genotyping. These Petri dishes were

foiled with aluminium paper, subjected to a heat shock of 30 min at 65°C and incubated at

27°C to induce germination (Silar, 2020). Germinated spores were counted on days 2, 3 and 7

to analyse the germination dynamics of small and large spores, i.e., the rates of germination

over time (Table 1). A few spores were excluded from the analysis when there was

contamination or two spores were observed germinating instead of one (see Table S4 with

raw data).

Graphical summaries and statistical analyses

All statistical analyses and graphical displays were carried out using the R software v 4.3.2

(R Core Team, 2023). A statistical significance level of 5% was considered for all statistical

tests. The linearity of the response and predictor relationship and the Gaussian distribution of

the residuals in all models (including random effects intercept and slope in the case of

random-effects models) were assessed visually using R built-in facilities (fitted vs. residual

values plot and QQ-plot, respectively).

For the growth experiment analyses in *S. tritetrasporum*, *S. tetrasporum* and *P. anserina*,

linear mixed-effect models were fitted, using the restricted maximum likelihood method from

the lme4 package, considering as fixed effects either ploidy (i.e., heterokaryon or

homokaryon) or the nuclear status (i.e., heterokaryon, *MAT1-1* homokaryon or *MAT1-2*

homokaryon), in addition to the growing condition, when applicable (i.e. light or dark for *S.*

tetrasporum, 18°C or 37°C for *P. anserina*). The interaction between the growing condition

and the ploidy or the nuclear status was also included in the models. The strain ID was

considered a random effect in all models, as well as the batch in the case of *S.*

tritetrasprorum. The statistical significance of fixed effects was assessed using F-tests with

the lmerTest package and a Satterthwaite approximation for degrees of freedom. The

emmeans package was used to estimate marginal means and their associated standard errors

and to estimate predefined pairwise contrasts, with degrees of freedom calculated using the

Kenward-Roger method. Obtained p-values from Student t-tests were adjusted using the

Holm method to account for multiple testing.

For the analysis testing the preferential loss of nuclei of one or the other mating type in

heterokaryotic mycelia in *S. tritetrasporum*, a generalized linear mixed-effects model with a

Binomial distribution was used. The model was designed to assess the proportion of plugs in

which nuclear loss occurred, with the genotype of the plug after the loss (i.e., *MAT1-1* or

MAT1-2) as a fixed effect and strain ID and batch as random effects. For this analysis, a

generalized linear mixed-effect model using the maximum likelihood method with the

Laplace approximation for random effects was fitted with the lme4 package. The statistical

significance of the fixed effect was assessed using a χ^2-test with the car package. The

ggeffect package was used to estimate marginal means and their associated standard errors.

The emmeans package was used to estimate a predefined pairwise contrast using a z-test.

For nucleus number analysis in the S strain of *P. anserina*, a generalised linear model with a

Poisson distribution was used, modelling the number of nuclei per article considering the

nuclear status (i.e., heterokaryon, *MAT1-1* homokaryon or *MAT1-2* homokaryon) as a fixed

effect. The statistical significance of the fixed effect was assessed using a χ^2-test with the car

package. The ggeffect package was used to estimate marginal means and their associated

standard errors. The emmeans package was used to estimate predefined pairwise contrasts

and obtained p-values from z-test were adjusted using the Holm method to account for

multiple testing. A proxy of the nucleus density was computed as the number of nuclei

divided by the size of the article, and a linear model was fitted, modelling the density of

nuclei per article considering the nuclear status (i.e., heterokaryon, *MAT1-1* homokaryon or

MAT1-2 homokaryon) as a fixed effect. The statistical significance of the fixed effect was

assessed using a F-test. The emmeans package was used to estimate marginal means and their

associated standard errors and to estimate predefined pairwise contrasts. Obtained p-values

from Student t-tests were adjusted using the Holm method to account for multiple testing.

For the germination experiment analysis in *S. tetrasporum*, a generalised linear mixed-effect

model with a Binomial distribution was used, modelling germination rate as a linear

combination of day (as a continuous predictor) and spore size (i.e. large or small), treated as

fixed effects, and strain ID and batch as random effects. For this analysis, a generalized linear

mixed-effect model using the maximum likelihood method with the Laplace approximation

for random effects was fitted with the lme4 package. The statistical significance of fixed

effects was assessed using χ^2-tests with the car package. The ggeffect package was used to

estimate marginal means and their associated standard errors. The emmeans package was

used to estimate a predefined pairwise contrast using a z-test.

418   Results

Genome-wide homozygosity except in the mating-type locus region, that carries likely

heterozygous deleterious mutations

To assess whether heterozygosity in the genomes of the strains used for experiments was

mainly located around the mating-type locus and whether signs of heterozygous deleterious

mutations could be detected, we used the available genomes of four and nine heterokaryotic

strains of *S. tetrasporum* and *S. tritetrasporum* (De Filippo et al., 2025). All strains displayed

heterozygosity around their mating-type locus, in a genomic region ranging from 1.38Mb to

1.47Mb in *S. tetrasporum* and 1.30Mb to 1.71Mb kb in *S. tritetrasporum* (Figs 2, S1 and S2).

As reported for the CBS815.71 *S. tetrasporum* reference strain (Vittorelli et al., 2023),

distinct evolutionary strata seemed to be present in all strains, i.e. segments with different

divergence levels (Figs 2, S1 and S2). We estimated the age of recombination suppression in

S. tetrasporum based on previous estimates of synonymous differences in the non-

recombining region in the reference genome CBS815.71 (Vittorelli et al., 2023, Hartmann et

al., 2021a) and synonymous substitutions rates in ascomycetes (Kasuga et al., 2002), to be

between 40 ky and 730 ky for the oldest stratum, and between 9 ky and 170 ky for the

youngest stratum.

Among the genomes of the 13 heterokaryotic strains analysed, only four showed some tracks

of heterozygosity in autosomes, corresponding to one *S. tetrasporum* strain (PSN710A; Fig

S1) and three *S. tritetrasporum* strains (PSN777A, PSN352A and PSN829C; Figs 2 and S2);

the heterozygous regions were short: on average, the strains with heterozygosity on the

autosomes had a total of 910 kb of heterozygous regions on their autosomes, i.e., only 3.6%

of the total autosomal genomic size. The nine other strains were homozygous genome-wide,

except in the non-recombining region near the mating-type locus (Figs S1 and S2).

For all strains analysed, the region around the mating-type locus also harbored heterozygous

non-synonymous mutations, especially changes in amino-acids, but also gains of stop codons

and losses of start codons (Figs 2, Fig S1 and Fig S2), suggesting that deleterious mutations

may be present at the heterozygous state. We detected no heterozygous SNP inducing the loss

of a stop codon or the gain of a start codon compared to the reference genome.

Growth advantage of heterokaryons

To investigate the presence of sheltered load in the non-recombining regions around the

mating-type locus, or at their margin, in *S. tetrasporum*, *S. tritetrasporum* and *P. anserina*,

experiments were performed to compare the fitness of heterokaryons to that of homokaryons

using strains as biological replicates in each species (10, 13 and 17 strains, respectively;

Table S1; Fig 1).

For *S. tetrasporum* and *S. tritetrasporum*, we used as fitness proxies the mycelium growth

speed under light at 22°C between day 6 and day 14. In addition, we used as fitness proxies

for *S. tetrasporum* the mycelium growth speed under dark at 22°C and assessed the spore

germination dynamics between day 2 and day 7. For *P. anserina*, we measured mycelium

growth speed between day 2 and day 6 under light at 18°C and 37°C. These conditions were

chosen for each species based on preliminary experiments or previous knowledge on the

different species, to use more or less stressful conditions. Table S2 gives the raw measures

for the growth experiment for which 8% of plates have been excluded on average per species

due to the presence of satellite colonies, contaminants or dryness.

In *S. tritetrasporum*, heterokaryons grew significantly faster than homokaryons under light at

22°C, by 3.7% (see Fig 3A showing raw data, Fig 3B showing model estimates, Table S5A).

When considering the two types of homokaryons separately in the statistical analysis, we

found that heterokaryons grew significantly faster than *MAT1-1* homokaryons, by 5.7%.

MAT1-2 homokaryons had a tendency to grow more slowly than heterokaryons, but the

difference was not significant (Fig 3A showing raw data, Fig 3C showing model estimates,

Table S5B).

In *S. tetrasporum*, heterokaryons also grew significantly faster than homokaryons under light

at 22°C, by 3.6% (see Fig 4A showing raw data, Fig 4B showing model estimates, Table

S5C; the global model was not significant, but individual tests for pairwise comparisons,

relying on unpooled variance between all conditions, indicate a significant difference

between homokaryons and heterokaryons for growth speed under light). The difference in

growth speed between homokaryons and heterokaryons was not significant under dark

conditions, although the tendency was similar (Fig 4A and B, Table S5C). When separating

the two homokaryons in the analysis, we found that heterokaryons grew significantly faster

than *MAT1-1* homokaryons under light at 22°C, by 5.3% (see Fig 4A showing raw data, Fig

4C showing model estimates, Table S5D). Heterokaryons had a tendency to also grow faster

than MAT1-2 homokaryons under both light and dark conditions, albeit the difference was

not significant (Figs 4A and C, Table S5D).

In *P. anserina*, heterokaryons had a tendency to grow faster than homokaryons at the two

temperatures, but no differences were significant when considering both types of

homokaryons together (see Fig 5A showing raw data, Fig 5B showing model estimates, Table

S5E). However, when considering the two types of homokaryons separately, heterokaryons

grew significantly faster than *MAT1-2* homokaryons at 18°C, by 7.9% (see Fig 5A showing

raw data, Fig 5C showing model estimates, Table S5F). The tendency was similar at 37°C but

not significantly so (Figs 5A and 5C, Table S5F).

Re-running all statistical analyses after removing the strains having likely experienced

laboratory culture (i.e., CBS370.69, CBS815.71 and the S strain) and those showing traces of

heterozygosity in autosomes (i.e. PSN710A, PSN777A, PSN352A and PSN829C) did not

change the conclusions: the same fixed effects and pairwise differences as above remained

significant (not shown). In *S. tetrasporum,* this even led to a significant effect of ploidy on

mycelium growth, and rendered the global model significant.

Heterokaryotic mycelia lost nuclei of one mating type in small proportions, and with no preferential loss for one mating type over the other

Given that some level of nuclear loss of one mating type has been reported to occur during heterokaryotic mycelium growth in the *S. tetrasporum* species complex (Vittorelli et al., 2023), while it does not occur in *P. anserina*, we investigated whether, and to what extent, such a phenomenon occurred in the *S. tritetrasporum* heterokaryotic mycelia used for growth experiments. For this goal, we genotyped the mating-type locus for nine plugs per Petri dish. Table S2 gives the PCR results, 3% of the plugs from plates of *S. tritetrasporum* heterokaryotic mycelium used for growth measures having been excluded because PCRs did not yield any band at all, for neither mating type.

We found that mycelia had remained mostly heterokaryotic, although we detected a single mating type in 27.7% of the plugs (Figs S3 and S4A). In these plugs where a single mating-type was detected, there was no significant over-representation of one mating type over the other one (see Fig S4A showing raw data, Fig S4B showing model estimates, Table S5G).

No differences in nuclei number or density between heterokaryotic and homokaryotic mycelia

As nucleus number per cell might vary between heterokaryotic and homokaryotic mycelia, then possibly impacting their growth rates, we investigated whether the number or the density of nuclei per cell differed between heterokaryons and each of the homokaryons in the reference *P. anserina* S strain. Table S3 gives the raw measures of nucleus counts and article lengths, articles being hyphal cells delimited by two septa.

The differences in the number and density of nuclei per article between heterokaryons and

each of the homokaryons in the *P. anserina* S strain were small and not significant (Fig S5

showing a picture of nucleus counts; Figs S6A and S6C showing raw data, Figs S6B and S6D

showing model estimates; Tables S5H and S5I).

No differences in germination rates between heterokaryotic and homokaryotic spores

We also investigated the existence of a heterokaryon fitness advantage in terms of spore

germination, using the same *S. tetrasporum* strains as for growth experiments. Table S4 gives

the raw measures for the germination experiment, for which two out of the 10 strains of *S.*

tetrasporum were excluded due to lack of perithecia and less than 1% of the spores were

excluded, due to contamination or the germination of two spores, meaning that two spores

had been deposited instead of one.

The comparison of the germination dynamics of small spores (i.e. homokaryotic spores)

versus large spores (i.e. mainly heterokaryotic spores; (Vittorelli et al., 2023)) in *S.*

tetrasporum did not reveal any evidence of heterozygous advantage. Indeed, although large

spores had a tendency to germinate at a higher rate than small spores, the difference in

germination rate between small and large spores was not significant (see Fig 6A showing raw

data, Fig 6B showing model predictions, Table S5J). Adding an interaction between the two

fixed effects, i.e. spore size and day, and/or considering the variable day as a factor in the

statistical model did not alter the previous conclusions (not shown).

Discussion

Likely presence of recessive deleterious mutations around the mating-type locus

Our fitness assay showed that heterokaryons were fitter than at least one homokaryon under light at 18°C in *P. anserina* and at 22°C in *S. tetrasporum* and *S. tritetrasporum*, suggesting the presence of recessive deleterious mutations in the non-recombining region around the mating-type locus, and/or tightly linked to it, being associated to a specific single mating-type allele. We indeed confirmed that *S. tetrasporum* and *S. tritetrasporum* strains were mostly homozygous genome-wide, except around the mating-type locus, as previously shown in *P. anserina* (Hartmann et al., 2021a). We further revealed the existence of heterozygous non-synonymous mutations in the non-recombining region, in particular changes in amino-acids, gains of stop codons and losses of start codons, that constitute potential deleterious mutations. Our findings thus indicate the existence of recessive deleterious mutations around the mating-type locus, that are expressed at the homokaryotic state, but sheltered at the heterokaryotic state, and that impact more homokaryons of a mating type than the alternative mating type, and only under particular conditions. Note that, even if the sheltered load was only detected at some temperature, under light and during growth (and not germination), it still means that sheltered load exists. The sheltered load caused 3-8% of decrease in mycelium growth speed, which represents a non-negligible fitness impact, especially for colonizing a dung before competitors or before it dries out. Our study is one of the rare experiments showing the existence of a sheltered load in sex-related chromosomes. Only a few studies could investigate a sheltered load so far, around self-incompatibility loci in plants (Stone, 2004, Llaurens et al., 2009, Le Veve et al., 2024) as this is one of the rare systems where mating-type loci can be rendered homozygous. To investigate the presence of a sheltered load, we leveraged another unique biological system, pseudo-homothallic fungi,

whose life cycle allows fitness comparison between individuals in which mutations around

the mating-type locus are sheltered versus exposed to selection, while the rest of the genome

is exposed to selection, being highly homozygous due to high selfing rates in nature. Previous

evidence of a sheltered load in fungal-like organisms include haplo-lethal alleles linked to

mating types in anther-smut fungi and oomycetes, in which one or the two mating types fail

to grow at the haploid stage or cause a segregation bias in progenies (Hood & Antonovics,

2000, Thomas et al., 2003, Oudemans et al., 1998, Judelson et al., 1995). The non-

recombining regions were, however, much older in these cases, i.e., million years old

(Duhamel et al., 2022, Duhamel et al., 2023), while we estimated here that recombination

suppression occurred in *S. tetrasporum* ca. 730 ky ago. Few other studies have compared

fitness proxies between homokaryons and heterokaryons in fungi. In three sibling species of

the species complex *Neurospora tetrasperma*, other Sordariales fungi with recombination

suppression around its mating-type locus, the quantity of asexual haploid spores (conidia)

produced by mycelia has been compared between heterokaryons and homokaryons (Meunier

et al., 2018). No evidence of sheltered load was found, but a single strain has been used for

these experiments in each of the sibling species (Meunier et al., 2018). Recessive deleterious

mutations have also been experimentally detected in a few autosomal supergenes (Jay et al.,

2021, Wielstra, 2020, Hill et al., 2023). Altogether, such direct experimental evidence of

sheltered load is important for understanding the evolution of non-recombining regions.

Sheltered load detected in homokaryons of a single mating type and not in all conditions

The lower fitness of homokaryons was not detected in all traits or all conditions, and was

significant in a single one of the homokaryons, in all three species tested. This may be

explained by the young age of the recombination suppression, having allowed the

accumulation of only few deleterious mutations so far, in a few genes, perhaps only involved

in tolerance to particular conditions, and by chance more in the flanking region of one of the

mating-type alleles. The genomic analyses in fact suggest differences between mating-type

chromosomes in terms of non-synonymous substitutions.

A possible non-negligible haploid phase can also purge, to some extent, recessive mutations

with large detrimental effects on some traits and in some specific conditions. Indeed,

although rare and with an unknown importance in nature, some homokaryotic spores are

produced in these fungi: in *S. tetrasporum*, 2% to 10% of asci with small (i.e. homokaryotic)

spores have been reported (Raju & Perkins, 1994, Vittorelli et al., 2023); in *P. anserina*, 5%

of spores are homokaryotic (Ament-Velásquez & Vogan, 2022). The loss of nuclei of one

mating type during mycelium growth, as shown here and previously (Vittorelli et al., 2023),

may also expose recessive deleterious mutations to selection.

The growth as haploids could be linked to laboratory conditions, as a few of the strains used

in our experiments have been maintained in laboratories for some time (in particular the *P.*

anserina S strain), where they can be cultivated as homokaryons. If these strains have

experienced purifying selection as homokaryons more often than wild strains, this may have

reduced fitness differences between homokaryons and heterokaryons in some conditions; the

effect would therefore be conservative regarding the differences that we detected. In addition,

stock strains are maintained at -80°C, from where they are taken for each experiment, likely

limiting this effect. Furthermore, removing these strains from analyses did not change our

results.

Another mechanism allowing the purge of deleterious mutations could be the occurrence of

very rare events of recombination in the MAT-proximal region, as reported in *P. anserina*

(Contamine et al., 1996). Such extremely rare events of recombination, also reported in sex

chromosomes in frogs (Rodrigues et al., 2018), has been proposed to indeed prevent strong

differentiation and high level of degeneration of sex chromosomes (Rodrigues et al., 2018).

An explanation for a stronger sheltered load on homokaryons of one mating type could be an

over-representation of nuclei of one of the two mating types in heterokaryons, exposing more

the genomic background of this mating type to selection. Similarly, if the genome of one of

the two nuclei in each cell, associated to a particular mating type, is expressed at a higher

level, as suggested in *N. tetrasperma* (Meunier et al., 2022), selection could act more strongly

on this mating-type's genomic background. The effects of higher accumulation of deleterious

mutations by chance in one mating type's background and of stronger selection on the other

mating type's background could be mutually reinforcing: the first partly recessive deleterious

mutations occurring by chance more in the non-recombining regions of one mating type may

make nuclei of this mating type less expressed or replicating more slowly within cells, being

then under-represented, leading to stronger selection on the genomic background of the other

mating-type.

Our experiments actually suggest that nuclear losses of one mating type occur in sectors of

heterokaryotic mycelia in *S. tritetrasporum*, as previously reported in the *S. tetrasporum*

species complex (Vittorelli et al., 2023), but none of the mating types showed a preferential

tendency to be lost. Such occasional losses of nuclei mean that some parts of the mycelium in

the heterokaryotic treatment may actually be homokaryotic. Such a phenomenon, if any effect

in our experimental design, would have attenuated differences between homokaryons and

heterokaryons, thereby possibly preventing the detection of lower homokaryon fitness in

some conditions or making their disadvantage appear weaker than it is. In addition, large

spores have been shown to be sometimes homokaryotic in *S. tetrasporum* (less than 13% of

large spores germinated as homokaryons, which may also be due to nucleus loss during

growth (Vittorelli et al., 2023)); this could have artefactually reduced the difference between

homokaryotic and heterokaryotic spores and have prevented the detection of a lower fitness

of homokaryons for spore germination dynamics.

Alternative hypotheses to sheltered load

There may be, in principle, alternative hypotheses to sheltered load explaining the lower

fitness of homokaryons. Although the species studied mainly reproduce by selfing and most

strains display genome-wide homozygosity (Grognet et al., 2014, Hartmann et al., 2021a,

Vittorelli et al., 2023), some rare events of outcrossing can generate short tracks of

heterozygosity in genomic regions far from the mating-type locus. The analysis here of the

genomes of 13 heterokaryotic strains however showed that most strains displayed genome-

wide homozygosity, except around the mating-type locus, in *S. tetrasporum* and *S.*

tritetrasporum. Removing the strains with short tracks of heterozygosity in autosomes and

the laboratory strains from statistical analyses did not affect the general results, indicating

that the lower fitness of homokaryons was not due to these autosomal heterozygous regions.

Similar patterns have previously been reported in *P. anserina*: only two out of ten strains had

small heterozygous regions in autosomes (Hartmann et al., 2021a). The panel of *P. anserina*

strains analysed in the previous study only had the S strain in common with the set of strains

used here, and the S strain displays genome-wide homozygosity except around its mating-

type locus (Grognet et al., 2014).

Other types of heterozygous advantage can also exist, although such advantages are rare in

nature and mostly related to host-parasite interactions (Gemmell & Slate, 2006), and

therefore unlikely in our experimental setting. The lower fitness of homokaryons could also

have been due, in principle, to a ploidy advantage for heterokaryons, for example related to

gene expression dosage, which may have been expected to be twice as high in heterokaryons

as in homokaryons. We, however, found no evidence for a higher number or higher density of

nuclei per cell in heterokaryons compared to homokaryons in the S strain of *P. anserina,*

which had never been investigated so far in this model organism. Furthermore, this

hypothesis would predict a lower fitness of both homokaryons compared to heterokaryons,

while we found significance only between one homokaryon and the heterokaryon.

Another explanation for differences in growth between heterokaryons and homokaryons

could be differences in their developmental programs. No data or observation are available to

our knowledge to support this hypothesis, however, and similar expression patterns have been

found in homokaryons and heterokaryons in *N. tetrasperma*, except in the non-recombining

region (Meunier et al., 2022). The gene expression differences in *N. tetrasperma* between

homokaryons and heterokaryons (Meunier et al., 2022) are indeed restricted to genes around

the mating-type locus, which is likely due to degeneration following less efficient selection

(Fontanillas et al., 2015) rather than different developmental programs resulting from

selection for different behaviours. Here too, this hypothesis would predict a heterokaryon

advantage over each of the homokaryons.

Conclusion

In conclusion, our findings based on both genome analyses and experiments suggest the

presence of a sheltered load, i.e., of recessive deleterious mutations at the heterozygous state

around the mating-type locus, associated to a single mating type, in three species of

Sordariales fungi with young regions of recombination suppression. Leveraging the

experimental assets of fungi, allowing cultivating separately haploid-like and diploid-like life

stages, in a system with two mating types permanently heterozygous and genome-wide

homozygosity except around the mating-type locus, our experiments provided one of the rare

direct experimental evidence of sheltered load around mating-compatibility loci, which is

crucial for our understanding of sex-related chromosome evolution.

Competing interests: The authors have no competing interests.

Acknowledgements: This work was supported by the EvolSexChrom ERC advanced grant
#832352 (H2020 European Research Council) to T.G. We acknowledge the ImagoSeine core
facility of the Institut Jacques Monod, member of the France BioImaging infrastructure
(ANR-10-INBS-04) and GIS-IBiSA and the support of the Region Ile-de-France (Sesame).
The funders had no role in study design, data collection and analysis, decision to publish, or
preparation of the manuscript. We thank Philippe Silar for isolating heterokaryotic strains
used here of *Podospora anserina*, *Schizothecium tetrasporum* and *S. tritetrasporum*, for
designing the primer pairs, for the picture of spores in Figure 1 and for advice during the
experiment design. We thank Asen Daskalov, Kamalraj Subban and Alexandra Granger-
Farbos for isolating *MAT1-1* and *MAT1-2* homokaryons of the *Podospora anserina* strains
used in this study. We thank Albane Thibeault for help with nucleus counts. We thank Jacqui
Shykoff for discussions on statistical analyses. We thank the collectors of soil and animal
faeces: Philippe Silar, Delphine Paumier, Yves Hurand, Pierre Defos du Rau, Jean-Marie
Ouary (Milles Traces association), Jérome Letty (Office français de la biodiversité), Paul Jay,
Stella McQueen, Karen Vanderwolf, Jacqui Shykoff, Lucas Bonometti and Pierre Gladieux.

Author contribution: Conceptualization: TG, FEH and LG. Data curation: LG, SB, EDF.
Formal analysis: LG, CL, EDF. Funding acquisition: TG. Investigation: LG, TG, FEH, SB,
EC and EDF. Supervision: TG and FEH. Visualization: LG and EDF. Writing: LG, TG.
Writing – review & editing: LG, TG, FEH, CL, EDF.

References

Ament-Velásquez, S. L. & Vogan, A. A. 2022. Podospora anserina. *Trends in Microbiology*
30: 1243-1244.
Andrews, S. F. A. Q. C. T. f. H. T. S. D. O. 2010. FastQC: A Quality Control Tool for High
Throughput Sequence Data [Online]. **Available online at:**
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.
Antonovics, J. & Abrams, J. Y. 2004. Intratetrad mating and the evolution of linkage
relationships. *Evolution* **58**: 702-709.

Bachtrog, D. 2013. Y-chromosome evolution : emerging insights into processes of Y-
chromosome degeneration. *Nature Reviews Genetics* **14**: 113-124.
Bergero, R. & Charlesworth, D. 2009. The evolution of restricted recombination in sex
chromosomes. *Trends in Ecology & Evolution* **24**: 94-102.
Billiard, S., López-Villavicencio, M., Hood, M. & Giraud, T. 2012. Sex, outcrossing and
mating types: unsolved questions in fungi and beyond. *J Evol Biol* **25**: 1020-1038.
Bolger, A. M., Lohse, M. & Usadel, B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**: 2114-2120.
Boucher, C., Nguyen, T.-S. & Silar, P. 2017. Species delimitation in the *Podospora anserina,*
P. pauciseta, P. comata species complex (Sordariales). *Cryptogamie, Mycologie* **38**:
485-506.
Branco, S., Badouin, H., Rodríguez de la Vega, R. C., Gouzy, J., Carpentier, F., Aguileta, G.,
Siguenza, S., Brandenburg, J.-T., Coelho, M. A., Hood, M. E. & Giraud, T. 2017.
Evolutionary strata on young mating-type chromosomes despite the lack of sexual
antagonism. *Proceedings of the National Academy of Sciences* **114**: 7067-7072.
Branco, S., Carpentier, F., Rodríguez de la Vega, R. C., Badouin, H., Snirc, A., Le Prieur, S.,
Coelho, M. A., de Vienne, D. M., Hartmann, F. E., Begerow, D., Hood, M. E. &
Giraud, T. 2018. Multiple convergent supergene evolution events in mating-type
chromosomes. *Nature Communications* **9**: 2000.
Carpentier, F., de la Vega, R. R. C., Jay, P., Duhamel, M., Shykoff, J. A., Perlin, M. H.,
Wallen, R. M., Hood, M. E. & Giraud, T. 2022. Tempo of degeneration across
independently evolved nonrecombining regions. *Molecular Biology and Evolution* **39**.
Charlesworth, D. 2017. Evolution of recombination rates between sex chromosomes.
Philosophical Transactions of the Royal Society B-Biological Sciences **372**.
Cingolani, P., Platts, A., Wang le, L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X. &
Ruden, D. M. 2012. A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster
strain w1118; iso-2; iso-3. *Fly (Austin)* **6**: 80-92.
Contamine, V., Lecellier, G., Belcour, L. & Picard, M. 1996. Premature death in Podospora
anserina: sporadic accumulation of the deleted mitochondrial genome, translational
parameters and innocuity of the mating types. *Genetics* **144**: 541-55.
De Filippo, E., Gautier, V., Lalanne, C., Levert, E., Chahine, E., Hartmann, F. E., Giraud, T.
& Silar, P. 2025. Huge genetic diversity in Schizothecium tetrasporum (Wint.). N.
Lundq.: delimitation of 18 sibling species and three clades through genome
sequencing. *BioRxiv*
https://www.biorxiv.org/content/10.1101/2025.03.27.645579v1.
Duhamel, M., Carpentier, F., Begerow, D., Hood, M. E., Rodriguez de la Vega, R. C. &
Giraud, T. 2022. Onset and stepwise extensions of recombination suppression are
common in mating-type chromosomes of *Microbotryum* anther-smut fungi. *Journal of*
Evolutionary Biology **35**: 1619-1634.
Duhamel, M., Hood, M. E., Rodriguez de la Vega, R. C. & Giraud, T. 2023. Dynamics of
transposable element accumulation in the non-recombining regions of mating-type
chromosomes in anther-smut fungi. *Nature Communications* **14**: 5692
Felsenstein, J. 1974. The evolutionary advantage of recombination. *Genetics* **78**: 737-756.
Fisher, R. A. 1930. *The genetical theory of natural selection*.
Fontanillas, E., Hood, M. E., Badouin, H., Petit, E., Barbe, V., Gouzy, J., De Vienne, D. M.,
Aguileta, G., Poulain, J., Wincker, P., Chen, Z., Toh, S. S., Cuomo, C. A., Perlin, M.
H., Gladieux, P. & Giraud, T. 2015. Degeneration of the nonrecombining regions in
the mating-type chromosomes of the anther-smut fungi. *Molecular Biology and*
Evolution **32**: 928-943.
Gemmell, N. J. & Slate, J. 2006. Heterozygote advantage for fecundity. *PLoS One* **1**: e125.
Giraud, T., Yockteng, R., Lopez-Villavicencio, M., Refregier, G. & Hood, M. E. 2008. The
mating system of the anther smut fungus, *Microbotryum violaceum*: selfing under
heterothallism. *Euk. Cell* **7**: 765-775.

Glémin, S., Bataillon, T., Ronfort, J., Mignot, A. & Olivieri, I. 2001. Inbreeding depression in
783  small populations of self-incompatible plants. *Genetics* **159**: 1217-29.
784 Grognet, P., Bidard, F., Kuchly, C., Tong, L. C. H., Coppin, E., Benkhali, J. A., Couloux, A.,
785  Wincker, P., Debuchy, R. & Silar, P. 2014. Maintaining two mating types: structure of
786  the mating type locus and its role in heterokaryosis in *Podospora anserina*. *Genetics*
787  **197**: 421-432.
788 Hartmann, F. E., Ament-Velasquez, S. L., Vogan, A. A., Gautier, V., Le Prieur, S.,
789  Berramdane, M., Snirc, A., Johannesson, H., Grognet, P., Malagnac, F., Silar, P. &
790  Giraud, T. 2021a. Size variation of the nonrecombining region on the mating-type
791  chromosomes in the fungal *Podospora anserina* species complex. *Molecular Biology*
792  *and Evolution* **38**: 2475-2492.
793 Hartmann, F. E., Duhamel, M., Carpentier, F., Hood, M. E., Foulongne-Oriol, M., Silar, P.,
794  Malagnac, F., Grognet, P. & Giraud, T. 2021b. Recombination suppression and
795  evolutionary strata around mating-type loci in fungi: documenting patterns and
796  understanding evolutionary and mechanistic causes. *New Phytologist* **229**: 2470-
797  2491.
798 Hill, J., Enbody, E. D., Bi, H., Lamichhaney, S., Lei, W., Chen, J., Wei, C., Liu, Y.,
799  Schwochow, D., Younis, S., Widemo, F. & Andersson, L. 2023. Low mutation load in
800  a supergene underpinning alternative male mating strategies in Ruff *(Calidris*
801  *pugnax).* **40**.
802 Hill, W. G. & Robertson, A. 1966. The effect of linkage on limits to artificial selection.
803  *Genetics Research* **8**: 269-294.
804 Hood, M. E. & Antonovics, J. 2000. Intratetrad mating, heterozygosity, and the maintenance
805  of deleterious alleles in *Microbotryum violaceum (= Ustilago violacea)*. *Heredity* **85**:
806  231-241.
807 Hough, J., Hollister, J. D., Wang, W., Barrett, S. C. H. & Wright, S. I. 2014. Genetic
808  degeneration of old and young y chromosomes in the flowering plant *Rumex*
809  *hastatulus. Proceedings of the National Academy of Sciences of the United States of*
810  *America* **111**: 7713-7718.
811 Hyde, K., Maharachchikumbura, S., Hongsanan, S., Samarakoon, M., Lucking, R., Pem, D.,
812  Harishchandra, D., Jeewon, R., Zhao, R.-L., Xu, J.-C., Liu, J.-K., Al-Sadi, A. M.,
813  Bahkali, A. H. & Elgorban, A. M. 2017. The ranking of fungi: a tribute to David L.
814  Hawksworth on his 70th birthday. *Fungal Divers.* **84**: 1–23.
815 Jay, P., Chouteau, M., Whibley, A., Bastide, H., Parrinello, H., Llaurens, V. & Joron, M.
816  2021. Mutation load at a mimicry supergene sheds new light on the evolution of
817  inversion polymorphisms. *Nature Genetics* **53**: 288-293.
818 Jay, P., Jeffries, D., Hartmann, F. E., Véber, A. & Giraud, T. 2024. Why do sex
819  chromosomes progressively lose recombination? *Trends in Genetics* **40**: 564-579.
820 Judelson, H. S., Spielman, L. J. & Shattock, R. C. 1995. Genetic mapping and non-
821  Mendelian segregation of mating type loci in the oomycete, Phytophthora infestans.
822  *Genetics* **141**: 503-12.
823 Kasuga, T., White, T. J. & Taylor, J. W. 2002. Estimation of Nucleotide Substitution Rates in
824  Eurotiomycete Fungi. *Molecular Biology and Evolution* **19**: 2318-2324.
825 Langmead, B. & Salzberg, S. L. 2012. Fast gapped-read alignment with Bowtie 2. *Nature*
826  *Methods* **9**: 357-359.
827 Le Veve, A., Genete, M., Lepers-Blassiau, C., Ponitzki, C., Poux, C., Vekemans, X., Durand,
828  E. & Castric, V. (2024) The genetic architecture of the load linked to dominant and
829  recessive self-incompatibility alleles in Arabidopsis halleri and A. lyrata. pp. eLife
830  Sciences Publications, Ltd.
831 Llaurens, V., Gonthier, L. & Billiard, S. 2009. The sheltered genetic load linked to the S locus
832  in plants: new insights from theoretical and empirical approaches in sporophytic self-
833  incompatibility. *Genetics* **183**: 1105-1118.
834 Ma, W.-J. & Veltsos, P. 2021. The diversity and evolution of sex chromosomes in frogs.
835  *Genes* **12**: 483.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M. & DePristo, M. A. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* **20**: 1297-303.

Mena-Alí, J. I., Keser, L. H. & Stephenson, A. G. 2009. The effect of sheltered load on reproduction in Solanum carolinense, a species with variable self-incompatibility. *Sex Plant Reprod* **22**: 63-71.

Menkis, A., Jacobson, D. J., Gustafsson, T. & Johannesson, H. 2008. The mating-type chromosome in the filamentous ascomycete *Neurospora tetrasperma* represents a model for early evolution of sex chromosomes. *PLoS Genet.* **4**: e1000030.

Meunier, C., Darolti, I., Reimegård, J., Mank, J. E. & Johannesson, H. 2022. Nuclear-specific gene expression in heterokaryons of the filamentous ascomycete Neurospora tetrasperma. *Proc Biol Sci* **289**: 20220971.

Meunier, C., Hosseini, S., Heidari, N., Maryush, Z. & Johannesson, H. 2018. Multilevel selection in the filamentous ascomycete *Neurospora tetrasperma*. *The American Naturalist* **191**: 290-305.

Muller, H. J. 1932. Some genetic aspects of sex. *The American Naturalist* **66**: 118-138.

Nicolas, M., Marais, G. A. B., Hykelova, V., Janousek, B., Laporte, V., Vyskot, B., Mouchiroud, D., Negrutiu, I., Charlesworth, D. & Monéger, F. 2004. A gradual process of recombination restriction in the evolutionary history of the sex chromosomes in dioecious plants. *PLoS Biology* **3**: e4.

Oudemans, P. V., Alexander, H. M., Antonovics, J., Altizer, S., Thrall, P. H. & Rose, L. 1998. The distribution of mating-type bias in natural populations of the anther-smut Ustilago violacea on *Silene alba* in Virginia. *Mycologia* **90**: 372-381.

R Core Team 2023. R: A Language and Environment for Statistical Computing. . *R Foundation for Statistical Computing, Vienna, Austria*.

Raju, N. B. & Perkins, D. D. 1994. Diverse programs of ascus development in pseudohomothallic species of Neurospora, Gelasinospora, and Podospora. *Developmental Genetics* **15**: 104-118.

Rice, W. R. & Chippindale, A. K. 2001. Sexual recombination and the power of natural selection. *Science* **294**: 555-559.

Rodrigues, N., Studer, T., Dufresnes, C. & Perrin, N. 2018. Sex-Chromosome Recombination in Common Frogs Brings Water to the Fountain-of-Youth. *Molecular Biology and Evolution* **35**: 942-948.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P. & Cardona, A. 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**: 676-682.

Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* **9**: 671-675.

Schwander, T., Libbrecht, R. & Keller, L. 2014. Supergenes and complex phenotypes. *Current Biology* **24**: R288-R294.

Shen, L., Chapeland-Leclerc, F., Ruprich-Robert, G., Chen, Q., Chen, S., Adnan, M., Wang, J., Liu, G. & Xie, N. 2022. Involvement of VIVID in white light-responsive pigmentation, sexual development and sterigmatocystin biosynthesis in the filamentous fungus Podospora anserina. *Environ Microbiol* **24**: 2907-2923.

Silar, P. 2020. *Podospora anserina*. hal-02475488.

Silar, P., Gautier, V., Lalanne, C., Tangthirasunun, N., Arthur, M., Hartmann, F. E. & Giraud, T. 2024. The Podospora anserina species complex in metropolitan and overseas France with description of a new species, Podospora reunionensis sp. nov. *submitted*.

Smith, J. M. & Haigh, J. 2009. The hitch-hiking effect of a favourable gene. *Genetical Research* **23**: 23-35.

Stone, J. 2004. Sheltered load associated with S-alleles in Solanum carolinense. *Heredity* **92**: 335-342.

Tezenas, E., Giraud, T., Véber, A. & Billiard, S. 2023. The fate of recessive deleterious or
overdominant mutations near mating-type loci under partial selfing. *Peer Community*
Journal **3**: e14.
Thomas, A., Shykoff, J., Jonot, O. & Giraud, T. 2003. Sex-ratio bias in populations of the
phytopathogenic fungus *Microbotryum violaceum* from several host species.
International Journal of Plant Sciences **164**: 641-647.
Uyenoyama, M. K. 2005. Evolution under tight linkage to mating type. *New Phytologist* **165**:
63-70.
Vittorelli, N., Rodríguez de la Vega, R. C., Snirc, A., Levert, E., Gautier, V., Lalanne, C., De
Filippo, E., Gladieux, P., Guillou, S., Zhang, Y., Tejomurthula, S., Grigoriev, I. V.,
Debuchy, R., Silar, P., Giraud, T. & Hartmann, F. E. 2023. Stepwise recombination
suppression around the mating-type locus in an ascomycete fungus with self-fertile
spores. *PLoS Genetics* **19**: e1010347.
Whittle, C. A., Sun, Y. & Johannesson, H. 2011. Degeneration in codon usage within the
region of suppressed recombination in the mating-type chromosomes of *Neurospora*
tetrasperma. *Eukaryotic Cell* **10**: 594-603.
Wielstra, B. 2020. Balanced lethal systems. *Current Biology* **30**: R742-R743.
Wright, A. E., Dean, R., Zimmer, F. & Mank, J. E. 2016. How to make a sex chromosome.
Nature Communications **7**: 12087.
Zakharov, I. A. 2005. [Intratetrad mating and its genetic and evolutionary consequences].
Genetika **41**: 508-19.
Zakharov, I. A. 2023. Intratetrad mating as the driving force behind the formation of sex
chromosomes in fungi. *Trends in Genetics and Evolution* **6**: doi:
10.24294/tge.v6i1.252.
915

918   **Table 1:** Number of strains from *Schizothecium tetrasporum, Schizothecium*

919   *tritetrasporum* and *Podospora anserina* used for experiments, summary of the conditions

920   (light and temperature) and measure time points for which fitness proxies (growth speed or

921   germination dynamics) were assessed according to species and strains. Information on strains

922   can be found in Table S1.

Growth speed				
Species	**Number of strains**	**Measure time points (days)**	**Temperature**	**Light exposure**
S. tetrasporum	10	6 / 14	22°C	Yes
S. tetrasporum	10	6 / 14	22°C	No
S. tritetrasporum	13	6 / 14	22°C	Yes
P. anserina	17	2 / 6	18°C	Yes
P. anserina	17	2 / 6	37°C	Yes
Germination dynamics				
Species	**Number of strains**	**Measurement time points (days)**		
S. tetrasporum	8	2 / 3 / 7		

Figure legends

Figure 1: Fitness experiments to test for the presence of a sheltered load in pseudo-

homothallic ascomycete fungi. A) Schema illustrating how the life cycle of pseudo-

homothallic fungi can be used to test for the existence of sheltered load (i.e. the presence of

heterozygous recessive deleterious mutations) in the non-recombining region around the

mating-type locus. Pseudo-homothallic ascomycete fungi produce heterokaryotic spores

(with nuclei of opposite mating types), and with therefore a main phase in their life cycle that

is heterokaryotic, i.e. diploid-like. **B)** Pictures of mycelium after six days of growth on Petri

dishes under light conditions at 22°C, with heterokaryons at left, *MAT1-2* homokaryons in the

middle and *MAT1-1* homokaryons at right, for *Schizothecium tritetrasporum* (strain

PSN935D), *Schizothecium tetrasporum* (strain PSN781A) and *Podospora anserina* (strain

Wa68), from top to bottom. Scale bar: 4 cm. **C)** Picture of *Schizothecium tetrasporum* spores

(strain CBS815.71)*,* with two small spores on the top right and two large spores on the

bottom left. The large spores have length and width of ca. 25 x 12 μm.

Figure 2: Genomic analyses of heterozygosity and mating-type specific putative

deleterious mutations (non-synonymous SNPs) along the chromosomes of the

***Schizothecium tritetrasporum* PSN829 strain*,* as an illustration of the situation with**

genome-wide homozygosity except a few heterozygous autosomal regions in addition to

the flanking region of the mating-type locus. A) Genome-wide per-gene density in

heterozygous SNPs. Each dot represents one gene, the Y axis corresponding to the mean

proportion of heterozygous SNPs per base pair for each gene, and the X axis to the genomic

position along the CBS815.71-sp3 assembly in megabases (Mb). The seven longest contigs in

the assembly are represented, by length order. B) Genome-wide per-gene density in

heterozygous non-synonymous SNPs. Each dot represents one gene, the Y axis

corresponding to the mean proportion of heterozygous non-synonymous SNPs per base pair;

these SNPs introduced either a missense mutation (i.e., change in amino-acid; blue dot), the

gain of a stop codon (pink dot) or the loss of a start codon (orange dot) compared to the

reference genome CBS815.71-sp3. The X axis corresponds to the genomic position along the

CBS815.71-sp3 assembly in megabases (Mb). Dots indicating a null value have been

removed for clarity. The mating-type locus (MAT) position is symbolized by a vertical red

line.

Figure 3: Mycelium growth speed of heterokaryons versus homokaryons in

***Schizothecium tritetrasporum.* A)** Mycelium growth speed (mm.day⁻¹) of heterokaryons (in

purple), *MAT1-1* homokaryons (in red) and *MAT1-2* homokaryons (in blue), of the three

replicates for each of two batches, for the 13 strains of *S. tritetrasporum,* between day 6 and

day 14, grown under light at 22°C. Boxplot elements: central line: median, box limits: 25th

and 75th percentiles, whiskers: 1.5× interquartile range. **B)** Model estimates of mycelium

growth speed for heterokaryons (in purple) and homokaryons (in grey) in *S. tritetrasporum.*

Dots and bars represent marginal means and standard errors estimated from a linear mixed-

effect model explaining growth speed using ploidy (i.e. heterokaryon or homokaryon) as a

fixed effect, and strain ID and batch as random effects. **C)** Model estimates of mycelium

growth speed for heterokaryons (in purple), *MAT1-1* homokaryons (in red) and *MAT1-2*

homokaryons (in blue) in *S. tritetrasporum.* Dots and bars represent marginal means and

standard errors estimated from a linear mixed-effect model explaining growth speed using the

nuclear status (i.e. *MAT1-1* homokaryon, *MAT1-2* homokaryon or heterokaryon) as a fixed

effect, and strain ID and batch as random effects.

P-values correspond to pairwise t-tests adjusted for multiple testing using the Holm method

(ns: non-significant; **: p<0.01; ***: p<0.001).

Figure 4: Mycelium growth speed of heterokaryons versus homokaryons in

***Schizothecium tetrasporum*. A)** Mycelium growth speed (mm.day⁻¹) of heterokaryons (in

purple), *MAT1-1* homokaryons (in red) and *MAT1-2* homokaryons (in blue), of the three

replicates of the 10 *S. tetrasporum* strains between day 6 and day 14, grown under light or

dark conditions at 22°C. Boxplot elements: central line: median, box limits: 25th and 75th

percentiles, whiskers: 1.5× interquartile range. **B)** Model estimates of mycelium growth speed

for heterokaryons (in purple) and homokaryons (in grey) in *S. tetrasporum* under light or dark

conditions. Dots and bars represent marginal means and standard errors estimated from a

linear mixed-effect model explaining growth speed using ploidy (i.e. homokaryon or

heterokaryon) and condition (i.e. light or dark) as fixed effects, with their interaction, and

strain ID as a random effect. **C)** Model estimates of mycelium growth speed for

heterokaryons (in purple), *MAT1-1* homokaryons (in red) and *MAT1-2* homokaryons (in blue)

in *S. tetrasporum* under light or dark conditions. Dots and bars represent marginal means and

standard errors estimated from a linear mixed-effect model explaining growth speed using

nuclear status (i.e. *MAT1-1* homokaryon, *MAT1-2* homokaryon or heterokaryon) and

condition (i.e. light or dark) as fixed effects, with their interaction, and strain ID as a random

effect.

P-values correspond to pairwise t-tests adjusted for multiple testing using the Holm method

(ns: non-significant; *: $p < 0.05$).

Figure 5: Mycelium growth speed of heterokaryons versus homokaryons in *Podospora*

***anserina*. A)** Mycelium growth speed (mm.day⁻¹) of heterokaryons (in purple), *MAT1-1*

homokaryons (in red) and *MAT1-2* homokaryons (in blue), of the three replicates of the 17 *P.*

anserina strains between day 2 and day 6, grown at 18°C or 37°C under light conditions.

Boxplot elements: central line: median, box limits: 25th and 75th percentiles, whiskers: $1.5\times$

interquartile range. **B)** Model estimates of mycelium growth speed for heterokaryons (in

purple) and homokaryons (in grey) in *P. anserina* at 18°C and 37°C. Dots and bars represent

marginal means and standard error estimated from a linear mixed-effect model explaining

growth speed using ploidy (i.e. homokaryon or heterokaryon) and condition (i.e. 18°C or

37°C) as fixed effects, with their interaction, and strain ID as a random effect.

C) Model estimates of mycelium growth speed of heterokaryons (in purple), *MAT1-1*

homokaryons (in red) and *MAT1-2* homokaryons (in blue) in *P. anserina* at 18°C and 37°C.

Dots and bars represent marginal means and standard errors estimated from a linear mixed-

effect model explaining growth speed using nuclear status (i.e. *MAT1-1* homokaryon, *MAT1-*

2 homokaryon or heterokaryon) and condition (i.e. 18°C or 37°C) as fixed effects, with their

interaction, and strain ID as a random effect.

P-values correspond to pairwise t-tests adjusted for multiple testing using the Holm method

(ns: non-significant; *: $p<0.05$).

1013

Figure 6: Spore germination dynamics in *Schizothecium tetrasporum,* depending on

their heterokaryotic versus homokaryotic nuclear status, as inferred from their size.

A) Germination rate on days 2, 3 and 7 for small spores (in purple) and large spores (in grey)

in the 10 spores for each of two batches of the eight *S. tetrasporum* strains. Symbol shapes

correspond to batches. Boxplot elements: central line: median, box limits: 25th and 75th

percentiles, whiskers: $1.5\times$ interquartile range. **B)** Model predictions of germination rate

dynamics for small spores (in purple) and large spores (in grey) in *S. tetrasporum*. Curves

and shaded error bands represent model prediction and standard errors from the fit of the

germination dynamics using a generalised linear mixed-effect model with a Binomial

distribution modelling germination rate as a linear combination of day (as a continuous

predictor) and spore size (i.e. large or small) treated as fixed effects, and strain ID and batch

as random effects (pairwise z-test between the two spore size levels: p=0.4459, non-

significant).

1027

1028

1029   **Supplementary Figure legends**

1030

Supplementary Figure 1: Genomic analyses of heterozygosity and mating-type specific

putative deleterious mutations (non-synonymous SNPs) along the chromosomes of four

strains of *Schizothecium tetrasporum*. The left column depicts the genome-wide per-gene

density in heterozygous SNPs for the four *S. tetrasporum* strains for which genomes were

available. Each dot represents one gene, the Y axis corresponding to the mean proportion of

heterozygous SNPs per base pair for each gene and the X axis to the genomic position along

the CBS815.71-sp3 assembly in megabases (Mb). The seven longest contigs in the assembly

are represented, by length order. The right column depicts the genome-wide per-gene density

in heterozygous non-synonymous SNPs. Each dot represents one gene, the Y axis

corresponding to the mean proportion of heterozygous non-synonymous SNPs per base pair;

these SNPs introduced either a missense mutation (i.e., a change in amino-acid; blue dot) or

the gain of a stop codon (pink dot) compared to the reference genome CBS815.71-sp3. The X

axis corresponds to the genomic position along the CBS815.71-sp3 assembly in megabases

(Mb). Dots indicating a null value have been removed for clarity. The mating-type locus

(MAT) position is indicated by a vertical red line.

1046

Supplementary Figure 2: Genomic analyses of heterozygosity and mating-type specific

possible deleterious mutations (non-synonymous SNPs) along the chromosomes of nine

strains of *Schizothecium tritetrasporum*. The left column depicts the genome-wide per-gene

density in heterozygous SNPs for the nine *S. tritetrasporum* strains for which genomes were

available. Each dot represents one gene, the Y axis corresponding to the mean proportion of

heterozygous SNPs per base pair for each gene and the X axis to the genomic position along

the CBS815.71-sp3 assembly in megabases (Mb). The seven longest contigs in the assembly

are represented, by length order. The right column depicts the genome-wide per-gene density

in heterozygous non-synonymous SNPs. Each dot represents one gene, the Y axis

corresponding to the mean proportion of heterozygous non-synonymous SNPs per base pair;

these SNPs introduced either a missense mutation (i.e., a change in amino-acid; blue dot), the

gain of a stop codon (pink dot) or the loss of a start codon (orange dot) compared to the

reference genome CBS815.71-sp3. The X axis corresponds to the genomic position along the

CBS815.71-sp3 assembly in megabases (Mb). Dots indicating a null value have been

removed for clarity. The mating-type locus (MAT) position is indicated by a vertical red line.

1062

Supplementary Figure S3: Mycelium growth speed of heterokaryons versus

homokaryons for 13 strains of *Schizothecium tritetrasporum*.

Mycelium growth speed (mm.day^{-1}) of heterokaryons (in purple), *MAT1-1* homokaryons (in

red) and *MAT1-2* homokaryons (in blue), of the three replicates for each of two batches, for

the 13 *S. tritetrasporum* strains, between day 6 and day 14, grown under light conditions at

22°C. Green dots highlight the heterokaryon replicates for which fewer than 50% of the nine

mycelium plugs from their Petri dish were genotyped heterokaryotic. Boxplot elements:

central line: median, box limits: 25th and 75th percentiles, whiskers: 1.5× interquartile range.

1071

Supplementary Figure S4: Nuclear loss in *Schizothecium tritetrasporum* heterokaryotic

mycelium.

A) Proportions of plugs with only *MAT1-1* (in red), only *MAT1-2* (in blue) or both MAT (in

purple) detected in *S. tritetrasporum* heterokaryotic mycelium after 14 days of growth.

Proportions were calculated out of nine plugs per plate that were genotyped by PCR in three

replicates per strain for each of two batches. Symbol shapes correspond to replicate plates.

Boxplot elements: central line: median, box limits: 25th and 75th percentiles, whiskers: 1.5×

interquartile range. B) Model estimates of proportions of plug with only the *MAT1-1* or the

MAT1-2 allele detected by PCR in *S. tritetrasporum* heterokaryotic mycelium. Dots and bars

represent marginal means and standard error estimated from a generalized linear mixed-effect

model with a Binomial distribution modelling plug proportion with nuclear loss using the

genotype of the plug after the loss (i.e. *MAT1-1* or *MAT1-2*) as a fixed effect and strain ID

and batch as random effects. The p-value corresponds to a pairwise t-test (ns: non-

significant).

1086

Supplementary Figure S5: Distribution of nuclei in vegetative hyphae of the *Podospora*

***anserina* S strain.** Epifluorescence microscopy pictures of vegetative hyphae of the

heterokaryon, the *MAT1-1* homokaryon and the *MAT1-2* homokaryon of the *P. anserina* S

strain. Left column: DAPI stained nuclei, images correspond to z-projections of three z-slices

separated by 1 µm. Right column: bright-field images corresponding to the slice in the middle

of the z-stack. Arrow heads: septa. Scale bar: 10 µm.

1093

Supplementary Figure S6: Number of nuclei and their density per article in

heterokaryotic versus homokaryotic mycelia in the *Podospora anserina* S strain.

A) Number of nuclei per article in heterokaryons (in purple), *MAT1-1* homokaryons (in red)

and *MAT1-2* homokaryons (in blue), in 30 articles of three replicate Petri dishes of the *P.*

anserina S strain at day 6 of growth, at 18°C under light. Boxplot elements: central line:

median, box limits: 25th and 75th percentiles, whiskers: 1.5× interquartile range.

B) Model estimates of nucleus number per article in heterokaryons (in purple), *MAT1-1*

homokaryons (in red) and *MAT1-2* homokaryons (in blue) of the *P. anserina* S strain. Dots

and bars represent marginal means and standard errors estimated from a generalised linear

model with a Poisson distribution explaining nucleus number per article using nuclear status

(i.e. *MAT1-1* homokaryon, *MAT1-2* homokaryon or heterokaryon) as fixed effects. P-values

correspond to pairwise z-tests adjusted for multiple testing using the Holm method (ns: non-

significant). **C)** Density of nuclei per article (nb of nuclei.μ^{-1}) in heterokaryons (in purple),

MAT1-1 homokaryons (in red) and *MAT1-2* homokaryons (in blue), in 30 articles of three

replicate Petri dishes of the *P. anserina* S strain at day 6 of growth, at 18°C under light.

Boxplot elements: central line: median, box limits: 25th and 75th percentiles, whiskers: 1.5×

interquartile range. **D)** Model estimates of the density of nuclei per article in heterokaryons

(in purple), *MAT1-1* homokaryons (in red) and *MAT1-2* homokaryons (in blue) of the *P.*

anserina S strain. Dots and bars represent marginal means and standard errors estimated from

a linear model explaining nucleus density per article using nuclear status (i.e. *MAT1-1*

homokaryon, *MAT1-2* homokaryon or heterokaryon) as fixed effects. P-values correspond to

pairwise F-tests adjusted for multiple testing using the Holm method (ns: non-significant).

1116

1117

1118   **Figure 1**

1119

1120

1121

1122

1123   **Figure 2**

1124

1125

1126

1127

Figure 3

1128

1129

1130

1131

1132

1133 **Figure 4**

1134

1135

1136

Figure 5

1137

1138

1139

1140

1141

1142

Figure 6

1143

1144

1145

1146   **Figure S1**

1148   **Figure S2**

1149

1150
1151 **Figure S3**

1152

1153

1154
1155   **Figure S4**

1156

1157

1158

1159

1160

1161

1162

1163

1164 **Figure S5**

1165

1166   **Figure S6**

1167