# High-coverage, massively parallel sequencing of single-cell genomes with CAP-seq

Mengzhen Li[a,#], Xuanpei Zhai[a,#], Jie Li[a,#], Shiyan Li[b,f], Yanan Du[a], Jian Zhang[a], Rong Zhang[a], Yuan Luo[c], Wu Wei[b,*] and Yifan Liu[a,d,e,*]

[a] *School of Physical Science and Technology, ShanghaiTech University, Shanghai 201210, China.*

[b] *Lingang Laboratory, Shanghai 200031, China.*

[c] *State Key Laboratory of Transducer Technology, Shanghai Institute of Microsystem and Information Technology, Chinese Academy of Sciences, Shanghai 200050, China.*

[d] *State Key Laboratory of Advanced Medical Materials and Devices, ShanghaiTech University, Shanghai 201210, China.*

[e] *Shanghai Clinical Research and Trial Center, Shanghai 201210, China.*

[f] *CAS Key Laboratory of Computational Biology, Shanghai Institute of Nutrition and Health, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai, 200031, China.*

* Corresponding author. Email: wuwei@lglab.ac.cn (W.W.), liuyf6@shanghaitech.edu.cn (Y.L.)

# These authors contribute equally to this work.

**Keywords:** single-cell, whole genome sequencing, microbe, microfluidics, metagenomics

**ABSTRACT**

Microorganisms play essential roles in ecosystems such as the Earth and human bodies. Studying these organisms relies on their genome information, but obtaining high-quality genomes has long been challenging due to technical limitations in genomic sequencing. Traditional genome sequencing methods are limited by the need to culture isolated strains, excluding unculturable taxa and restricting the study of complex communities. Metagenomics bypasses this but lacks single-cell resolution and often misses rare, critical species. Droplet-based single-cell genomics offers high-throughput genome library preparation but faces challenges like requiring complex microfluidic setups and low genome coverage. To address these challenges, we introduce CAP-seq, a high-throughput single-cell genomic sequencing method with markedly improved genome coverage and ease of use. CAP-seq employs semi-permeable compartments that allow reagent exchange while retaining large DNA fragments, enabling efficient genome processing. This innovation results in higher-quality single amplified genomes (SAGs) and significantly improves resolution. In validation tests with simple and complex microbial communities, CAP-seq yielded high-quality SAGs with over 50% genome coverage, capturing rare taxa more accurately and providing detailed insights into strain-level variation. CAP-seq thus offers a scalable, high-resolution solution for microbial genomic analysis, overcoming the limitations of droplet-based single-cell genomics and enhancing the study of complex microbial ecosystems.

**MAIN**

Our planet hosts a vast array of organisms, with microorganisms (primarily bacteria, archaea, and fungi) dominating in both abundance and diversity[1-3]. Current estimates suggest the existence of up to a trillion microbial species, most of which remain uncharacterized[3-5]. This remarkable genomic diversity constitutes a rich repository of genetic material[6]. For decades, researchers have been sequencing microbial genomes to understand individual microbe and their communities[7,8] (e.g., human microbiota), while leveraging the resulting sequence libraries[9] to discover novel drug candidates[10,11], molecular tools[12], and biosynthetic pathways[13,14].

Classical genome sequencing of microorganisms relies on pure culture of isolated strains to amplify their DNA[15], but this method excludes unculturable taxa, which can occupy a significant portion of a given community[3]. It is also labor-intensive, limiting its throughput and suitability for profiling complex ecosystems. To overcome these issues, cultivation-free methods like metagenomics have been widely adopted[16,17]. However, these techniques compromise single-cell or strain-level resolution, depending on *in-silico* binning to assemble metagenomic "genomes"[18-20]. In addition, these methods are often biased towards abundant groups during the library preparation[21,22] and binning steps[23], missing low-abundance taxa that are critical to community function[24,25].

To tackle the challenges in community-wide approaches, efforts have been devoted to single-cell genomics[26]. This technique isolates individual cells and then prepares their genomes independently through processes such as cell sorting, lysis, and whole-genome amplification[26]. While this approach offers valuable insights, it remains limited by resource availability, resulting in low throughput. Recently, the workflow has been implemented in droplet microfluidics to enable thousands of single-cell genome libraries to be prepared simultaneously[27,28]. However, this requires multiple (e.g., five) advanced microfluidic modules and experimentations, posing significant technical challenges for non-experts and thus limiting broad use. In addition, droplets act as isolated environments, preventing purification and buffer exchange[29]. It is thus difficult to optimize sequential reactions, such as lysis, genome amplification, fragmentation, and

indexing, in the same droplet. As a result, the quality of single amplified genomes (SAGs) is often low, with typical genome coverage under 5%, restricting our understanding of microbial communities.

Here, we introduce single-cell genomic sequencing using compartments with adjusted permeability (CAP-seq), a high-throughput method that achieves high-coverage SAGs with only two microfluidic steps. As outlined in Fig. 1a, CAP-seq begins by encapsulating individual microbial cells into semi-permeable compartments, referred to as CAPs (the first microfluidic step). These compartments are designed with an engineered hydrogel shell that allows the selective transport of smaller molecules[30] (e.g., enzymes, dNTPs, buffers) while retaining large DNA fragments (>600 bp), Fig. 1b. This unique feature enables efficient washing and reagent exchange throughout the process. In the subsequent steps, various assays are conducted on the CAP-encapsulated cells, all in bulk solutions, including cell lysis, whole-genome amplification, and DNA fragmentation.

Once the genomic DNA is processed, the CAPs, now containing single-cell genomes, are co-encapsulated with DNA barcode beads in droplets (second microfluidic step). This step assigns each genome a unique cell barcode. Afterward, the CAPs are extracted from the droplets, washed, and dissolved to release the barcoded DNA fragments (~1 kb). These fragments are then further amplified and prepared for nanopore sequencing. Finally, the sequenced reads are categorized into individual SAGs based on their cell barcodes, yielding high-coverage genomes with significantly improved throughput and resolution. CAP-seq thus overcomes limitations in droplet single-cell genomics, providing a streamlined, scalable approach for studying microbial communities with enhanced genome recovery.

We validated CAP-seq using a 1:1 mixture of *Escherichia coli* and *Bacillus subtilis* cells. In a representative experiment, around 30,000 cells were processed with the CAP-seq workflow. The resulting library was sequenced by the Oxford nanopore sequencer, yielding 84.5 million raw reads. Quality control and our in-house binning pipeline identified 48.5 million reads with full-length and correct cell barcodes, which were sorted

into 5,484 groups of ≥1,000 reads, including 1,607 groups with ≥10,000 reads (Fig. 1c). After subsequent barcode trimming, the read lengths ranged from 300 to 1,500 nt, with an average length of 898 nt (Fig. 1d). Organism-specific mapping (Fig. 1e) revealed that 56.4% of barcode groups were assigned to *E. coli* (98.8% reads on average mapped to the *E. coli* genome), and 39.1% were mapped to *B. subtilis* (average of 98.5% mapped reads). A small fraction (4.5%) of groups were mixed. These results validate the CAP-seq workflow, demonstrating its capability for high-throughput, long-read sequencing of single microbial cells.

Next, we tested CAP-seq with a more complex mock microbial community composed of four bacterial species with varied Gram polarities at uneven ratios (Fig. 2a). Sequencing around 30,000 cells and bioinformatic binning resulted in 3,490 barcode groups containing over 1,000 reads, 91% of which had a purity (percent reads mapped to a dominant genome) of over 95%, and 82.6% with over 99% purity (Fig. 2b). CAP-seq's species profile closely matched the community composition, capturing rare species more accurately than metagenomic sequencing, which was biased towards the more abundant species (Fig. 2c). Specifically, *S. aureus* and *K. pneumoniae*, which accounted for only 0.3% of the metagenomic reads, were accurately represented by CAP-seq, with 173 (5.4%) and 13 (0.4%) SAGs assigned to each species, respectively. These results highlight CAP-seq's ability to provide an unbiased microbial population profile, capturing rare taxa with single-cell accuracy, making it more effective for profiling complex microbial communities.

After mapping them to reference genomes, we calculated the genome completeness of 3,177 high-quality SAGs. Since genome completeness is depth-dependent, we plotted coverage versus sequencing depth (Fig. 2d). At 10X depth, most SAGs achieved over 50% genome coverage, regardless of Gram status. This demonstrates CAP-seq's ability to generate high-completeness SAGs with moderate sequencing efforts. The improved coverage is likely due to the CAPs' semi-permeable nature, enabling efficient washing, buffer exchange, and sustained reagent supply, which enhance key biochemical steps, especially whole-genome amplification, during library preparation.

We further examined the *E. coli* groups, which contained two distinct strains: BL21 and MACH1 (Fig. 2a). These strains were engineered to carry specific plasmid-encoded genes: BL21 harbored AmpR, while MACH1 carried sfGFP and CmR. We profiled these genes across 3,210 *E. coli* SAGs and observed distinct patterns in their relative abundance (Fig. 2e). The abundance of sfGFP and CmR were highly correlated, being prevalent in SAGs where AmpR was absent. In the remaining SAGs, only AmpR was detected, with neither sfGFP nor CmR present. These distinct gene abundance patterns enabled us to distinguish between the two strains. As a result, 10 SAGs were assigned to MACH1 and the rest to BL21. These findings confirm CAP-seq's ability to resolve intraspecific heterogeneity with single-cell resolution, enabling new avenues of research, such as tracking mobile genetic elements across individual cells and mining genes directly from SAGs.

In conclusion, CAP-seq enables the massively parallel sequencing of single-cell genomes. Compared to existing single-cell genomic methods, CAP-seq offers significantly improved coverage of SAGs and greater ease of use. We expect that CAP-seq will have broad applications across various fields related to microorganisms, paving the way for microbial single-cell genomics and genome-resolved metagenomics.

## METHODS

### Device design and fabrication

Microfluidic devices were designed using AutoCAD (Autodesk) and fabricated using polydimethylsiloxane (PDMS)-based soft lithography as previously described. Before use, the devices were treated with Aquapel (PPG Industries) to render the channel surfaces hydrophobic.

### Generation and characterization of CAPs

The CAPs were generated using a flow-focusing microfluidic chip (Supplementary Fig. and Supplementary Movie 1) by emulsifying an aqueous two-phase system (ATPS) consisting of dextran and acrylate-modified poly(ethylene glycol) diacrylate (PEGDA).

Upon emulsification, phase separation within the droplets resulted in a capsular configuration, with PEGDA forming the outer layer. The CAPs were collected in a tube and irradiated under a UV lamp (200 W) for 4 min to gel the outer PEGDA phase. The composition of the ATPS is provided in Supplementary Table X, and the details of its preparation are described elsewhere. The applied flow rates were provided in Supplementary Table X.

The semi-permeability of the CAPs was characterized using a customized PCR within the CAPs, where we amplified DNA fragments of varying lengths from encapsulated *E. coli* genomic DNA. This experiment determined the threshold size of dsDNA products that can be trapped in the CAPs. Detailed procedures for the characterization can be found in our previous reports.

**Cell culture**

*E. coli* (strains BL21 and MACH1), *B. subtilis* (strain ster164), *S. aureus* (strain RN4220), and *K. pneumoniae* (strain KP_1.6366) were separately cultured in 5 mL of LB medium at 37°C overnight in a shaking incubator. The cell concentration was determined by manually counting serial dilutions of the liquid culture on plastic slides (ThermoFisher Scientific, catalog no. C10228) using a microscope.

**Stool sample**

Stool samples were collected at the initial diagnosis of *Clostridioides difficile* infection (CDI), before treatment with antibiotics (vancomycin) or fecal microbiota transplantation (FMT), and at weeks 1, 2, 3, and 4 post-treatment. All samples were collected at the Department of Gastroenterology, Hepatology and Nutrition, Shanghai Children's Hospital, and stored at 4 °C for immediate usage. For long-term storage, the samples were mixed with 25% v/v glycerol and stored at -80°C.

To extract the cells, the samples were resuspended in PBS containing 0.1% Pluronic F68 (ThermoFisher, 24040032) by vigorous vortexing. Large debris was then removed by centrifugation at 4°C at 35g for 20 minutes. The cell-containing supernatant was transferred to a new tube, washed twice with PBS + 1% Pluronic F68, and resuspended

in 100 µL of PBS + 1% Pluronic F68. Cell concentration was determined by manual counting.

**CAP-seq procedure**

**Cell encapsulation.** Cell samples were resuspended in PBS buffer and spiked in the Dextran-rich solution (the inner phase of CAP) to a final cell density of 600 CFU/µL. This led to a positive rate of ~5% to minimize doublets formation, according to the Poisson statistics. The CAPs were then generated using microfluidics and collected in a 1.5 mL centrifugal tube. The collected emulsion was irradiated to gel the outer PEGDA phase. The CAPs were then extracted from the oil by adding an equal volume of 20% Perfluoroctane (PFO) in HFE-7500 and washed three times with Dulbecco's PBS (DPBS) buffer containing 0.1% v/v Tween-20.

**Cell lysis.** To lyse the encapsulated cells, 500 µL of dense-packed CAPs were mixed with an equal volume of 20 mM TE, 100 mM NaCl, 10 mM EDTA, 1% Triton X-100, 1800 U lysozyme and 20 U lysostaphin. The mixture was incubated at 37 °C for 2 hours. The CAPs were then washed with PBS buffer containing 10 mM EDTA for three times and mixed with 500 µL of 100 mM NaCl, 10 mM EDTA, 1% SDS, and 1mg proteinase K. The mix was incubated at 55°C for 30 min. The CAPs were eventually washed with a washing buffer containing 10 mM HEPES, 2% Tween, 20 mM EDTA and 0.1 mM NaCl for three times and resuspended in a DPBS buffer containing 0.1% Tween-20.

**Whole genome amplification.** After lysis, 300 µL of dense-packed CAPs were mixed with an equal volume of MDA mix containing 2X Phi29 reaction buffer, 30 µL Phi29 polymerase, 60 µL Pyrophosphatase, 20 mM dNTP, 0.25% v/v Triton X-100 and 0.25% v/v Pluronic F68. The MDA reaction was incubated at 30°C for 2 hours and terminated at 65°C (10 min). The post-MDA CAPs were washed with the washing buffer thrice and resuspended in a DPBS buffer containing 0.1% Tween-20.

**Tagmentation.** 200 µL of the post-MDA CAPs were mixed with an equal volume of tagmentation solution containing 80 µL Tag buffer and 40 µL Tag Enzyme 50 (aliquot with nuclease-free water). The mix was incubated at 55°C for 10 min. Afterward, the

CAPs were washed with the washing buffer thrice and resuspended in a DPBS buffer containing 0.1% Tween-20.

**Genome Barcoding.** To barcode the tagmented genomes, the CAPs were co-encapsulated with barcode beads using a custom bead-pairing device. The barcode bead (DTT-dissolvable polyacrylamide hydrogel bead) carried ssDNA primers containing a unique barcode sequence and an adapter sequence designed to bind to the genome fragments. The bead fabrication process has been described elsewhere. The beads used in this study were custom-ordered from DP-Bio, China. After merging, a resulting droplet contained a CAP, a unique barcoding primer (a dissolved bead), and a linear amplification solution. For linear amplification, the droplets were collected in PCR tubes and thermal cycled with the following recipe: 72°C for 5 min, 98°C for 3 min, and 15 cycles of 98°C for 10s, 60°C for 30s and 72°C for 5 min. The applied flow rates and reagents in the barcoding step were detailed in xxxx.

**CAP dissolution.** After barcoding, the droplets were broken by adding an equal volume of 20% PFO in HFE-7500. The released CAPs were washed three times using TE buffer containing 0.1% Tween-20. Next, the CAPs were then dissolved by mixing 20 µL dense-packed CAPs (containing ~9,000 cells) and 40 µL 0.8 M NaOH and incubated at 55°C for 10 min. Subsequently, the resulting solution was neutralized by adding 40 µL 0.8 M CH$_3$COOH. The DNA products were then purified by magnetic beads and dissolved in 15 µL nuclease-free water.

**Bulk PCR and sequencing.** The barcoded genomic DNA library was further amplified by PCR. The PCR reaction contained 15 µL purified linear amplification product, 25 µL KAPA hot-start HiFi polymerase mix, and 5 µL each of the forward and reverse primers. The PCR condition was 98°C for 3 min, and 16 cycles of 98°C for 10s, 60°C for 30s and 72°C for 5 min. The PCR product was purified using Ampure beads, examined using a Bioanalyzer, and then used for Nanopore sequencing library preparation following the manufacturer's protocol (https://nanoporetech.com/products/prepare/dna-library-preparation). The final DNA library was sequenced on an Oxford Nanopore sequencer

(GridION MK1 with P2 solo) using a PromethION Flow Cell. All the primer sequences for CAP-seq are detailed in Supplementary Table X.

**Bioinformatics**

**Initial quality control.** Raw sequencing data in POD5 format were base-called using Dorado in sup mode to generate high-quality sequence reads. The resulting reads were filtered with NanoFilt to remove low-quality sequences (quality score < 15) and reads with lengths outside the range of 20 to 60,000 bases.

**Barcode identification and demultiplexing.** Cell barcodes were identified by locating a known adapter sequence (GTCTCGTGGGCTCGG) using the Python pairwise2.align.localms function from the Bio module, with alignment parameters set as pairwise2.align.localms(seq1, seq2, 2, -1, -1, -1). Matches with an alignment score of 26 or higher were considered successful, and the adapter position was used to further locate the associated barcode sequence. Extracted barcodes were validated against a predefined whitelist of expected sequences. Reads with valid barcodes and complete library structures were annotated by appending the barcode sequence to their header. Following barcode annotation, adapter sequences were removed using Cutadapt. Demultiplexing was then performed with BBMap's demuxbyname.sh script, generating individual FASTQ files, each corresponding to a single SAG.

To distinguish genuine cell-associated barcode groups from background noise, the read count distribution of all the groups was analyzed. A knee plot (For example, Fig. 1c) was generated to identify a threshold barcode count (inflection point of the knee plot). Barcode groups beyond this threshold were considered valid SAGs and retained for further analysis.

**Purity and coverage assessment.** Purity and genome coverage metrics were calculated to evaluate the quality of each valid SAG in the sequencing of the *E. coli* and *B. subtilis* mixture and mock community. Purity was determined as the proportion of reads within a SAG mapped to the dominant species, based on alignment results from

Minimap2. Genome coverage was quantified using Samtools by calculating the percentage of the reference genome covered by aligned reads.

## AUTHOR CONTRIBUTIONS

Y. Liu conceived the idea of CAP-seq; J. Li, and R. Zhang developed the CAP-seq workflow supervised by Y. Liu; J. Zhang and Y. Luo assisted in the method development; Z. Song assisted the CAP-seq library preparations; M. Li and Y. Du performed CAP-seq sequencing of the samples included in this study; X. Li collected the stool samples; S. Li and X. Zhai built the bioinformatic pipeline; Y. Liu, S. Li, X. Zhai and W. Wei analyzed the data; Y. Liu, W. Wei, and T. Zhang supervised the microbiome study; Y. Liu prepared the figures and wrote the manuscript.

## ACKNOWLEDGEMENT

The authors would like to thank Dr. Jian Li for providing *E. coli* and *B. subtilis* strains, Dr. Quanjiang Ji for *S. aureus* and *K. pneumoniae* strains, Dr. Lichun Jiang for assistance in computation resources, and Drs. Jingwei Zhang, Ruyuan Song, and Suwen Zhao for many helpful discussions. Part of the bioinformatic analysis was conducted at the HPC Platform of ShanghaiTech University.

This work is financially supported by the start-up funding of ShanghaiTech University, the National Natural Science Foundation of China (32300081), the Shanghai Science and Technology Committee (23QA1406600), the ShanghaiTech AI4S Initiative (SHTAI4S202404) and ShanghaiTech Development Fund (SPST-YSFZ-2024-01).

## COMPETING INTERESTS

Y. Liu, R. Zhang, J. Li, and Y. Du are co-inventors on a patent application filed by ShanghaiTech University based on the CAP-seq workflow described in this article. The other authors declare no competing interests.

# Figures

**Figure 1. CAP-seq enables high-throughput and long-read sequencing of single-cell genomes.**
(a) Conceptual workflow of CAP-seq. The workflow generates amplified and barcoded single-cell genomic fragments suitable for nanopore sequencing, with only two microfluidic steps required. (b) Key features of CAP, a kind of semi-permeable hydrogel compartments. Through permeability engineering, the CAPs retain long DNA fragments (> 600 bp) while being permeable to other reagents used in the workflow. This allows for repeated washing and buffer exchange, enabling multistep processing of the individual genomes being retained. (c-d) Validation of CAP-seq using a 1:1 mixture of *Escherichia coli* and *Bacillus subtilis* cells. (c) Read distribution across different

barcode groups. (d) The length profile of all the reads after quality control, binning, and barcode trimming. (e) Scatter plot depicts the number of *E. coli* and *B. subtilis* reads associated with each barcode group.

**Figure 2. Cap-seq provides an unbiased profile of rare microbial populations and generates high-coverage single amplified genomes.** (a) A mock microbial community to examine CAP-seq. (b) Purity distribution of all SAGs from the mock community sample, with 91% SAGs exceeding 95%. (c) Relative abundance of microbial species derived from CAP-seq and metagenomic shotgun sequencing. Compared with metagenomics, CAP-seq is more capable of capturing rare taxa, with single-cell accuracy. (d) Genome coverage of 3,490 SAGs against the sequencing depth, separately plotted by species. (e) Relative gene abundance of AmpR, sfGFP, and CmR across 3,210 *E. coli* SAGs.

## References

1. Jiao, J.Y. et al. Microbial dark matter coming to light: challenges and opportunities. *National Science Review* **8**(2021).
2. Rinke, C. et al. Insights into the phylogeny and coding potential of microbial dark matter. *Nature* **499**, 431-437 (2013).
3. Yarza, P. et al. Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nature Reviews Microbiology* **12**, 635-645 (2014).
4. Dykhuizen, D.E. Santa Rosalia revisited: Why are there so many species of bacteria? *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology* **73**, 25-33 (1998).
5. Parte, A.C., Carbasse, J.S., Meier-Kolthoff, J.P., Reimer, L.C. & Göker, M. List of Prokaryotic names with Standing in Nomenclature (LPSN) moves to the DSMZ. *International Journal of Systematic and Evolutionary Microbiology* **70**, 5607-5612 (2020).
6. Chaffron, S., Rehrauer, H., Pernthaler, J. & von Mering, C. A global network of coexisting microbes from environmental and whole-genome sequence data. *Genome Research* **20**, 947-959 (2010).
7. Qin, J.J. et al. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* **464**, 59-70 (2010).
8. Methé, B.A. et al. A framework for human microbiome research. *Nature* **486**, 215-221 (2012).
9. Lok, C. Mining the Microbial Dark Matter. *Nature* **522**, 270-273 (2015).
10. Medema, M.H., de Rond, T. & Moore, B.S. Mining genomes to illuminate the specialized chemistry of life. *Nature Reviews Genetics* **22**, 553-571 (2021).
11. Pavlopoulos, G.A. et al. Unraveling the functional dark matter through global metagenomics. *Nature* **622**, 594-602 (2023).
12. Altae-Tran, H. et al. Uncovering the functional diversity of rare CRISPR-Cas systems with deep terascale clustering. *Science* **382**, eadi1910 (2023).
13. Wilkinson, B. & Micklefield, J. Mining and engineering natural-product biosynthetic pathways. *Nature Chemical Biology* **3**, 379-386 (2007).
14. Chiang, C.Y., Ohashi, M. & Tang, Y. Deciphering chemical logic of fungal natural product biosynthesis through heterologous expression and genome mining. *Natural Product Reports* **40**, 89-127 (2023).
15. Goig, G.A. et al. Whole-genome sequencing of directly from clinical samples for high-resolution genomic epidemiology and drug resistance surveillance: an observational study. *Lancet Microbe* **1**, e175-e183 (2020).
16. Liu, Y.X. et al. A practical guide to amplicon and metagenomic analysis of microbiome data. *Protein & Cell* **12**, 315-330 (2021).
17. Ustick, L.J. et al. Metagenomic analysis reveals global-scale patterns of ocean nutrient limitation. *Science* **372**, 287-291 (2021).
18. Nissen, J.N. et al. Improved metagenome binning and assembly using deep variational autoencoders. *Nature Biotechnology* **39**, 555-560 (2021).

19. Mattock, J. & Watson, M. A comparison of single-coverage and multi-coverage metagenomic binning reveals extensive hidden contamination. *Nature Methods* **20**, 1170-1173 (2023).
20. Yue, Y. et al. Evaluating metagenomics tools for genome binning with real metagenomic datasets and CAMI datasets. *BMC Bioinformatics* **21**, 334 (2020).
21. Poulsen, C.S., Ekstrom, C.T., Aarestrup, F.M. & Pamp, S.J. Library Preparation and Sequencing Platform Introduce Bias in Metagenomic-Based Characterizations of Microbiomes. *Microbiology Spectrum* **10**(2022).
22. Nearing, J.T., Comeau, A.M. & Langille, M.G.I. Identifying biases and their potential solutions in human microbiome studies. *Microbiome* **9** (2021).
23. Sun, Z. et al. Challenges in benchmarking metagenomic profilers. *Nature Methods* **18**, 618-626 (2021).
24. Jin, H. et al. Hybrid, ultra-deep metagenomic sequencing enables genomic and functional characterization of low-abundance species in the human gut microbiome. *Gut Microbes* **14**(2022).
25. Pust, M.M. & Tümmler, B. Bacterial low-abundant taxa are key determinants of a healthy airway metagenome in the early years of human life. *Computational and Structural Biotechnology Journal* **20**, 175-186 (2022).
26. Llorens-Rico, V., Simcock, J.A., Huys, G.R.B. & Raes, J. Single-cell approaches in human microbiome research. *Cell* **185**, 2725-2738 (2022).
27. Zheng, W.S. et al. High-throughput, single-microbe genomics with strain resolution, applied to a human gut microbiome. *Science* **376**, eabm1483 (2022).
28. Lan, F., Demaree, B., Ahmed, N. & Abate, A.R. Single-cell genome sequencing at ultra-high-throughput with microfluidic droplet barcoding. *Nature Biotechnology* **35**, 640-646 (2017).
29. Moragues, T. et al. Droplet-based microfluidics. *Nature Reviews Methods Primers* **3**, 32 (2023).
30. Li, L.Y. et al. Permeability-Engineered Compartmentalization Enables In Vitro Reconstitution of Sustained Synthetic Biology Systems. *Advanced Science* **9**(2022).