1

Computational anatomy: the cerebellar microzone computation, a

hypothesis

Mike Gilbert[1] *, Anders Rasmussen[2]

[1]School of Psychology, College of Life and Environmental Sciences, University of

Birmingham, Birmingham B15 2TT, UK   ORCID: 0000-0003-4415-6915

[2]Department of Experimental Medical Science, Lund University, BMC F10, 22184 Lund,

Sweden   ORCID: 0000-0003-4477-369X

*Correspondence: m.gilbert.1@bham.ac.uk;

Abstract

The cerebellum is a large brain structure. It is poorly understood how the cerebellum codes and processes information. It is generally assumed that the representation of information in the brain is in high resolution and replicable with high fidelity, and that those features make it precise and characterise the inputs to sophisticated computations. Replicability is an important assumption of neural network learning models, including cerebellar models, which depend on replicability because they rely on pattern memory. We challenge those ideas. We argue that the cerebellar microzone computation is the aggregate of simple unit computations and a passive effect of anatomy, unaided and unlearned, which we are able to recreate *in silico* in the form of a network of unit functions. Unit functions are attributed to identified anatomical hardware. The functions themselves are linear, with evidence. In this form, we are able to model large cell networks in high relevant definition, without simplifying or omitting parameters that affect the computation. Conversion of inputs to output of the system is mediated by a chain of rate codes, and not supplanted by learning. Unit populations are organised functionally in layers that span the network. A learning algorithm and replicability are unnecessary to explain the evidence.

Keywords cerebellum; theory; model; computation; anatomy; microzone; network; code

Lay summary

It is poorly understood – i.e., not understood – how the cerebellum represents information and uses it to generate motor outputs, and how outputs are coordinated with each other and with behaviour to control movement. We try to answer those questions.

The paper is in two main parts. In the first, we model the cerebellar network computation, and in particular the microzone computation, the middle layer of the network. We propose the computation is unlearned, simply a passive effect of cell morphologies and network geometry. This is still sufficient to distil a single output rate – motor output of the cerebellum – from 80,000 input signals.

In the second part, we propose that the topography of input and output of cerebellar motor circuits could automate unconscious control of motor sequences. To illustrate how that could work, we model swimming. We propose that the cerebellum connects proprioception-motor loops so that movement itself can drive coordinated motor output with only sparse (go, no go) executive input. The role of the cerebellum and functional topography of connections may remain today a variation of the same principles.

1. Introduction

Most of the mass and volume of the cerebellum is made up of the cerebellar cortex,

which is divided into an inner and outer layer, the granular layer and molecular layer.

The molecular layer is divided functionally into long, thin strips called microzones [1-3].

Microzones lie at right angles to the main portion of the axons of granule cells, called

parallel fibres, which themselves lie parallel to each other and the cerebellar surface. A

microzone is populated by a few hundred Purkinje cells. Purkinje cells receive contact

in passing from parallel fibres and are the sole outputs of the cerebellar cortex (Fig 1).

Firing rates of the output cells of the cerebellum are bidirectionally sensitive to

Purkinje cell rates [4]. This provides the main control of output cell rates.

Neuronal firings rates are thought to represent values received as inputs to network

computations. We present the simple idea that there is a passive, unlearned

computational effect of microzone anatomy. It is contentious partly because it

conflicts with traditional learning models, and partly because it depends on linear

communication between cells, for which there is significant but incomplete evidence.

Do we need a rethink? We believe probably yes, partly because the old view has not

aged well and partly to account for new evidence. In the classic cerebellar learning

model [5-7], the chief instrument of the network computation is the learned

modification of parallel fibre synaptic weights. The 'purpose' is that remembered

patterns – i.e., received by training-modified synapses – return a learned response.

The learned response displaces the naive response to unprocessed input rates.

However, experimental confirmation of the predicted role of synaptic modification in

motor learning did not materialise [8-11], and naturally-occurring grades of synaptic

depression were not found [12-14].

There is a significant body of evidence that rate information is linearly conserved in

signalling between cerebellar neurons [15-18], including synaptic signalling in the

molecular layer [19-22], and that cells are adapted for faithful transmission [23-29].

Firing rates linearly code behavioural and task metrics 'at all levels of the cerebellar

circuit' [30 p.239, citing 30 references]. Neurobiology is highly adapted to control for

biophysical and other noise. There are cerebellar examples of exactly this, where cell-

to-cell communication is adapted to control for variables that are not code, i.e., to

conserve fidelity of linear transmission of rate information [15-18, 23, 24, 27-29, 31-

33].

Perhaps surprisingly, it is not obvious how to account for linear signalling as part of a

computational mechanism. Also, it is probably still the mainstream view that synaptic

weights will be involved somehow. We propose here a mechanism of the microzone

computation in which anatomy has an unlearned computational role. This is passive,

simply a result of cell morphologies and network geometry. Nonetheless, it is able,

plausibly, to run uninterrupted calculations in parallel, in real time, and doesn't need

synaptic weights.

We propose – as artificial neural network models do – that signals traffic flows through

computational units organised in layers, but with the difference that:

1) Our units are matched with physiological hardware (and physical counterparts are

not synapses);

2) Units form functional populations whose size and spatial dimensions are a

consequence of known physiology; and

3) Unit functions are not conjecture, they are linear, with the basis in evidence of

linear signalling, cited in context.

Note that we do not propose to be mathematically interesting. The heart of the

proposal is that the physiological computation is in fact not sophisticated, against

expectation – at least, there is a plausible argument that this is the case, which we

present as an alternative to current models. For that reason, physiology has a high

profile in the paper. The sophistication of the biological design is found in the fact that

physiology is adapted to isolate computationally relevant parameters (and

computations themselves) from interference by other variables.

Fig 1. Schematic of data flow in the cerebellum.

A Sagittal section of the rat cerebellum (obtained by multi-photon microscopy, courtesy Tom Deerinck). The outer layer, the molecular layer, of the cerebellar cortex is stained green and the inner layer, the granular layer, is blue. **B** and **C** Purkinje cells in a sagittal slice (two-photon laser scanner micrograph, courtesy Mike Häusser). The Purkinje cell arbour is large from this view, filling the molecular layer vertically. It is flattened in the plane orthogonal to parallel fibres. **D** Top. Schematic of a parallel fibre view (i.e., the

same view as a–c) of a row of 80 Purkinje cell territories, but straightened out. The molecular layer is divided functionally into long, thin strips called microzones. At a mid-range estimate, a microzone is 75–100 Purkinje cells long and 4 wide (measuring 1,500–2,000 x 150 µm). Inset. A short section viewed from the cerebellar surface. **E** Simplified circuit wiring, highly preserved across vertebrates. Red: glutamate; blue: GABA. Brown boxes contain information about network connections. Microzones lie at right angles to parallel fibres. Tens of millions of parallel fibres pass through a microzone, making contact in passing (if at all) on Purkinje cells and interneurons. END CAPTION

2. Methods

2.1 The principles of the computation

The cerebellar network (as distinct from organisation into microcircuits that include a cell group in the inferior olive) is formed of a microzone (the middle layer), the region of the granular layer which supplies parallel fibre input to it (the input layer), and a cell group in the deep cerebellar nuclei which includes the output cells of the network (which are also the output cells of the cerebellum, the outer layer) (Fig 2). For non-cerebellar physiologists, a layer of a network has a different meaning to the same word used in granular layer and molecular layer. For example, a microzone, the middle layer

of a network, is a division of the molecular layer, a stratum of the cerebellar cortex.

Each successive network layer – input, middle, output – contains a smaller number of

cells than the last. The input layer contains millions. The number of output cells may

be around 50.

The cerebellum, we argue, exploits statistical effects of taking random samples. Say a

population of values (with any frequency distribution) is randomly sampled. Samples

are independent, sometimes called sampling with replacement – an illustration

sometimes used is that coloured balls taken from a bag are replaced in the bag

between samplings. There are n samples, sample size m. If we take the mean (or sum)

of each sample, and n is large, the sample means have a normal or near-normal

distribution, by the central limit theorem. (This works even if m is a range.) The new

distribution is centred on the same mean as the sampled distribution, but narrower.

Large values of n and m give a precisely-shaped new distribution. Modest values give

an untidier (but still normal-tending) shape. However, if the new distribution is then

itself independently randomly sampled, and the process is repeated, the results

nonetheless become more focused, and the relationship of the means of successive

distributions is well preserved. We will argue that the cerebellar microzone provides

(part of) the physical hardware for the cerebellar network to distil output from input in

this way, noting that at each step the relationship of the means includes a constant.

2.2 The relationship of physiology and model parameters

In this contention, the neurophysiology of a microzone is the physical form of data organised in layers. A data layer spans a network, so that it extends from end to end of a microzone. Data are computed by units. In its computational form, a unit is a linear function. In its physical form it is the component (a cell or part of a cell) that executes the unit function and returns a value. The biophysical form of a value depends on the layer. A unit randomly samples the afferent data layer. During behaviour, units receive a constant turnover of inputs and compute ceaselessly, so that the value coded by a unit is an analogue signal.

This is a third use of the word layer, again with a different meaning. Here, a layer is a layer of computations. Units in their physical form themselves form several layers in a single microzone, but that is immaterial to the computation. A microzone is the computational equivalent of a plane at right angles to the cerebellar surface and parallel fibres, where all cells sample the same convergent parallel fibre signals.

We argue that the microzone computation is the passive effect of anatomy which we are able to recreate *in silico* in the form of a network of unit functions. Functions are attributed to identified anatomical hardware and the functions themselves are evidenced. The evidence is discussed in context, in the main text. We simulate a moment in time, and more exactly the computation performed on a very short period

of inputs. In this form, we are able to model large cell networks in high relevant

definition, without simplifying assumptions. Put another way, if we are correct about

the physical form of units, and we are also correct that unit functions are linear and

about the physical dimensions of data layers, the model includes all computationally

relevant parameters. The sophistication of the system is biological, not computational.

It lies in the fact (as noted) that neurophysiology is adapted to exclude an effect of

other variables, including interference by (a redundant effect of) biophysical variables

which execute transmission of signals.

The simulation provides a demonstration in principle that processing by anatomy

provides a mechanism that:

1) Autonomously strictly regulates the fraction of active parallel fibres that converge at

any time on a microzone; and

2) Converts mossy fibre firing rates received by the input layer of a network into well

predicted coherent firing of middle-layer, microzone-grouped Purkinje cells.

Units are connected by known anatomy. Information about anatomy is detailed and

not contentious. There is some variation between species, and indeed between

estimates, but network architecture is strongly conserved, suggesting that variation

does not affect the form of the computation. The number of active inputs to a unit is in

some cases randomly variable with time. However, we can simulate that by, for each

unit, selecting the number at random from the reported range (with a distributed

probability if we know what that is).

It is unnecessary to know which individual cells contact each other because

(biologically and in the model) contact is at random inside microzone sectors defined

by a combination of topography and scaffold-like architecture of the cerebellar cortex.

Biologically, a sector is a volume with spatial dimensions. *In silico*, we can represent

that as a population of unit functions. We know from anatomy (or can derive) the size

(and shape) of unit populations, population ratios, and convergence and divergence

ratios.

202   2.3 Assumptions

203   2.3.1 THE GRANULE CELL CODE

We have previously simulated the granular layer mechanism which converts mossy

fibre signals into granule cell and therefore parallel fibre signals [34]. The granular

layer methodology and a review of the evidence appears in Supplemental Materials.

We use the granular layer mechanism to generate parallel fibre input to the microzone

computation. Previously, we modelled a sagittal strip of the granular layer. In this

paper, we extend that to the larger region (around 40 strips) that supplies parallel fibre

input to the microzone that equally bisects the region viewed from the cerebellar

surface – i.e., the input and middle layers of a network, respectively. The outputs of

the input layer mechanism are the homeostatic regulation of the fraction of active

input layer granule cells, and therefore the fraction of active parallel fibres which

converge on the central microzone, and the parallel fibre code.

In this paper, the workings of the granular layer mechanism are effectively an

assumption. The parallel fibre code has invariant features, regardless of the

information it contains. Information is coded at collective level, in the joint activity of

cells which are active at the same time. More exactly, it is coded in the frequency

distribution of the rates at which those cells fire, and more exactly still this reflects the

status of cells in a moment of time where they intersect an orthogonal plane.

Therefore, it is coded in the frequency distribution of active inputs to the central

microzone.

Firing rates are randomly distributed among active parallel fibres and active parallel

fibres are a random subset of the general population. Accordingly, any random sample

of active inputs is (at any moment) a random sample of the same distribution, and

because all sectors of a microzone receive a large number of active inputs at any time,

all sectors receive the same code, as one. In this paper, we treat the form of the

parallel fibre code as an assumption. The reader may, if they wish, read the granular

layer paper, but if not, they are not at a disadvantage.

granule cell – up to ~ 35m parallel fibres have input to a microzone

~80,000 mossy fibres have input to the pink zone

strips

files

6 mm – a microzone receives parallel fibre input from a range of 3 mm on each side

20 mm

microzone – contains ~400 PCs and 8,000 stellate cells

A file ≈ the width of parallel fibre input to a single PC

3mm

Expanded coronal X-section

To deep nuclei ~ 50 output cells

parallel fibre

Fig 2. Schematic of a network.

The number of cells in each network layer diminishes from the input layer to the middle

layer to the output layer. We use functional and nominal divisions of a network to

quantify the proposals. A 'strip' is a functional division of the granular layer lying parallel

to microzones (white stripes are also strips). A 'file' is a nominal division of the granular

layer lying at right angles to strips, with approximately the width of a Purkinje cell. A

'field' is the intersection of a strip and a file; it is also a nominal division, merely a device

used for modelling. END CAPTION

It is a final but important feature that code in this form mimics sampling with

replacement. Purkinje cells and molecular layer interneurons are both severely

flattened in the plane orthogonal to parallel fibres. Both receive contact from a

random sample of passing parallel fibres, and therefore receive contact from a random

sample that are active. They each receive, therefore, individually, at any moment, a

random sample of what is effectively the same distribution of parallel fibre rates,

which is independent of the sample received by any other cell, notwithstanding that in

most cases they receive contact from a sample of a different population of cells.

2.3.2 PARALLEL FIBRE SYNAPTIC MEMORY

The traditional learning model, which centrally features synaptic memory, has not

received confirmation by experimental work. However, there is still a mainstream

feeling that the computation will turn out to involve synaptic weights. In this view,

core assumptions are retained, and the aim of further work is to understand how

plasticity at multiple sites is related, and related to function. Historically, there has

been strong emphasis on researching learning [35, 36]. It is regarded as naive to have a

different opinion.

We think it is unnecessary to interpret experimentally induced learning as evidence of

functionally graded synaptic weights. There is now a range of evidence – both in the

granular layer, reviewed in Supplemental Materials, and in the molecular layer, cited

and discussed in context, in the main text – that firing rates are linearly related, and

that synaptic transmission is adapted for linear transmission, and of linear

relationships between firing rates and behavioural metrics (suggesting linearity is not

merely an artefact).

One of the present authors has previously proposed a learning outcome which is

consistent with linear transmission. In that proposal, the result of learned parallel fibre

synaptic modification is to polarise synaptic weights such that all synapses either

transmit robustly (working synapses) or do not transmit at all (silent synapses). This

was first proposed, with a synaptic mechanism, here [37]. The learning outcome is the

ratio of working to silent synapses. Working synapses are a random subset of the

population. There is no pattern memory. The effect on the response of the

postsynaptic cell is independent of the pattern in which parallel fibres are active and

inactive. Following training, all permutations are received at the same ratio of working

and silent synapses regardless of whether or not any particular pattern was involved in

training, even if it has never been received before. The effect of learning is to make a

gain adjustment to the network computation.

We make it a further assumption in this paper that the learning outcome has these

features. Therefore, it is unnecessary to include learning here.

The learned ratio is the same on all Purkinje cells in the same microzone. Microzones

are defined topographically by their climbing fibre input [1, 2] which is received from a

discrete cell group in the contralateral inferior olive. In response to stimulation,

climbing fibres in a group fire together, so that a microzone that receives the output of

a group receives climbing fibre tuition as a volley. Discharge synchrony of a climbing

fibre group is enhanced by dendritic electrotonic coupling of olivary cells [38] and is

seen in anaesthetised mice in response to sensory stimuli [3]. Coherence of complex

spikes between functionally-grouped Purkinje cells (Purkinje cells reliably respond to

climbing fibre discharge with a complex spike) is low when it includes spontaneous

discharge but high in response to sensory stimuli [39]. This provides the substrate for

microzone-grouped Purkinje cells to receive the same training, so that they learn as a

unit.

So, all Purkinje cells receive, at working synapses at any moment, a large random

sample of parallel fibre firing rates (and therefore the same code) and the same

number of signals (with no effect of the pattern).

2.4 Regulation of the fraction of active parallel fibres

2.4.1 A CASE FOR SPARSE CODE (SMALL FRACTION)

To simulate the computational effect of random sampling, we need to know or derive

sample sizes (the number of inputs to each type of unit), and therefore in turn what

fraction of parallel fibres is active at any time. There is a long-standing idea that the

fraction of parallel fibres that are active at any time is regulated [6], fixed [5], and low

(termed sparse code). We disagree with the theoretical reasons, but agree the code is

sparse, on these grounds.

1) Low density is stable (Fig 3). Randomly fluctuating density would be noise. Stability

may be enhanced by feedback via granule cell ascending axon contact on Golgi cell

basal dendrites [25] and parallel fibre contact on Golgi cell apical dendrites, acting

partly through gap junctions [40].

2) Low density is energy efficient. Granule cells are very numerous, and they are not

myelinated (so spike propagation is costly).

3) The reported number of coincidentally active inputs to a stellate cell is in single

figures [41], consistent with low density of co-active parallel fibres (see Table 1).

4) It is thought that the pattern of active and inactive parallel fibres, and the pattern in

which firing rates are distributed among active cells, are decorrelated – effectively,

quasi-randomised. Low density enhances practical decorrelation, that is, if

decorrelation is not perfectly random – as it may not be [42] – low density increases

the randomness of the pattern of synaptic activations received by targets.

For these reasons, we calibrated the granular layer model to convert inputs to a

network into a low fraction of active granule cells. We chose 0.5% active because it has

support from (as far as we know) the only estimate of the number of co-active inputs

to a stellate cell (per section 2.5). We suspect it may be higher – perhaps 1% – but a

very low fraction is a harder test that an anatomical computation is feasible (because

sample size is smaller).

2.4.2 HOW STRICT IS REGULATION OF THE FRACTION OF ACTIVE PARALLEL FIBRES?

It would cause interference with the computation if the fraction of active parallel fibres

varied randomly between sectors of a microzone, or with time. There is no

experimental evidence that the fraction is stable or that it is not stable. The granular

layer computation simulates the interaction of excitatory input to the cerebellum that

is received by granule cells and by granular layer interneurons called Golgi cells, and

concurrent inhibition of granule cells by Golgi cells (Supplementary Materials). We

adapted the granular layer model to generate parallel fibre input to a microzone. One

of the outputs of the model is the number of active granule cells per unit volume of

the granular layer.

The cerebellar cortex is divided only functionally into microzones and networks.

Anatomically, it appears as a seamless carpet of cells. To calculate the number of

active granule cells we divided it nominally into sagittal strips (the direction of

microzones) and bands which cross strips at right angles (the direction of parallel

fibres). Parallel fibre input to a microzone is supplied by 20 strips either side. These are

crossed by 100 bands, making a total of 4,100 200 µm x 150 µm 'fields' where a band

crosses a strip (the extra 100 is the strip of the granular layer under the middle

microzone). So, there are 100 fields in a strip and 41 in a band.

Together, strips and bands represent a region of the granular layer with dimensions

based on a 3 mm range of parallel fibres in each direction from the granule cell soma

[43, 44] and a microzone dimensions of 150 μm x 2,000 μm viewed from the cerebellar

surface [2, 3, 45].

The results are shown in Figure 3. The number of active granule cells in each field gives

the number in each file. The number in a file is equal to the number of active parallel

fibres at the intersection of that file with the middle microzone. Local regulation of the

number of active granule cells in each field in a strip (Fig 3 columns 2 and 5) becomes

still tighter regulation of the number of active parallel fibres in a file (Fig 3 columns 3

and 6), a statistical effect of cerebellar cortical architecture. Density is independent of

the frequency distribution of mossy fibre rates (Fig 3 columns 1 and 4). At low density

of active granule cells, the regulated density of active parallel fibres is hyper strict.

Physiological regulation may be still stricter, because the number of active granule

cells in each field may not vary as much as it does in the simulation. Local feedback via

the granule cell ascending axon [25, 46] may condense the physiological range of field-

to-field variation. Feedback via parallel fibres [5, 40] would (additionally) reduce the

range of file-to-file variation. These are not represented in the simulation.

So, by our calculation, data are not corrupted by noisy variation of the fraction of

parallel fibres that are active, because the fraction is strictly regulated by the granular

layer computation.

Fig 3. Recoding in the granular layer regulates the fraction of active parallel fibres.

Columns 1 and 4. Firing rates received as input to the granular layer region that supplies

parallel fibre input to a microzone. Data represent inputs to a sagittal strip that traverses

the region, selected at random. Inputs to each strip were independently randomly

generated with a uniform and discontinuous distribution, respectively. Data represent a

moment of time. **Columns 2 and 5.** The percentage of granule cells that fire in each field

of a single strip, again selected at random. **Columns 3 and 6.** The percentage of granule

cell that fire in each band, therefore also the percentage of active parallel fibres in each

band. **Top row.** ~30% of granule cells are active. **Bottom row**. Golgi cell inhibition of

granule cells is adjusted by a constant so that a lower fraction of granule cells fire,

around 0.5%. y axes are scaled to the data (so the contraction of the data range in the

bottom row is even greater than it appears). The shape of the sampled distribution

(columns 1 and 4) does not affect the results, suggesting that regulation of analogue

inputs at changing rates has the same result. END CAPTION

2.5 Functional anatomy

Cell anatomies have a function in the computation. This is not incidental but a main

driver of their physical form and the scaffold-like geometry of the contact they make

on each other. Cell morphologies, cell ratios, orientation to each other, spatial

organisation, the shape and size of dendritic and axonal territories, convergence and

divergence ratios, and the probabilities that they make and receive contact, all

contribute to function. We describe them here, in the methods section, because we

derive parameters from anatomy.

The number of Purkinje cells in a microzone probably varies. We use 400. Around

350,000 parallel fibres pass through the dendritic field of a single Purkinje cell, of

which one in two make contact, normally at a single synapse (sometimes two, average

1.24) [47]. Purkinje cells are severely flattened in the plane orthogonal to parallel

fibres and to the cerebellar surface, filling the molecular layer vertically. As noted,

microzones lie across the path of parallel fibres at right angles, so that parallel fibres

pass through by the shortest distance.

Purkinje cells receive inhibition from interneurons, stellate cells and in some

vertebrate groups, basket cells. We do not consider basket cells. The stellate cell

territory is sagittally flattened in the plane orthogonal to parallel fibres. Stellate cells

form planar networks [48] that occupy the space (about 40 μm) between Purkinje cells

[49]. The stellate cell dendritic territory varies with the depth of the cell body in the

molecular layer. For superficially located ('outer-level') cells it is around 80 x 80 μm. At

deeper level ('inner level') it is larger, around 120 x 120 μm [49 pp.217-221]. The

number of parallel fibres which pass through an inner-level stellate cell territory is

given by $a/((b \text{ x } c)/120^2)$, assuming parallel fibres are uniformly distributed, where a is

the number that pass through a Purkinje cell territory (350,000 [50]), and b and c are

the dimensions of the Purkinje cell territory in the same plane (200 x 300 μm), giving

~84,000. If 0.5% are active in the general population, around 420 of those are active.

It has been estimated that a stellate cell receives contact from several hundred parallel

fibres [49, 51], which we take as 1,000. Working with these estimates, 1 in 84 make

contact (i.e., 1,000 out of 84,000), so the probability that an active cell makes contact

is 1/84 = ~0.0119. Accordingly, contact by x (out of 420) active cells has a probability

given by

405
$$\frac{420!}{x!\,(420 - x)!} * 0.0119^x * (1 - 0.0119)^{420-x}$$

Probabilities are shown in Table 1 for the range x = 0 to 12, where x is the number of

active parallel fibres that make contact and p is the probability of that number. The

distribution of probabilities is in good agreement with the modest number – single

figures – suggested by 'two to eight substantial EPSPs' [41 p.9628].

Table 1 The probability, p, that a stellate cell receives contact from x active parallel

fibres

x	0	1	2	3	4	5	6	7	8	9	10	11	12
p	.0066	.0331	.0836	.1403	.1762	.1765	.147	.1047	.0651	.0359	.0178	.008	.0033

413 **Caption.** Table 1 shows the distributed probabilities of the number of active inputs to

414 inner-level stellate cell.

3. Results

A discussion of related evidence is included in the results section. We believe it gains by context.

3.1 Stellate cells

We divided cells in a microzone into nominal 'sectors'. A sector is the size of the intersection of a band and a microzone, so a sector and the field below it together form a column of the cerebellar cortex. Cells in a sector randomly sample parallel fibre rates in that band. Like fields, sectors *are simply a device*. They do not represent a separation of data streams or that there is any difference or compartmentalisation of the way data are processed.

The granular layer model returns the firing rates of granule cells in each field, and therefore the frequency distribution of parallel fibre rates in each file, and so rates received by each sector of the middle microzone, as noted. We next used that to simulate random sampling of parallel fibre firing rates by stellate cells in the middle microzone. There are an estimated 16 stellate cells for each Purkinje cell [52], meaning 80 stellate cells per sector (there are four Purkinje cells per sector so five layers of stellate cells are needed to inhibit them from both sides). Sample size was randomly generated with the probabilities in Table 1. This gave the number of active inputs, and input rates, to each stellate cell in each sector.

Molecular layer interneurons, which include stellate cells, reflect 'granule cell input

with linear changes in firing rate' [21 p.6]. We assume that stellate cell firing is a linear

function of the mean of input rates, which we take to be the mean itself (the sum

works as well). This is not to suggest that stellate cell firing rates are equal to the mean

of input rates, but that they are some linear function of the mean. As the physiological

function is not known, and our purpose is only to demonstrate a statistical relationship

of data layers, and not to calculate actual firing rates, we use a normalised scale. The

results are shown in Figure 4.

To put that in the language we used earlier, each stellate cell is the physical form of a

notional unit, and the afferent data set is the frequency distribution of parallel fibre

firing rates. Granule cells fire in short high-frequency bursts [21, 53, 54], and only few

are active at any time. But the high turnover of active cells means the distribution is

constantly changing. Sample size fluctuates; it varies with time and, at any time, the

number of active inputs to a unit has a distributed probability with a result that is

independent of other units. The unit computation is a linear function. In physiological

conditions, inputs are computed ceaselessly and in parallel by all units – 80 per sector,

all sectors, all (at any moment) effectively sampling the same distribution of parallel

fibre rates, notwithstanding that they all receive contact from different parallel fibres.

The outputs – each an analogue signal – become the data set sampled by the next

layer of units.

Fig 4. The microzone computation.

SC: stellate cell; PC: Purkinje cell; compt: PC dendritic compartment; mf: mossy fibre; pf: parallel fibre. **A** Mean and SD of mossy fibre input to 41 nominal strips of the granular layer. Strips are 150 microns wide and together make up the region supplying parallel fibre input to the microzone that equally bisects it, indicated by a dashed grey line. The morphology of mossy fibres means that all locations of a strip receive a large random sample of input rates to the whole strip. **B** Frequency distribution of mossy fibre rates received by a single nominal band, 200 microns wide. Bands cross strips at right angles, the direction of parallel fibres. Granule cell firing rates in a band are accordingly parallel fibre rates received by the microzone where they intersect. As bands cross all strips, they all receive the same distribution of mossy fibre rates. There are 100 bands. **C** Parallel fibre rates in a randomly selected band. The frequency distribution of parallel fibres rates is the same in all bands. This is always true, and does not depend on strips receiving the same mossy fibre distribution. **D** Parallel fibre rates received as input to stellate cells in the central microzone, obtained by randomly sampling parallel fibre rates (80 times per file). Sample size was randomly generated with Table 1 probabilities. The y axis range is 60 normalised Hz, down from a mossy fibre range of 250 Hz. **E** Outputs of the stellate cell function. Data represent 16 cells per sector. A sector is the intersection of the central microzone and a band (hence 1,600 data points). There is a further contraction of the data range. **F** The Purkinje cell dendritic arbour was nominally divided into 16 compartments, each randomly sampling local outputs of the stellate cell

function. Data are the sample means, shown for a single Purkinje cell per sector in the

central microzone (so there are 1,600 values, 16 compartments x 1 Purkinje cell x 100

sectors). **G** A Purkinje cell receives inhibition from two planar networks of stellate cells,

one each side. Step (F) was repeated, and the data averaged for each compartment. **H**

Mean of the compartment means calculated for each Purkinje cell in the central

microzone (so 400 data points, 4 Purkinje cells per field x 100 fields), representing

somatic integration of dendritic compartment data. **I** The same data but with the y axis

range further shortened. A uniform distribution of mossy fibre rates with a range of 50–

300 Hz in (A) becomes a very narrow Purkinje cell range with a normal distribution. **J** The

output of a microzone is channelled down onto a smaller group of cells which include

the output cells of the network. Each output cell receives input from a random sample

of 30–50 Purkinje cells [55]. Data are the sample means for 50 output cells. **K–T** We

repeated (A)–(J) but with normally distributed mossy fibre rates (SD 40), with the same

result. We also obtained the same results (not shown) with a discontinuous mossy fibre

distribution (range 50–300 Hz). END CAPTION

Figure 4 data represent a moment of time. We used values to represent mossy fibre

signals that drive parallel fibre input to a microzone, as before. Input rates to each strip

were independently randomly generated with a uniform (Fig 4A) and normal (Fig 4K)

distribution. Data are displayed as the mean and SD of input to each strip. Firing rates

received by each field were derived from input to strips (Supplementary Materials),

and input to bands – a row of fields crossing strips at right angles – were derived from

input to fields. Input to a randomly selected band is displayed as a histogram (Fig 4B

and L). Some very high mossy fibre rates have been recorded but the typical

physiological range may be lower [53, 56]. In this figure, strips all receive the same

range (50–300 Hz). Therefore, so do all bands.

The range of parallel fibre rates is much smaller (shown for a randomly selected band

in Fig 4C and M). This is a result, by design, of the statistical effect of random sampling

by granule cells, and of regulation of the fraction of granule cells which fire (Fig 3),

because the number is controlled by an inhibitory threshold. This limits firing to cells

which receive rates at the higher end of the mossy fibre range.

The parallel fibre range is the same in all bands, so this is the range that stellate cells

all sample (Fig 4D and N). We show data for 16 stellate cells per sector (the number

that sit between two Purkinje cells), so this step generates around 7,200 values (16

stellate cells x 100 sectors x the weighted mean of distributed probabilities). Signals

received by each stellate cell are then passed through the stellate cell unit function.

The data range is further narrowed (Fig 4E and O), from an original range of 250 Hz

down to around 20 Hz. The outputs of the unit functions become the sample

distribution in the next step.

There is direct evidence of a linear relationship of granule cell input and stellate cell

rates [21] but as far as we know not confirmation, or data about the sensitivity or

latency of the response. However, fidelity of transmission may be assisted. GABA

spillover from stellate cell synapses enhances parallel fibre synaptic transmission to

stellate cells [57, 58]. 'It is well established that parallel fibers express functional

GABA$_A$Rs, that their activation depolarizes the axon, and that they increase release

probability at parallel fiber synapses' [59 p.16924, 60].

525   3.2 Purkinje cells

In our simulation, a field contains 4 Purkinje cells. A Purkinje cell receives contact from

stellate cells from both of its flat sides, ~16 each side, and a smaller number of stellate

cells in the two sagittally-neighbouring fields. Inhibition from neighbouring fields is

from a smaller number of stellate cells because not all of them extend their main axon

far enough to reach, and some of them extend their main axon the wrong way. We

represent those conditions as variables. The stellate cell main axon gives rise to

ascending and descending axon collaterals. The vertical range of collaterals and the

horizontal range of the main axon both increase with the depth of the cell body in the

molecular layer [49, 61]. Because of their more limited horizontal range, outer-level

stellate cells are excluded in the simulation from contacting Purkinje cells in

neighbouring fields.

The planar Purkinje cell arbour is large and branches profusely, and stellate cell contact

is dispersed across it. We divide the Purkinje cell arbour nominally into sixteen 50 x 75

µm compartments. Input to a compartment is not confined to (or even necessarily

received from all of) the nearest stellate cells. It is unknown how many stellate cells

innervate a Purkinje cell compartment. Most stellate cells afferent to a Purkinje cell are

in range of most compartments, but not all make contact, and fewer make significant

contact which dominates the postsynaptic effect. We assume six make significant

contact. To represent that in the model, each compartment receives contact from a

random sample of stellate cells which flank a Purkinje cell. Sample size is six. Sampling

is independent of sampling by other compartments, i.e., innervation of a compartment

by a stellate cell does not exclude innervation by the same cell of other compartments.

Here, a notional compartment is a computational unit, and the afferent data set is a

(random, effectively) subset of the local outputs of the stellate cell function. The unit

computation is again a linear function. Spatial and temporal integration of inputs are

absent from the computation because inputs are not discrete. Compartment

membrane voltage is a linear function of stellate cell rates afferent to that

compartment. In its physical form, the postsynaptic effect – the output of the

computation – is an analogue signal. In its computational form, a compartment

continually recalculates inputs.

There is support for a linear effect. It has been reported that Purkinje cell firing is a

linear function of synaptic excitation, with inhibition blocked in slices [19, 20]. In mice

during locomotion, linear relationships are reported between granule cell rates and

interneuron firing, the balance between excitation and inhibition of Purkinje cells and

depolarising or hyperpolarising dendritic membrane voltage, and between membrane

voltage and bidirectional modulation of Purkinje cell simple spike output [21]. The

granule cell → interneuron → Purkinje cell pathway drives a fall in the firing rate of

Purkinje cells as a linear function of the strength of input from granule cells [22].

To declutter the data, we show unit outputs for a single Purkinje cell in each sector (so

1,600 values) (Fig 4F and P). Data are now in a range of around 10 Hz on a normalised

scale (switching calibration would add a layer of conjecture because the actual

function is unreported).

A Purkinje cell receives inhibition from two stellate cell networks, one each side. We

repeated the steps above for the other flanking network, then took the mean for each

compartment (Fig 4G and Q).

At the next step, dendritic signals are integrated at the Purkinje cell soma. Stellate cells

contact Purkinje cells on smooth dendrites [49]. Smooth dendrites extend into all

compartments, mitigating the effect of filtering and distance on dendritic signalling.

Recording from smooth dendrites, locomotion-dependent modulation of dendritic

membrane voltage 'linearly transforms into bidirectional modulation of PC SSp

[Purkinje cell simple spike] output' [21 p.9]. Dendritic signals are a sustained state (we

moot). Passive modulation of somatic charge is fast and bidirectional, and reflects the

average of dendritic states, which transforms into a linear effect on firing. To represent

that, we took the mean of compartment membrane voltages for each Purkinje cell in

all sectors, so that this step returned 400 values (4 x 100 Purkinje cells; Fig 4H, I, R and

S). There is a further contraction of the data range. This then becomes the data set

sampled by the next layer of computational units, with the physical form of a cell

group in the deep cerebellar nuclei, which includes the output cells of the network (not

part of the microzone computation but included in the Discussion).

The result is a progressive contraction of the data range received as inputs to the

system, a cell network. At each step, data are coded in a frequency distribution. There

is a linear relationship of the mean of each distribution with any other distribution.

With physiological parameters, this is sufficient to distil a single firing rate even from

an input range of 250 Hz. This result was independent of the input distribution (tested:

uniform, normal and discontinuous).

4. Discussion

4.1 Summary of ideas

We make the assumptions stated earlier. A mechanism and evidence of the granular

layer computation [34] and microzone memory [37] have appeared previously.

The microzone computation is a computational effect of anatomy. This is not

incidental but a main driver of cell morphologies, the three-dimensional space they

occupy and the scaffold-like geometry of the connections they make. Communication

of cells is linear. Cells are highly adapted for faithful transmission [23-29].

Neurobiology is customised to control for biophysical and other noise. 'Although

cerebellar neurons and synapses have nonlinear intrinsic properties [62-66], these

properties are disengaged or compensated in ways that keep the cerebellar network

operating in a linear regime [29, 67-70]' [30]. Anatomy and linearity combine. The

result is a passive, unlearned computational effect of simply passing signals through

the network. At each step, data are coded in a frequency distribution. Anatomy

provides the physical hardware to randomly sample the afferent data layer. Sampling

is a facsimile of sampling with 'replacement'. Samples are passed through the

biological form of linear functions to generate the next data layer, which is in turn itself

sampled by the next layer, and so on. Unit computations in a layer are made in

parallel, continuously, so that unit outputs are constantly refreshed, i.e., analogue. The

612 result is a progressive reduction of the data range but preservation of a linear

613 relationship of the means, by the central limit theorem.

615 4.2 Context related to theory

The dominant cerebellum model is the supervised learning model, a group of models

and variants which share the idea that the cerebellum implements a learning algorithm

which teaches memories that are stored as synaptic changes [5-7, 71]. The algorithm

depends on the model. This idea has been widely influential on neural network

modelling generally. Following training, a pattern of inputs that is received at training-

modified synapses generates a learned response. Instruction is externally sourced,

hence 'supervised'. The model does not need to conjecture how mossy fibre rates and

granule cell rates are related because it is an assumption that the desired output of the

system is what the algorithm teaches.

As noted, after a promising early start [72], there have been problems with the

evidence [8-13]. It is academically possible but difficult to reconcile learning models

with linear transmission because it is the purpose of teaching in learning models to

displace a purely rate-coded response (and replace it with a learned one).

Our solution is to jettison the twin pillars of the traditional model – pattern memory

stored as tuned synaptic weights, and a learning algorithm needed to teach them –

and to propose instead that input and output of the system are connected by a chain

of rate codes. Our unit functions are linear. Units are not an analogue of synapses.

Output is an unlearned response to input rates and the mechanism of the network

computation is anatomy. Many neural network learning models share the core

assumptions of the traditional cerebellar model. We disagree with them also in these

ways.

638 4.3 Context related to function

Network modelling is normally led by a high-level idea, which can be a function or

problem for which the model is the mechanism or a solution. The order is function →

model (the third step is to consult the evidence for support). These are the first two

levels of Marr's levels of analysis. It is not contentious that a mis-step at level one is

fatal. We have the reverse problem. We have a mechanism without a function. The

high-level step is at the end, an inference from the mechanism.

The cerebellum is necessary for skilled execution and coordination of movement. We

reason that in motor control, the cerebellum may process feedback to inform motor

output. Cerebellospinal neurons provide 'a direct link between the cerebellum and

spinal substrates for motor coordination' [73]. The output pathway is reciprocated by

spinocerebellar neurons, providing a substrate for direct motor control, which spinal

locomotor networks alone cannot accomplish reliably or with finesse [74]. Neurons of

the spinocerebellar tracts are excited monosynaptically by group Ia and group Ib

afferents that originate in muscle spindles and tendon organs respectively [75].

As noted, we have proposed that the granule cell code conflates mossy fibre rate

information in long thin strips, so that a unit of granule cell information has those

minimum dimensions. We must suppose, if we are correct, that disregard of resolution

below that scale is functional. To conjecture, the many-to-one conversion of input to

output rates may distil a precise readout from individually untidy data, which include

proprioceptive data. Biologically made components, including muscle fibres, stretch

receptors and neurons, are not precision-made standard units, as models tend to

assume. Speaking generally, it is well established that neuron performance is noisy –

there is inherent biological variability from cell to cell and even of firing of the same

cell at different times in the same conditions. Muscle fibres come in different types,

vary individually, and have different tensile properties. Synaptic transmission is

randomly variable, even at the same synapse, and so on. Virtually all recordings

provide examples.

The biological solution – we suggest – is to conflate the data using a statistical effect of

random sampling. Output of a muscle is pooled for this purpose. Input to the

cerebellum is accordingly handled with disregard for resolution below a

topographically-defined scale. It may be that large cell groups in the cerebellum are a

representation of multicellular structures, which include muscles, which themselves

generate signals which together code indivisible information. Hyper-conflation of data

is a *function* of cerebellar data processing.

Correspondence between a code group in the cerebellum and a muscle may not be

limited to the input pathway. 'Motor neurons are clustered in columnar, spinal nuclei

called motor neuron pools (or motor nuclei). All of the motor neurons in a motor

neuron pool innervate a single muscle, and all motor neurons that innervate a

particular muscle are contained in the same motor neuron pool. Thus, there is a one-

to-one relationship between a muscle and a motor neuron pool.'[1] Spinal neurons

terminate profusely in a motor neuron pool. In principle, this permits a one-to-one

relationship of a cerebellar network and a motor pool, and through a motor pool, a

one-to-one relationship with a muscle.

Graded control of muscle fibres is provided because motor neurons are different sizes

(in mammals there are three sizes). Each step up in size selectively innervates more

powerfully contractile (but more exhaustible) muscle fibres. Larger motor neurons

require stronger depolarising current to fire, so weak input signals excite only small

motor neurons and the least powerful muscle fibres. Stronger signals cause more

---

[1] Reproduced with permission by McGovern Medical School, University of Texas

motor neurons to be recruited, and therefore more muscle fibres, including fibres that

contract with more force.

Credibly, these are the input and output pathways of a proprioception-motor loop and

feedback about movement is processed to inform motor output. The effect and

function of processing is to distil precision from individually imprecise data. Inform

means input rates code and *directly drive* motor output. The appearance of the

sacrifice of resolution is in fact what a collective code looks like – single signals have no

meaning for the cerebellum, which is designed to conflate data below minimum

dimensions. The cerebellum does not need to 'know' what data it is working with

because code groups are defined by topography.

Code groups are strips in our terminology, but in complex animals there is no need to

think that strips in a network all have the same width, or that networks are all the

same size. The idea that a muscle may drive motor output to another muscle is a

candidate to explain where drive comes from, if it is not central pattern generators. In

higher vertebrates, it is likely that this simple principle has acquired complex

organisation. But it is readily imagined in a simpler system.

The vertebrate cerebellum first appeared in a marine environment. It is probably not

contentious that early vertebrate swimming, before efficient teleosts, was by

undulating body movements (anguilliform). Almost the entire musculature of fish is

made up of a bilaterally symmetrical row of paired axial muscle segments, termed

myomeres. Movement through the water is produced by a repeating sequence of

muscle contractions which cause lateral body flexion. In anguilliform swimming,

alternating left and right head turns set up a herring-bone pattern of high pressure

which the animal rides by making itself an undulating shape. A sagittally-flattened tail

may contribute thrust.

Modern muscles generate information constantly, in muscle spindles, specialised

muscle fibres. Group II afferent neurons signal muscle length. Group Ia afferent

neurons fire at a rate that varies with muscle length but in addition varies

proportionally to the rate of muscle extension and (more modestly) inversely

proportionally to the rate of muscle contraction. We observe simply that these are the

makings of self-drive. Say that in a vertebrate ancestor, input to a row of

topographically-defined strips of the granular layer arose from a corresponding row of

contralateral myomeres, with neighbouring strips receiving the output of neighbouring

myomeres. Say also that these provide drive via parallel fibres to the microzone which

equally bisects the group, and that this controls motor output to the ipsilateral muscle

segment opposite the middle of the row providing input. A network is therefore the

junction of multiple input pathways and a single output pathway.[2] Passive muscle

extension might in this way be used to drive muscle contractions on the other side of

---

[2] To elaborate, neighbouring networks connected by parallel fibres send output to neighbouring myomeres. This connectivity is mirrored in the contralateral cerebellum, which has motor output to the other half of the body.

the body. Contractions would be automatically phase-locked and wavelength sensitive.

The reason that can't be mediated by spinal reflexes is that it would drive a positive

feedback loop. To address that, it is routed through the cerebellum. The solution is

parallel fibres.

Parallel fibres add a second step to the sacrifice of resolution. Strip data only retain

their integrity – i.e., are only segregated – in granule cell ascending axons, while signals

travel towards the cerebellar surface. Beyond that, after the granule cell axon divides

in two, and parallel fibre signals are randomly mixed up, within modal strata [49, 53,

76]. On the face of it, this only seems to spectacularly compound the corruption of

clean data available downstream to microzones. However, it is able to code phase and

wavelength because of topography. By averaging strip codes to generate output, the

amplitude of output is made subject to constraints that it can't exceed. The cap varies

with the number of muscle segments in a wavelength of the body and the number of

strip codes that are averaged to generate an output signal.

We can visualise this by varying input rates to each strip. In our illustration, the

number of strip codes is constant and the number of muscle segments in a body

wavelength is varied (Fig 5). Decreasing the amount of a body wavelength represented

in signals that are received as input to a network, increases output amplitude (and

sensitivity), and vice versa. Note that output is phase sensitive because it is coded in

input.

Fig 5. In theory, parallel fibres allow mutual exchange of drive between antagonist muscles without positive feedback

A Input rates to a network were varied from strip to strip, so that a plot of the mean and SD of input rates to each strip is sine-wave-like. The frequency distribution received by strips individually is normal and narrow (SD = 12.5 Hz, all strips). The central strip is indicated by the vertical dashed line. We imagined that in a locomotor cycle, a single strip receives input which varies sinusoidally at these rates. All strips receive the same cycle. Next-neighbouring strips are a step out of phase. Each pair of panels – 1 and 2, 3 and 4 and so on – represents a moment in time, but at a different phase of the cycle, advanced 90 degrees each time. Input to the network is shown in the left panel of a pair. This was passed through the network computation to derive output. Output of the computation is shown in the right panel. This represents firing of deep cerebellar neurons (DCNs), the network output cells. Output is the same in all phases. Parallel fibres perfectly dampen output. **B** The same but input rates represent 3/4 of a wavelength and the phase shift between panel pairs is 45 degrees. Output has become sensitive to cycle phase but is still dampened. **C** Input rates represent 1/2 a wavelength. Amplitude of the output range is now around twice the range in (B). **D** Input rates represent 1/4 of a wavelength. The output range increases again. Amplitude now approaches the input range (and remains phase sensitive) but can't exceed a maximum which is set by a linear relationship with input rates. END CAPTION

There are two physiological adaptations (not included in the simulation) that may

mean the response receives phase-biased weighting. Parallel fibres make about half as

many synapses distally than proximally, and swellings where contact is made contain

fewer synaptic densities [77]. As a result, there is a stronger influence of nearer

granule cells. Second, Purkinje cells receive significant granule cell ascending axon

contact [78]. Ascending axon synapses are resistant to long-term depression, and have

low uncertainty of release probabilities [79]. Accordingly, excitation of Purkinje cells

receives phase weighting from ascending axons of directly underlying granule cells.

We observe finally that this suggests how motor outputs may be coordinated by

topography of proprioception-motor loops which are connected by parallel fibres.

Immediately neighbouring microzones receive almost all the same parallel fibre input,

made up of all the same strip codes except one. Input to their next nearest neighbours

differs by two strip codes, and so on. The strip code is a frequency distribution.

Compounding strip codes as parallel fibre signals creates a frequency distribution

centred on the mean of the strip code means. So, microzones receive drive which is

automatically coordinated with their near neighbours because it has the same range

which is centred on the mean of most of the same signals. The relationship is closest

with close neighbours and weaker with further microzones, diminishing with distance.

Because firing of Purkinje cells – and therefore in turn output cells – is coded in parallel

fibre rates, output of neighbours is automatically coordinated by the relationship of

inputs.

Acknowledgements

Funding: This work was supported by the Leverhulme Trust [grant number ECF-2022-

079] (Mike Gilbert); the Swedish Research Council [grant number 2020-01468]; the

Per-Eric och Ulla Schybergs Foundation [grant number 42630]; and the Crafoord

Foundation [grant numbers 20180704, 20200729, 20220776, & 20230655] (Anders

Rasmussen).

Author contributions

M.G. researched, conceived and developed the ideas, performed the analysis, wrote

the code, prepared the figures and wrote the manuscript. A.R. provided expertise and

feedback, and reviewed and provided editorial comments on the manuscript.

REFERENCES

1. Garwicz, M., C.F. Ekerot, and H. Jorntell, *Organizational principles of cerebellar*
neuronal circuitry. News in Physiological Sciences, 1998. **13**: p. 28-34.
2. Oscarsson, O., *Functional units of the cerebellum - sagittal zones and*
microzones. Trends in Neurosciences, 1979. **2**: p. 143-145.
3. Ozden, I., et al., *Reliable coding emerges from coactivation of climbing fibers in*
microbands of cerebellar Purkinje neurons. J Neurosci, 2009. **29**(34): p. 10463-73.
4. Pedroarena, C.M. and C. Schwarz, *Efficacy and short-term plasticity at GABAergic*
synapses between Purkinje and cerebellar nuclei neurons. J Neurophysiol, 2003.
89(2): p. 704-15.
5. Albus, J.S., *A Theory of Cerebellar Function.* Mathematical Biosciences, 1971.
10(1-2): p. 25-61.
6. Marr, D., *A theory of cerebellar cortex.* J Physiol, 1969. **202**(2): p. 437-70.
7. Fujita, M., *Adaptive filter model of the cerebellum.* Biol Cybern, 1982. **45**(3): p.
195-206.
8. Welsh, J.P., et al., *Normal motor learning during pharmacological prevention of*
Purkinje cell long-term depression. Proc Natl Acad Sci U S A, 2005. **102**(47): p.
17166-71.
9. Schonewille, M., et al., *Reevaluating the role of LTD in cerebellar motor learning.*
Neuron, 2011. **70**(1): p. 43-50.
10. ten Brinke, M.M., et al., *Evolving Models of Pavlovian Conditioning: Cerebellar*
Cortical Dynamics in Awake Behaving Mice. Cell Rep, 2015. **13**(9): p. 1977-88.
11. Boele, H.-J., et al., *Impact of parallel fiber to Purkinje cell long-term depression is*
unmasked in absence of inhibitory input. Science Advances, 2018. **4**(10).
12. Isope, P. and B. Barbour, *Properties of unitary granule cell-->Purkinje cell*
synapses in adult rat cerebellar slices. J Neurosci, 2002. **22**(22): p. 9668-78.
13. Ekerot, C.F. and H. Jörntell, *Parallel fibre receptive fields of Purkinje cells and*
interneurons are climbing fibre-specific. Eur J Neurosci, 2001. **13**(7): p. 1303-10.
14. Hesslow, G., et al., *Classical conditioning of motor responses: what is the learning*
mechanism? Neural Netw, 2013. **47**: p. 81-7.
15. Rancz, E.A., et al., *High-fidelity transmission of sensory information by single*
cerebellar mossy fibre boutons. Nature, 2007. **450**(7173): p. 1245-8.
16. Ritzau-Jost, A., et al., *Ultrafast action potentials mediate kilohertz signaling at a*
central synapse. Neuron, 2014. **84**(1): p. 152-163.
17. Duguid, I., et al., *Control of cerebellar granule cell output by sensory-evoked Golgi*
cell inhibition. Proc Natl Acad Sci U S A, 2015. **112**(42): p. 13099-104.
18. Duguid, I., et al., *Tonic inhibition enhances fidelity of sensory information*
transmission in the cerebellar cortex. J Neurosci, 2012. **32**(32): p. 11132-43.
19. Walter, J.T. and K. Khodakhah, *The linear computational algorithm of cerebellar*
Purkinje cells. J Neurosci, 2006. **26**(50): p. 12861-72.
20. Walter, J.T. and K. Khodakhah, *The advantages of linear information processing for*
cerebellar computation. Proc Natl Acad Sci U S A, 2009. **106**(11): p. 4471-6.
21. Jelitai, M., et al., *Dendritic excitation-inhibition balance shapes cerebellar output*
during motor behaviour. Nat Commun, 2016. **7**: p. 13722.

22.   Park, S.M., E. Tara, and K. Khodakhah, *Efficient generation of reciprocal signals by*
843   *inhibition.* J Neurophysiol, 2012. **107**(9): p. 2453-62.

844   23.   Delvendahl, I. and S. Hallermann, *The Cerebellar Mossy Fiber Synapse as a Model*
845   *for High-Frequency Transmission in the Mammalian CNS.* Trends Neurosci, 2016.
846   **39**(11): p. 722-737.

847   24.   Saviane, C. and R.A. Silver, *Fast vesicle reloading and a large pool sustain high*
848   *bandwidth transmission at a central synapse.* Nature, 2006. **439**(7079): p. 983-7.

849   25.   Cesana, E., et al. *Novel granule cell-Golgi cell excitatory input in the cerebellar*
850   *granular layer.* in *FENS Abstr.* 2010.

851   26.   Moore, J.W. and D.E.J. Blazis, *11 - Simulation of a Classically Conditioned*
852   *Response: A Cerebellar Neural Network Implementation of the Sutton–Barto–*
853   *Desmond Model*, in *Neural Models of Plasticity*, J.H. Byrne and W.O. Berry, Editors.
854   1989, Academic Press. p. 187-207.

855   27.   Sargent, P.B., et al., *Rapid vesicular release, quantal variability, and spillover*
856   *contribute to the precision and reliability of transmission at a glomerular synapse.*
857   J Neurosci, 2005. **25**(36): p. 8173-87.

858   28.   Turecek, J., S.L. Jackman, and W.G. Regehr, *Synaptic Specializations Support*
859   *Frequency-Independent Purkinje Cell Output from the Cerebellar Cortex.* Cell
860   Reports, 2016. **17**(12): p. 12.

861   29.   Turecek, J., S.L. Jackman, and W.G. Regehr, *Synaptotagmin 7 confers frequency*
862   *invariance onto specialized depressing synapses.* Nature, 2017. **551**: p. 503.

863   30.   Raymond, J.L. and J.F. Medina, *Computational Principles of Supervised Learning in*
864   *the Cerebellum.* Annu Rev Neurosci, 2018. **41**: p. 233-253.

865   31.   DiGregorio, D.A., et al., *Desensitization properties of AMPA receptors at the*
866   *cerebellar mossy fiber granule cell synapse.* J Neurosci, 2007. **27**(31): p. 8344-57.

867   32.   Mapelli, L., S. Solinas, and E. D'Angelo, *Integration and regulation of glomerular*
868   *inhibition in the cerebellar granular layer circuit.* Front Cell Neurosci, 2014. **8**: p.
869   55.

870   33.   Galliano, E., P. Mazzarello, and E. D'Angelo, *Discovery and rediscoveries of Golgi*
871   *cells.* J Physiol, 2010. **588**(Pt 19): p. 3639-55.

872   34.   Gilbert, M. and A. Rasmussen, *The cerebellum converts input data into a hyper*
873   *low-resolution granule cell code with spatial dimensions: a hypothesis.* bioRxiv,
874   2024: p. DOI: 10.1101/2023.07.14.548987 ver 9,
875   https://www.biorxiv.org/content/10.1101/2023.07.14.548987v9.

876   35.   Hansel, C., D.J. Linden, and E. D'Angelo, *Beyond parallel fiber LTD: the diversity of*
877   *synaptic and non-synaptic plasticity in the cerebellum.* Nat Neurosci, 2001. **4**(5):
878   p. 467-75.

879   36.   Gao, Z., B.J. van Beugen, and C.I. De Zeeuw, *Distributed synergistic plasticity and*
880   *cerebellar learning.* Nature Reviews Neuroscience, 2012. **13**(9): p. 619-635.

881   37.   Gilbert, M., *Gating by Memory: a Theory of Learning in the Cerebellum.* The
882   Cerebellum, 2021. **21**(6): p. 926-943.

883   38.   De Gruijl, J.R., et al., *Inferior Olive: All Ins and Outs*, in *Handbook of the*
884   *Cerebellum and Cerebellar Disorders*, S.J.D. Manto M., Rossi F., Gruol D.L.,
885   Koibuchi N., Editor. 2012, Springer. p. 1013-1058.

39.   Ju, C., et al., *Neurons of the inferior olive respond to broad classes of sensory*
input while subject to homeostatic control. J Physiol, 2019. **597**(9): p. 2483-2514.

40.   Gilbert, M. and A. Rasmussen, *Gap Junctions May Have A Computational Function*
In The Cerebellum: A Hypothesis. The Cerebellum, 2024. **23**(5): p. 1903-1915.

41.   Jörntell, H. and C.F. Ekerot, *Receptive field plasticity profoundly alters the*
cutaneous parallel fiber synaptic input to cerebellar interneurons in vivo. J
Neurosci, 2003. **23**(29): p. 9620-31.

42.   Wilms, C.D. and M. Häusser, *Reading out a spatiotemporal population code by*
imaging neighbouring parallel fibre axons in vivo. Nature Communications, 2015.
6: p. 6464.

43.   Brand, S., A.L. Dahl, and E. Mugnaini, *The length of parallel fibers in the cat*
cerebellar cortex. An experimental light and electron microscopic study. Exp Brain
Res, 1976. **26**(1): p. 39-58.

44.   Mugnaini, E., *The length of cerebellar parallel fibers in chicken and rhesus*
monkey. J Comp Neurol, 1983. **220**(1): p. 7-15.

45.   Ramirez, J.E. and B.M. Stell, *Calcium Imaging Reveals Coordinated Simple Spike*
Pauses in Populations of Cerebellar Purkinje Cells. Cell Reports, 2016. **17**(12): p.
3125-3132.

46.   Cesana, E., et al., *Granule cell ascending axon excitatory synapses onto Golgi*
cells implement a potent feedback circuit in the cerebellar granular layer. J
Neurosci, 2013. **33**(30): p. 12430-46.

47.   Napper, R.M. and R.J. Harvey, *Number of parallel fiber synapses on an individual*
Purkinje cell in the cerebellum of the rat. J Comp Neurol, 1988. **274**(2): p. 168-77.

48.   Häusser, M. and B.A. Clark, *Tonic synaptic inhibition modulates neuronal output*
pattern and spatiotemporal synaptic integration. Neuron, 1997. **19**(3): p. 665-78.

49.   Palay, S.L. and V. Chan-Palay, *Cerebellar cortex : cytology and organization.* 1974,
Berlin: Springer.

50.   Harvey, R.J. and R.M. Napper, *Quantitative study of granule and Purkinje cells in*
the cerebellar cortex of the rat. J Comp Neurol, 1988. **274**(2): p. 151-7.

51.   Carter, A.G. and W.G. Regehr, *Quantal events shape cerebellar interneuron firing.*
Nat Neurosci, 2002. **5**(12): p. 1309-18.

52.   Llinás, R.R. and M.N. Negrello. *Cerebellum.* Scholarpedia, 10(1):4606 2015 [cited
2017 06/01]; Available from: http://www.scholarpedia.org/article/Cerebellum.

53.   Jörntell, H. and C.F. Ekerot, *Properties of somatosensory synaptic integration in*
cerebellar granule cells in vivo. J Neurosci, 2006. **26**(45): p. 11786-97.

54.   van Beugen, B.J., et al., *High Frequency Burst Firing of Granule Cells Ensures*
Transmission at the Parallel Fiber to Purkinje Cell Synapse at the Cost of Temporal
Coding. Frontiers in Neural Circuits, 2013. **7**: p. 95.

55.   Person, A.L. and I.M. Raman, *Synchrony and neural coding in cerebellar circuits.*
Frontiers in Neural Circuits, 2012. **6**: p. 97.

56.   van Kan, P.L., A.R. Gibson, and J.C. Houk, *Movement-related inputs to*
intermediate cerebellum of the monkey. J Neurophysiol, 1993. **69**(1): p. 74-94.

57.   Stell, B.M., et al., *Activation of presynaptic GABA(A) receptors induces glutamate*
release from parallel fiber synapses. J Neurosci, 2007. **27**(34): p. 9022-31.

58. Dellal, S.S., R. Luo, and T.S. Otis, *GABAA receptors increase excitability and*
conduction velocity of cerebellar parallel fiber axons. J Neurophysiol, 2012.
107(11): p. 2958-70.
59. Pugh, J.R. and C.E. Jahr, *Activation of Axonal Receptors by GABA Spillover*
Increases Somatic Firing. The Journal of Neuroscience, 2013. **33**(43): p. 16924-
16929.
60. Pugh, J.R. and C.E. Jahr, *Axonal GABAA receptors increase cerebellar granule cell*
excitability and synaptic activity. J Neurosci, 2011. **31**(2): p. 565-74.
61. Sultan, F. and J.M. Bower, *Quantitative Golgi study of the rat cerebellar molecular*
layer interneurons using principal component analysis. J Comp Neurol, 1998.
393(3): p. 353-73.
62. Hoebeek, F.E., et al., *Differential olivo-cerebellar cortical control of rebound*
activity in the cerebellar nuclei. Proc Natl Acad Sci U S A, 2010. **107**(18): p. 8410-5.
63. Ishikawa, T., M. Shimuta, and M. Häusser, *Multimodal sensory integration in single*
cerebellar granule cells in vivo. eLife, 2015. **4**: p. e12916.
64. Jahnsen, H., *Electrophysiological characteristics of neurones in the guinea-pig*
deep cerebellar nuclei in vitro. J Physiol, 1986. **372**: p. 129-47.
65. Loewenstein, Y., et al., *Bistability of cerebellar Purkinje cells modulated by*
sensory stimulation. Nat Neurosci, 2005. **8**(2): p. 202-11.
66. Person, A.L. and I.M. Raman, *Purkinje neuron synchrony elicits time-locked*
spiking in the cerebellar nuclei. Nature, 2012. **481**(7382): p. 502-505.
67. Alvina, K., et al., *Questioning the role of rebound firing in the cerebellum*. Nat
Neurosci, 2008. **11**(11): p. 1256-8.
68. Rössert, C., et al., *Model cerebellar granule cells can faithfully transmit modulated*
firing rate signals. Front Cell Neurosci, 2014. **8**: p. 304.
69. Schonewille, M., et al., *Purkinje cells in awake behaving animals operate at the*
upstate membrane potential. Nat Neurosci, 2006. **9**(4): p. 459-61; author reply
461.
70. van Vreeswijk, C. and H. Sompolinsky, *Chaos in neuronal networks with balanced*
excitatory and inhibitory activity. Science, 1996. **274**(5293): p. 1724-6.
71. Albus, J.S. *The Marr and Albus theories of the cerebellum-two early models of*
associative memory. in *Digest of Papers. COMPCON Spring 89. Thirty-Fourth IEEE*
Computer Society International Conference: Intellectual Leverage. 1989.
72. Ito, M. and M. Kano, *Long-lasting depression of parallel fiber-Purkinje cell*
transmission induced by conjunctive stimulation of parallel fibers and climbing
fibers in the cerebellar cortex. Neurosci Lett, 1982. **33**(3): p. 253-8.
73. Sathyamurthy, A., et al., *Cerebellospinal Neurons Regulate Motor Performance*
and Motor Learning. Cell Rep, 2020. **31**(6): p. 107595.
74. Parker, D., *Complexities and uncertainties of neuronal network function*. Philos
Trans R Soc Lond B Biol Sci, 2006. **361**(1465): p. 81-99.
75. Sengul, G. and C. Watson, *Chapter 8 - Ascending and Descending Pathways in the*
Spinal Cord, in *The Rat Nervous System (Fourth Edition)*, G. Paxinos, Editor. 2015,
Academic Press: San Diego. p. 115-130.

76. Zhang, W. and D.J. Linden, *Calcium influx measured at single presynaptic boutons*
of cerebellar granule cell ascending axons and parallel fibers. Cerebellum, 2012.
11(1): p. 121-31.
77. Pichitpornchai, C., J.A. Rawson, and S. Rees, *Morphology of parallel fibres in the*
cerebellar cortex of the rat: an experimental light and electron microscopic study
with biocytin. J Comp Neurol, 1994. **342**(2): p. 206-20.
78. Gundappa-Sulur, G., E. De Schutter, and J.M. Bower, *Ascending granule cell axon:*
an important component of cerebellar cortical circuitry. J Comp Neurol, 1999.
408(4): p. 580-96.
79. Sims, R.E. and N.A. Hartell, *Differences in transmission properties and*
susceptibility to long-term depression reveal functional specialization of
ascending axon and parallel fiber synapses to Purkinje cells. J Neurosci, 2005.
25(12): p. 3246-57.