1   **Computational anatomy: the cerebellar microzone computation**

2   **Mike Gilbert[1] *, Anders Rasmussen[2]**

[1]School of Psychology, College of Life and Environmental Sciences, University of

Birmingham, Birmingham B15 2TT, UK   ORCID: 0000-0003-4415-6915

[2]Department of Experimental Medical Science, Lund University, BMC F10, 22184 Lund,

Sweden   ORCID: 0000-0003-4477-369X

*Correspondence: m.gilbert.1@bham.ac.uk;

Acknowledgements

Funding: This work was supported by the Leverhulme Trust [grant number ECF-2022-

079] (Mike Gilbert); the Swedish Research Council [grant number 2020-01468]; the

Per-Eric och Ulla Schybergs Foundation [grant number 42630]; and the Crafoord

Foundation [grant numbers 20180704, 20200729, 20220776, & 20230655] (Anders

Rasmussen).

Author contributions

15   M.G. researched, conceived and developed the ideas, performed the analysis, wrote

16   the code, prepared the figures and wrote the manuscript. A.R. provided expertise and

17   feedback, and reviewed and provided editorial comments on the manuscript.

Abstract

The cerebellum is a large brain structure. Most of the mass and volume of the

cerebellum is made up by the cerebellar cortex. The outer layer of the cerebellar

cortex is divided functionally into long, thin strips called microzones. We argue that the

cerebellar microzone computation is the aggregate of simple unit computations and a

passive effect of anatomy, unaided and unlearned, which we recreate *in silico.* This is

likely to polarise opinion. In the traditional view, data processing by the cerebellum

(stated very briefly) is the effect of learned synaptic changes. However, this has

become difficult to reconcile with evidence that rate information is linearly conserved

in cerebellar signalling. We present an alternative interpretation of cell morphologies

and network architecture in the light of linear communication. Parallel fibre synaptic

memory has a supporting role in the network computation.

Keywords cerebellum; theory; model; computation; anatomy; microzone; network;

code

Lay summary

It is not fully understood how the cerebellum represents information and uses it to generate motor outputs, and how outputs are coordinated to control movement. We model the cerebellar network computation, with a focus on the microzone computation, the middle layer of the network. The main proposal is the contentious idea that the core computation is a passive and unlearned effect of neuroanatomy, unit functions are linear, and the computation is mathematically unsophisticated.

To provide functional context, we propose that the topography of input and output of cerebellar motor circuits could be a functional extension of modular cerebellar wiring which automates motor sequences. To illustrate how that could work, we present a simple model of anguilliform (eel-like) swimming.

1. Introduction

Most of the mass and volume of the cerebellum is made up of the cerebellar cortex,

which is divided into an inner and outer layer, the granular layer and molecular layer.

The molecular layer is divided functionally into long, thin strips called microzones [1-3].

Microzones lie at right angles to the main portion of the axons of granule cells, called

parallel fibres, which themselves lie parallel to each other and the cerebellar surface. A

microzone is populated by a few hundred Purkinje cells. Purkinje cells receive contact

in passing from parallel fibres and are the sole outputs of the cerebellar cortex. Firing

rates of the output cells of the cerebellum are bidirectionally sensitive to Purkinje cell

rates [4]. This provides the main control of output cell rates.

At a mid-range estimate, a microzone is ~150 μm wide and 15–20 mm long. Purkinje

cells are sagittally flattened in the plane orthogonal to parallel fibres but present a

large target – an estimated 350,000 parallel fibres pass through the dendritic arbour of

a single Purkinje cell [5, 6], and many millions through a microzone. Parallel fibres have

a range of ~3 mm in both directions from the granule cell soma [7, 8], so that a

microzone receives parallel fibre input from both sides, from a large region. A

microzone accordingly defines a functional network made up of the microzone (the

middle layer of the network), the region of the granular layer from which it receives

parallel fibre input (the input layer of the network), and the nuclear group it has

output to (the output layer of the network).

There is growing evidence that rate information is linearly conserved in signalling

between cerebellar neurons [9-12], including synaptic signalling in the molecular layer

[13-16], and that cells are adapted for faithful transmission [17-23]. Firing rates linearly

code behavioural and task metrics 'at all levels of the cerebellar circuit' [24 p.239,

citing 30 references]. Neurobiology is highly adapted to control for biophysical and

other noise. There are cerebellar examples of this, where cell-to-cell communication is

adapted to conserve fidelity of linear transmission of rate information and control for

other variables [9-12, 17, 18, 21-23, 25-27].

This paper is an attempt to account for detailed neuroanatomy, network organisation

and linear signalling. The main proposals are that i) the core computation is a passive

and unlearned effect of neuroanatomy; ii) the network is the physical form of

functionally-layered linear unit functions; and iii) the computation is mathematically

unsophisticated. Unit functions are attributed to identified anatomical hardware,

connected by known anatomy, and evidenced. In this form, we are able to model large

cell networks in high computationally-relevant definition. Model parameter values

represent coded data and not biophysics. Firing rates are an example, where complex

biophysics code a value. We claim that this step allows the model to be biophysically

simplified without simplifying the computation.

To provide functional context, we incorporate the network computation in a simple

model of anguilliform (eel-like) swimming, a candidate proto-vertebrate wiring plan.

We discuss learning in the Discussion section. We propose that parallel fibre synaptic

plasticity is in a complementary role to the network computation, with the function of

making a pattern-blind, whole-microzone gain adjustment.

While there is a computational element, this is not primarily a computational paper.

Many neural network models present a computational mechanism that executes the

function of the network. The function is a proposal and becomes an assumption of the

model. In practice, these steps may be conflated, but the direction of the argument is

function → mechanism (the first two levels of Marr's three levels of analysis). We work

in reverse. We infer a mechanism from the evidence and then infer function from the

mechanism.

This is significant. In our approach, the simulation provides a test of the performance

of the system. The purpose is to demonstrate proof of concept, as opposed to being

the form of a mathematical generalisation. 'Proof' does not refer to mathematical

proof, but a practical demonstration, using examples.

We do not propose or claim to be mathematically interesting. The sophistication of the

system lies in its biological implementation, isolating the relationship of parameters

that code information and controlling for other variables, including biophysical noise. A

note on terminology: a layer of a network has a different meaning to layers of the

cerebellar cortex. For example, a microzone, the middle layer of a network, is a

division of the molecular layer, a stratum of the cerebellar cortex.

Sections 2 and 3 describe the derivation of parallel fibre input to a microzone. Section

4 describes the proposed microzone computation. Section 5 describes a simple system

which incorporates the network computation.

2. A network of linear functions

2.1 Computational anatomy

We have previously simulated the granular layer mechanism which converts mossy

fibre signals into granule cell and therefore parallel fibre signals [28]. In the granular

layer model, the granular layer is divided functionally into long thin strips – 'ranks' –

which mirror microzonal organisation. We envisage that anatomy is the physical form

of a network of computational units. A unit takes inputs and returns output. Inputs and

output represent values (that is, numbers). Physically, a unit is the site of biophysical

events which relate inputs and output – the unit function. Unit functions are linear –

output is a linear function of inputs. A review of evidence of linear signalling in the

granular layer appears in Supplementary materials.

The physiological form of a unit depends on the layer/unit type. Units are physically

intermingled in both a rank and in a microzone, but organised functionally in layers.

The outputs of a layer provide the data sampled (i.e., received as inputs) by units in the

next layer. Computations run continuously, so unit outputs are constantly refreshed –

i.e., analogue.

All layers of unit functions span the network from side to side (because the population

of a layer occupies either the volume of a rank or a microzone). Information is coded

collectively in the concurrent outputs of the whole population of a layer. It is coded as

a frequency distribution, so that a whole layer codes information that is both

indivisible (cannot be subdivided into smaller units of information) and yet coded in

any random sample of unit output values.

134   2.2 How are units connected?

Units are connected by anatomy, so we know what the connections are, because

anatomy has been reported in detail. The number of inputs to a unit is usually

anatomically randomly variable in a reported range. In some cases, we simulate that

by generating the number with a distributed probability, and in others we use the

mean.

It is unnecessary to know which individual cells contact each other because

(biologically and in the model) contact is at random inside topographically defined

dimensions. This has the asset for modelling that a target population can be

represented by randomly sampling data received as input to a location.

Biologically, a location is a volume with spatial dimensions. *In silico*, we can represent

that as a population of unit functions. We know from anatomy (or can derive) the size

of unit populations, population ratios, and convergence and divergence ratios. These

provide model parameters and the values they take.

2.3 How does that 'look' in the granular layer?

Stated in short form, the input layer of a network receives input from what may be

around 80,000 mossy fibres. Inputs are passed through a chain of iterative random

samplings (simplified: Golgi cells sample mossy fibre rates and glomeruli sample Golgi

cell rates) to calculate inhibition of granule cells at each of 700 glomeruli per field, in

4,100 fields (into which we nominally divide the input layer of the network; Fig 1).

Outputs are passed to the granule cell computation to derive the independent

outcome of comparing inhibition and randomly sampled excitation of each dendrite of

around 9,000 granule cells per field, field by field, to obtain the number that fire in

each field, and the rates they fire at on a normalised scale. The proportion of granule

cells which fire is well-predicted by probability, which is controlled by the relationship

of inhibition and the mean of mossy fibre rates. At collective level, granule cell firing

rates are well-predicted by mossy fibre rates, and have a focused range centred on the

mean.

162 2.4 In this paper, the form of the granule cell/parallel fibre code

163 is an assumption

We use the granular layer mechanism to generate parallel fibre input to the microzone

computation. Previously, we modelled a single rank [28]. In this paper, we extend that

to the larger region (around 40 ranks) that supplies parallel fibre input to the

microzone that equally bisects the region viewed from the cerebellar surface.

In this paper, the workings of the granular layer mechanism are effectively an

assumption. Recoding in the granular layer generates a constantly changing pattern of

active parallel fibres but with invariant features, regardless of the information it

contains. Information is coded at collective level, in the joint activity of cells which are

active at the same time. More exactly, it is coded in the frequency distribution of firing

rates where active cells intersect a sagittal plane. Firing rates are randomly distributed

among active parallel fibres and active parallel fibres are a random subset of the

general population. Accordingly, any random sample of firing rates is (at any moment)

a random sample of the same distribution. Functionally, a microzone is the

physiological form of a sagittal plane. So, because all sectors of a microzone receive a

large number of active inputs at any time, all sectors receive the same code at the

same time.

2.5 The key importance of mossy fibre morphology

2.5.1 MOSSY FIBRE TERMINAL MORPHOLOGY, 'RANKS' AND 'FIELDS'

Mossy fibre morphology has a computational effect, we submit. Mossy fibres give rise

to sagittally-extending collaterals, the direction of the long axis of microzones.

Collaterals branch terminally, and each branch ends in a cluster of terminals, so that a

single cell terminates in a spaced-out row of terminal clusters which are always lined

up in the same direction (parallel to microzones but in the granular layer) [29-32].

Clusters are separated by a randomly variable distance typically not less than 200 µm

[29, 30] measured from edge to edge of clusters (so they are separated by a greater

distance measured from centre to centre). Terminals are small compared to the

average volume occupied by a cluster, and intimately and randomly intermingled with

terminals of other mossy fibres.

Accordingly, mossy fibres do not terminate at a point but a region with dimensions:

the field occupied by a terminal cluster. A single mossy fibre gives rise to several

terminal branches and each of those to several terminals, such that a sagittal strip of

the granular layer receives multiple copies of every signal at each of multiple locations.

The dimensions which enclose a cluster are variable (partly because it depends on the

number of terminals in a cluster), averaging 200 μm sagittally x 150 μm mediolaterally

[30]. A sagittal row of 100 cluster fields ('fields') has the dimensions of a rank viewed

from the cerebellar surface (Fig 1). Fields are nominal, a modelling device. Ranks are

functional divisions – the minimum dimensions at which the cerebellum makes any

effort to tell signals apart, we submit.

2.5.2 TERMINAL BRANCHING

Mossy fibre terminal morphology has important computational significance (for a full

discussion see [28]). Terminal branching of mossy fibre collaterals simulates

independent random sampling, by fields, of input rates to a rank, also called sampling

with 'replacement'. If each mossy fibre could terminate in only one field, a mossy fibre

'sampled' by one field clearly cannot also be sampled by any others. So, samples are

not independent. Termination by a single mossy fibre in a random number of terminal

branches, at multiple locations, removes that restriction.

In theory, sampling with replacement would mean any number of fields can receive

the same signal, up to all of them. Clearly this is not true – in reality there is an

anatomical limit. However, even if there was real replacement, the number of repeat

samplings is constrained by a probability distribution. If the number of terminal

branches per mossy fibre has the same probability distribution, branching has the

same result as if firing rates received by each field in a rank were in fact a random

sample with replacement of firing rates received by the rank as a whole.

Expanded sagittal X-section

A

Granule cell

B

~9,000 granule cells per field,

350,000 per file

File: a *nominal* division of the granular layer

Ranks are functional divisions

6 mm – a microzone receives parallel fibre input from a range of 3 mm on each side

41 'ranks' x 100 'files' = 4,100 'fields'

20 mm – microzones vary, perhaps widely

C

Expanded coronal X-section

Central microzone contains ~4 x 100 PCs and ~8,000 stellate cells

Parallel fibre, granule cell axon, 3mm range

~180 mossy fibres innervate a field

~80,000 supply the whole of the chequered pink area, the input layer of the network

400 Purkinje cells

Output of a microzone is sent to a smaller group of cells in the deep cerebellar nuclei. ~50 cells carry the output of the network

As many as 35m parallel fibres innervate a microzone, 350,000 each file

Fig 1 Schematic of a network

Schematic of a region of the flattened cerebellar cortex made up of the input (pink) and

middle (green) layers of the cerebellar network, showing relative dimensions of

functional and nominal subdivisions. **A** Light blue: molecular layer; pink: granular layer;

chequered pattern: files in cross-section; dark blue blocks: the dimensions of the volume

occupied by an average mossy fibre terminal cluster. In reality, clusters do not fit neatly

into fields or files, and terminals are randomly intermingled with terminals of other

mossy fibres. There are around 700 terminals in total per field; a single mossy fibre

contributes less than 1%. **B B**oth the molecular and granular layers are functionally

divided into long, thin sagittal strips defined by the way input maps to the cerebellum.

Granular layer strips are 'ranks'. A microzone receives parallel fibre input from around

40 ranks, 20 each side. 'Files' are nominal divisions, simply a modelling device to quantify

the granular layer computation and in doing so to derive the number of active granule

cells that provide input to a microzone sector, and the rates they fire at. Only a single

microzone is shown. Each microzone defines its own network, so the input layers of near

neighbouring networks overlap very greatly. **C** The output of a microzone is the

continuous firing of Purkinje cells which converge onto an associated nuclear group.

2.5.3 TERMINAL CLUSTERING

Mossy fibre terminal clustering simulates independent random sampling, by individual

cells, of input rates to a field. The principle is the same as with terminal branching, but

at local scale. Mossy fibre terminal clustering, and random intermingling with

terminals of other mossy fibres, means that contact on any particular cell does not

mean other cells cannot also receive the same signal. Again, to approximate

independence, it is only necessary for the number of repeat selections of a signal to

have the same probability distribution as if there was actual replacement.

2.5.4 THE COMBINED EFFECT OF TERMINAL BRANCHING AND CLUSTERING

The combined effect is a biological facsimile of simultaneous random sampling with

replacement by *individual* granule cells and Golgi cells of firing rates received as input

to *a rank*, notwithstanding that single cells individually receive fixed contact from a

tiny fraction of mossy fibres. Note that it is the population of mossy fibre *rates* that are

hypothetically sampled with replacement, and not *signals.* The population of signals

includes copies of rates because of branching and clustering.

As a result, all granule cells in a rank randomly sample, at any time, the same

distribution, and information represented by the output of a rank – i.e., granule cell

rates – is coded in any random sample of signals [28]. Since all sectors of a microzone

receive parallel fibre input from all the same ranks, microzone-grouped cells all

simultaneously randomly sample, with simulated replacement, the whole population

of parallel fibre rates received by that microzone. Independent random sampling by

topographically-defined cell groups is the cornerstone of the network computation.

2.5.5 WHY IS THAT KEY?

Say a population of values (with any frequency distribution) is randomly sampled with

replacement. There are *n* samples, sample size *m*. If we take the mean (or sum) of each

sample, and *n* is large, the sample means have a normal or near-normal distribution,

by the central limit theorem. (This works even if *m* is a range.) The new distribution is

centred on the same mean as the sampled distribution, but narrower.

Large values of *n* and *m* give a nearly-normally-shaped new distribution. Modest values

give a normal-tending distribution. However, if the new distribution is then itself

independently randomly sampled, and the process is repeated, the results become

more focused, and the relationship of the means of successive distributions is well

preserved. We will argue that the cerebellar microzone provides (part of) the physical

hardware for the cerebellar network to distil output from input in this way, noting that

at each step the relationship of the means is adjusted by a constant.

A precis of the ideas in this section and a schematic discussion of merits and limitations

appears in Figure 2.

2.6 Derivation of mossy fibre rates used in the simulation

We use the simulation to test if an anatomical computation is feasible. For that, it is

necessary to generate input to the system that reflects mossy fibre terminal branching

and terminal clustering, and also to generate input to each field which reflects the fact

that cluster size is randomly variable and most clusters straddle field limits, so that

some terminals fall inside and some outside. The derivation of mossy fibre rates

received as input to a rank and to fields is described in Supplementary materials.

THE PRINCIPLES

RANK THE GRANULAR LAYER IS DIVIDED FUNCTIONALLY INTO SAGITTAL STRIPS – 'RANKS' – THAT MIRROR MICROZONES.

NETWORK A NETWORK IS MADE UP OF A MICROZONE, THE RANKS THAT PROVIDE PARALLEL FIBRE INPUT TO IT, AND THE DEEP NUCLEAR GROUP IT HAS OUTPUT TO.

CODE GROUPS BOTH RANKS AND A MICROZONES ARE POPULATED BY COMPUTATIONAL UNITS. UNITS ARE INTERMINGLED PHYSICALLY BUT ARE ORGANISED FUNCTIONALLY INTO LAYERS.

LINEAR UNITS OUTPUT OF A UNIT IS A LINEAR FUNCTION OF INPUTS. THE PHYSICAL FORM OF A UNIT AND THE FUNCTION DEPEND ON THE UNIT TYPE/LAYER.

CODE INFORMATION IS CODED (AT ANY MOMENT) COLLECTIVELY IN THE OUTPUTS OF A CODE GROUP.

ANALOGUE DATA AT MOST STEPS, THE BIOPHYSICAL FORM OF INFORMATION IS A SUSTAINED STATE.

FAUX SAMPLING ANATOMY SIMULATES RANDOM SAMPLING BY A SINGLE UNIT OF THE WHOLE OF OUTPUT OF THE AFFERENT CODE GROUP. AT MOST STEPS, SAMPLING IS WITH SIMULATED 'REPLACEMENT'.

ITERATIVE SAMPLING ITERATIVE SAMPLINGS (EACH CODE GROUP SAMPLES THE OUTPUT OF THE ONE BEFORE) CONVERT INPUT TO THE NETWORK INTO V. NARROWLY FOCUSSED OUTPUT

LEARNING PARALLEL FIBRE SYNAPTIC PLASTICITY PROVIDES A GLOBAL GAIN ADJUSTMENT TO THE NETWORK COMPUTATION.

FIG 2

WE INFER A MECHANISM FROM THE EVIDENCE AND FUNCTION FROM THE MECHANISM. THIS HAS LED TO SOME UNEXPECTED CONCLUSIONS. THERE ARE PROS AND CONS OF THIS APPROACH.

FOR

THE PROPOSALS ATTEMPT TO ACCOUNT FOR DETAILED ANATOMY This includes cell morphologies, network geometry, cell ratios, the shape and size of dendritic and axonal territories, convergence and divergence ratios, contact at random by all cell types on a subset of in-range targets, microzonal organisation, synaptic adaptations, and multiple copies of some signals.

LINEAR UNITS We interpret neurophysiology as a network of linear functions. This provides a functional explanation of evidence of linear relationships between cerebellar cell firing rates, firing rates and behavioural metrics, and that synapses are adapted to conserve rate information.

SOLUTION TO HIGH-LEVEL PROBLEMS? The ideas may explain: 1. How the cerebellum processes data 'blind'. 2. Coherent firing and functional definition of large cell groups. 3. How variable performance of non-standard biological units is mitigated.

RISK MANAGEMENT In the classic Marr 'three levels' approach, the model/mechanism is justified by an assumption 'up front' about the function attributed to it. If the assumption is wrong the model fails. Inferring function from the mechanism does not involve that risk.

AGAINST

THE PROPOSALS DO NOT ACCOUNT FOR SOME OTHER EVIDENCE Giving precedence to anatomical evidence means the proposals can't also explain evidence of multi-site synaptic plasticity (except parallel fibre synaptic learning/memory).

SIMPLISTIC? Brain computations are expected to be powerful. Data are represented by complex biophysical variables. Is it credible that computations are straightforward?

A CONFLICT OF EVIDENCE It is difficult to reconcile evidence of linear signalling with (some) evidence of synaptic learning. Giving priority to either makes it difficult to account for the other.

DO THOSE PROBLEMS NEED SOLVING? An alternative explanation is that the cerebellum is sighted by external supervision, information is coded at single cell level and not in joint activity of cell groups, and biological units perform well.

BIAS CONFIRMATION? Both directions of analysis risk being self-fulfilling. Using inferred function as a supporting argument is circular.

3. Regulation of parallel fibre activity

3.1 A case for sparse code (small fraction)

To simulate the computational effect of random sampling of parallel fibre signals, we need to know or derive sample sizes (the number of inputs to each type of unit), and therefore in turn what fraction of parallel fibres is active at any time. There is a long-standing idea that the fraction of parallel fibres that are active at any time is regulated [33], fixed [34], and low (termed sparse code). We disagree with the theoretical reasons, but agree the code is sparse, on these grounds.

1) Low density is stable (by our calculation – Fig 3). Randomly fluctuating density would be noise. Stability may be enhanced by feedback via granule cell ascending axon contact on Golgi cell basal dendrites [19] and parallel fibre contact on Golgi cell apical dendrites, acting partly through gap junctions [35].

2) Low density is energy efficient. Granule cells are very numerous, and they are not myelinated (so spike propagation is costly).

3) The reported number of coincidentally active inputs to a stellate cell is in single figures [36], consistent with low density of co-active parallel fibres (see Table 1).

4) It is thought that the pattern of active and inactive parallel fibres, and the pattern in which firing rates are distributed among active cells, are decorrelated – effectively,

quasi-randomised. Targets receive contact from a subset of passing parallel fibres –

Purkinje cells receive contact from 1 in 2 [37] and stellate cells from something like 1 in

80 (section 4). Low density enhances practical decorrelation: if decorrelation is not

perfectly random – as it may not be [38] – low density increases the randomness of the

pattern of synaptic activations received by parallel fibre targets.

For these reasons, we calibrated the granular layer model to convert input to the

system into a low fraction of active granule cells. We chose 0.5% active because it has

support from (as far as we know) the only estimate of the number of co-active inputs

to a stellate cell (per section 4). We suspect it may be higher – perhaps 1% – but a very

low fraction is a more rigorous test of our proposals (because sample sizes are

smaller).

310 3.2 Simulated regulation of the fraction of active parallel fibres

As noted, we extended the granular layer model to simulate recoding of input to a

network. One of the outputs of the model is the number of active granule cells per unit

volume of the granular layer. We use fields as the unit of volume. A mediolateral row

of fields is a 'file' (Fig 1). The input layer of a network is accordingly crossed in one

direction by 41 ranks (20 either side of the middle microzone and one directly

beneath) and at right angles by 100 files, making 4,100 fields.

The number of active granule cells in each field gives the number in each file. The

number in a file is equal to the number of active parallel fibres in that file that

converge on the middle microzone. The results are shown in Figure 3. Local regulation

of the number of active granule cells in each field in a rank (Fig 3 columns 2 and 5)

becomes still tighter regulation of the number of active parallel fibres in a file (Fig 3

columns 3 and 6), a statistical effect of cerebellar cortical architecture. Density is

independent of the frequency distribution of mossy fibre rates (Fig 3 columns 1 and 4).

At low density of active granule cells, the regulated density of active parallel fibres is

very strict.

Physiological regulation may be still stricter, that is, the number of active granule cells

in each field may not vary as much as it does in the simulation. Local feedback via the

granule cell ascending axon [19, 39] may condense the physiological range of field-to-

field variation. Feedback via parallel fibres [34, 35] may in addition reduce the range of

file-to-file variation. These are not represented in the simulation.

Fig 3. Recoding in the granular layer regulates the fraction of active parallel fibres.

Columns 1 and 4. Firing rates received as input to the granular layer region that supplies parallel fibre input to a microzone. Data represent inputs to a single rank, selected at random. Inputs to each rank were independently randomly generated with a uniform and discontinuous distribution, respectively. The method of generating mossy fibre rates received by each field (in all ranks) is described in Supplementary materials. Data represent a short period, effectively a moment of time. **Columns 2 and 5.** The

So, by our calculation, data are not corrupted by noisy variation of the fraction of

active parallel fibres, because the fraction is strictly regulated. The primary regulator is

the granular layer computation. This has functional significance in two ways. The first is

that it means the distributed probability is unchanging of the number of active inputs

received concurrently by parallel fibre targets. The second is that it means Purkinje

cells receive a fixed number (but high turnover) of active inputs, so that learning can

influence Purkinje cell firing by controlling the number of signals that Purkinje cells

receive (vs learned changes of graded synaptic weights). We develop this idea in the

Discussion.

4. The microzone computation

4.1 Stellate cells

4.1.1 SAMPLE SIZE

Purkinje cells receive inhibition from interneurons, stellate cells (some vertebrate

groups have a second phenotype not included here). The stellate cell territory is

sagittally flattened in the plane orthogonal to parallel fibres. Stellate cells form planar

networks [40] that occupy the space (about 40 μm) between Purkinje cells [41]. The

stellate cell dendritic territory varies with the depth of the cell body in the molecular

layer. For superficially located ('outer-level') cells it is around 80 x 80 μm. At deeper

level ('inner level') it is larger, around 120 x 120 μm [41 pp.217-221]. The number of

parallel fibres which pass through an inner-level stellate cell territory is given by $a/((b$ x

$c)/120^2)$, assuming parallel fibres are uniformly distributed, where a is the number

that pass through a Purkinje cell territory (350,000 [6]), and b and c are the dimensions

of the Purkinje cell territory in the same plane (200 x 300 μm), giving ∼84,000. If 0.5%

are active in the general population, around 420 of those are active.

It has been estimated that a stellate cell receives contact from several hundred parallel

fibres [41, 42], which we take as 1,000. Working with these estimates, 1 in 84 make

contact (i.e., 1,000 out of 84,000), so the probability that an active cell makes contact

378   is 1/84 = ~0.0119. Accordingly, contact by x (out of 420) active cells has a probability

379   given by

380
$$\frac{420!}{x!\,(420-x)!} * 0.0119^x * (1-0.0119)^{420-x}$$

Probabilities are shown in Table 1 for the range x = 0 to 12, where x is the number of

active parallel fibres that make contact and p is the probability of that number. The

distribution of probabilities is in good agreement with the modest number – single

figures – suggested by 'two to eight substantial EPSPs' [36 p.9628].

Table 1 The probability, p, that a stellate cell receives contact from x active parallel

fibres

x	0	1	2	3	4	5	6	7	8	9	10	11	12
p	.0066	.0331	.0836	.1403	.1762	.1765	.147	.1047	.0651	.0359	.0178	.008	.0033

Caption. Table 1 shows the distributed probabilities of the number of active inputs to

an inner-level stellate cell.

4.1.2 SIMULATED SAMPLING AND INTEGRATION

We notionally divided the middle layer of a network (the middle microzone in Fig 1)

into 'sectors'. Sectors are the same size as fields viewed from the cerebellar surface so

that a sector and the field below it together form a column of the cerebellar cortex.

Purkinje cells and stellate cells in a sector randomly sample parallel fibre rates in the

file that intersects a microzone at that location. (Like fields, sectors are notional and do

not represent functional segregation of data.)

The granular layer model returns the firing rates of granule cells in each field, and

therefore the frequency distribution of parallel fibre rates in each file, and accordingly

rates received by each sector. There are an estimated 16 stellate cells for each Purkinje

cell [43], meaning 80 stellate cells per sector (there are four Purkinje cells per sector so

five layers of stellate cells are needed to inhibit them from both sides). Sample size

was randomly generated with the probabilities in Table 1. This gave the number of

active inputs, and input rates, to each stellate cell in each sector.

Molecular layer interneurons, which include stellate cells, reflect 'granule cell input

with linear changes in firing rate' [15 p.6]. We assume that stellate cell firing is a linear

function of the mean of input rates, which we take as the mean itself (the sum works

as well). As the physiological function is not known, and our purpose is only to

demonstrate a statistical relationship of data layers (i.e., the outputs of units functions

functionally organised into layers), and not to calculate actual firing rates, we use a

normalised scale. The results are shown in Figure 4.

In this proposal, each stellate cell is the physical form of a notional unit, and the

afferent data set is the frequency distribution of parallel fibre firing rates. Physically,

and in the simulation, units are limited to sampling only signals in that file. Few parallel

fibres are active at any time. But the high turnover of active cells means that sample

415 size varies with time and, at any time, the number of active inputs to a unit has a

416 distributed probability. The simulation generates sample size for each cell with the

417 Table 1 probabilities. Firing rates received by a stellate cell are an independent random

418 sample. The unit computation is a linear function. In physiological conditions, inputs

419 are computed in parallel by all units (in all sectors) concurrently. All (at any moment)

420 effectively sample the distribution of parallel fibre rates received by the whole

421 microzone (because all sectors receive the same distribution), notwithstanding that

422 they all receive contact from different parallel fibres. The outputs – each an analogue

423 signal – become the data set sampled by the next layer of units.

424 The fast turnover of active parallel fibres means that units receive input at time-

425 varying rates notwithstanding that granule cells typically fire in short bursts [15, 44, 45]

426 at a rate which (as far as we know) does not vary in a burst. Accordingly (and because

427 stellate cells fire intrinsically), unit outputs are analogue.

428 Stellate cell data generated by the simulation and shown in Figure 4 are the population

429 of parallel fibre rates received as input to 16 stellate cells per sector (the number that

430 sit between two Purkinje cells), so around 7,200 values (Fig 4D and N), and the outputs

431 of the unit function used by the simulation to represent those cells (1,600 values, Fig

432 4E and O). It forms part of the proposal that a stellate cell is the physical form of the

433 unit function.

Fig 4. The microzone computation.

SC: stellate cell; PC: Purkinje cell; compt: PC dendritic compartment; mf: mossy fibre; pf: parallel fibre; 'rank' and 'file' are defined in Fig 2. **A** Mean and SD of mossy fibre input to 41 ranks of the granular layer. Ranks are 150 microns wide and together make up the region supplying parallel fibre input to the microzone that equally bisects it, indicated by a dashed grey line. The morphology of mossy fibres means that all locations of a rank receive a large random sample of input rates to the whole strip. **B** Frequency distribution of mossy fibre rates received by a single file, 200 microns wide. Files cross ranks at right angles. Granule cell firing rates in a file are accordingly parallel fibre rates received by the middle microzone. As files cross all ranks, all middle-microzone sectors receive the same distribution of mossy fibre rates. There are 100 files. **C** Parallel fibre rates in a randomly selected file. The frequency distribution of parallel fibres rates is the same in all files. This is always true, and does not depend on ranks receiving the same mossy fibre distribution. **D** Parallel fibre rates received as input to 16 stellate cells per sector (out of 80, for illustration), obtained by randomly sampling parallel fibre rates (16 times per file, generating sample size randomly with Table 1 probabilities). The y axis range is 60 normalised Hz, down from a mossy fibre range of 250 Hz. **E** Outputs of the stellate cell unit functions which represent the cells in D. The program derives input to, and output of, all stellate cells, generating 8,000 data points, but the data are thinned for presentation. **F** We notionally divided the Purkinje cell dendritic arbour into 16 compartments per cell. Each compartment randomly samples a subset of outputs of the

stellate cell unit function which represents the network of stellate cells afferent to that

cell. Data are the sample means, shown for a single Purkinje cell per middle-microzone

sector (so there are 1,600 values, 16 compartments x 1 Purkinje cell x 100 sectors). **G** A

Purkinje cell receives inhibition from two local stellate cell networks, one on each flank.

Step F was repeated, and the data averaged for each compartment. **H** Mean of the

compartment means calculated for each Purkinje cell in the middle microzone (so 400

data points, 4 Purkinje cells per field x 100 fields), representing somatic integration of

dendritic compartment data. **I** The same data but with the y axis range shortened to 10

normalised Hz. **J** The output of a microzone is channelled down onto a smaller cell group

in the deep cerebellar nuclei, which includes the output cells of the network. Each

output cell receives input from a random sample of 30–50 Purkinje cells [46]. There are

~50 output cells in a nuclear group. Data are the sample means. **K–T** We repeated (A)–

(J) but with normally distributed mossy fibre rates (SD 40), with the same result. We also

obtained the same results (not shown) with a discontinuous mossy fibre distribution

(range 50–300 Hz). END CAPTION

There is direct evidence of a linear relationship of granule cell input and stellate cell

rates [15] but as far as we know not confirmation, or data about the sensitivity or

latency of the response. However, fidelity of transmission may be assisted. GABA

spillover from stellate cell synapses enhances parallel fibre synaptic transmission to

stellate cells [47, 48]. 'It is well established that parallel fibers express functional

GABA$_A$Rs, that their activation depolarizes the axon, and that they increase release

probability at parallel fiber synapses' [49 p.16924, 50]. High release probability hones a

rate-proportional response.

4.2 Purkinje cells

4.2.1 PURKINJE CELL DENDRITIC COMPARTMENTS RECEIVE INNERVATION FROM AN

INDEPENDENT RANDOM SAMPLE OF STELLATE CELLS AFFERENT TO THE CELL

Around 350,000 parallel fibres pass through the dendritic field of a single Purkinje cell,

of which one in two make contact, normally at a single synapse (sometimes two,

average 1.24) [5]. Purkinje cells are severely flattened in the plane orthogonal to

parallel fibres and to the cerebellar surface, filling the molecular layer vertically, so

that parallel fibres pass through by the shortest distance.

A microzone is ~100 Purkinje cells long and 4 wide (400 is consistent with reported

convergence and divergence ratios onto 50 excitatory nuclear projection cells [rats:

46]. Purkinje cells are inhibitory and fire spontaneously [40, 51-53] at a rate adjusted

by simultaneous direct excitation from parallel fibres and inhibition by stellate cells.

In the simulation, a field contains 4 Purkinje cells. A Purkinje cell receives contact from

stellate cells from both of its flat sides, ~16 each side, and a smaller number of stellate

cells in the two sagittally-neighbouring fields. Lateral inhibition from neighbouring

fields is from a smaller number of stellate cells because not all of them extend their

main axon far enough to reach, and some of them extend their main axon the wrong

way. We represent those conditions as variables in the simulation. The vertical range

of axon collaterals and the horizontal range of the main axon both increase with the

depth of the stellate cell body in the molecular layer [41, 54]. Because of their more

limited horizontal range, outer-level stellate cells are excluded in the simulation from

contacting Purkinje cells in neighbouring fields.

The Purkinje cell arbour is large and branches profusely, and stellate cell contact is

dispersed across it. The local postsynaptic effect is therefore a response to local inputs.

We divide the Purkinje cell arbour nominally into sixteen 50 x 75 μm compartments.

For morphological reasons, input to a compartment is not confined to (or even

necessarily always received from) the nearest stellate cells, but each compartment is in

range of most stellate cells. As a result, each compartment receives contact from a

near approximation of a random subset of stellate cells afferent to a Purkinje cell.

It is unknown how many stellate cells innervate a Purkinje cell compartment. We take

it as 6, around 20% of the afferent network, and that local dendritic polarisation is a

linear function of the mean firing rate of stellate cells afferent to a compartment.

Sampling by a compartment is independent: innervation of a compartment by a

stellate cell does not exclude innervation by the same cell of other compartments.

Physically, contact is limited by stellate cell shape and range to a small group of local

cells. However, the stellate cell network afferent to a Purkinje cell could in theory be

made up of any random sample of stellate cells in the same microzone, with the same

result at population level.

4.2.2 SIMULATED SAMPLING AND INTEGRATION

A notional compartment is a computational unit. Each compartment randomly samples

a subset of local stellate cells. For this purpose, stellate cells are represented by

outputs of the stellate cell unit function. A subset is made up of 16 of the 80 units that

represent stellate cells in that sector, plus up to a further 8 in each of the sagittally

neighbouring sectors. Compartment membrane voltage is a linear function of stellate

cell rates afferent to that compartment. In its physical form, a compartment

continually recalculates inputs (the simulation represents a moment of time). Dendritic

signals are a sustained state, we propose, mitigating a randomly variable effect of

spatiotemporal integration of discrete signals – a noise reduction strategy.

To declutter the data, we show unit outputs (representing Purkinje cell dendritic

compartments) for a single Purkinje cell per sector (so 1,600 values in total) (Fig 4F and

P). Data are by this step in a range of around 10 Hz on a normalised scale. This is down

from an original mossy fibre range of 250 Hz.

A Purkinje cell receives inhibition from two stellate cell networks, one each side. We

repeated the steps above for the other flanking network, then took the mean for each

compartment (Fig 4G and Q).

At the next step, dendritic signals are integrated at the Purkinje cell soma. Integration

is a linear function of compartment data, calculated for each Purkinje cell, in all

sectors, so that this step returned 400 values (4 Purkinje cells x 100 sectors; Fig 4H, I, R

and S). Modulation of somatic charge is fast and bidirectional, and reflects the average

of dendritic states, which transforms into a linear effect on firing.

The result is a progressive contraction of the data range received as inputs to the

system. At each step, data are coded in a frequency distribution. There is a linear

relationship of the mean at each step with all other steps. This result is independent of

the shape of the mossy fibre distribution (we also tested a discontinuous distribution

with the same results).

Linear functions are atypical for dendritic models. What is the evidence? Purkinje cell

firing is a linear function of synaptic excitation, with inhibition blocked [13, 14]. The

granule cell → interneuron → Purkinje cell pathway drives a fall in the firing rate of

Purkinje cells as a linear function of the strength of input from granule cells [16]. In

mice during locomotion, there is a linear relationship between granule cell rates and

interneuron firing, and also between excitation and inhibition of Purkinje cells and

depolarising or hyperpolarising dendritic membrane voltage. Stellate cells contact

Purkinje cells on smooth dendrites [41]. Smooth dendrites extend into all

compartments, mitigating the effect of filtering and distance on dendritic signalling.

Recording from smooth dendrites, locomotion-dependent modulation of dendritic

membrane voltage 'linearly transforms into bidirectional modulation of PC SSp

[Purkinje cell simple spike] output' [15 p.9].

The effect of the form of the parallel fibre code is to simulate simultaneous

independent random sampling by single cells of the whole population of parallel fibre

rates which converge on a microzone (summarised in Fig 5). This mirrors functionally

simulated random sampling by granular layer units of the whole population of mossy

fibre rates received as input to a rank.

568   **Fig 5. The effect of the nature of the granule cell code on sampling in the middle**

569   **layer of the cerebellar network.**

ANATOMY SIMULATES RANDOM SAMPLING BY SINGLE CELLS OF PARALLEL FIBRE RATES RECEIVED BY A WHOLE MICROZONE, WITH REPLACEMENT

GRANULE CELS

CODE

As all fields randomly sample mossy fibre input to a rank, and inhibition of granule cells is synchronised (narrowly constrained to the same range in all fields), all fields return the same frequency distribution of granule cell firing rates.

This has two consequences.

1. Granule cell encoded information has the dimensions of a rank.

2. Below that scale, the same information is coded in any random sample of signals.

REGULATION

The probability that a granule cell fires depends on the relationship of the strength of inhibition and the mean of mossy fibre rates received by a field.

Since, at any time, inhibition is synchronised in all fields and granule cells all randomly sample the same frequency distribution of mossy fibre rates, the probability is the same in all fields

Inhibition of granule cells is proportional to the mean of mossy fibre rates. Therefore, the probability is unchanging.

Because the number of granule cells is large even in a single field, the proportion that fire is well predicted by probability.

Therefore, the regulated density of active parallel fibres is fixed.

STELLATE CELLS

PHYSICALLY SAMPLED POPULATION

A stellate cell receives contact from a few hundred parallel fibres, perhaps 1,000. These are a random sample of the much larger population that pass nearby. As active cells are always in a random pattern, the fraction of cells which make contact that are active at any time is the same as in the general population, and a stellate cell receives a random sample of firing rates.

EFFECTIVELY SAMPLED POPULATION

Anatomy, however, has the notional result that a single stellate cell randomly samples the whole population of active parallel fibres that pass through a microzone, with 'replacement' – ~360,000 signals assuming 1% are active.

This is because all sectors of a microzone receive the same frequency distribution of parallel fibre rates – not a replica, but the same shape and bandwidth, Also, all sectors receive the same regulated density of active cells.

It is functionally immaterial (i.e., codes nothing) which parallel fibres are active or in what pattern, both in the general population and of those which contact a stellate cell.

PURKINJE CELLS

PHYSICALLY SAMPLED POPULATION

Purkinje cell dendritic compartments each receive contact from a random sample of the population of stellate cells afferent to the cell. Contact is from a tiny fraction of the stellate cell population of a microzone, and an unchanging group of cells.

EFFECTIVELY SAMPLED POPULATION

Those cells are themselves, however, notionally a random sample with 'replacement' of the microzone population (at the same depth of the molecular layer).

That is, the result of the microzone population would be the same if stellate cells were reshuffled and a stellate cell could be in more than one place (so that sampling by compartments was both random and independent).

This is because stellate cells are interchangeable because they all(as good as) randomly sample the same distribution of parallel fibre rates.

SOMATIC INTEGRATION

Purkinje cell somatic membrane voltage is a linear function of the mean of dendritic compartment states.

Physically, these are dendritic compartments of the same cell.

But compartment states are (what amounts to) a random sample of the general population of compartment states.

5. A proto-vertebrate wiring plan?

5.1 Recap

We have argued that the microzone computation is a computational effect of

anatomy. This is not incidental but a candidate to account for cell morphologies, the

three-dimensional space they occupy and the scaffold-like geometry of the

connections they make. There is evidence that communication of cerebellar cells is

linear [9-12]. Cells are highly adapted for faithful transmission [17-23]. Neurobiology is

customised to control for biophysical and other noise. 'Although cerebellar neurons

and synapses have nonlinear intrinsic properties [55-59], these properties are

disengaged or compensated in ways that keep the cerebellar network operating in a

linear regime [23, 60-63]' [24]. Firing rates linearly code metrics of behaviour [24].

Anatomy and linearity combine functionally. The result is a passive, unlearned

computational effect of simply passing signals through the network.

To this point, we have a mechanism without a function. How, if at all, does a

microzone discriminate between input from different ranks? Why do neighbouring

microzones receive most of the same parallel fibre signals, and how is their output

functionally related and coordinated?

5.2 Proprioception-motor loops?

The cerebellum is necessary for skilled execution and coordination of movement. We reason that in motor control, the cerebellum may process feedback to directly drive motor output.

It is probably not contentious that early vertebrate swimming, before efficient teleosts, was by undulating body movements (anguilliform). Almost the entire musculature of fish is made up of a bilaterally symmetrical row of paired axial muscle segments, termed myomeres. Movement through the water is produced by a repeating sequence of muscle contractions which cause lateral body flexion. Alternating left and right head turns set up a herring-bone pattern of high pressure which the animal rides by making itself an undulating shape.

Muscles generate analogue signals, in muscle spindles, specialised muscle fibres. Group II afferent neurons signal muscle length. Group Ia afferent neurons fire at a rate that varies with muscle length but in addition varies proportionally to the rate of muscle extension and (more modestly) inversely proportionally to the rate of muscle contraction. Say that in a vertebrate ancestor, input to a row of topographically-defined strips of the cerebellar granular layer arose from a corresponding row of contralateral myomeres, with neighbouring strips receiving the output of neighbouring myomeres. Say also that these provide drive via parallel fibres to the microzone which

equally bisects the group, and that this controls motor output to the ipsilateral muscle

segment opposite the middle of the row providing input. A network in this wiring plan

is the junction of a proprioception-motor loop with multiple input pathways and a

single output pathway. Neighbouring networks connected by parallel fibres send output

to neighbouring myomeres. This connectivity is mirrored in the contralateral cerebellum,

which has motor output to the other half of the body.

This is a candidate proto-vertebrate wiring plan for anguilliform swimming. Fish muscle

fibres are anchored to membranes that separate myomeres. Undulations of the body

cause the length of muscle fibres and the rate of change of the length of muscle fibres

to vary in a rhythmically and smoothly time-varying manner. Input to a rank

accordingly has a sine-wave-like signature. Input to neighbouring ranks is a small step

out of phase. The size of the step depends on number of muscle segments in a full

'cycle' of the trunk. Passive muscle extension might in this way be used to drive muscle

contractions on the other side of the body. Contractions would be automatically

phase-locked and wavelength sensitive.

623   5.3 Why are loops connected by parallel fibres?

624   Parallel fibres coordinate motor outputs and propagate motor sequences, in this

625   system. We briefly take these ideas in turn.

5.3.1 COORDINATION OF MOTOR OUTPUTS

Cerebellar rank data are only segregated in granule cell ascending axons, while signals

travel towards the cerebellar surface. Beyond that, after the granule cell axon divides

in two, parallel fibre signals are randomly intermingled (within modal strata [41, 44,

64]). This means a microzone is unable to discriminate between signals from different

ranks on the face of the data, compounding the cerebellar sacrifice of resolution. The

sacrifice is in two steps. In the first, recoding in ranks does not discriminate between

mossy fibre signals. In the second, a microzone does not discriminate between ranks.

This is not a cost but functional. It means that outputs are coordinated by topography.

Immediately neighbouring microzones receive almost all the same parallel fibre input,

from all the same ranks except one. Input to their next nearest neighbours differs by

two ranks, and so on. We have argued that firing of Purkinje cells is a linear function of

parallel fibre rates and synchronised in a microzone. Therefore, microzones receive

drive which is closely coordinated with their neighbours. The relationship is closest

with near neighbours and weaker with further microzones, diminishing with distance.

The output of a network – a synchronised firing rate or strongly focussed firing rates of

perhaps 50 nuclear output cells – is thus automatically coordinated between parallel-

fibre-connected networks by the relationship of inputs.

Figure 6 illustrates **i)** input and output of a network are constrained to a phase-

sensitive linear relationship, and **ii)** the relationship of the number of muscle segments

in a body wave and the sensitivity of the response.

5.3.2 SELF-DRIVING MOTOR SEQUENCES?

Motor sequences are self-driving. In anguilliform swimming, a wave of muscle

contraction passes down each side of the body, 180 degrees out of phase, initiated by

alternating left and right head turns. Passive extension of neck muscles contralateral to

the direction of a head turn generates mossy fibre input to a corresponding row of

ipsilateral ranks, and indirect input via parallel fibres to a longer row of microzones,

both with strength proportional to the amount and rate of extension, with a rostro-

caudal gradient. This drives motor output at related strength to a caudally-extending

row of ipsilateral muscle segments, causing them to contract with proportional power.

Contractions in turn cause contralateral muscle extension to propagate backwards,

and so on, generating a spiral of signals generated by movement which are received as

input to the cerebellum, and motor outputs.

The result of a single head turn is short-lived and ineffectual. However, alternating left

and right head turns initiate backwards propagating waves of contraction – self-drive –

assisted by passive undulation (modern anguilliform swimming remains in part

passively powered).

Fig 6. In a model of swimming, the network computation is phase sensitive

A Firing rates received as input to a network were varied from rank to rank, so that a plot of the mean and SD of input rates to each rank is sine-wave-like. The frequency distribution received by a single rank is normal and narrow (SD = 12.5 Hz, all ranks). The central rank is indicated by the vertical dashed line. In a locomotor cycle, a single rank receives input which varies sinusoidally with time. All ranks receive the same cycle but a step out of phase with their neighbours. Each pair of panels – 1 and 2, 3 and 4 and so on – represents a moment of time, but a different phase of the cycle, advanced 90 degrees each time. Input (to a network) is shown on the left, output on the right. Output represents firing of the output cells of the network. Output is the same in all phases: parallel fibres perfectly dampen positive feedback. **B** The same, but input rates represent 3/4 of a wavelength and the phase shift between panel pairs is 45 degrees. Output has become sensitive to cycle phase but is still dampened. **C** Input rates represent 1/2 a wavelength. The phase shift is 45 degrees. The output range is now around twice the range in (B). **D** Input rates represent 1/4 of a wavelength (45 degree phase shift). The output range increases again, now approaching the input range (and remains phase sensitive). END CAPTION

6. Discussion

It is clear that the cerebellum is involved in learned behaviours. These can be sophisticated. When rats are trained to press a lever, the striatum receives efferent copies of cortical motor signals. Following training, the action becomes practiced and highly stereotyped. It now seems that that the basal ganglia contribute execution-related elements of the learned behaviour, such as vigour, commitment [65, 66] and kinematics [67] (vs. action selection). If cortico-striatal signals are blocked following training, performance is unimpaired [68]. If, instead, the striatum is lesioned, the learned behaviour is lost. While lesioned animals still engage with the task, they revert to their former, less practiced movements, seen early in training [67]. The same result is obtained if the pathway from the thalamus to the striatum is silenced after learning [68], indicating that the cerebellum is necessary to execute the acquired behaviour. Animals trained over a long period are able to perform the well-practiced action sequence even if the basal ganglia is then lesioned, suggesting that learned changes in the cerebellum are sufficient to execute the behaviour independently.

The changes in the cerebellum that underpin learned behaviour are not fully understood. Historically, interest has focussed on synaptic plasticity, and artificially intelligent networks often use synaptic modification to store learned changes. However, the loss and formation of synapses, growth or reduction of axons, and cell death, neurogenesis and migration, may also provide the substrate of change.

We envisage parallel fibre synaptic changes in a different role. In this view, the

outcome of training under climbing fibre tuition is to polarise parallel fibre synaptic

weights such that all synapses either transmit robustly (working synapses) or do not

transmit at all (silent synapses). The mechanism was first proposed here [69] and is

also discussed here [70]. The learning outcome is the ratio of working to silent

synapses ('synaptic ratio'). In naturally occurring conditions, silent synapses can make

up a large proportion of the synaptic population [71, 72]. Working synapses are a

random subset of the population. There is no pattern memory (the response does not

depend on the pattern of synaptic activations). Following training, all patterns are

received at the same ratio of working to silent synapses regardless of the pattern,

including patterns that are not involved in training. Microzone-grouped Purkinje cells

are trained and learn as a unit, so they all have the same synaptic ratio.

The function of the ratio is a gain adjustment to the anatomical computation, at

network level. The adjustment is made by learned control of the number of signals

that Purkinje cells receive. Because the fraction of active parallel fibres is strictly

regulated, and active cells are a random (though ceaselessly changing) subset of the

general population, the number is a proportion of active cells which depends on the

synaptic ratio. This remains indefinitely adjustable by further training but is fixed on a

behavioural timescale, and pattern-blind. All Purkinje cells in a microzone make the

same adjustment. With this feature, individual gain of unit computations does not

need to be taught or encoded in cell genotypes, and it can be tuned by performance.

What controls the synaptic ratio? Briefly, it depends on the relationship between

parallel fibre rates received by a microzone, and the strength of excitatory drive to the

inferior olive that evokes instruction signals. This is because the synaptic ratio

modulates the strength of nucleo-olivary inhibitory feedback, which in turn determines

whether drive evokes an instruction signal (in this model). The learning outcome is the

synaptic ratio trained by the effect of feedback on the ratio of paired (with a climbing

fibre signal) to unpaired parallel fibre synaptic activations.

This mechanism predicts that excitatory input to the cerebellum and to the olivary

group is from the same or a related source, so that they have correctly related

strength. In this contention, instruction signals are an automated part of a single

system that extends outside the cerebellum (vs. executive supervision from an

independent source).

REFERENCES

1. Garwicz, M., C.F. Ekerot, and H. Jorntell, *Organizational principles of cerebellar neuronal circuitry.* News in Physiological Sciences, 1998. **13**: p. 28-34.
2. Oscarsson, O., *Functional units of the cerebellum - sagittal zones and microzones.* Trends in Neurosciences, 1979. **2**: p. 143-145.
3. Ozden, I., et al., *Reliable coding emerges from coactivation of climbing fibers in microbands of cerebellar Purkinje neurons.* J Neurosci, 2009. **29**(34): p. 10463-73.
4. Pedroarena, C.M. and C. Schwarz, *Efficacy and short-term plasticity at GABAergic synapses between Purkinje and cerebellar nuclei neurons.* J Neurophysiol, 2003. **89**(2): p. 704-15.
5. Napper, R.M. and R.J. Harvey, *Number of parallel fiber synapses on an individual Purkinje cell in the cerebellum of the rat.* J Comp Neurol, 1988. **274**(2): p. 168-77.
6. Harvey, R.J. and R.M. Napper, *Quantitative study of granule and Purkinje cells in the cerebellar cortex of the rat.* J Comp Neurol, 1988. **274**(2): p. 151-7.
7. Mugnaini, E., *The length of cerebellar parallel fibers in chicken and rhesus monkey.* J Comp Neurol, 1983. **220**(1): p. 7-15.
8. Brand, S., A.L. Dahl, and E. Mugnaini, *The length of parallel fibers in the cat cerebellar cortex. An experimental light and electron microscopic study.* Exp Brain Res, 1976. **26**(1): p. 39-58.
9. Rancz, E.A., et al., *High-fidelity transmission of sensory information by single cerebellar mossy fibre boutons.* Nature, 2007. **450**(7173): p. 1245-8.
10. Ritzau-Jost, A., et al., *Ultrafast action potentials mediate kilohertz signaling at a central synapse.* Neuron, 2014. **84**(1): p. 152-163.
11. Duguid, I., et al., *Control of cerebellar granule cell output by sensory-evoked Golgi cell inhibition.* Proc Natl Acad Sci U S A, 2015. **112**(42): p. 13099-104.
12. Duguid, I., et al., *Tonic inhibition enhances fidelity of sensory information transmission in the cerebellar cortex.* J Neurosci, 2012. **32**(32): p. 11132-43.
13. Walter, J.T. and K. Khodakhah, *The linear computational algorithm of cerebellar Purkinje cells.* J Neurosci, 2006. **26**(50): p. 12861-72.
14. Walter, J.T. and K. Khodakhah, *The advantages of linear information processing for cerebellar computation.* Proc Natl Acad Sci U S A, 2009. **106**(11): p. 4471-6.
15. Jelitai, M., et al., *Dendritic excitation-inhibition balance shapes cerebellar output during motor behaviour.* Nat Commun, 2016. **7**: p. 13722.
16. Park, S.M., E. Tara, and K. Khodakhah, *Efficient generation of reciprocal signals by inhibition.* J Neurophysiol, 2012. **107**(9): p. 2453-62.
17. Delvendahl, I. and S. Hallermann, *The Cerebellar Mossy Fiber Synapse as a Model for High-Frequency Transmission in the Mammalian CNS.* Trends Neurosci, 2016. **39**(11): p. 722-737.
18. Saviane, C. and R.A. Silver, *Fast vesicle reloading and a large pool sustain high bandwidth transmission at a central synapse.* Nature, 2006. **439**(7079): p. 983-7.
19. Cesana, E., et al. *Novel granule cell-Golgi cell excitatory input in the cerebellar granular layer.* in *FENS Abstr.* 2010.
20. Moore, J.W. and D.E.J. Blazis, *11 - Simulation of a Classically Conditioned Response: A Cerebellar Neural Network Implementation of the Sutton–Barto–*

Desmond Model, in *Neural Models of Plasticity*, J.H. Byrne and W.O. Berry, Editors.
1989, Academic Press. p. 187-207.
21. Sargent, P.B., et al., *Rapid vesicular release, quantal variability, and spillover*
contribute to the precision and reliability of transmission at a glomerular synapse.
J Neurosci, 2005. **25**(36): p. 8173-87.
22. Turecek, J., S.L. Jackman, and W.G. Regehr, *Synaptic Specializations Support*
Frequency-Independent Purkinje Cell Output from the Cerebellar Cortex. Cell
Reports, 2016. **17**(12): p. 12.
23. Turecek, J., S.L. Jackman, and W.G. Regehr, *Synaptotagmin 7 confers frequency*
invariance onto specialized depressing synapses. Nature, 2017. **551**: p. 503.
24. Raymond, J.L. and J.F. Medina, *Computational Principles of Supervised Learning in*
the Cerebellum. Annu Rev Neurosci, 2018. **41**: p. 233-253.
25. DiGregorio, D.A., et al., *Desensitization properties of AMPA receptors at the*
cerebellar mossy fiber granule cell synapse. J Neurosci, 2007. **27**(31): p. 8344-57.
26. Mapelli, L., S. Solinas, and E. D'Angelo, *Integration and regulation of glomerular*
inhibition in the cerebellar granular layer circuit. Front Cell Neurosci, 2014. **8**: p.
55.
27. Galliano, E., P. Mazzarello, and E. D'Angelo, *Discovery and rediscoveries of Golgi*
cells. J Physiol, 2010. **588**(Pt 19): p. 3639-55.
28. Gilbert, M. and A. Rasmussen, *The cerebellum converts input data into a hyper*
low-resolution granule cell code with spatial dimensions: a hypothesis. Royal
Society Open Science, 2025, in press (in the meantime an earlier draft is available
at bioRxiv DOI: 10.1101/2023.07.14.548987 ver 9,
https://www.biorxiv.org/content/10.1101/2023.07.14.548987v9).
29. Wu, H.S., I. Sugihara, and Y. Shinoda, *Projection patterns of single mossy fibers*
originating from the lateral reticular nucleus in the rat cerebellar cortex and nuclei.
J Comp Neurol, 1999. **411**(1): p. 97-118.
30. Sultan, F. and D. Heck, *Detection of sequences in the cerebellar cortex: numerical*
estimate of the possible number of tidal-wave inducing sequences represented. J
Physiol Paris, 2003. **97**(4-6): p. 591-600.
31. Shinoda, Y. and I. Sugihara, *Axonal Trajectories of Single Climbing and Mossy Fiber*
Neurons in the Cerebellar Cortex and Nucleus, in *Handbook of the Cerebellum*
and Cerebellar Disorders, S.J.D. Manto M., Rossi F., Gruol D.L., Koibuchi N.,
Editor. 2013, Springer, Dordrecht.
32. Shinoda, Y., et al., *The entire trajectory of single climbing and mossy fibers in the*
cerebellar nuclei and cortex. Prog Brain Res, 2000. **124**: p. 173-86.
33. Marr, D., *A theory of cerebellar cortex.* J Physiol, 1969. **202**(2): p. 437-70.
34. Albus, J.S., *A Theory of Cerebellar Function.* Mathematical Biosciences, 1971.
10(1-2): p. 25-61.
35. Gilbert, M. and A. Rasmussen, *Gap Junctions May Have A Computational Function*
In The Cerebellum: A Hypothesis. The Cerebellum, 2024. **23**(5): p. 1903-1915.
36. Jörntell, H. and C.F. Ekerot, *Receptive field plasticity profoundly alters the*
cutaneous parallel fiber synaptic input to cerebellar interneurons in vivo. J
Neurosci, 2003. **23**(29): p. 9620-31.

37.   Harvey, R.J. and R.M. Napper, *Quantitative studies on the mammalian cerebellum*.
Prog Neurobiol, 1991. **36**(6): p. 437-63.
38.   Wilms, C.D. and M. Häusser, *Reading out a spatiotemporal population code by*
imaging neighbouring parallel fibre axons in vivo. Nature Communications, 2015.
6: p. 6464.
39.   Cesana, E., et al., *Granule cell ascending axon excitatory synapses onto Golgi*
cells implement a potent feedback circuit in the cerebellar granular layer. J
Neurosci, 2013. **33**(30): p. 12430-46.
40.   Häusser, M. and B.A. Clark, *Tonic synaptic inhibition modulates neuronal output*
pattern and spatiotemporal synaptic integration. Neuron, 1997. **19**(3): p. 665-78.
41.   Palay, S.L. and V. Chan-Palay, *Cerebellar cortex : cytology and organization*. 1974,
Berlin: Springer.
42.   Carter, A.G. and W.G. Regehr, *Quantal events shape cerebellar interneuron firing*.
Nat Neurosci, 2002. **5**(12): p. 1309-18.
43.   Llinás, R.R. and M.N. Negrello. *Cerebellum*. Scholarpedia, 10(1):4606 2015 [cited
2017 06/01]; Available from: http://www.scholarpedia.org/article/Cerebellum.
44.   Jörntell, H. and C.F. Ekerot, *Properties of somatosensory synaptic integration in*
cerebellar granule cells in vivo. J Neurosci, 2006. **26**(45): p. 11786-97.
45.   van Beugen, B.J., et al., *High Frequency Burst Firing of Granule Cells Ensures*
Transmission at the Parallel Fiber to Purkinje Cell Synapse at the Cost of Temporal
Coding. Frontiers in Neural Circuits, 2013. **7**: p. 95.
46.   Person, A.L. and I.M. Raman, *Synchrony and neural coding in cerebellar circuits*.
Frontiers in Neural Circuits, 2012. **6**: p. 97.
47.   Stell, B.M., et al., *Activation of presynaptic GABA(A) receptors induces glutamate*
release from parallel fiber synapses. J Neurosci, 2007. **27**(34): p. 9022-31.
48.   Dellal, S.S., R. Luo, and T.S. Otis, *GABAA receptors increase excitability and*
conduction velocity of cerebellar parallel fiber axons. J Neurophysiol, 2012.
107(11): p. 2958-70.
49.   Pugh, J.R. and C.E. Jahr, *Activation of Axonal Receptors by GABA Spillover*
Increases Somatic Firing. The Journal of Neuroscience, 2013. **33**(43): p. 16924-
16929.
50.   Pugh, J.R. and C.E. Jahr, *Axonal GABAA receptors increase cerebellar granule cell*
excitability and synaptic activity. J Neurosci, 2011. **31**(2): p. 565-74.
51.   Zhou, H., et al., *Cerebellar modules operate at different frequencies*. Elife, 2014.
3: p. e02536.
52.   Cerminara, N.L. and J.A. Rawson, *Evidence that climbing fibers control an intrinsic*
spike generator in cerebellar Purkinje cells. J Neurosci, 2004. **24**(19): p. 4510-7.
53.   Raman, I.M. and B.P. Bean, *Ionic currents underlying spontaneous action*
potentials in isolated cerebellar Purkinje neurons. J Neurosci, 1999. **19**(5): p. 1663-
74.
54.   Sultan, F. and J.M. Bower, *Quantitative Golgi study of the rat cerebellar molecular*
layer interneurons using principal component analysis. J Comp Neurol, 1998.
393(3): p. 353-73.
55.   Hoebeek, F.E., et al., *Differential olivo-cerebellar cortical control of rebound*
activity in the cerebellar nuclei. Proc Natl Acad Sci U S A, 2010. **107**(18): p. 8410-5.

56.   Ishikawa, T., M. Shimuta, and M. Häusser, *Multimodal sensory integration in single*
872   *cerebellar granule cells in vivo.* eLife, 2015. **4**: p. e12916.

873   57.   Jahnsen, H., *Electrophysiological characteristics of neurones in the guinea-pig*
874   *deep cerebellar nuclei in vitro.* J Physiol, 1986. **372**: p. 129-47.

875   58.   Loewenstein, Y., et al., *Bistability of cerebellar Purkinje cells modulated by*
876   *sensory stimulation.* Nat Neurosci, 2005. **8**(2): p. 202-11.

877   59.   Person, A.L. and I.M. Raman, *Purkinje neuron synchrony elicits time-locked*
878   *spiking in the cerebellar nuclei.* Nature, 2012. **481**(7382): p. 502-505.

879   60.   Alvina, K., et al., *Questioning the role of rebound firing in the cerebellum.* Nat
880   Neurosci, 2008. **11**(11): p. 1256-8.

881   61.   Rössert, C., et al., *Model cerebellar granule cells can faithfully transmit modulated*
882   *firing rate signals.* Front Cell Neurosci, 2014. **8**: p. 304.

883   62.   Schonewille, M., et al., *Purkinje cells in awake behaving animals operate at the*
884   *upstate membrane potential.* Nat Neurosci, 2006. **9**(4): p. 459-61; author reply
885   461.

886   63.   van Vreeswijk, C. and H. Sompolinsky, *Chaos in neuronal networks with balanced*
887   *excitatory and inhibitory activity.* Science, 1996. **274**(5293): p. 1724-6.

888   64.   Zhang, W. and D.J. Linden, *Calcium influx measured at single presynaptic boutons*
889   *of cerebellar granule cell ascending axons and parallel fibers.* Cerebellum, 2012.
890   **11**(1): p. 121-31.

891   65.   Dudman, J.T. and J.W. Krakauer, *The basal ganglia: from motor commands to the*
892   *control of vigor.* Curr Opin Neurobiol, 2016. **37**: p. 158-166.

893   66.   Yttri, E.A. and J.T. Dudman, *Opponent and bidirectional control of movement*
894   *velocity in the basal ganglia.* Nature, 2016. **533**(7603): p. 402-406.

895   67.   Dhawale, A.K., et al., *The basal ganglia control the detailed kinematics of learned*
896   *motor skills.* Nature Neuroscience, 2021. **24**(9): p. 1256-1269.

897   68.   Wolff, S.B.E., R. Ko, and B.P. Ölveczky, *Distinct roles for motor cortical and*
898   *thalamic inputs to striatum during motor skill learning and execution.* Sci Adv,
899   2022. **8**(8): p. eabk0231.

900   69.   Gilbert, M., *Gating by Memory: a Theory of Learning in the Cerebellum.* The
901   Cerebellum, 2021. **21**(6): p. 926-943.

902   70.   Gilbert, M. and A. Rasmussen, *What if...? Rethinking the role of learning in neural*
903   *networks.* 2025. HAL. https://hal.science/hal-04863982.

904   71.   Isope, P. and B. Barbour, *Properties of unitary granule cell-->Purkinje cell*
905   *synapses in adult rat cerebellar slices.* J Neurosci, 2002. **22**(22): p. 9668-78.

906   72.   Ekerot, C.F. and H. Jörntell, *Parallel fibre receptive fields of Purkinje cells and*
907   *interneurons are climbing fibre-specific.* Eur J Neurosci, 2001. **13**(7): p. 1303-10.

908