Rapid production of modified cattle and sheep from spermatid-like stem cells

Lei Yang[1, 2, 4, 5, *], Anqi Di[1, 4], Lishuang Song[1, 4], Xuefei Liu[1], Di Wu[1], Song Wang[1, 3], Zhenting Hao[1], Lige Bu[1], Chunling Bai[1], Guanghua Su[1], Zhuying Wei[1], Li Zhang[1], Zhonghua Liu[3], Shaorong Gao[2, 5, *], and Guangpeng Li[1, 5, *]

1 State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock, College of Life Sciences, Inner Mongolia University, Hohhot, China.

2 Frontier Science Center for Stem Cell Research, School of Life Sciences and Technology, Tongji University, Shanghai, China.

3 Key Laboratory of Animal Cellular and Genetics Engineering of Heilongjiang Province, College of Life Science, Northeast Agricultural University, Harbin, China

4 These authors contributed equally: Lei Yang, Anqi Di, Lishuang Song

5 These authors jointly supervised this work: Lei Yang, Shaorong Gao, Guangpeng Li

* e-mail: leiyang@imu.edu.cn; gaoshaorong@tongji.edu.cn; gpengli@imu.edu.cn

Abstract

Haploid androgenetic stem cells (haSCs) are revolutionary resources for studying animal traits and greatly impact livestock breeding, but livestock haSCs have yet to be obtained. Here, we report the derivation of cattle and sheep haSCs using a novel FACE medium, and these cells harbor formative pluripotency features. Notably, knockout cattle and sheep can be efficiently obtained upon injection of the protaminized haSCs with targeted gene mutations into oocytes.

Main

Haploid androgenetic stem cells (haSCs) can support the production of viable transgenic rodents (mice and rats) by oocyte intracytoplasmic haSCs injection (iCHI) [1-3]. Currently, somatic cell nuclear transfer (SCNT) allows CRISPR/Cas-edited cells to give rise to produce genetically modified livestock. However, the low efficiency of CRISPR-SCNT dramatically limits its application. Thus, CRISPR-iCHI holds excellent promise in agrobiotechnology, but livestock haSCs have not been achieved.

To this end, we generated bovine haploid androgenetic (ha) embryos by two different methods: (i) injecting sperm into enucleated oocytes; (ii) removing the female pronucleus from zygotes (Fig.1a, Supplementary Fig.1a). Expectedly, sperm underwent actively DNA demethylate in ha embryos [2] (Supplementary Fig.1b). The blastocyst rate of the injecting and removing approach was comparable (31.16% vs. 30.21%; Supplementary Fig.1d,e, Table 1), indicating that the reprogramming of sperm by ooplasm was not affected by the manipulations. The ha-blastocysts were further cultivated feeder cells under "naïve" or "primed" conditions with mouse, human, and bovine diploid SCs to derive bovine haSCs (b-haSCs; Supplementary Fig.2a). From 261 ha-blastocyst, we only obtained outgrowths consisting of "master-regulators" OCT4$^+$SOX2$^+$ in LCDM, CTFR, and bEPSC medium, but couldn't establish stable cell lines (no more than eight passages; Supplementary Fig.2b-d).

To long-term maintenance of the b-haSCs, we screened the different combinations of

inhibitor and cytokine in the above successful culturing medium (11 hits that target ten

signal-pathways; Supplementary Fig.3a). Cells cultured in FACE [FGF2::Activin-

A::CHIR99021::IWR-1-endo] and FACX [FAC::XAV939] combinations formed colonies,

while in others appeared differentiated or did not form colonies (Supplementary

Fig.3b,c). Both IWR-1-endo and XAV939 act as Wnt/β-catenin inhibitors and could be

used interchangeably [4, 5], thus [FACX] and [FACE] are essentially the same medium.

We further titrated Activin-A and obtained cell lines that could be propagated for more

than 80 passages in low Activin-A, and haploid cells could be enriched by

fluorescence-activated cell sorting (FACS; Fig.1b-d, Supplementary Fig.2e, 3d, 4a-c).

We also succeeded in establishing ovine haSCs (o-haSCs) by FACE medium, which

expressed pluripotency marks similar to that of b-haSCs (Fig.1b-d, Supplementary

Fig.1c-e, 2e, 4b,c, Table 1). Consistent with the rodent haSCs [1-3], no ruminant haSCs

with the Y chromosome (Supplementary Fig.4b,d). Strikingly, originally obtained b- and

o-haSCs contain nearly 20% haploid cells, which was higher than that of murine haSCs

[1-3] (< 5%; Fig.1c). Overall, the novel FACE medium can be used for both b- and o-

haSCs derivation.

Inner cell mass (ICM) of the blastocyst, during gastrulation, is through three

pluripotency states known as naïve, formative, and primed states [6] (Supplementary

Fig.6a). It is well known that, under the naïve state, mouse haSCs (m-haSCs) tend to

rapidly diploidization [1, 2], whereas the ruminant haSCs do not. These results prompted

us to determine the pluripotency state of ruminant haESCs. To this end, we analysed

the transcriptome by bulk RNA sequencing (buRNA-seq) and found that the b-haSCs

couldn't belong to any stage of pre-implantation embryos [7, 8] (Supplementary Fig.5a,b).

Meanwhile, b-haSCs also segregated from the recently reported diploid primed and

expanded SCs [9, 10] (Supplementary Fig.5c,d), implying that b-haSCs enriches the

features of formative pluripotency (Supplementary Fig.5e).

To test this hypothesis, we performed single cell (sc) RNA-seq at the gastrulation

period of bovine epiblast from *in vivo* diploid embryos (E8~14; Supplementary Fig.6b,c).

Expectedly, b-haSCs were clustered with the embryos at E10 (Supplementary

Fig.6b,c). Indeed, we identified three types of cells in b-haSCs: 83.8% formative, 12.1%

primed, and a few 4.1% naïve (Fig.1e,f, Supplementary Fig.7a,b). To further dissect

the epigenetic features, we performed CUT&Tag assay (Supplementary Fig.8a). The

H3K4me3 and H3K27me3 at the promoter of formative genes showed higher and

lower in b-haSCs than primed b-SCs, respectively. Also, these modifications of primed

genes exhibited the opposite trend (Fig.1g, Supplementary Fig.8b). In agreement, the

H3K4me3 and H3K27me3 clustered profiles segregated from primed b-SCs

(Supplementary Fig.8c). The presence of H3K4me3 but not H3K27me3 in the

promoter of basic-pluripotency genes, including OCT4, SOX2, and SALL4, is common

to both b-haSCs and primed diploid b-SCs (Supplementary Fig.8d).

We further undertook a cross-species comparison, and found that o-haSCs clustered

closer with b-haSCs than with primed o-SCs (Supplementary Fig.7c). Formative-

related regulators were both present in bovine, ovine, and murine formative cells

(Fig.1h). The mRNA and protein levels of the key formative markers, including

DNMT3A, DNMT3B, and OTX2, were comparable among each species

(Supplementary Fig.7d,e). We thus conclude that b- and o-haSCs harbor molecular

features of formative pluripotency.

Expectedly, both b- and o-haSCs could differentiate into three germ-line with around

16% haploid cells both *in vitro* and *in vivo*, which is much higher than that of m-haSCs

[1] (Supplementary Fig.9a-d, 10a-d). Both b- and o-haSCs could localised to ICM

positions of the intra-/inter-species chimeric blastocysts (Supplementary Fig.11a-d).

Following embryo transfer into the foster mouse, we detected b-haSCs (but not o-

haSCs) gave rise to live inter-species chimeras at E18.5 (albino mouse for the host;

Fig.1i, Supplementary Fig.12a). However, cattle-mouse chimera only reached 2-day-

old with colored eyes (Supplementary Fig.12b,c), suggesting that xenogeneic cells

cannot maintain the metabolic environment for each other.

We next performed the iCHI experiment to replace sperm, by injecting the b- and o-

haSCs into bovine and ovine oocytes, respectively [2] (Supplementary Fig.13a,b). The

injected b- and o-haSCs formed a pseudo-pronucleus and underwent DNA

demethylation, which is consistent with *in vitro* fertilization (IVF) embryos

(Supplementary Fig.13c,d). Similar to the embryos derived by SCNT, most b- and o-

iCHI embryos could not develop to the blastocyst stage (Fig.2a, Supplementary

Fig.14a, Table 2). We and other previous studies revealed aberrant DNA and histone

methylation is a barrier for iCHI and SCNT [11, 12]. We thus performed pyrosequencing

and whole genome bisulfite sequencing (WGBS) on b- and o-haSCs, and detected

that the differentially methylated regions (DMRs) of parental imprints maintained

proper status as in sperm, including H19, IGF2, and DIO3 (Fig.2b, Supplementary

Fig.15a-c).

Unlike the asymmetric in IVF zygote, H3K4/K9/K27me3 in all pronuclei of b- and o-

iCHI embryos (Fig.2c, Supplementary Fig.14b). Injection with relative demethylases

KDM5B/4D/6A mRNA into b- and o-iCHI zygote, could not correct symmetric errors

(Supplementary Fig.16a-c, Table 2). Moreover, KDM5B/4D/6A overexpression

accelerated self-diploidization of ruminant haSCs (Supplementary Fig.16d). These

results prompted us to find another method to rescue the ruminant iCHI embryos.

As protamine (Pro) can exclude histone modifications during spermatogenesis [13], we

used the non-integrated episomal vector to express the PRM1 gene transiently. After

Pro-induction in b- and o-haSCs, we observed their nuclei compacted into spermatid-

like structures concomitant with H3K4/9/27me3 removal (Supplementary Fig.17a,b).

Importantly, the symmetric errors in b- and o-iCHI embryos can be amended by using

protaminized b- and o-haSCs as donors, leading iCHI blastocyst rate was comparable

to that of IVF embryos (Fig.2a,c, Supplementary Fig.14a,b, Table 2).

To maximize the utility of the Pro-iCHI approach, we chose to edit the Myostatin (MSTN)

gene, whose mutation in animals causes a much greater muscle mass known as

"double-muscling" [14]. Although the CRISPR prime editor (PE) has been applied in mice,

it remains to be tested whether PE could be applied in large animals [15]. Genomic

integration of exogenous DNA increases the risk of biosafety. We thus developed an

episomal plasmid-based PE (ePE) to generate the gene-edited livestock with

integration-free. Indeed, the ePE enables the production of nearly 100% short deletion

(11 bp) and nucleotide substitution (G>A) in MSTN loci of the b- and o-haSCs,

respectively (Supplementary Fig.18a-e).

We further tested whether the Pro-iCHI approach can produce live animals.

Consequently, one MSTN-edited premature lamb with a much smaller size was

obtained at 14 weeks (0.39%, 1/256; Fig.2d, Supplementary Table 3). Of the bovine

Pro-iCHI embryos, 14.3% (1/7) were born as full-term (287 days) calf using WT b-

haSCs as donor cells. Remarkably, we obtained two calves (13.3%, 2/15) with "double-

muscle" phenotype using MSTN-edited b-haSCs (Fig.2e, Supplementary Fig.19a-e,

Table 3).

Here, we show that FACE medium supports the establishment of b- and o-haSCs. As

a proof-of-concept, we also developed a novel Pro-iCHI::ePE system to rapidly

produce trait-improved cattle with exogene-free effects.

References

1. Li, W. et al. Androgenetic haploid embryonic stem cells produce live transgenic mice. *Nature* **490**, 407-411 (2012).
2. Yang, H. et al. Generation of genetically modified mice by oocyte injection of androgenetic haploid embryonic stem cells. *Cell* **149**, 605-617 (2012).
3. Li, W. et al. Genetic modification and screening in rat using haploid embryonic stem cells. *Cell Stem Cell* **14**, 404-414 (2014).
4. Lu, J. et al. Structure-activity relationship studies of small-molecule inhibitors of Wnt response. *Bioorg Med Chem Lett* **19**, 3825-3827 (2009).
5. Huang, S.M. et al. Tankyrase inhibition stabilizes axin and antagonizes Wnt signalling. *Nature* **461**, 614-620 (2009).
6. Smith, A. Formative pluripotency: the executive phase in a developmental continuum. *Development* **144**, 365-373 (2017).
7. Graf, A. et al. Fine mapping of genome activation in bovine embryos by RNA sequencing. *Proc Natl Acad Sci U S A* **111**, 4139-4144 (2014).
8. Jiang, Z. et al. Transcriptional profiles of bovine in vivo pre-implantation development. *BMC Genomics* **15**, 756 (2014).
9. Bogliotti, Y.S. et al. Efficient derivation of stable primed pluripotent embryonic stem cells from bovine blastocysts. *Proc Natl Acad Sci U S A* **115**, 2090-2095 (2018).
10. Zhao, L. et al. Establishment of bovine expanded potential stem cells. *Proc Natl Acad Sci U S A* **118** (2021).
11. Zhong, C. et al. CRISPR-Cas9-Mediated Genetic Screening in Mice with Haploid Embryonic Stem Cells Carrying a Guide RNA Library. *Cell Stem Cell* **17**, 221-232 (2015).
12. Yang, L., Song, L., Liu, X., Bai, L. & Li, G. KDM6A and KDM6B play contrasting roles in nuclear transfer embryos revealed by MERVL reporter system. *EMBO Rep* **19** (2018).
13. Czernik, M., Iuso, D., Toschi, P., Khochbin, S. & Loi, P. Remodeling somatic nuclei via exogenous expression of protamine 1 to create spermatid-like structures for somatic nuclear transfer. *Nat Protoc* **11**, 2170-2188 (2016).
14. Rodgers, B.D. & Garikipati, D.K. Clinical, agricultural, and evolutionary biology of myostatin: a comparative review. *Endocr Rev* **29**, 513-534 (2008).
15. Anzalone, A.V. et al. Search-and-replace genome editing without double-strand breaks or donor DNA. *Nature* **576**, 149-157 (2019).

Methods

Animals and chemicals

All animal procedures were performed under the ethical guidelines of the State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock, College of Life Sciences, Inner Mongolia University (IMU). Specific pathogen-free-grade B6D2F1 (C57BL/6 × DBA/2), CD1, and Kun-Ming (KM) mice were purchased from Laboratory Animal Research Center (Inner Mongolia University) or Vital River Laboratories Co., Ltd. Livestock are raised at the experimental farm of IMU and processed in commercial slaughterhouses. Mice were housed in specific pathogen-free conditions with a 12-h dark/12-h light cycle. Unless otherwise indicated, all chemicals were purchased from Sigma-Aldrich Chemical Reagent Co., Ltd.

Collection and *in vitro* maturation of oocytes

Collection and *in vitro* maturation (IVM) of ruminant oocytes were performed following the published protocols with slight modifications [13, 16]. Briefly, bovine or ovine ovaries from the slaughterhouse were transported to the laboratory. Collect cumulus-oocyte complexes (COCs) and transfer them to the IVM medium for 24 h in 5% CO_2 with humidified atmosphere at 38.5 °C [basic M199 medium supplemented with 2 mM glutamine, 300 µM, sodium pyruvate, 100 µM cysteamine, follicle-stimulating hormone (FSH; 50 ng/mL), luteinizing hormone (LH; 3 mg/mL), 1 µg/mL estradiol, and 0.04% gentamicin solution].

Generation of haploid androgenetic embryos

Sperm of Angus cattle or Dorper sheep were prepared for manipulation according to the published protocols [17-19]. The oocytes or zygotes were stained with 5 µg/mL Hoechst 33342 for 10 min to visualize spindle or pronuclei by a short 1-2 sec exposure to the ultraviolet (UV) light. Option A for injecting strategy, one-step SCNT was performed [20, 21], in which somatic cells were replaced by sperms. Briefly, the MII oocyte spindle was adjusted to 8 to 10 o'clock, and one donor sperm was injected into the nearby plasma. Immediately, the spindle was aspirated into the injection pipette and removed from the MII oocyte using the Piezo micromanipulator (Primetech) in the M2 medium containing 5 µg/mL cytochalasin B (CB) on a 37 °C heating stage of an inverted microscope (Nikon). After manipulation, the androgenetic embryos were activated in 5 µM ionomycin for 5 min and then incubated in 2 mM 6-dimethylaminopurine (6-DMAP) and 7.5 µg/mL CB in Ca^{2+}-free KSOM-AA for 5 h. Option B for removing strategy, the zygotes were produced by *in vitro* fertilization (IVF) as previously described [22]. Briefly, the zygotes were cultured in G1 medium for 7-8 h after IVF. Female pronuclei, which were near the second polar body, were removed

from zygotes in the M2 medium containing 5 μg/mL CB using a blunt Piezo-driven
pipette. Following activation or removing, the androgenetic embryos were cultured in
G1 medium (Vitrolife) for 72 h, and then transferred into G2 medium (Vitrolife) for
further culturing at 38.5 °C, humidity to saturation, 5% CO_2, 5% O_2, and 90% N_2. The
androgenetic embryos were allowed to develop into blastocysts that were used for
haSCs derivation.
239

Further culture of blastocysts using different SCs medium

The blastocysts with good morphology were selected to generate SCs. In brief, zona
pellucida (ZP) were removed from blastocysts by treatment with 5 mg/mL of pronase
for 3-4 min under an inverted microscope with a warming stage. After removing ZP,
blastocysts were plated on mitomycin C (MMC) treated mouse embryonic fibroblasts
(MEFs) in the different mediums for SCs derivation, including 2i/L [23], 4i [24], t2iL+Gö [25],
5i/L/A [26], ABCL [27], LCDM [28], EPSC [29, 30], CTFR [9, 31], bEPSC [10]. After 10 days in culture,
the outgrowths were manually picked in separate wells of a 12-well dish seeded with
MMC-treated MEFs for further cultivation to avoid contamination of trophoblasts.
249

Further culture of blastocysts using different small-molecules

Individual denuded blastocysts were placed in separate wells of a 12-well dish seeded
with a monolayer of MMC-treated MEFs, and cultured in basal N2B27 [32] or mTeSR [9]
medium supplemented with 15% knockout serum replacement (KSR; Thermo), and
different small-molecule compounds (Supplementary information, Fig.S3a). The small-
molecules are used at the following concentrations: 3 μM CHIR99021 (C;
Selleckchem), 2 μM Dimethindene maleate (D; Tocris), 2 μM Minocycline
hydrochloride (M; Santacruz), 10 ng/mL LIF (L; Millipore), 2.5 μM IWR-1-endo (E;
Selleckchem), 10 ng/mL FGF2 (F; Peprotech), 5 μM XAV939 (X; Selleckchem), 0.3
μM WH-4-023 (W; Selleckchem), 1 ng/mL TGFβ (T; Peprotech), 20 ng/mL Activin-A (A;
Peprotech), 50 ng/mL BMP4 (B; R&D systems), 50 μg/mL Vitamin C (V; Selleckchem).
All small-molecules are dissolved in dimethyl sulfoxide (DMSO). After 10 days in
culture, the outgrowths were manually picked to avoid contamination of trophoblasts.
263

Derivation and routine culture of haSCs

After removing ZP, each blastocyst was plated into one well of a 96-well plate with the
MMC-treated MEFs feeder in FACE medium at 37 °C, humidity to saturation, 5% CO_2,
5% O_2, and 90% N_2. The outgrowths were formed after 10-14 days, the medium was
changed by half every 1-2 days, and passaged on the MEFs feeder followed by
mechanically separating into several pieces. Cell expansion was carried out from 48-
well to 6-well plates with feeder cells and then to 6-well plates for routine culture.
Established haSCs were cultured on the MMC-treated MEFs feeder in FACE medium,
with daily media change. For passaging, the cells were disaggregated using TrypLE

(Thermo) or Accutase (Stemcell) and passaged onto new feeder layers at a split ratio
of 1:10 or 1:20 every 3-5 days. To enhance the survival of haSCs, 10 µM Y-27632
(ROCK inhibitor, Stemcell) was recommended to be added into the FACE medium for
24 h after the passage.
277
The basal medium for FACE is mTeSR1 (Stemcell, 85850 based media) [33] or N2B27
[32] [total 500 mL was prepared as follows: 240 mL DMEM/F12 (Thermo), 240 mL
neurobasal (Thermo), 2.5 mL N2 supplement (Thermo), 5 mL B27 supplement
(Thermo), 1% GlutaMAX (Thermo), 1% nonessential amino acids (Thermo), 0.1 mM
β-mercaptoethanol (Thermo), 1% penicillin-streptomycin (Thermo), and 15% KSR
(Thermo)]. To prepare the FACE medium, small molecules and cytokines were added
in the basal medium at the following final concentrations: 10 ng/mL FGF2 (F;
Peprotech), 10 ng/mL Activin-A (A; Peprotech), 3 µM CHIR99021 (C; Selleckchem),
and 2.5 µM IWR-1-endo (E; Selleckchem).
287

FACS-enrichment of haploid cells
To sort haploid cells, we used fluorescence activating cell sorter (FACS) according to
the published protocol with slight modifications [34-36]. Briefly, haSCs were dissociated
into single cells with TrypLE (Thermo) or Accutase (Stemcell) and then incubated with
15 µg/mL Hoechst 33342 for 15 min in a 37 °C water bath. Following centrifugation,
cells were rinsed with Dulbecco's phosphate-buffered saline (PBS) for removal of the
Hoechst 33342, and resuspended in FACE medium with 10 µM Y-27632.
Subsequently, the stained cells were through 40 µm cell strainer and the haploid cells
were sorted using FACSAria II (BD biosciences). After enrichment, the haploid cells
were plated on fresh MEFs feeder in FACE medium containing 10 µM Y-27632. After
24 h, haploid cells were cultured in FACE medium without Y-27632. Usually, haESC
clones were formed within 10 days. For cell analysis, after fixation in 70% ethanol, cells
were digested by 20 mg/ml RNase A and stained with 50 mg/ml propidium iodide (PI).
Then, the haploid cells were analyzed using CytoFLEX LX (Beckman). Analytic flow
profiles were performed using FlowJo software (Ashland).
303

Immunofluorescence staining
Immunofluorescence (IF) was performed as described previously [20]. Briefly, the cells
were fixed with 4% paraformaldehyde (PFA) in PBS at RT for 1 h, and permeabilised
in PBS supplemented with 0.2% Triton X-100 at RT for 45 min. The cells were
incubated with primary antibodies in PBS that contained 3% blocker BSA (Thermo) at
4 °C overnight, then incubated with secondary antibodies for 1 h at RT in the dark. For
5mC and 5hmC staining, permeabilised embryos were incubated in 4 N HCl treatment
for 10 min at room temperature, then the oocytes were immediately transferred to Tris-
HCl (pH 8.5) for 10 min to neutralise the residual HCl. Finally, the cells were mounted
in the anti-fade solution with DAPI (Thermo) and compressed with a cover-slip. All
samples were observed by a laser scanning microscope (Nikon). The antibody

information was presented in Supplementary Table 4.
316

Western blot

Western blot (WB) was performed as described previously [12]. In brief, cells were lysed
in RIPA lysis buffer (Beyotime) with protease inhibitor (Thermo) and phosphatase
inhibitor (Thermo). Whole-cell lysates were separated by sodium dodecyl sulfate-
polyacrylamide gel electrophoresis (SDS-PAGE). Following electrophoresis (90 V for
30 min, 30–50 V overnight), the proteins were transferred from the gel to 0.2 mm PVDF
membranes (Bio-sharp), and blocked for 2 h at RT in TBST buffer containing 5% BSA
or 5% fat-free milk. Films were incubated with the primary antibodies in TBST buffer
containing 3% BSA at 4 °C overnight. After washing three times in TBST, PVDF
membranes were incubated with secondary antibodies for 1 h at RT. Then, bound
antibodies were detected with SuperSignal West Femto Substrate Trial Kit (Thermo).
The antibody information was presented in Supplementary Table 4.
329

Karyotype analysis

Before karyotype analysis, haSCs were incubated with 0.1 µg/mL KaryoMAX Colcemid
(Thermo) in FACE medium for 3-4 h, and mechanically separated from the MEFs
feeder using pipette tips. Then, haSCs were disaggregated into single cells by 0.05%
trypsin-EDTA (Thermo) at 37 °C for 10 min. After centrifugation, the pellet was re-
suspended in hypotonic solution (0.56% KCL) at room temperature (RT) for 10 min.
Cells were then fixed with ice-cold fixative (3:1 methanol: acetic acid) at RT for 5 min
three times. Finally, haSCs suspension was dropped onto an ice-cold fixative washed
slide and stained with 4′,6-diamidino-2-phenylindole (DAPI; Thermo). For each cell
line, more than 30 metaphases were examined.
340

Chimeric assay

The chimeric assay was performed as previously described with slight modification [20].
In brief, haSCs were used 1 day before passaging, which showed an optimal
undifferentiated morphology. For chimeric mice, ten haSCs were microinjected into
CD1 or KM blastocysts using a blunt Piezo-driven pipette. Then, the reconstructed
blastocysts were transplanted into the uterus of pseudo-pregnant mice at 3.5 days
post-coitum (dpc). For chimeric blastocyst, six haSCs labelled by EGFP fluorescence
were microinjected into an 8-cell stage B6D2F1 embryo. The reconstructed embryos
were cultured in FACE medium for 4-5 h, then changed into the G1/G2 medium (1:1;
Vitrolife) to obtain the chimeric blastocysts.
351

Embryoid body formation assay

Embryoid body (EB) formation was performed as previously described with slight
modification [34]. Briefly, haSCs were dissociated into single cells with TrypLE (Thermo)
and cultured for 7 days on ultra-low attachment plates in Iscove's modified Dulbecco's
medium (IMDM; Thermo) supplemented with 1% nonessential amino acids (Thermo),
1% penicillin-streptomycin (Thermo), and 15% FBS (Gibco). Subsequently, Ebs were
collected and dissociated by 0.05% trypsin-EDTA (Thermo) at 37 °C for 10 min. After

enrichment, cells were plated on atrigel-coated slides for an additional 3 days before IF staining.

Teratoma formation assay

Teratoma formation was performed as previously described [20]. In brief, nearly 10^9 haSCs were subcutaneously injected into the hind limbs of 8-week-old nude mice. After 5 weeks, fully formed teratomas were dissected and fixed with PBS containing 4% PFA, then embedded in paraffin, sectioned, and stained with hematoxylin and eosin (HE) for histological analysis.

Intracytoplasmic haSCs injection

Angus cattle and Dorper sheep ovaries were obtained from commercial slaughterhouses, and transported to the laboratory within 2 h at 28 °C in sterile saline solution (8.5 mg/mL NaCl) with a thermal container. The oocytes, which possessed at least three layers of cumulus, were selected for IVM according to the methods described above. Intracytoplasmic haSCs injection (iCHI) was performed following the published protocols with slight modifications [1, 2, 34]. Briefly, haSCs were arrested metaphase by culturing in FACE medium containing 0.05 µg/mL demecolcine for 12 h. Following centrifugation, cells were rinsed three times with PBS for removal of the demecolcine, and resuspended in the M2 medium containing 10 µM Y-27632 and 1.5% (w/v) polyvinylpyrrolidone (PVP). Each nucleus from haSCs was directly injected into an IVM oocyte using the Piezo micromanipulator (Primetech) in the M2 medium containing 5 µg/mL cytochalasin B (CB) on a 37 °C heating stage of an inverted microscope (Nikon). The reconstructed embryos were cultured in KSOM-AA medium for 1 hr and then activated using ionomycin followed by exposing them to 10 µg/mL cycloheximide (CHX) according to the methods described above.

Embryo Transfer

For bovine embryo transfer, blastocysts were stored in the TCM-199 medium (Thermo) supplemented with 15% FBS (Gibco). The iCHI blastocysts were transferred via the non-surgical transcervical method as previously described [37]. Briefly, a single blastocyst was transferred into the uterine horn of hormonally synchronised (CIDR progesterone + cloprostenol + gonadorelin) nulliparous heifer at day 7, estrus day 0 = day of micromanipulation). For ovine embryo transfer, 2-cell or 4-cell stage embryos were stored in TCM-199 medium (Thermo) supplemented with 10% BSA. The ovine embryos were transferred via surgical method using a TomCat cathether (Sovereign). Eight iCHI embryos were transferred into the oviduct of hormonally synchronized (CIDR progesterone + FSH + cloprostenol) recipient ewes at day 2 (estrus day 0 = day of micromanipulation). Pregnancy detection was diagnosed on day 35 and 50 of bovine and ovine embryonic development, respectively, using transrectal ultrasound EVO scanner (5.0MHz linear probe, E.I. Medical Imaging).

Reverse transcription-quantitative PCR

The reverse transcription PCR and real-time quantitative PCR reaction were performed using HiScript II One Step qRT-PCR SYBR Green Kit (Vazyme), and signals were detected with Applied Biosystems QuantStudio 3 (Thermo). Analysis of relative gene expression was normalized to the internal GAPDH gene, and measured using the $2^{(\Delta\Delta Ct)}$ method. The primer information was presented in Supplementary Table 5.

Bisulphite sequencing

Genomic DNA was extracted using the tissue and blood DNA extraction kit (Qiagen) and treated with the Methylamp DNA modification kit (EpiGentek). To obtain sperm genomic DNA, samples were pretreated with dithiothreitol (DTT) for 3 hr, following proteinase K lysis and phenol-chloroform extraction. The bisulfite conversion was performed as previously published studies [38]. The PCR products were cloned into vectors using TA/Blunt-Zero Cloning Kit (Vazyme), and individual clones were sequenced. The primer information was presented in Supplementary Table 5.

Plasmid construction

The coding sequence of Cas9 (H840A) :: M-MLVrt (D200N, T306K, W313F, T330P, L603W), derived from Prime editor 2 system (PE2, Addgene plasmid #132775) [15], was codon-optimized based on the general codon usage in cattle and sheep then synthesized by commercial company. The above vector was designated boPE2 and ovPE2. Indeed, "all-in-one" episomal PE plasmid is the same as epiCRISPR [39] except that the Cas9 was replaced by boPE2 or ovPE2. All CRISPR editor plasmids used in this work were assembled using the USER cloning method as previously described [15]. DNA amplification was conducted by Phanta Max Super-Fidelity DNA Polymerase (Vazyme). Integrate the synthesised DNA into the designated positions on the vector using ClonExpress Ultra One Step Cloning Kit V2 (Vazyme). The pegRNA targeting gene was designed by the online program (http://pegfinder.sidichenlab.org) [40] based on GenBank sequence.

Genome editing

Genome editing was performed as previously described with slight modification [39]. In brief, approximately 10^4 haSCs were disassociated using TrypLE (Thermo), and cultured in a 6-well plate with FACE medium for 24 h. A total of 2 μg ePE plasmid were transfected into haSCs using Lipomaster 3000 Transfection Reagent (Vazyme) following the manufacturer's protocol. After transfection 72 h, haSCs were selected by puromycin (0.2-0.5 μg/mL). For analysis of single cell-derived clones, the haSCs were disaggregated into single cells at day 15 post-transfection, and plated them onto new feeder layers with puromycin-free FACE medium for 14 days. Finally, individual

441 colonies were picked and genotyped. The pegRNA information was presented in
442 Supplementary Table 6.

WGS and data processing

The whole genome sequencing (WGS) was performed as described previously [34,41].
Briefly, libraries were generated from 1 mg of genomic DNA using the TruSeq DNA
Sample Preparation Kit (Illumina), and sequenced on the HiSeq 4000 system (Illumina)
by Lc-Bio technologies Co., Ltd. For data processing, the copy number variation (CNV)
detection was performed as previously described [42].
The template genome was downloaded from NCBI RefSeq, including cattle (ARS-
UCD2.0, https://www.ncbi.nlm.nih.gov/datasets/genome/GCA_002263795.4/) and
sheep genome (ARS-UI_Ramb_v3.0,
https://www.ncbi.nlm.nih.gov/datasets/genome/GCA_016772045.2/).

WGBS and data processing

The whole genome bisulfite sequencing (WGBS) was performed according to
previously [34, 43]. Briefly, genomic DNA was extracted from haSCs using the DNeasy B
lood & Tissue extraction kit (Qiagen) following end repair and A-tailing. The DNA was
fragmented into around 200-300 bp by ultrasonicator (Covaris). Bisulfite-converted
DNA was captured using EZ-96 DNA Methylation-DirectTM MagPrep (Zymo)
according to the manufacturer's instructions. Library sequencing was performed on the
HiSeq 4000 system (Illumina) by Lc-Bio Technologies Co., Ltd. The cleaned data were
aligned to the cattle (ARS-UCD2.0) reference with Bismark/Bowtie2 mode.
Bismark2summary command was used to calculate the genome CpG methylation level.
RnBeads R package was used to calculate the percentage of methylation.

CUT&Tag and data processing

The CUT&Tag was performed using the Hyperactive Universal CUT&Tag Assay Kit for
Illumina (Vazyme) according to the manufacturer's instructions. Paired-end
sequencing was further performed on the HiSeq 4000 system (Illumina) by Frasergen

Co., Ltd. The data processing was performed as described previously [44]. Briefly,

paired-end reads were aligned using Bowtie2 (2.2.5) with options: -local-very-
sensitive-local–no-unal–no-mixed–no-discordant-phred33 -I 10 -X 700. For peak
calling, the parameters used were macs2 callpeak -t input_file -p 1e-5 -f BEDPE/BED
-keep-dup all -n out_name. The antibody information was presented in Supplementary
Table 4.

buRNA-seq and data processing

478 The bulk RNA sequencing (buRNA-seq) was followed by previously published studies
479 [20, 35]. Briefly, total RNA was extracted from haSCs using TRIzol Plus RNA Purification
480 Kit (Thermo) following the manufacturer's instructions, and the RNA integrity was
481 checked using Bioanalyzer 2100 (Agilent). The total cDNA library was then amplified
482 by 18–20 cycles for library construction, which was performed following the

manufacturer's instructions (Illumina, USA). Sequencing was further performed on the HiSeq 4000 system (Illumina) by Frasergen Co., Ltd. After removing low-quality reads and adapters, the raw reads were mapped to the cattle (ARS-UCD2.0) or sheep (ARS-UI_Ramb_v3.0) genome using Tophat (v1.3.3) with default parameters [45]. Expression levels for all RefSeq transcripts were quantified to fragments per kilobase of exon model per million mapped reads (FPKM) using Cufflinks (v1.2.0) [46]. Gene Ontology (GO) analyses were performed using R package GO.db and Database for Annotation, Visualization, and Integrated Discovery (DAVID) [47].

scRNA-seq and data processing

The single-cell RNA sequencing (scRNA-seq) was followed by previously published studies [43, 48]. Briefly, haSCs or E8~E14 embryos were dissociated into single cells with TrypLE (Thermo) and resuspended in 1 Ca^{2+}-Mg^+ free PBS containing 0.04% BSA. Next, 300-1000 living cells were loaded on a Chromium Single Cell Controller (10x Genomics) to generate single-cell gel beads in emulsion (GEMs) cDNAs. Then, recovered cDNAs were used to construct a library using Single Cell 3' Library and Gel Bead Kit V2 (10x Genomics) according to the manufacturer's instructions. Library sequencing was performed on the HiSeq 4000 system (Illumina) with pair-end 150 bp (PE150) by Lc-Bio technologies Co., Ltd. The raw sequencing data were aligned and quantified using the Cell Ranger (10x Genomics) pipeline against the cattle (ARS-UCD2.0) reference genome. The Seurat R package was used to merge the scRNA-seq and cell clustering. Gene expression was calculated by the LogNormalize method in the NormalizeData function. RunPCA function was used in principal component analysis (PCA) with the most variable genes. Cell clusters were identified by a shared nearest neighbor (SNN) modularity optimisation based on a clustering algorithm with the FindClusters function. The *t*-SNE (*t*-distributed stochastic neighbor embedding) dimensionality reduction was generated using the RunTSNE function and the UMAP (uniform manifold approximation and projection) was performed using the RunUMAP function. Differentially expressed genes were identified using FindAllMarkers, and the top 30 differentially expressed genes (DEGs) were marked as signature genes. Heatmaps were plotted by the DoHeatmap function.

Data availability

All WGBS, CUT&Tag, buRNA-Seq, scRNA-seq, WGS data are available through GEO under the accession numbers GSE250491, GSE250493, GSE250494, GSE250495, and GSE250496. In addition, the SuperSeries GSE250497 links all the above data sets together.

Acknowledgements

We thank Hong Wang (Inner Mongolia University) for technical help. This study was supported by the National Natural Science Foundation of China (32360837, 32341052), Scientific and Technological Innovation 2030 (2023ZD0404803), Inner Mongolia Open

Competition Projects (2022JBGS0025), Inner Mongolia Science and Technology

Leading Team (2022LJRC0006), Inner Mongolia Science and Technology Major

Projects (2021ZD0009, 2021ZD0008, 2022ZD0008, 2023KJHZ0028), Inner Mongolia

Young Talents Projects (NJYT23138), Inner Mongolia Natural Science Foundation

(2023MS03004), Central Government Guides Development (2022ZY0212), National

Agricultural Science and Technology Project (NK2022130203), Collaborative

Innovation among Universities in Hohhot (XTCX2023-06), Ministry of Education

Engineering Centre Project (JYBGCSYS2022).

Author contributions

535 L.Y., A.D., X.L., L.S., D.W., S.W., Z.H., L.B., C.B., G.S., Z.W., L.Z. performed

536 experiments. L.Y., G.L., S.G. designed experiments, L.Y. and G.L. wrote the

537 manuscript.

Competing interests

540 The authors declare no competing interests.

References

16. Ross, P.J. & Cibelli, J.B. Bovine somatic cell nuclear transfer. *Methods Mol Biol* **636**, 155-177 (2010).

17. Czernik, M., Iuso, D., Toschi, P., Khochbin, S. & Loi, P. Remodeling somatic nuclei via exogenous expression of protamine 1 to create spermatid-like structures for somatic nuclear transfer. *Nat Protoc* **11**, 2170-2188 (2016).

18. Zhang, J. et al. Dissecting the molecular features of bovine-arrested eight-cell embryos using single-cell multi-omics sequencingdagger. *Biol Reprod* **108**, 871-886 (2023).

19. Alberio, R., Motlik, J., Stojkovic, M., Wolf, E. & Zakhartchenko, V. Behavior of M-phase synchronized blastomeres after nuclear transfer in cattle. *Mol Reprod Dev* **57**, 37-47 (2000).

20. German, S.D., Lee, J.H., Campbell, K.H., Sweetman, D. & Alberio, R. Actin depolymerization is associated with meiotic acceleration in cycloheximide-treated ovine oocytes. *Biol Reprod* **92**, 103 (2015).

21. Yang, L. et al. Transient Dux expression facilitates nuclear transfer and induced pluripotent stem cell reprogramming. *EMBO Rep* **21**, e50054 (2020).

22. Zhou, Q. et al. Generation of fertile cloned rats by regulating oocyte activation. *Science* **302**, 1179 (2003).

23. Kinoshita, M. et al. Pluripotent stem cells related to embryonic disc exhibit common self-renewal requirements in diverse livestock species. *Development* **148** (2021).

24. Ying, Q.L. et al. The ground state of embryonic stem cell self-renewal. *Nature* **453**, 519-523 (2008).

25. Irie, N. et al. SOX17 is a critical specifier of human primordial germ cell fate. *Cell* **160**, 253-268 (2015).

26. Takashima, Y. et al. Resetting transcription factor control circuitry toward ground-state pluripotency in human. *Cell* **158**, 1254-1269 (2014).

27. Theunissen, T.W. et al. Systematic identification of culture conditions for induction and maintenance of naive human pluripotency. *Cell Stem Cell* **15**, 471-487 (2014).

28. Bao, S. et al. Derivation of hypermethylated pluripotent embryonic stem cells with high potency. *Cell Res* **28**, 22-34 (2018).

29. Yang, Y. et al. Derivation of pluripotent stem cells with in vivo embryonic and extraembryonic potency. *Cell* **169**, 243-257 e225 (2017).

30. Yang, J. et al. Establishment of mouse expanded potential stem cells. *Nature* **550**, 393-397 (2017).

31. Gao, X. et al. Establishment of porcine and human expanded potential stem
cells. *Nat Cell Biol* **21**, 687-699 (2019).
32. Bogliotti, Y.S. et al. Efficient derivation of stable primed pluripotent embryonic
stem cells from bovine blastocysts. *Proc Natl Acad Sci U S A* **115**, 2090-2095
(2018).
33. Vilarino, M. et al. Derivation of sheep embryonic stem cells under optimized
conditions. *Reproduction* **160**, 761-772 (2020).
34. Zhao, L. et al. Establishment of bovine expanded potential stem cells. *Proc*
Natl Acad Sci U S A **118** (2021).
35. De Los Angeles, A., Okamura, D. & Wu, J. Highly efficient derivation of
pluripotent stem cells from mouse preimplantation and postimplantation
embryos in serum-free conditions. *Methods Mol Biol* **2005**, 29-36 (2019).
36. Ludwig, T.E. et al. Feeder-independent culture of human embryonic stem
cells. *Nat Methods* **3**, 637-646 (2006).
37. Zhang, X.M. et al. In vitro expansion of human sperm through nuclear
transfer. *Cell Res* **30**, 356-359 (2020).
38. Zhong, C. et al. Generation of human haploid embryonic stem cells from
parthenogenetic embryos obtained by microsurgical removal of male
pronucleus. *Cell Res* **26**, 743-746 (2016).
39. Elling, U. et al. Derivation and maintenance of mouse haploid embryonic stem
cells. *Nat Protoc* **14**, 1991-2014 (2019).
40. Yang, L., Song, L., Liu, X., Bai, L. & Li, G. KDM6A and KDM6B play
contrasting roles in nuclear transfer embryos revealed by MERVL reporter
system. *EMBO Rep* **19** (2018).
41. Yang, H. et al. Generation of genetically modified mice by oocyte injection of
androgenetic haploid embryonic stem cells. *Cell* **149**, 605-617 (2012).
42. Li, W. et al. Androgenetic haploid embryonic stem cells produce live
transgenic mice. *Nature* **490**, 407-411 (2012).
43. Owen, J.R. et al. One-step generation of a targeted knock-in calf using the
CRISPR-Cas9 system in bovine zygotes. *BMC Genomics* **22**, 118 (2021).
44. Hajkova, P. et al. DNA-methylation analysis by the bisulfite-assisted genomic
sequencing method. *Methods Mol Biol* **200**, 143-154 (2002).
45. Anzalone, A.V. et al. Search-and-replace genome editing without double-
strand breaks or donor DNA. *Nature* **576**, 149-157 (2019).
46. Xie, Y. et al. An episomal vector-based CRISPR/Cas9 system for highly
efficient gene knockout in human pluripotent stem cells. *Sci Rep* **7**, 2320
(2017).
47. Chow, R.D., Chen, J.S., Shen, J. & Chen, S. A web tool for the design of
prime-editing guide RNAs. *Nat Biomed Eng* **5**, 190-194 (2021).

48.   Patch, A.M. et al. Whole-genome characterization of chemoresistant ovarian
620   cancer. *Nature* **521**, 489-494 (2015).

621   49.   Xie, C. & Tammi, M.T. CNV-seq, a new method to detect copy number
622   variation using high-throughput sequencing. *BMC Bioinformatics* **10**, 80
623   (2009).

624   50.   Shen, H. et al. Mouse totipotent stem cells captured and maintained through
625   spliceosomal repression. *Cell* **184**, 2843-2859 e2820 (2021).

626   51.   Kaya-Okur, H.S. et al. CUT&Tag for efficient epigenomic profiling of small
627   samples and single cells. *Nat Commun* **10**, 1930 (2019).

628   52.   Trapnell, C., Pachter, L. & Salzberg, S.L. TopHat: discovering splice junctions
629   with RNA-seq. *Bioinformatics* **25**, 1105-1111 (2009).

630   53.   Trapnell, C. et al. Transcript assembly and quantification by RNA-seq reveals
631   unannotated transcripts and isoform switching during cell differentiation. *Nat*
632   *Biotechnol* **28**, 511-515 (2010).

633   54.   Huang da, W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative
634   analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*
635   **4**, 44-57 (2009).

636   55.   Zhao, T. et al. Single-cell RNA-seq reveals dynamic early embryonic-like
637   programs during chemical reprogramming. *Cell Stem Cell* **23**, 31-45 e37
638   (2018).

 Figure legends

Fig.1 | *De novo* derivation of ruminant haSCs with formative features.

a. Schematic overview of haSCs derivation, haploid embryos were generated by

injecting sperm into enucleated oocytes.

b. Morphology of ruminant haSCs derived from haploid blastocysts.

c. The effects of FACE medium on haploidy maintenance of ruminant haSCs with

FACS enrichment.

d. Immunofluorescence images showing that ruminant haSCs expressed core

pluripotency markers (SOX2 and OCT4) and formative pluripotency marker

(OTX2).

e. UMAP projection of b-haSCs, colored by pluripotency-associated marker genes.

f. Heatmap and bubble displaying the expression of formative and primed marker

genes in different samples.

g. Heatmap of H3K4me3 and H3K27me3 signals in b-haSCs and primed-b-SCs.

h. Ternary plot showing cross-species comparison of orthologous transcription

factors, formative-pluripotency related genes.

i. Chimeric mouse generated by EGFP-labeled ruminant haSCs at E18.5.

Scale bars: b, 100 µm; d, 25 µm; n ≥ 3 independent experiments; B, bovine; O,

ovine; M, mouse.

Fig.2 | Generation of MSTN-knockout ruminants using the protaminized haSCs.

a. Morphology of ruminant blastocysts derived by iCHI, Pro-iCHI, and IVF.

b. Bisulfite pyrosequencing analysis of paternally imprinted regions. The filled and

open squares represent methylated and unmethylated CpG sites, respectively.

c. Immunostaining for H3K4me3, H3K9me3, and H3K27me3 of ruminant blastocysts

derived by iCHI, Pro-iCHI, and IVF.

d. Image of ovine fetus at 98 days from MSTN-edited o-haSCs.

e. Representative images of WT and MSTN-edited cattle through the Pro-iCHI

approach.

Scale bars: a, 100 µm; c, 25 µm; n ≥ 3 independent experiments; B, bovine; O,

ovine; ♀, female; ♂, male.

Figure 1

Figure 2

a

B — IVF | iCHI | Pro-iCHI
Blastocyst rate: 68.8%   27.1%   64.9%

O — IVF | iCHI | Pro-iCHI
Blastocyst rate: 53.7%   24.2%   51.2%

b

B — IGF2/H19-DMR
haESCs   90.2%
sperm   92.2%
fibroblast   51.0%

O — IGF2/H19-DMR
haESCs   80.7%
sperm   92.1%
fibroblast   53.5%

c

B — IVF | iCHI | Pro-iCHI
DAPI K4me3
DAPI K9me3
DAPI K27me3

O — IVF | iCHI | Pro-iCHI
DAPI K4me3
DAPI K9me3
DAPI K27me3

d

O — MSTN G > A

e

B — WT | MSTN$^{+/-}$ #1 | #2