Progressive overfilling of readily releasable pool underlies short-term facilitation

at recurrent excitatory synapses in layer 2/3 of the rat prefrontal cortex

Jiwoo Shin[1,2], Seung Yeon Lee[2], Yujin Kim[1,2*], Suk-Ho Lee[1,2,*].

1. Department of Brain and Cognitive Science, College of Natural Sciences, Seoul National

University.   2. Department of Physiology, Seoul National University College of Medicine.

03080, Seoul, South Korea.

corresponding authors

Yujin Kim Ph.D. (kim.y@snu.ac.kr)

Cell Physiology Lab

Department of Physiology

Seoul National Univ. College of Medicine

Suk-Ho Lee M.D., Ph.D. (leesukho@snu.ac.kr)

Cell Physiology Lab

Department of Physiology

Seoul National Univ. College of Medicine

103 Daehakro, 03080, Seoul, South Korea

Phone) +82-2-740-8222

Abstract

Short-term facilitation of recurrent excitatory synapses within the cortical network has been proposed to support persistent activity during working memory tasks, yet the underlying mechanisms remain poorly understood. We characterized short-term plasticity at the local excitatory synapses in layer 2/3 of the rat medial prefrontal cortex and studied its presynaptic mechanisms. Low-frequency stimulation induced slowly developing facilitation, whereas high-frequency stimulation initially induced strong depression followed by rapid facilitation. This non-monotonic delayed facilitation after a brief depression resulted from a high vesicular fusion probability and slow activation of Ca^{2+}-dependent vesicle replenishment, which led to the overfilling of release sites beyond their basal occupancy. Pharmacological and gene knockdown (KD) experiments revealed that the facilitation was mediated by phospholipase C/diacylglycerol signaling and synaptotagmin 7 (Syt7). Notably, Syt7 KD abolished facilitation and slowed the refilling rate of vesicles with high fusion probability. Furthermore, Syt7 deficiency in layer 2/3 pyramidal neurons impaired the acquisition of trace fear memory and reduced c-Fos activity. In conclusion, Ca^{2+}- and Syt7-dependent overfilling of release sites mediates synaptic facilitation at L2/3 recurrent excitatory synapses and contributes to temporal associative learning.

Introduction

Synapses in the mammalian central nervous system undergo activity-dependent modulation of synaptic weight during repetitive stimulation, known as short-term synaptic plasticity (STP). STP is mediated by presynaptic mechanisms that regulate the release probability (p_r), size of the readily releasable pool (RRP), and kinetics of vesicle replenishment of RRP (Zucker and Regehr, 2002; Jackman and Regehr, 2017; Neher and Brose, 2018). Short-term depression has been observed at synapses with a high baseline p_r, primarily because of the rapid depletion of releasable vesicles during high-frequency stimulation (HFS) (Lin et al., 2022). Conversely, vesicle depletion is less pronounced at synapses with a low baseline p_r, and cumulative increases in p_r during HFS are considered to mediate short-term facilitation (Rozov et al., 2001; Pan and Zucker, 2009; Aldahabi et al., 2024). Central synapses exhibit a wide spectrum of STP depending on different combinations of presynaptic factors mediating short-term depression and facilitation.

Previously, facilitation was modeled as an increase in p_r after an action potential (AP) and its slow decay during the inter-spike interval (ISI), resulting in a cumulative increase in p_r during HFS (Markram et al., 1998; Dittman et al., 2000). Recent studies have suggested that the number of docking (or release) sites in an active zone is limited and only partially occupied by releasable vesicles in the resting states (Miki et al., 2016; Pulido and Marty, 2018; Malagon et al., 2020; Lin et al., 2022). Therefore, p_r is determined by the vesicular fusion probability (p_v) and release site occupancy (p_{occ}). However, distinguishing the individual contributions of p_v and p_{occ} to p_r is challenging (Silva et al., 2021; Neher, 2024). It is also unclear whether facilitation is mediated by activity-dependent increases in the p_v, p_{occ}, or both.

Two recent models for vesicle priming and release view the 'priming' as a two-step reversible

process that leads to a release-ready state. The release-ready vesicles could be vesicles in a

tightly docked state (TS) or those occupying specialized docking sites (DS), and are supplied

from vesicles in a loosely docked state (LS) or those occupying a distinct replacement sites

(RS), which are called LS/TS model ((Neher, 2024); (Aldahabi et al., 2024)) and RS/DS model

((Miki et al., 2016); (Silva et al., 2021)), respectively. Common to both models is the

assumption that release-ready vesicles constitute only a fraction of all docked vesicles under

resting conditions and that this fraction is up- and down-regulated by synaptic activity.

Synaptotagmin-7 (Syt7) was recently found to play a crucial role in facilitation at different

types of central synapses (Jackman et al., 2016; Chen et al., 2017; Martinetti et al., 2022). In

addition to its role in facilitation, Syt7 accelerates refilling of the RRP after depletion and

mediates asynchronous release (Bacaj et al., 2013; Liu et al., 2014; Jackman et al., 2016;

Tawfik et al., 2021). Syt7-dependent synaptic facilitation is interpreted as an activity-

dependent increase in p_v (Turecek et al., 2016; Jackman and Regehr, 2017; Turecek et al., 2017;

Norman et al., 2023). However, it is unclear whether Syt7 contributes to facilitation through

a Ca^{2+}-dependent increase in p_{occ} (i.e., overfilling). Overfilling has been proposed as a

mechanism supporting facilitation at the Drosophila neuromuscular junction (Kobbersmed et

al., 2020), yet whether this process also occurs at the mammalian central synapses mediated

by Syt7 has not been studied.

Facilitation at recurrent excitatory synapses in cortical recurrent networks may play a crucial

role in maintaining working memory (Mongillo et al., 2008; Mongillo et al., 2012; Hansel and

Mato, 2013). Although layer 2/3 (L2/3) pyramidal cells (PCs) are highly interconnected and

represent task-relevant information during a delay period in working memory tasks (Fujisawa

et al., 2008; Ozdemir et al., 2020), most studies on local excitatory recurrent synapses in the

neocortex have focused on layer 5 (L5) exhibiting short-term depression (Reyes and Sakmann,

1999; Hempel et al., 2000; Yoon et al., 2020). Further, the synaptic properties of excitatory

synapses in L2/3 remain poorly understood.

In the present study, we found that HFS of local excitatory synapses in L2/3 of the medial

prefrontal cortex (mPFC) induced initial strong depression followed by delayed facilitation,

whereas low-frequency stimulation induced slow monotonous facilitation. This unique form

of STP could be explained by 1) high p_v of RRP vesicles at rest and throughout a train

stimulation; 2) the low resting vesicular occupancy of RRP and its activity-dependent increase.

These conditions led to delayed facilitation through progressive overfilling of release sites

during HFS with high p_v vesicles beyond the resting occupancy. Syt7 knockdown (KD) in L2/3

PCs completely abolished these synaptic enhancements by inhibiting the Ca^{2+}-dependent

increase in release site occupancy. Finally, behavioral tests for trace fear conditioning showed

that Syt7 KD impaired the acquisition of trace fear memory and reduced c-Fos expression in

the mPFC. Collectively, these results support that Ca^{2+}- and Syt7-dependent overfilling of

release sites mediates synaptic facilitation at L2/3 recurrent excitatory synapses and

contributes to temporal associative learning.

Results

Frequency dependence of STP at local excitatory synapses in L2/3 of the prelimbic cortex

We examined STP profiles at recurrent excitatory synapses between L2/3 PCs and at synapses from L2/3-PCs to fast-spiking interneurons (FSINs) in the prelimbic area of the rat mPFC. The PCs and FSINs were identified based on their distinct intrinsic properties and morphologies (Fig. S1). To stimulate presynaptic PCs, L2/3 PCs were transfected with a plasmid encoding oChIEF using *in utero* electroporation (IUE; Fig. 1A; Lee et al., 2024). Photostimulation of the oChIEF-expressing cell soma at 40 Hz consistently evoked single APs with high fidelity throughout the 600-pulse trains (Fig. S2A-B), which was consistent with little use-dependent inactivation of oChIEF (Lin et al., 2009; Hass and Glickfeld, 2016).

Given that IUE transfected approximately 10-20% of L2/3 PCs (Fig. 1A), we made a whole-cell patch on a non-transfected L2/3 PC or an FSIN (Fig. 1C). To stimulate a minimal number of excitatory axon fibers, EPSCs were evoked using minimal optical stimulation (Fig. S3A; 470 nm, 3-4 μm in radius and 3-5 ms in duration). EPSCs evoked by minimal stimulation were examined to determine the uniformity of the kinetic properties (Fig. S3B-C). Moreover, we reassessed the kinetics of EPSCs during trains *post hoc* to ensure uniformity. EPSCs were evoked by 20-pulse trains at 4 different frequencies (5, 10, 20, and 40 Hz) to cover the range of firing rates of PCs observed *in vivo* during working memory tasks (Baeg et al., 2003). Excitatory synapses from PCs onto both other PCs and FSINs (PC–PC and PC–FSIN) showed strong (~2-fold enhancement) and lasting (several seconds) short-term facilitation at 20 Hz and lower

frequencies (Fig. 1D-E). The PC-PC synapses exhibited stronger facilitation than did the PC-

FSIN synapses.

Steady-state EPSCs at the PC-PC synapses were not significantly different from 5 to 20 Hz (Fig.

1F). Frequency invariance at the PC-PC synapses suggests that Ca^{2+}-dependent synaptic

facilitation may counteract vesicle depletion in a frequency-dependent manner (Turecek et

al., 2017; Lin et al., 2022). Despite facilitation at all frequencies, the paired pulse ratio (PPR)

rapidly decreased as the ISI decreased (Fig. 1G, I). At 40 Hz stimulation, both PC-PC and PC-

FSIN synapses initially underwent strong depression then followed by facilitation in a few

stimuli (Fig. 1Dd, Ed). Such delayed facilitation after paired pulse depression (PPD) has not

been previously observed in central synapses except for a slight increase in EPSC at the

cerebellar synapses under artificially depolarized conditions (Pulido and Marty, 2018).

We confirmed that optically evoked EPSCs were strictly dependent on AP generation (Fig.

S4A), suggesting that EPSCs are evoked by optical stimulation of presynaptic axon fibers

rather than by direct depolarization of presynaptic boutons. Additionally, dual patch-clamp

techniques showed that the STP of optically evoked EPSCs did not differ from the electrically

evoked STP of EPSCs at the PC-FSIN synapses (Fig. S4B). Finally, we confirmed that AMPA

receptor (AMPAR) desensitization was not responsible for PPD at 40 Hz, ruling out the

possible effects of postsynaptic factors on STP (Fig. S5A), and that AMPAR saturation was not

significant during facilitation (Fig. S5B).

Delayed facilitation results from slow activation of Ca^{2+}-dependent vesicle replenishment at a

constantly high vesicular fusion probability

The strong PPD at 40 Hz suggests a high vesicular fusion probability (p_v) and likely tight

coupling of the releasable vesicles to a Ca^{2+} source. To test this hypothesis, we examined the

effects of EGTA, a slow calcium chelator, on synaptic transmission at the PC-PC synapses. The

baseline EPSC amplitude (EPSC$_1$) was not affected by incubation of the slice in the bath

solution containing 50 µM EGTA-AM for longer than 20 min, supporting tight coupling.

Moreover, the PPR was decreased to 0.19, and subsequent facilitation slowed but was still

distinct (Fig. 2A-B). These findings indicate that the initial p_v should be higher than 0.81

considering vesicle recruitment during ISI. The effects of EGTA-AM were confirmed in the

same cell by measuring the baseline EPSC and PPR before and after applying EGTA-AM to the

bath. EPSC$_1$ remained unchanged, while PPR was reduced by half (Fig. S6).

Notably, the application of EGTA-AM markedly slowed facilitation at 40 Hz (Fig. 2A,C). The

frequency invariance of steady-state EPSCs (Fig. 1F) and the slower facilitation at 40 Hz in the

presence of EGTA (Fig. 2A) imply a Ca^{2+}-dependent increase in the refilling rate of the RRP

and/or an increase in the p_v of reluctant vesicles in light of previous studies (Hosoi et al., 2007;

Liu et al., 2014; Ritzau-Jost et al., 2014; Turecek et al., 2016; Ritzau-Jost et al., 2018; Kusick et

al., 2020). Moreover, facilitation was accelerated at higher stimulation frequencies. The plot

of the rate constant for the increase in EPSCs (denoted as $\mathbf{k}_{STF}$, the rate constant for the

development of short-term facilitation) as a function of the stimulation frequency (f_{stim})

revealed a linear relationship (Fig. 2C). $\mathbf{k}_{STF}$ in the presence of EGTA-AM was distinctly below

this relationship, supporting the Ca^{2+}-dependent acceleration of facilitation.

k_{STF} was also accelerated when the stimulation frequency was increased from 5 Hz to 40 Hz

(Fig. 2D; 1.27 ± 0.18/s to 24.27 ± 3.44/s). Notably, the second 40-Hz train stimulation led to

strong PPD, followed by accelerated facilitation (Fig. 2D,E). The strong PPD suggests that the

vesicular fusion probability kept high during the 5-Hz stimulation, arguing against the

possibility for an increase in p_v. Both two-step sequential priming models, LS/TS and RS/DS

models, are suitable to interpret our data and we will refer to the different populations of

vesicles as TS and LS vesicles, respectively (Taschenberger et al., 2016; Neher and Brose, 2018;

Lin et al., 2022; Neher, 2024). Within this framework, vesicles are are released from TS with

high p_v both in the resting state and during facilitation, and the delayed facilitation is seen as

a result of Ca^{2+}-dependent progressive increase in the occupancy of TS vesicles on docking

sites.

Release sites have low baseline occupancy and these are increased during facilitation and post-

tetanic augmentation

Given that p_v was greater than 0.8, the two-fold increase in EPSC during 20 pulse train

stimulations could not be explained by an increase in p_v. Recent studies have shown that

vesicle release sites (N) are limited in number and are partially occupied at rest (Miki et al.,

2016; Pulido and Marty, 2018; Malagon et al., 2020; Lin et al., 2022). Under such a high p_v,

two-fold synaptic facilitation should be attributed to an increase in p_{occ}. The vesicle

replenishment rate is accelerated during HFS in a Ca^{2+}-dependent manner (Fig.2A-C; Hosoi et

al., 2007). This may lead to the overfilling of release sites beyond their basal vesicular

occupancy, which in turn may increase EPSC under high p_v. To test this hypothesis, we

determined the quantal parameters at the L2/3 local excitatory synapses. We first measured

the quantal size from asynchronous release events after replacing extracellular Ca^{2+} with Sr^{2+},

as previously described (Yoon et al., 2020) (Fig. S7A-B). The mean and coefficient of variance

of the quantal sizes were 24 pA and 0.25 at PC-PC synapses, respectively, and were 30 pA and

0.31 at PC-FSIN synapses, (Fig. S7C).

Next, we performed a variance-mean (V-M) analysis of phasic EPSCs at excitatory synapses

during train stimulation at 5 Hz and 40 Hz (Scheuss et al., 2002). Applying multiple-probability

fluctuation analysis (MPFA) using intra-site quantal variability of 0.2 (Valera et al., 2012), the

V-M plot was best fitted by the number of release sites (N) of 5.3 and the quantal size (q) of

25 pA at the PC-PC synapses (Fig. 3A). The p_r for $EPSC_1$ was calculated as 0.32. We performed

the same analysis for the PC-FSIN synapses and found N = 5.3, q = 35 pA, and p_r for $EPSC_1$ =

0.31 (Fig. 3B). Our estimate for N was smaller but comparable to the estimate (6.9) at

excitatory synapses in L2/3 of the mouse barrel cortex (Holler et al., 2021).

The p_r estimate was clearly smaller than the possible minimum value for p_v (0.81) estimated

from the PPR data (Fig. 2B). Given that p_r is a product of p_{occ} and p_v (Neher, 2024), a p_r smaller

than p_v indicates the partial occupancy of release sites in the resting state. If synaptic

facilitation were mediated by the recruitment of new release sites, it would be accompanied

by an increase in the variance of augmented EPSCs (Valera et al., 2012; Kobbersmed et al.,

2020). However, the augmented EPSCs during facilitation followed the same parabola in the

V-M plot (Fig. 3A,B). Therefore, facilitation appears to be mediated by the progressive

overfilling of a finite number of release sites beyond the basal occupancy. In the next section,

we show that p_v is close to unity based on a synaptic failure analysis (Fig. S8). Assuming this,

the baseline p_r at the first pulse estimated from the fitted parabolic curve can be largely

regarded as a baseline p_{occ} (c.a. 30%), and the two-fold increase in EPSCs during facilitation

can be attributed to an increase of up to 60% in p_{occ}.

We investigated the time course of recovery from facilitation and post-tetanic augmentation

(PTA) by applying dual-train stimulations (30 pulses at 40 Hz) separated by different inter-

burst intervals (IBIs; Fig. 3C,D). The initial EPSCs in the second train were enhanced by more

than twofold, and followed by a strong PPD, suggesting that fusion probability stays high in

the IBIs and thus synaptic transmission is carried by TS vesicles. The decay time course of PTA

was characterized by two phases (initially fast and later slow; Fig. 3E), which may reflect a

two-phase Ca^{2+} decay or two distinct recovery processes that occur sequentially. The variance

of potentiated EPSCs measured at IBIs of 0.1, 0.2, and 0.5 s was not different from that of

EPSCs at the peak of facilitation (filled circles, Fig. 3A-B). Given that the same RRP mediates

both facilitation and augmentation and that post-tetanic EPSCs is carried by TS vesicles similar

to the baseline EPSCs, this plot suggests that PTA, similar to short-term facilitation, is

mediated by an increase in the TS vesicle occupancy.

Failure rate analysis suggests that vesicular fusion probability is closed to one

The vesicle dynamics of STP has been explained using a simple refilling model (Hosoi et al.,

2007; Neher and Sakaba, 2008). This model assumes that the vesicles are reversibly docked

to a limited number of release sites (N) with forward and reverse rate constants (denoted as

k_1 and b_1, respectively). Under the framework of the simple refilling model (see *Materials and*

Methods), the rate constant for the refilling of the RRP after depletion by the first pulse is the

sum of the baseline k_1 (k_1) and b_1. This is estimated as 23/s from the plot of PPR as a function

of ISIs (Fig. S8A). From the baseline occupancy of 0.3 and $p_{occ} = k_1 / (k_1 + b_1)$, we estimated k_1

to be 6.9/s.

When paired APs were applied with an ISI of 20 ms, the failure rate at 1^{st} pulse, $P(F_1)$, was

10.6%, and the probability of two consecutive failures, $P(F_1, F_2)$, was 6.2% (Fig. S8A). From the

equation $P(F_1) = (1 - p_r)^N$, p_r was calculated as 0.312 when N = 6, comparable to p_r (= 0.32)

estimated from the V-M analysis (Fig. 3A). We calculated $P(F_1, F_2)$ under the framework of a

simple refilling model, with p_v and k_1 set as free parameters and other parameters set

according to the following relationships: $P(F_1) = (1 - p_r)^N$, $p_r = p_v \cdot p_{occ}$, and $p_{occ} = k_1 / (k_1 + b_1)$.

The calculated values for $P(F_1, F_2)$ and their difference from the observed value (6.2%) are

shown in the plane of k_1 *vs.* p_v in Figure S8B-C. As previously mentioned, k_1 was greater than

6/s. Under this high k_1 condition, the difference between the calculated and observed values

of $P(F_1,F_2)$ was minimal when $p_v = 1$ (Fig. S8C).

Pharmacological manipulations reveal specific molecular identities underlying vesicle loading

processes

Our results suggest that the baseline occupancy of TS vesicles is low and that the activity-

dependent progressive overfilling of docking sites with TS vesicles is responsible for short-

term facilitation and augmentation. To elucidate the specific molecular link between synaptic

enhancement and Ca^{2+}-dependent overfilling, we examined the effects of several drugs

known to affect vesicle dynamics. We first examined the roles of phospholipase C (PLC) and

diacylglycerol (DAG) by applying 5 μM U73122, a PLC inhibitor, or 20 μM 1-oleoyl-2-acetyl-sn-

glycerol (OAG), a DAG analog. The STP at 40 Hz and PTA at 0.5-s interval were examined before

and after applying each drug to the bath. For both drugs, the facilitation and augmentation

were reduced (Fig. 4A-B). U73122 did not affect the basal EPSC amplitude ($EPSC_1$) but resulted

in a stronger PPD, slower facilitation, and lower augmentation (Fig. 4A). These effects of

U73122 suggest the involvement of Ca^{2+}-induced PLC activation in progressive overfilling. In

contrast, OAG increased $EPSC_1$ while maintaining a pronounced PPD (Fig. 4B), indicating an

increase in the TS vesicle pool size. Meanwhile, OAG slowed k_{STF} and reduced augmentation,

indicating occlusion of activity-dependent synaptic facilitation. Phorbol esters and DAG are

thought to accelerate priming through Munc13 activation (Rhee et al., 2002; Taschenberger

et al., 2016; Aldahabi et al., 2022); they also shorten the bridges that interlink docked vesicles

with the active zone (AZ) membrane (Papantoniou et al., 2023). Collectively, these suggest

that PLC- and DAG-mediated signaling play a role in the overfilling of docking sites underlying

short-term facilitation and augmentation.

Although the acceleration of Ca^{2+}-dependent vesicle refilling supports rapid facilitation at 40

Hz, the later part of the delayed facilitation underwent a slight depression, which was more

pronounced during the second burst (Fig. 3D, Fig. 4). Sustained synaptic transmission is

limited by the spatial availability of docking sites because it is undermined by accumulation

of exocytic remainings in the presynaptic AZ during HFS (Neher, 2010; Haucke et al., 2011).

To verify this, we examined the effects of dynasore (100 μM), a specific inhibitor of

endocytosis (Macia et al., 2006). The treatment with dynasore did not influence $EPSC_1$ or the

early phase of facilitation, but it exacerbated depression during the later phase and

attenuated PTA (Fig. 4C). This suggests that late synaptic transmission during the 40-Hz train

is limited by site clearance.

Given the involvement of actin polymerization in the multiple steps of vesicle docking and

recycling, we examined the effects of 20 μM latrunculin B (LatB), an inhibitor of actin

polymerization. The effects of LatB were complicated: (1) $EPSC_1$ was decreased, whereas PPR

was increased (Fig. 4Da) and (2) facilitation and augmentation were reduced (Fig. 4Db). The

latter effect may result from defects in vesicle replenishment rates, either through the

impaired physical movement of vesicles or disrupted site clearance (Sakaba and Neher, 2003;

Lee et al., 2012; Hallermann and Silver, 2013; Miki et al., 2016), as Ca^{2+}-dependent vesicle

refilling mediates facilitation and augmentation (Figs. 2–3). Meanwhile, the former effect

might be attributed to a shift in equilibrium in the priming states towards a loosely docked

state (LS) at rest. It is unknown whether actin affects vesicular fusogenicity in neurons,

although it facilitates multiple steps of vesicle fusion by enhancing membrane tension in

neuroendocrine cells (Wu and Chan, 2022).

Short-term facilitation at both types of local excitatory synapses in L2/3 is abolished by Syt7

knockdown

We examined whether Syt7 mediates facilitation at the PC-PC and PC-FSIN synapses in L2/3

of the mPFC. To this end, oChIEF and shRNA against Syt7 (shSyt7) were co-expressed in L2/3

PCs using IUE. Syt7 KD resulted in a complete loss of facilitation at all tested stimulation

frequencies (Fig. 5A-B). Short-term depression and paired-pulse depression were more

pronounced as the stimulation frequency increased, and the frequency invariance of steady-

state EPSC was abolished, implying little contribution of Ca^{2+}-dependent vesicle recruitment

(Fig. 5C-F). The lack of facilitation could not be attributed to the failure of APs in presynaptic

axons, as optical stimulation elicited reliable APs in Syt7 KD PCs during the 40-Hz train (Fig.

S2C). Syt7 KD had little effect on the basal properties, including the peak amplitude,

rise/decay time, and time-to-peak, of single AP-induced EPSCs (Fig. S3D-E). Moreover, Syt7

KD did not significantly affect the intrinsic properties of neurons (Fig. S9).

To examine whether Syt7 KD had any nonautonomous effects on postsynaptic cells, we

measured the quantal size (q) from asynchronous events in the presence of Sr^{2+} and found

that the mean quantal size remained unaltered at Syt7 KD synapses (Fig. S7C). This indicated

that Syt7 did not influence postsynaptic parameters, including the density and kinetic

properties of AMPARs. To test the specificity of shSyt7, we examined whether co-expression

of the shRNA-resistant form of Syt7 restored synaptic facilitation in both excitatory synapse

types. Presynaptic overexpression of shRNA-resistant Syt7 under the CAG promoter rescued

the effect of shRNA at both PC-PC and PC-FSIN synapses (Fig. S10). These results underscore

the crucial role of presynaptic Syt7 in mediating short-term facilitation at local excitatory

synapses in L2/3 of the mPFC.

Syt7 KD synapses exhibit complementary changes in the number of release sites and their

vesicle occupancy

To estimate the quantal parameters in Syt7 KD synapses, we performed V-M analysis of EPSCs

evoked by train stimulation (Fig. 6A-B). Fitting a parabola to the V-M plot revealed higher

baseline p_{occ} in Syt7 KD synapses than in wild-type (WT) synapses for both synapse types (KD,

0.68 and 0.42; WT, 0.32 and 0.31 for PC-PC and PC-FSIN synapses, respectively). The P_{occ}

monotonously decreased during a train stimulation, as expected from short-term depression.

Moreover, the number of release sites (N) was markedly lower in Syt7 KD synapses (N = 3.5)

than in WT synapses (N = 5.3; Fig. 3A). Next, we examined the PTA induced by the same

protocol as in Figure 3 (two 40-Hz train stimulations separated by variable IBIs) at Syt7 KD

synapses (Fig. 6C). The 40 Hz train-induced synaptic depression quickly recovered during IBIs,

but no PTA was observed at Syt7 KD synapses, indicating its crucial role in augmentation (Fig.

6D-E).

TS vesicle recovery following depletion is accelerated by Syt7

Syt7 is recognized not only for its crucial role in facilitation, but also for mediating Ca^{2+}-

dependent replenishment of releasable vesicles (Liu et al., 2014; Tawfik et al., 2021). Given

that facilitation at L2/3 excitatory synapses is driven by progressive overfilling (Figs. 2-3), the

loss of facilitation at Syt7 KD synapses in the present study implies that Syt7 plays a key role

in the Ca^{2+}-dependent acceleration of refilling and overfilling of the docking sites with TS

vesicles. To test this hypothesis, we examined the recovery kinetics of EPSCs after TS vesicles

were depleted by 3 pulses at 40 Hz in wild-type (WT) and Syt7 KD synapses (Fig. 7). Strong

PPD was observed not only in the first burst, but also in the second burst. This suggested that

EPSCs in the second burst were mediated by TS vesicles, similar to those in the first burst (Fig.

7A,D). Therefore, $EPSC_1$ in the second burst can be interpreted as a release from TS vesicles

recovering during a given IBI. The recovery time course of $EPSC_1$ in the second burst indicated

full recovery of TS vesicles within 100 ms in both excitatory synapse types in WT rats (Fig.

7C,F). However, this recovery was remarkably slower in Syt7 KD synapses, requiring 5 s for

full recovery, indicating that TS vesicle recovery was greatly accelerated by Syt7. Overall,

synaptic facilitation and augmentation at local excitatory synapses in L2/3 can be ascribed to

progressive activity-dependent increases in the occupancy of TS vesicles (Figs. 2-3), in which

Syt7 plays a key role.

Behavioral consequences of Syt7 KD in L2/3 PCs of the mPFC

We previously proposed that the disparity in STP between the PC-PC and PC-IN synapses

results in activity-dependent changes in the ratio of synaptic weights at the PC-PC and PC-IN

synapses (J_{ee}/J_{ie}). This may in turn have a profound influence on persistent activity in a

recurrent network by biasing the E-I balance (Yoon et al., 2020). Syt7 KD not only eliminated

facilitation, but also the activity-dependent increase in the J_{ee}/J_{ie} ratio in the L2/3 network

(Fig. S11). This suggests the possibility that Syt7 contributes to the persistent activity during

working memory tasks in the L2/3 recurrent network. Trace fear conditioning (tFC) requires

associative learning between an auditory cue (conditioned stimulus, CS) and temporally

separate aversive events (unconditioned stimulus, US). The formation of trace fear memory

requires the prelimbic area of the mPFC and hippocampus. mPFC activity during the trace

interval is thought to temporarily hold CS information, enabling the network to associate

temporally discontinuous events (i.e., temporal associative learning) (Gilmartin et al., 2013).

Synaptic facilitation and augmentation are implicated in temporary memory retention in

recurrent networks (Mongillo et al., 2008; Mongillo et al., 2012).

Given that Syt7 is crucial for synaptic facilitation (Fig. 5) and augmentation (Fig. 6) in prelimbic

L2/3 recurrent excitatory synapses, we tested whether trace fear memory formation was

impaired in rats with depleted Syt7 transcripts, specifically in the L2/3 PCs of the mPFC. Syt7-

targeted shRNA (shSyt7) or scrambled shRNA (Scr) constructs were transfected bilaterally into

the L2/3 PCs of the mPFC using IUE (Fig. 8A). Both the Scr-transfected control and shSyt7-

transfected rats underwent tFC training (Fig. 8B). The following day, the rats were subjected

to a tone test in which trace fear memory formation was assessed from the freezing response

to a tone alone in a distinct context. Compared to the control group, the shSyt7-transfected

group exhibited significantly lower levels of freezing behavior in response to auditory cues

during the first four trials, suggesting an impairment in trace memory formation (Fig. 8C).

The tone test was repeated the following day to test for extinction memory formation. Syt7

KD animals exhibited much less freezing to the CS (+1d) than did WT animals. Considering the

Rescorla-Wagner model (Yau and McNally, 2023), our results suggest that the strength of CS-

US associative memory was weaker in KD animals than in WT animals; thus, the acquisition

of trace fear memory was impaired by Syt7 KD in the L2/3 PCs of the mPFC.

Next, we assessed the locomotor activity and anxiety levels of rats using the open field test

(OFT) and elevated plus maze test (EPM). Scr controls and Syt7 KD rats showed similar

exploratory patterns, displaying comparable total distance moved and time spent in the

periphery of the OFT or the open/closed arms of the EPM (Fig. S12). Previous studies have

suggested that the acquisition of trace fear memory requires elevated trace activity in

prelimbic PCs (Gilmartin et al., 2013). Given that c-Fos, an immediate early gene, is

upregulated following patterned activities in neurons, its expression is considered a reliable

indicator of recent neuronal activation (Fields et al., 1997; Guzowski et al., 2005). To

investigate the effect of Syt7 on neuronal activity *in vivo* during tFC, animals were sacrificed

at the time of peak c-Fos protein expression, approximately 90 min after completion of the

behavioral task (Arime and Akiyama, 2017).

To elucidate the activation patterns of specific neuronal populations, we performed

immunohistochemical co-labeling of c-Fos and GAD67 (a GABAergic interneuron marker)

upon tFC acquisition. The density of c-Fos(+) cells and the percentage of GAD67(+) cells among

c-Fos(+) cells in all prelimbic layers were significantly lower in Syt7 KD rats than in control rats

(Fig. 8F-G). Additionally, analysis of the proportion of active cells among transfected L2/3 PCs

(c-Fos+/GFP+) revealed that Syt7 KD L2/3 PCs were significantly more active than Scr-

transfected PCs (Fig. 8H). This unexpected hyperactivity of Syt7 KD PCs may be attributed to

the high reciprocal connectivity between PCs and FSINs in the neocortex (Holmgren et al.,

2003; Otsuka and Kawaguchi, 2009), as PCs with short-term depression would transmit less

input to FSINs, which in turn would exert less feedback inhibition on reciprocally connected

PCs. Collectively, these findings show that the slow activation of vesicle refilling supports

various forms of facilitation at excitatory synapses in PFC L2/3, which is essential for neuronal

activity during temporal associative learning.

Discussion

The present study characterized STP at local excitatory synapses in the L2/3 network of the

rat mPFC at physiological extracellular $[Ca^{2+}]$. These synapses displayed initial strong

depression and then delayed facilitation at 40 Hz, while monotonous slowly developing

facilitation was observed at lower frequencies. Our study suggests that such delayed

facilitation after a brief depression results from a high p_v (Figs. 2-3 and Fig. S8), along with

Ca^{2+}-dependent delayed acceleration of vesicle refilling and overfilling (Figs. 2–3). The two-

fold increase in EPSC during 20 pulse train stimulations under the condition of such a high p_v

suggests a Ca^{2+}-dependent increase in the vesicle replenishment rate during a train; this then

leads to overfilling of release sites beyond their incomplete basal occupancy (Fig. 3A,B). The

high p_v and strong PPD did not result from artificial bouton stimulation, AMPAR

desensitization, or any potential distortion caused by optogenetic methods (Figs. S4 and S5).

Additionally, similar to short-term facilitation, PTA could be explained by an increase in the

TS vesicle occupancy (Fig. 3C-E), consistent with previous studies showing post-tetanic

increases in the p_{occ} and/or vesicle pool size at various types of synapses (Lee et al., 2010;

Vandael et al., 2020; Tran et al., 2023; Silva et al., 2024). These findings suggest that release

sites are partially occupied at rest and that a progressive overfilling of release sites with TS

vesicles underlies facilitation and augmentation at intracortical L2/3 excitatory synapses.

Progressive overfilling of docking sites

Conventional models for facilitation assume an increase in p_v after an AP of RRP vesicles,

which had been in a low_p_v or reluctant state at rest (Dittman et al., 2000; Turecek et al.,

2016). This facilitation model is unlikely to explain our results, because TS vesicles are

predominantly released not only at rest but also in the facilitated state during or after train

stimulation, as evidenced by the strong PPD (ISI, 25 ms) during a 5-Hz train (Fig. 2D-E) and at

different intervals after a 40-Hz/30 pulse trains (Fig. 3D). Moreover, the PPD (ISI, 25 ms)

remained pronounced during the early phase of facilitation (Fig. 7), arguing against the

possibility of slowly increasing the p_v of reluctant vesicles. These results suggested that

facilitation was mediated by an activity-dependent progressive increase in occupancy of TS

vesicles (TS occupancy). Facilitation through an increase in TS occupancy has been termed

frequency facilitation, differentiating it from paired pulse facilitation that relies on a residual

calcium-dependent transient increase in the p_v of releasable vesicles (Neher, 2024).

Assuming an increase in TS occupancy, to attain a two-fold increase in EPSC (Fig. 1), the

baseline occupancy of the docking sites by TS vesicles should be low, as estimated from the

V-M plot (c.a. 30% in Fig. 3). However, it is unclear whether the remaining docking sites that

are not occupied by TS vesicles are truly vacant or are occupied by reluctant LS vesicles that

are not released by the AP or AP trains. TS vesicles can be supplied by the transformation of

LS to TS vesicles, both of which reside at the same docking site (LS/TS model; reviewed in

Neher, 2024). Alternatively, TS vesicles can be supplied from a distinct replacement pool

(RS/DS model; reviewed in Silva et al., 2021). Further studies, including ultrastructural

analyses of active zones, are required to address this issue.

Given the crucial role of Syt7 in facilitation (Fig. 5), the facilitation mechanisms are closely

related to how Syt7 mediates synaptic facilitation. Based on the slow kinetics of Syt7, previous

computational modeling studies have proposed that Syt7 may mediate short-term facilitation

through an activity-dependent increase in p_v (Turecek et al., 2016; Jackman and Regehr, 2017;

Turecek et al., 2017; Norman et al., 2023). The present study, however, found that both

facilitation and augmentation were mediated by overfilling of the release sites (Figs. 2–3).

Moreover, Syt7-KD synapses displayed slower recovery of TS vesicles (Fig. 7). These results

suggest that Syt7 is essential for activity-dependent overfilling of docking sites with TS vesicles.

Whereas the baseline EPSC was not altered (Fig. S3), and complementary changes in the

number of docking sites and their baseline occupancy were observed in Syt7 KD synapses (Fig.

6). These results raise a possibility that Syt7 may play a role in providing additional vacant

docking sites that can be overfilled during facilitation. It remains to be elucidated whether the

decrease in the number of docking sites at Syt7 KD synapses is related to the subcellular

localization of Syt7 to the plasma membrane of the active zone (Sugita et al., 2001; Vevea et

al., 2021).

STP model in light of known Ca^{2+} binding kinetics of Syt7

Syt7 acts as an upstream Ca^{2+} sensor that facilitates activity-dependent replenishment (Liu et

al., 2014; Bacaj et al., 2015; Chen et al., 2017; Tawfik et al., 2021). The present study shows

that EPSCs are mediated by TS vesicles both at rest and during activity-dependent facilitation.

Releasable vesicles are homogeneous, and Syt7-dependent overfilling is responsible for the

activity-dependent enhancement of EPSCs. Thus, we tested whether slowly developing

facilitation could be replicated within the framework of a simple refilling model based on the

known Ca^{2+} binding kinetics of Syt7 (Brandt et al., 2012). To this end, we made following

assumptions: (1) a Ca^{2+}-dependent increase in the forward refilling rate (k_1) requires a full

Ca^{2+}-bound form of Syt7, to which multiple Ca^{2+} ions can bind (two or three Ca^{2+} ions on each

C2A and/or C2B domain) (Brandt et al., 2012; Chon et al., 2015; Voleti et al., 2017); (2)

sequential Ca^{2+} binding steps are synergistic through a cooperativity factor, *b*; (3) Syt7 senses

AP-induced local $[Ca^{2+}]$ similar to that estimated at the calyx of Held (a Gaussian function with

a full width at half maximum value of 0.2 ms and a peak of 40 μM) (Wang et al., 2008); and

(4) residual $[Ca^{2+}]$ follows a mono-exponential function with an amplitude of 1 μM and decay

time constant of 50 ms (Jackson and Redman, 2003).

Under these assumptions and the framework of the simple refilling model, slow and delayed

facilitation could be replicated by modeling k_1 as $k_1 = k_{1,b} + K_{1,max} \times$ [Syt7: n Ca^{2+}], where [Syt7:

n Ca^{2+}] represents the fraction of fully Ca^{2+}-bound Syt7 (Fig. 9A-C). Syt7 was slowly activated

over the course of the stimulus train, suggesting that the cooperative binding of multiple Ca^{2+}

ions progressively increased the putative active form of Syt7 because of its slow membrane

binding/unbinding kinetics. Nevertheless, the late phase of facilitation at high frequencies

was predicted to be higher in the simulation than in the experiment. As expected, based on

the notion that sustained release during HFS is limited by site clearance in the active zone (Fig.

4C), the model/data discrepancy became more pronounced at higher frequencies.

Incorporating endocytic terms into the model may mitigate late-phase discrepancies.

Moreover, removal of the catalytic function of Syt7 in the model effectively accounted for the

complete loss of facilitation observed in Syt7 KD (Fig. 9D).

Possible mechanisms underlying the depression-facilitation sequence

Cooperative binding of multiple Ca^{2+} ions to Syt7 predicts delayed activation and a progressive

increase in vesicle refilling during train stimulations, resulting in delayed facilitation.

Combining this with a high p_v could replicate not only the slow facilitation at 5 Hz, but also

the unique depression-facilitation sequence observed at 40 Hz (Fig. 9). As simulated by Pulido

and Marty (2018), delayed facilitation can be reproduced by a high p_v in combination with a

low baseline occupancy of replacement sites and subsequent Ca^{2+}-dependent refilling.

However, whether this model is also able to replicate slow facilitation at low frequencies

remains to be tested. An alternative, but not exclusive, possibility for delayed facilitation is

that the presynaptic global Ca^{2+} itself may undergo a slow buildup during a train due to the

saturation of endogenous Ca^{2+}-binding proteins.

Facilitation requires not only Syt7, but also activation of PLC-DAG signaling

To gain insight into the specific molecular mechanisms underlying the vesicle-loading

processes, we examined the effects of pharmacological inhibitors targeting PLC and DAG (Fig.

4). Both a PLC inhibitor and a DAG analog reduced facilitation and augmentation. This

suggests that the progressive overfilling of docking sites with TS vesicles is supported by the

PLC-DAG signaling pathway and probably downstream Munc13 (Rhee et al., 2002; Lou et al.,

2008). Phorbol esters have been recently shown to increase the TS fraction of docked vesicles

by enhancing vesicle priming without affecting Ca^{2+} transients or p_v (Taschenberger et al.,

2016; Aldahabi et al., 2022; Papantoniou et al., 2023; Aldahabi et al., 2024). In line with these

findings, signaling involving PLC and DAG may support Ca^{2+}-dependent overfilling of TS

vesicles by enhancing molecular priming processes.

Roles of facilitation and augmentation in working memory and related tasks

Previous theoretical studies have proposed that working memory in a recurrent network can

be maintained in the form of augmentation at synapses (Mongillo et al., 2008) and/or

persistent activity of neurons in a memory ensemble. This is supported by the facilitation of

recurrent excitatory synapses or disparity in STP between excitatory and inhibitory synapses.

However, whether these short-term enhancements in synaptic efficacy contribute to working

memory-related behaviors has not been examined. tFC is a temporal associative learning task

requiring temporary retention of CS information during a trace period. The mPFC and

hippocampus are crucial for tFC, and tFC is interfered with by a load of working memory and

attention distraction, implying that the tFC shares the same neural substrate with other

working memory tasks. In this study, tFC in WT and Syt7 KD animals revealed that Syt7

deficiency in the L2/3 pyramidal neurons of the mPFC impaired the acquisition of trace fear

memory and reduced c-Fos expression associated with tFC training (Fig. 8). Given that

synaptic enhancements in prelimbic L2/3 recurrent excitatory synapses can be attributed to

an increase in p_{occ}, these behavioral results suggest that presynaptic high Ca^{2+} transients are

temporally memorized in the presynaptic terminal as an increase in TS occupancy.

Conclusions

Progressive overfilling of synaptic vesicles with a high fusion probability supports the

facilitation and augmentation of recurrent excitatory synapses within the L2/3 network of the

mPFC and is essential for temporal associative learning.

Materials and Methods

Animals/Subjects

All experiments were carried out on Sprague-Dawley rats of either sex at p28-43. Rats were maintained in temperature-controlled rooms on a constant 12 h light/12 h dark cycle and were given access to water and food ad libitum. All animal procedures were approved and performed in accordance with the Institutional Animal Care and Use Committee at Seoul National University (SNU200522-1).

In utero electroporation (IUE)

Pregnant Sprague Dawley rats on embryonic day (E) 17.5 were deeply anesthetized with 5% (v/v) isoflurane for the duration of surgery. The uterine horns were exposed by laparotomy and wet with warm sterile phosphate-buffered saline (PBS). The plasmids (0.5-2 µg/µl) together with the 0.01% fast green dye were injected into either the left or the right lateral ventricle or bilateral ventricles of embryos through a thin glass capillary (Narishige PC-10). The embryo's head was carefully held between the forceps-shaped electrodes of 10 mm diameter (CUY65010; NEPAGENE, Japan). Electroporation pulses (50 V for 50 ms) were delivered five times at intervals of 150 ms using a square-wave generator (ECM830; BEX, United States). The uterine horns were placed back into the abdominal cavity and abdominal muscles and skin were sutured. After the surgery, the animals were recovered under the infrared lamp. For knockdown experiments in slice electrophysiology, the shRNA constructs were co-electroporated with the pCAG-oChIEF-tdTomato, because of the consistent co-

expression of the two constructs in a significant proportion of the transfected cells in vivo

(Bony et al., 2013). Pups with robust fluorescence signals in the prefrontal cortex of both

hemispheres were used for behavior experiments.

Preparation of vector constructs

For an optic stimulation, pCAG-oChIEF-tdTomato was transfected by IUE. Short hairpin RNA

(shRNA) against Syt7 sequence (GATCTACCTGTCCTGGAAGAG) was expressed under the

control of an U6 promoter of pAAV-U6-GFP vector (Cell Biolabs, #VPK-413). This shRNA was

shown to effectively knockdown Syt7 in cell cultures and in vivo by previous reports from

more than one research group (Bacaj et al., 2013; Li et al., 2017). Scrambled control sequence

(TCGCATAGCGTATGCCGTT) was designed by shuffling the recognition region sequences and

a BLAST analysis confirmed that the control shRNA sequence had no target gene. For rescue

experiments, the cDNA for rat Syt7 (NM_021659) rendered resistant to the shRNA were

inserted in the KD vector and their expression was driven by the CAG promoter; the vector

also contained a self-cleaving P2A peptide sequence followed by EGFP, which ensures

visualization of transfected cells. All the constructs above were verified by DNA sequencing.

Plasmid DNA was purified and concentrated under endotoxin free conditions (Qiagen

EndoFree Maxi Kit).

 Slice preparation

 Acute brain slices were prepared by using a VT1200S Vibratome (Leica Microsystems) after

 isoflurane anesthesia and decapitation. Coronal sections of the mPFC (300 μm in thickness)

 were cut in ice-cold artificial cerebrospinal fluid (aCSF) composed of (in mM) 125 NaCl, 3.2

 KCl, 25 $NaHCO_3$, 1.25 NaH_2PO_4, 20 D-glucose, 2 NaPyr, 1 $MgCl_2$, and 1.3 $CaCl_2$ (Yoon et al.,

 2020), bubbled with 95% O_2 and CO_2 (pH = 7.3; 310 mOsm). Slices were allowed to recover

 for a minimum of 30 minutes at 36 °C, and subsequently maintained at room temperature

 until recording. Composition of aCSF was identical throughout the preparation, incubation,

 and recording periods unless otherwise stated. For the recordings, the mPFC slices were

 continuously perfused with oxygenated aCSF at a flow rate of 1.5 ml/min, and the

 temperature was maintained at 31–32 °C using an in-line temperature controller (Warner

 Instruments).

 Electrophysiology

 Whole-cell patch-clamp recordings were performed from pyramidal cells and fast-spiking

 interneurons in layer 2/3 of the prelimbic mPFC. Neuronal subtypes were initially selected

 based on morphology under visual guidance by using an upright microscope equipped with

 infrared differential interference contrast (IR-DIC) optics (BX51WI, Olympus), and were

 further identified by their electrophysiological properties (see Supplementary Fig. 1; van

 Aerde and Feldmeyer, 2015; Tremblay et al., 2016). To analyze these intrinsic properties in

 current-clamp experiments, we measured the following parameters: (1) resting membrane

potential (RMP), (2) input resistance (R_{in}), (3) sag ratio, and (4) the F-I curve, which plots firing

frequencies (F) against the amplitude of injected currents (I). Both cell types exhibited a low

sag ratio below 0.1, consistent with findings from other studies (van Aerde and Feldmeyer,

2015; Yoon et al., 2020). PCs and FSINs were easily distinguished by their unique firing

frequency, Rin, spike adaptation, RMP and morphological characteristics including the shape

of soma and the thickness of apical dendrites. For experiments involving oChIEF-expressing

neurons in WT or Syt7 KD rats, whole-cell recordings were performed from the fluorescently

labeled pyramidal neurons in layer 2/3 of the prelimbic cortex using suitable filter sets. Patch

pipettes (tip resistance of 2–3 MΩ) were pulled from borosilicate glass and filled with (in mM)

130 K-gluconate, 8 KCl, 10 HEPES, 0.2 EGTA, 20 Na_2Pcr, 0.3 Na_2GTP, 4 MgATP. Intracellular

solutions were adjusted to pH 7.25 and 300–310 mOsm. Neurons were voltage clamped at

−78 mV and series resistance (Rs) were continuously monitored during the EPSCs recordings.

Experiments were discarded before analysis if the Rs changed by 20% or larger deviation of

baseline value. Measurements were not corrected for Rs compensation, bridge balance, or

liquid junction potential. Electrophysiological data were low-pass-filtered at 1 kHz (Bessel)

and acquired at 20 kHz using a Multiclamp 700B amplifier paired with Digidata 1440A digitizer

and Clampex 10.2 software (Molecular Devices).

Synaptic stimulation

IUE resulted in sparse expression of channelrhodopsin (10-20%) (Fig. 1A). Exploiting the

sparse expression, we employed a minimal optical stimulation method to activate a single or

very few excitatory synapses in layer 2/3. The optic minimal stimulation was performed by

our published protocol with minor modifications (Yoon et al., 2020). This approach aimed to

minimize the risk of synaptic contamination, which could arise from engaging extra terminals

that were initially inactive (depolarized below threshold) but became active by temporal

summation during high-frequency trains. It also helped prevent the buildup of depolarization

at the synaptic terminals, which might influence the probability of release, as noted by

(Jackman et al., 2014). To achieve this, we employed a collimated digital micromirror device

(DMD)-coupled LED (Polygon400; Mightex Systems) to confine 470 nm blue light to a small

area with a radius of approximately 3–8 μm (typically 3–4 μm), as measured at the focal plane

of a 60× water immersion objective [numerical aperture (NA) = 1.0; LUMPlanFL, Olympus] (Fig.

S3A). To find the minimal stimulation area, the radius of illumination area was increased from

2 μm by 50% at each step (Fig. S3A), and selected the smallest illumination area that elicited

EPSCs. Setting the duration of illumination between 3 and 5 ms (typically 5 ms),

photostimulation onto oChIEF-expressing cells reliably induced action potentials (APs) during

600-pulse trains (see Fig. S2).

STP at various stimulus frequencies (5, 10, 20, and 40 Hz) was examined using 20-pulse trains.

Trials of train stimulation were repeated with a prolonged interval of 30 s to avoid change in

baseline properties. We observed no systematic changes in the amplitudes of the first EPSC

of each train, nor did we find significant differences in the paired-pulse ratio (PPR) across

stimulation trials in the same cell. For each stimulus frequency, approximately ten to twenty

consecutive traces were collected and then averaged across trials to yield a mean EPSC trace.

For recovery experiments, two consecutive trains of 3 or 30 stimuli at 40 Hz, with varying time

intervals (0.1-10 s), were applied every 10 s or 60 s, respectively, at both excitatory synapses.

For 600-stimulus trains at 10Hz, the trial was delivered only once for each cell to avoid

possible after-effects caused by prolonged synaptic stimulation (Hirsch and Crepel, 1990).

Prior to the stimulus trains, the EPSCs were normalized to the mean amplitude of the baseline.

For the baseline measurements, paired pulses with various ISIs (20-200 ms) were delivered

every 3 s (1/3 Hz) at least 20 times.

For all recordings, picrotoxin (PTX, 10 μM) was included in the recording solution to isolate

glutamatergic responses. mPFC PC terminals were entirely glutamatergic as the application

of 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX, 20 μM), an AMPA receptor antagonist,

completely abolished the monosynaptic events evoked by photostimulation. A low

concentration of PTX was employed to minimize the relieving effects on spike suppression

caused by blockade of GABAa receptors. We confirmed that no synaptic response was elicited

by electrical stimulation in the presence of both 10 μM PTX and 20 μM CNQX.

In Fig. S4A, 1 μM TTX (1 mM stock solution) and 0.1 mM 4-AP (500 mM stock solution) were

added to the bath solution to block action potentials and restore presynaptic glutamate

release, respectively (Fig. S4A). In Fig. S5A, 50 μM of cyclothiazide, a positive allosteric

modulator of the AMPA receptors, was included to minimize rapid desensitization of the

postsynaptic receptors. In Fig. S5B, 0.5 mM kynurenate was added to bath solution to prevent

possible saturation of AMPA receptors, especially during maximal synaptic facilitation. For

pharmacological manipulations of synaptic vesicle dynamics (Fig. 3), drugs related to vesicle

supply including U73122, OAG, LatB, EGTA-AM, and dynasore at indicated concentrations

were added to the recording solution during baseline measurements or PTA/recovery

experiments (IBI = 0.5 s). Chemicals were from Sigma-Aldrich, Merck, or Tocris.

Measurements of quantal size

Experiments for measuring asynchronous release were performed as previously described

(Yoon et al., 2020). In brief, we replaced extracellular Ca^{2+} entirely with 2 mM Sr^{2+} to induce

asynchronous release. A single stimulus was delivered at 3-second intervals over 100

repetitions, and asynchronous release events were detected during a 100-ms window starting

25 ms after stimulus onset for each trial. The same procedure was also executed under sham

conditions (0% light intensity) to serve as a control, thus mitigating contamination from

nonspecific spontaneous events and potential errors associated with in the detection

algorithm. q was determined as the mean value from the first Gaussian function after fitting

the EPSC distribution with either a single or double Gaussian function (Fig. S8).

Calculation of probability for double failure

The failure rate at the 1st pulse when initial RRP size (n_1) = i can be calculated as: $P(F_1 \mid n_1=i)$

= $(1 - p_v)^i$. The probability that initial RRP size is i was calculated as: $P(n_1=i) = C(N, i) * p_{occ}^i *$

$(1-p_{occ})^{(N-i)}$, where p_{occ} = mean release site occupancy, and N = total number of release sites.

Under the conditions of no vesicle release at the 1st pulse (F_1) and $n_1 = i$, the probability for

the second failure is

$$P(F_2 \mid n_1=i, F_1) = \Sigma_{n2=0..N} P(n_2 \mid n_1=i, F_1) (1 - p_v)^{n2} .$$

, where n_2 = number of vesicles just before the 2nd pulse. When forward and backward rate

constants are k_1 and b_1, respectively, $P(n_2 \mid n_1=i, F1)$ is calculated as follows:

683     If $n_2 > i$, $P(n_2 \mid n_1 = i, F1) = C(N - i, n_2 - i) [1 - \exp(-k_1 \Delta t)]^{(n_2 - i)}$.

684     If $n_2 < i$, $P(n_2 \mid n_1 = i, F1) = C(i, i - n_2) [1 - \exp(-b_1 \Delta t)]^{(i - n_2)}$.

685     If $n_2 = i$, $P(n_2 \mid n_1 = i, F1) = \exp[-i\, k1\, \Delta t\,] \exp[-i\, b_1\, \Delta t\,]$.

686  , where $\Delta t = $ ISI. By the Bayes rule,

687     $P(F_1, F_2, n_1 = i) = P(F_2 \mid n_1 = i, F_1)\, P(F_1 \mid n_1 = i)\, P(n_1 = i)$.

688  Thus, the probability for two consecutive failures can be calculated as:

689     $P(F_1, F_2) = \Sigma_{i=0..N}\, P(F_2 \mid n_1 = i, F_1)\, P(F_1 \mid n_1 = i)\, P(n_1 = i)$.

690  $P(F_1)$ and N are observed values. The k_1 and p_v were set free variables. Other parameters were

691  set according to following relationships: $P(F_1) = (1 - p_r)^N$, $p_r = p_v\, p_{occ}$, and $p_{occ} = k_1 / (k_1 + b_1)$.

693  *Behavioral tests*

694  **Trace fear conditioning**: For trace fear conditioning (tFC) (Gilmartin et al., 2013), individual

695  rats were placed in the training chamber (30.5 x 25.4 x 30.5 cm; Coulbourn Instruments)

696  consisting of metal grid floor connected to an electrical stimulator (H10-11R-TC-SF, Coulbourn

697  Instruments) for delivery of a footshock. After a 5 min habituation period, rats received five

698  pairings of a 20 s tone conditioned stimulus (CS, 2.5 kHz, 75 dB), followed by empty 20s trace

699  period and a unconditioned stimulus (US; 1 s, 0.7 mA footshock) with a pseudorandom inter-

700  trial interval (ITI) of 240 ± 20 s. During this training session, rats learned to associate the

701  auditory CS with the shock US. The next day, the rats were tested for memory of CS-US trace

702  association (called tone test). The acquisition of fear learning, often indicated by the extent

703  of freezing to the CS, was assessed in a novel chamber (a white hexagonal enclosure) in a

704  dimly lit room. During this tone (or retrieval) test, rats were first allowed to explore the new

context for 5 min (habituation), followed by twelve 20 s CS presentations with a variable ITI

of 120 ± 20 s. This entire session was repeated on the next day for assessment of extinction

learning. Identities of the chambers were also determined with the presence of distinct odors;

the training and testing chambers were cleaned with 70% ethanol and 1% acetic acid

respectively before and after each session. For all experimental sessions, the activity of

animals in the chambers was recorded at 30 frames per second using the EthoVision XT

(Noldus Information Technology, Wageningen, Netherlands), and stored as a video file. The

freezing behavior, defined as behavioral immobility other than respiratory movement, was

assessed using an open-source video analysis pipeline ezTrack (Pennington et al., 2019). The

percentage of freezing was calculated as a total duration of freezing divided by the total

duration of observation.

Elevated Plus Maze: Elevated plus maze (EPM) test was performed using a plus-shaped

apparatus elevated 60 cm above the ground. The maze consisted of two open arms (50 x 10

x 0.5 cm (H)), two closed arms (50 x 10 x 40 cm (H)) and a center area (10 x 10 cm). Rats were

individually placed on the center area, facing an open arm, and the path of the animal was

recorded with a video camera for 5 min. We analyzed the total distance traveled for 5 min

and time spent in each arm. Animals were tracked offline using the open-source tracking

system ezTrack (Pennington et al., 2019).

Open field test: Open field exploration test (OFT) was performed in an open field apparatus

in a dimly lit room. The open field consisted of a white plastic board (1.20 m in diameter)

surrounded by white plastic walls (50 cm in height). Individual rats were placed in a center

zone (0.6 m in diameter), and allowed to freely explore the apparatus for 10 min. Time spent

in the center zone and total distance traveled were analyzed for initial 5 min. Animals were

tracked offline using the open-source tracking system ezTrack (Pennington et al., 2019).

Histology and immunohistochemistry

At the end of the behavioral experiments, rats were anesthetized using isoflurane inhalation

and transcardially perfused with PBS followed by 4% paraformaldehyde (PFA, T&I, Korea).

Brains were extracted and post-fixed in 4% PFA solution for 24 h. Perfused brains were sliced

at 100 μm for estimation of GFP expression or 50 μm for immunohistochemistry. For analysis

of c-Fos expression, rats were returned to their home cages immediately after the training

session and perfused approximately 90 minutes later (Arime and Akiyama, 2017). Brains were

then cryopreserved, frozen, and sectioned coronally. Immunofluorescence staining was

conducted on free-floating slices using a rabbit monoclonal anti c-fos (1:1000, Cell Signaling

Technology, cat# 2250s) and mouse monoclonal anti-GAD67 (1:500, Millipore, cat# mab5406).

Fluorescent secondary antibodies included Cy5-conjugated goat anti-rabbit IgG (1:1000,

Abcam, cat# ab97077) and Cy3-conjugated goat anti-mouse IgG (1:1000, Abcam, cat#

ab97035). Brain slices were permeabilized and blocked with PBS containing 0.5% Triton X-100

and 5% normal goat serum (NGS) for 1h at room temperature. Slices were incubated

overnight at 4 °C with primary antibodies, followed by 4 h incubation with secondary

antibodies at room temperature. Slices were then mounted with Antifading Mounting

medium (Abcam, cat# ab104139). Confocal images were acquired using a Leica TCS SP8

microscope set to the same laser intensity and acquisition parameters to compare the

immunosignals in sections from Scr and Syt7 KD groups. For cell counting and colocalization,

fluorescent protein-positive cells were automatically quantified with spot-detection

algorithms in Imaris 9.5 (Oxford instruments).

Data analysis

Data were analyzed and presented using ClampFit (Molecular Devices), Igor Pro

(Wavemetrics), MATLAB (Mathworks), and Prism (GraphPad). Data points and error bars

represent the mean ± standard error of the mean. All statistical analyses were conducted

using two-tailed comparisons. Sample sizes and statistical tests for each group are detailed in

the figure legends.

Numerical integration of the simple refilling model

This model posits reversible docking of vesicles at finite release sites (N_{max}) with forward (k_1)

and reverse rate constants (b_1) (Hosoi et al., 2007; Neher and Sakaba, 2008).

$$\underset{b_1}{\overset{k_1}{\rightleftarrows}}\ n\ \overset{p_v}{\rightarrow}$$

According to this scheme, $dn/dt = k_1\ u - b_1\ n - p_v\ n\ \delta(t - t_{AP})$, where u is the number of

unoccupied sites ($= N_{max} - n$); t_{AP} is the timing of AP firing; and δ is the Dirac delta function.

The p_v was assumed to be unity (Fig. S8). The sum of basal k_1 ($= k_{1,b}$) and b_1 was be estimated

as 23/s from the dependence PPR on ISIs (Fig. S8Ab). From the baseline occupancy of 0.3 and

$p_{occ} = k_1 / (k_1 + b_1)$, we could estimate baseline k_1 ($k_{1,b}$) and b_1 as 6.9/s, and 16.1/s, respectively.

Ca^{2+}- and Syt7-dependent increase of k_1 was modeled as $k_1 = k_{1,b} + K_{1,max} [Syt7: n\ Ca^{2+}]$, where

$[Syt7: n\ Ca^{2+}]$ represents the fraction of full Ca^{2+}-bound Syt7. For deterministic simulations,

we time-integrated a set of differential equations for each model using Euler methods with a

time step of 1 ms. All calculations were performed on the platform of Matlab (R2022b,

Mathworks, USA).

Acknowledgements

775   We thank Dr. E. Neher and Dr. A. Marty for critical reading of this manuscript and

776   invaluable comments.

Author contributions

779   All experiments were carried out at cell physiology lab under Y.K and S.H.L's supervision.

780   Conception or design of the work: J.S. and S.H.L. Acquisition, analysis or interpretation of data

781   for the work: J.S, Y.K. and L.S.H. Drafting the work or revising it critically for important

782   intellectual content: J.S., S.Y.L, Y.K. and L.S.H.

Funding

785   This study was supported by grants from the National Research Foundation of Korea (RS-

786   2024-333669 to S-H. Lee; 2021R1I1A1A01059646 to Y. Kim), and Seoul National University

787   Hospital (2024).

References

Aldahabi M, Neher E, Nusser Z (2024) Different states of synaptic vesicle priming explain target cell type-dependent differences in neurotransmitter release. Proc Natl Acad Sci U S A 121:e2322550121.

Aldahabi M, Balint F, Holderith N, Lorincz A, Reva M, Nusser Z (2022) Different priming states of synaptic vesicles underlie distinct release probabilities at hippocampal excitatory synapses. Neuron 110:4144-4161 e4147.

Arime Y, Akiyama K (2017) Abnormal neural activation patterns underlying working memory impairment in chronic phencyclidine-treated mice. PLoS One 12:e0189287.

Bacaj T, Wu D, Burre J, Malenka RC, Liu X, Sudhof TC (2015) Synaptotagmin-1 and -7 Are Redundantly Essential for Maintaining the Capacity of the Readily-Releasable Pool of Synaptic Vesicles. PLoS Biol 13:e1002267.

Bacaj T, Wu D, Yang X, Morishita W, Zhou P, Xu W, Malenka RC, Sudhof TC (2013) Synaptotagmin-1 and synaptotagmin-7 trigger synchronous and asynchronous phases of neurotransmitter release. Neuron 80:947-959.

Baeg EH, Kim YB, Huh K, Mook-Jung I, Kim HT, Jung MW (2003) Dynamics of population code for working memory in the prefrontal cortex. Neuron 40:177-188.

Bony G, Szczurkowska J, Tamagno I, Shelly M, Contestabile A, Cancedda L (2013) Non-hyperpolarizing GABAB receptor activation regulates neuronal migration and neurite growth and specification by cAMP/LKB1. Nat Commun 4:1800.

Brandt DS, Coffman MD, Falke JJ, Knight JD (2012) Hydrophobic contributions to the membrane docking of synaptotagmin 7 C2A domain: mechanistic contrast between isoforms 1 and 7. Biochemistry 51:7654-7664.

Chen C, Satterfield R, Young SM, Jr., Jonas P (2017) Triple Function of Synaptotagmin 7 Ensures Efficiency of High-Frequency Transmission at Central GABAergic Synapses. Cell Rep 21:2082-2089.

Chon NL, Osterberg JR, Henderson J, Khan HM, Reuter N, Knight JD, Lin H (2015) Membrane Docking of the Synaptotagmin 7 C2A Domain: Computation Reveals Interplay between Electrostatic and Hydrophobic Contributions. Biochemistry 54:5696-5711.

Dittman JS, Kreitzer AC, Regehr WG (2000) Interplay between facilitation, depression, and residual calcium at three presynaptic terminals. J Neurosci 20:1374-1385.

Fields RD, Eshete F, Stevens B, Itoh K (1997) Action potential-dependent regulation of gene expression: temporal specificity in ca2+, cAMP-responsive element binding proteins, and mitogen-activated protein kinase signaling. J Neurosci 17:7252-7266.

Fujisawa S, Amarasingham A, Harrison MT, Buzsaki G (2008) Behavior-dependent short-term assembly dynamics in the medial prefrontal cortex. Nat Neurosci 11:823-833.

Gilmartin MR, Miyawaki H, Helmstetter FJ, Diba K (2013) Prefrontal activity links nonoverlapping events in memory. J Neurosci 33:10910-10914.

Guzowski JF, Timlin JA, Roysam B, McNaughton BL, Worley PF, Barnes CA (2005) Mapping behaviorally relevant neural circuits with immediate-early gene expression. Curr Opin Neurobiol 15:599-606.

Hallermann S, Silver RA (2013) Sustaining rapid vesicular release at active zones: potential
roles for vesicle tethering. Trends Neurosci 36:185-194.

Hansel D, Mato G (2013) Short-term plasticity explains irregular persistent activity in working
memory tasks. J Neurosci 33:133-149.

Hass CA, Glickfeld LL (2016) High-fidelity optical excitation of cortico-cortical projections at
physiological frequencies. J Neurophysiol 116:2056-2066.

Haucke V, Neher E, Sigrist SJ (2011) Protein scaffolds in the coupling of synaptic exocytosis
and endocytosis. Nat Rev Neurosci 12:127-138.

Hempel CM, Hartman KH, Wang XJ, Turrigiano GG, Nelson SB (2000) Multiple forms of short-
term plasticity at excitatory synapses in rat medial prefrontal cortex. J Neurophysiol
83:3031-3041.

Hirsch JC, Crepel F (1990) Use-dependent changes in synaptic efficacy in rat prefrontal
neurons in vitro. J Physiol 427:31-49.

Holler S, Kostinger G, Martin KAC, Schuhknecht GFP, Stratford KJ (2021) Structure and
function of a neocortical synapse. Nature 591:111-116.

Holmgren C, Harkany T, Svennenfors B, Zilberter Y (2003) Pyramidal cell communication
within local networks in layer 2/3 of rat neocortex. J Physiol 551:139-153.

Hosoi N, Sakaba T, Neher E (2007) Quantitative analysis of calcium-dependent vesicle
recruitment and its functional role at the calyx of Held synapse. J Neurosci 27:14286-
14298.

Jackman SL, Regehr WG (2017) The Mechanisms and Functions of Synaptic Facilitation.
Neuron 94:447-464.

Jackman SL, Beneduce BM, Drew IR, Regehr WG (2014) Achieving high-frequency optical
control of synaptic transmission. J Neurosci 34:7704-7714.

Jackman SL, Turecek J, Belinsky JE, Regehr WG (2016) The calcium sensor synaptotagmin 7 is
required for synaptic facilitation. Nature 529:88-91.

Jackson MB, Redman SJ (2003) Calcium dynamics, buffering, and buffer saturation in the
boutons of dentate granule-cell axons in the hilus. J Neurosci 23:1612-1621.

Kobbersmed JRL, Grasskamp AT, Jusyte M, Böhme MA, Ditlevsen S, Sørensen JB, Walter AM
(2020) Rapid regulation of vesicle priming explains synaptic facilitation despite
heterogeneous vesicle:Ca2+ channel distances. eLife 9:e51032.

Kusick GF, Chin M, Raychaudhuri S, Lippmann K, Adula KP, Hujber EJ, Vu T, Davis MW,
Jorgensen EM, Watanabe S (2020) Synaptic vesicles transiently dock to refill release
sites. Nat Neurosci 23:1329-1338.

Lee HR, Choi SH, Lee SH (2024) Differential involvement of mitochondria in post-tetanic
potentiation at intracortical excitatory synapses of the medial prefrontal cortex. Cereb
Cortex 34.

Lee JS, Ho WK, Lee SH (2010) Post-tetanic increase in the fast-releasing synaptic vesicle pool
at the expense of the slowly releasing pool. J Gen Physiol 136:259-272.

Lee JS, Ho WK, Lee SH (2012) Actin-dependent rapid recruitment of reluctant synaptic vesicles
into a fast-releasing vesicle pool. Proc Natl Acad Sci U S A 109:E765-774.

Li YC, Chanaday NL, Xu W, Kavalali ET (2017) Synaptotagmin-1- and Synaptotagmin-7-
Dependent Fusion Mechanisms Target Synaptic Vesicles to Kinetically Distinct
Endocytic Pathways. Neuron 93:616-631 e613.

Lin JY, Lin MZ, Steinbach P, Tsien RY (2009) Characterization of engineered channelrhodopsin
variants with improved properties and kinetics. Biophys J 96:1803-1814.
Lin KH, Taschenberger H, Neher E (2022) A sequential two-step priming scheme reproduces
diversity in synaptic strength and short-term plasticity. Proc Natl Acad Sci U S A
119:e2207987119.
Liu H, Bai H, Hui E, Yang L, Evans CS, Wang Z, Kwon SE, Chapman ER (2014) Synaptotagmin 7
functions as a Ca2+-sensor for synaptic vesicle replenishment. Elife 3:e01524.
Lou X, Korogod N, Brose N, Schneggenburger R (2008) Phorbol esters modulate spontaneous
and Ca2+-evoked transmitter release via acting on both Munc13 and protein kinase C.
J Neurosci 28:8257-8267.
Macia E, Ehrlich M, Massol R, Boucrot E, Brunner C, Kirchhausen T (2006) Dynasore, a cell-
permeable inhibitor of dynamin. Dev Cell 10:839-850.
Malagon G, Miki T, Tran V, Gomez LC, Marty A (2020) Incomplete vesicular docking limits
synaptic strength under high release probability conditions. Elife 9.
Markram H, Wang Y, Tsodyks M (1998) Differential signaling via the same axon of neocortical
pyramidal neurons. Proc Natl Acad Sci U S A 95:5323-5328.
Martinetti LE, Bonekamp KE, Autio DM, Kim HH, Crandall SR (2022) Short-Term Facilitation of
Long-Range Corticocortical Synapses Revealed by Selective Optical Stimulation. Cereb
Cortex 32:1932-1949.
Miki T, Malagon G, Pulido C, Llano I, Neher E, Marty A (2016) Actin- and Myosin-Dependent
Vesicle Loading of Presynaptic Docking Sites Prior to Exocytosis. Neuron 91:808-823.
Mongillo G, Barak O, Tsodyks M (2008) Synaptic theory of working memory. Science
319:1543-1546.
Mongillo G, Hansel D, van Vreeswijk C (2012) Bistability and spatiotemporal irregularity in
neuronal networks with nonlinear synaptic transmission. Phys Rev Lett 108:158101.
Neher E (2010) What is Rate-Limiting during Sustained Synaptic Activity: Vesicle Supply or the
Availability of Release Sites. Front Synaptic Neurosci 2:144.
Neher E (2024) Interpretation of presynaptic phenotypes of synaptic plasticity in terms of a
two-step priming process. J Gen Physiol 156.
Neher E, Sakaba T (2008) Multiple roles of calcium ions in the regulation of neurotransmitter
release. Neuron 59:861-872.
Neher E, Brose N (2018) Dynamically Primed Synaptic Vesicle States: Key to Understand
Synaptic Short-Term Plasticity. Neuron 100:1283-1291.
Norman CA, Krishnakumar SS, Timofeeva Y, Volynski KE (2023) The release of inhibition model
reproduces kinetics and plasticity of neurotransmitter release in central synapses.
Commun Biol 6:1091.
Otsuka T, Kawaguchi Y (2009) Cortical inhibitory cell types differentially form intralaminar and
interlaminar subnetworks with excitatory neurons. J Neurosci 29:10533-10540.
Ozdemir AT, Lagler M, Lagoun S, Malagon-Vina H, Lasztoczi B, Klausberger T (2020)
Unexpected Rule-Changes in a Working Memory Task Shape the Firing of Histologically
Identified Delay-Tuned Neurons in the Prefrontal Cortex. Cell Rep 30:1613-1626 e1614.
Pan B, Zucker RS (2009) A general model of synaptic transmission and short-term plasticity.
Neuron 62:539-554.

Papantoniou C, Laugks U, Betzin J, Capitanio C, Ferrero JJ, Sanchez-Prieto J, Schoch S, Brose
N, Baumeister W, Cooper BH, Imig C, Lucic V (2023) Munc13- and SNAP25-dependent
molecular bridges play a key role in synaptic vesicle priming. Sci Adv 9:eadf6222.

Pennington ZT, Dong Z, Feng Y, Vetere LM, Page-Harley L, Shuman T, Cai DJ (2019) ezTrack:
An open-source video analysis pipeline for the investigation of animal behavior. Sci
Rep 9:19979.

Pulido C, Marty A (2018) A two-step docking site model predicting different short-term
synaptic plasticity patterns. J Gen Physiol 150:1107-1124.

Reyes A, Sakmann B (1999) Developmental switch in the short-term modification of unitary
EPSPs evoked in layer 2/3 and layer 5 pyramidal neurons of rat neocortex. J Neurosci
19:3827-3835.

Rhee JS, Betz A, Pyott S, Reim K, Varoqueaux F, Augustin I, Hesse D, Sudhof TC, Takahashi M,
Rosenmund C, Brose N (2002) Beta phorbol ester- and diacylglycerol-induced
augmentation of transmitter release is mediated by Munc13s and not by PKCs. Cell
108:121-133.

Ritzau-Jost A, Jablonski L, Viotti J, Lipstein N, Eilers J, Hallermann S (2018) Apparent calcium
dependence of vesicle recruitment. J Physiol 596:4693-4707.

Ritzau-Jost A, Delvendahl I, Rings A, Byczkowicz N, Harada H, Shigemoto R, Hirrlinger J, Eilers
J, Hallermann S (2014) Ultrafast action potentials mediate kilohertz signaling at a
central synapse. Neuron 84:152-163.

Rozov A, Burnashev N, Sakmann B, Neher E (2001) Transmitter release modulation by
intracellular Ca2+ buffers in facilitating and depressing nerve terminals of pyramidal
cells in layer 2/3 of the rat neocortex indicates a target cell-specific difference in
presynaptic calcium dynamics. J Physiol 531:807-826.

Sakaba T, Neher E (2003) Involvement of actin polymerization in vesicle recruitment at the
calyx of Held synapse. J Neurosci 23:837-846.

Scheuss V, Schneggenburger R, Neher E (2002) Separation of presynaptic and postsynaptic
contributions to depression by covariance analysis of successive EPSCs at the calyx of
Held synapse. J Neurosci 22:728-739.

Silva M, Tran V, Marty A (2021) Calcium-dependent docking of synaptic vesicles. Trends
Neurosci 44:579-592.

Silva M, Tran V, Marty A (2024) A maximum of two readily releasable vesicles per docking site
at a cerebellar single active zone synapse. Elife 12.

Sugita S, Han W, Butz S, Liu X, Fernandez-Chacon R, Lao Y, Sudhof TC (2001) Synaptotagmin
VII as a plasma membrane Ca(2+) sensor in exocytosis. Neuron 30:459-473.

Taschenberger H, Woehler A, Neher E (2016) Superpriming of synaptic vesicles as a common
basis for intersynapse variability and modulation of synaptic strength. Proc Natl Acad
Sci U S A 113:E4548-4557.

Tawfik B, Martins JS, Houy S, Imig C, Pinheiro PS, Wojcik SM, Brose N, Cooper BH, Sorensen
JB (2021) Synaptotagmin-7 places dense-core vesicles at the cell membrane to
promote Munc13-2- and Ca(2+)-dependent priming. Elife 10.

Tran V, Silva M, Marty A (2023) Prioritized docking of synaptic vesicles provided by a rapid
recycling pathway. iScience 26:106366.

Tremblay R, Lee S, Rudy B (2016) GABAergic Interneurons in the Neocortex: From Cellular
Properties to Circuits. Neuron 91:260-292.

Turecek J, Jackman SL, Regehr WG (2016) Synaptic Specializations Support Frequency-
Independent Purkinje Cell Output from the Cerebellar Cortex. Cell Rep 17:3256-3268.

Turecek J, Jackman SL, Regehr WG (2017) Synaptotagmin 7 confers frequency invariance onto
specialized depressing synapses. Nature 551:503-506.

Valera AM, Doussau F, Poulain B, Barbour B, Isope P (2012) Adaptation of granule cell to
Purkinje cell synapses to high-frequency transmission. J Neurosci 32:3267-3280.

van Aerde KI, Feldmeyer D (2015) Morphological and physiological characterization of
pyramidal neuron subtypes in rat medial prefrontal cortex. Cereb Cortex 25:788-805.

Vandael D, Borges-Merjane C, Zhang X, Jonas P (2020) Short-Term Plasticity at Hippocampal
Mossy Fiber Synapses Is Induced by Natural Activity Patterns and Associated with
Vesicle Pool Engram Formation. Neuron 107:509-521 e507.

Vevea JD, Kusick GF, Courtney KC, Chen E, Watanabe S, Chapman ER (2021) Synaptotagmin 7
is targeted to the axonal plasma membrane through gamma-secretase processing to
promote synaptic vesicle docking in mouse hippocampal neurons. Elife 10.

Voleti R, Tomchick DR, Sudhof TC, Rizo J (2017) Exceptionally tight membrane-binding may
explain the key role of the synaptotagmin-7 C(2)A domain in asynchronous
neurotransmitter release. Proc Natl Acad Sci U S A 114:E8518-E8527.

Wang LY, Neher E, Taschenberger H (2008) Synaptic vesicles in mature calyx of Held synapses
sense higher nanodomain calcium concentrations during action potential-evoked
glutamate release. J Neurosci 28:14450-14458.

Wu LG, Chan CY (2022) Multiple Roles of Actin in Exo- and Endocytosis. Front Synaptic
Neurosci 14:841704.

Yau JO, McNally GP (2023) The Rescorla-Wagner model, prediction error, and fear learning.
Neurobiol Learn Mem 203:107799.

Yoon JY, Lee HR, Ho WK, Lee SH (2020) Disparities in Short-Term Depression Among Prefrontal
Cortex Synapses Sustain Persistent Activity in a Balanced Network. Cereb Cortex
30:113-134.

Zucker RS, Regehr WG (2002) Short-term synaptic plasticity. Annu Rev Physiol 64:355-405.

Figure 1. Frequency-dependence of short-term synaptic plasticity at prelimbic L2/3

excitatory synapses.

(**A**) In utero electroporation following injection of the plasmid (CAG-oChIEF-tdTomato) into

the ventricle of an embryo (E17.5). (**B**) Representative images showing specific expression of

oChIEF-tdTomato in L2/3 pyramidal cells after IUE. The red fluorescence of tdTomato clearly

visualizes oChIEF-expressing cell bodies in L2/3 and axons in L5. Scale bar: 1 mm, 100 μm, 50

μm from left to right. (**C**) Recording schematic showing photostimulation of oChIEF-expressing

axon fibers of transfected PCs (*filled triangles*) and a whole-cell recording from non-

transfected PCs (*empty triangles*) or FSINs (*empty circles*). A collimated DMD-coupled LED was

used to confine the area of excitation (typically 3–4 μm in diameter) to a small region (*blue*

circle) near the soma. (**D, E**) Representative traces for EPSCs averaged over 10 trials at each

frequency (*left*) and average amplitudes of baseline-normalized EPSCs (*right*) during 20-pulse

trains at frequencies from 5 to 40 Hz at PC-PC (*D*; n = 12, 9, 21, 10 cells for 5, 10, 20, 40 Hz,

respectively) and PC-FSIN (*E*; n = 8, 9, 10, 10) synapses. Each data point was normalized to the

average of the first EPSC. (**F**) Baseline-normalized amplitudes of steady-state EPSCs ($EPSC_{ss}$;

black symbols) and synaptic efficacy ($EPSC_{ss} \times f$; *red symbols*) as a function of stimulation

frequency (*f*) at PC–PC synapses. $EPSC_{ss}$ was measured from the average of last 5 EPSCs from

the 20-pulse trains (n = 12, 9, 21, 10). (**G**) Paired pulse ratio (PPR) as a function of inter-spike

intervals (n = 25, 9, 25, 22, 43 for 200, 100, 50, 25, 20 ms ISI, respectively) at PC-PC synapses.

(**H**) $EPSC_{ss}$ and synaptic efficacy at PC-FSIN (n = 8, 9, 10, 10). (**I**) PPR at PC-FSIN (n = 8, 9, 10,

15, 34). *Gray symbols*, individual data.

Figure 2. Delayed facilitation results from slow activation of Ca^{2+}-dependent vesicle replenishment at a constantly high vesicular fusion probability

(**A**) Representative traces (*left*) and mean baseline-normalized amplitudes (*right*) of EPSCs evoked by 30-pulse trains at 40 Hz in control (n = 21; *black*) and in the presence of 50 µM EGTA-AM (n = 14; *green*). (**B**) Mean values for the first EPSC amplitude (EPSC$_1$, *left*) and PPR (*right*) from the experiments displayed in (A). *Gray symbols*, individual data. (**C**) Plot of rate constants for short-term facilitation (k$_{STF}$) as a function of stimulation frequency (*f$_{stim}$*), showing a linear relationship. The linear regression line (*black*) is shown fitted to k$_{STF}$ values, estimated from Figure 1. (**D**) Representative EPSC traces (*left*) and average of baseline-normalized EPSCs (*right*) evoked by 12-pulse stimulation at 5 Hz, followed by 40 Hz 7-pulse train (n = 12). Note that slowly developing facilitation was converted to rapid facilitation after strong PPD. (**E**) Mean values for PPR at 40 Hz. The baseline PPR was reproduced from Figure 1G and the PPR during 5 Hz train was calculated as (13th EPSC) / (12th EPSC). *Gray symbols*, individual data. All statistical data are represented as mean ± S.E.M.; n.s. = not significant; **, P＜0.01; unpaired t-test.

Figure 3. Low baseline occupancy of release sites and its increase during facilitation and post-tetanic augmentation

(**A, B**) *Left*, Representative EPSCs evoked by 5 and 40 Hz train stimulation (*black*, 5 Hz; *purple*, 40 Hz). *Right*, Variance-mean plots of EPSCs amplitude from averaged EPSCs recorded at PC-PC (A; n = 12, 17 for 5 and 40 Hz, respectively) and PC-FSIN (B; n = 8, 10) synapses. The data were fitted using multiple-probability fluctuation analysis (MPFA). Error bars are omitted for clarity. The 1st EPSC of 5 Hz train (*broken line*) was used to estimate the resting level of p_{occ}. *Filled circles* were measured from post-tetanic augmented EPSCs (*A*, n = 12, 12, 9; *B*, n = 7, 7, 7 for 0.1, 0.2, 0.5 s IBIs, respectively). (**C-E**) Post-tetanic augmentation (PTA) experiments at PC-PC (*top*) and PC-FSIN (*bottom*) synapses. (**C**) Representative traces for EPSCs evoked by double 40 Hz train stimulations separated by 0.5 s. (**D**) Mean baseline-normalized amplitudes of EPSCs evoked by double 40 Hz trains at different inter-burst intervals (IBIs, 0.1, 0.2, 0.5, 1, 2, 5, 10 s). *Upper*, PC-PC synapse (n = 12, 12, 21, 11, 11, 16, 11 from short to long IBIs,

respectively). *Lower*, PC-FSIN synapse (n = 10, 9, 8, 9, 9, 7, 9). (**E**) PTA time course. The

baseline-normalized amplitudes of 1^{st} EPSC from the 2^{nd} train were plotted as a function of

IBIs.

Figure 4. Pharmacological experiments reveal specific molecular mechanisms underlying

vesicle loading processes

(**A-C**) (**a**) Representative EPSC traces (*left*) and mean baseline-normalized EPSCs (*right*) evoked

by double 40 Hz train stimulations separated by 0.5 s inter-burst interval (IBI) in control and

in the presence of 5 μM U73122 (*Aa*, n = 7, *orange*), 20 μM OAG (*Ba*, n = 12, *cyan*) or 100 μM

dynasore (*Ca*, n = 10, *blue*). (**b**) Mean values for baseline EPSCs (EPSC$_1$, *left*) and augmentation

(*right*) from the experiments shown in corresponding *a* panel. (**Da**) Representative EPSC

traces evoked by paired pulses (*left*) and mean values for baseline EPSC amplitude (*middle*)

and PPR (*right*) before and after applying 20 μM LatB (n = 6). (**Db**) Representative EPSC traces

(*left, upper*) and average of normalized EPSCs (*left, lower*) evoked by double 40 Hz train

stimulation separated by 0.5 s in control (n = 21) and in 20 μM LatB conditions (n = 16; *pink*).

Right, Mean values for augmentation in control and LatB conditions. *Gray symbols*, individual

data. All statistical data are represented as mean ± S.E.M., *, P＜0.05; **, P＜0.01; ***, P＜

0.001; unpaired or paired t-test; n.s. = not significant.

Figure 5. STF at both types of local excitatory synapses is abolished by Syt7 KD

(**A, B**) Representative EPSC traces (*left*) and mean baseline-normalized amplitudes of EPSCs (*right*) evoked by 20-pulse trains at 5 to 40 Hz. STP was measured at PC-PC (*A*; n = 10, 9, 12, 9) and PC-FSIN (*B*; n = 9, 8, 10, 9) synapses, in which presynaptic Syt7 transcripts were depleted (Syt7 KD). Syt7 KD pyramidal cells are indicated as *black triangles* on the top. For comparison, STP in WT synapses is reproduced from Figure 1 (*dotted lines*). Same frequency color codes were used as in Figure 1. (**C**) Baseline-normalized amplitudes of steady-state EPSC (EPSC$_{ss}$; *black symbols*) and synaptic efficacy (EPSC$_{ss}$ × *f*; *red symbols*) as a function of

82   stimulation frequency (f) at PC-PC synapses. $EPSC_{ss}$ was defined as the average of last 5 EPSC

83   amplitudes from 20-pulse trains. (**D**) PPR as a function of inter-spike intervals (n = 10, 9, 12,

84   18, 26). (**E**) $EPSC_{ss}$ and synaptic efficacy at PC-FSIN synapses. (**F**) PPR at PC-FSIN synapses (n =

85   9, 8, 10, 12, 24). *Gray symbols*, individual data.

Figure 6. Syt7 KD synapses exhibit complementary changes in the number of release sites and their vesicle occupancy

(**A, B**) *Left*, Representative traces of EPSCs evoked by 5 and 40 Hz train stimulations (*black*, 5 Hz; *purple*, 40 Hz). *Right*, Variance-mean plots of EPSCs amplitude from averaged EPSCs recorded at PC-PC (*A*; n = 9, 15, 12, 10, 9) and PC-FSIN (*B*; n = 4, 14, 10, 9, 8) synapses in which presynaptic Syt7 has been knocked down. The data were fitted using MPFA and error bars are omitted for clarity. The mean 1st EPSC amplitude of 5 Hz train (*dotted line*) was used for estimation of baseline p_{occ}. *Broken lines* indicate MPFA fits to variance-mean plot of WT synapses reproduced from Figure 2. (**C-E**) Recovery experiments at PC-PC (*top*) and PC-FSIN (*bottom*) synapses in which presynaptic Syt7 was knocked-down. (**C**) Representative EPSCs evoked by double 40 Hz train stimulations separated by 0.5 s. (**D**) Mean baseline-normalized amplitudes of EPSCs evoked by double 40 Hz trains at PC-PC (n = 12, 10, 9, 9, 10, 9, 8) and PC-FSIN (n = 10, 9, 10, 11, 8, 9, 10) synapses at different interburst intervals (IBIs). (**E**) PTA time

101   course. Baseline-normalized amplitudes of 1st EPSC from the 2nd burst were plotted as a

102   function of various IBIs. *Dotted lines* indicate augmented EPSCs in the WT reproduced from

103   Figure 2.

Figure 7. Recovery of TS vesicles following depletion is accelerated by Syt7

(**A-C**). Recovery experiments at PC-PC synapses in WT (*A*) and Syt7 KD (*B*). Mean baseline-normalized amplitudes of EPSCs evoked by two consecutive 3-pulse 40 Hz trains in WT (n = 16, 13, 11, 11, 11, 9 from short to long IBIs, respectively) and KD (n = 13, 11, 12, 8, 10, 11) synapses at different IBIs (0.1, 0.2, 0.5, 1, 2, 5 s). *Inset*, representative traces of EPSCs evoked by two consecutive 3-pulse 40 Hz train stimulations separated by 0.5 s. (**D-F**). Recovery experiments at PC-FSIN synapses in WT (*D*) and Syt7 KD (*E*). Mean baseline-normalized amplitudes of EPSCs evoked by two consecutive 3-pulse 40 Hz trains in WT (n = 9, 11, 11, 9, 10, 13) and KD (n = 9, 12, 9, 7, 10, 9) synapses at different IBIs. *Inset*, representative traces of EPSCs evoked by two consecutive 3-pulse 40 Hz train stimulations separated by 0.5 s. (**C, F**) Recovery time course. Baseline-normalized amplitudes of 1st EPSC from the 2nd burst were plotted as a function of various IBIs.

Figure 8. Behavioral effects of Syt7 deficiency in L2/3 PCs of the mPFC

(**A**) *Top*, Representative images showing bilateral expression of U6-GFP in L2/3 of PCs after

IUE at E17.5. Scale bar: 1 mm, 100 μm. *Bottom*, Schematic of trace fear conditioning and

extinction (tone test) protocol. (**B**) Freezing behavior of control (expressing scrambled shRNA,

Scr) and Syt7 KD rats during acquisition of tFC. (**C**) Freezing ratio during tone tests on following

days. Data are shown as average freezing during T1–T4, T5–T8, or T9–T12 (T, trials). The

freezing on T1-T4 was significantly lower in KD rats suggesting that formation of trace memory

was impaired in Syt7 KD rats (n = 10, 11 for Ctrl and KD, respectively; P = 0.0007, F(1, 19) =

16.43, two-way repeated measures ANOVA; P = 0.0064, 0.0211, 0.0064, 0.0064; Holm-Sidak

test). (**D**) Representative images of c-Fos immunoreactivity in the prelimbic cortex of control

or Syt7 KD rats 90 min after tFC acquisition. c-Fos (*left*, *red*, Cy5) and GAD67 (*right*, *cyan*, Cy3)

were immunostained in the same brain slice expressing U6-GFP (*middle*; *green*). Scale bar, 50

µm. (**E**) Exemplar images of c-Fos positive neurons expressing GFP (*top*) or GAD67 (*bottom*)

in control (*left*) or Syt7 KD rats (*right*). Scale bar, 50 µm. (**F-H**) Effects of Syt7 KD on c-Fos

density (*F*) and percentage of c-Fos positive neurons co-labeled with GAD67 (*G*) or GFP (*H*) in

L2/3 or L5 of prelimbic cortex (n = 9, 9 for Ctrl and KD, respectively). *Open symbols*, individual

data. All statistical data are represented as mean ± S.E.M.; ****, $P < 0.0001$; unpaired t-test.

Figure 9. STP model in light of known Ca²⁺ binding kinetics of Syt7

(**A-D**) *Left*, Schematic of allosteric calcium binding to Syt7. The number of Ca^{2+} bound to Syt7

was denoted as # in 'S#' in the reaction scheme. k_{on} = 7/μM/s, k_{off} = 10/s. *Middle*, Simulated

changes of k_1 (*black*) in response to 5 Hz train of Ca^{2+} transients (*light blue traces*). The priming

step of the simple refilling model was assumed to be catalyzed by full Ca^{2+}-bound from of Syt7.

Accordingly, Ca^{2+}-dependent increase in k_1 was calculated as $K_{1,max}$ multiplied by a fraction of

full Ca^{2+}-bound form of Syt7. We assumed that local $[Ca^{2+}]_i(t)$ follows a Gaussian function: $(1/\sigma$

$\sqrt{2\pi})$ exp$[-(t-t_p)^2/2\sigma^2]$, in which t_p = 0.25 ms and σ = 0.085 ms. *Right*, Fits of the Syt7 model to

the STP data. To fit this model to the STP data, $K_{1,max}$ was set to 300/s (*A*), 220/s (*B*), and 180/s

(*C*). Cooperativity factor (*b*) was set to 0.35 (*A*), 0.2 (*B*), and 0.05 (*C*). k_1 is set to be constant

for Syt7 KD (*D*). Same frequency color codes were used as in Figure 1.

16