Progressive overfilling of readily releasable pool underlies short-term facilitation

at recurrent excitatory synapses in layer 2/3 of the rat prefrontal cortex

Jiwoo Shin[1,2,3], Seung Yeon Lee[2], Yujin Kim[1,2,3*], Suk-Ho Lee[1,2,3*]

1. Department of Brain and Cognitive Sciences, College of Natural Sciences, Seoul National

University. 2. Department of Physiology, Seoul National University College of Medicine. 03080,

Seoul, South Korea. 3. Neuroscience Research Institute, Seoul National University College of

Medicine, Seoul 03080, Republic of Korea.

corresponding authors

Yujin Kim Ph.D. (kim.y@snu.ac.kr)

Cell Physiology Lab

Department of Physiology

Seoul National Univ. College of Medicine

Suk-Ho Lee M.D., Ph.D. (leesukho@snu.ac.kr)

Cell Physiology Lab

Department of Physiology

Seoul National Univ. College of Medicine

103 Daehakro, 03080, Seoul, South Korea

Phone) +82-2-740-8222

Abstract

Short-term facilitation of recurrent excitatory synapses within the cortical network has been proposed to support persistent activity during working memory tasks, yet the underlying mechanisms remain poorly understood. We characterized short-term plasticity at the local excitatory synapses in layer 2/3 of the rat medial prefrontal cortex and studied its presynaptic mechanisms. Low-frequency stimulation induced slowly developing facilitation, whereas high-frequency stimulation initially induced strong depression followed by rapid facilitation. This non-monotonic delayed facilitation after a brief depression resulted from a high vesicular fusion probability and slow activation of Ca^{2+}-dependent vesicle replenishment, which led to the overfilling of release sites beyond their basal occupancy. Pharmacological and gene knockdown (KD) experiments revealed that the facilitation was mediated by phospholipase C/diacylglycerol signaling and synaptotagmin 7 (Syt7). Notably, Syt7 KD abolished facilitation and slowed the refilling rate of vesicles with high fusion probability. Furthermore, Syt7 deficiency in layer 2/3 pyramidal neurons impaired the acquisition of trace fear memory and reduced c-Fos activity. In conclusion, Ca^{2+}- and Syt7-dependent overfilling of release sites mediates synaptic facilitation at layer 2/3 recurrent excitatory synapses and contributes to temporal associative learning.

Introduction

Synapses in the mammalian central nervous system undergo activity-dependent modulation of synaptic weight during repetitive stimulation, known as short-term synaptic plasticity (STP). STP is mediated by presynaptic mechanisms that regulate the release probability (p_r), size of the readily releasable pool (RRP), and kinetics of vesicle replenishment of RRP (Zucker and Regehr, 2002; Jackman and Regehr, 2017; Neher and Brose, 2018). Short-term depression has been observed at synapses with a high baseline p_r, primarily because of the rapid depletion of releasable vesicles during high-frequency stimulation (HFS) (Lin et al., 2022). Conversely, vesicle depletion is less pronounced at synapses with a low baseline p_r, and cumulative increases in p_r during HFS are considered to mediate short-term facilitation (Rozov et al., 2001; Pan and Zucker, 2009; Aldahabi et al., 2024). Central synapses exhibit a wide spectrum of STP depending on different combinations of presynaptic factors mediating short-term depression and facilitation.

Previously, facilitation was modeled as an increase in p_r after an action potential (AP) and its slow decay during the inter-spike interval (ISI), resulting in a cumulative increase in p_r during HFS (Markram et al., 1998; Dittman et al., 2000). Recent studies have suggested that the number of docking (or release) sites in an active zone is limited and only partially occupied by releasable vesicles in the resting states (Miki et al., 2016; Pulido and Marty, 2018; Malagon et al., 2020; Lin et al., 2022). Therefore, p_r is determined by the vesicular fusion probability (p_v) and release site occupancy (p_{occ}). However, distinguishing the individual contributions of p_v and p_{occ} to p_r is challenging (Silva et al., 2021; Neher, 2024). It is also unclear whether facilitation is mediated by activity-dependent increases in the p_v, p_{occ}, or both.

Two recent models for vesicle priming and release view the 'priming' as a two-step reversible

process that leads to a release-ready state. The release-ready vesicles could be vesicles in a

tightly docked state (TS) or those occupying specialized docking sites (DS), and are supplied

from vesicles in a loosely docked state (LS) or those occupying a distinct replacement sites

(RS), which are called LS/TS model (Aldahabi et al., 2024; Neher, 2024) and RS/DS model (Miki

et al., 2016; Silva et al., 2021), respectively. Common to both models is the assumption that

release-ready vesicles constitute only a fraction of all docked vesicles under resting conditions

and that this fraction is up- and down-regulated by synaptic activity.

Synaptotagmin-7 (Syt7) was recently found to play a crucial role in facilitation at different

types of central synapses (Jackman et al., 2016; Chen et al., 2017; Martinetti et al., 2022). In

addition to its role in facilitation, Syt7 accelerates refilling of the RRP after depletion and

mediates asynchronous release (Bacaj et al., 2013; Liu et al., 2014; Jackman et al., 2016;

Tawfik et al., 2021). Syt7-dependent synaptic facilitation is interpreted as an activity-

dependent increase in p_v (Turecek et al., 2016; Jackman and Regehr, 2017; Turecek et al., 2017;

Norman et al., 2023). However, it is unclear whether Syt7 contributes to facilitation through

a Ca^{2+}-dependent increase in p_{occ} (i.e., overfilling). Overfilling has been proposed as a

mechanism supporting facilitation at the Drosophila neuromuscular junction (Kobbersmed et

al., 2020), yet whether this process also occurs at the mammalian central synapses mediated

by Syt7 has not been studied.

Each of layer 2/3 (L2/3) and layer 5 (L5) of neocortex displays intralaminar excitatory synapses

between pyramidal cells comprising a recurrent network (Holmgren et al., 2003; Thomson

and Lamy, 2007). Short-term facilitation at recurrent excitatory synapses in a cortical network

may play a crucial role in maintaining working memory (Mongillo et al., 2008; Mongillo et al.,

2012; Hansel and Mato, 2013). Although L2/3 pyramidal cells (PCs) represent task-relevant

information during a delay period in working memory tasks (Fujisawa et al., 2008; Ozdemir et

al., 2020), most studies on local excitatory recurrent synapses in the neocortex have focused

on L5 exhibiting short-term depression (Reyes and Sakmann, 1999; Hempel et al., 2000; Yoon

et al., 2020). Further, the synaptic properties of excitatory synapses in L2/3 remain poorly

understood.

In the present study, we found that HFS of local excitatory synapses in L2/3 of the medial

prefrontal cortex (mPFC) induced initial strong depression followed by delayed facilitation,

whereas low-frequency stimulation induced slow monotonous facilitation. This unique form

of STP could be explained by 1) high p_v of RRP vesicles at rest and throughout a train

stimulation; 2) the low resting vesicular occupancy of RRP and its activity-dependent increase.

These conditions led to delayed facilitation through progressive overfilling of release sites

during HFS with high p_v vesicles beyond the resting occupancy. Syt7 knockdown (KD) in L2/3

PCs completely abolished these synaptic enhancements by inhibiting the Ca^{2+}-dependent

increase in release site occupancy. Finally, behavioral tests for trace fear conditioning showed

that Syt7 KD impaired the acquisition of trace fear memory and reduced c-Fos expression in

the mPFC. Collectively, these results support that Ca^{2+}- and Syt7-dependent overfilling of

release sites mediates synaptic facilitation at L2/3 recurrent excitatory synapses and

contributes to temporal associative learning.

Results

Frequency dependence of STP at local excitatory synapses in L2/3 of the prelimbic cortex

We examined STP profiles at recurrent excitatory synapses between L2/3 PCs and at synapses from L2/3-PCs to fast-spiking interneurons (FSINs) in the prelimbic area of the rat mPFC. The PCs and FSINs were identified based on their distinct intrinsic properties and morphologies (*Figure 1—figure supplement 1*). To stimulate presynaptic PCs, L2/3 PCs were transfected with a plasmid encoding oChIEF using *in utero* electroporation (IUE; *Figure 1A*; Lee et al., 2024). Photo-stimulation of the oChIEF-expressing cell soma at 40 Hz consistently evoked single APs with high fidelity throughout the 600-pulse trains (*Figure 1—figure supplement 2A-B*), which was consistent with little use-dependent inactivation of oChIEF (Lin et al., 2009; Hass and Glickfeld, 2016).

Given that IUE transfected approximately 10-20% of L2/3 PCs (*Figure 1A*), we made a whole-cell patch on a non-transfected L2/3 PC or an FSIN (*Figure 1C*). To stimulate a minimal number of excitatory axon fibers, EPSCs were evoked using minimal optical stimulation (*Figure 1—figure supplement 3A*; 470 nm, 3-4 μm in radius and 3-5 ms in duration). EPSCs evoked by minimal stimulation were examined to determine the uniformity of the kinetic properties (*Figure 1—figure supplement 3B-C*). Moreover, we reassessed the kinetics of EPSCs during trains *post hoc* to ensure uniformity. EPSCs were evoked by 20-pulse trains at 4 different frequencies (5, 10, 20, and 40 Hz) to cover the range of firing rates of PCs observed *in vivo* during working memory tasks (Baeg et al., 2003). Excitatory synapses from PCs onto both other PCs and FSINs (PC–PC and PC–FSIN) showed strong (~2-fold enhancement) and lasting

(several seconds) short-term facilitation at 20 Hz and lower frequencies (*Figure 1D-E*). The

PC-PC synapses exhibited stronger facilitation than did the PC-FSIN synapses.

Steady-state EPSCs at the PC-PC synapses were not significantly different from 5 to 20 Hz

(*Figure 1F*). Frequency invariance at the PC-PC synapses suggests that Ca^{2+}-dependent

synaptic facilitation may counteract vesicle depletion in a frequency-dependent manner

(Turecek et al., 2017; Lin et al., 2022). Despite facilitation at all frequencies, the paired pulse

ratio (PPR) rapidly decreased as the ISI decreased (*Figure 1G, I*). At 40 Hz stimulation, both

PC-PC and PC-FSIN synapses initially underwent strong depression then followed by

facilitation in a few stimuli (*Figure 1Dd, Ed*). Such delayed facilitation after paired pulse

depression (PPD) has not been previously observed in central synapses except for a slight

increase in EPSC at the cerebellar synapses under artificially depolarized conditions (Pulido

and Marty, 2018).

We confirmed that optically evoked EPSCs were strictly dependent on AP generation (*Figure*

1—figure supplement 4A), suggesting that EPSCs are evoked by optical stimulation of

presynaptic axon fibers rather than by direct depolarization of presynaptic boutons.

Additionally, dual patch-clamp techniques showed that the STP of optically evoked EPSCs did

not differ from the electrically evoked STP of EPSCs at the PC-FSIN synapses (*Figure 1—figure*

supplement 4B). Finally, we confirmed that AMPA receptor (AMPAR) desensitization was not

responsible for PPD at 40 Hz, ruling out the possible effects of postsynaptic factors on STP

(*Figure 1—figure supplement 5A*), and that AMPAR saturation was not significant during

facilitation (*Figure 1—figure supplement 5B*).

Delayed facilitation results from slow activation of Ca^{2+}-dependent vesicle replenishment at a constantly high vesicular fusion probability

The strong PPD at 40 Hz suggests a high vesicular fusion probability (p_v) and likely tight coupling of the releasable vesicles to a Ca^{2+} source. To test this hypothesis, we examined the effects of EGTA, a slow calcium chelator, on synaptic transmission at the PC-PC synapses. The baseline EPSC amplitude ($EPSC_1$) was not affected by incubation of the slice in the bath solution containing 50 μM EGTA-AM for longer than 20 min, supporting tight coupling. Moreover, the PPR was decreased to 0.19, and subsequent facilitation slowed but was still distinct (*Figure 2A-B*). These findings indicate that the initial p_v should be higher than 0.81 considering vesicle recruitment during ISI. The effects of EGTA-AM were confirmed in the same cell by measuring the baseline EPSC and PPR before and after applying EGTA-AM to the bath. $EPSC_1$ remained unchanged, while PPR was reduced by half (*Figure 2—figure supplement 1*).

Notably, the application of EGTA-AM markedly slowed facilitation at 40 Hz (*Figure 2A, C*). The frequency invariance of steady-state EPSCs (*Figure 1F*) and the slower facilitation at 40 Hz in the presence of EGTA (*Figure 2A*) imply a Ca^{2+}-dependent increase in the refilling rate of the RRP and/or an increase in the p_v of reluctant vesicles in light of previous studies (Hosoi et al., 2007; Liu et al., 2014; Ritzau-Jost et al., 2014; Turecek et al., 2016; Ritzau-Jost et al., 2018; Kusick et al., 2020). Moreover, facilitation was accelerated at higher stimulation frequencies. The plot of the rate constant for the increase in EPSCs (denoted as $\mathbf{k}_{STF}$, the rate constant for the development of short-term facilitation) as a function of the stimulation frequency (f_{stim})

revealed a linear relationship (*Figure 2C*). k_{STF} in the presence of EGTA-AM was distinctly

below this relationship, supporting the Ca^{2+}-dependent acceleration of facilitation.

k_{STF} was also accelerated when the stimulation frequency was increased from 5 Hz to 40 Hz

(*Figure 2D*; 1.27 ± 0.18/s to 24.27 ± 3.44/s). Notably, the second 40-Hz train stimulation led

to strong PPD, followed by accelerated facilitation (*Figure 2D-E*). The strong PPD suggests that

the vesicular fusion probability kept high during the 5-Hz stimulation, arguing against the

possibility for an increase in p_v. Both two-step sequential priming models, LS/TS and RS/DS

models, are suitable to interpret our data and we will refer to the different populations of

vesicles as TS and LS vesicles, respectively (Taschenberger et al., 2016; Neher and Brose, 2018;

Lin et al., 2022; Neher, 2024). Within this framework, vesicles are released from TS with high

p_v both in the resting state and during facilitation, and the delayed facilitation is seen as a

result of Ca^{2+}-dependent progressive increase in the occupancy of TS vesicles on docking sites.

Presynaptic calcium measurements at axonal boutons of L2/3 pyramidal cells

To test contribution of Ca^{2+} channel regulation to STP (Dobrunz et al., 1997; Jackman and

Regehr, 2017), we estimated total amount of Ca^{2+} increments during 40 Hz AP trains at axonal

boutons of L2/3 PCs in the mPFC. To this end, we measured AP-evoked Ca^{2+} transients (AP-

CaTs) using two-dye ratiometry techniques, and estimated calcium binding ratio from AP-CaTs

(*Figure 2—figure supplement 2A-C*). Measuring the amplitudes and decay time constants of

AP-CaTs at different concentrations of Fluo-5F (150, 250 or 500 μM), and plotting them as a

function of calcium binding ratio of Fluo-5F ($κ_B$), we estimated the amplitude of AP-CaTs (A_{Ca}

= 1.16 μM), Ca^{2+} decay time constant (τ_c = 43 ms) in the absence exogenous Ca^{2+} buffer, and

endogenous static Ca^{2+} binding ratio (κ_S = 96.4 or 77.3 estimated from *Figure 2—figure*

supplement 2D-E, respectively; *See Materials and Methods*). The resting $[Ca^{2+}]$ ($[Ca^{2+}]_{rest}$) was

measured as 50 nM (*Figure 2—figure supplement 2F*). Estimating the total Ca^{2+} increments

for the first four AP-CaTs in a 40 Hz train according to Equation 2 (*Materials and Methods*),

we found no evidence for inactivation or facilitation of calcium influx during a 40 Hz train

(*Figure 2—figure supplement 2G-I*), arguing against the possibility that Ca^{2+} channel

inactivation and facilitation contribute to strong PPD and delayed facilitation during a 40 Hz

train.

Release sites have low baseline occupancy and these are increased during facilitation and post-

tetanic augmentation

Given that p_v was greater than 0.8, the two-fold increase in EPSC during 20 pulse train

stimulations could not be explained by an increase in p_v. Recent studies have shown that

vesicle release sites (N) are limited in number and are partially occupied at rest (Miki et al.,

2016; Pulido and Marty, 2018; Malagon et al., 2020; Lin et al., 2022). Under such a high p_v,

two-fold synaptic facilitation should be attributed to an increase in p_{occ}. The vesicle

replenishment rate is accelerated during HFS in a Ca^{2+}-dependent manner (*Figure 2A-C*; Hosoi

et al., 2007). This may lead to the overfilling of release sites beyond their basal vesicular

occupancy, which in turn may increase EPSC under high p_v. To test this hypothesis, we

determined the quantal parameters at the L2/3 local excitatory synapses. We first measured

the quantal size from asynchronous release events after replacing extracellular Ca^{2+} with Sr^{2+},

as previously described (Yoon et al., 2020) (*Figure 3—figure supplement 1A-B*). The mean

and coefficient of variance of the quantal sizes were 24 pA and 0.25 at PC-PC synapses,

respectively, and were 30 pA and 0.31 at PC-FSIN synapses (*Figure 3—figure supplement 1C*).

Next, we performed a variance-mean (V-M) analysis of phasic EPSCs at excitatory synapses

during train stimulation at 5 Hz and 40 Hz (Scheuss et al., 2002). Applying multiple-probability

fluctuation analysis (MPFA) using intra-site quantal variability of 0.2 (Valera et al., 2012), the

V-M plot was best fitted by the number of release sites (N) of 5.3 and the quantal size (q) of

25 pA at the PC-PC synapses (*Figure 3A*). The p_r for $EPSC_1$ was calculated as 0.32. We

performed the same analysis for the PC-FSIN synapses and found N = 5.3, q = 35 pA, and p_r

for $EPSC_1$ = 0.31 (*Figure 3B*). Our estimate for N was smaller but comparable to the estimate

(6.9) at excitatory synapses in L2/3 of the mouse barrel cortex (Holler et al., 2021).

The p_r estimate was clearly smaller than the possible minimum value for p_v (0.81) estimated

from the PPR data (*Figure 2B*). Given that p_r is a product of p_{occ} and p_v (Neher, 2024), a p_r

smaller than p_v indicates the partial occupancy of release sites in the resting state. If synaptic

facilitation were mediated by the recruitment of new release sites, it would be accompanied

by an increase in the variance of augmented EPSCs (Valera et al., 2012; Kobbersmed et al.,

2020). However, the augmented EPSCs during facilitation followed the same parabola in the

V-M plot (*Figure 3A-B*). Therefore, facilitation appears to be mediated by the progressive

overfilling of a finite number of release sites beyond the basal occupancy. In the next section,

we show that p_v is close to unity based on high double failure rate upon paired pulse

stimulation (*Figure 3—figure supplement 2*) and little effect of extracellular [Ca^{2+}] ([Ca^{2+}]$_o$)

elevation on baseline EPSC amplitudes (*Figure 3—figure supplement 3*). Assuming this, the

baseline p_r at the first pulse estimated from the fitted parabolic curve can be largely regarded

as a baseline p_{occ} (c.a. 30%), and the two-fold increase in EPSCs during facilitation can be

attributed to an increase of up to 60% in p_{occ}.

We investigated the time course of recovery from facilitation and post-tetanic augmentation

(PTA) by applying dual-train stimulations (30 pulses at 40 Hz) separated by different inter-

burst intervals (IBIs; *Figure 3C-D*). The initial EPSCs in the second train were enhanced by more

than twofold, and followed by a strong PPD, suggesting that fusion probability stays high in

the IBIs and thus synaptic transmission is carried by TS vesicles. The decay time course of PTA

was characterized by two phases (initially fast and later slow; *Figure 3E*), which may reflect a

two-phase Ca^{2+} decay or two distinct recovery processes that occur sequentially. The variance

of potentiated EPSCs measured at IBIs of 0.1, 0.2, and 0.5 s was not different from that of

EPSCs at the peak of facilitation (filled circles, *Figure 3A-B*). Given that the same RRP mediates

both facilitation and augmentation and that post-tetanic EPSCs is carried by TS vesicles similar

to the baseline EPSCs, this plot suggests that PTA, similar to short-term facilitation, is

mediated by an increase in the TS vesicle occupancy.

High double failure rate and little effect of $[Ca^{2+}]_o$ elevation on baseline EPSC amplitudes

suggest that vesicular fusion probability is close to one

The vesicle dynamics of STP has been explained using a simple refilling model (Hosoi et al.,

2007; Neher and Sakaba, 2008). This model assumes that the vesicles are reversibly docked

to a limited number of release sites (N) with forward and reverse rate constants (denoted as

k_1 and b_1, respectively). Under the framework of the simple refilling model (see *Materials and*

Methods), the rate constant for the refilling of the RRP after depletion by the first pulse is the

sum of the baseline k_1 and b_1. This is estimated as 23/s from the plot of PPR as a function of

ISIs (*Figure 3—figure supplement 2A*). From the baseline occupancy of 0.3 and $p_{occ} = k_1 / (k_1$

$+ b_1)$, we estimated k_1 to be 6.9/s.

When paired APs were applied with an ISI of 20 ms, the failure rate at 1^{st} pulse, $P(F_1)$, was

10.6%, and the probability of two consecutive failures, $P(F_1, F_2)$, was 6.2% (*Figure 3—figure*

supplement 2A). From the equation $P(F_1) = (1 - p_r)^N$, p_r was calculated as 0.312 when N = 6,

comparable to p_r (= 0.32) estimated from the V-M analysis (*Figure 3A*). We calculated $P(F_1, F_2)$

under the framework of a simple refilling model, with p_v and k_1 set as free parameters and

other parameters set according to the following relationships: $P(F_1) = (1 - p_r)^N$, $p_r = p_v \cdot p_{occ}$,

and $p_{occ} = k_1 / (k_1 + b_1)$. The calculated values for $P(F_1, F_2)$ and their difference from the

observed value (6.2%) are shown in the plane of k_1 *vs.* p_v in *Figure 3—figure supplement 2B-*

C. Given that k_1 was greater than 5/s, the difference between the calculated and observed

values of $P(F_1,F_2)$ was minimal when $p_v = 1$ (*Figure 3—figure supplement 2C*).

To find optimal values for k_1 and p_v that best explain observed probabilities shown in *Figure*

3—figure supplement 2Ac, a cost function was implemented as the sum of squared errors

between observed and predicted probability values for the four combinations of success and

failure of 1st and 2nd EPSCs as described in *Materials and Methods*. The minimum of the cost

function (5.63×10^{-5}) was found at $k_1 = 5.21$/s and $p_v = 0.999$, which predicted $P_{11} = 41.5\%$, P_{10}

= 47.9%, P_{01} = 4.93%, and P_{00} = 5.67% (subscript 1 and 0 denote success and failure,

respectively).

To further validate that baseline p_v is near to unity, we examined EPSC changes following

elevation of $[Ca^{2+}]_o$ from 1.3 to 2.5 mM. According to the fourth-power relationship between

synaptic responses and $[Ca^{2+}]_o$ (Equation 3 in (Dodge Jr and Rahamimoff, 1967)), a 3.24-fold

increase in the EPSC amplitude was expected. However, the EPSC amplitude increased only

1.23-fold in average, a change that was not statistically significant (*Figure 3—figure*

supplement 3A-B). This small response suggests that p_v is already saturated at rest, limiting

the dynamic range for further enhancement through increased calcium influx. Recent

morphological and functional studies revealed that elevation of $[Ca^{2+}]_o$ induces an increase in

the number of TS or docked vesicles to a similar extent as our observation (Kusick et al., 2020;

Lin et al., 2025), raising a possibility that an increase in p_{occ} is responsible for the 1.23-fold

increase in EPSC at high $[Ca^{2+}]_o$. A slight but significant reduction in PPR was observed under

high $[Ca^{2+}]_o$ too. An increase in p_{occ} is thought to be associated with that in the baseline vesicle

refilling rate. While PPR is always reduced by an increase in p_v, the effects of refilling rate to

PPR is complicated. For example, PPR can be reduced by both a decrease (*Figure 2—figure*

supplement 1) and an increase (Lin et al., 2025) in the refilling rate induced by EGTA-AM and

PDBu, respectively. Thus, the slight reduction in PPR is not contradictory to the possible

contribution of p_{occ} to the high $[Ca^{2+}]_o$ effects. Notably, when slices were pre-incubated with

50 μM EGTA-AM, elevating extracellular $[Ca^{2+}]$ from 1.3 to 2.5 mM produced no significant

change in either baseline EPSC amplitude or PPR (*Figure 3—figure supplement 3C-D*),

supporting that the modest Ca^{2+}-dependence of baseline EPSCs and PPR in the absence of

EGTA is primarily mediated by a change in p_{occ} rather than p_v.

Pharmacological manipulations reveal specific molecular identities underlying vesicle loading

processes

Our results suggest that the baseline occupancy of TS vesicles is low and that the activity-

dependent progressive overfilling of docking sites with TS vesicles is responsible for short-

term facilitation and augmentation. To elucidate the specific molecular link between synaptic

enhancement and Ca^{2+}-dependent overfilling, we examined the effects of several drugs

known to affect vesicle dynamics. We first examined the roles of phospholipase C (PLC) and

diacylglycerol (DAG) by applying 5 μM U73122, a PLC inhibitor, or 20 μM 1-oleoyl-2-acetyl-sn-

glycerol (OAG), a DAG analog. The STP at 40 Hz and PTA at 0.5-s interval were examined before

and after applying each drug to the bath. For both drugs, the facilitation and augmentation

were reduced (*Figure 4A-B*). U73122 did not affect the basal EPSC amplitude (EPSC$_1$) but

resulted in a stronger PPD, slower facilitation, and lower augmentation (*Figure 4A*). These

effects of U73122 suggest the involvement of Ca^{2+}-induced PLC activation in progressive

overfilling. In contrast, OAG increased EPSC$_1$ while maintaining a pronounced PPD (*Figure 4B*),

indicating an increase in the TS vesicle pool size. Meanwhile, OAG slowed k_{STF} and reduced

augmentation, indicating occlusion of activity-dependent synaptic facilitation. Phorbol esters

and DAG are thought to accelerate priming through Munc13 activation (Rhee et al., 2002;

Taschenberger et al., 2016; Aldahabi et al., 2022); they also shorten the bridges that interlink

docked vesicles with the active zone (AZ) membrane (Papantoniou et al., 2023). Collectively,

these suggest that PLC- and DAG-mediated signaling play a role in the overfilling of docking

sites underlying short-term facilitation and augmentation.

Although the acceleration of Ca^{2+}-dependent vesicle refilling supports rapid facilitation at 40

Hz, the later part of the delayed facilitation underwent a slight depression, which was more

pronounced during the second burst (*Figure 3D and Figure 4*). Sustained synaptic

transmission is limited by the spatial availability of docking sites because it is undermined by

accumulation of exocytic remainings in the presynaptic AZ during HFS (Neher, 2010; Haucke

et al., 2011). To verify this, we examined the effects of dynasore (100 μM), a specific inhibitor

of endocytosis (Macia et al., 2006). The treatment with dynasore did not influence $EPSC_1$ or

the early phase of facilitation, but it exacerbated depression during the later phase and

attenuated PTA (*Figure 4C*). This suggests that late synaptic transmission during the 40-Hz

train is limited by site clearance.

Given the involvement of actin polymerization in the multiple steps of vesicle docking and

recycling, we examined the effects of 20 μM latrunculin B (LatB), an inhibitor of actin

polymerization. The effects of LatB were complicated: (1) $EPSC_1$ was decreased, whereas PPR

was increased (*Figure 4Da*) and (2) facilitation and augmentation were reduced (*Figure 4Db*).

The latter effect may result from defects in vesicle replenishment rates, either through the

impaired physical movement of vesicles or disrupted site clearance (Sakaba and Neher, 2003;

Lee et al., 2012; Hallermann and Silver, 2013; Miki et al., 2016), as Ca^{2+}-dependent vesicle

refilling mediates facilitation and augmentation (*Figure 2 and Figure 3*). Meanwhile, the

former effect might be attributed to a shift in equilibrium in the priming states towards a

loosely docked state (LS) at rest. It is unknown whether actin affects vesicular fusogenicity in

neurons, although it facilitates multiple steps of vesicle fusion by enhancing membrane

tension in neuroendocrine cells (Wu and Chan, 2022).

Short-term facilitation at both types of local excitatory synapses in L2/3 is abolished by Syt7

knockdown

We examined whether Syt7 mediates facilitation at the PC-PC and PC-FSIN synapses in L2/3

of the mPFC. To this end, oChIEF and shRNA against Syt7 (shSyt7) were co-expressed in L2/3

PCs using IUE. Syt7 KD resulted in a complete loss of facilitation at all tested stimulation

frequencies (*Figure 5A-B*). Short-term depression and paired-pulse depression were more

pronounced as the stimulation frequency increased, and the frequency invariance of steady-

state EPSC was abolished, implying little contribution of Ca^{2+}-dependent vesicle recruitment

(*Figure 5C-F*). The lack of facilitation could not be attributed to the failure of APs in

presynaptic axons, as optical stimulation elicited reliable APs in Syt7 KD PCs during the 40-Hz

train (*Figure 1—figure supplement 2C*). Syt7 KD had little effect on the basal properties,

including the peak amplitude, rise/decay time, and time-to-peak, of single AP-induced EPSCs

(*Figure 1—figure supplement 3D-E*). Moreover, Syt7 KD did not significantly affect the

intrinsic properties of neurons (*Figure 5—figure supplement 1*).

To examine whether Syt7 KD had any nonautonomous effects on postsynaptic cells, we

measured the quantal size (q) from asynchronous events in the presence of Sr^{2+} and found

that the mean quantal size remained unaltered at Syt7 KD synapses (*Figure 3—figure*

supplement 1C). This indicated that Syt7 did not influence postsynaptic parameters, including

the density and kinetic properties of AMPARs. To test the specificity of shSyt7, we examined

whether co-expression of the shRNA-resistant form of Syt7 restored synaptic facilitation in

both excitatory synapse types. Presynaptic overexpression of shRNA-resistant Syt7 under the

CAG promoter rescued the effect of shRNA at both PC-PC and PC-FSIN synapses (*Figure 5—*

figure supplement 2). These results underscore the crucial role of presynaptic Syt7 in

mediating short-term facilitation at local excitatory synapses in L2/3 of the mPFC.

Syt7 KD synapses exhibit complementary changes in the number of release sites and their

vesicle occupancy

To estimate the quantal parameters in Syt7 KD synapses, we performed V-M analysis of EPSCs

evoked by train stimulation (*Figure 6A-B*). Fitting a parabola to the V-M plot revealed higher

baseline p_{occ} in Syt7 KD synapses than in wild-type (WT) synapses for both synapse types (KD,

0.68 and 0.42; WT, 0.32 and 0.31 for PC-PC and PC-FSIN synapses, respectively). The p_{occ}

monotonously decreased during a train stimulation, as expected from short-term depression.

Moreover, the number of release sites (N) was markedly lower in Syt7 KD synapses (N = 3.5)

than in WT synapses (N = 5.3; *Figure 3A*). Next, we examined the PTA induced by the same

protocol as in *Figure 3* (two 40-Hz train stimulations separated by variable IBIs) at Syt7 KD

synapses (*Figure 6C*). The 40 Hz train-induced synaptic depression quickly recovered during

IBIs, but no PTA was observed at Syt7 KD synapses, indicating its crucial role in augmentation

(*Figure 6D-E*).

TS vesicle recovery following depletion is accelerated by Syt7

Syt7 is recognized not only for its crucial role in facilitation, but also for mediating Ca^{2+}-dependent replenishment of releasable vesicles (Liu et al., 2014; Tawfik et al., 2021). Given that facilitation at L2/3 excitatory synapses is driven by progressive overfilling (*Figure 2 and Figure 3*), the loss of facilitation at Syt7 KD synapses in the present study implies that Syt7 plays a key role in the Ca^{2+}-dependent acceleration of refilling and overfilling of the docking sites with TS vesicles. To test this hypothesis, we examined the recovery kinetics of EPSCs after TS vesicles were depleted by 3 pulses at 40 Hz in wild-type (WT) and Syt7 KD synapses (*Figure 7*). Strong PPD was observed not only in the first burst, but also in the second burst. This suggested that EPSCs in the second burst were mediated by TS vesicles, similar to those in the first burst (*Figure 7A, D*). Therefore, $EPSC_1$ in the second burst can be interpreted as a release from TS vesicles recovering during a given IBI. The recovery time course of $EPSC_1$ in the second burst indicated full recovery of TS vesicles within 100 ms in both excitatory synapse types in WT rats (*Figure 7C, F*). However, this recovery was remarkably slower in Syt7 KD synapses, requiring 5 s for full recovery, indicating that TS vesicle recovery was greatly accelerated by Syt7. Overall, synaptic facilitation and augmentation at local excitatory synapses in L2/3 can be ascribed to progressive activity-dependent increases in the occupancy of TS vesicles (*Figure 2 and Figure 3*), in which Syt7 plays a key role.

Behavioral consequences of Syt7 KD in L2/3 PCs of the mPFC

We previously proposed that the disparity in STP between the PC-PC and PC-IN synapses

results in activity-dependent changes in the ratio of synaptic weights at the PC-PC and PC-IN

synapses (J_{ee}/J_{ie}). This may in turn have a profound influence on persistent activity in a

recurrent network by biasing the E-I balance (Yoon et al., 2020). Syt7 KD not only eliminated

facilitation, but also the activity-dependent increase in the J_{ee}/J_{ie} ratio in the L2/3 network

(*Figure 8—figure supplement 1*). This suggests the possibility that Syt7 contributes to the

persistent activity during working memory tasks in the L2/3 recurrent network. Trace fear

conditioning (tFC) requires associative learning between an auditory cue (conditioned

stimulus, CS) and temporally separate aversive events (unconditioned stimulus, US). The

formation of trace fear memory requires the prelimbic area of the mPFC and hippocampus.

mPFC activity during the trace interval is thought to temporarily hold CS information, enabling

the network to associate temporally discontinuous events (i.e., temporal associative learning)

(Gilmartin et al., 2013). Synaptic facilitation and augmentation are implicated in temporary

memory retention in recurrent networks (Mongillo et al., 2008; Mongillo et al., 2012).

Given that Syt7 is crucial for synaptic facilitation (*Figure 5*) and augmentation (*Figure 6*) in

prelimbic L2/3 recurrent excitatory synapses, we tested whether trace fear memory

formation was impaired in rats with depleted Syt7 transcripts, specifically in the L2/3 PCs of

the mPFC. Syt7-targeted shRNA (shSyt7) or scrambled shRNA (Scr) constructs were

transfected bilaterally into the L2/3 PCs of the mPFC using IUE (*Figure 8A*). Both the Scr-

transfected control and shSyt7-transfected rats underwent tFC training (*Figure 8B*).

Consistent with Gilmartin et al. (2013), in which prefrontal persistent firing was

optogenetically inhibited, WT and KD animals exhibited similar freezing behavior during this

acquisition phase. The following day, the rats were subjected to a tone test in which trace

fear memory formation was assessed from the freezing response to a tone alone in a distinct

context. Compared to the control group, the shSyt7-transfected group exhibited significantly

lower levels of freezing behavior in response to auditory cues during the first four trials,

suggesting an impairment in trace memory formation (*Figure 8C*).

The tone test was repeated the following day to test for extinction memory formation. Syt7

KD animals exhibited much less freezing to the CS (+1d) than did WT animals. Considering the

Rescorla-Wagner model (Yau and McNally, 2023), our results suggest that the strength of CS-

US associative memory was weaker in KD animals than in WT animals; thus, the acquisition

of trace fear memory was impaired by Syt7 KD in the L2/3 PCs of the mPFC.

Next, we assessed the locomotor activity and anxiety levels of rats using the open field test

(OFT) and elevated plus maze test (EPM). Scr controls and Syt7 KD rats showed similar

exploratory patterns, displaying comparable total distance moved and time spent in the

periphery of the OFT or the open/closed arms of the EPM (*Figure 8—figure supplement 2*).

Previous studies have suggested that the acquisition of trace fear memory requires elevated

trace activity in prelimbic PCs (Gilmartin et al., 2013). Given that c-Fos, an immediate early

gene, is upregulated following patterned activities in neurons, its expression is considered a

reliable indicator of recent neuronal activation (Fields et al., 1997; Guzowski et al., 2005). To

investigate the effect of Syt7 on neuronal activity *in vivo* during tFC, animals were sacrificed

at the time of peak c-Fos protein expression, approximately 90 min after completion of the

behavioral task (Arime and Akiyama, 2017).

To elucidate the activation patterns of specific neuronal populations, we performed

immunohistochemical co-labeling of c-Fos and GAD67 (a GABAergic interneuron marker)

upon tFC acquisition. The density of c-Fos(+) cells and the percentage of c-Fos(+) cells among

GAD67(+) cells in all prelimbic layers were significantly lower in Syt7 KD rats than in control

rats (*Figure 8F-G*). Additionally, analysis of the proportion of active cells among transfected

L2/3 PCs (c-Fos+/GFP+) revealed that Syt7 KD L2/3 PCs were significantly more active than

Scr-transfected PCs (*Figure 8H*). This unexpected hyperactivity of Syt7 KD PCs may be

attributed to the high reciprocal connectivity between PCs and FSINs in the neocortex

(Holmgren et al., 2003; Otsuka and Kawaguchi, 2009), as PCs with short-term depression

would transmit less input to FSINs, which in turn would exert less feedback inhibition on

reciprocally connected PCs. Collectively, these findings show that the slow activation of

vesicle refilling supports various forms of facilitation at excitatory synapses in PFC L2/3, which

is essential for neuronal activity during temporal associative learning.

Discussion

The present study characterized STP at local excitatory synapses in the L2/3 network of the

rat mPFC at physiological extracellular $[Ca^{2+}]$. These synapses displayed initial strong

depression and then delayed facilitation at 40 Hz, while monotonous slowly developing

facilitation was observed at lower frequencies. Our study suggests that such delayed

facilitation after a brief depression results from a high p_v (*Figure 2 and Figure 3 and Figure*

3—figure supplement 2), along with Ca^{2+}-dependent delayed acceleration of vesicle refilling

and overfilling (*Figure 2 and Figure 3*). The two-fold increase in EPSC during 20 pulse train

stimulations under the condition of such a high p_v suggests a Ca^{2+}-dependent increase in the

vesicle replenishment rate during a train; this then leads to overfilling of release sites beyond

their incomplete basal occupancy (*Figure 3A-B*). The high p_v and strong PPD did not result

from artificial bouton stimulation, AMPAR desensitization, or any potential distortion caused

by optogenetic methods (*Figure 1—figure supplements 4 and 5*). Additionally, similar to

short-term facilitation, PTA could be explained by an increase in the TS vesicle occupancy

(*Figure 3C-E*), consistent with previous studies showing post-tetanic increases in the p_{occ}

and/or vesicle pool size at various types of synapses (Lee et al., 2010; Vandael et al., 2020;

Tran et al., 2023; Silva et al., 2024). These findings suggest that release sites are partially

occupied at rest and that a progressive overfilling of release sites with TS vesicles underlies

facilitation and augmentation at intracortical L2/3 excitatory synapses.

Definition of vesicular release probability in the present and previous studies

Previous studies for post-tetanic augmentation (Stevens and Wesseling, 1999; Garcia-Perez

and Wesseling, 2008) observed invariance of the RRP size after tetanic stimulation. In these

studies, the RRP size was estimated by hypertonic sucrose solution or as the sum of EPSCs

evoked 20 Hz/60 pulses train (denoted as 'RRP$_{hyper}$'). Because reluctant vesicles can be quickly

converted to TS vesicles (16/s) and are released during a train (Lee et al., 2012), it is likely that

the RRP size measured by these methods encompasses both LS and TS vesicles. In contrast,

we assert high p_v based on the observation of strong PPD, failure rates upon paired

stimulations at ISI of 20 ms (*Figure 2 and Figure 3—figure supplement 2*). Given that single

AP-induced vesicular release occurs from TS vesicles but not from LS vesicles, p_v in the present

study indicates the fusion probability of TS vesicles. From the same reasons, p_{occ} denotes the

occupancy of release sites by TS vesicles. Note that our study does not provide direct clue

whether release sites are occupied by LS vesicles that are not tapped by a single AP, although

an increase in the LS vesicle number may accelerate the recovery of TS vesicles. As suggested

in Neher (2024), even if the number of LS plus TS vesicles are kept constant, an increase in

p_{occ} (occupancy by TS vesicles) would be interpreted as an increase in 'vesicular release

probability' if it was measured based on RRP_{hyper}.

By the way, it should be noted that increases in both p_v and p_{occ} may contribute to PTA. Post-

tetanic potentiation (PTP) and phorbol esters had similar effects on baseline EPSCs

(Taschenberger et al., 2016), and PDBu increased both the TS vesicle pool size (1.9-fold) and

fusion probability of TS vesicles (1.3-fold) at the calyx synapses (Lin et al., 2025). Therefore, it

is possible that PTP may increase both p_{occ} and p_v of TS vesicles, although it has not been yet

directly tested. Moreover, acceleration of the latency in the hypertonicity-induced vesicle

release raises a possibility for an increase in fusion probability of TS vesicles (Stevens and

Wesseling, 1999; Garcia-Perez and Wesseling, 2008), although it is still controversial that the

acceleration of the release latency represents a reduction in the activation energy barrier for

vesicle fusion (Schotten et al., 2015).

Progressive overfilling of docking sites

Conventional models for facilitation assume an increase in p_v after an AP of RRP vesicles,

which had been in a low p_v or reluctant state at rest (Dittman et al., 2000; Turecek et al., 2016).

This facilitation model is unlikely to explain our results, because TS vesicles are predominantly

released not only at rest but also in the facilitated state during or after train stimulation, as

evidenced by the strong PPD (ISI, 25 ms) during a 5-Hz train (*Figure 2D-E*) and at different

intervals after a 40-Hz/30 pulse trains (*Figure 3D*). Moreover, the PPD (ISI, 25 ms) remained

pronounced during the early phase of facilitation (*Figure 7*), arguing against the possibility of

slowly increasing the p_v of reluctant vesicles. These results suggested that facilitation was

mediated by an activity-dependent progressive increase in occupancy of TS vesicles (TS

occupancy). Facilitation through an increase in TS occupancy has been termed frequency

facilitation, differentiating it from paired pulse facilitation that relies on a residual calcium-

dependent transient increase in the p_v of releasable vesicles (Neher, 2024).

Assuming an increase in TS occupancy, to attain a two-fold increase in EPSC (*Figure 1*), the

baseline occupancy of the docking sites by TS vesicles should be low, as estimated from the

V-M plot (c.a. 30% in *Figure 3*). However, it is unclear whether the remaining docking sites

that are not occupied by TS vesicles are truly vacant or are occupied by reluctant LS vesicles

that are not released by the AP or AP trains. TS vesicles can be supplied by the transformation

of LS to TS vesicles, both of which reside at the same docking site (LS/TS model; reviewed in

Neher, 2024). Alternatively, TS vesicles can be supplied from a distinct replacement pool

(RS/DS model; reviewed in Silva et al., 2021). Further studies, including ultrastructural

analyses of active zones, are required to address this issue.

Given the crucial role of Syt7 in facilitation (*Figure 5*), the facilitation mechanisms are closely

related to how Syt7 mediates synaptic facilitation. Based on the slow kinetics of Syt7, previous

computational modeling studies have proposed that Syt7 may mediate short-term facilitation

through an activity-dependent increase in p_v (Turecek et al., 2016; Jackman and Regehr, 2017;

Turecek et al., 2017; Norman et al., 2023). The present study, however, found that both

facilitation and augmentation were mediated by overfilling of the release sites (*Figure 2 and*

Figure 3). Moreover, Syt7-KD synapses displayed slower recovery of TS vesicles (*Figure 7*).

These results suggest that Syt7 is essential for activity-dependent overfilling of docking sites

with TS vesicles. Whereas the baseline EPSC was not altered (*Figure 1—figure supplement 3*),

and complementary changes in the number of docking sites and their baseline occupancy

were observed in Syt7 KD synapses (*Figure 6*). These results raise a possibility that Syt7 may

play a role in providing additional vacant docking sites that can be overfilled during facilitation.

It remains to be elucidated whether the decrease in the number of docking sites at Syt7 KD

synapses is related to the subcellular localization of Syt7 to the plasma membrane of the

active zone (Sugita et al., 2001; Vevea et al., 2021).

STP model in light of known Ca^{2+} binding kinetics of Syt7

Syt7 acts as an upstream Ca^{2+} sensor that facilitates activity-dependent replenishment (Liu et

al., 2014; Bacaj et al., 2015; Chen et al., 2017; Tawfik et al., 2021). The present study shows

that EPSCs are mediated by TS vesicles both at rest and during activity-dependent facilitation.

Releasable vesicles are homogeneous, and Syt7-dependent overfilling is responsible for the

activity-dependent enhancement of EPSCs. Thus, we tested whether slowly developing

facilitation could be replicated within the framework of a simple refilling model based on the

known Ca^{2+} binding kinetics of Syt7 (Brandt et al., 2012). To this end, we made following

assumptions: (1) a Ca^{2+}-dependent increase in the forward refilling rate (k_1) requires a full

Ca^{2+}-bound form of Syt7, to which multiple Ca^{2+} ions can bind (two or three Ca^{2+} ions on each

C2A and/or C2B domain) (Brandt et al., 2012; Chon et al., 2015; Voleti et al., 2017); (2)

sequential Ca^{2+} binding steps are synergistic through a cooperativity factor, *b*; (3) Syt7 senses

AP-induced local $[Ca^{2+}]$ similar to that estimated at the calyx of Held (a Gaussian function with

a full width at half maximum value of 0.2 ms and a peak of 40 μM) (Wang et al., 2008); and

(4) residual $[Ca^{2+}]$ follows a mono-exponential function with an amplitude of 1 μM and decay

time constant of 50 ms (Jackson and Redman, 2003).

Under these assumptions and the framework of the simple refilling model, slow and delayed

facilitation could be replicated by modeling k_1 as $k_1 = k_{1,b} + K_{1,max} \times [Syt7: n\ Ca^{2+}]$, where [Syt7:

n Ca^{2+}] represents the fraction of fully Ca^{2+}-bound Syt7 (*Figure 9A-C*). Syt7 was slowly

activated over the course of the stimulus train, suggesting that the cooperative binding of

multiple Ca^{2+} ions progressively increased the putative active form of Syt7 because of its slow

membrane binding/unbinding kinetics. Nevertheless, the late phase of facilitation at high

frequencies was predicted to be higher in the simulation than in the experiment. As expected,

based on the notion that sustained release during HFS is limited by site clearance in the active

zone (*Figure 4C*), the model/data discrepancy became more pronounced at higher

frequencies. Incorporating endocytic terms into the model may mitigate late-phase

discrepancies. Moreover, removal of the catalytic function of Syt7 in the model effectively

accounted for the complete loss of facilitation observed in Syt7 KD (*Figure 9D*).

Conventional STF models involving cumulative increase in p_v of reluctant vesicles

Conventional models for facilitation assume a post-AP residual Ca^{2+}-dependent step increase

in p_v of RRP (Dittman et al., 2000) or reluctant vesicles (Turecek et al., 2016). Given that p_v of

TS vesicles is close to one in the present study, an increase in p_v of TS vesicles cannot account

for facilitation. The possibility for activity-dependent increase in fusion probability of LS

vesicles (denoted as $p_{v,LS}$) should be considered in two ways depending on whether LS and TS

vesicles reside in distinct pools or in the same pool. Notably, strong PPD at short ISI implies

that $p_{v,LS}$ is near zero at the resting state. Whereas LS vesicles do not contribute to baseline

transmission, short-term facilitation (STF) may be mediated by cumulative increase in p_v of LS

vesicles that reside in a distinct pool. Because the increase in $p_{v,LS}$ during facilitation recruits

new release sites (increase in N), the variance of EPSCs should become larger as stimulation

frequency increases, resulting in upward deviation from a parabola in the V-M plane, as

shown in recent studies (Valera et al., 2012; Kobbersmed et al., 2020). This prediction is not

compatible with our results of V-M analysis (*Figure 3*), showing that EPSCs during STF fell on

the same parabola regardless of stimulation frequencies. Therefore, it is unlikely that an

increase in fusion probability of reluctant vesicles residing in a distinct release pool mediates

STF in the present study. For the latter case, in which LS and TS vesicles occupy in the same

release sites, it is hard to distinguish a step increase in fusion probability of LS vesicles from a

conversion of LS vesicles to TS. Nevertheless, our results do not support the possibility for a

gradual increase in $p_{v,LS}$ that occurs in parallel with STF. Strong PPD, indicative of high p_v, was

consistently found not only in the baseline (*Figure 2 and Figure 2—figure supplement 1*) but

also during post-tetanic augmentation phase (*Figure 3D*) and even during the early

development of facilitation (*Figure 2D-E and Figure 7*), arguing against the gradual increase

in $p_{v,LS}$. One may argue that STF may be mediated by a drastic step increase of $p_{v,LS}$ from zero

to one, but it is not distinguishable from conversion of LS to TS vesicles.

Possible mechanisms underlying the depression-facilitation sequence

Cooperative binding of multiple Ca^{2+} ions to Syt7 predicts delayed activation and a progressive

increase in vesicle refilling during train stimulations, resulting in delayed facilitation.

Combining this with a high p_v could replicate not only the slow facilitation at 5 Hz, but also

the unique depression-facilitation sequence observed at 40 Hz (*Figure 9*). As simulated by

Pulido and Marty (2018), delayed facilitation can be reproduced by a high p_v in combination

with a low baseline occupancy of replacement sites and subsequent Ca^{2+}-dependent refilling.

However, whether this model is also able to replicate slow facilitation at low frequencies

remains to be tested. An alternative, but not exclusive, possibility for delayed facilitation is

that the presynaptic global Ca^{2+} itself may undergo a slow buildup during a train due to the

saturation of endogenous Ca^{2+}-binding proteins.

Facilitation requires not only Syt7, but also activation of PLC-DAG signaling

To gain insight into the specific molecular mechanisms underlying the vesicle-loading

processes, we examined the effects of pharmacological inhibitors targeting PLC and DAG

(*Figure 4*). Both a PLC inhibitor and a DAG analog reduced facilitation and augmentation. This

suggests that the progressive overfilling of docking sites with TS vesicles is supported by the

PLC-DAG signaling pathway and probably downstream Munc13 (Rhee et al., 2002; Lou et al.,

2008). Phorbol esters have been recently shown to increase the TS fraction of docked vesicles

by enhancing vesicle priming without affecting Ca^{2+} transients or p_v (Taschenberger et al.,

2016; Aldahabi et al., 2022; Papantoniou et al., 2023; Aldahabi et al., 2024). In line with these

findings, signaling involving PLC and DAG may support Ca^{2+}-dependent overfilling of TS

vesicles by enhancing molecular priming processes.

Roles of facilitation and augmentation in working memory and related tasks

Previous theoretical studies have proposed that working memory in a recurrent network can

be maintained in the form of augmentation at synapses (Mongillo et al., 2008) and/or

persistent activity of neurons in a memory ensemble (Mongillo et al., 2012). The latter can be

supported by short-term facilitation of recurrent excitatory synapses or disparity in STP

between excitatory and inhibitory synapses (Mongillo et al., 2012; Yoon et al., 2020). However,

whether these short-term enhancements in synaptic efficacy contribute to working memory-

related behaviors has not been examined. tFC is a temporal associative learning task requiring

temporary retention of CS information during a trace period. The mPFC and hippocampus are

crucial for tFC, and tFC is interfered with by a load of working memory and attention

distraction, implying that the tFC shares the same neural substrate with other working

memory tasks (Raybuck and Lattal, 2014). In the present study, tFC in WT and Syt7 KD animals

revealed that Syt7 deficiency in the L2/3 PCs of the mPFC impaired the acquisition of trace

fear memory and reduced c-Fos expression associated with tFC training (*Figure 8*).

The acquisition of trace fear memory was impaired by inhibition of persistent activity in mPFC

during trace period (Gilmartin et al., 2013). The similar deficit observed in Syt7 KD animals is

consistent with the hypothesis that STF provides bi-stable ensemble activity in a recurrent

network (Mongillo et al., 2012). Nevertheless, alternative mechanisms may be responsible for

the behavioral deficit. Not only recurrent network but also long-range loop between the

mPFC and the mediodorsal (MD) thalamus play a critical role in maintaining persistent activity

within the mPFC especially for a delay period longer than 10 s (Bolkan et al., 2017). Prefrontal

L2/3 is heavily innervated by MD thalamus, and L2/3-PCs subsequently relay signals to L5

cortico-thalamic (CT) neurons (Collins et al., 2018). Given that L2/3 is an essential component

of the PFC-thalamic loop, loss of STF at recurrent synapses between L2/3 PCs may lead to

insufficient L2/3 inputs to L5 CT neurons and failure in the reverberant PFC-MD thalamic

feedback loop. Therefore, not only L2/3 recurrent network but also its output to downstream

network should be considered as a possible network mechanism underlying behavioral deficit

caused by Syt7 KD L2/3.

In summary, activity-dependent synaptic facilitation and augmentation at prelimbic L2/3

recurrent excitatory synapses can be attributed to overfilling of release sites with TS vesicles.

Given that the lack of short-term facilitation in L2/3 PCs impairs trace fear memory acquisition,

temporary maintenance of CS information during trace period may be supported by an

increase in the TS occupancy in presynaptic terminals of CS-representing neuronal ensembles.

Materials and Methods

Animals/Subjects

All experiments were carried out on Sprague-Dawley rats of either sex at p28-43. Rats were maintained in temperature-controlled rooms on a constant 12 h light/12 h dark cycle and were given access to water and food ad libitum. All animal procedures were approved and performed in accordance with the Institutional Animal Care and Use Committee at Seoul National University (SNU200522-1).

In utero electroporation (IUE)

Pregnant Sprague Dawley rats on embryonic day (E) 17.5 were deeply anesthetized with 5% (v/v) isoflurane for the duration of surgery. The uterine horns were exposed by laparotomy and wet with warm sterile phosphate-buffered saline (PBS). The plasmids (0.5-2 µg/µl) together with the 0.01% fast green dye were injected into either the left or the right lateral ventricle or bilateral ventricles of embryos through a thin glass capillary (Narishige PC-10). The embryo's head was carefully held between the forceps-shaped electrodes of 10 mm diameter (CUY65010; NEPAGENE, Japan). Electroporation pulses (50 V for 50 ms) were delivered five times at intervals of 150 ms using a square-wave generator (ECM830; BEX, United States). The uterine horns were placed back into the abdominal cavity and abdominal muscles and skin were sutured. After the surgery, the animals were recovered under the infrared lamp. For knockdown experiments in slice electrophysiology, the shRNA constructs were co-electroporated with the pCAG-oChIEF-tdTomato, because of the consistent co-

expression of the two constructs in a significant proportion of the transfected cells in vivo

(Bony et al., 2013). Pups with robust fluorescence signals in the prefrontal cortex of both

hemispheres were used for behavior experiments.

Preparation of vector constructs

For an optic stimulation, pCAG-oChIEF-tdTomato was transfected by IUE. Short hairpin RNA

(shRNA) against Syt7 sequence (GATCTACCTGTCCTGGAAGAG) was expressed under the

control of an U6 promoter of pAAV-U6-GFP vector (Cell Biolabs, #VPK-413). This shRNA was

shown to effectively knockdown Syt7 in cell cultures and in vivo by previous reports from

more than one research group (Bacaj et al., 2013; Li et al., 2017). Scrambled control sequence

(TCGCATAGCGTATGCCGTT) was designed by shuffling the recognition region sequences and

a BLAST analysis confirmed that the control shRNA sequence had no target gene. For rescue

experiments, the cDNA for rat Syt7 (NM_021659) rendered resistant to the shRNA were

inserted in the KD vector and their expression was driven by the CAG promoter; the vector

also contained a self-cleaving P2A peptide sequence followed by EGFP, which ensures

visualization of transfected cells. All the constructs above were verified by DNA sequencing.

Plasmid DNA was purified and concentrated under endotoxin free conditions (Qiagen

EndoFree Maxi Kit).

Slice preparation

Acute brain slices were prepared by using a VT1200S Vibratome (Leica Microsystems) after

isoflurane anesthesia and decapitation. Coronal sections of the mPFC (300 μm in thickness)

were cut in ice-cold artificial cerebrospinal fluid (aCSF) composed of (in mM) 125 NaCl, 3.2

KCl, 25 $NaHCO_3$, 1.25 NaH_2PO_4, 20 D-glucose, 2 NaPyr, 1 $MgCl_2$, and 1.3 $CaCl_2$ (Yoon et al.,

2020), bubbled with 95% O_2 and CO_2 (pH = 7.3; 310 mOsm). Slices were allowed to recover

for a minimum of 30 minutes at 36 °C, and subsequently maintained at room temperature

until recording. Composition of aCSF was identical throughout the preparation, incubation,

and recording periods unless otherwise stated. For the recordings, the mPFC slices were

continuously perfused with oxygenated aCSF at a flow rate of 1.5 ml/min, and the

temperature was maintained at 31–32 °C using an in-line temperature controller (Warner

Instruments).

Electrophysiology

Whole-cell patch-clamp recordings were performed from pyramidal cells and fast-spiking

interneurons in layer 2/3 of the prelimbic mPFC. Neuronal subtypes were initially selected

based on morphology under visual guidance by using an upright microscope equipped with

infrared differential interference contrast (IR-DIC) optics (BX51WI, Olympus), and were

further identified by their electrophysiological properties (see *Figure 1—figure supplement*

1; van Aerde and Feldmeyer, 2015; Tremblay et al., 2016). To analyze these intrinsic

properties in current-clamp experiments, we measured the following parameters: (1) resting

membrane potential (RMP), (2) input resistance (R_{in}), (3) sag ratio, and (4) the F-I curve, which

plots firing frequencies (F) against the amplitude of injected currents (I). Both cell types

exhibited a low sag ratio below 0.1, consistent with findings from other studies (van Aerde

and Feldmeyer, 2015; Yoon et al., 2020). PCs and FSINs were easily distinguished by their

unique firing frequency, Rin, spike adaptation, RMP and morphological characteristics

including the shape of soma and the thickness of apical dendrites. For experiments involving

oChIEF-expressing neurons in WT or Syt7 KD rats, whole-cell recordings were performed from

the fluorescently labeled pyramidal neurons in layer 2/3 of the prelimbic cortex using suitable

filter sets. Patch pipettes (tip resistance of 2–3 MΩ) were pulled from borosilicate glass and

filled with (in mM) 130 K-gluconate, 8 KCl, 10 HEPES, 0.2 EGTA, 20 Na_2Pcr, 0.3 Na_2GTP, 4

MgATP. Intracellular solutions were adjusted to pH 7.25 and 300–310 mOsm. Neurons were

voltage clamped at −78 mV and series resistance (Rs) were continuously monitored during

the EPSCs recordings. Experiments were discarded before analysis if the Rs changed by 20%

or larger deviation of baseline value. Measurements were not corrected for Rs compensation,

bridge balance, or liquid junction potential. Electrophysiological data were low-pass-filtered

at 1 kHz (Bessel) and acquired at 20 kHz using a Multiclamp 700B amplifier paired with

Digidata 1440A digitizer and Clampex 10.2 software (Molecular Devices).

Synaptic stimulation

IUE resulted in sparse expression of channelrhodopsin (10-20%) (*Figure 1A*). Exploiting the

sparse expression, we employed a minimal optical stimulation method to activate a single or

very few excitatory synapses in layer 2/3. The optic minimal stimulation was performed by

our published protocol with minor modifications (Yoon et al., 2020). This approach aimed to

minimize the risk of synaptic contamination, which could arise from engaging extra terminals

that were initially inactive (depolarized below threshold) but became active by temporal

summation during high-frequency trains. It also helped prevent the buildup of depolarization

at the synaptic terminals, which might influence the probability of release, as noted by

(Jackman et al., 2014). To achieve this, we employed a collimated digital micromirror device

(DMD)-coupled LED (Polygon400; Mightex Systems) to confine 470 nm blue light to a small

area with a radius of approximately 3–8 µm (typically 3–4 µm), as measured at the focal plane

of a 60× water immersion objective [numerical aperture (NA) = 1.0; LUMPlanFL, Olympus]

(*Figure 1—figure supplement 3A*). To find the minimal stimulation area, the radius of

illumination area was increased from 2 µm by 50% at each step (*Figure 1—figure supplement*

3A), and selected the smallest illumination area that elicited EPSCs. Setting the duration of

illumination between 3 and 5 ms (typically 5 ms), photostimulation onto oChIEF-expressing

cells reliably induced action potentials (APs) during 600-pulse trains (see *Figure 1—figure*

supplement 2).

STP at various stimulus frequencies (5, 10, 20, and 40 Hz) was examined using 20-pulse trains.

Trials of train stimulation were repeated with a prolonged interval of 30 s to avoid change in

baseline properties. We observed no systematic changes in the amplitudes of the first EPSC

of each train, nor did we find significant differences in the paired-pulse ratio (PPR) across

stimulation trials in the same cell. For each stimulus frequency, approximately ten to twenty

consecutive traces were collected and then averaged across trials to yield a mean EPSC trace.

For recovery experiments, two consecutive trains of 3 or 30 stimuli at 40 Hz, with varying time

intervals (0.1-10 s), were applied every 10 s or 60 s, respectively, at both excitatory synapses.

For 600-stimulus trains at 10Hz, the trial was delivered only once for each cell to avoid

possible after-effects caused by prolonged synaptic stimulation (Hirsch and Crepel, 1990).

Prior to the stimulus trains, the EPSCs were normalized to the mean amplitude of the baseline.

For the baseline measurements, paired pulses with various ISIs (20-200 ms) were delivered

every 3 s (1/3 Hz) at least 20 times.

For all recordings, picrotoxin (PTX, 10 µM) was included in the recording solution to isolate

glutamatergic responses. mPFC PC terminals were entirely glutamatergic as the application

of 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX, 20 µM), an AMPA receptor antagonist,

completely abolished the monosynaptic events evoked by photostimulation. A low

concentration of PTX was employed to minimize the relieving effects on spike suppression

caused by blockade of GABAa receptors. We confirmed that no synaptic response was elicited

by electrical stimulation in the presence of both 10 µM PTX and 20 µM CNQX.

In *Figure 1—figure supplement 4A*, 1 µM TTX (1 mM stock solution) and 0.1 mM 4-AP (500

mM stock solution) were added to the bath solution to block action potentials and restore

presynaptic glutamate release, respectively (*Figure 1—figure supplement 4*). In *Figure 1—*

figure supplement 5A, 50 µM of cyclothiazide, a positive allosteric modulator of the AMPA

receptors, was included to minimize rapid desensitization of the postsynaptic receptors. In

Figure 1—figure supplement 5B, 0.5 mM kynurenate was added to bath solution to prevent

possible saturation of AMPA receptors, especially during maximal synaptic facilitation. For

pharmacological manipulations of synaptic vesicle dynamics (*Figure 4*), drugs related to

vesicle supply including U73122, OAG, LatB, EGTA-AM, and dynasore at indicated

concentrations were added to the recording solution during baseline measurements or

PTA/recovery experiments (IBI = 0.5 s). Chemicals were from Sigma-Aldrich, Merck, or Tocris.

Measurements of quantal size

Experiments for measuring asynchronous release were performed as previously described

(Yoon et al., 2020). In brief, we replaced extracellular Ca^{2+} entirely with 2 mM Sr^{2+} to induce

asynchronous release. A single stimulus was delivered at 3-second intervals over 100

repetitions, and asynchronous release events were detected during a 100-ms window starting

25 ms after stimulus onset for each trial. The same procedure was also executed under sham

conditions (0% light intensity) to serve as a control, thus mitigating contamination from

nonspecific spontaneous events and potential errors associated with in the detection

algorithm. q was determined as the mean value from the first Gaussian function after fitting

the EPSC distribution with either a single or double Gaussian function (*Figure 3—figure*

supplement 1).

Estimation of Ca^{2+} kinetics at axonal boutons

Cytosolic [Ca^{2+}] at single axonal boutons was estimated by employing the two-dye, dual-

excitation wavelength technique as described by (Oheim et al., 1998). This method involves

measuring the fluorescence ratio of a Ca^{2+}-sensitive dye to that of a Ca^{2+}-insensitive dye to

correct for dye concentrations. We used Fluo-5F at concentrations of 150, 250, or 500 µM

(dissociation constant, K_d = 2.3 µM) as the calcium indicator dye and Alexa Flour 555 (50 µM)

as the reference dye. The fluorophores were from Invitrogen. Both dyes were co-loaded into

presynaptic terminals of layer 2/3 pyramidal cells in mPFC through whole-cell patch clamp.

Before data acquisition, the dyes were allowed to diffuse to terminal boutons for at least 30

minutes after the patch break-in, a duration sufficient for the equilibrium of dye

concentrations between the soma and its axonal terminals.

Calcium imaging was performed by sequentially exciting Fluo-5F and AF555 at wavelengths

of 473 and 559 nm, respectively, using a confocal laser-scanning microscope (FV1200;

Olympus, Japan) equipped with a 60× water immersion objective (NA, 0.9; LUMPlanFl;

Olympus, Tokyo, Japan). Once a single axonal bouton was identified, lines for scanning were

drawn across boutons, oriented perpendicular to the axon (*Figure 2—figure supplement 2A-*

B). After baseline fluorescence was measured, line scanning was proceeded while a single or

train APs were induced by applying current pulses. Upon stimulation, green fluorescence but

not red one increased along the vertical array of lines across the bouton (*Figure 2—figure*

supplement 2B-C). Measurements were obtained under current-clamp conditions, with the

resting potential regularly monitored. The scanning speed was adjusted to 600-700 Hz by

controlling the line length. Line scan for measuring fluorescent transients in a bouton was

repeated at least 3 times, and then averaged to enhance the signal-to-noise ratio. To mitigate

photobleaching, minimal intensity of the laser power and maximal pinhole size were used.

Emitted light was filtered through 490/590 or 575/675 bandpass filters before detection by

photomultiplier tubes for green or red fluorescence, respectively.

Fluorescence measurements were converted to $[Ca^{2+}]_i$ using the ratio of background-

subtracted fluorescence ($F = F_0 - F_b$) from two dyes. The fluorescence ratio ($R = F_{green}/F_{red}$) in

data traces were converted to $[Ca^{2+}]_i$ using the following equation:

$$[Ca^{2+}]_i = K_d \times (R-R_{min}) / (R_{max}-R). \tag{1}$$

Calibration parameters were obtained using an in-cell calibration protocol (Helmchen, 2011),

wherein the minimum (R_{min}) and maximum ratio (R_{max}) values were established through the

use of intracellular solutions containing 10 mM EGTA or 10 mM $[Ca^{2+}]$, respectively.

When Ca^{2+} is extruded with a rate constant, γ, after an AP-induced short pulse of Ca^{2+} influx

to a single compartment, $[Ca^{2+}]_i$ initially rises to A_{Ca} and then undergoes a mono-exponential

decay with a time constant (τ_{Ca}). Under this framework, A_{Ca} and τ_{Ca} depend on Ca^{2+} binding

ratios of endo- and exogenous Ca^{2+} buffers (denoted as κ_S and κ_B, respectively) as following

equations (Neher, 1995; Kim et al., 2003):

$$1/A_{Ca} = (1 + \kappa_S + \kappa_B) / \Delta[Ca^{2+}]_T. \tag{2}$$

$$\tau_{Ca} = (1 + \kappa_S + \kappa_B) / \gamma. \tag{3}$$

, where $\Delta[Ca^{2+}]_T$ is an increment in total $[Ca^{2+}]$ (free plus bound) elicited by an AP. Ca^{2+} binding

ratio indicates the ratio of concentration changes in the Ca^{2+}-buffer complex relative to free

$[Ca^{2+}]$ change, and linearized κ_B is calculated from the dye concentration and the change from

$[Ca^{2+}]_1$ to $[Ca^{2+}]_2$ as follows:

$$\kappa_B = \Delta[CaB] / \Delta[Ca^{2+}] = [B]_t K_d / [([Ca^{2+}]_1+K_d) ([Ca^{2+}]_2+K_d)]. \tag{4}$$

The total dye concentration, $[B]_t$, was taken as [Fluo-5F] in the patch pipette. With κ_B

determined from Equation 4 and τ_{Ca} from the decay of a Ca^{2+} transient, κ_S can be estimated

by plotting τ versus κ_B and applying a linear fit to the plot, which gives the x-intercept as $-(1 +$

κ_S). Once determined, κ_S can be used to calculate $\Delta[Ca^{2+}]_T$.

Optimization of k_1 and p_v from failure rates

Let n_0 = the number of docked SVs before arrival of a first AP; n_1 = the number of docked SVs

remaining after a first AP (i.e. $n_0 - n_1$ vesicles are released); n_2 = the number of docked SVs

just before arrival of a 2nd AP (ISI = 20 ms; *Figure 3—figure supplement 2*). If the number of

release sites (N), forward and reverse rates for vesicle docking (k_1 and b_1, respectively), mean

release site occupancy (p_{occ}) and vesicular fusion probability upon an AP (p_v) are given, we can

predict $P(n_0, n_1, n_2)$, the probability for $[n_0, n_1, n_2]$, as following steps.

1) Probability for a synapse harbouring n_0 docked SVs just before arrival of a first AP is

calculated as:

$$P(n_0) = C(N, n_1)\ p_{occ}^{n0} \times (1 - p_{occ})^{(N-n0)}.$$

2) Probability that n_1 vesicles remain after a first AP when the initial number of docked SVs is

n_0 $[P(n_1|n_0)]$ is calculated as:

$$P(n_1|n_0) = C(n_0, n_1) \times (1-p_v)^{n1} \times p_v^{(n0-n1)},\ \text{if } n_0 \geq n_1\ ,$$

$$P(n_1|n_0) = 0,\ \text{if } n_0 < n_1.$$

3) When forward and backward rate constants are k_1 and b_1, respectively, probability that n_2

vesicles are docked just before arrival of a 2nd AP when n_1 SVs remain after a 1st AP ($\Delta t = 20$

ms) $[P(n_2 | n_1)]$ is predicted as follows: Assuming that docking and undocking follow a Poisson

process,

$$P(n_2 \mid n_1) = C(N - n_1, n_2 - n_1) [1 - \exp(-k_1 \, \Delta t)]^{(n_2 - n_1)}, \text{ if } n_2 > n_1 .$$

$$P(n_2 \mid n_1) = C(n_1, n_1 - n_2) [1 - \exp(-b_1 \, \Delta t)]^{(n_1 - n_2)}, \text{ if } n_2 < n_1 .$$

$$P(n_2 \mid n_1) = \exp[-n_1 k_1 \, \Delta t] \exp[- n_1 b_1 \, \Delta t], \text{ if } n_2 = n_1 .$$

4) Combining probabilities calculated from above three steps, we can calculate $P(n_0, n_1, n_2)$:

$$P(n_0, n_1, n_2) = P(n_0) \times P(n_1 \mid n_0) \times P(n_2 \mid n_1)$$

5) Once $P(n_0, n_1, n_2)$ is given, we can predict probability for double failure (P_{00}) and probability

for the 1^{st} success and the 2^{nd} failure (P_{10}) as:

$$P_{00} = \Sigma_{n0=0..N} \, \Sigma_{n1 = n0} \, \Sigma_{n2 = 0..N} \quad p(n_0, n_1, n_2) (1 - p_v)^{n_2} .$$

$$P_{10} = \Sigma_{n0=0..N} \, \Sigma_{n1 \neq n0} \, \Sigma_{n2 = 0..N} \quad p(n_0, n_1, n_2) (1 - p_v)^{n_2} .$$

6) Probability for the 1^{st} failure and the 2^{nd} success (P_{01}) and probability for double success

(P_{11}) were calculated as:

$$P_{01} = P(F_1) - P(F_1, F_2)$$

$$P_{11} = P(S_1) - P(S_1, F_2)$$

, in which $P(F_1)$ and $P(S_1)$ are probability for the 1^{st} failure and the 1^{st} success, respectively,

and these are calculated as:

$$P(F_1) = P(n_0) P(F_1 \mid n_0) = \Sigma_{n0=0..N} P(n_0) (1 - p_v)^{n_0} ,$$

$$P(S_1) = 1 - P(F_1).$$

To optimize k_1 and p_v, we used the Matlab routine, *fmincon*, in which the sum of squared

errors (= observed value minus predicted value) was set as a cost function to be minimized.

Lower and upper bounds for k_1 and p_v were set as [4, 8] and [0.8, 1], respectively. Release

probability (p_r) was calculated from the observed 1^{st} failure rate, $P(F_1)$, as

$$p_r = 1 - [P(F_1)]^{(1/N)}$$

N was assumed to be six as shown in *Figure 3*. Other parameters (b_1, and p_{occ}) were set

according to following relationships:

895                       $$p_{occ} = p_r / p_v .$$

896                       $$b_1 = k_1 (1 - p_{occ})/p_{occ} .$$

898   *Calculation of probability for double failure*

899   Double failure probability, P_{00}, can be more easily predicted. By the Bayes rule,

900               $$P(F_1, F_2, n_1) = P(F_2 \mid n_1, F_1) \, P(F_1 \mid n_1) \, P(n_1).$$

901   , where F_1 and F_2 means 1st and 2nd failure. Moreover, and $n_0 = n_1$ because of the 1st failure.

902   Thus, the probability for two consecutive failures can be calculated as:

903            $$P(F_1, F_2) = \Sigma_{n1=0..N} \, P(F_2 \mid n_1, F_1) \, P(F_1 \mid n_1) \, P(n_1).$$

$P(F_1)$ and N are observed values. For calculation in *Figure 3—figure supplement 2*, k_1 and p_v

were set free variables. Other parameters were set under the following constrains: $P(F_1) = (1$

$- p_r)^N$, $p_r = p_v \, p_{occ}$, and $p_{occ} = k_1 / (k_1 + b_1)$.

Behavioral tests

Trace fear conditioning: For trace fear conditioning (tFC) (Gilmartin et al., 2013), individual

rats were placed in the training chamber (30.5 x 25.4 x 30.5 cm; Coulbourn Instruments)

consisting of metal grid floor connected to an electrical stimulator (H10-11R-TC-SF, Coulbourn

Instruments) for delivery of a footshock. After a 5 min habituation period, rats received five

pairings of a 20 s tone conditioned stimulus (CS, 2.5 kHz, 75 dB), followed by empty 20s trace

period and a unconditioned stimulus (US; 1 s, 0.7 mA footshock) with a pseudorandom inter-

trial interval (ITI) of 240 ± 20 s. During this training session, rats learned to associate the

auditory CS with the shock US. The next day, the rats were tested for memory of CS-US trace

association (called tone test). The acquisition of fear learning, often indicated by the extent

of freezing to the CS, was assessed in a novel chamber (a white hexagonal enclosure) in a

dimly lit room. During this tone (or retrieval) test, rats were first allowed to explore the new

context for 5 min (habituation), followed by twelve 20 s CS presentations with a variable ITI

of 120 ± 20 s. This entire session was repeated on the next day for assessment of extinction

learning. Identities of the chambers were also determined with the presence of distinct odors;

the training and testing chambers were cleaned with 70% ethanol and 1% acetic acid

respectively before and after each session. For all experimental sessions, the activity of

animals in the chambers was recorded at 30 frames per second using the EthoVision XT

(Noldus Information Technology, Wageningen, Netherlands), and stored as a video file. The

freezing behavior, defined as behavioral immobility other than respiratory movement, was

assessed using an open-source video analysis pipeline ezTrack (Pennington et al., 2019). The

percentage of freezing was calculated as a total duration of freezing divided by the total

duration of observation.

Elevated Plus Maze: Elevated plus maze (EPM) test was performed using a plus-shaped

apparatus elevated 60 cm above the ground. The maze consisted of two open arms (50 x 10

x 0.5 cm (H)), two closed arms (50 x 10 x 40 cm (H)) and a center area (10 x 10 cm). Rats were

individually placed on the center area, facing an open arm, and the path of the animal was

recorded with a video camera for 5 min. We analyzed the total distance traveled for 5 min

and time spent in each arm. Animals were tracked offline using the open-source tracking

system ezTrack (Pennington et al., 2019).

Open field test: Open field exploration test (OFT) was performed in an open field apparatus

in a dimly lit room. The open field consisted of a white plastic board (1.20 m in diameter)

surrounded by white plastic walls (50 cm in height). Individual rats were placed in a center

zone (0.6 m in diameter), and allowed to freely explore the apparatus for 10 min. Time spent

in the center zone and total distance traveled were analyzed for initial 5 min. Animals were

tracked offline using the open-source tracking system ezTrack (Pennington et al., 2019).

Histology and immunohistochemistry

At the end of the behavioral experiments, rats were anesthetized using isoflurane inhalation

and transcardially perfused with PBS followed by 4% paraformaldehyde (PFA, T&I, Korea).

Brains were extracted and post-fixed in 4% PFA solution for 24 h. Perfused brains were sliced

at 100 μm for estimation of GFP expression or 50 μm for immunohistochemistry. For analysis

of c-Fos expression, rats were returned to their home cages immediately after the training

session and perfused approximately 90 minutes later (Arime and Akiyama, 2017). Brains were

then cryopreserved, frozen, and sectioned coronally. Immunofluorescence staining was

conducted on free-floating slices using a rabbit monoclonal anti c-fos (1:1000, Cell Signaling

Technology, cat# 2250s) and mouse monoclonal anti-GAD67 (1:500, Millipore, cat# mab5406).

Fluorescent secondary antibodies included Cy5-conjugated goat anti-rabbit IgG (1:1000,

Abcam, cat# ab97077) and Cy3-conjugated goat anti-mouse IgG (1:1000, Abcam, cat#

ab97035). Brain slices were permeabilized and blocked with PBS containing 0.5% Triton X-100

and 5% normal goat serum (NGS) for 1h at room temperature. Slices were incubated

overnight at 4 °C with primary antibodies, followed by 4 h incubation with secondary

antibodies at room temperature. Slices were then mounted with Antifading Mounting

medium (Abcam, cat# ab104139). Confocal images were acquired using a Leica TCS SP8

microscope set to the same laser intensity and acquisition parameters to compare the

immunosignals in sections from Scr and Syt7 KD groups. For cell counting and colocalization,

fluorescent protein-positive cells were automatically quantified with spot-detection

algorithms in Imaris 9.5 (Oxford instruments).

Data analysis

Data were analyzed and presented using ClampFit (Molecular Devices), Igor Pro

(Wavemetrics), MATLAB (Mathworks), and Prism (GraphPad). Data points and error bars

represent the mean ± standard error of the mean. All statistical analyses were conducted

using two-tailed comparisons. Sample sizes and statistical tests for each group are detailed in

the figure legends.

Numerical integration of the simple refilling model

This model posits reversible docking of vesicles at finite release sites (N_{max}) with forward (k_1)

and reverse rate constants (b_1) (Hosoi et al., 2007; Neher and Sakaba, 2008).

$$\overset{k_1}{\underset{b_1}{\rightleftharpoons}} n \overset{p_v}{\rightarrow}$$

According to this scheme, $dn/dt = k_1 u - b_1 n - p_v n \delta(t - t_{AP})$, where u is the number of

unoccupied sites (= $N_{max} - n$); t_{AP} is the timing of AP firing; and δ is the Dirac delta function.

The p_v was assumed to be unity (*Figure 3—figure supplement 2*). The sum of basal k_1 (= $k_{1,b}$)

and b_1 was be estimated as 23/s from the dependence PPR on ISIs (*Figure 3—figure*

supplement 2Ab). From the baseline occupancy of 0.3 and $p_{occ} = k_{1,b} / (k_{1,b} + b_1)$, we could

estimate baseline k_1 and b_1 as 6.9/s, and 16.1/s, respectively. Ca^{2+}- and Syt7-dependent

increase of k_1 was modeled as $k_1 = k_{1,b} + K_{1,max} [Syt7: n Ca^{2+}]$, where $[Syt7: n Ca^{2+}]$ represents

the fraction of full Ca^{2+}-bound Syt7. For deterministic simulations, we time-integrated a set

of differential equations for each model using Euler methods with a time step of 1 ms. All

calculations were performed on the platform of Matlab (R2022b, Mathworks, USA).

Acknowledgements

We thank Dr. E. Neher and Dr. A. Marty for critical reading of this manuscript and

invaluable comments.

Author contributions

All experiments were carried out at cell physiology lab under Y.K and S.H.L's supervision.

Conception or design of the work: J.S. and S.H.L. Acquisition, analysis or interpretation of data

for the work: J.S, Y.K. and L.S.H. Drafting the work or revising it critically for important

intellectual content: J.S., S.Y.L, Y.K. and L.S.H.

Funding

This study was supported by grants from the National Research Foundation of Korea (RS-

2024-333669 to S-H. Lee; 2021R1I1A1A01059646 to Y. Kim), and Seoul National University

Hospital (2024).

1003

1004

References

Aldahabi M, Neher E, Nusser Z (2024) Different states of synaptic vesicle priming explain target cell type-dependent differences in neurotransmitter release. Proc Natl Acad Sci U S A 121:e2322550121.

Aldahabi M, Balint F, Holderith N, Lorincz A, Reva M, Nusser Z (2022) Different priming states of synaptic vesicles underlie distinct release probabilities at hippocampal excitatory synapses. Neuron 110:4144-4161 e4147.

Arime Y, Akiyama K (2017) Abnormal neural activation patterns underlying working memory impairment in chronic phencyclidine-treated mice. PLoS One 12:e0189287.

Bacaj T, Wu D, Burre J, Malenka RC, Liu X, Sudhof TC (2015) Synaptotagmin-1 and -7 Are Redundantly Essential for Maintaining the Capacity of the Readily-Releasable Pool of Synaptic Vesicles. PLoS Biol 13:e1002267.

Bacaj T, Wu D, Yang X, Morishita W, Zhou P, Xu W, Malenka RC, Sudhof TC (2013) Synaptotagmin-1 and synaptotagmin-7 trigger synchronous and asynchronous phases of neurotransmitter release. Neuron 80:947-959.

Baeg EH, Kim YB, Huh K, Mook-Jung I, Kim HT, Jung MW (2003) Dynamics of population code for working memory in the prefrontal cortex. Neuron 40:177-188.

Bolkan SS, Stujenske JM, Parnaudeau S, Spellman TJ, Rauffenbart C, Abbas AI, Harris AZ, Gordon JA, Kellendonk C (2017) Thalamic projections sustain prefrontal activity during working memory maintenance. Nat Neurosci 20:987-996.

Bony G, Szczurkowska J, Tamagno I, Shelly M, Contestabile A, Cancedda L (2013) Non-hyperpolarizing GABAB receptor activation regulates neuronal migration and neurite growth and specification by cAMP/LKB1. Nat Commun 4:1800.

Brandt DS, Coffman MD, Falke JJ, Knight JD (2012) Hydrophobic contributions to the membrane docking of synaptotagmin 7 C2A domain: mechanistic contrast between isoforms 1 and 7. Biochemistry 51:7654-7664.

Chen C, Satterfield R, Young SM, Jr., Jonas P (2017) Triple Function of Synaptotagmin 7 Ensures Efficiency of High-Frequency Transmission at Central GABAergic Synapses. Cell Rep 21:2082-2089.

Chon NL, Osterberg JR, Henderson J, Khan HM, Reuter N, Knight JD, Lin H (2015) Membrane Docking of the Synaptotagmin 7 C2A Domain: Computation Reveals Interplay between Electrostatic and Hydrophobic Contributions. Biochemistry 54:5696-5711.

Collins DP, Anastasiades PG, Marlin JJ, Carter AG (2018) Reciprocal Circuits Linking the Prefrontal Cortex with Dorsal and Ventral Thalamic Nuclei. Neuron 98:366-379 e364.

Dittman JS, Kreitzer AC, Regehr WG (2000) Interplay between facilitation, depression, and residual calcium at three presynaptic terminals. J Neurosci 20:1374-1385.

Dobrunz LE, Huang EP, Stevens CF (1997) Very short-term plasticity in hippocampal synapses. Proceedings of the national academy of sciences 94:14843-14847.

Dodge Jr FA, Rahamimoff R (1967) Co-operative action of calcium ions in transmitter release at the neuromuscular junction. The Journal of physiology 193:419-432.

Fields RD, Eshete F, Stevens B, Itoh K (1997) Action potential-dependent regulation of gene
expression: temporal specificity in ca2+, cAMP-responsive element binding proteins,
and mitogen-activated protein kinase signaling. J Neurosci 17:7252-7266.
Fujisawa S, Amarasingham A, Harrison MT, Buzsaki G (2008) Behavior-dependent short-term
assembly dynamics in the medial prefrontal cortex. Nat Neurosci 11:823-833.
Garcia-Perez E, Wesseling JF (2008) Augmentation controls the fast rebound from depression
at excitatory hippocampal synapses. J Neurophysiol 99:1770-1786.
Gilmartin MR, Miyawaki H, Helmstetter FJ, Diba K (2013) Prefrontal activity links
nonoverlapping events in memory. J Neurosci 33:10910-10914.
Guzowski JF, Timlin JA, Roysam B, McNaughton BL, Worley PF, Barnes CA (2005) Mapping
behaviorally relevant neural circuits with immediate-early gene expression. Curr Opin
Neurobiol 15:599-606.
Hallermann S, Silver RA (2013) Sustaining rapid vesicular release at active zones: potential
roles for vesicle tethering. Trends Neurosci 36:185-194.
Hansel D, Mato G (2013) Short-term plasticity explains irregular persistent activity in working
memory tasks. J Neurosci 33:133-149.
Hass CA, Glickfeld LL (2016) High-fidelity optical excitation of cortico-cortical projections at
physiological frequencies. J Neurophysiol 116:2056-2066.
Haucke V, Neher E, Sigrist SJ (2011) Protein scaffolds in the coupling of synaptic exocytosis
and endocytosis. Nat Rev Neurosci 12:127-138.
Helmchen F (2011) Calibration protocols for fluorescent calcium indicators. Cold Spring Harb
Protoc 2011:980-984.
Hempel CM, Hartman KH, Wang XJ, Turrigiano GG, Nelson SB (2000) Multiple forms of short-
term plasticity at excitatory synapses in rat medial prefrontal cortex. J Neurophysiol
83:3031-3041.
Hirsch JC, Crepel F (1990) Use-dependent changes in synaptic efficacy in rat prefrontal
neurons in vitro. J Physiol 427:31-49.
Holler S, Kostinger G, Martin KAC, Schuhknecht GFP, Stratford KJ (2021) Structure and
function of a neocortical synapse. Nature 591:111-116.
Holmgren C, Harkany T, Svennenfors B, Zilberter Y (2003) Pyramidal cell communication
within local networks in layer 2/3 of rat neocortex. J Physiol 551:139-153.
Hosoi N, Sakaba T, Neher E (2007) Quantitative analysis of calcium-dependent vesicle
recruitment and its functional role at the calyx of Held synapse. J Neurosci 27:14286-
14298.
Jackman SL, Regehr WG (2017) The Mechanisms and Functions of Synaptic Facilitation.
Neuron 94:447-464.
Jackman SL, Beneduce BM, Drew IR, Regehr WG (2014) Achieving high-frequency optical
control of synaptic transmission. J Neurosci 34:7704-7714.
Jackman SL, Turecek J, Belinsky JE, Regehr WG (2016) The calcium sensor synaptotagmin 7 is
required for synaptic facilitation. Nature 529:88-91.
Jackson MB, Redman SJ (2003) Calcium dynamics, buffering, and buffer saturation in the
boutons of dentate granule-cell axons in the hilus. J Neurosci 23:1612-1621.

Kim MH, Lee SH, Park KH, Ho WK, Lee SH (2003) Distribution of K+-dependent Na+/Ca2+
exchangers in the rat supraoptic magnocellular neuron is polarized to axon terminals.
J Neurosci 23:11673-11680.

Kobbersmed JRL, Grasskamp AT, Jusyte M, Böhme MA, Ditlevsen S, Sørensen JB, Walter AM
(2020) Rapid regulation of vesicle priming explains synaptic facilitation despite
heterogeneous vesicle:Ca2+ channel distances. eLife 9:e51032.

Kusick GF, Chin M, Raychaudhuri S, Lippmann K, Adula KP, Hujber EJ, Vu T, Davis MW,
Jorgensen EM, Watanabe S (2020) Synaptic vesicles transiently dock to refill release
sites. Nat Neurosci 23:1329-1338.

Lee HR, Choi SH, Lee SH (2024) Differential involvement of mitochondria in post-tetanic
potentiation at intracortical excitatory synapses of the medial prefrontal cortex. Cereb
Cortex 34.

Lee JS, Ho WK, Lee SH (2010) Post-tetanic increase in the fast-releasing synaptic vesicle pool
at the expense of the slowly releasing pool. J Gen Physiol 136:259-272.

Lee JS, Ho WK, Lee SH (2012) Actin-dependent rapid recruitment of reluctant synaptic vesicles
into a fast-releasing vesicle pool. Proc Natl Acad Sci U S A 109:E765-774.

Li YC, Chanaday NL, Xu W, Kavalali ET (2017) Synaptotagmin-1- and Synaptotagmin-7-
Dependent Fusion Mechanisms Target Synaptic Vesicles to Kinetically Distinct
Endocytic Pathways. Neuron 93:616-631 e613.

Lin JY, Lin MZ, Steinbach P, Tsien RY (2009) Characterization of engineered channelrhodopsin
variants with improved properties and kinetics. Biophys J 96:1803-1814.

Lin KH, Taschenberger H, Neher E (2022) A sequential two-step priming scheme reproduces
diversity in synaptic strength and short-term plasticity. Proc Natl Acad Sci U S A
119:e2207987119.

Lin KH, Ranjan M, Lipstein N, Brose N, Neher E, Taschenberger H (2025) Number and relative
abundance of synaptic vesicles in functionally distinct priming states determine
synaptic strength and short-term plasticity. The Journal of Physiology.

Liu H, Bai H, Hui E, Yang L, Evans CS, Wang Z, Kwon SE, Chapman ER (2014) Synaptotagmin 7
functions as a Ca2+-sensor for synaptic vesicle replenishment. Elife 3:e01524.

Lou X, Korogod N, Brose N, Schneggenburger R (2008) Phorbol esters modulate spontaneous
and Ca2+-evoked transmitter release via acting on both Munc13 and protein kinase C.
J Neurosci 28:8257-8267.

Macia E, Ehrlich M, Massol R, Boucrot E, Brunner C, Kirchhausen T (2006) Dynasore, a cell-
permeable inhibitor of dynamin. Dev Cell 10:839-850.

Malagon G, Miki T, Tran V, Gomez LC, Marty A (2020) Incomplete vesicular docking limits
synaptic strength under high release probability conditions. Elife 9.

Markram H, Wang Y, Tsodyks M (1998) Differential signaling via the same axon of neocortical
pyramidal neurons. Proc Natl Acad Sci U S A 95:5323-5328.

Martinetti LE, Bonekamp KE, Autio DM, Kim HH, Crandall SR (2022) Short-Term Facilitation of
Long-Range Corticocortical Synapses Revealed by Selective Optical Stimulation. Cereb
Cortex 32:1932-1949.

Miki T, Malagon G, Pulido C, Llano I, Neher E, Marty A (2016) Actin- and Myosin-Dependent
Vesicle Loading of Presynaptic Docking Sites Prior to Exocytosis. Neuron 91:808-823.

Mongillo G, Barak O, Tsodyks M (2008) Synaptic theory of working memory. Science 319:1543-1546.

Mongillo G, Hansel D, van Vreeswijk C (2012) Bistability and spatiotemporal irregularity in neuronal networks with nonlinear synaptic transmission. Phys Rev Lett 108:158101.

Neher E (1995) The use of fura-2 for estimating Ca buffers and Ca fluxes. Neuropharmacology 34:1423-1442.

Neher E (2010) What is Rate-Limiting during Sustained Synaptic Activity: Vesicle Supply or the Availability of Release Sites. Front Synaptic Neurosci 2:144.

Neher E (2024) Interpretation of presynaptic phenotypes of synaptic plasticity in terms of a two-step priming process. J Gen Physiol 156.

Neher E, Sakaba T (2008) Multiple roles of calcium ions in the regulation of neurotransmitter release. Neuron 59:861-872.

Neher E, Brose N (2018) Dynamically Primed Synaptic Vesicle States: Key to Understand Synaptic Short-Term Plasticity. Neuron 100:1283-1291.

Norman CA, Krishnakumar SS, Timofeeva Y, Volynski KE (2023) The release of inhibition model reproduces kinetics and plasticity of neurotransmitter release in central synapses. Commun Biol 6:1091.

Oheim M, Naraghi M, Muller TH, Neher E (1998) Two dye two wavelength excitation calcium imaging: results from bovine adrenal chromaffin cells. Cell Calcium 24:71-84.

Otsuka T, Kawaguchi Y (2009) Cortical inhibitory cell types differentially form intralaminar and interlaminar subnetworks with excitatory neurons. J Neurosci 29:10533-10540.

Ozdemir AT, Lagler M, Lagoun S, Malagon-Vina H, Lasztoczi B, Klausberger T (2020) Unexpected Rule-Changes in a Working Memory Task Shape the Firing of Histologically Identified Delay-Tuned Neurons in the Prefrontal Cortex. Cell Rep 30:1613-1626 e1614.

Pan B, Zucker RS (2009) A general model of synaptic transmission and short-term plasticity. Neuron 62:539-554.

Papantoniou C, Laugks U, Betzin J, Capitanio C, Ferrero JJ, Sanchez-Prieto J, Schoch S, Brose N, Baumeister W, Cooper BH, Imig C, Lucic V (2023) Munc13- and SNAP25-dependent molecular bridges play a key role in synaptic vesicle priming. Sci Adv 9:eadf6222.

Pennington ZT, Dong Z, Feng Y, Vetere LM, Page-Harley L, Shuman T, Cai DJ (2019) ezTrack: An open-source video analysis pipeline for the investigation of animal behavior. Sci Rep 9:19979.

Pulido C, Marty A (2018) A two-step docking site model predicting different short-term synaptic plasticity patterns. J Gen Physiol 150:1107-1124.

Raybuck JD, Lattal KM (2014) Bridging the interval: theory and neurobiology of trace conditioning. Behav Processes 101:103-111.

Reyes A, Sakmann B (1999) Developmental switch in the short-term modification of unitary EPSPs evoked in layer 2/3 and layer 5 pyramidal neurons of rat neocortex. J Neurosci 19:3827-3835.

Rhee JS, Betz A, Pyott S, Reim K, Varoqueaux F, Augustin I, Hesse D, Sudhof TC, Takahashi M, Rosenmund C, Brose N (2002) Beta phorbol ester- and diacylglycerol-induced augmentation of transmitter release is mediated by Munc13s and not by PKCs. Cell 108:121-133.

Ritzau-Jost A, Jablonski L, Viotti J, Lipstein N, Eilers J, Hallermann S (2018) Apparent calcium dependence of vesicle recruitment. J Physiol 596:4693-4707.

Ritzau-Jost A, Delvendahl I, Rings A, Byczkowicz N, Harada H, Shigemoto R, Hirrlinger J, Eilers J, Hallermann S (2014) Ultrafast action potentials mediate kilohertz signaling at a central synapse. Neuron 84:152-163.

Rozov A, Burnashev N, Sakmann B, Neher E (2001) Transmitter release modulation by intracellular Ca2+ buffers in facilitating and depressing nerve terminals of pyramidal cells in layer 2/3 of the rat neocortex indicates a target cell-specific difference in presynaptic calcium dynamics. J Physiol 531:807-826.

Sakaba T, Neher E (2003) Involvement of actin polymerization in vesicle recruitment at the calyx of Held synapse. J Neurosci 23:837-846.

Scheuss V, Schneggenburger R, Neher E (2002) Separation of presynaptic and postsynaptic contributions to depression by covariance analysis of successive EPSCs at the calyx of Held synapse. J Neurosci 22:728-739.

Schotten S, Meijer M, Walter AM, Huson V, Mamer L, Kalogreades L, ter Veer M, Ruiter M, Brose N, Rosenmund C (2015) Additive effects on the energy barrier for synaptic vesicle fusion cause supralinear effects on the vesicle fusion rate. Elife 4:e05531.

Silva M, Tran V, Marty A (2021) Calcium-dependent docking of synaptic vesicles. Trends Neurosci 44:579-592.

Silva M, Tran V, Marty A (2024) A maximum of two readily releasable vesicles per docking site at a cerebellar single active zone synapse. Elife 12.

Stevens CF, Wesseling JF (1999) Augmentation is a potentiation of the exocytotic process. Neuron 22:139-146.

Sugita S, Han W, Butz S, Liu X, Fernandez-Chacon R, Lao Y, Sudhof TC (2001) Synaptotagmin VII as a plasma membrane Ca(2+) sensor in exocytosis. Neuron 30:459-473.

Taschenberger H, Woehler A, Neher E (2016) Superpriming of synaptic vesicles as a common basis for intersynapse variability and modulation of synaptic strength. Proc Natl Acad Sci U S A 113:E4548-4557.

Tawfik B, Martins JS, Houy S, Imig C, Pinheiro PS, Wojcik SM, Brose N, Cooper BH, Sorensen JB (2021) Synaptotagmin-7 places dense-core vesicles at the cell membrane to promote Munc13-2- and Ca(2+)-dependent priming. Elife 10.

Thomson AM, Lamy C (2007) Functional maps of neocortical local circuitry. Frontiers in neuroscience 1:48.

Tran V, Silva M, Marty A (2023) Prioritized docking of synaptic vesicles provided by a rapid recycling pathway. iScience 26:106366.

Tremblay R, Lee S, Rudy B (2016) GABAergic Interneurons in the Neocortex: From Cellular Properties to Circuits. Neuron 91:260-292.

Turecek J, Jackman SL, Regehr WG (2016) Synaptic Specializations Support Frequency-Independent Purkinje Cell Output from the Cerebellar Cortex. Cell Rep 17:3256-3268.

Turecek J, Jackman SL, Regehr WG (2017) Synaptotagmin 7 confers frequency invariance onto specialized depressing synapses. Nature 551:503-506.

Valera AM, Doussau F, Poulain B, Barbour B, Isope P (2012) Adaptation of granule cell to Purkinje cell synapses to high-frequency transmission. J Neurosci 32:3267-3280.

van Aerde KI, Feldmeyer D (2015) Morphological and physiological characterization of
pyramidal neuron subtypes in rat medial prefrontal cortex. Cereb Cortex 25:788-805.
Vandael D, Borges-Merjane C, Zhang X, Jonas P (2020) Short-Term Plasticity at Hippocampal
Mossy Fiber Synapses Is Induced by Natural Activity Patterns and Associated with
Vesicle Pool Engram Formation. Neuron 107:509-521 e507.
Vevea JD, Kusick GF, Courtney KC, Chen E, Watanabe S, Chapman ER (2021) Synaptotagmin 7
is targeted to the axonal plasma membrane through gamma-secretase processing to
promote synaptic vesicle docking in mouse hippocampal neurons. Elife 10.
Voleti R, Tomchick DR, Sudhof TC, Rizo J (2017) Exceptionally tight membrane-binding may
explain the key role of the synaptotagmin-7 C(2)A domain in asynchronous
neurotransmitter release. Proc Natl Acad Sci U S A 114:E8518-E8527.
Wang LY, Neher E, Taschenberger H (2008) Synaptic vesicles in mature calyx of Held synapses
sense higher nanodomain calcium concentrations during action potential-evoked
glutamate release. J Neurosci 28:14450-14458.
Wu LG, Chan CY (2022) Multiple Roles of Actin in Exo- and Endocytosis. Front Synaptic
Neurosci 14:841704.
Yau JO, McNally GP (2023) The Rescorla-Wagner model, prediction error, and fear learning.
Neurobiol Learn Mem 203:107799.
Yoon JY, Lee HR, Ho WK, Lee SH (2020) Disparities in Short-Term Depression Among Prefrontal
Cortex Synapses Sustain Persistent Activity in a Balanced Network. Cereb Cortex
30:113-134.
Zucker RS, Regehr WG (2002) Short-term synaptic plasticity. Annu Rev Physiol 64:355-405.
1238

Figure 1. Frequency-dependence of short-term synaptic plasticity at prelimbic L2/3

excitatory synapses.

(**A**) In utero electroporation following injection of the plasmid (CAG-oChIEF-tdTomato) into

the ventricle of an embryo (E17.5). (**B**) Representative images showing specific expression of

oChIEF-tdTomato in L2/3 pyramidal cells after IUE. The red fluorescence of tdTomato clearly

visualizes oChIEF-expressing cell bodies in L2/3 and axons in L5. Scale bar: 1 mm, 100 μm, 50

μm from left to right. (**C**) Recording schematic showing photostimulation of oChIEF-expressing

axon fibers of transfected PCs (*filled triangles*) and a whole-cell recording from non-

transfected PCs (*empty triangles*) or FSINs (*empty circles*). A collimated DMD-coupled LED was

used to confine the area of excitation (typically 3–4 μm in diameter) to a small region (*blue*

circle) near the soma. (**D, E**) Representative traces for EPSCs averaged over 10 trials at each

frequency (*left*) and average amplitudes of baseline-normalized EPSCs (*right*) during 20-pulse

trains at frequencies from 5 to 40 Hz at PC-PC (*D*; n = 12, 9, 21, 10 cells for 5, 10, 20, 40 Hz,

respectively) and PC-FSIN (*E*; n = 8, 9, 10, 10) synapses. Each data point was normalized to the

average of the first EPSC. (**F**) Baseline-normalized amplitudes of steady-state EPSCs (EPSC$_{ss}$;

black symbols) and synaptic efficacy (EPSC$_{ss}$ × *f*; *red symbols*) as a function of stimulation

frequency (*f*) at PC–PC synapses. EPSC$_{ss}$ was measured from the average of last 5 EPSCs from

the 20-pulse trains (n = 12, 9, 21, 10). (**G**) Paired pulse ratio (PPR) as a function of inter-spike

intervals (n = 25, 9, 25, 22, 43 for 200, 100, 50, 25, 20 ms ISI, respectively) at PC-PC synapses.

(**H**) EPSC$_{ss}$ and synaptic efficacy at PC-FSIN (n = 8, 9, 10, 10). (**I**) PPR at PC-FSIN (n = 8, 9, 10,

15, 34). *Gray symbols*, individual data.

Figure 2. Delayed facilitation results from slow activation of Ca²⁺-dependent vesicle replenishment at a constantly high vesicular fusion probability

(**A**) Representative traces (*left*) and mean baseline-normalized amplitudes (*right*) of EPSCs evoked by 30-pulse trains at 40 Hz in control (n = 21; *black*) and in the presence of 50 µM EGTA-AM (n = 14; *green*). (**B**) Mean values for the first EPSC amplitude (EPSC$_1$, *left*) and PPR (*right*) from the experiments displayed in (A). *Gray symbols*, individual data. (**C**) Plot of rate constants for short-term facilitation (k_{STF}) as a function of stimulation frequency (f_{stim}), showing a linear relationship. The linear regression line (*black*) is shown fitted to k_{STF} values, estimated from *Figure 1*. (**D**) Representative EPSC traces (*left*) and average of baseline-normalized EPSCs (*right*) evoked by 12-pulse stimulation at 5 Hz, followed by 40 Hz 7-pulse train (n = 12). Note that slowly developing facilitation was converted to rapid facilitation after strong PPD. (**E**) Mean values for PPR at 40 Hz. The baseline PPR was reproduced from *Figure 1G* and the PPR during 5 Hz train was calculated as (13th EPSC) / (12th EPSC). *Gray symbols*, individual data. All statistical data are represented as mean ± S.E.M.; n.s. = not significant; **, P ＜ 0.01; unpaired t-test.

Figure 3. Low baseline occupancy of release sites and its increase during facilitation and

post-tetanic augmentation

(**A, B**) *Left*, Representative EPSCs evoked by 5 and 40 Hz train stimulation (*black*, 5 Hz; *purple*,

40 Hz). *Right*, Variance-mean plots of EPSCs amplitude from averaged EPSCs recorded at PC-

PC (A; n = 12, 17 for 5 and 40 Hz, respectively) and PC-FSIN (B; n = 8, 10) synapses. The data

were fitted using multiple-probability fluctuation analysis (MPFA). Error bars are omitted for

clarity. The 1st EPSC of 5 Hz train (*broken line*) was used to estimate the resting level of p_{occ}.

Filled circles were measured from post-tetanic augmented EPSCs (*A*, n = 12, 12, 9; *B*, n = 7, 7,

7 for 0.1, 0.2, 0.5 s IBIs, respectively). (**C-E**) Post-tetanic augmentation (PTA) experiments at

PC-PC (*top*) and PC-FSIN (*bottom*) synapses. (**C**) Representative traces for EPSCs evoked by

double 40 Hz train stimulations separated by 0.5 s. (**D**) Mean baseline-normalized amplitudes

of EPSCs evoked by double 40 Hz trains at different inter-burst intervals (IBIs, 0.1, 0.2, 0.5, 1,

2, 5, 10 s). *Upper*, PC-PC synapse (n = 12, 12, 21, 11, 11, 16, 11 from short to long IBIs,

respectively). *Lower*, PC-FSIN synapse (n = 10, 9, 8, 9, 9, 7, 9). (**E**) PTA time course. The

baseline-normalized amplitudes of 1^{st} EPSC from the 2^{nd} train were plotted as a function of

IBIs.

Figure 4. Pharmacological experiments reveal specific molecular mechanisms underlying

vesicle loading processes

(**A-C**) (**a**) Representative EPSC traces (*left*) and mean baseline-normalized EPSCs (*right*) evoked

by double 40 Hz train stimulations separated by 0.5 s inter-burst interval (IBI) in control and

in the presence of 5 μM U73122 (*Aa*, n = 7, *orange*), 20 μM OAG (*Ba*, n = 12, *cyan*) or 100 μM

dynasore (*Ca*, n = 10, *blue*). (**b**) Mean values for baseline EPSCs (EPSC$_1$, *left*) and augmentation

(*right*) from the experiments shown in corresponding *a* panel. (**Da**) Representative EPSC

traces evoked by paired pulses (*left*) and mean values for baseline EPSC amplitude (*middle*)

and PPR (*right*) before and after applying 20 μM LatB (n = 6). (**Db**) Representative EPSC traces

(*left, upper*) and average of normalized EPSCs (*left, lower*) evoked by double 40 Hz train

stimulation separated by 0.5 s in control (n = 21) and in 20 μM LatB conditions (n = 16; *pink*).

Right, Mean values for augmentation in control and LatB conditions. *Gray symbols*, individual

data. All statistical data are represented as mean ± S.E.M., *, $P < 0.05$; **, $P < 0.01$; ***, $P <$

0.001; unpaired or paired t-test; n.s. = not significant.

Figure 5. STF at both types of local excitatory synapses is abolished by Syt7 KD

(**A, B**) Representative EPSC traces (*left*) and mean baseline-normalized amplitudes of EPSCs (*right*) evoked by 20-pulse trains at 5 to 40 Hz. STP was measured at PC-PC (*A*; n = 10, 9, 12, 9) and PC-FSIN (*B*; n = 9, 8, 10, 9) synapses, in which presynaptic Syt7 transcripts were depleted (Syt7 KD). Syt7 KD pyramidal cells are indicated as *black triangles* on the top. For comparison, STP in WT synapses is reproduced from *Figure 1* (*dotted lines*). Same frequency color codes were used as in *Figure 1*. (**C**) Baseline-normalized amplitudes of steady-state EPSC (EPSC$_{ss}$; *black symbols*) and synaptic efficacy (EPSC$_{ss}$ × *f*; *red symbols*) as a function of

81 stimulation frequency (f) at PC-PC synapses. $EPSC_{ss}$ was defined as the average of last 5 EPSC

82 amplitudes from 20-pulse trains. (**D**) PPR as a function of inter-spike intervals (n = 10, 9, 12,

83 18, 26). (**E**) $EPSC_{ss}$ and synaptic efficacy at PC-FSIN synapses. (**F**) PPR at PC-FSIN synapses (n =

84 9, 8, 10, 12, 24). *Gray symbols*, individual data.

Figure 6. Syt7 KD synapses exhibit complementary changes in the number of release sites and their vesicle occupancy

(**A, B**) *Left*, Representative traces of EPSCs evoked by 5 and 40 Hz train stimulations (*black*, 5 Hz; *purple*, 40 Hz). *Right*, Variance-mean plots of EPSCs amplitude from averaged EPSCs recorded at PC-PC (*A*; n = 9, 15, 12, 10, 9) and PC-FSIN (*B*; n = 4, 14, 10, 9, 8) synapses in which presynaptic Syt7 has been knocked down. The data were fitted using MPFA and error bars are omitted for clarity. The mean 1st EPSC amplitude of 5 Hz train (*vertical broken line*) was used for estimation of baseline p$_{occ}$. Dashed parabolas indicate MPFA fits to variance-mean plot of WT synapses reproduced from *Figure 3*. (**C-E**) Recovery experiments at PC-PC (*top*) and PC-FSIN (*bottom*) synapses in which presynaptic Syt7 was knocked-down. (**C**) Representative EPSCs evoked by double 40 Hz train stimulations separated by 0.5 s. (**D**) Mean baseline-normalized amplitudes of EPSCs evoked by double 40 Hz trains at PC-PC (n = 12, 10, 9, 9, 10, 9, 8) and PC-FSIN (n = 10, 9, 10, 11, 8, 9, 10) synapses at different interburst intervals (IBIs).

100   (E) Recovery time course. Baseline-normalized amplitudes of 1^{st} EPSC from the 2^{nd} burst were

101   plotted as a function of various IBIs. *Dotted lines* indicate augmented EPSCs in the WT

102   reproduced from *Figure 3*.

Figure 7. Recovery of TS vesicles following depletion is accelerated by Syt7

(**A-C**). Recovery experiments at PC-PC synapses in WT (*A*) and Syt7 KD (*B*). Mean baseline-normalized amplitudes of EPSCs evoked by two consecutive 3-pulse 40 Hz trains in WT (n = 16, 13, 11, 11, 11, 9 from short to long IBIs, respectively) and KD (n = 13, 11, 12, 8, 10, 11) synapses at different IBIs (0.1, 0.2, 0.5, 1, 2, 5 s). *Inset*, representative traces of EPSCs evoked by two consecutive 3-pulse 40 Hz train stimulations separated by 0.2 s. (**D-F**). Recovery experiments at PC-FSIN synapses in WT (*D*) and Syt7 KD (*E*). Mean baseline-normalized amplitudes of EPSCs evoked by two consecutive 3-pulse 40 Hz trains in WT (n = 9, 11, 11, 9, 10, 13) and KD (n = 9, 12, 9, 7, 10, 9) synapses at different IBIs. *Inset*, representative traces of EPSCs evoked by two consecutive 3-pulse 40 Hz train stimulations separated by 0.2 s. (**C, F**) Recovery time course. Baseline-normalized amplitudes of 1st EPSC from the 2nd burst were plotted as a function of various IBIs.

Figure 8. Behavioral effects of Syt7 deficiency in L2/3 PCs of the mPFC

(**A**) *Top*, Representative images showing bilateral expression of U6-GFP in L2/3 of PCs after IUE at E17.5. Scale bar: 1 mm, 100 μm. *Bottom*, Schematic of trace fear conditioning and extinction (tone test) protocol. (**B**) Freezing behavior of control (expressing scrambled shRNA, *Scr*) and Syt7 KD rats during acquisition of tFC. (**C**) Freezing ratio during tone tests on following

days. Data are shown as average freezing during T1–T4, T5–T8, or T9–T12 (T, trials). The

freezing on T1-T4 was significantly lower in KD rats suggesting that formation of trace memory

was impaired in Syt7 KD rats (n = 10, 11 for Ctrl and KD, respectively; $P = 0.0007$, $F_{(1, 19)} =$

16.43, two-way repeated measures ANOVA; $P = 0.0064, 0.0211, 0.0064, 0.0064$; Holm-Sidak

test). (**D**) Representative images of c-Fos immunoreactivity in the prelimbic cortex of control

or Syt7 KD rats 90 min after tFC acquisition. c-Fos (*left*, *red*, Cy5) and GAD67 (*right*, *cyan*, Cy3)

were immunostained in the same brain slice expressing U6-GFP (*middle*; *green*). Scale bar, 50

μm. (**E**) Exemplar images of c-Fos positive neurons expressing GFP (*top*) or GAD67 (*bottom*)

in control (*left*) or Syt7 KD rats (*right*). Scale bar, 50 μm. (**F-H**) Effects of Syt7 KD on c-Fos

density (*F*) and percentage of c-Fos positive neurons co-labeled with GAD67 (*G*) or GFP (*H*) in

L2/3 or L5 of prelimbic cortex (n = 9, 9 for Ctrl and KD, respectively). *Open symbols*, individual

data. All statistical data are represented as mean ± S.E.M.; ****, $P < 0.0001$; unpaired t-test.

Figure 9. STP model in light of known Ca²⁺ binding kinetics of Syt7

(**A-D**) *Left*, Schematic of allosteric calcium binding to Syt7. The number of Ca²⁺ bound to Syt7 was denoted as # in 'S#' in the reaction scheme. k_{on} = 7/μM/s, k_{off} = 10/s. *Middle*, Simulated changes of k_1 (*black*) in response to 5 Hz train of Ca²⁺ transients (*light blue traces*). The priming step of the simple refilling model was assumed to be catalyzed by full Ca²⁺-bound form of Syt7. Accordingly, Ca²⁺-dependent increase in k_1 was calculated as $K_{1,max}$ multiplied by a fraction of full Ca²⁺-bound form of Syt7. We assumed that local $[Ca^{2+}]_i(t)$ follows a Gaussian function: $(1/\sigma \sqrt{2\pi}) \exp[-(t-t_p)^2/ 2\sigma^2]$, in which t_p = 0.25 ms and σ = 0.085 ms. *Right*, Fits of the Syt7 model to the STP data. To fit this model to the STP data, $K_{1,max}$ was set to 300/s (*A*), 220/s (*B*), and 180/s (*C*). Cooperativity factor (*b*) was set to 0.35 (*A*), 0.2 (*B*), and 0.05 (*C*). k_1 is set to be constant for Syt7 KD (*D*). Same frequency color codes were used as in *Figure 1*.

Supplementary Figures

Progressive overfilling of readily releasable pool underlies short-term facilitation

at recurrent excitatory synapses in layer 2/3 of the rat prefrontal cortex

Jiwoo Shin[1,2,3], Seung Yeon Lee[2], Yujin Kim[1,2,3*], Suk-Ho Lee[1,2,3*]

Figure 1—figure supplement 1. Intrinsic membrane properties of mPFC neurons.

(**A**) Neuronal subtypes in mPFC layer 2/3. Cells were characterized based on their

electrophysiological properties (n = 66, 64 for PC and FSIN, respectively). Representative

voltage responses to current injection (-200 or +400 pA) of pyramidal cells (*left*) and fast-

spiking interneurons (*right*). (**B-D**) Mean resting membrane potential (*B*), input resistance (*C*),

and Sag ratio (*D*) measured in layer 2/3 PCs and FSINs. *Gray symbols*, individual data. (**E**)

Current-spike relationship in each cell type. All statistical data are represented as mean ±

S.E.M.; n.s. = not significant; ****, P＜0.0001; unpaired t-test or two-way ANOVA.

Figure 1—figure supplement 2. Test for consistency of light stimulation in oChIEF-expressing

L2/3 pyramidal cells.

(**A**) Representative images of an oChIEF-expressing neuron. Whole-cell recordings were

performed from the fluorescently labeled neurons in layer 2/3 of the prelimbic cortex. *Left*,

IR-DIC image (60×) of the pyramidal cell under whole-cell patch clamp. *Right*, fluorescence

image of td-Tomato in the same cell excited at 530 nm wavelength. (**B**) Current-clamp

recordings of action potentials evoked by 600 light pulse trains from an oChIEF-expressing

L2/3 cell. For somatic stimulation, we used the typical light stimulation conditions used in STP

experiments (5 ms in duration and 4 µm in radius; see *Methods*). Uniform action potentials

were produced with high fidelity, albeit not without compromise towards the end of the train

in the case of 40 Hz stimulation. Note that these results are similar to previously reported

values from neocortical neurons (Lin et al., 2009; Yoon et al., 2020). (**C**) Same experiments as

(*B*) but in Syt7 KD pyramidal cells that express oChIEF in L2/3 of the mPFC.

Figure 1—figure supplement 3. Properties of EPSCs evoked by minimal stimulation.

(**A**) Optical minimal stimulation. A collimated DMD-coupled LED was used to confine the area of excitation to a small region near the soma. To stimulate a minimal number of excitatory axon fibers, the radius of illumination area was increased from 2 µm by 50% at each step, and selected the smallest illumination area that elicits EPSCs (Yoon et al., 2020) before starting each recording session. The illumination area was in the range of 3 to 8 µm in radius, when measured at the focal plane (typically 3–4 µm) of a 60× water immersion objective (*blue circle in the photograph*). (**B-E**) Properties of minimally stimulated baseline EPSCs (failure-excluded). Medians of frequency distributions are presented. Decay times below 1 ms were excluded from analysis. From left to right, representative traces (100 pulses), EPSC amplitude, rise time (20-80%), decay time (80-20%), and time to peak (from stimulus onset) from WT PC-PC (*B*, n = 81), WT PC-FSIN (*C*, n = 77), Syt7 KD PC-PC (*D*, n = 72), and Syt7 KD PC-FSIN (*E*, n = 78) synapses.

Figure 1—figure supplement 4. Validation of STP of optically evoked EPSCs

(**A**) The high p_v at PC-PC synapses may result from direct photo-stimulation of axon boutons.

To test if optically evoked EPSCs depend on APs, the effect of 1 μM tetrodotoxin (TTX) was

tested. Applying 1 μM tetrodotoxin (TTX) and additionally 0.1 mM 4-aminopyridine, a D-type

K channel blocker abolished optically evoked EPSCs (n = 8). Even when repolarization was

hindered by subsequent addition of 100 μM 4-aminopyridine (4-AP), a potassium channel

blocker, EPSCs were not rescued. These findings suggest that the light-evoked responses are

dependent on APs of presynaptic axon fibers rather than photo-induced depolarization of

presynaptic boutons (Little and Carter, 2013). (**B**) Comparison of STP evoked by optical

stimulation with that by dual patch techniques. To test validity of STP of photostimulation-

evoked EPSCs, we compared STP of unitary EPSCs evoked by dual whole-cell patch at PC-FSIN

synapses. Both of EPSC1 and STP at 40 Hz recorded by using paired whole-cell recordings were

not significantly different from those evoked by photo-stimulation, supporting the validity of

our optogenetic methods. (**Ba**) Representative EPSC traces evoked by 20-pulse 40 Hz trains stimulation at pyramidal cell to fast spiking interneuron (PC-FSIN) synapses. A train of APs in a presynaptic PC (*top*), evoked EPSCs in a FSIN by dual whole-cell patch clamp (*middle*) or by optical stimulation (*bottom*) under same conditions. (**Bb**) Mean values for baseline-normalized amplitudes of EPSCs evoked by 40 Hz train at PC-FSIN synapses. *Opto*, optical stimulation (n = 10); *Dual*, dual patch techniques (n = 5). STP evoked by the two techniques were not significantly different (two-way repeated measures ANOVA, $F_{(1, 13)} < 0.01$, P = 0.955). (**Bc**) Mean values for the first EPSC in different stimulation methods (P = 0.5724, unpaired t-test). *Open circles*, individual data. All statistical data are represented as mean ± S.E.M.

Figure 1—figure supplement 5. Test for AMPAR desensitization and saturation

(**A**) Representative traces of EPSCs evoked by paired-pulse photostmulation at 20 ms ISI (*left*)

and mean values for paired pulse ratio (PPR, *right*) before and after applying 50 μM

cyclothiazide (CTZ, n = 8), an AMPA receptor (AMPAR) desensitization inhibitor. CTZ induced

no significant change in PPR arguing against the possibility that strong PPD is caused by

AMPAR desensitization. *Gray symbols*, individual data. (**B**) It has been shown that multi-

vesicular release occurs at recurrent excitatory synapses in neocortical L2/3 (Holler et al.,

2021). Whereas a single vesicle release is insufficient to saturate postsynaptic AMPARs, an

increase in the number of vesicular release during facilitation may saturate AMPAR resulting

in underestimation of facilitation. To test this possibility, facilitation at 20 Hz was measured

in the presence of 0.5 mM kynurenic acid (Kyn), a fast competitive antagonist (Diamond and

Jahr, 1997; Wadiche and Jahr, 2001). Fast blockade of AMPA receptors did not have significant

effects on short-term facilitation at excitatory synapses in the mPFC L2/3, suggesting that

AMPAR saturation may not be significant during facilitation. (**Ba**) Representative EPSC traces

(*left*) and average of baseline-normalized EPSCs (*right*) evoked by 40 Hz 20-pulse trains in

control and in the presence of 0.5 mM kynurenic acid (Kyn). Although facilitation was slightly

increased by Kyn, it was not statistically significant (n = 12; two-way repeated measures

ANOVA, F (1, 22) = 0.575, P = 0.456). The slightly higher facilitation may result from reduced

blocking efficiency of Kyn upon increased glutamate concentration in the synaptic cleft (Tong

and Jahr, 1994; Meyer et al., 2001). (**Bb**) Mean values for the baseline EPSC (*left*) and the

steady-state EPSC (*right*) before and after applying 0.5 mM kynurenate as shown in *Ba*. The

baseline EPSC was reduced by Kyn to 43.5% of control. *Gray symbols*, individual data. All

statistical data are represented as mean ± S.E.M.; n.s. = not significant; ***, P＜0.001; paired

t-test.

Figure 2—figure supplement 1. Effects of EGTA-AM on synaptic transmission

(**A**) Representative traces (*top*) and time courses of normalized amplitudes of the first EPSCs

(*bottom*) during paired-pulse photostimulation at 20 ms ISI in control and in the presence of

50 μM EGTA-AM (n = 8). The green horizontal bar indicates application of 50 μM EGTA-AM.

(**B**) Mean values for the first EPSC (*left*) and PPR (*right*) before and after EGTA-AM application.

Gray symbols, individual data. All statistical data are represented as mean ± S.E.M.; n.s. = not

significant; *, P＜0.05; paired t-test.

Figure 2—figure supplement 2. Presynaptic calcium kinetics at boutons of layer 2/3 pyramidal cells

(**A-C**) A mixture of Fluo-5F (150, 250 or 500 μM) and Alexa Flour 555 was loaded on their axonal boutons of L2/3-PCs through whole-cell patch techniques for dual excitation ratiometric measurements of the cytosolic $[Ca^{2+}]$ at axonal boutons. (**A**) The long-branched stretch of axon from L2/3 PCs visualized through maximum fluorescence intensity projection using confocal laser scanning of AF555 perfused via a patch clamp pipette. Scale bar, 10 μm. (**B**) *Top*, the selected bouton for recordings, among the three, is displayed with the corresponding scanning line. Scale bar, 5 μm. *Bottom*, Line-scan images for fluorescence of Fluo-5F and AF555. Scale bars, 50 ms and 2 μm. (**C**) Baseline-normalized fluorescence changes evoked by an AP. Fluorescence intensities were spatially averaged over the bouton shown in

(*B*), background-subtracted and normalized to the baseline value (dF/F$_0$). *Inset*, Ca^{2+} transient estimated from dF/F$_0$. Scale bar, 50 ms and 100 nM. (**D, E**) The kinetic parameters of an AP-CaT were determined from a plot for the inverse of AP-evoked [Ca^{2+}] increment (1/Δ[Ca^{2+}]) *vs.* calcium binding ratio (κ_B; Eq. 4 in *Material and Methods*). (**D**) Reciprocal plot of the AP-induced [Ca^{2+}] changes (1/Δ[Ca^{2+}]) vs. corresponding calcium binding constants (κ_B) according to Eq. 2 (n = 14). A linear fit to the plot gave the amplitude of AP-CaT (A$_{Ca}$ = 1.16 μM) from the y-intercept, representing the amplitude of free [Ca^{2+}] increase in the absence exogenous Ca^{2+} buffer. The calcium binding ratio of endogenous static Ca^{2+} buffers (κ_S) was estimated as 96 from the x-intercept of the linear regression based on Eq. 2 in *Material and Methods*. Linear regression line, P = 0.035. (**E**) Plot of [Ca^{2+}] decay time constant versus κ_B (n = 16). The time constant for [Ca^{2+}] decay from mono-exponential fit was plotted vs. κ_B, and a linear regression based on Eq. 3 yielded the κ_S value of 77 from the x-intercept. The y-intercept was 43 ms, an estimate for a dye-independent value for Ca^{2+} decay time constant (τ_c) (Eq. 3 in *Material and Methods*). Linear regression line, P = 0.011. (**F**) Resting [Ca^{2+}] measured from basal fluorescence ratio using Eq. 1 in *Material and Methods* as a function of [Fluo-5F] in patch pipettes. The horizontal line indicates 50 nM. The mean [Ca^{2+}]$_{rest}$ at three different concentrations of Ca^{2+} indicator dye were not different and measured as 50 nM, suggesting that the dye did not influence [Ca^{2+}]$_{rest}$. (**G-I**) Strong PPD and the subsequent STF may be caused by Ca^{2+} channel inactivation and Ca^{2+}-dependent facilitation (CDF) of calcium influx into presynaptic boutons, respectively. To test these possibilities, we measured presynaptic Ca^{2+} increments evoked by each of successive APs applied in a train at 40 Hz. (**G**) Fluorescence signals from a selected bouton with a scanning line. *Scale bars*, 3 μm (*top*), 100 ms and 2 μm (*bottom*). (**H**) Normalized fluorescence (dF/F$_0$) responses to a train of APs at 40 Hz. (**I**) Total Ca^{2+} increments (Δ[Ca^{2+}]$_T$; free plus bound form) are plotted against the number of APs in the train (n = 13). Δ[Ca^{2+}]$_T$ were calculated based on Eq. 2 with κ_S set to 90. No significant change was observed for the estimates of Δ[Ca^{2+}]$_T$ at each of first 4 APs during 40 Hz train, indicating that contributions of Ca^{2+} channel inactivation and CDF to STP are unlikely. Mean ± S.E.M., One-way ANOVA; n.s. = not significant.

Figure 3—figure supplement 1. Quantal size estimated from Sr²⁺-induced asynchronous

release.

(**A**) Representative traces (100 pulses) from PC-PC synapses. Black bars indicate the detection

window (100 ms) for release events in sham stimulation (*left bar*) and photostimulation (*right*

bar). (**B**) Representative distribution of asynchronous EPSCs at PC-PC synapses. *Left*, after

photostimulation. *Middle*, before stimulation. *Right*, difference between left and middle

panels. The difference distribution was fitted with a Gaussian function (m = 24.93, σ = 6.09).

(**C**) Comparisons of estimates for quantal size (q) between WT and Syt7 KD synapses. From

left to right: PC-PC (n = 16, 13 for WT and KD, respectively), PC-FSIN (n = 16, 15). *Gray symbols*,

individual data. All data are represented as mean ± S.E.M., unpaired t-test; n.s. = not

significant.

Figure 3—figure supplement 2. Analysis of double failure rate supports high p_v at excitatory

synapses

(**Aa**) Schematic of simple refilling model. (**Ab**) An exponential fit to the PPR plot as a function

of ISIs reproduced from *Figure 1G*. *Fitted function*, PPR = 1.1 [1 − exp(− t/43.5 ms)]. (**Ac**)

Representative traces for four combinations of success and failure of 1st and 2nd EPSCs

evoked by paired pulses with ISI of 20 ms at PC-PC synapses, showing successes in both pulses

(1_1), failures in both pulses (0_0), and mixed responses (1_0; 0_1). On the right side of each

trace, corresponding observation probability is shown in percentages (mean ± S.E.M. n = 57).

(**B-C**) Plots of the calculated P(F1, F2) values (*B*) and their differences (*C*) from the observed

value (6.2%) under the simple refilling model are displayed on the plane of k_1 versus p_v.

Figure 3—figure supplement 3. Minimal EPSC increase upon elevated [Ca²⁺]ₒ indicates

saturated pᵥ at baseline

(**A**) Representative time courses of normalized amplitudes of the first EPSCs during paired-

pulse photo-stimulation at 20 ms ISI under control conditions (1.3 mM [Ca²⁺]ₒ) and elevated

extracellular [Ca²⁺] (2.5 mM; n = 8). (**B**) Mean values for the first EPSC (*left*) and PPR (*right*)

under the two conditions. (**C-D**) Experiments of (A-B) were repeated in the slices that were

pre-incubated in aCSF containing 50 µM EGTA-AM (n = 20). Note that the modest extracellular

[Ca²⁺]-dependence of the first EPSC and PPR observed in (A-B) was abolished by 50 µM EGTA-

AM pre-incubation. *Gray symbols*, individual data. All statistical data are represented as mean

± S.E.M.; n.s. = not significant; *, P < 0.05; paired t-test.

Figure 5—figure supplement 1. Little effects of Syt7 KD on intrinsic properties of neurons

Whole-cell recordings were performed from the fluorescently labeled pyramidal neurons in

the prelimbic L2/3 of WT (n = 30) or Syt7 KD (n = 16) rats. (**A**) Resting membrane potential. (**B**)

Input resistance. (**C**) Sag ratio. *Filled triangles*, individual data. (**D**) Current-spike frequency

relationship in WT and KD cells. All statistical data are represented as mean ± S.E.M., unpaired

t-test or two-way ANOVA; n.s. = not significant.

Figure 5—figure supplement 2. Presynaptic expression of Syt7 rescues facilitation at L2/3

excitatory synapses

(**A, D**) Representative amplitudes of EPSC trains evoked by 600-pulse 10 Hz trains (flanked by

2 baseline periods comprised of 40 pulses each) at PC-PC (*A*) and PC-FSIN (*D*) synapses, in

which presynaptic WT (*left, gray triangles*) pyramidal cells and those transfected with plasmid

encoding shRNA against Syt7 (sh-Syt7, KD, *middle, black triangles*) or sh-Syt7 plus Syt7

insensitive to shRNA (rescue, *right, red triangles*). (**B, E**) STP of EPSCs evoked by 600-pulse

trains at 10 Hz at L2/3 PC-PC (*B*; n = 9, 10, 7; WT, Syt7 KD, Rescue, respectively) and PC-FSIN

(*E*; n = 12, 10, 8) synapses. Each data point represents an average from a moving window of

20 pulses, normalized to the baseline EPSC. (**C, F**) Mean values of 41-60th pulses from *B* and *E*.

Mean ± S.E.M.; n.s. = not significant; **, P＜0.01; ***, P＜0.001; ****, P＜0.0001; one-way

ANOVA with Tukey's post-hoc test.

Figure 8—figure supplement 1. Activity-dependent increase in the synaptic weight ratio at

PC-PC to at PC-FSIN connections (J_{ee}/J_{ie}) is abolished by Syt7 KD

(**A, C**) Average amplitudes of baseline-normalized EPSCs during 20-pulse trains at frequencies

from 5 to 40 Hz at PC-PC (*left*) and PC-FSIN (*right*) synapses in WT (*A*) and Syt7 KD (*C*). Data

are reproduced from *Figure 1* (*A*) and *Figure 5* (*C*). (**B, D**) Relative synaptic weights of

excitatory input to PCs over that onto FSINs (*Jee/Jie*) in WT (*B*) and Syt7 KD (*D*). Note that

excitatory influence (*Jee/Jie*) to the recurrent network increases as the firing number and

frequency increase in WT but not in Syt7 KD.

Figure 8—figure supplement 2. Locomotor activity and anxiety levels in Syt7 KD rats using

OFT and EPM

(**A, C**) Representative tracking trace of the elevated plus maze (*A*) and open field test (*C*). (**B,**

D) Mean values for total distance moved (*left*) and time spent (*right*) in each arm (*B*) or in

periphery zone (*D*) in control (n = 13) or Syt7 KD rats (n = 9). Mean ± S.E.M.; Unpaired t-test

or two-way ANOVA; n.s. = not significant.

References

Diamond JS, Jahr CE (1997) Transporters buffer synaptically released glutamate on a submillisecond time scale. J Neurosci 17:4672-4687.

Holler S, Kostinger G, Martin KAC, Schuhknecht GFP, Stratford KJ (2021) Structure and function of a neocortical synapse. Nature 591:111-116.

Lin JY, Lin MZ, Steinbach P, Tsien RY (2009) Characterization of engineered channelrhodopsin variants with improved properties and kinetics. Biophys J 96:1803-1814.

Little JP, Carter AG (2013) Synaptic mechanisms underlying strong reciprocal connectivity between the medial prefrontal cortex and basolateral amygdala. J Neurosci 33:15333-15342.

Meyer AC, Neher E, Schneggenburger R (2001) Estimation of quantal size and number of functional active zones at the calyx of Held synapse by nonstationary EPSC variance analysis. J Neurosci 21:7889-7900.

Tong G, Jahr CE (1994) Multivesicular release from excitatory synapses of cultured hippocampal neurons. Neuron 12:51-59.

Wadiche JI, Jahr CE (2001) Multivesicular release at climbing fiber-Purkinje cell synapses. Neuron 32:301-313.

Yoon JY, Lee HR, Ho WK, Lee SH (2020) Disparities in Short-Term Depression Among Prefrontal Cortex Synapses Sustain Persistent Activity in a Balanced Network. Cereb Cortex 30:113-134.