A cell and transcriptome atlas of the human arterial vasculature

Quanyi Zhao*[1], Albert Pedroza*[2], Disha Sharma*[1], Wenduo Gu[1], Alex Dalal[2], Chad Weldy[1,3], William Jackson[1], Daniel Yuhang Li[1], Yana Ryan[1], Trieu Nguyen[1], Rohan Shad[2], Brian T. Palmisano[1], João P. Monteiro[1], Matthew Worssam[1], Alexa Berezwitz[1], Meghana Iyer[1], Huitong Shi[1], Ramendra Kundu[1], Lasemahang Limbu[2], Juyong Brian Kim[1,3], Anshul Kundaje[4], Michael Fischbein[2], Robert Wirka[5], Thomas Quertermous[1,3#], Paul Cheng[1,3#]

[1]Division of Cardiovascular Medicine, Stanford University School of Medicine, Stanford, CA 94305

[2]Department of Cardiothoracic Surgery, Stanford University School of Medicine, Stanford, CA 94305

[3]Cardiovascular Institute, Stanford University School of Medicine, Stanford, CA 94305

[4]Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305

[5]Division of Cardiology, Departments of Medicine and Cell Biology and Physiology, McAllister Heart Institute, University of North Carolina, Chapel Hill

*Contributed equally

#Co-Correspondence:

Paul Cheng
Chengpa@stanford.edu
Falk Cardiovascular Research Building
CV 279, 870 Quarry Road
Stanford, CA 94305

Thomas Quertermous
Tomq1@stanford.edu
Falk Cardiovascular Research Building
CV 297, 870 Quarry Road
Stanford, CA 94305

SUMMARY:

Vascular beds show different propensities for different vascular pathologies, yet mechanisms

explaining these fundamental differences remain unknown. We sought to build a transcriptomic,

cellular, and spatial atlas of human arterial cells across multiple different arterial segments to

understand this phenomenon. We found significant cell type-specific segmental heterogeneity.

Determinants of arterial identity are predominantly encoded in fibroblasts and smooth muscle

cells, and their differentially expressed genes are particularly enriched for vascular disease-

associated loci and genes. Adventitial fibroblast-specific heterogeneity in gene expression

coincides with numerous vascular disease risk genes, suggesting a previously unrecognized

role for this cell type in disease risk. Adult arterial cells from different segments cluster not by

anatomical proximity but by embryonic origin, with differentially regulated genes heavily

influenced by developmental master regulators. Non-coding transcriptomes across arterial cells

contain extensive variation in lnc-RNAs expressed in cell type- and segment-specific patterns,

rivaling heterogeneity in protein coding transcriptomes, and show enrichment for non-coding

genetic signals for vascular diseases.

Keywords
single-cell, vascular development, vascular cell populations, arterial segments, non-coding RNA,
vascular diseases

INTRODUCTION:

Human vascular diseases are the leading cause of morbidity and mortality worldwide[1]. Different segments of the human aorta and its branches have seemingly identical morphology and structures, yet have dramatically distinct propensity for different vascular diseases[2-4]. Atherosclerotic disease frequently influences coronaries and carotids, while relatively sparing the descending thoracic aorta[5,6]. Hereditary aneurysmal diseases frequently affect the aortic root and ascending aorta but not other aortic segments[7,8]. Environmental triggers such as tobacco smoking and methamphetamine abuse disproportionally bring on pathologies in the infra-renal segments of the aorta and pulmonary arteries respectively[9,10]. Even inflammatory or infectious aortopathies, such as Takayasu's vasculitis, Kawasaki's diseases, or syphilis aortitis, have well documented segmental preferences[11-13]. While the differences in disease propensity are well documented, the underlying causal mechanisms are less clear. Prevalent hypotheses include differential exposure to life-long hemodynamic forces, along with distinct cellular lineages from different embryonic origins[14,15]. Understanding the cellular and molecular basis for the differences between these arteries may open the door to therapies for a variety of vascular diseases.

Lineage tracing data primarily from several animal models suggest a distinct pattern of embryonic origins for the different arteries that appears conserved through vertebrates[15-18]. The coronary artery is thought to derive from several sources[19,20]. The endothelium from coronary arteries can be traced to two distinct origins, the sinus venosus and the endocardium. Coronary endothelium is thought to instruct the surrounding mesenchyme, which consists of cells originating from the proepicardial organ, to turn into smooth muscle cells. [21,22]. The ascending aorta is thought to be made primarily of at least two populations of embryonic progenitor cells, arising from both the second heart field as well as the neural crest[21,23]. The two populations of cells mix in a distinct pattern, ending before the arch, where most smooth muscle cells are thought to come from neural crest cells of the 4th pharyngeal arch. The carotid artery arises from the 3rd pharyngeal arch. The pulmonary artery (PA) arises from the 6th pharyngeal arch, the thoracic descending aorta is thought to arise from the somitic mesoderm, and splenic mesoderm thought to be responsible for the iliac and femoral arteries. These distinct developmental origins have long been postulated to be causal for risk related to various diseases, as different segments have been shown to have differential expression of selected genes as studied with bulk RNA isolation[24,25]. Transplantation experiments in mammals have suggested arterial segments retain the disease propensities of their original site, suggesting intrinsic behavior of segments accounts for their disease propensity[26,27]. However, whether the same embryological developmental pattern exists in humans is unknown. In addition, the extent to which developmental origin influences cellular gene expression and behavior of adult vessels remains elusive.

To gain insight into these fundamental observations on differential arterial disease risk, we generated a human arterial cellular atlas spanning multiple segments of the arterial circulation. Single cell transcriptomic analysis revealed that different arterial segments exhibit distinct gene expression patterns in the adult. The distinct transcriptomic signatures were most discrepant between fibroblast and smooth muscle cells (SMC) from different vascular beds, while gene expression was relatively similar between endothelial cells and residential

99   inflammatory cells between the vascular beds. Taken together, these data suggest that vascular
100   identity is encoded mostly in the vascular fibroblast and smooth muscle cell populations.
101
RESULTS:
To generate a baseline human vascular cellular atlas simultaneously from multiple arterial sites
from the same person, we collaborated with the organ donor network to collect arterial tissues
from organ donors. Up to 9 distinct arterial sites (ascending thoracic aorta (ASC), aortic arch
(ARCH), descending thoracic aorta (DESC), infra-renal abdominal aorta (INFRARENAL), iliac
artery (ILIAC), coronary artery (RCA), carotid artery (CAROTID), pulmonary artery (PA), and
aortic root (ROOT)) were chosen to be evaluated due to their different embryonic origins and
disease propensities. Available segments were collected using identical surgical techniques by
a dedicated team of cardiothoracic surgeons to minimalize surgical variability (Figure 1A) and
processed in an identical manner from 6 different patients. Tissues were dissociated into single
cells using an enzymatic digestion approach we developed to overcome the dense vascular
connective tissue [28-30]. This method provided for vascular smooth muscle cells to be the
predominant population captured, matching histological cellular composition ratios, while
preserving single-cell RNA (scRNA) quality and library complexity. Not all sites were collected
from each patient as certain segments were retained along with organs that were utilized for
patient care. The digested cells were then sorted for viability and then loaded onto the 10x
Genomics platform to perform single-cell transcriptomic capture. Libraries that were made were
then sequenced at a targeted depth of at least 50,000 reads per cell, and the resultant data
processed using the Seurat platform. The dataset was then filtered to remove poor quality cells
and doublets. The overall dataset contained 187,958 cells. Individual cell gene expression was
then normalized based on read depth and mitochondrial/ribosomal fraction of reads, to allow for
comparison between datasets. The resultant data was then dimensionally reduced onto a 2D
representation using uniform manifold approximation and projection (UMAP) for ease of
visualization.
The cell type identities were then identified first via unsupervised low dimensional
clustering of the transcriptomic data. To avoid over-clustering, we initiated the analysis using
very low-resolution clustering, which enables the identification of known distinct cell types, such
as endothelial cells and smooth muscle cells. Visualization of the combined scRNA data
revealed the presence of all the expected major cell types in the human arteries, namely
endothelial cells (EC), smooth muscle cells (SMC), fibroblasts, microvascular cells and immune
cells. These cell types are represented using existing canonical markers of each population
(Figures 1B, S1A). The largest component of the variance in the dataset came from markers of
different cell types, though the data contained a small level of expected inter-person variation
such as gender-specific differences (Figure S1B). For ease of visualization and to focus on
biology that exists between vascular beds instead of inter-person variations, we performed
batch correction between different donors using Single-cell-transform (SCT)-based integration.
These batch corrections allow us to better visualize the biology in question[31]. The resultant data
are presented here (Figures 1C, S1C), and will be used as the main dimensional-reduction
framework for data visualization.
141
Cells of the Human Arteries

We proceeded to identify each of these specific populations of cells found in our scRNAseq data.
Currently, the cellular populations in the vessels are often defined by positive staining of one or
two marker genes (i.e. *CDH5* for endothelial cells), which relies on the intrinsic assumption of
"marker genes" being specific, and can fail to capture unexpected heterogeneity[17,32-35]. This had
led to significant debate regarding the presence and identity of certain cell types, such as
vascular mesenchymal stem cells, endothelial progenitor cells, adventitial progenitor cells,
pericytes, fibrocytes, fibromyocytes, modulated smooth muscle cells, fibroblasts etc[17,28,32-36].
Discrete populations of cell types should have distinct transcriptomic signatures at rest. The
single cell approach allows for identification of different populations of cells that exist in the
human arteries a priori utilizing only the data encoded in the global transcriptomes instead of
individual marker genes. Thus, we first sought to identify the location and specific markers for
different subpopulations of cells that seem readily apparent based on clustering of the cells'
global transcriptome using a very conservative clustering resolution. We subsequently applied
the canonical "cell type markers" that have been classically utilized to identify SMC, EC,
immune cells, fibroblast, pericytes, and various progenitor cells to identify each of these arterial
cell populations, to test both the specificity of the markers as well as identify cellular
heterogeneity within marker populations. Each of the sub-populations were then located to
arterial sections via complementary approaches using RNAscope of specific markers identified
from the dataset or mapped spatially with Slideseq (Figures 1D, S1D). Each one of these
populations of cells was then compared across the vessel beds to understand what encodes
arterial identity. The differentially expressed genes were then assessed for their relevance to
human disease by enrichment analysis against human vascular genetics data. In addition, this
dataset was then compared with existing murine datasets to identify differences between the
human and murine arterial vasculature. Lastly, utilizing the rich information captured with this
technique, we created an additional non-coding RNA atlas, to understand the variability and
diversity of non-coding transcripts across different vessel beds and cell types.
169
Smooth Muscle Cells of Great Arteries
Histologically, the largest population of cells in the great arteries are medial SMC and their
intimal derivatives that arise from phenotypic modulation of SMC. These cells make up the
majority (47.2% in RCA and 53.7% in AORTA) of the arterial wall[28,36,37]. While existing scRNA
datasets of arteries in the past have demonstrated inflammatory cells being the most numerous
populations of isolated cells[28,38,39], our more vigorous tissue digestion protocol allowed for
identification and isolation of SMC as by far the most numerous cell type, matching histological
observations. SMC were classically identified using expression of mature smooth muscle
contractile apparatus genes such as SMC heavy chain (*MYH11*) or calponin 1 (*CNN1*) (Figures
2A, 2B). Visualizing the population using several canonical SMC marker genes revealed a
continuous gradient of SMC maturity in healthy vessels as marked by a gradient of expression
of each of these "SMC-specific" markers. The medial and intimal SMC (as evidenced by
RNAscope and spatial transcriptome), exist in a distinct gradient of hierarchical maturity,
$MYOCD^+>CNN1^+>MYH11^+>ACTA2^+>TAGLN^+$ population. (Figure 2A). It is important to note
that commonly applied smooth muscle markers (smooth muscle actin/*ACTA2*, and
transgelin/*TAGLN*/SM22α) are expressed quite broadly, including robust expression of *ACTA2*
and *TAGLN* in fibroblast and endothelial populations. This has significant implications regarding

lineage tracing models using *Tagln* or *Acta2* loci as the driver for Cre-recombinase lineage
labeling, and perhaps explains the distinct results obtained using different Cre-recombinase in
the existing literature.
The high depth of sequencing has allowed for the identification of two large
transcriptomically distinct populations of SMC based on very conservative clustering of cells in
all vessel beds. Both populations of SMC contain a contractile population, expressing mature
SMC markers, and a more "phenotypically modulated" population. We first investigated the
smaller population of cells expressing mature SMC markers.
Beyond medial arterial SMC, the other cell population that has been described
previously to express mature SMC markers in the vessel wall are SMC -like cells associated
with arterioles and pericytes, which are thought to represent mural cells of the microvasculature.
Indeed, staining for specific contractile SMC marker *CNN1* revealed positive cells in both the
media and small vessels in adventitia (Figure 2B). Pericytes are a heterogenous population of
cells classically identified using expression of markers *RGS5* and *NG2/CSPG4*[34,40]. Interestingly,
these "canonical pericyte markers" are not confined to one transcriptomically distinct population,
and are frequently non-overlapping (Figures 2C, S2A), suggesting that cells previously thought
to be pericytes in different studies may represent different heterogeneous populations of cells,
including contractile SMC. Differentially expressed gene analysis across these two large SMC
populations revealed 587 significantly differentially expressed genes between these two clusters
of SMC populations (Figure 2D, Table S1), with pathway analysis demonstrating distinct
biological functions (Figures 2E, S2B). For example, these data suggest very distinct pattern of
expression of adrenergic receptors, with alpha adrenergic receptor 2A (*ADRA2A*) (Figure 2F)
found only in the microvascular population whereas *ADRA2C* is found more in the medial SMC)
(Figure S2C). Leveraging RNA in situ hybridization and spatial transcriptome with Slideseq, for
markers unique for this microvascular populations identified with scRNAseq, revealed that this
population of cells resides exclusively in the adventitia (Figures 2F, S2D). While these cells do
express some of the classical "pericyte" markers, due to the potential imprecise nature of these
markers as described above, we will call this population of cells "adventitial microvasculature"
cells.
The larger SMC population, which is the most abundant cell type in the arteries, the
markers map to the cells in the media and intima of the vessel. There was one subcluster of the
population where the mature SMC markers were lowly expressed, and another sublcuster
where the expression of contractile markers was lost altogether. To identify where these cells
were located, we performed spatial transcriptomic profiling using Slideseq in conjunction with
RNAscope for specific markers for this population. Consistent with previous work, one of the
most specifically expressed genes in these SMC subclusters included *TNFRSF11B* and *FN1*.
The less "mature SMC" cells localized to the intima (Figures 2G, 2H, S2E). In conjunction with
Myh11-cre SMC lineage tracing in murine models[28,41], this lends further evidence to the notion
that arterial intima, which is present in human but not mice, are likely smooth muscle derived,
arising from media SMC in a process analogous to neointimal formation in murine
atherosclerosis and intimal hyperplasia models.
228
Adventitial Fibroblast in Human Arteries

The second most abundant population of cells isolated in our dataset are adventitial fibroblasts,
which form a continuum in the transcriptomic space with smooth muscle cells. This population of
cells did not express mature SMC markers (Figures 1C, 2A), but did express high levels of
extracellular matrix components and modifiers (Figure 3A). Marker gene RNAscope studies for
this population, along with spatial transcriptomics, localized this population exclusively to the
adventitia (Figures 3B, 1E). Based on marker gene expression, we assigned this population to
be vascular adventitial fibroblast. Surprisingly, while it has been described that the adventitial
fibroblast is a heterogeneous population, they appeared to be relatively homogenous in
transcriptomic space within the same vascular segments. There were additional endothelial
cells and other microvascular cells that are in the adventitia, which makes it more heterogenous
than other layers, but the fibroblast themselves within each segment appeared to be one large
population. However, there did appear to be significant heterogeneity across the adventitial
fibroblasts among the different vascular beds (Figure S3A). Commonly used canonical fibroblast
marker genes[32,42], including fibroblast specific protein (FSP1/*S100A4*), *PDGFRA*, *COL1A1*,
GLI1, were expressed relatively diffusely in several different populations of cells (Figure S3B).
This likely explains part of the heterogeneity of results that comes from different lineage tracing /
manipulation Cre models in murine studies[43-45]. Using the scRNAseq data, we additionally
identified *PI16*, *SFRP2*, and *MFAP5* as specific marker genes for this adventitial fibroblast
population (Figure 3C, Table S2). Of note, genes highly expressed specifically in the adventitial
fibroblast population were enriched for genes known to cause genetic aortopathies[46-48],
including *FBN1*, *TGFBR2*, *LOX*, *FBLN1*, and *MFAP5* (Figure 3D). This suggests a potentially
prominent role for this population of cells in the etiology of aortopathies and other
vasculopathies.
253
Endothelial Cells
Endothelial cells heterogeneity across different types of vessels has been well described,
including at the single cell level in the organ specific endothelial atlases[35,49,50]. Even at a specific
arterial site, there is evidence of different populations of endothelial cells as determined by
single cell RNA sequencing of murine aorta[35]. Well established markers for endothelial cells
(*CDH5*, *VWF, CD31*) fall within two large distinct clusters (Figures 4A, S4A). A single layer of
endothelial cells is found in the intimal surface of the arteries, but frequently many more are
found in the adventitia associated with the adventitial micro-vasculature. In our data, the
microvascular endothelial cells appear to be the most abundant population of endothelial cells
(Figure 4B). The biggest determinant of the transcriptome of the endothelial cells appears to be
which region of the vascular wall (intima/adventitia) these cells occupy (Figure S4B). The larger
of the two clusters constitutes the majority of the endothelial cells ("Endo1"), and can be further
subdivided into additional subclusters (Figure 4C). One subcluster expresses markers
previously found to be associated with venous and capillary endothelium, (for example, *ACKR1*),
whereas another subcluster expresses *HEY1*, a Notch signaling responsive gene previously
associated with arterial endothelium (Figure 4D). Consistent with existing literature, the cluster
that expresses *ACKR1* is found exclusively in the microvasculature in the adventitia, whereas
the cluster that expresses *HEY1*, a key gene associated with arterial endothelial specification[51],
is found both in the adventitia as well as on the luminal surfaces. These data suggest that these
EC reside in arterioles in the adventitia as well as on the vascular surfaces (Figures 4E, S4C).

Arterial endothelial cells have been implicated in endothelial to mesenchymal transition
(EndoMT), whereby these cells start to express mesenchymal genes[52,53]. This is observed even
in our relatively healthy vessels, with expression of SMC marker genes *TAGLN* and *ACTA2*
readily found in small groups of endothelial cells at a low level (Figures 2A, 4F). Of note, while it
has been suggested that EndoMT may induce endothelial cell transition to smooth muscle cells,
we do not detect significant levels of any of the more selective, mature SMC genes (i.e. *CNN1*
or *MYH11*) being expressed in the endothelial subset. Lastly, using existing markers for
lymphatic endothelial cells (*PROX1, LYVE1*)[54,55], we can observe a small population in the
adventitial EC as well, though they consist of a very small minority (Figure 4F).
The second cluster of endothelial cells ("Endo2") are notably transcriptomically distinct
from the remaining endothelial cells. Interestingly, even within the same segment, this
population of cells appears to express different *HOX* genes compared to the other larger
endothelial population, suggesting they likely come from distinct origins (Figure 4G). This
endothelial cell population expresses a large number of distinct markers, many with known
biological function and vascular disease relevance, including *ITLN1, NRG1*[56-58] (Figure 4H). The
Endo2 population is found in all the different vascular territories, and accounts for a ~10% of the
total endothelial population. Interestingly, one of the most specific markers, *ITLN1*, is known to
be upregulated after large vascular ischemic events[59-61]. RNAscope using *ITLN1* revealed that
this population exists exclusively on the luminal surface, and are not found in the adventitia
similar to the other endothelial (Endo1) population (Figures 4I, 4J). Compared to previously
reported data, the marker gene for this population appears distinct from the different endothelial
populations reported in murine aortic endothelial cells[35,50]. However, the same distinct
population of endothelial cells appears to be present in other human large artery single cell
datasets[38,62]. Differentially expressed (DE) gene comparison between the two endothelial cell
populations (Table S3) revealed that they likely have distinct functions. Compared to the
numerically more abundant Endo1 population, the Endo2 population enriches for genes
associated with TGFb/BMP signaling, angiogenesis and SMC crosstalk, as well as a strong
disease association with atherosclerosis (Figures 4K, S4D). These findings suggest this Endo2
population may be uniquely involved in human vascular diseases.
303
Macrophages And Lymphocytes:
While classically not considered part of the vessel wall, emerging evidence suggests that
inflammatory cells play an important part in vascular homeostasis[11,33,62-64]. Our data further
supports this notion, as even though these donor samples were obtained from relatively healthy
vessels without clinically overt diseases, there is a significant number of macrophages and
lymphocytes present in all the vascular beds. There is a trend towards a slight increase in the
total fraction of immune cells in the atherosclerosis prone arterial beds (coronary, infra-renal
aorta) (Figure S5A). This may reflect early atherosclerosis, but there are no inflammatory cells
or overt plaque seen in the existing intima. The majority of the macrophages and lymphocytes
are found in the adventitia. Clusters of B and T cells can also occasionally be found in organized
centers of what appears to be tertiary lymphoid organs that abut the adventitia (Figure 1D).
Putative computational analysis of cell-to-cell crosstalk using CellChat[65] demonstrates
significant cross talk between the adventitial fibroblast, smooth muscle, and the immune cell

populations (Figure S5B). These findings suggest a potential role for resident vascular mural
cells in coordinating the vascular immune response.
319
Determinants of Arterial Identity
To understand differences between arterial segments, we first tried to determine whether there
was a consistent relative composition of different cell types within different arteries. Using the
low clustering resolution which identified the canonical populations of vascular cells, we
discovered that with identical digestion protocols, the relative proportion of cell populations were
mostly similar across different arterial beds (Figure S6A). There is a slight increase in
inflammatory cells in the vascular beds that are more prone to atherosclerosis, such as the
coronary artery, and infra-renal segment of the aorta (Figure S5A). However, there were no
large differences in cellular composition across arterial segments, similar to histological studies.
329   Visualization of a large dataset using UMAP accentuates the drivers of the variance
within the dataset[66,67]. Thus, visualization of the data in this manner also enables us to get a
sense of relative heterogeneity of each population of cells across the different vascular beds.
Simply visualizing this dataset coded by segmental origins, it became readily apparent that
there is significant heterogeneity between adventitial fibroblasts and smooth muscle cells
among the different vascular beds. However, there were relatively fewer differences between
the microvascular, inflammatory and endothelial populations across different vessel beds, as
demonstrated on the UMAP (Figure 5A). Because dimensional reduction such as UMAP can
distort true distances between populations[66,68], we repeated the analysis by looking at the
number of differentially expressed genes between the ascending and descending arteries, and
between the coronary and descending pairs (Figures S6B, S6C), as well as the mean Euclidean
distance[69] between the average global gene expression of each cell population across the
vascular beds (Figures S6D, S6E). Consistent with the UMAP visualization, both of these
analyses revealed fewer differences between endothelial cells and macrophages across
different vascular beds compared to fibroblast and smooth muscle cells. Interestingly, the same
analysis comparing Endo1 and Endo2 population showed much smaller difference between
Endo1 population but significantly more difference in the Endo2 population across vascular
beds (Figure S6E). These data suggest that the main determinant of arterial identities, at least
transcriptomically, appears to be held mostly by fibroblast, smooth muscle cells, and possibly
the smaller subset of intimal endothelial cells, Endo2 population, and much less so in the
remaining endothelial cells, microvasculature, and inflammatory cells.
350
Prominent Roles of Embryonic Origin in Adult Vascular Identity
We next investigated the main differences between SMC and fibroblasts from different vascular
beds. Examination of the differentially expressed genes across SMC and Fibroblasts of different
vascular beds revealed a large number of developmentally important transcription factors[28,70-75],
including *HOX* family genes, *ISL1*, *MEF2C*, *TCF21, HAND2, GATA4* etc. (Figure 5C). We
hypothesize, therefore, that embryonic origins are the dominant factor in vascular identity. We
regenerated a UMAP visualization focusing on the top transcriptomic variances that exist in
each individual population. This UMAP readily revealed distinct transcriptomic sub-clusters of
Fibroblast and SMC (Figures S7A, S7B). Consistent with our hypothesis, the proximity of
different populations of fibroblast by either Euclidean distance or visualization in UMAP space

appears to correspond to each segments presumed embryonic origin reported in animal models
(Figures 5D, 5E, S7C). Cells from arteries of different caliber and flow with similar embryonic
origins (i.e.aortic arch, and ascending aorta/carotid arteries) cluster closer together than their
connected counterparts (such as descending thoracic artery to aortic arch). This suggests that
embryonic origins of each of these arterial segments play a dominant role in adult gene
expression. Curiously, this developmental pattern is less preserved in Identical analysis of
macrophage, microvasculature, and Endo1, suggesting that in these cells the resting gene
expression may be less influenced by their embryonic origin (Figures S7D, S7E, S7F).
Many of the differentially regulated genes found on the list has been known to be
regulated by these transcription factors in different processes. To unbiasedly determine if these
lowly expressed developmental master-regulators continue to play an important role in adult
gene expression globally, we investigated whether the binding sites for these transcription
factors are enriched in differentially expressed genes between segments compared to other
variable genes in the dataset, focusing on the ascending and descending thoracic aorta as the
main test case. Transcription factors *ISL1*, *HAND2*, *MEF2C* are expressed in the second heart
field and neural crest cells which give rise to the ascending aorta, whereas *HOX* genes are
expressed in the somitic mesoderm which gives rise to the descending thoracic aorta (Figures
5F, S7G). Consistent with these transcription factors playing an important role, we saw a very
strong enrichment of ISL1, HAND2, MEF2C, and HOX binding sites in the regulatory regions of
differentially expressed genes in fibroblast and SMC across these vessel beds of distinct
embryonic origin (Figures 5G, S7H). These data suggest the embryonic origin and the
persistent expression of these critical transcription factors into adulthood plays a critical role in
arterial identity and their global gene expression in adult.
384
Adult Vascular Hox Gene Expression Patterns
While *HOX* genes' roles are best elucidated in development, as one of the most potent cell fate
and patterning signals[70,76], their role in adult tissue identity and disease is less well understood.
It has been shown that different arterial segments express different *HOX* genes by bulk
RNAseq[25]. Given the impressive enrichment of *HOX* gene binding motifs in regulatory regions of
most differentially expressed genes, we leveraged our dataset to better understand *HOX* gene
expression across the different cell types within each segment of the aorta.
It appears that adult *HOX* gene expression patterns from the same vessel segment are
conserved in adulthood and thus mirror their embryonic origin (Figures 6A, S8A). Descending
thoracic aorta, abdominal aorta, and iliac arteries each expresses distinct subsets of *HOX*
genes that are conserved from one person to the next (Figure S8B). However, we observed that
within the different cell types of a segment, each cell type expresses a different subset of *HOX*
genes. For example, in descending thoracic aorta, *HOXD8* is most highly expressed only by
Endo1, *HOX A4/A5* predominantly by Endo2, and *HOXA3* in SMC, and *HOX A5/B7* in fibroblast.
While smooth muscle and microvasculature are transcriptionally and phenotypically similar, and
AdvFib and microvasculature are geographically close, they express very distinct *HOX* genes
patterns (Figures 6A, S8C). On the other hand, adventitial fibroblast *HOX* gene expression is
quite similar to those of the medial SMC. This potentially reflects these two populations of cells
(medial SMC and AdvFib) have more closely shared embryonic origins compared to those of
microvasculature in the adventitia. As further evidence that the *HOX* genes likely play an

important role in adult arterial identity, we clustered the cells and segments based on *HOX*
genes alone. Hierarchical clustering of SMC and fibroblast based on *HOX* genes alone mirrors
hierarchical clustering of whole transcriptomes, and readily reflects each segment's
developmental origin (Figures 6B, 6C). Given that most immune cells are likely formed uniformly
in the marrow and migrate into each segment, unsurprisingly there is less distinction between
them. While the endothelial cells are some of the earliest to form during embryonic development
of the great vessels, with each segments' endothelial cells having distinct origins, they are
significantly less heterogeneous compared to neighboring segments of different embryonic
origin, with the potential exception of the Endo2 population (Figure S8D). Those populations
that differ from one vascular bed to another tend to have a very distinct pattern of *HOX* gene
expression, whereas cell types with similar *HOX* gene expressions tend to have globally more
similar transcriptomes. The persistence of this very distinct pattern of *HOX* gene expression
across individuals suggests that conserved transcriptomic regulation of these genes in
adulthood, along with enrichment in HOX binding motifs in differentially expressed genes noted
above suggests critical roles for *HOX* genes in cell type specific adult segmental specific gene
expression.
421
Enrichment of Vascular Disease Rare and Common Variant Genetic Signals in Segmental
Transcriptomic Differences
Our original hypothesis was that the transcriptomic differences between these vascular beds are
important for differences in disease propensities, and that the single cell atlas will help identify
novel pathologic populations and genetic programs. To determine if these identified
transcriptomic differences play a role in adult diseases, we then asked whether the differentially
regulated genes were enriched in human signals of genetic diseases. We again focused our
attention on the difference between ascending and descending thoracic aortic segments, which
have a known strong propensity for aortic aneurysms[77,78]. While these disease segments are
subject to changes in flow, there was relatively few differences between the global endothelial
cell expression of cells from these two vessel beds (Table S4). However, there were
significantly more transcriptomic differences between fibroblast and SMC from these different
connected arterial segments. Most notably, there were a large number of differentially regulated
genes in the fibroblast population that have known roles in inherited aortopathies[79]. A number of
mendelian aortopathies involve haploinsufficiency of critical ECM components, including fibrilin
1 (*FBN1*), lysyl oxidase (*LOX*), microfibril associated protein 4 (*MFAP4*), microfibril associated
protein 5 (*MFAP5*), and collagen 3A1 (*COL3A1*). Pathological decreased expression of these
genes is known to be causal for thoracic aortic aneurysms, with a strong preference for affecting
the root and ascending aorta. Interestingly, the expression of each of these genes is
significantly lower in the ascending than the descending aorta (Figure 7A). Closer examination
revealed that differences in expression were seen nearly exclusively in the adventitial fibroblasts,
and not in the medial SMC (Figure 7B). Conversely, *PRKG1* is a gene whose gain of function is
known to increase the risk of aneurysm[80-82], also has higher expression in the aortic root and
ascending compared to descending (Figure 7C).
To determine if other arterial segment specific vascular traits showed segregation of
genetic signals, we subsequently performed the same analysis on common genetic variants
associated with human aortic diameter. Most GWAS lead variants are located in non-coding

regions of the genome, likely exerting its effect on gene dosages. Thus, GWAS "hits" enrich for
genes whose dose are likely to contribute to pathology, with the added benefit that GWAS "hits"
do not hold bias towards known pathways. We leveraged recently published GWAS on aortic
diameter[83], and asked how often GWAS genes for aortic diameter are differentially expressed in
different populations of cells in aneurysm prone (i.e. ascending) vs resistant (i.e. descending)
segments. For those within the non-coding region of the genome, we used the putative target
gene suggested in the original manuscript, which mostly reflected the closest gene. From the list
of GWAS genes, we then determined how often these dose sensitive genes were differentially
expressed at the individual cell level in ascending and descending aorta. After Bonferroni
correction for multiple hypothesis testing, our analysis revealed 31 diameter sensitive genes are
differentially expressed in ascending vs descending fibroblasts, and 18 diameter sensitive
genes are differentially expressed in smooth muscle cells, and 5, 3, and 4 genes are
differentially expressed in microvasculature, macrophage, and endothelial cells, respectively
(Table S5). These analyses suggest a large proportion of segment-specific modifiers of aortic
propensity for aneurysm act through fibroblast and SMC, with surprisingly little transcriptomic
differences in the endothelial cells. These findings confirm the known role of SMC in aneurysms,
while suggesting a prominent unrecognized role of AdvFib in the same disease process.
We subsequently performed the same analysis in atherosclerotic disease and observed
a similar pattern. Coronary artery and carotid arteries are known to be more prone to
atherosclerosis whereas descending thoracic aorta is less so[84]. Thus, these vessel beds are
used for the atherosclerosis disease propensity comparisons to identify causal cells and causal
genes for the reasons for the differences in disease propensity. A significant subset of common
variant associated genes for atherosclerotic disease (in this case, coronary artery disease) are
differentially expressed in the different population of cells in the vessel wall (Figures 7D, 7E,
S9A). Our analysis revealed 45 CAD GWAS genes differentially expressed in fibroblast, 30 in
SMC, 11 in microvasculature, 15 in Endo1, 7 in Endo2, and 27 in macrophages (Table S6).
Functional analysis of enriched genes in each population of cells revealed angiogenesis related
biological processes are significantly enriched in SMC, and PDGF and VEGF signaling enriched
in the fibroblast population (Figure 7F). These analyses suggest that segment specific
differences in vascular disease propensities appears to be mostly encoded in fibroblast and
SMC, and much less so in endothelial cells.
480
Sex-specific differences
At a population level, there is a distinct difference in the susceptibility for vascular diseases
between patients of different biological sex[85-88]. Coronary artery disease and Marfan syndrome,
for example, are known to have worse vascular manifestation in men compared to women[89,90],
whereas Takayasu arteritis or spontaneous coronary artery dissections are much more common
in women than men[91,92]. The underlying cause of these sex specific differences is poorly
understood. While many of the causes of the sex specific differences may be related to
numerous systemic and other confounding factors that are unrelated to the cell biology at the
vessel wall, this dataset containing an equal number of men versus women, offers an
opportunity to explore potential differences between male and female arteries. We performed
differential expression analysis comparing different cellular populations in our dataset. This
analysis revealed different cell-type specific transcriptomes between male and female sex.

Unsurprisingly, many sex chromosome specific genes, such as *XIST*, were differentially
regulated in all cells. The analysis revealed 119, 176, 201, 79, 196, 281, 172 sex-specific
differentially expressed (DE) genes between SMC, fibroblast, macrophage, Endo1,
microvasculature, Endo2, and lymphocytes respectively (Table S7). Unlike cross-segmental
identities that appears to lie more in fibroblast and SMC, sex-specific differences were seen
across all cell types and are more profound in endothelial and immune cells. This parallels the
observations of sex specific differences observed in incidence of vasculitis and vascular
endothelial dysfunction. Sex differences in different populations appears to govern different
aspects of vascular biology and different diseases. Pathway analysis with the GO biological
process performed on each of these DE genes revealed distinct biological process enrichments
(Table S8). Different cell type DE gene overlap with OMIM data revealed distinct vascular
disease associations with known sex preferences (i.e. SMC association with Ehler-Danlos, and
EC association with macular degeneration) (Table S9).
506
Of mice and men
Our dataset enables us to compare cellular transcriptomes and cell populations of healthy
human arteries with murine ones. One of the biggest differences observed between human and
murine arteries is the presence of an intima at baseline. It is well-established that even a healthy
human has a significant layer of intimal cells as early as the second or third decade of life[93-95].
This is well reflected in this dataset. While healthy murine vessels do not contain a population
contiguous with the fibroblast population[28-30], we see a population of cells in healthy human
vessels that appears to be correspond to "phenotypically modulated smooth muscle cells" in
mice (Figure S10A). These modulated SMC are found almost exclusively in the human arterial
intima, and appear to comprise the majority of healthy intimal cells (Figure 2H). Beyond the
differences in the intimal cells that exist in the resting vessel, a comparison of the human and
mouse single cell RNA datasets processed in a similar manner reveals notable differences in a
few cellular populations. There is a significantly larger population of microvascular cells,
associated with the vaso-vasorum in human arteries compared to murine tissue. This finding is
expected given the increased thickness of human arterial tissues. As a result, the majority of
endothelial cells isolated from human arteries actually lines the microvasculature that supplies
the vessels instead of those on the luminal surface. In addition, there appears to be two
transcriptionally distinct subsets of endothelial cells in the human data, whereas the endothelial
populations appear less distinct in mice. While heterogeneity in endothelial cells in murine
arteries has also been described, no previously described population of endothelial cells in mice
matches the "Endo2" population found in humans[28,29,35].
A larger number of immune cells (both monocyte/macrophages and lymphocytes) are
present in donor human vasculature without overt disease compared to young/healthy mice.
This may reflect subclinical disease in these donors after decades of hyperlipidemia and
hypertension or rather fundamental differences between murine and human vasculature[29,30].
Lastly, while pro-calcific progenies of the SMC often accompany intimal formation or other
vascular damage in murine models, the same population/markers (*COL2A1*, *IBSP*) are not
found in healthy human arteries with extensive intima (Figure S10B), which may suggest a
different process causing calcification in mice compared to humans.
536

Cell type- and Segment- Specific Non-coding Transcriptome

537
While existing analysis of scRNAseq data has mostly focused on protein coding genes, there is
an increasing recognition that non-coding transcripts such as long non-coding RNAs (lncRNAs)
play an important role in health and disease[96-98]. Taking advantage of the fact that a significant
portion of lncRNAs is poly-adenylated and thus captured by the 10x platform, we asked whether
lncRNAs also follow segmental and cell-type specific expression by aligning these sequencing
data to a library of known lncRNAs. Given the library preparation of 10x does not allow for
strand specific capture, one cannot readily determine if a read is from the coding or antisense
strand of a transcript. To remove any potential confounding reads from protein coding genes,
we leveraged a reference lncRNA transcriptome that eliminated any segments of overlap with
any protein coding mRNAs. We will refer to this as the sc-lncRNA transcriptome.

This dataset contains 53,247 unique non-coding transcripts that are found in the
vascular cells (compared to 18,088 protein coding genes). Despite our stringent filtering, on
average, ~1601 (median 1284) non-coding transcripts were detected in each cell, encoding an
average of ~397 (median 323) lncRNAs per cell. This represents ~10% in number of genes or
reads compared to the protein coding genome. Differential gene expression analysis of these
data found that a significant portion of the sc-lncRNA transcriptome also exhibited cell-type and
segmental specific expression (Figures 8A, 8B).

Employing these data, we investigated how much of the cell type specific identities are
encoded in this limited subset of the non-coding transcriptome. To accomplish this, we
performed unbiased clustering of the sc-lncRNA transcriptome using standard methods, along
with UMAP based visualization. Utilizing no protein coding gene data, analyzing sc-lncRNA
transcriptome alone, standard unbiased expression clustering approach could readily segregate
the dataset into cell type specific clusters that recapitulate those identified using the entire
transcriptome (Figure 8C). Comparison of cell type identities assigned via unsupervised low-
resolution clustering to those identified by protein coding transcriptome demonstrated that ~92%
of the cells were clustered in identical ways using sc-lncRNA data alone without any protein
coding transcripts. The cells exhibiting discordance were primarily situated at the interfaces
between distinct cell types as visualized on the UMAP (Figure S11A). Hierarchical clustering
based on sc-lncRNA data itself in SMC and fibroblast could recapitulate the developmental
origin (Figure 8D). This suggests that a significant portion of the identities of each cell types,
behavior, and embryological origin are embedded within the non-coding transcriptome.

The majority of common variants that are identified in humans that carry heritable risk of
vascular diseases lie within the non-coding region of the genome[99-101]. One hypothesis has
been that some of these non-coding regions of the genome may overlap with functional
lncRNAs and alter their function. To test this hypothesis, we asked whether these vascular cell
specific lncRNAs enriched for genome-wide significant GWAS SNPs for different vascular
conditions. Interestingly, there appears to be a significant enrichment of GWAS SNPs for
coronary artery diseases that lie within the exons of lncRNAs compared to expected (18
lncRNAs with 158 MVP SNPs) (Figure 8E, Table S10) [(cite MVP)]. We subsequently
intersected the differentially expressed lncRNAs with the loci from the GWAS catalog, revealing
a significant intersection of these lncRNAs with loci associated with coronary artery disease
(CAD) and heart disease (Figure S11B). This finding is consistent with results obtained with
analysis of the Million Veterans Program CAD GWAS [100].

581
582
DISCUSSION

By performing cellular transcriptomic and spatial localization of donor arteries across a variety of
arterial segments, we created a resource for the vascular research community to better
understand the cellular composition and distribution of gene expression programs across
human arteries. We identified major cellular populations characterized not by individual markers,
but by entire transcriptomic profiles and spatial localizations. We contribute to the existing
literature of single cell characterization of human arteries by improving optimal digestion of
highly fibrous arterial tissue to obtain high quality single cell RNAseq data superior to the
cellular populations and ratios observed by histology[28,30]. These techniques also enabled us to
achieve deeper characterization of vascular cell transcriptomes, including lowly expressed
transcription factors, to better elucidate their biology. We hope this dataset can serve as a
useful reference to aid the cardiovascular research community in future single cell
characterization of vascular conditions.

596    Utilizing entire cellular transcriptomes instead of specific marker genes to define cellular
heterogeneity, this dataset both validates and challenges existing approaches to study vascular
diseases in murine and cellular models. One immediate benefit of this approach is the ability to
identify specific markers empirically in an unbiased manner for populations of cells that are
transcriptomically distinct. This large number of cell-type specific markers can then be utilized to
enable other vascular discoveries, including generating more specific murine Cre-lox models to
understand individual cell populations. Furthermore, our data can help reconcile seemingly
paradoxical findings of existing cellular and lineage tracing models. For example, our analyses
have revealed surprising heterogeneity and limitations of many commonly used vascular
population markers (*RGS5*, *TAGLN*, *GLI1*, *ACTA2* etc), which may explain some of the
contradictory findings in animal models that arise from modification using different "marker"
genes[102]. By combining single cell RNAseq data with complementary spatial transcriptomic
techniques, we were able to localize transcriptomically unique populations that reside in
different parts of the vessel wall. This includes identification of multiple "smooth muscle marker"
positive populations, and multiple subsets of endothelial cells with distinct transcriptomes and
localizations. This extensive heterogeneity that exists in healthy vessels may help explain
conflicting findings that sometimes occur from in vitro studies of primary human arterial cells, as
in vitro "arterial SMC" or "arterial endothelial" cells using canonical markers may be isolated
from one or multiple populations of cells. Our dataset may also be used to deconvolute the
population(s) of vascular endothelial cells or SMC[103,104].

616    One of our main goals with this dataset was to identify underlying differences in the
biology and disease propensities across different human arterial segments. Current
understanding of such differences mainly attributes them to flow characteristics and
hemodynamics[105-107]. Indeed, turbulent flow, laminar flow, and static flow are known to create
significant alterations in transcriptomic profiles of endothelial cells that are relevant to
diseases[108,109]. On the other hand, numerous pieces of evidence also exist suggesting that
developmental influences may alter vascular disease propensity[110-112], such as the frequently
observed changes in atherosclerotic disease lesion quality and severity near junctions of
vessels segments deriving from distinct developmental origins[113]. This nature vs nurture debate

of vascular disease remains unsettled. Our analysis of healthy vessels prior to initiation of
disease, yet still subjected to decades of differential flow patterns, reveals relatively few
differences between the transcriptomes of endothelial cells across different vascular segments.
However, adventitial fibroblasts and medial smooth muscle cells are clearly responsible for the
main transcriptomic differences across different vascular beds. Differentially expressed genes in
these populations of cells enrich for both common and rare variants of vascular diseases.
Furthermore, the regulatory regions of these differentially regulated genes significantly enrich for
transcription factor binding sites of developmentally critical master regulators. These findings
suggest that developmental origins of these vessels play a large role in different disease
propensities at least equally and maybe more so than decades of differences in vascular flow
and pressure. Developmental master regulators may therefore possess key pathogenic roles.
Given the clinical observation of strong associations of different arterial segments for diseases
within the same person[5,78,93,114], modulation of these developmental master regulators' function
may be sufficient to alter disease development or progression. Lessons learned from
developmental biology and additional discoveries aimed to better understand these factors'
exact functions are attractive targets for future therapies.
One of the largest populations of cells identified from this dataset are adventitial
fibroblasts. This population of cells carries remarkable heterogeneity across vascular segments
with different disease predilections, including a large number of genetic signals for aneurysmal
diseases and atherosclerosis. Yet, this population of cells has traditionally not been the focus of
pathogenesis of vascular diseases. Further evaluation of this population of cells and their roles
in vascular pathologies are warranted.
Analysis of the human endothelial population reveals significant heterogeneity as
previously demonstrated in murine vascular single cell analysis. However, unlike murine
models[35,50], the majority of human endothelial cells in the vessel walls reside in the
microvasculature of the adventitia. Consistent with distinct developmental origins and early
arterial/venous/lymphatic endothelial fate commitment demonstrated in animal
models[14,15,18,35,50,54,55], the most significant determinant of the endothelial transcriptome is the
type of vessel it covers. A distinct population of relatively rare endothelial cells is located at the
luminal surface of the great arteries. The lack of an equivalent endothelial cell population in
murine single cell vascular data suggests the function of this endothelial population may not be
accurately measured in murine models. Given the enrichment of this population of cells for
distinct disease-related transcriptomic programs, further dedicated studies of this population of
cells may yield distinct insight into their pathophysiological function.
Murine disease models with SMC lineage tracing have identified "de-differentiated" and
"phenotypically-modulated" SMC as key pathogenic cells in a variety of vascular
conditions[17,28,36,37]. The importance of this population of cells is well-established through a
variety of lineage tracing and manipulation models of different diseases[28,41,115,116]. However, in
macroscopically healthy vessels as obtained in these studies, an apparent de-differentiated
SMC subset is universally present across all vascular segments and populates the intima. This
population of cells expresses a similar global transcriptomic program as those SMC-lineage
traced cells that have lost their contractile genes in murine models. This potentially reflects the
presence of subclinical diseases in all these donors given years of relative hyperlipidemia due to
exposure to a western diet[117]. Alternatively, another explanation for this observation is that SMC

phenotypic modulation itself is part of the normal physiological spectrum and not necessarily
pathologic. Instead, the specific ways which seemingly similar SMC populations phenotypically
modulate in different ways may govern the quality and burden of disease. This notion is further
supported by the fact that there appears to be particularly large global transcriptomic differences
in the "modulated" /intimal SMC population observed across the vascular beds, more so than
"contractile" smooth muscle cells. This same population of cells also carries more genetic
signals of vascular diseases than other populations. Existing data are unfortunately inadequate
to discern these two possibilities. Additional characterization of a larger number of healthy and
diseased vascular tissues at different ages may help elucidate the answer.
678   Lastly, while the focus of pathogenic factors for vascular disease is often on protein
coding genes[83,99-101], analysis of non-protein-coding transcriptomes of the vascular beds reveals
surprisingly rich information and heterogeneity. Global analysis of lncRNAs alone can readily
identify the main vascular wall cell types and differentially expressed lncRNA across vascular
beds disproportionally enrich for non-coding variants that influence vascular disease risk. While
the analysis is preliminary, the enrichment of genetic signals suggests these differentially
enriched lncRNAs may contain critical regulators of disease risk.
685

ACKNOWLEDGMENTS

This work is made possible by the generosity of our tissue donors who donated their organs to save others' lives and expand our scientific understanding of the human vasculature. This work was supported by Human Cell Atlas grant (ZF2019-002437) from the Chan Zuckerberg Foundation (T.Q., P.C); National Institutes of Health grants K08HL153798 (P.C.), K08HL153798-S1(HuBMAP)(P.C.), R01HL171045 (T.Q), R01HL134817 (T.Q.), R01HL139478 (T.Q)., R01HL156846 (T.Q.), R01HL151535 (T.Q.), R01HL158525 (T.Q.), UM1 HG011972 (T.Q.), U01HG011762 (T.Q.), R01HL151535 (J.B.K), F32HL165854 (B.T.P), F32HL165819 (D.Y.L.); American Heart Association 23SCISA1144703(P.C), 24SCEFIA1248386 (P.C.), 20CDA35310303 (P.C.), 24CDA1272805 (B.T.P.); the Tobacco-Related Disease Research Program T32IR5352 (J.B.K), T32IR5240 (J.B.K);

AUTHOR CONTRIBUTIONS

P.C., T.Q., and A.K conceived and supervised the study. Q.Z., D.S., and P.C performed bioinformatic analyses. P.C., A.P., W.G., A.D., C.W., W.J., A.B. and R.W. performed single cell captures, spatial transcriptomics, histology, and IHC. D.Y.L., H.S., and R.W. helped analyses. A.P., A.D., R.S. L.L. M.F. performed human tissue collection, D.Y.L., Y.R., T.N., R.S., B.T.P., J.P.M., M.W., A.B., M.I., R.K., L.L., helped conducting experiments. P.C., T.Q., and Q.Z. interpreted data and wrote the manuscript. R.W., M.F., and J.B.K contributed to experiments and manuscript. The authors read and approved the final manuscript.

DECLARATION OF INTERESTS

A.K. is on the scientific advisory board of SerImmune, AINovo, TensorBio, Arcadia, Inari and OpenTargets; and has financial stake in Illumina, DeepGenomics, Immunai and Freenome.

SUPPLEMENTAL INFORMATION

Figures S1–S11, Word file containing additional figure.
Figure S1. Single cell RNA and spatial Slide-Seq of human arteries, related to Figure 1.
Figure S2. Smooth muscle cells lineage, related to Figure 2.
Figure S3. Fibroblast in different embryonic origins, related to Figure 3.
Figure S4. Endothelial cell populations across vessels, related to Figure 4.
Figure S5. Macrophages and lymphocytes.
Figure S6. Determinants of arterial identity.
Figure S7. The roles of embryonic origin in different cell types, related to Figure 5.
Figure S8. Vascular *HOX* gene expression patterns, related to Figure 6.
Figure S9. GWAS genes enrichment in different cell types, related to Figure 7.
Figure S10. Smooth muscle cell modulation in human arteries.
Figure S11. Concordant of lncRNA cell type annotation and GWAS catalog enrichment, related to Figure 8.
Table S1-S11. Excel files containing additional data too large to fit in a PDF.
Table S1. Differentially expressed genes between these two clusters of SMC populations, related to Figure 2.
Table S2. Specific markers defined adventitial fibroblast population, related to Figure 3.

742
743

FIGURE LEGENDS

Figure 1. Single cell RNA and spatial landscape of human arteries.
(A) Schematic diagram of the 8 distinct human arterial sites collection strategy.
(B) Workflow of the experimental design and the sample processing for scRNA-seq and Slide-
seq.
(C) RNA expression of existing canonical markers of each cell type population.
(D) UMAP visualization and clustering of cells labeled by cell type.
(E) Spatial dim plot visualization of the right coronary artery (RCA), annotated by cell type (left).
The right panel presents a split view, illustrating the positions of individual cell types on the
same slide.
See also Figure S1.

Figure 2. Smooth muscle cells of great arteries.
(A) RNA expression of mature smooth muscle contractile genes *MYOCD*, *CNN1*, *MYH11*,
ACTA2 and *TAGLN*, shown on the UMAPs.
(B) Immuno-fluorescence staining of CNN1 (green) and DAPI (blue) in ascending aorta.
(C) UMAP showing the *RGS5* RNA expression level.
(D) The differentially expressed genes between the two main clusters of SMC population,
ranked by the average log2 fold change level.
(E) Dot plot showing the z scores of enriched biological processes in the differentially expressed
gene between the two clusters of SMC population.
(F) The UMAPs showing the RNA expression of the unique makers in adventitial
microvasculature cell cluster (left panels). RNAscope validated the expression of these markers
with specific probes (right panels).
(G) RNA expression of medial and intimal smooth muscle genes *MYH11*, *FN1* and *TNFRSF11B*,
shown on the UMAPs.
(H) Spatial localization of the cells expressed *MYH11* and *FN1* in RCA (left, middle). And the
RNAscope staining of *TNSRSF11B* expression (right).

See also Figure S2 and Table S1.
774
Figure 3. Adventitial fibroblast in human arteries.
(A) The scores derived from the expression levels of collagen (middle) and extracellular matrix
genes (right) depicted on UMAP plots were analyzed alongside cell type annotations (left).
(B) *PI16* RNA expression levels on RNAseq UMAP plot (left), Slide-seq dim plot alongside the
fibroblast localization in RCA (middle), and the RNAscope staining with probes (right).
(C) The differentially expressed genes between fibroblast cells and all other cell types, plotted
by their percent of expressed cells and the average log2 fold changes.
(D) The violin plots illustrate the specific expression of the genes *FBN1*, *TGFBR2*, *LOX*, *FBLN1*,
and *MFAP5* within the adventitial fibroblast population.
See also Figure S3 and Table S2.
785
Figure 4. Distinct endothelial cell populations across vessels.
(A) Two large distinct clusters of endothelial cells shown on the UMAP.
(B) Endothelial cells identified by the canonical marker *CDH5* RNAscope staining (left), and the
endothelial cells localization on the Slide-seq dim plot of aorta (right).
(C) The sub-clustering of endothelial 1 population.
(D) The UMAP (top panels) and violin (bottom panels) plots showing the expression of cluster-
specific genes *ACKR1* (left panels) and *HEY1* (right panels) in endothelial1 cells.
(E) The expression of cluster-specific genes *ACKR1* (left) and *HEY1* (right) are stained by
RNAscope probes in ascending aorta.
(F) The UMAP (left panels) and violin (right panels) plots showing the expression of SMC genes
TAGLN and *ACTA2* (left panels), or lymphatic genes *PROX1* and *LYVE1* (right panels) in
endothelial1 cells.
(G) Heatmap showing the z scores of *HOX* genes expression in each of the vessel segments in
both endothelial 1 and 2.
(H) Endothelial 2 specific genes *ITLN1* and *NRG1* expression levels on RNAseq UMAP plot.
(I) RNAscope staining of *ITLN1* gene expression ascending aorta.
(K) The co-staining of *ITLN1* (blue) and *CDH5* (red) expression in ascending aorta.
(L) Dot plot showing the z scores of enriched biological processes in the differentially expressed
gene between endothelial 1 and 2.
See also Figure S4 and Table S3.
806
Figure 5. Prominent roles of embryonic origin in adult vascular identity.
(A) The heterogeneity of each population of cells across the different vascular beds on UMAP.
(B) A split view UMAP of the segmental origins presents the heterogeneity of each cell types.
(C) Heatmap showing the z scores of *HOX* genes expression in each of the vessel segments in
both SMC and fibroblast.
(D) Dendrogram Clustering using Euclidean distance calculated by all expressed genes
showing the correlation of each segment with its anticipated embryonic lineage within fibroblasts.
(E) Heatmap showing the percentage of each segment depicted within the distinct UMAP
clusters.

(F) Violin plots showing the differential expression patterns of transcription factors *ISL1*, *HAND2*,
MEDF2C, and *HOX* genes in fibroblast of ascending and descending sites.
(G) Enrichment of ISLET1, HAND2, MEF2C, and HOX binding motifs on the promoter regions of
differentially expressed genes between ascending and descending fibroblasts, with
macrophages serving as controls.
See also Figure S7.
822
Figure 6. Adult vascular *HOX* gene expression patterns.
(A) Heatmap showing the log10 normalized expression values of *HOX* genes across the
different vascular segment in fibroblast, SMC, endothetial1, endothelial2, microvasculature, or
macrophage.
(B) Dendrogram clustering using Euclidean distance calculated by all *HOX* genes showing the
correlation of each segment with its anticipated embryonic lineage in SMC (bottom) or
fibroblasts (top).
(C) The heatmap illustrates the Pearson correlation coefficients among all arterial segments,
derived from the expression values of all *HOX* genes. These segments are clustered according
to Euclidean distance in SMC (left) or fibroblast (right).
See also Figure S8.
834
Figure 7. Enrichment of vascular disease variants in segmental transcriptomic
differences.
(A) Dot plot showing the differential expression patterns of mendelian aortopathies genes *FBN1*,
FBNL1, *LOX*, *MFAP4*, *MFAP5*, and *COL3A1* genes in fibroblast of ascending and descending
sites.
(B) Bar plot showing the differential expression significances of these mendelian aortopathies
genes *FBN1*, *FBNL1*, *LOX*, *MFAP4*, *MFAP5*, and *COL3A1* between ascending and descending
sites in fibroblast.
(C) Violin plots showing the differential expression patterns of aneurysm risk in fibroblast (left) or
SMC (right) of ascending and descending sites.
(D) The bar graph displays the count of Million Veteran Program (MVP) GWAS genes that are
differentially expressed in each cell type when comparing the coronary artery and carotid with
descending segments.
(E) Genetic association between the MVP GWAS genes and the differentially expressed genes
across the coronary artery, carotid, and descending segments within each cell type, ranked by
the number of overlapping genes.
(F) Heatmap showing the z scores of fold enrichment of biological processes in the differentially
expressed genes across the coronary artery, carotid, and descending segments within each cell
type.
See also Figure S9 and Table S4, S5, S6.
855
Figure 8. Cell type and segment specific non-coding transcriptome.
(A) Heatmap showing the z scores of non-coding RNAs expression in each of the cell types.
(B) Heatmap showing the z scores of non-coding RNAs expression in each of the vascular
segments.

(C) UMAP visualization and clustering of cells using the sc-lncRNA transcriptome, annotated
based on either whole transcriptome-derived cell types (left) or lncRNA-derived cell type
identification (right).
(D) Dendrogram clustering using Euclidean distance calculated by all expressed lncRNAs
showing the correlation of each segment with its anticipated embryonic lineage in SMC (top) or
fibroblasts (bottom).
(E) The dot plot (top) illustrating the MVP SNPs positioned directly within the loci of differentially
expressed lncRNAs. The violin plot displays the expression levels of these intersected lncRNAs.
See also Figure S11 and Table S10.
869

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

● **KEY RESOURCES TABLE**

● **RESOURCE AVAILABILITY**

○ Lead contact

○ Materials availability

○ Data and code availability

● **SUBJECT DETAILS**

○ Human subjects

● **METHOD DETAILS**

○ Human Tissue Collection

○ Histology

○ Single-Cell RNA Sequencing

○ Analysis of Single-Cell Sequencing Data

○ Spatial Slide-Seq

○ Analysis of Spatial Slide-Seq Data

○ Functional and Motif Analyses

○ Correlation Analyses

○ GWAS Enrichment Analyses

○ Cell-cell Communications

○   Lnc-RNA Analyses

●   **QUANTIFICATION AND STATISTICAL ANALYSIS**

REFERENCES

1. Tsao, C.W., Aday, A.W., Almarzooq, Z.I., Anderson, C.A.M., Arora, P., Avery, C.L., Baker-Smith, C.M., Beaton, A.Z., Boehme, A.K., Buxton, A.E., et al. (2023). Heart Disease and Stroke Statistics-2023 Update: A Report From the American Heart Association. Circulation *147*, e93-e621. 10.1161/CIR.0000000000001123.

2. Cooke, J.P., and Meng, S. (2020). Vascular Regeneration in Peripheral Artery Disease. Arterioscler Thromb Vasc Biol *40*, 1627-1634. 10.1161/ATVBAHA.120.312862.

3. De Backer, J., Bondue, A., Budts, W., Evangelista, A., Gallego, P., Jondeau, G., Loeys, B., Pena, M.L., Teixido-Tura, G., van de Laar, I., et al. (2020). Genetic counselling and testing in adults with congenital heart disease: A consensus document of the ESC Working Group of Grown-Up Congenital Heart Disease, the ESC Working Group on Aorta and Peripheral Vascular Disease and the European Society of Human Genetics. Eur J Prev Cardiol *27*, 1423-1435. 10.1177/2047487319854552.

4. Donato, A.J., Machin, D.R., and Lesniewski, L.A. (2018). Mechanisms of Dysfunction in the Aging Vasculature and Role in Age-Related Disease. Circ Res *123*, 825-848. 10.1161/CIRCRESAHA.118.312563.

5. Poredos, P., Poredos, P., and Jezovnik, M.K. (2018). Structure of Atherosclerotic Plaques in Different Vascular Territories: Clinical Relevance. Curr Vasc Pharmacol *16*, 125-129. 10.2174/1570161115666170227103125.

6. DeBakey, M.E., Lawrie, G.M., and Glaeser, D.H. (1985). Patterns of atherosclerosis and their surgical significance. Ann Surg *201*, 115-131. 10.1097/00000658-198502000-00001.

7. Morisaki, T., and Morisaki, H. (2016). Genetics of hereditary large vessel diseases. J Hum Genet *61*, 21-26. 10.1038/jhg.2015.119.

8. Loeys, B.L., Schwarze, U., Holm, T., Callewaert, B.L., Thomas, G.H., Pannu, H., De Backer, J.F., Oswald, G.L., Symoens, S., Manouvrier, S., et al. (2006). Aneurysm syndromes caused by mutations in the TGF-beta receptor. N Engl J Med *355*, 788-798. 10.1056/NEJMoa055695.

9. Kim, J.B., Zhao, Q., Nguyen, T., Pjanic, M., Cheng, P., Wirka, R., Travisano, S., Nagao, M., Kundu, R., and Quertermous, T. (2020). Environment-Sensing Aryl Hydrocarbon Receptor Inhibits the Chondrogenic Fate of Modulated Smooth Muscle Cells in Atherosclerotic Lesions. Circulation *142*, 575-590. 10.1161/CIRCULATIONAHA.120.045981.

10. Kondo, T., Nakano, Y., Adachi, S., and Murohara, T. (2019). Effects of Tobacco Smoking on Cardiovascular Disease. Circ J *83*, 1980-1985. 10.1253/circj.CJ-19-0323.

11. Maleszewski, J.J. (2015). Inflammatory ascending aortic disease: perspectives from pathology. J Thorac Cardiovasc Surg *149*, S176-183. 10.1016/j.jtcvs.2014.07.046.

12. Rajasekaran, K., Duraiyarasan, S., Adefuye, M., Manjunatha, N., and Ganduri, V. (2022). Kawasaki Disease and Coronary Artery Involvement: A Narrative Review. Cureus *14*, e28358. 10.7759/cureus.28358.

13. Yuan, S.M. (2018). Syphilitic aortic aneurysm. Z Rheumatol *77*, 741-748. 10.1007/s00393-018-0519-1.

14. Noden, D.M. (1989). Embryonic origins and assembly of blood vessels. Am Rev Respir Dis *140*, 1097-1103. 10.1164/ajrccm/140.4.1097.

15. Saha, M.S., Cox, E.A., and Sipe, C.W. (2004). Mechanisms regulating the origins of the vertebrate vascular system. J Cell Biochem *93*, 46-56. 10.1002/jcb.20196.

16. Leonard, E.V., Figueroa, R.J., Bussmann, J., Lawson, N.D., Amigo, J.D., and Siekmann, A.F. (2022). Regenerating vascular mural cells in zebrafish fin blood vessels are not

943   derived from pre-existing mural cells and differentially require Pdgfrb signalling for their
944   development. Development *149*. 10.1242/dev.199640.

945 17.   Miano, J.M., Fisher, E.A., and Majesky, M.W. (2021). Fate and State of Vascular
946   Smooth Muscle Cells in Atherosclerosis. Circulation *143*, 2110-2116.
947   10.1161/CIRCULATIONAHA.120.049922.

948 18.   Rasanen, M., Sultan, I., Paech, J., Hemanthakumar, K.A., Yu, W., He, L., Tang, J., Sun,
949   Y., Hlushchuk, R., Huan, X., et al. (2021). VEGF-B Promotes Endocardium-Derived
950   Coronary Vessel Development and Cardiac Regeneration. Circulation *143*, 65-77.
951   10.1161/CIRCULATIONAHA.120.050635.

952 19.   Sharma, B., Chang, A., and Red-Horse, K. (2017). Coronary Artery Development:
953   Progenitor Cells and Differentiation Pathways. Annu Rev Physiol *79*, 1-19.
954   10.1146/annurev-physiol-022516-033953.

955 20.   Red-Horse, K., Ueno, H., Weissman, I.L., and Krasnow, M.A. (2010). Coronary arteries
956   form by developmental reprogramming of venous cells. Nature *464*, 549-553.
957   10.1038/nature08873.

958 21.   Majesky, M.W. (2007). Developmental basis of vascular smooth muscle diversity.
959   Arterioscler Thromb Vasc Biol *27*, 1248-1258. 10.1161/ATVBAHA.107.141069.

960 22.   High, F.A., Lu, M.M., Pear, W.S., Loomes, K.M., Kaestner, K.H., and Epstein, J.A.
961   (2008). Endothelial expression of the Notch ligand Jagged1 is required for vascular
962   smooth muscle development. Proc Natl Acad Sci U S A *105*, 1955-1959.
963   10.1073/pnas.0709663105.

964 23.   Sato, Y. (2013). Dorsal aorta formation: separate origins, lateral-to-medial migration, and
965   remodeling. Dev Growth Differ *55*, 113-129. 10.1111/dgd.12010.

966 24.   Dobnikar, L., Taylor, A.L., Chappell, J., Oldach, P., Harman, J.L., Oerton, E., Dzierzak,
967   E., Bennett, M.R., Spivakov, M., and Jorgensen, H.F. (2018). Disease-relevant
968   transcriptional signatures identified in individual smooth muscle cells from healthy mouse
969   vessels. Nat Commun *9*, 4567. 10.1038/s41467-018-06891-x.

970 25.   Trigueros-Motos, L., Gonzalez-Granado, J.M., Cheung, C., Fernandez, P., Sanchez-
971   Cabo, F., Dopazo, A., Sinha, S., and Andres, V. (2013). Embryological-origin-dependent
972   differences in homeobox expression in adult aorta: role in regional phenotypic variability
973   and regulation of NF-kappaB activity. Arterioscler Thromb Vasc Biol *33*, 1248-1256.
974   10.1161/ATVBAHA.112.300539.

975 26.   Haimovici, H., Maier, N., and Strauss, L. (1959). Fate of aortic homografts in
976   experimental canine atherosclerosis. II. Study of fresh abdominal aortic implants into
977   abdominal aorta. AMA Arch Surg *78*, 239-245. 10.1001/archsurg.1959.04320020061010.

978 27.   Haimovici, H., and Maier, N. (1964). Fate of Aortic Homografts in Canine Atherosclerosis.
979   3. Study of Fresh Abdominal and Thoracic Aortic Implants into Thoracic Aorta: Role of
980   Tissue Susceptibility in Atherogenesis. Arch Surg *89*, 961-969.
981   10.1001/archsurg.1964.01320060029006.

982 28.   Wirka, R.C., Wagh, D., Paik, D.T., Pjanic, M., Nguyen, T., Miller, C.L., Kundu, R., Nagao,
983   M., Coller, J., Koyano, T.K., et al. (2019). Atheroprotective roles of smooth muscle cell
984   phenotypic modulation and the TCF21 disease gene as revealed by single-cell analysis.
985   Nat Med *25*, 1280-1289. 10.1038/s41591-019-0512-5.

986 29.   Cheng, P., Wirka, R.C., Kim, J.B., Kim, H.J., Nguyen, T., Kundu, R., Zhao, Q., Sharma,
987   D., Pedroza, A., Nagao, M., et al. (2022). Smad3 regulates smooth muscle cell fate and
988   mediates adverse remodeling and calcification of the atherosclerotic plaque. Nat
989   Cardiovasc Res *1*, 322-333. 10.1038/s44161-022-00042-8.

990 30.   Cheng, P., Wirka, R.C., Shoa Clarke, L., Zhao, Q., Kundu, R., Nguyen, T., Nair, S.,
991   Sharma, D., Kim, H.J., Shi, H., et al. (2022). ZEB2 Shapes the Epigenetic Landscape of
992   Atherosclerosis. Circulation *145*, 469-485. 10.1161/CIRCULATIONAHA.121.057789.

993   31.   Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-
994   cell RNA-seq data using regularized negative binomial regression. Genome Biol *20*, 296.
995   10.1186/s13059-019-1874-1.
996   32.   Soliman, H., Tung, L.W., and Rossi, F.M.V. (2021). Fibroblast and Myofibroblast
997   Subtypes: Single Cell Sequencing. Methods Mol Biol *2299*, 49-84. 10.1007/978-1-0716-
998   1382-5_4.
999   33.   Hume, D.A., Millard, S.M., and Pettit, A.R. (2023). Macrophage heterogeneity in the
1000   single-cell era: facts and artifacts. Blood *142*, 1339-1347. 10.1182/blood.2023020597.
1001   34.   Mae, M.A., He, L., Nordling, S., Vazquez-Liebanas, E., Nahar, K., Jung, B., Li, X., Tan,
1002   B.C., Chin Foo, J., Cazenave-Gassiot, A., et al. (2021). Single-Cell Analysis of Blood-
1003   Brain Barrier Response to Pericyte Loss. Circ Res *128*, e46-e62.
1004   10.1161/CIRCRESAHA.120.317473.
1005   35.   Kalucka, J., de Rooij, L., Goveia, J., Rohlenova, K., Dumas, S.J., Meta, E., Conchinha,
1006   N.V., Taverna, F., Teuwen, L.A., Veys, K., et al. (2020). Single-Cell Transcriptome Atlas
1007   of Murine Endothelial Cells. Cell *180*, 764-779 e720. 10.1016/j.cell.2020.01.015.
1008   36.   Muhl, L., Mocci, G., Pietila, R., Liu, J., He, L., Genove, G., Leptidis, S., Gustafsson, S.,
1009   Buyandelger, B., Raschperger, E., et al. (2022). A single-cell transcriptomic inventory of
1010   murine smooth muscle cells. Dev Cell *57*, 2426-2443 e2426.
1011   10.1016/j.devcel.2022.09.015.
1012   37.   Allahverdian, S., Chaabane, C., Boukais, K., Francis, G.A., and Bochaton-Piallat, M.L.
1013   (2018). Smooth muscle cell fate and plasticity in atherosclerosis. Cardiovasc Res *114*,
1014   540-550. 10.1093/cvr/cvy022.
1015   38.   Hu, Z., Liu, W., Hua, X., Chen, X., Chang, Y., Hu, Y., Xu, Z., and Song, J. (2021).
1016   Single-Cell Transcriptomic Atlas of Different Human Cardiac Arteries Identifies Cell
1017   Types Associated With Vascular Physiology. Arterioscler Thromb Vasc Biol *41*, 1408-
1018   1427. 10.1161/ATVBAHA.120.315373.
1019   39.   Crnkovic, S., Valzano, F., Fliesser, E., Gindlhuber, J., Thekkekara Puthenparampil, H.,
1020   Basil, M., Morley, M.P., Katzen, J., Gschwandtner, E., Klepetko, W., et al. (2022).
1021   Single-cell transcriptomics reveals skewed cellular communication and phenotypic shift
1022   in pulmonary artery remodeling. JCI Insight *7*. 10.1172/jci.insight.153471.
1023   40.   Armulik, A., Genove, G., and Betsholtz, C. (2011). Pericytes: developmental,
1024   physiological, and pathological perspectives, problems, and promises. Dev Cell *21*, 193-
1025   215. 10.1016/j.devcel.2011.07.001.
1026   41.   Shankman, L.S., Gomez, D., Cherepanova, O.A., Salmon, M., Alencar, G.F., Haskins,
1027   R.M., Swiatlowska, P., Newman, A.A., Greene, E.S., Straub, A.C., et al. (2015). KLF4-
1028   dependent phenotypic modulation of smooth muscle cells has a key role in
1029   atherosclerotic plaque pathogenesis. Nat Med *21*, 628-637. 10.1038/nm.3866.
1030   42.   Kramann, R., Schneider, R.K., DiRocco, D.P., Machado, F., Fleig, S., Bondzie, P.A.,
1031   Henderson, J.M., Ebert, B.L., and Humphreys, B.D. (2015). Perivascular Gli1+
1032   progenitors are key contributors to injury-induced organ fibrosis. Cell Stem Cell *16*, 51-
1033   66. 10.1016/j.stem.2014.11.004.
1034   43.   Guerrero-Juarez, C.F., Dedhia, P.H., Jin, S., Ruiz-Vega, R., Ma, D., Liu, Y., Yamaga, K.,
1035   Shestova, O., Gay, D.L., Yang, Z., et al. (2019). Single-cell analysis reveals fibroblast
1036   heterogeneity and myeloid-derived adipocyte progenitors in murine skin wounds. Nat
1037   Commun *10*, 650. 10.1038/s41467-018-08247-x.
1038   44.   Xie, T., Wang, Y., Deng, N., Huang, G., Taghavifar, F., Geng, Y., Liu, N., Kulur, V., Yao,
1039   C., Chen, P., et al. (2018). Single-Cell Deconvolution of Fibroblast Heterogeneity in
1040   Mouse Pulmonary Fibrosis. Cell Rep *22*, 3625-3640. 10.1016/j.celrep.2018.03.010.
1041   45.   Philippeos, C., Telerman, S.B., Oules, B., Pisco, A.O., Shaw, T.J., Elgueta, R., Lombardi,
1042   G., Driskell, R.R., Soldin, M., Lynch, M.D., and Watt, F.M. (2018). Spatial and Single-

Cell Transcriptional Profiling Identifies Functionally Distinct Human Dermal Fibroblast
Subpopulations. J Invest Dermatol *138*, 811-825. 10.1016/j.jid.2018.01.016.

46. Pedroza, A.J., Tashima, Y., Shad, R., Cheng, P., Wirka, R., Churovich, S., Nakamura, K.,
Yokoyama, N., Cui, J.Z., Iosef, C., et al. (2020). Single-Cell Transcriptomic Profiling of
Vascular Smooth Muscle Cell Phenotype Modulation in Marfan Syndrome Aortic
Aneurysm. Arterioscler Thromb Vasc Biol *40*, 2195-2211.
10.1161/ATVBAHA.120.314670.

47. Renner, S., Schuler, H., Alawi, M., Kolbe, V., Rybczynski, M., Woitschach, R.,
Sheikhzadeh, S., Stark, V.C., Olfe, J., Roser, E., et al. (2019). Next-generation
sequencing of 32 genes associated with hereditary aortopathies and related disorders of
connective tissue in a cohort of 199 patients. Genet Med *21*, 1832-1841.
10.1038/s41436-019-0435-z.

48. Sawada, H., Katsumata, Y., Higashi, H., Zhang, C., Li, Y., Morgan, S., Lee, L.H., Singh,
S.A., Chen, J.Z., Franklin, M.K., et al. (2022). Second Heart Field-Derived Cells
Contribute to Angiotensin II-Mediated Ascending Aortopathies. Circulation *145*, 987-
1001. 10.1161/CIRCULATIONAHA.121.058173.

49. Naito, H., Iba, T., and Takakura, N. (2020). Mechanisms of new blood-vessel formation
and proliferative heterogeneity of endothelial cells. Int Immunol *32*, 295-305.
10.1093/intimm/dxaa008.

50. Hennigs, J.K., Matuszcak, C., Trepel, M., and Korbelin, J. (2021). Vascular Endothelial
Cells: Heterogeneity and Targeting Approaches. Cells *10*. 10.3390/cells10102712.

51. Fischer, A., Schumacher, N., Maier, M., Sendtner, M., and Gessler, M. (2004). The
Notch target genes Hey1 and Hey2 are required for embryonic vascular development.
Genes Dev *18*, 901-911. 10.1101/gad.291004.

52. Dong, G., Huang, X., Xu, Y., Chen, R., and Chen, S. (2023). Mechanical stress induced
EndoMT in endothelial cells through PPARgamma downregulation. Cell Signal *110*,
110812. 10.1016/j.cellsig.2023.110812.

53. Li, Y., Lui, K.O., and Zhou, B. (2018). Reassessing endothelial-to-mesenchymal
transition in cardiovascular diseases. Nat Rev Cardiol *15*, 445-456. 10.1038/s41569-
018-0023-y.

54. Arroz-Madeira, S., Bekkhus, T., Ulvmar, M.H., and Petrova, T.V. (2023). Lessons of
Vascular Specialization From Secondary Lymphoid Organ Lymphatic Endothelial Cells.
Circ Res *132*, 1203-1225. 10.1161/CIRCRESAHA.123.322136.

55. Jalkanen, S., and Salmi, M. (2020). Lymphatic endothelial cells of the lymph node. Nat
Rev Immunol *20*, 566-578. 10.1038/s41577-020-0281-x.

56. Hedhli, N., Kalinowski, A., and K, S.R. (2014). Cardiovascular effects of neuregulin-
1/ErbB signaling: role in vascular signaling and angiogenesis. Curr Pharm Des *20*, 4899-
4905. 10.2174/1381612819666131125151058.

57. Odiete, O., Hill, M.F., and Sawyer, D.B. (2012). Neuregulin in cardiovascular
development and disease. Circ Res *111*, 1376-1385.
10.1161/CIRCRESAHA.112.267286.

58. Guclu-Geyik, F., Erkan, A.F., Ozuynuk, A.S., Ekici, B., and Coban, N. (2022). Val109Asp
polymorphism in Intelectin 1 gene is associated with coronary artery disease severity in
women. Turk Kardiyol Dern Ars *50*, 34-45. 10.5543/tkda.2022.21003.

59. Gu, N., Dong, Y., Tian, Y., Di, Z., Liu, Z., Chang, M., Jia, X., Qian, Y., and Zhang, W.
(2017). Anti-apoptotic and angiogenic effects of intelectin-1 in rat cerebral ischemia.
Brain Res Bull *130*, 27-35. 10.1016/j.brainresbull.2016.12.006.

60. Luo, X.L., Jiang, Y., Li, Q., Yu, X.J., Ma, T., Cao, H., Ke, M.X., Zhang, P., Tan, J.L.,
Gong, Y.S., et al. (2023). hESC-Derived Epicardial Cells Promote Repair of Infarcted
Hearts in Mouse and Swine. Adv Sci (Weinh) *10*, e2300470. 10.1002/advs.202300470.

61.   Menzel, J., di Giuseppe, R., Biemann, R., Wittenbecher, C., Aleksandrova, K., Pischon,
T., Fritsche, A., Schulze, M.B., Boeing, H., Isermann, B., and Weikert, C. (2016).
Omentin-1 and risk of myocardial infarction and stroke: Results from the EPIC-Potsdam
cohort study. Atherosclerosis *251*, 415-421. 10.1016/j.atherosclerosis.2016.06.003.
62.   Litvinukova, M., Talavera-Lopez, C., Maatz, H., Reichart, D., Worth, C.L., Lindberg, E.L.,
Kanda, M., Polanski, K., Heinig, M., Lee, M., et al. (2020). Cells of the adult human heart.
Nature *588*, 466-472. 10.1038/s41586-020-2797-4.
63.   Zhang, X., McDonald, J.G., Aryal, B., Canfran-Duque, A., Goldberg, E.L., Araldi, E., Ding,
W., Fan, Y., Thompson, B.M., Singh, A.K., et al. (2021). Desmosterol suppresses
macrophage inflammasome activation and protects against vascular inflammation and
atherosclerosis. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2107682118.
64.   Zhang, Q., Liu, Q., Feng, S., Li, X., and Jiang, Q. (2024). Non-Coding RNAs: Novel
Regulators of Macrophage Homeostasis in Ocular Vascular Diseases. Biomolecules *14*.
10.3390/biom14030328.
65.   Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.H., Myung, P.,
Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using
CellChat. Nat Commun *12*, 1088. 10.1038/s41467-021-21246-9.
66.   Becht, E., McInnes, L., Healy, J., Dutertre, C.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F.,
and Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using
UMAP. Nat Biotechnol. 10.1038/nbt.4314.
67.   Zhai, Z., Lei, Y.L., Wang, R., and Xie, Y. (2022). Supervised capacity preserving
mapping: a clustering guided visualization method for scRNA-seq data. Bioinformatics
38, 2496-2503. 10.1093/bioinformatics/btac131.
68.   Liu, J., and Vinck, M. (2022). Improved visualization of high-dimensional data using the
distance-of-distance transformation. PLoS Comput Biol *18*, e1010764.
10.1371/journal.pcbi.1010764.
69.   Ultsch, A., and Lotsch, J. (2022). Euclidean distance-optimized data transformation for
cluster analysis in biomedical data (EDOtrans). BMC Bioinformatics *23*, 233.
10.1186/s12859-022-04769-w.
70.   Gorski, D.H., and Walsh, K. (2000). The role of homeobox genes in vascular remodeling
and angiogenesis. Circ Res *87*, 865-872. 10.1161/01.res.87.10.865.
71.   Swift, M.R., and Weinstein, B.M. (2009). Arterial-venous specification during
development. Circ Res *104*, 576-588. 10.1161/CIRCRESAHA.108.188805.
72.   Barnes, R.M., Firulli, B.A., VanDusen, N.J., Morikawa, Y., Conway, S.J., Cserjesi, P.,
Vincentz, J.W., and Firulli, A.B. (2011). Hand2 loss-of-function in Hand1-expressing cells
reveals distinct roles in epicardial and coronary vessel development. Circ Res *108*, 940-
949. 10.1161/CIRCRESAHA.110.233171.
73.   Maruyama, K., Miyagawa-Tomita, S., Haneda, Y., Kida, M., Matsuzaki, F., Imanaka-
Yoshida, K., and Kurihara, H. (2022). The cardiopharyngeal mesoderm contributes to
lymphatic vessel development in mouse. Elife *11*. 10.7554/eLife.81515.
74.   Materna, S.C., Sinha, T., Barnes, R.M., Lammerts van Bueren, K., and Black, B.L.
(2019). Cardiovascular development and survival require Mef2c function in the
myocardial but not the endothelial lineage. Dev Biol *445*, 170-177.
10.1016/j.ydbio.2018.12.002.
75.   Crispino, J.D., Lodish, M.B., Thurberg, B.L., Litovsky, S.H., Collins, T., Molkentin, J.D.,
and Orkin, S.H. (2001). Proper coronary vascular development and heart
morphogenesis depend on interaction of GATA-4 with FOG cofactors. Genes Dev *15*,
839-844. 10.1101/gad.875201.
76.   Hubert, K.A., and Wellik, D.M. (2023). Hox genes in development and beyond.
Development *150*. 10.1242/dev.192476.

77.   Vapnik, J.S., Kim, J.B., Isselbacher, E.M., Ghoshhajra, B.B., Cheng, Y., Sundt, T.M., 3rd,
1144   MacGillivray, T.E., Cambria, R.P., and Lindsay, M.E. (2016). Characteristics and
1145   Outcomes of Ascending Versus Descending Thoracic Aortic Aneurysms. Am J Cardiol
1146   *117*, 1683-1690. 10.1016/j.amjcard.2016.02.048.
1147   78.   Huang, Y., Schaff, H.V., Bagameri, G., Pochettino, A., DeMartino, R.R., Todd, A., and
1148   Greason, K.L. (2024). Differential expansion and outcomes of ascending and
1149   descending degenerative thoracic aortic aneurysms. J Thorac Cardiovasc Surg *167*,
1150   918-926 e913. 10.1016/j.jtcvs.2022.03.032.
1151   79.   Chou, E.L., and Lindsay, M.E. (2020). The genetics of aortopathies: Hereditary thoracic
1152   aortic aneurysms and dissections. Am J Med Genet C Semin Med Genet *184*, 136-148.
1153   10.1002/ajmg.c.31771.
1154   80.   Isselbacher, E.M., Lino Cardenas, C.L., and Lindsay, M.E. (2016). Hereditary Influence
1155   in Thoracic Aortic Aneurysm and Dissection. Circulation *133*, 2516-2528.
1156   10.1161/CIRCULATIONAHA.116.009762.
1157   81.   Shalhub, S., Regalado, E.S., Guo, D.C., Milewicz, D.M., and Montalcino Aortic, C.
1158   (2019). The natural history of type B aortic dissection in patients with PRKG1 mutation
1159   c.530G>A (p.Arg177Gln). J Vasc Surg *70*, 718-723. 10.1016/j.jvs.2018.12.032.
1160   82.   Gago-Diaz, M., Blanco-Verea, A., Teixido, G., Huguet, F., Gut, M., Laurie, S., Gut, I.,
1161   Carracedo, A., Evangelista, A., and Brion, M. (2016). PRKG1 and genetic diagnosis of
1162   early-onset thoracic aortic disease. Eur J Clin Invest *46*, 787-794. 10.1111/eci.12662.
1163   83.   Pirruccello, J.P., Chaffin, M.D., Chou, E.L., Fleming, S.J., Lin, H., Nekoui, M., Khurshid,
1164   S., Friedman, S.F., Bick, A.G., Arduini, A., et al. (2022). Deep learning enables genetic
1165   analysis of the human thoracic aorta. Nat Genet *54*, 40-51. 10.1038/s41588-021-00962-
1166   4.
1167   84.   Khoury, Z., Gottlieb, S., Stern, S., and Keren, A. (1997). Frequency and distribution of
1168   atherosclerotic plaques in the thoracic aorta as determined by transesophageal
1169   echocardiography in patients with coronary artery disease. Am J Cardiol *79*, 23-27.
1170   10.1016/s0002-9149(96)00670-4.
1171   85.   Allais, G., Chiarle, G., Sinigaglia, S., Airola, G., Schiapparelli, P., and Benedetto, C.
1172   (2018). Estrogen, migraine, and vascular risk. Neurol Sci *39*, 11-20. 10.1007/s10072-
1173   018-3333-2.
1174   86.   Klouche, M. (2006). Estrogens in human vascular diseases. Ann N Y Acad Sci *1089*,
1175   431-443. 10.1196/annals.1386.032.
1176   87.   Lin, G.M., Yano, Y., and Hoshide, S. (2017). Sex Differences in the Association between
1177   Traditional Vascular Risk Factors and Subclinical Carotid Atherosclerosis in Taiwan. J
1178   Atheroscler Thromb *24*, 673-674. 10.5551/jat.ED068.
1179   88.   Liu, P.P., and Fukuoka, M. (2007). Sex hormones as novel risk biomarkers for
1180   atherosclerosis in peripheral vascular disease. J Am Coll Cardiol *50*, 1077-1079.
1181   10.1016/j.jacc.2007.05.031.
1182   89.   Yahagi, K., Davis, H.R., Arbustini, E., and Virmani, R. (2015). Sex differences in
1183   coronary artery disease: pathological observations. Atherosclerosis *239*, 260-267.
1184   10.1016/j.atherosclerosis.2015.01.017.
1185   90.   Nucera, M., Heinisch, P.P., Langhammer, B., Jungi, S., Mihalj, M., Schober, P., Luedi,
1186   M.M., Yildiz, M., and Schoenhoff, F.S. (2022). The impact of sex and gender on aortic
1187   events in patients with Marfan syndrome. Eur J Cardiothorac Surg *62*.
1188   10.1093/ejcts/ezac305.
1189   91.   Shimizu, Y., and Murohara, T. (2024). Takayasu Arteritis in Terms of Disease Duration
1190   and Sex Differences. Circ J *88*, 295-296. 10.1253/circj.CJ-23-0900.
1191   92.   Brundage, J.N., Mason, C.P., Wadsworth, H.A., Finuf, C.S., Nelson, J.J., Ronstrom,
1192   P.J.W., Jones, S.R., Siciliano, C.A., Steffensen, S.C., and Yorgason, J.T. (2022).

Regional and sex differences in spontaneous striatal dopamine transmission. J Neurochem *160*, 598-612. 10.1111/jnc.15473.

93. Crouse, J.R., 3rd, Craven, T.E., Hagaman, A.P., and Bond, M.G. (1995). Association of coronary disease with segment-specific intimal-medial thickening of the extracranial carotid artery. Circulation *92*, 1141-1147. 10.1161/01.cir.92.5.1141.

94. Nakashima, Y., Wight, T.N., and Sueishi, K. (2008). Early atherosclerosis in humans: role of diffuse intimal thickening and extracellular matrix proteoglycans. Cardiovasc Res *79*, 14-23. 10.1093/cvr/cvn099.

95. Milutinovic, A., Suput, D., and Zorc-Pleskovic, R. (2020). Pathogenesis of atherosclerosis in the tunica intima, media, and adventitia of coronary arteries: An updated review. Bosn J Basic Med Sci *20*, 21-30. 10.17305/bjbms.2019.4320.

96. Shi, H., Nguyen, T., Zhao, Q., Cheng, P., Sharma, D., Kim, H.J., Brian Kim, J., Wirka, R., Weldy, C.S., Monteiro, J.P., and Quertermous, T. (2023). Discovery of Transacting Long Noncoding RNAs That Regulate Smooth Muscle Cell Phenotype. Circ Res *132*, 795-811. 10.1161/CIRCRESAHA.122.321960.

97. Leung, A., and Natarajan, R. (2014). Noncoding RNAs in vascular disease. Curr Opin Cardiol *29*, 199-206. 10.1097/HCO.0000000000000054.

98. Simion, V., Haemmig, S., and Feinberg, M.W. (2019). LncRNAs in vascular biology and disease. Vascul Pharmacol *114*, 145-156. 10.1016/j.vph.2018.01.003.

99. Sollis, E., Mosaku, A., Abid, A., Buniello, A., Cerezo, M., Gil, L., Groza, T., Gunes, O., Hall, P., Hayhurst, J., et al. (2023). The NHGRI-EBI GWAS Catalog: knowledgebase and deposition resource. Nucleic Acids Res *51*, D977-D985. 10.1093/nar/gkac1010.

100. Djousse, L., Ho, Y.L., Nguyen, X.T., Gagnon, D.R., Wilson, P.W.F., Cho, K., Gaziano, J.M., and Program, V.A.M.V. (2018). DASH Score and Subsequent Risk of Coronary Artery Disease: The Findings From Million Veteran Program. J Am Heart Assoc *7*. 10.1161/JAHA.117.008089.

101. Nikpay, M., Goel, A., Won, H.H., Hall, L.M., Willenborg, C., Kanoni, S., Saleheen, D., Kyriakou, T., Nelson, C.P., Hopewell, J.C., et al. (2015). A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. Nat Genet *47*, 1121-1130. 10.1038/ng.3396.

102. Chakraborty, R., Saddouk, F.Z., Carrao, A.C., Krause, D.S., Greif, D.M., and Martin, K.A. (2019). Promoters to Study Vascular Smooth Muscle. Arterioscler Thromb Vasc Biol *39*, 603-612. 10.1161/ATVBAHA.119.312449.

103. Avila Cobos, F., Alquicira-Hernandez, J., Powell, J.E., Mestdagh, P., and De Preter, K. (2020). Benchmarking of cell type deconvolution pipelines for transcriptomics data. Nat Commun *11*, 5650. 10.1038/s41467-020-19015-1.

104. Wang, X., Park, J., Susztak, K., Zhang, N.R., and Li, M. (2019). Bulk tissue cell type deconvolution with multi-subject single-cell expression reference. Nat Commun *10*, 380. 10.1038/s41467-018-08023-x.

105. Gaudio, L.T., Caruso, M.V., De Rosa, S., Indolfi, C., and Fragomeni, G. (2018). Different Blood Flow Models in Coronary Artery Diseases: Effects on hemodynamic parameters. Annu Int Conf IEEE Eng Med Biol Soc *2018*, 3185-3188. 10.1109/EMBC.2018.8512917.

106. Govindaraju, K., Badruddin, I.A., Viswanathan, G.N., Ramesh, S.V., and Badarudin, A. (2013). Evaluation of functional severity of coronary artery disease and fluid dynamics' influence on hemodynamic parameters: A review. Phys Med *29*, 225-232. 10.1016/j.ejmp.2012.03.008.

107. Nixon, A.M., Gunel, M., and Sumpio, B.E. (2010). The critical role of hemodynamics in the development of cerebral vascular disease. J Neurosurg *112*, 1240-1253. 10.3171/2009.10.JNS09759.

1242   108.   Chiu, J.J., and Chien, S. (2011). Effects of disturbed flow on vascular endothelium:
1243   pathophysiological basis and clinical perspectives. Physiol Rev *91*, 327-387.
1244   10.1152/physrev.00047.2009.
1245   109.   Wang, C., Baker, B.M., Chen, C.S., and Schwartz, M.A. (2013). Endothelial cell sensing
1246   of flow direction. Arterioscler Thromb Vasc Biol *33*, 2130-2136.
1247   10.1161/ATVBAHA.113.301826.
1248   110.   Langley-Evans, S.C., and McMullen, S. (2010). Developmental origins of adult disease.
1249   Med Princ Pract *19*, 87-98. 10.1159/000273066.
1250   111.   Thornburg, K.L. (2015). The programming of cardiovascular disease. J Dev Orig Health
1251   Dis *6*, 366-376. 10.1017/S2040174415001300.
1252   112.   Tian, X., Pu, W.T., and Zhou, B. (2015). Cellular origin and developmental program of
1253   coronary angiogenesis. Circ Res *116*, 515-530. 10.1161/CIRCRESAHA.116.305097.
1254   113.   MacFarlane, E.G., Parker, S.J., Shin, J.Y., Kang, B.E., Ziegler, S.G., Creamer, T.J.,
1255   Bagirzadeh, R., Bedja, D., Chen, Y., Calderon, J.F., et al. (2019). Lineage-specific
1256   events underlie aortic root aneurysm pathogenesis in Loeys-Dietz syndrome. J Clin
1257   Invest *129*, 659-675. 10.1172/JCI123547.
1258   114.   Gailloud, P. (2022). The segmentation of the vertebral artery: An ambiguous anatomical
1259   concept. Interv Neuroradiol *28*, 765-772. 10.1177/15910199211063275.
1260   115.   Liu, M., and Gomez, D. (2019). Smooth Muscle Cell Phenotypic Diversity. Arterioscler
1261   Thromb Vasc Biol *39*, 1715-1723. 10.1161/ATVBAHA.119.312131.
1262   116.   Bentzon, J.F., and Majesky, M.W. (2018). Lineage tracking of origin and fate of smooth
1263   muscle cells in atherosclerosis. Cardiovasc Res *114*, 492-500. 10.1093/cvr/cvx251.
1264   117.   Egusa, G., Watanabe, H., Ohshita, K., Fujikawa, R., Yamane, K., Okubo, M., and Kohno,
1265   N. (2002). Influence of the extent of westernization of lifestyle on the progression of
1266   preclinical atherosclerosis in Japanese subjects. J Atheroscler Thromb *9*, 299-304.
1267   10.5551/jat.9.299.

1273
Figure 1. Single cell RNA and spatial landscape of human arteries.
(A) Schematic diagram of the 8 distinct human arterial sites collection strategy.
(B) Workflow of the experimental design and the sample processing for scRNA-seq and Slide-
seq.
(C) RNA expression of existing canonical markers of each cell type population.
(D) UMAP visualization and clustering of cells labeled by cell type.
(E) Spatial dim plot visualization of the right coronary artery (RCA), annotated by cell type (left).
The right panel presents a split view, illustrating the positions of individual cell types on the
same slide.
See also Figure S1.
1284

Figure 2. Smooth muscle cells of great arteries.

(A) RNA expression of mature smooth muscle contractile genes *MYOCD*, *CNN1*, *MYH11*, *ACTA2* and *TAGLN*, shown on the UMAPs.

(B) Immuno-fluorescence staining of CNN1 (green) and DAPI (blue) in ascending aorta.

(C) UMAP showing the *RGS5* RNA expression level.

(D) The differentially expressed genes between the two main clusters of SMC population, ranked by the average log2 fold change level.

(E) Dot plot showing the z scores of enriched biological processes in the differentially expressed gene between the two clusters of SMC population.

(F) The UMAPs showing the RNA expression of the unique makers in adventitial
microvasculature cell cluster (left panels). RNAscope validated the expression of these markers
with specific probes (right panels).
(G) RNA expression of medial and intimal smooth muscle genes *MYH11*, *FN1* and *TNFRSF11B*,
shown on the UMAPs.
(H) Spatial localization of the cells expressed *MYH11* and *FN1* in RCA (left, middle). And the
RNAscope staining of *TNSRSF11B* expression (right).
See also Figure S2 and Table S1.
1303

Figure 3

1304
Figure 3. Adventitial fibroblast in human arteries.
(A) The scores derived from the expression levels of collagen (middle) and extracellular matrix
genes (right) depicted on UMAP plots were analyzed alongside cell type (left).
(B) *PI16* RNA expression levels on RNAseq UMAP plot (left), Slide-seq dim plot alongside the
fibroblast localization in RCA (middle), and the RNAscope staining with probes (right).
(C) The differentially expressed genes between fibroblast cells and all other cell types, plotted
by their percent of expressed cells and the average log2 fold changes.
(D) The violin plots illustrate the specific expression of the genes *FBN1*, *TGFBR2*, *LOX*, *FBLN1*,
and MFAP5 within the adventitial fibroblast population.
See also Figure S3 and Table S2.

Figure 4

Figure 4. Distinct endothelial cell populations across arteries.

(A) Two large distinct clusters of endothelial cells shown on the UMAP.

(B) Endothelial cells identified by the canonical marker CDH5 RNAscope staining (left), and the endothelial cells localization on the Slide-seq dim plot of aorta (right).

(C) The sub-clustering of endothelial 1 population.

(D) The UMAP (top panels) and violin (bottom panels) plots showing the expression of cluster-specific genes *ACKR1* (left panels) and *HEY1* (right panels) in endothelial1 cells.

(E) The expression of cluster-specific genes *ACKR1* (left) and *HEY1* (right) are stained by

RNAscope probes in ascending aorta.

(F) The UMAP (left panels) and violin (right panels) plots showing the expression of SMC genes

TAGLN and *ACTA2* (left panels), or lymphatic genes *PROX1* and *LYVE1* (right panels) in

endothelial1 cells.

(G) Heatmap showing the z scores of *HOX* genes expression in each of the vessel segments in

both endothelial 1 and 2.

(H) Endothelial 2 specific genes *ITLN1* and *NRG1* expression levels on RNAseq UMAP plot.

(I) RNAscope staining of *ITLN1* gene expression in ascending aorta

(K) The co-staining of *ITLN1* (blue) and *CDH5* (red) expression in ascending aorta

(L) Dot plot showing the z scores of enriched biological processes in the differentially expressed

gene between endothelial 1 and 2.

See also Figure S4 and Table S3.

1337

1338
Figure 5. Prominent roles of embryonic origin in adult vascular identity.
(A) The heterogeneity of each population of cells across the different vascular beds on UMAP.
(B) A split view UMAP of the segmental origins presents the heterogeneity of each cell types.
(C) Heatmap showing the z scores of *HOX* genes expression in each of the vessel segments in both SMC and fibroblast.
(D) Dendrogram Clustering using Euclidean distance calculated by all expressed genes showing the correlation of each segment with its anticipated embryonic lineage within fibroblasts.
(E) Heatmap showing the percentage of each segment depicted within the distinct UMAP clusters.
(F) Violin plots showing the differential expression patterns of transcription factors *ISL1*, *HAND2*, *MEDF2C*, and *HOX* genes in fibroblast of ascending and descending sites.

1350   (G) Enrichment of ISLET1, HAND2, MEF2C, and HOX binding motifs on the promoter regions of

1351   differentially expressed genes between ascending and descending fibroblasts, with

1352   macrophages serving as controls.

1353   See also Figure S7.

1354

1355

Figure 6. Adult vascular *HOX* gene expression patterns.

(A) Heatmap showing the log10 normalized expression values of *HOX* genes across the

different vascular segment in fibroblast, SMC, endothetial1, endothelial2, microvasculature, or

macrophage.

(B) Dendrogram clustering using Euclidean distance calculated by all *HOX* genes showing the

correlation of each segment with its anticipated embryonic lineage in SMC (bottom) or

fibroblasts (top).

(C) The heatmap illustrates the Pearson correlation coefficients among all arterial segments,

derived from the expression values of all *HOX* genes. These segments are clustered according

to Euclidean distance in SMC (left) or fibroblast (right).

See also Figure S8.

1367

Figure 7

1368
Figure 7. Enrichment of vascular disease variants in segmental transcriptomic differences.
(A) Dot plot showing the differential expression patterns of mendelian aortopathies genes *FBN1*, *FBNL1*, *LOX*, *MFAP4*, *MFAP5*, and *COL3A1* genes in fibroblast of ascending and descending sites.
(B) Bar plot showing the differential expression significances of these mendelian aortopathies genes *FBN1*, *FBNL1*, *LOX*, *MFAP4*, *MFAP5*, and *COL3A1* between ascending and descending sites in fibroblast.
(C) Violin plots showing the differential expression patterns of aneurysm risk in fibroblast (left) or SMC (right) of ascending and descending sites.
(D) The bar graph displays the count of Million Veteran Program (MVP) GWAS genes that are differentially expressed in each cell type when comparing atherosclerosis prone (coronary / carotid) with relatively atherosclerosis resistant descending aorta.
(E) Genetic association between the MVP GWAS genes and the differentially expressed genes across the coronary artery, carotid, and descending segments within each cell type, ranked by the number of overlapping genes.

1385   (F) Heatmap showing the z scores of fold enrichment of biological processes in the differentially
1386   expressed genes across the coronary artery, carotid, and descending segments within each cell
1387   type.
1388   See also Figure S9 and Table S4, S5, S6.
1389

1390

Figure 8. Cell type and segment specific non-coding transcriptome.

(A) Heatmap showing the z scores of non-coding RNAs expression in each of the cell types.

(B) Heatmap showing the z scores of non-coding RNAs expression in each of the vascular

segments.

(C) UMAP visualization and clustering of cells using the sc-lncRNA transcriptome, annotated

based on either whole transcriptome-derived cell types (left) or lncRNA-derived cell type

identification (right).

(D) Dendrogram clustering using Euclidean distance calculated by all expressed lncRNAs

showing the correlation of each segment with its anticipated embryonic lineage in SMC (top) or

fibroblasts (bottom).

(E) The dot plot (top) illustrating the MVP SNPs positioned directly within the loci of differentially

expressed lncRNAs. The violin plot displays the expression levels of these intersected lncRNAs.

See also Figure S11 and Table S10.

1404