Anisotropy of object nonrigidity: High-level perceptual consequences of cortical anisotropy

Akihito Maruya & Qasim Zaidi*

Graduate Center for Vision Research, State University of New York, 33 West 42nd St, New York, NY 10036.

*Corresponding author: qz@sunyopt.edu

AUTHOR CONTRIBUTIONS

AM and QZ designed the study. AM programmed and ran the experiments. AM & QZ analyzed and modeled the results. AM & QZ wrote the paper.

The authors declare no competing interests.

Acknowledgements: NEI grants EY035085 & EY035838

Abstract

We present a surprising anisotropy in perceived object nonrigidity, a complex, higher-level perceptual phenomenon, and explain it unexpectedly by the distribution of low-level neural properties in primary visual cortex. We examined the visual interpretation of two rigidly connected rotating circular rings. At speeds where observers predominantly perceived rigid rotation of the rings rotating horizontally, observers perceived only nonrigid wobbling when the image was rotated by 90°. Additionally, vertically rotating rings appeared narrower and longer compared to their physically identical horizontally rotating counterparts. We show that these perceived shape changes can be decoded from V1 outputs by incorporating documented anisotropies in orientation selectivity. We then show that even when the shapes are matched, the increased nonrigidity persists in vertical rotations, suggesting that motion mechanisms also play a role. By incorporating cortical anisotropies into optic flow computations, we show that the kinematic gradients (Divergence, Curl, Deformation) for vertical rotations align more with physical nonrigidity, while those for horizontal rotations align closer to rigidity, indicating that cortical anisotropies contribute to the orientation dependence of the perception of nonrigidity. Our results reveal how high-level percepts are shaped by low-level anisotropies, which raises questions about their evolutionary significance, particularly regarding shape constancy and motion perception.

Introduction

We present a remarkable instance of variations in a complex higher-level percept explained directly from the distribution of low-level neural properties in primary visual cortex by using a combination of mathematical derivations and computational simulations to reproduce the results of psychophysical experiments. The higher-level percept we demonstrate is an unexpected anisotropy in object nonrigidity, and the low-level properties are documented cortical anisotropies in numbers and tuning widths of orientation selective and direction selective neurons. The video in Figure 1A shows two connected Styrofoam rings on a rotating turntable. Most observers report that the two rings are rigidly joined and rotating together, although sometimes the top ring is reported as moving independently from the bottom ring and rolling or wobbling, seemingly defying physical plausibility. In the video in Figure 1B, the right ring is almost always reported as moving independently, wobbling against the other ring. The context reveals that one video is just a 90° rotation of the other, so clearly an explanation is needed for the increase in nonrigidity.

3D-shape-from-motion models typically assume that objects are rigid, which simplifies the math[1] but organisms generally change shape to move, eat, observe, mate, etc., so articulated and elastic motions are common in the real world[2], and human observers are as good at discerning 3D shapes of nonrigid objects as of rigid objects[3]. Maruya and Zaidi (2024) showed that a rotating rigid object with two salient parts, such as the object in Fig 1, can appear rigid or nonrigid depending on speed of rotation and the presence of salient features[4]. At moderate to high speeds, the nonrigidity is predictable from the motion-energy flow, whereas at slow speeds, feature tracking dominates[5,6], if trackable features are present [7,8,9,10,11,12]. The change in nonrigidity when the image is rotated by 90° could be due to anisotropy in shape or motion mechanisms. We investigate both as possible consequences of documented anisotropies in the population distribution of orientation and direction selective neurons.

To function successfully in the objective reality of the physical world, an

organism's perceptions must lead to accurate inferences of attributes such as shapes,

sizes, locations, distances, motions, colors, textures, and kinds[13]. The earliest

discovered cave drawings[14] suggest that shape is the most important attribute for

humans to identify objects and animals. In mathematics, the shape of a rigid object is a

geometric attribute that remains unchanged across variations in location, rotation, and

scale[15], which corresponds to viewpoint invariance for a visual system. The shape of an

object would not be a useful attribute if the mathematical definition was violated when

an object was viewed from different angles. Unfortunately shape constancy is often

violated. For example, horizontal parallelogram balconies on a tall building are seen as

rectangles tilted up or down depending on viewpoint[16]. Such violations have been

attributed to a tendency to assume right angles[16,17], to try to maximize compactness or

symmetry[18], but the neural substrates of such processes are completely unknown. A

more understandable kind of shape-constancy violation for 3D objects is caused by

simply rotating the object in the frontal-plane. Cohen and Zaidi showed that the

perceived depth of a wedge defined by shape-from-texture cues is a function of the

orientation of the shape[19]. They linked this to a 2D anisotropy in the perception of an

angle as a function of its orientation and showed that the 2D effect could be explained

by a model based on the documented anisotropy in number and selectivity of orientation

tuned cells in primary visual cortex of mammals which show the greatest concentration

and narrower tuning for horizontal orientations, then vertical, and then oblique[20] The

anisotropy in primary visual cortex has been linked to encoding the statistical

distribution of orientations in the world[21,22], thus suggesting a common cause for the

perceived effects of object orientation and the "oblique effect" where observers exhibit

superior performance for stimuli presented along the cardinal (vertical and horizontal)

axes as compared to oblique axes[23]. The cortical anisotropy has now been well

documented for primate brains[24], where the anisotropy seems to increase from V1 to

V4, raising the question of why evolution leads to brain properties that give more

importance to the detection/discrimination of isolated orientations than to preserving

perceived shapes across image rotations.

The neural bases of anisotropy of motion processing and perception is less well

understood. Psychophysical anisotropies favoring cardinal directions over oblique

directions have been shown for direction-discrimination but not for direction-detection of

translating stimuli[25, 26, 27, 28, 29]. The anisotropies that have been documented in area

MT[30, 31] seem to not be systematic, although the number of cells measured is much less

than in the V1 studies. There are some reports of direction selectivity for object motion

in V4[32], where there are documented anisotropies[24], but the two issues have not been

studied jointly. Given that most V1 simple cells receiving Magnocellular input from LGN

are direction selective[33], in our modeling, we assume that the direction preference and

tuning of cells in mammalian striate cortex could be assumed to correspond to

orientation preference along the axis orthogonal to the motion direction[20]. To our

knowledge, anisotropies in motion based higher level perception, such as 3D-shape-

from-motion or rigidity versus nonrigidity, have not been studied previously.

Shape and nonrigidity anisotropies are also easy to see when we switch from

physical objects to computer graphics and orient the object to be symmetrical around

the vertical or horizontal axis, such as the videos of the ring-pairs in Figures 1C – 1H.

Figures 1C & 1D in the 2nd row are perspective projections of rigidly rotating

configurations and Figures 1E & 1F in the 3rd row are perspective projections of rings

that wobble and slide nonrigidly against each other. When attention is directed to the

connecting joint, rigidity and nonrigidity are both easy to see in the octagonal rings

because the vertices at the junction of the two rings either stay together or slide past

each other. The circular rings, however, are without features, so rigidly rotating and

wobbling rings generate identical retinal images, and could be seen as rigidly rotating or

nonrigidly wobbling, depending on how the images are processed by the visual system.

If the image is rotated by 90 degrees, the nonrigidity becomes much more pronounced

for both types of rings (Figures 1G & 1H in the 4th row). Figures in the 2nd and 4th rows

show a single frame of horizontally or vertically rotating rings, where the intersection of

the two rings is in the center of the image. The vertically elongated rings appear

noticeably narrower and longer than the horizontal ones. By turning the page 90

degrees, the reader can ascertain that this shape illusion is a function of object orientation.

Are the shape changes responsible for the increased nonrigidity, and is it due to a cortical anisotropy? We first show that these perceived shape changes can be decoded from V1 outputs, by considering anisotropies in the number and tuning widths of orientation-selective cells. Then we show empirically that when observers widened the vertically rotating ellipses or elongated the horizontally rotating ellipses so that the shapes matched, the perceived nonrigidity was still greater for the vertical rotation. We next incorporated cortical anisotropies into motion flow computations. The estimated motion fields were decomposed into velocity gradients of divergence, curl, and deformation. Rigidity of rotating 2D shapes can be tested by showing that only the curl component is non-zero[34], but this is not true for the projections of the 3D rings, so we had to compare all three gradients generated by physical rotation versus physical wobbling. The gradients for vertical rotation of the rings more closely matched physical wobbling, while the gradients for horizontal rotation fell between physical wobbling and rotation, suggesting that hardwired cortical anisotropies can explain the increase in perceived nonrigidity from horizontal to vertical rotation.

Results

We will first test whether the static shape anisotropy can account for the nonrigidity anisotropy, and if that does not account for most of the illusion, we will then examine velocity anisotropy.

Shape Anisotropy

Figures 2A & 2B present the two orientations of the rings in the phase that best demonstrates that the vertical rings look narrower and longer than the horizontal rings. This illusion is similar to anisotropy for an elongated diamond (Figure 2C & 2D). The diamonds have identical shape but the vertical diamond looks narrower and longer. In Figure 2C & 2D, the angles ϕ_h and ϕ_v are physically equal but ϕ_v looks narrower (the subscript *h* indicates angles centered around the horizontal axis, while the subscript *v* indicates angles centered around the vertical axis). Similarly, ψ_h looks wider despite ψ_h and ψ_v being equal angles. As a result, the vertical diamond looks narrower. Cohen and Zaidi showed that flat 2D angles look broader when the central axis is obliquely oriented compared to vertical and modeled this as an effect of decoding the outputs of V1 cells, where those tuned to horizontal orientations were larger in number and narrower in tuning than those tuned to oblique orientations[19]. We now extend this to comparing vertical versus horizontally centered angles to explain the diamond illusion.

Cortical Anisotropy

Figure 3A shows the orientation tuning widths of V1 simple cells we used in our model, centered at the preferred orientation, where 0° and ±180° represent horizontal preferred orientation, and ±90° represent vertical orientation, representing a summary of cat cortex measurements made with drifting gratings[20]. The heights of the curves reflect the relative number of cells tuned to each direction[20]. Cardinal orientations exhibit a higher number of cells and narrower tuning width. However, cells preferring vertical orientation are fewer and broader tuned than cells preferring horizontal orientation. The left two panels in Figure 3B show the Li et al (2003) curves plotted in terms of preferred orientation by averaging across opposite motion directions[20]. There are more recent and more extensive data on primate V1 and V4[24], but we did not use these because they provide only relative numbers, whereas Cohen & Zaidi found that tuning widths were

more important than numbers, and our simulations reproduced that effect[19]. The

simulated orientation tuning curves use von Mises probability density functions to

generate the tuning curve for all orientations θ:

$$f(\theta|\mu_i, k_i) \propto \exp(k_i \cos(\theta - \mu_i)) \tag{1}$$

Where μ_i is the preferred orientation of the i-th cell and k_i sets the tuning width. Since

we will be decoding angles, two orientations 180° apart are considered distinct, so

Equation 1 covers 360°, unlike most fits of the von Mises function[35]. As a control, we

also decoded angles from a cortex with isotropic distribution of orientation preference

and tuning (Figure 3B).

Perceived angle decoded from cortical responses

In Figure 2, angles around the horizontal axis (γ_h) are perceived to be wider than equal

angles around the vertical axis (γ_v) and we decoded the difference between them for

angles ranging from 10-170 degrees from an anisotropic cortex as well as an isotropic

cortex. We assume that each angle γ is formed by two lines θ_a and θ_b, to which the

response of n_i neurons with preferred orientation μ_i are r_{ai} and r_{bi} :

$$r_{ai} = n_i f(\theta_a|\mu_i, k_i)$$

$$r_{bi} = n_i f(\theta_b|\mu_i, k_i) \tag{2}$$

Each response is normalized by a divisive gain control:

$$R_{ai} = \frac{r_{ai}}{g_i}, R_{bi} = \frac{r_{bi}}{g_i} \tag{3}$$

$$\text{where } g_i = \sqrt{r_{ai}^2 + r_{bi}^2 + c}$$

In primary visual cortex, neuronal responses are suppressed by the presence of

surrounding stimuli. This suppression is orientation-specific, with the strongest

suppression occurring when the stimuli in the receptive field and the surround share a

similar orientation, and the weakest suppression occurring when they are orthogonal to

each other[36, 37,38]. The divisive gain control mechanism, involving self-normalization or

dividing by the weighted sum of all other neurons (where the weight is highest for

neurons with the same orientation preference), can achieve this neural interaction[39].

The line orientations, θ_a and θ_b, are decoded using vector sums[40, 41]:

11
$$\widehat{\theta_a} = Ang\left(\sum_i R_{ai}\boldsymbol{l}\right), \widehat{\theta_b} = Ang\left(\sum_i R_{bi}\boldsymbol{l}\right) \qquad (4)$$

14 Where $\boldsymbol{l}$ is the unit vector that points to the preferred orientation. Then, the decoded

15 angle is simply calculated from the two decoded line orientations:

17
$$\widehat{\gamma} = mod\left(\widehat{\theta_a} - \widehat{\theta_b}, 180°\right) \qquad (5)$$

Right two panels in Figure 3B show decoded angles for the anisotropic cortex (red) and

the isotropic cortex (black). Figure 3C displays the decoded difference $(\widehat{\gamma_h} - \widehat{\gamma_v})$

between an angle centered around the horizontal axis and a physically identical angle

around the vertical axis, for the anisotropic cortex (red) and the isotropic cortex (black).

Decoded responses from the isotropic cortex show no difference between the estimated

horizontal and vertical angles. However, when decoded from the anisotropic cortex,

angles around the horizontal axis γ_h are estimated to be wider than physically equal

angles around the vertical axis γ_v which is consistent with what is observed in Figure 2C

& 2D. The difference varies with the magnitude of the angle but is greater than 5

degrees for all but very obtuse angles. We present this simulation to show that the

illusory shape anisotropy in diamonds can be attributed to anisotropy in cortical

populations. The estimated difference for a physical angle of 90° also explains why a

square rotated 45° appears as a diamond which is elongated vertically[42]. Similar, but

more complex decoding would yield a similar conclusion for the ellipses in Figure 2A &

2B. The qualitative inference is sufficient for our purposes, because the anisotropy

assumptions from cat cortex are qualitatively similar but not quantitatively identical to

measurements from primate cortex. Without the normalization in Equation 3, the

magnitude of the decoded difference is appreciably smaller. Cohen & Zaidi (2007)

reproduced perceived differences in vertical versus oblique angles with maximum

likelihood decoding by explicitly including cross-orientation suppression[19].

Effect of shape matching on perceived nonrigidity

Maruya & Zaidi (2024) showed that horizontally rotating elliptical rings with physical

aspect ratios greater than 1.0 (narrower than circular rings) are perceived as more

nonrigid than horizontally rotating elliptical rings with physical aspect ratios smaller than

1.0 (wider than circular), so the shape difference is a potential explanation for the

nonrigidity anisotropy[4]. To test whether this explanation is sufficient, we had observers

adjust shapes for vertical and horizontal rotations until they matched, and then tested if

matching projected shapes abolishes the nonrigidity anisotropy.

We presented pairs of vertically and horizontally rotating rings and asked

observers to match their shapes by stretching one of the pairs horizontally. Note that

horizontal stretching makes the horizontally rotating rings narrower, and the vertically

rotating rings wider, thus both counter the illusory percept of greater elongation for the

vertically rotating rings. In Figure 4A, the 1st and 3rd rows show horizontal stretches of

the 2D images ranging from 0% to 50%, while the 2nd and 4th rows illustrate 3D

physical stretches of the rings, i.e. simulated stretches of the 3D rings observed as

stretches of the projected 2D images of the rings. Figures 4B and 4C present

histograms of the horizontal stretch in the image domain (red) and the physical domain

(blue) that match the shapes of horizontally rotating rings (left) and vertically rotating

rings (right). To match shapes, observers stretched the horizontal rings and the vertical

rings in the horizontal direction by approximately 20-30%. Individual results are shown

in Supplementary Figure S1.

Next we presented selected pairs of rotating rings in random order and 4

observers reported which pair looked more independent and nonrigid: (i) original circular

horizontal versus circular vertical rings, (ii) physically stretched horizontal rings versus

circular vertical rings, (iii) physically stretched vertical rings versus circular horizontal

rings, (iv) stretched horizontal rings in the image domain versus circular vertical rings,

and (v) stretched vertical rings in the image domain versus circular horizontal rings.

Figure 4D shows the probability of observers reporting vertically rotating rings as more

nonrigid. For the original pair, observers perceived vertically rotating rings as more

nonrigid 95.0% of the time (black). Matching the shapes perceptually reduced the

nonrigidity anisotropy by up to 40%, with a greater reduction observed in image-stretch

(red bars) compared to physical stretch (blue bars). Individual results are shown in

Supplementary Figure S2. In an ancillary forced choice experiment, observers picked

the image stretch as a better shape match than the physical stretch (61%), possibly

because the shape anisotropy is an illusory stretch of the image. However, the

nonrigidity anisotropy remains significantly above chance levels despite the shape

matches so the perceived shape does not account for almost 60% of the effect. As a

critical test, we took the most stretched horizontal and vertical rings (Figure 4A

Rightmost rings in 2nd and 4th row) and asked 4 observers to choose which pair was

more nonrigid when rotated. Despite the horizontal rings being much narrower than

vertical rings, the opposite of the original condition, observers picked the vertical rings

as more nonrigid on 80% of the comparisons, opening the possibility that anisotropy of

low-level motion detectors also plays a role.

Nonrigidity anisotropy from cortical anisotropy in direction selectivity

As we state in the introduction, since the cat cortical anisotropy was measured with

drifting gratings[20], we make the reasonable assumption that the anisotropic distribution

of numbers and tuning widths also represents direction selective cells. To simulate the

effect of spatial anisotropy on optic flow, we computed local motion energy for each

pixel of the videos. Each motion-energy unit consisted of a quadrature pair of 3D Gabor

filters[43, 44, 45, 46, 47, 48, 49,50]:

$$G_{i,odd}(x,y,t) = Q_i\exp\left(-\frac{(x-c_{x,i})^2 + (y-c_{y,i})^2}{2\sigma_{s,i}{}^2} - \frac{(t-c_{t,i})^2}{2\sigma_{t,i}{}^2}\right)$$
$$\times \sin(2\pi\omega_{x0}x + 2\pi\omega_{y0}y + 2\pi\omega_{t0}t) \tag{6}$$

$$G_{i,even}(x,y,t) = Q_i\exp\left(-\frac{(x-c_{x,i})^2 + (y-c_{y,i})^2}{2\sigma_{s,i}{}^2} - \frac{(t-c_{t,i})^2}{2\sigma_{t,i}{}^2}\right)$$
$$\times \cos(2\pi\omega_{x0}x + 2\pi\omega_{y0}y + 2\pi\omega_{t0}t) \tag{6'}$$

Where Q_i is the normalization constant; and $c_{x,i}$ and $c_{y,i}$ are the spatial means of the

Gaussian envelopes, and $c_{t,i}$ is the temporal mean; $\sigma_{s,i}$ and $\sigma_{t,i}$ are the spatial and

temporal standard deviations of the Gaussian envelopes, respectively; and ω_{x0}, ω_{y0},

and ω_{t0} are the spatial and temporal frequencies of the sine component of the Gabor

filter, referred to as the preferred spatial and temporal frequencies where:

$$\omega_{x0} = F_0\sin\eta_t\cos\eta_s; \quad \omega_{y0} = F_0\sin\eta_t\sin\eta_s; \quad \omega_{t0} = F_0\cos\eta_t \tag{7}$$

η_s and η_t are spatial and temporal orientations (8 spatial orientations from 0 to $7/8\pi$ and

5 temporal orientations from $-\pi/4$ to $\pi/4$) and F_0 is the frequency (0.28 and 0.14

cycles/pixel). For the anisotropic cortex, σ is adjusted so that the half-height orientation

bandwidths $\Delta\omega_{\frac{1}{2}}$ are matched to cat cortex estimates[20] by the following equation[51]:

$$\Delta\omega_{\frac{1}{2}} \approx 2\tan^{-1}\left(\frac{0.5}{F_0\sigma}\right) \tag{8}$$

Each filter $G_{i,even}$ and $G_{i,odd}$ was convolved with the video $I(x,y,t)$. The responses of the

quadrature pair were squared and summed to produce a phase-independent response

m_i. Figure 5 A and B illustrate the direction tuning curves for isotropic and anisotropic

cortices, weighted by the number of cells. In the anisotropic cortex, cells that prefer a

vertically moving bar (either upwards or downwards) exhibit narrower tuning and are

more numerous compared to cells that prefer oblique or horizontally moving bars.

To obtain the best estimate of velocity $(\hat{u}, \hat{v})$ per point using the whole family of filters,

we computed optic flow by minimizing a weighted average of the combined filter

responses[52]:

$$(\hat{u}, \hat{v}) = argmin_{(u,v)} \left(\sum_i n_i (m_i - \overline{m_i} \frac{\mathfrak{N}_i(u,v)}{\overline{\mathfrak{N}_i(u,v)}})^2 \right) \quad\quad (9)$$

$$\mathfrak{N}_i(u,v) = \exp\left(-\frac{4\pi^2 \sigma_s^4 \sigma_t^2 (u\omega_{x0} + v\omega_{y0} + \omega_{t0})^2}{u^2 \sigma_s^2 \sigma_t^2 + v^2 \sigma_s^2 \sigma_t^2 + \sigma_s^4} \right) \quad\quad (9')$$

Where $\mathfrak{N}_i(u,v)$ is the predicted response of filter i to velocity (u,v), m_i is the motion

energy output, and $\overline{m_i}$ and $\overline{\mathfrak{N}_i(u,v)}$ are the sum of $\mathfrak{N}_i$ and the sum of m_i for filters

sharing the same spatial orientation. Figure 5C shows optic flow fields for different

conditions of rotation of a single ring, horizontal rotation $\mathbb{R}_y$ around the y-axis (left

column), and vertical rotation $\mathbb{R}_x$ around the x-axis (right column), for the isotropic

cortex (top row), and the anisotropic cortex (bottom row). The color of the vector

corresponds to the direction and the length represents the magnitude of the estimated

velocity. Most of the optic flow vectors from the isotropic cortex are orthogonal to the

contour. In contrast, for the anisotropic cortex, the optic flow vectors shift towards the

horizontal direction, as indicated by the red and turquoise colors.

To understand what this horizontal shift in the velocity field implies for perceived

rotation versus wobble (which adds rotation $\mathbb{R}_z$ around the z-axis), we derived the

velocity field for different combinations of physical rotation and wobbling of a single ring,

with the wobbling weight k ranging from $k = 0 \sim rotation$ to $k = 1 \sim wobbling$. We began

with the vector $\vec{P}(\theta) = (cos\theta, sin\theta, 0)^T$ which defines the position of every point on a ring

lying on the $X - Z$ plane for each angular coordinate of the circular ring θ (Figure 5D) ,

and calculated $\overrightarrow{P_k}(\theta, t)$ for every combination of rotation and wobble given by $\mathbb{R}_y$ and $\mathbb{R}_z$

the rotational matrices around the Y and Z axis respectively, with the angular velocity ω and the tilt of the ring, τ:

$$\overrightarrow{P_k}(\theta, t) = Proj(\mathbb{R}_y(\omega t)\mathbb{R}_z(\tau)\mathbb{R}_y(-k\omega t)\vec{P}(\theta)) = (u(\theta, t), v(\theta, t))^T \qquad (10)$$

$Proj(\cdot)$ is a projection function and we used the orthographic projection. Then, the velocity field F_k is calculated by the partial derivative with respect to time:

$$F_k = \frac{\partial}{\partial t}\overrightarrow{P_k}(\theta, t) \qquad (11)$$

Figure 5E illustrates the velocity fields for horizontal rotation (top) and vertical (bottom), obtained with wobbling weights (k) increasing from left to right, from pure rotation to pure wobbling. For horizontal rotation (top row), horizontal vectors are predominant when $k = 0$ and vertical vectors at $k = 1$, with the opposite for the vertical rotation (bottom row). These velocity fields serve as templates, and the cosine similarity between these derived templates and the estimated optic flows in Figure 5C is calculated at each point and time for both the isotropic and anisotropic cortex (Figures 5F & 5G) and then averaged over all points for a complete cycle. The blue trace represents mean cosine similarity for horizontal rotation, and the red trace for vertical rotation. As expected, for flows calculated from an isotropic cortex, cosine similarities are the same for horizontal and vertical rotations with the best fit at k = 0.84. For flows from the anisotropic cortex, horizontal and vertical rotations result in different cosine similarities. The horizontal rotation cosine similarity curve is relatively flat from k values 0.0 to 0.7 with a peak around 0.4, while the vertical rotation curve is steep and peaks at a higher k, close to 0.9. Thus, if generated velocity fields were compared to velocity field templates for physical rotation and wobbling, it would, explain why observers perceive more nonrigidity when the two rings rotate vertically. However, having all possible templates is biologically implausible, so next we explore how a more biologically plausible factoring of the velocity fields can lead to similar comparisons that predict the nonrigidity anisotropy.

Nonrigidity anisotropy from biologically plausible filters for differential invariants

In kinematics, which describes the motion of objects without considering the forces that

cause them to move, motion fields are analyzed in terms of the local differential

invariants, divergence, curl, and deformation after excluding translation parallel to the

image plane, and these analyses have been useful in understanding object motion from

optic flow[34, 53, 54, 55]. For the single ring with some combination of rotation and wobble,

the divergence field is perpendicular to the contour and captures expansion/contraction

of the projection (Figure 6A); the curl field is tangential to the contour and captures

rotation; the deformation fields captures shear caused by the difference between

orthogonal axes in contraction/expansion or shear and can be combined into one

measure giving direction and magnitude of deformation[55]. Besides providing

understandable summary measures, these operators also have biological validity.

Zhang et al. (1993) demonstrated that position-independent detectors of curl and

divergence can emerge through simple Hebbian learning, from MT-like receptors to

MST-like receptors[56]. Physiological studies on nonhuman primates and imaging studies

on humans have identified neurons in areas MST, STPa, and STS that preferentially

respond to the curl and div components of optic flow[57, 58, 59, 60]. In addition, neural activity

sensitive to def has been found in dorsal MST[62], and human psychophysical

experiments have demonstrated that perceived slant is proportional to def [53, 62, 63].

To derive divergence, curl, and deformation fields for physical rotation and

physical wobbling, the dot product of the velocity fields derived in Equation 11 as a

function of k (illustrated in Figure 5E) was taken with each operator corresponding to

those illustrated in Figure 6A, and then integrated across space to obtain one number

for each invariant per phase of motion, where i is the basis vector (1,0) and j is (0,1) :

Divergence:

$$\int F_k \cdot \left(\frac{dv}{d\theta} \boldsymbol{i} - \frac{du}{d\theta} \boldsymbol{j} \right) d\theta \tag{12}$$

Curl:

$$\int F_k \cdot \left(\frac{du}{d\theta} \boldsymbol{i} + \frac{dv}{d\theta} \boldsymbol{j} \right) d\theta \tag{13}$$

Deformation:

$$Def1 = \int F_k \cdot \left(-\frac{du}{d\theta} \boldsymbol{i} + \frac{dv}{d\theta} \boldsymbol{j} \right) d\theta \, ,$$
$$Def2 = \int F_k \cdot \left(\frac{dv}{d\theta} \boldsymbol{i} + \frac{du}{d\theta} \boldsymbol{j} \right) d\theta \tag{14}$$

These are combined into one measure of the magnitude of Def:

$$Def = \sqrt{Def1^2 + Def2^2} \tag{14'}$$

Figure 6B displays the values of derived integrated gradients for divergence, curl, and deformation from left to right as a function of the motion phase. The colors represent different wobbling weights (k), with warmer colors indicating more wobbling and cooler colors representing more rotation. For all three gradients, the velocity field for rotation exhibits lower variance across different motion phases because the local velocity vectors consistently point in horizontal directions, so changes in the value result solely from alterations in shape due to projection. The wobbling field exhibits higher variance because the projected shapes and thus the local velocities change more over time. The value of Divergence varies from positive to negative representing divergence versus convergence, and the amplitude is greater for wobble because of the additional axis of rotation. Curl for the rotational velocity field is high, but almost constant, reflecting the minimal change in projected shape, whereas the greater projected shape changes for wobbling lead to smaller values of curl but a larger amplitude of variation. The derived

deformation estimates have a similar variation. Taken together, the three derived gradients for the physical motions provide templates for distinguishing between relative weights of rotation and wobble, so next we estimated the values of the gradients for the optic flow fields for the single rotating ring computed from the isotropic and anisotropic cortex (Figure 5C). For cortical isotropy (Figure 6C), the divergence, curl, and deformation are identical for vertical and horizontal orientation (red symbols represent divergence, green represent curl, blue represent deformation). The dashed smooth curve in the corresponding color indicates the best fit of the derived physical motion. For both orientations, the best fits occur at a wobbling weight of $k = 0.78$, consistent with the computational results in Maruya and Zaidi [4] using CNNs that showed that motion-energy flows support percepts of wobble, because they predominantly signal motion orthogonal to contours. The cortical anisotropy distorts the optic flow so that the optimal fit for vertical rotation ($k = 0.91$) corresponds closer to physical wobbling than the fit ($k = 0.48$) for horizontal rotation. The analysis in terms of differential invariants thus provides an explanation for the anisotropy of nonrigidity that matches the one provided by template matching.

To further test the efficacy of this model, we tested if it could predict reduced anisotropy due to horizontal shape stretching in the shape matching experiment. We took the videos of the rotating single ring and applied horizontal image stretches ranging from 0% to 50%, and calculated the optic flows generated by the isotropic and anisotropic cortex similar to that shown in Figure 5C. Then we calculated the three differential invariants at each image stretch and estimated the optimal k as in Figures 6C and 6D for horizontal rotation and vertical rotation. For both isotropic (Figure 6E) and anisotropic (Figure 6F) cortices, increased stretching during horizontal rotation (red) results in a higher amount of wobbling, while increased stretching during vertical rotation (blue) leads to a reduced amount of wobbling, consistent with the experimental result that reduced anisotropy is observed with a greater amount of horizontal stretch. However, for the anisotropic cortex vertical rotation maintains higher nonrigidity across different amounts of stretch, which aligns with the results in Figure 4D, while the isotropic cortex predicts the opposite. Moreover, the isotropic cortex simulations predict least anisotropy at 0% stretch and most at 50%, which is the opposite of the

experimental results, while the anisotropic predicts that the anisotropy reduces with

stretch but does not disappear, consistent with the experimental results. Taken together,

the simulations from the anisotropic cortex explain the experimental results.

Experimental methods

<u>Stimulus generation</u>

Using Python, we created two circular rings connected rigidly at a 60 degree angle,

rotating around either a vertical axis or a horizontal axis oblique to both rings. The joint's

speed when passing fronto-parallel was 6°/s, with a ring diameter of 3 degrees of visual

angle (dva), resulting in an angular speed of 0.3 cycles per second (cps). We presented

pairs of vertically and horizontally rotating rings and asked observers to match their

shapes. The rings could be stretched horizontally from 0% to 50%, either in the image

domain or the physical domain. Horizontal stretches in the image domain transformed

the projected ellipses for the horizontal rotation into narrower and longer ellipses, and

into wider and shorter ellipses for the vertically rotating axes. In the physical domain,

stretches transformed the circular rings into elongated ellipses, and transformations in

the projected images were similar to those for the image stretches. Figure 4A shows

examples of elongated rings as a function of horizontal stretch in the image domain (first

and third rows) and the physical domain (second and fourth rows). The matching was

done while the rings were rotating in the same phase.

<u>Stimulus display</u>

The videos were displayed on a VIEWPixx/3D (VPixx Technologies, Saint-Bruno-de-

Montarville, QC, Canada) at 120 Hz. PsychoPy was used to display the stimulus and

run the experiment. The data were analyzed using Python. An observer's viewing

position was fixed with a chin rest to ensure the video was viewed at the same elevation

as the camera position.

<u>Psychophysical procedures</u>

Observers were asked to use the mouse to select the best-matched shape. After three

trials, the median stretch was taken, and each matched shape in the image and

physical domains was presented side by side for the observer to select the most similar

shape. Finally, we presented a vertically rotating ring pair and a horizontally rotating ring

pair side by side, and observers were asked to select the most nonrigidly moving pair.

There were 5 conditions as shown in Figure 4D. The set of conditions was repeated 40

times. Measurements were made by four observers with normal or corrected-to-normal

vision. Observers gave written informed consent. All experiments were conducted in

compliance with a protocol approved by the institutional review board at SUNY College

of Optometry, adhering to the tenets of the Declaration of Helsinki.

Discussion

Despite the abundance of nonrigid 3D objects and organisms in the world, and

despite the evolutionary advantage of being able to judge their actions from shape

deformations, nonrigid motion perception is an understudied phenomenon[3, 64, 65, 66, 67, 68,

70, 71, 72, 73]. Maruya and Zaidi[4] showed that the horizontal rotation of two rigidly

connected circular 3D rings can appear either rigid or nonrigid (rings wobbling and

rolling independently) if the rotational speed is sufficient to activate motion-energy cells.

They found that the nonrigid perception arises from populations of motion energy cells

signaling motion orthogonal to the contour. The nonrigidity can be decreased and even

abolished if the rings contain salient features that can be used for feature-tracking and if

the shape does not support a prior for spinning motion[4]. We strengthen this link by

showing that the perceived nonrigidity is strongly affected by the cortical anisotropy for

motion-direction selective neurons.

The nonrigidity anisotropy we demonstrate is as interesting as it is unexpected.

We show that the hard-wired cortical anisotropy is powerful enough in modifying the

optic flow from the vertically rotating rings towards the horizontal directions to counter

the effects of salient features that evoke feature-tracking and overcome the effects of shape-based priors in the horizontal rotation (Video in Figure 1H).

A model that explains the complete percept of the 3D rings is beyond the scope of this paper, so we concentrate on the perception of nonrigidity versus rigidity and find surprisingly that variations in this high-level percept can be explained by anisotropies in the neural population, possibly even at the primary visual cortex. That certain variations in a high-level percept such as rigidity are a function of hardwired low-level neural properties is remarkable. Other than Maxwell's success in identifying cone photoreceptors from color matches, there are almost always too many degrees of freedom to model anything definitive about neural mechanisms from psychophysical results. In addition, it seems that there is little information about higher-level neural processes in lower areas[74]. We may have succeeded because we picked a single higher-level aspect of a percept and a well-documented property of visual cortex, so our strategy may be worth repeating.

To link motion-energy to anisotropy we found a combination of computer simulations and mathematical derivations to be extremely useful. For calculating the optic flow for rings rotating horizontally and vertically we used banks of motion-energy filters, and adjusted their spatial envelopes to compare the effects of V1 cortical anisotropy to isotropy. To understand the physical motions that best describe the optic flows, we mathematically derived the velocity fields of rings undergoing different weighted combinations of rotation and wobbling motion (given by the parameter k), and correlated them with the optic flows. For optic flows generated from an isotropic cortex, the best fitted weights were the same for horizontal and vertical rotation. However, the change in the optic flow caused by an anisotropic cortex predicted different physical combinations for horizontal and vertical rotations. Vertical rotation aligned closely with physical wobbling, while horizontal rotation aligned closer to physical rotation, matching the perceptions reported by our observers. We realized that correlating different templates for different motions to the optic flow from an object is computationally expensive and biologically implausible, not to mention that the templates would not generalize across object shapes. Therefore, we explored a more efficient and plausible model using differential invariants.

Differential invariants from kinematics have the advantage of not being restricted

to precise shapes, similar to direction-selective filters being more general than matching

specific features across frames, and they can be calculated with combinations of

biologically plausible filters. Consequently, we derived the differential invariants

mathematically for the physical combinations of rotation and wobbling and plotted their

values as a function of motion phase. Then we computed the values of the differential

invariants given by the optic flows obtained from isotropic and anisotropic cortex and

compared them to the values for different combinations of rotation and wobbling. The

results strengthen our contention that the anisotropy in nonrigidity arises from the

hardwired cortical anisotropy in numbers and tuning widths of direction selective

neurons. From our previous work, we learned that the optic flows from motion-energy

units can lead to seeing rigid objects as nonrigid, while the presence of trackable

features increases the probability of rigid percepts[4]. Figure 1H demonstrates that the

cortical anisotropy biases the optic flow enough to overcome the rigidity signals

provided by trackable features.

In summary, our study demonstrates that vertically rotating rings appear more

nonrigid than horizontally rotating rings. The shape of the vertically rotating rings seems

narrower and longer compared to the horizontally rotating rings. By matching the

shapes, observers were able to reduce perceived anisotropy by up to 30%, yet vertically

rotating rings were still perceived as more nonrigid after shapes were matched. We

incorporated anisotropy into our motion-energy computations and found that the

extracted optic flow patterns from anisotropic cortex align with templates for physical

wobbling for the vertically rotated rings but not for the horizontally rotated rings.

Matching to derived differential invariants as a biologically plausible model,

strengthened our conclusions.

Appendix

Stimulus Generation and projection

We created 3D circular, rotating/wobbling rings by applying the equations for rotation

along the $X, Y, and\ Z$ axes in a 3D space to all points on the rendered object. The

rotational matrix around each axis is given by:

$$\mathbb{R}_X(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & cos\theta & -sin\theta \\ 0 & sin\theta & cos\theta \end{pmatrix} \tag{A1}$$

$$\mathbb{R}_y(\theta) = \begin{pmatrix} cos\theta & 0 & sin\theta \\ 0 & 1 & 0 \\ -sin\theta & 0 & cos\theta \end{pmatrix} \tag{A2}$$

$$\mathbb{R}_Z(\theta) = \begin{pmatrix} cos\theta & -sin\theta & 0 \\ sin\theta & cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \tag{A3}$$

13   If $\vec{P}(\theta) = (cos\theta, sin\theta, 0)^T$ is the initial location of a 3-D point on the object lying on the

14   XY plane with radius 1, the position of the point $(\vec{P_r})$ on a rotating ring inclined at an

15   angle of τ from the ground plane and angular velocity of ω is expressed as:

$$\vec{P_r} = \mathbb{R}_y(\omega t)\mathbb{R}_Z(\tau)\vec{P}(\theta) \tag{A4}$$

19   The position of the point on the wobbling object will be:

$$\vec{P_w} = \mathbb{R}_y(\omega t)\mathbb{R}_Z(\tau)R_Z(-\omega t)\vec{P}(\theta) \tag{A5}$$

The operations of the rotational matrix for wobbling and rotation are illustrated in Figure

5 D. In Equation 10, if $k = 0$ in $\mathbb{R}_z(-k\omega t),\ then\ \mathbb{R}_z(-k\omega t)$ is the identity matrix. Thus,

Equation 10 provides a smooth transition from rotation to wobbling. Then, we projected

each point in perspective for the stimulus, but for the analysis, we used orthographic

projection. For the physical stretch, $\vec{P}(\theta) = (a * cos\theta, b * sin\theta, 0)^T$ a and b were changed

to make changes in aspect ratio that led to narrower or wider ellipses. Image stretches

were performed on the projected image.

To compute velocity fields, the time derivative was taken as shown in Equation 11.

6
$$F_k = \frac{\partial}{\partial t} \overrightarrow{P_k}(\theta, t)$$

8
$$= \left(S(0,1,0)R_y(\omega t)R_z(\tau)R_z(-k\omega t) + R_y(\omega t)R_z(\tau)S(0,-1,0)R_z(-k\omega t) \right) \overrightarrow{P_k}(\theta, t) \quad (A6)$$

11 where $S(0,1,0)$ is a skew symmetric matrix $\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}$.

29 \

Equation symbol lookup table

Symbol	Description
$\phi_v, \phi_h, \psi_v, \psi_h$	Angles in two elongated diamonds depicted in Figure 2C and D
k_i, μ_i	Kappa sets the tuning width and Mu the mean of the von Mises function for the i-th neuron
γ_h, γ_v	Angles around the horizontal axis and the vertical axis
θ_a, θ_b	Orientation of two lines that form an angle of γ
n_i	Number of cells for i-th neuron
r_{ai}, r_{bi}	Feedforward response of i-th cell given orientations of θ_a and θ_b respectively
g_i	Divisive gain control of i-th cell
R_{ai}, R_{bi}	Normalized response of i-th cell
$\widehat{\theta_a}, \widehat{\theta_b}$	Decoded orientations
$\widehat{\gamma_v}, \widehat{\gamma_h}$	Decoded angles given γ_h and γ_v respectively
$G_{i,odd}, G_{i,even}$	Odd and even 3D Gabor filters of i-th cell
Q_i	Normalization constant for the Gabor filter
$c_{x,i}, c_{y,i}, c_{t,i}$	Spatial and temporal means of the Gabor filter
$\sigma_{s,i}, \sigma_{t,i}$	Spatial and temporal envelopes of the Gabor filter
$\omega_{x0}, \omega_{y0}, \omega_{t0}$	Spatial and temporal frequencies of the sine component of the Gabor filter
η_s, η_t	Spatial and temporal orientations of the Gabor filter
F_0	Frequency of the Gabor filter
$\Delta\omega_{\frac{1}{2}}$	Half-height orientation bandwidths
$\hat{u}, \hat{v}$	The best estimate of local velocity components
$\mathfrak{N}_i(u, v)$	Predicted response of filter i to velocity (u, v)
m_i	i-th motion energy output
$\overline{m_i}, \overline{\mathfrak{N}_i}(u, v)$	The sum of $\mathfrak{N}_i$ and the sum of m_i for filters sharing the same spatial orientation
$\mathbb{R}_y, \mathbb{R}_z$	Rotational matrices around Y and Z axis respectively

$\vec{P}(\theta)$	The position of a point on the ring lying on the $X - Z$ plane
τ	Tilt of the ring
ω	Angular velocity of the rotation of the ring
k	The wobbling weight $(k = 0 \sim rotation, k = 1 \sim wobbling)$
$Proj(\cdot)$	The projection function
$\overrightarrow{P_k}(\theta, t)$	The position of a point at a space and time for a moving ring with the wobbling weight of k
$u(\theta, t), v(\theta, t)$	The x and y components for $\overrightarrow{P_k}(\theta, t)$
F_k	The velocity field for the moving ring with the wobbling weight of k

References

1. Tomasi, C., & Kanade, T. (1992). Shape and motion from image streams under orthography: a factorization method. International Journal of Computer Vision, 9, 137-154.

2. Aggarwal, J. K., Cai, Q., Liao, W., & Sabata, B. (1998). Nonrigid motion analysis: Articulated and elastic motion. Computer Vision and Image Understanding, 70(2), 142-156.

3. Jain, A., & Zaidi, Q. (2011). Discerning nonrigid 3D shapes from motion cues. Proceedings of the National Academy of Sciences, 108(4), 1663-1668.

4. Maruya, A., & Zaidi, Q. (2024). Perceptual transitions between object rigidity and non-rigidity: Competition and cooperation among motion energy, feature tracking, and shape-based priors. *Journal of Vision*, *24*(2), 3-3.

5. Lu, Z. L., & Sperling, G. (2001). Three-systems theory of human visual motion perception: review and update. JOSA A, 18(9), 2331-2370.

6. Zaidi, Q., & DeBonet, S. J. (2000). Motion energy versus position tracking: Spatial, temporal and chromatic parameters. Vision Research, 40(26), 3613-3635.

7. Shiffrar, M., & Pavel, M. (1991). Percepts of rigid motion within and across apertures. Journal of Experimental Psychology: Human Perception and Performance, 17(3), 749.

8. Lorenceau, J., & Shiffrar, M. (1992). The influence of terminators on motion integration across space. Vision Research, 32(2), 263–273.

9. Mach, E. (1886). Beiträge zur Analyse der Empfindungen. Jena: Gustav Fischer. In: Williams, C. M. (Trans.), English translation: Contributions to the analysis of the sensations (p. 1897). Chicago: The Open Court.

10. Weiss, Y., & Adelson, E. H. (2000). Adventures with gelatinous ellipses—constraints on models of human motion analysis. Perception, 29(5), 543–566.

11. Rokers, B., Yuille, A., & Liu, Z. (2006). The perceived motion of a stereokinetic stimulus. Vision Research, 46(15), 2375–2387.

12. Vezzani, S., Kramer, P., & Bressan, P. (2014). Stereokinetic effect, kinetic depth effect, and structure from motion. In Wagemans, J. (Ed.), The Oxford handbook of perceptual organization (pp. 521–540). Oxford, UK: Oxford Library of Psychology.

13. Burge, T. (2010). Origins of objectivity. Oxford University Press.

14. Oktaviana, A. A., Joannes-Boyau, R., Hakim, B., Burhan, B., Sardi, R., Adhityatama, S., ... & Aubert, M. (2024). Narrative cave art in Indonesia by 51,200 years ago. Nature, 1-5.

15. Kendall D, Barden D, Carne T, Le H (1999) Shape and shape theory (Wiley, New York).

16. Griffiths, A. F., & Zaidi, Q. (2000). Perceptual assumptions and projective distortions in a three-dimensional shape illusion. Perception, 29(2), 171-200.

17. Dvoeglazova, M., & Sawada, T. (2024). A role of rectangularity in perceiving a 3D shape of an object. Vision Research, 221, 108433.

18. Pizlo, Z., & De Barros, J. A. (2021). The concept of symmetry and the theory of perception. Frontiers in Computational Neuroscience, 15, 681162.

19. Cohen EH, Zaidi Q. Fundamental failures of shape constancy resulting from cortical anisotropy. J Neurosci. 2007 Nov 14; 27(46):12540-5. doi: 10.1523/JNEUROSCI.4496-07.2007. PMID: 18003832; PMCID: PMC2843153.

20. Li B, Peterson MR, Freeman RD (2003) Oblique effect: a neural basis in the visual cortex. J Neurophysiol 90:204–217.

21. Girshick, A. R., Landy, M. S., & Simoncelli, E. P. (2011). Cardinal rules: visual orientation perception reflects knowledge of environmental statistics. Nature Neuroscience, 14(7), 926-932.

22. Harrison, W. J., Bays, P. M., & Rideaux, R. (2023). Neural tuning instantiates prior expectations in the human visual system. Nature Communications, 14(1), 5320.

23. Appelle, S. (1972). Perception and discrimination as a function of stimulus orientation: The "oblique" effect in man and animals. Psychological Bulletin, 78, 266–278.

24. Fang, C., Cai, X., & Lu, H. D. (2022). Orientation anisotropies in macaque visual areas. Proceedings of the National Academy of Sciences, 119(15), e2113407119.

25. Coletta, N. J., Segu, P., & Tiana, C. L. (1993). An oblique effect in parafoveal motion perception. Vision Research, 33(18), 2747-2756.

26. Gros, B. L., Blake, R., & Hiris, E. (1998). Anisotropies in visual motion perception: a fresh look. JOSA A, 15(8), 2003-2011.

27. Dakin, S. C., Mareschal, I., & Bex, P. J. (2005). An oblique effect for local motion: Psychophysics and natural movie statistics. Journal of vision, 5(10), 9-9.

28. Steven C. Dakin, Isabelle Mareschal, Peter J. Bex; An oblique effect for local motion: Psychophysics and natural movie statistics. Journal of Vision 2005; 5(10):9. https://doi.org/10.1167/5.10.9.

29. Rokem, A., & Silver, M. A. (2009). A model of encoding and decoding in V1 and MT accounts for motion perception anisotropies in the human visual system. Brain research, 1299, 3-16.

30. Maunsell, J. H., & Van Essen, D. C. (1983). Functional properties of neurons in middle temporal visual area of the macaque monkey. I. Selectivity for stimulus direction, speed, and orientation. Journal of neurophysiology, 49(5), 1127-1147.

31. Churchland, A. K., Gardner, J. L., Chou, I. H., Priebe, N. J., & Lisberger, S. G. (2003). Directional anisotropies reveal a functional segregation of visual motion processing for perception and action. Neuron, 37(6), 1001-1011.

32. Bigelow, A., Kim, T., Namima, T., Bair, W., & Pasupathy, A. (2023). Dissociation in neuronal encoding of object versus surface motion in the primate brain. Current Biology, 33(4), 711-719.

33. Chariker, L., Shapley, R., Hawken, M., & Young, L. S. (2022). A computational model of direction selectivity in Macaque V1 cortex based on dynamic differences between ON and OFF pathways. Journal of Neuroscience, 42(16), 3365-3380.

34. Todorović, D. (1993). Analysis of two-and three-dimensional rigid and nonrigid motions in the stereokinetic effect. JOSA A, 10(5), 804-826.

35. Swindale, N. V. (1998). Orientation tuning curves: empirical description and estimation of parameters. Biological cybernetics, 78, 45-56.

36. DeAngelis, G. C., Freeman, R. D., & Ohzawa, I. Z. U. M. I. (1994). Length and width tuning of neurons in the cat's primary visual cortex. Journal of neurophysiology, 71(1), 347-374.

37. Slllito, A. M., Grieve, K. L., Jones, H. E., Cudeiro, J., & Davls, J. (1995). Visual cortical mechanisms detecting focal orientation discontinuities. Nature, 378(6556), 492-496.

38. Shushruth, S., Nurminen, L., Bijanzadeh, M., Ichida, J. M., Vanni, S., & Angelucci, A. (2013). Different orientation tuning of near-and far-surround suppression in macaque primary visual cortex mirrors their tuning in human perception. Journal of Neuroscience, 33(1), 106-119.

39. Schwartz, O., & Simoncelli, E. P. (2001). Natural signal statistics and sensory gain control. Nature Neuroscience, 4(8), 819-825.

40. Georgopoulos, A. P., Schwartz, A. B., & Kettner, R. E. (1986). Neuronal population coding of movement direction. Science, 233(4771), 1416-1419.

41. Schwartz, O., Sejnowski, T. J., & Dayan, P. (2009). Perceptual organization in the tilt illusion. Journal of Vision, 9(4), 19-19.

42. Mach, E. (1897). Contributions to the Analysis of the Sensations. Open Court.

43. Watson, A. B., & Ahumada, A. (1983). A look at motion in the frequency domain. Silicon Valley, CA: Ames Research Center, National Aeronautics and Space Administration.

44. Adelson, E. H., & Bergen, J. R. (1985). Spatiotemporal energy models for the perception of motion. Journal of the Optical Society of America A, 2(2), 284–299.

45. Watson, A. B., & Ahumada, A. J. (1985). Model of human visual-motion sensing. Journal of the Optical Society of America A, 2(2), 322–342.

46. Van Santen, J. P., & Sperling, G. (1985). Elaborated Reichardt detectors. Journal of the Optical Society of America A, 2(2), 300–321.

47. Rust, N. C., Mante, V., Simoncelli, E. P., & Movshon, J. A. (2006). How MT cells analyze the motion of visual patterns. Nature Neuroscience, 9(11), 1421–1431.

48. Bradley, D. C., & Goyal, M. S. (2008). Velocity computation in the primate visual system. Nature Reviews Neuroscience, 9(9), 686–695.

49. Nishimoto, S., & Gallant, J. L. (2011). A three-dimensional spatiotemporal receptive field model explains responses of area MT neurons to naturalistic movies. Journal of Neuroscience, 31(41), 14551–14564.

50. Nishimoto, S., Vu, A. T., Naselaris, T., Benjamini, Y., Yu, B., & Gallant, J. L. (2011). Reconstructing visual experiences from brain activity evoked by natural movies. Current Biology, 21(19), 1641–1646.

51. Movellan, J. R. (2002). Tutorial on Gabor filters. Open source document, 40, 1-23.

52. Heeger, D. J. (1988). Optical flow using spatiotemporal filters. International journal of computer vision, 1, 279-302.

53. Koenderink, J. J., & van Doorn, A. J. (1976). Local structure of movement parallax of the plane. JOSA, 66(7), 717-723.

54. Longuet-Higgins, H. C., & Prazdny, K. (1980). The interpretation of a moving retinal image. Proceedings of the Royal Society of London. Series B. Biological Sciences, 208(1173), 385-397.

55. Koenderink, J. J. (1986). Optic flow. Vision Research, 26(1), 161-179.

56. Zhang, K., Sereno, M. I., & Sereno, M. E. (1993). Emergence of position-independent detectors of sense of rotation and dilation with Hebbian learning: an analysis. Neural Computation, 5(4), 597-612.

57. Duffy, C. J., & Wurtz, R. H. (1997). Medial superior temporal area neurons respond to speed patterns in optic flow. Journal of Neuroscience, 17(8), 2839-2851.

58. Saito, H. A., Yukie, M., Tanaka, K., Hikosaka, K., Fukada, Y., & Iwai, E. (1986). Integration of direction signals of image motion in the superior temporal sulcus of the macaque monkey. Journal of Neuroscience, 6(1), 145-157.

59. Lappe, M., Bremmer, F., Pekel, M., Thiele, A., & Hoffmann, K. P. (1996). Optic flow processing in monkey STS: a theoretical and experimental approach. Journal of Neuroscience, 16(19), 6265-6285.

60. Anderson, K. C., & Siegel, R. M. (1999). Optic flow selectivity in the anterior superior temporal polysensory area, STPa, of the behaving monkey. Journal of Neuroscience, 19(7), 2681-2692.

61. Sugihara, H., Murakami, I., Shenoy, K. V., Andersen, R. A., & Komatsu, H. (2002). Response of MSTd neurons to simulated 3D orientation of rotating planes. Journal of Neurophysiology, 87(1), 273-285.

62. Freeman, T. C., Harris, M. G., & Meese, T. S. (1996). On the relationship between deformation and perceived surface slant. Vision Research, 36(2), 317-IN1.

63. Domini, F., & Caudek, C. (2003). 3-D structure perceived from dynamic information: A new theory. Trends in Cognitive Sciences, 7(10), 444-449.

64. He, D.; Nguyen, D.T.; Ogmen, H.; Nishina, S.; Yazdanbakhsh, A. Perception of Rigidity in Three- and Four-Dimensional Spaces. Front. Psychol. 2023, 14, 1180561.

65. Ullman, S. (1984). Maximizing rigidity: the incremental recovery of 3-D structure from rigid and nonrigid motion. Perception, 13(3), 255-274. doi: 10.1068/p130255. PMID: 6514511.

66. Sinha, P., & Poggio, T. (1996). Role of learning in three-dimensional form perception. Nature, 384(6608), 460-463.

67. Johansson, G. (1973). Visual perception of biological motion and a model for its analysis. Perception & psychophysics, 14, 201-211.

68. Choi, R., Feldman, J., & Singh, M. (2024). Perceptual Biases in the Interpretation of Non-Rigid Shape Transformations from Motion. Vision, 8(3), 43.

69. Sparrow, J. E., & Stine, W. W. (1998). The perceived rigidity of rotating eight-vertex geometric forms: extracting nonrigid structure from rigid motion. Vision Research, 38(4), 541-556.

70. Cohen, E. H., Jain, A., & Zaidi, Q. (2010). The utility of shape attributes in deciphering movements of non-rigid objects. Journal of Vision, 10(11), 29-29.

71. Chuang, L. L., Vuong, Q. C., & Bülthoff, H. H. (2012). Learned non-rigid object motion is a view-invariant cue to recognizing novel objects. Frontiers in Computational Neuroscience, 6, 26.

72. Braunstein, M. L., Hoffman, D. D., & Pollick, F. E. (1990). Discriminating rigid from nonrigid motion: Minimum points and views. Perception & Psychophysics, 47(3), 205-214.

73. Vuong, Q. C., Friedman, A., & Read, J. C. (2012). The relative weight of shape and non-rigid motion cues in object perception: A model of the parameters underlying dynamic object discrimination. Journal of Vision, 12(3), 16-16.

74. Xiao, W., Sharma, S., Kreiman, G., & Livingstone, M. S. (2024). Feature-selective responses in macaque visual cortex follow eye movements during natural vision. Nature Neuroscience, 1-10.

Figure 1. Anisotropy of a rotating ring illusion videos. A: Two Styrofoam rings glued together at an angle are seen to rotate together on the turntable. B: When rotated 90°, one ring appears to move independently and wobble against the ring on the turntable. C & D: Two circular (C) and octagonal (D) rings physically rotate together with a fixed connection at the junction. E & F: The two pairs of rings physically move independently and wobble against each other. For the circular rings, E is indistinguishable from C. However, with trackable vertices, F is discernable from D. G & H are the same as C & D except that they are rotated 90° and they both look nonrigidly connected.

Figure 2. Anisotropy of the shape of the ring. A and B correspond to snapshots from Figure 1E and 1F respectively. Despite being physically identical shapes, B is perceived as vertically elongated and narrower in comparison to A. C and D: The shape illusion is similar for two elongated diamonds and can be explained by differences in perceived angles: ϕ_h is perceived to be wider than ϕ_v despite being physically equal, and ψ_h is perceived to be wider than ψ_v.

Figure 3. Cortical anisotropy. A: Orientation tuning curves of V1 simple cells were simulated by von Mises distributions to match the anisotropic and isotropic cortices and weighted by the number of cells (See text for references). B: Detailed parameters for the von Mises distributions are shown on the left two panels for the anisotropic (red) and isotropic (black) cortex. Decoded angles around the vertical and horizontal diverging axes respectively are shown on the right two panels. C: The decoded angle difference ($\hat{\gamma}_H - \hat{\gamma}_V$) from the anisotropic cortex (red) and the isotropic cortex (dotted black). For the isotropic cortex, there is no difference in the decoded angles for γ_V and γ_H. However, for the anisotropic cortex, γ_H is decoded to be broader than γ_V.

Horizontal stretch

A

0% stretch   50% stretch

Image stretch

Physical stretch

Image stretch

Physical stretch

B

Matching horizontally rotating shapes

C

Matching vertically rotating shapes

D

Figure 4. Effect of image and physical stretch on perceived nonrigidity. A: Examples of rings elongated to match the shape of orthogonally oriented circular rings. Subjects were allowed to stretch rings in image domain (first and third rows) or physically (second and fourth rows), ranging from 0% (left column) to 50% (right column). B: Histograms representing the extent of horizontal stretch in the image domain (red) and physical domain (blue) for vertically rotating rings adjusted to match the shape of physically circular horizontally rotating rings. C: Histograms representing the extent of horizontal stretch in the image domain (red) and physical domain (blue) for horizontally rotating rings adjusted to match the shape of physically circular vertically rotating rings. D: Probability of observers reporting vertically rotating rings as more nonrigid. Original: Two pairs of circular rings (black). H indicates that horizontally rotating rings were stretched to match shapes and V indicates that vertically rotating rings were stretched, either in the image (red) or physically before projection (blue). Max: Both vertical and horizontal rings were stretched maximally like in the last column of Panel A, so horizontally rotating rings were perceived as narrower and longer than vertically rotating rings.

Figure 5. Comparing optic flows from isotropic cortex and anisotropic cortex to templates for rotation and wobbling. A: Tuning curves for direction selective cells reflecting documented anisotropies in width and number of cells. B: Tuning curves for direction selective cells for an isotropic cortex. C: Optic flow fields generated by isotropic cortex (top row) and anisotropic cortex (bottom row) for physical horizontal rotation (left column) and vertical rotation (right column). Anisotropic cortex generates vectors pointing horizontally. D: Depiction of rotation and wobbling axes. E: Physical velocity field for horizontal (top) and vertical (bottom) rotations mixed with physical wobbling (Weights k=0-1). F & G Cosine similarity of ME optic flows from isotropic (F) and anisotropic (G) cortex to different templates (k=0: physical rotation, k=1: physical wobbling, shown in E). Isotropic cortex leads to no difference between best fitting k for horizontal (blue) and vertical (red) rotation. However, cortical anisotropy results in different best fitting k: more wobbling for vertical rotation.

A

Divergence   Curl   Deformation 1   Deformation 2

B

Gradients of rotation-wobbling

Div

Curl

Def

1 (wobbling)

k

0 (rotation)

Motion Phase (Deg)

C

Isotropic tuning

Vertical   Horizontal

Gradients

Divergence
Curl
Deformation

$k = 0.78$ $k = 0.78$

D

Anisotropic tuning

Vertical   Horizontal

Gradients

$k = 0.91$ $k = 0.48$

Motion Phase (Deg)

E

Isotropic Cortex

Best k

H-rotation
V-rotation

Amount of H-stretch

F

Anisotropic Cortex

Best k

H-rotation
V-rotation

Amount of H-stretch

Figure 6. Gradients of ME flow from the rotating ring align with percepts. A: Examples of differential invariants. B: Gradients of rotation to wobbling, divergence (left), curl (middle), and deformatiosn1 (right) as a function of rotational phase (0°-180°: half a cycle). These gradients illustrate that with increasing emphasis on wobbling from k=0 (blue curves) to k=1 (red curves), variability increases across all three gradients, while rotational fields consistently exhibit higher values of curl and deformation. C: Isotropic Tuning Gradients for Vertical (left) and Horizontal (right) Rotation: Similar gradients are observed for both orientations, with optimal fitting by curves at a wobbling weight of k=0.78. D: Upon adjusting filter numbers and tuning widths to match V1 anisotropies before velocity field calculations, gradients for vertical rotation closely correspond to physical wobbling (k=0.91), while those for horizontal rotation indicate lower levels of wobbling (k=0.48). E & F : Best k as a function of image stretch. E for isotropic cortex and F for anisotropic cortex. A greater amount of stretch leads to increased wobbling during horizontal rotation (red) and reduced wobbling during vertical rotation (blue). For an anisotropic cortex, the higher amount of wobbling during vertical rotation is maintained, consistent with experimental results. Furthermore, at maximum stretch, the isotropic and anisotropic cortices predict different outcomes: the isotropic cortex suggests that horizontal rotation should be perceived as more nonrigid, while the anisotropic cortex suggests that vertical rotation should appear more nonrigid.

Figure S1. Individual results for the shape matching experiment. For each observer, left histograms represent the extent of horizontal stretch in the image domain (red) and physical domain (blue) for vertically rotating rings adjusted to match the shape of physically circular horizontally rotating rings. Right histograms represent the extent of horizontal stretch in the image domain (red) and physical domain (blue) for horizontally rotating rings adjusted to match the shape of physically circular vertically rotating rings.

Figure S2. Individual results for the nonrigidity comparison. Probability of each observer reporting vertically rotating rings as more nonrigid. Original: Two pairs of circular rings (black). H indicates that horizontally rotating rings were stretched to match shapes, and V indicates that vertically rotating rings were stretched, either in the image (red) or physically before projection (blue). Max: Both vertical and horizontal rings were stretched maximally like in the last column of Panel A in Figure 4, so horizontally rotating rings were perceived as narrower and longer than vertically rotating rings