Intergenerational transmission of the structure of the auditory cortex and reading skills

Olga Kepinska[1,2*], Florence Bouhali[3], Giulio Degano[4], Raphael Berthele[5], Hiroko Tanaka[6,7], Fumiko Hoeft[8,9,10], Narly Golestani[1,2,4]

1. Brain and Language Lab, Cognitive Science Hub, University of Vienna, Vienna, Austria.
2. Department of Behavioral and Cognitive Biology, Faculty of Life Sciences, University of Vienna, Vienna, Austria.
3. Aix Marseille Univ, CNRS, CRPN, Marseille, France.
4. Department of Psychology, Faculty of Psychology and Educational Sciences, University of Geneva, Geneva, Switzerland.
5. Institute of Multilingualism, University of Fribourg, Fribourg, Switzerland.
6. Department of Pediatrics, College of Medicine, University of Arizona, Tucson, AZ, USA.
7. Banner University Medical Center – Tucson, Tucson, AZ, USA.
8. Department of Psychological Sciences, University of Connecticut, Storrs, CT, USA.
9. Brain Imaging Research Center, University of Connecticut, Storrs, CT, USA.
10. Departments of Mathematics, Neuroscience, Psychiatry, Educational Psychology, Pediatrics, Computer Science and Engineering, University of Connecticut, Storrs, CT, USA.

* corresponding author

ABSTRACT. High-level cognitive skill development relies on genetic and environmental factors, tied to brain structure and function. Inter-individual variability in language and music skills has been repeatedly associated with the structure of the auditory cortex: the shape, size and asymmetry of the transverse temporal gyrus (TTG) or gyri (TTGs). TTG is highly variable in shape and size, some individuals having one single gyrus (also referred to as Heschl's gyrus, HG) while others presenting duplications (with a common stem or fully separated) or higher-order multiplications of TTG. Both genetic and environmental influences on children's cognition, behavior, and brain can to some to degree be traced back to familial and parental factors. In the current study, using a unique MRI dataset of parents and children (135 individuals from 37 families), we ask whether the anatomy of the auditory cortex is related to reading skills, and whether there are intergenerational effects on TTG(s) anatomy. For this, we performed detailed, automatic segmentations of HG and of additional TTG(s), when present, extracting volume, surface area, thickness and shape of the gyri. We tested for relationships between these and reading skill, and assessed their degree of familial similarity and intergenerational transmission effects. We found that volume and area of all identified left TTG(s) combined was positively related to reading scores, both in children and adults. With respect to intergenerational similarities in the structure of the auditory cortex, we identified structural brain similarities for parent-child pairs of the 1st TTG (Heschl's gyrus, HG) (in terms of volume, area and thickness for the right HG, and shape for the left HG) and of the lateralization of all TTG(s) surface area for father-child pairs. Both the HG and TTG-lateralization findings were significantly more likely for parent-child dyads than for unrelated adult-child pairs. Furthermore, we established characteristics of parents' TTG that are related to better reading abilities in children: fathers' small left HG, and a small ratio of HG to planum temporale. Our results suggest intergenerational transmission of specific structural features of the auditory cortex; these may arise from genetics and/or from shared environment.

1   Introduction

The development of high-level cognitive skills invariably depends on a complex interplay between genetic and environmental factors, and is related to brain structural and functional indices. In the context of brain-behavior relations, inter-individual variability in reading, language, and musical abilities has been repeatedly associated with the structure of the auditory cortex, and in particular with the shape, size and asymmetry of the transverse temporal gyrus (TTG) or gyri (TTGs), if there are several (Benner et al., 2017; Turker et al., 2021). The TTG(s) is (are) located within the Sylvian fissure, on the superior surface of the superior temporal gyrus. The most anterior TTG, or the only TTG in the case of a single gyrus, is also known as Heschl's gyrus (HG) and includes the primary auditory cortex (von Economo and Horn, 1930). Additional TTGs, if present, are part of the planum temporale (PT), which houses the secondary auditory cortex, although the mapping between cytoarchitecture and cross anatomy is not one-to-one. The TTG(s) show(s) high variability in shape and size between hemispheres and individuals: there may be a single gyrus (HG), duplications (with a common stem or fully separated) or multiplications of the TTGs (Geschwind and Levitsky, 1968; Marie et al., 2015). This variability in shape but also in size, has in turn, been frequently associated with individual differences in a range of auditory, language and music-related skills, and with a number of disorders. These include phonetic learning skill (Golestani et al., 2007a, 2002) and expertise (Golestani et al., 2011), linguistic pitch learning ability (Wong et al., 2008), musical expertise (Benner et al., 2017) and language aptitude (Ramoser et al., *under review*; Turker et al., 2019, 2017). These individual differences can arise from influences both in predisposition and in experience-dependent plasticity, as suggested by earlier work in phonetics experts (Golestani et al., 2011), by our recent work in multilinguals (Kepinska et al., 2023), and in the neurocognitive model of language aptitude proposed by Turker and colleagues (2021). TTG(s) anatomy has also been associated with pathological conditions such as tinnitus (Schneider et al., 2009) and schizophrenia (Takahashi et al., 2022), as well as with (poor) pre-reading and reading skills (Altarelli et al., 2014; Blockmans et al., 2023; Kuhl et al., 2020; Leonard et al., 2001; Serrallach et al., 2016; Sutherland et al., 2012). The latter is the focus of the present study.

Reading skill and disorder have been associated with variations in several structural features of the TTG(s) (gyrification patterns, volume, thickness, surface area, asymmetry, etc.), and as such, there is no clear consensus on which specific anatomical feature is most explanatory of individual differences in reading ability. For example, dyslexia has been associated with differences in gross shape, with more TTG duplications in dyslexic children in the right hemisphere only (Altarelli et al., 2014; Serrallach et al., 2016, with the former study showing these results only in boys), whereas a similar result has previously been found in the left hemisphere, in adults (Leonard et al., 2001). Kuhl and colleagues (2020) showed that a higher local gyrification index around the left HG (reflecting the presence of more involutions and buried

cortex) in pre-reading children was predictive of a dyslexia diagnosis three years later. The predictive value

of HG surface area and of TTG duplication patterns for later reading was also recently demonstrated by

Blockmans and colleagues (2023)). They found that greater surface area of the left HG and of left and right

PT, and more left TTG multiplications in pre-reading children were associated with better word reading

three years later. With respect to TTG(s) cortical thickness, Clark et al. (2014) found that pre-reading 6-

year-olds with thinner HG cortex were more likely to develop dyslexia, while Ma et al. (2015) reported that

school-aged children with dyslexia had a thicker right HG and surrounding areas (including superior

temporal gyrus and PT). While some inconsistencies remain, these findings overall align with observations

of pervasive deficits in the representation and processing of speech sounds in dyslexia. Such deficits are

considered by many to be one of the factors at the origin of dyslexia, and the above results show that

these can be traced back to auditory cortex neuroanatomical features (Linkersdörferet al., 2012). Indeed,

Sutherland et al. (2012) found a higher gray-matter probability in the left HG as a neuroanatomical marker

underlying the link between literacy and auditory processing in the temporal domain (again, a relationship

that was present in boys but not in girls).

Both genetic and environmental influences on children's cognition, behavior, and brains can, to some

extent, be traced back to familial and parental factors. Van Bergen (2014) suggests that "both parents

confer [dyslexia] liability via intertwined genetic and environmental pathways", and points to

intergenerational designs as a promising avenue of research into the etiology of the disorder. Given that

TTG morphology has been associated with experiential factors such as bi- and multilingualism (Kepinska

et al., 2023; Ressel et al., 2012) but that it has also been shown to have moderate to high heritability (Eyler

et al., 2012; Grasby et al., 2020), studying the anatomical features of the TTG(s) across different family

members and generations may allow us to better understand the familial transmission (via shared

environment and/or of genes) of brain structure and of associated phenotypes. Here, we address this

question using a unique brain imaging and behavioral dataset from parents and their children.

Understanding the intergenerational, familial transmission of brain structure by examining familial similarity

may shed light on the mechanisms of inheritance of complex behavioral traits (Ho et al., 2016), although

similarities cannot be pinpointed to genetic inheritance specifically since they could also arise from shared

experiential influences. Indeed, studies are beginning to unravel the concordance of brain structural indices

across family members, including mother-daughter similarities in whole-brain sulcal morphology (Ahtam et

al., 2021), corticolimbic circuitry (Dimanova et al., 2023; Yamagata et al., 2016), reading network anatomy

(Fehlbaum et al., 2022), and parent-child similarity in whole-brain functional and structural measures

(Takagi et al., 2021).

Due to the variability in its morphology, the TTG(s) has been notoriously difficult to automatically segment accurately, with template-based analyses largely ignoring individual variability in its shape (Dalboni da Rocha et al., 2020). Recent developments in cortical segmentation of the TTG(s) (Dalboni da Rocha et al., 2023, 2020) provide an anatomically precise, reliable and automatic (i.e., without manual segmentation or subjective decisions) method for delineating the region, and deriving continuously quantified measures describing its shape and size (indexed by volume, surface area and thickness). The present study capitalizes on these recent efforts in cortical segmentation. We aim to determine the degree of familial similarity and intergenerational transmission effects on the anatomy of the auditory cortex in a group of 135 individuals from 37 families, and to identify the precise anatomical features of the TTG(s) that are related to reading ability in children and adults, and across generations (i.e., are there any specific anatomical features of the parents' TTG(s) that predict children's reading ability?).

2 Methods

2.1 Participants

MRI data from a total of 135 individuals, including 72 children (M_{age} = 8.81, SD = 2.11, 35 female), and 63 adults (M_{age} = 42.5, SD = 5.14, 33 female) were analyzed. In total, there were 37 biological families (65 mother-child, 60 father-child dyads; most families had more than one child), see Table S1 for an overview of the families. Twelve more children were scanned but their data were excluded due to excessive motion and bad segmentation (see Section 2.4 for details). The majority of the participants were right-handed (112:16:3 right:left:ambidextrous ratio); handedness was dummy coded for the analyses, and in 4 cases where the information was missing it was replaced by the mean. Parents' socioeconomic status (SES) was indexed by the years of education they completed ($M_{mothers}$ = 17.08, SD = 2.06; $M_{fathers}$ = 16.80, SD = 2.07); for children, we used an average of the years of completed education of both parents (M_{SES} = 16.94, SD = 1.74).

2.2 Behavioral data

Reading and reading-related test scores were available for 129 participants (70 children and 59 adults). These included:

(1) Sight Word Efficiency (T-SWE) subtest from the Test of Word Reading Efficiency (TOWRE) (Torgesen et al., 1999), in which as many as possible increasingly difficult items from a list of high frequency words including irregular grapheme to phoneme mappings have to be read aloud within 45 seconds;

(2) Phonemic Decoding Efficiency (T-PDE) subtest from TOWRE (Torgesen et al., 1999), in which as many as possible increasingly difficult items from a list of high frequency words including pseudo-words have to be read aloud within 45 seconds;

(3) Rapid Automatized Naming and Rapid Alternating Stimulus Tests (RAN/RAS) - Letters (RN-LTR)

subtest (Wolf and Denckla, 2005), where participants are asked to quickly name fifty individual letters

aloud;

(4) Woodcock Reading Mastery Tests Revised-Normative Update (WRMT) (Woodcock, 1998): Word

Identification (WRMT-WID), consisting of untimed reading of increasingly difficult individual low

frequency words;

(5) WRMT Word Attack (WRMT-WA) consisting of untimed reading of increasingly difficult individual

pseudo-words.

Standard scores[1] on individual tests were used in the analyses reported in Sections 3.2, 3.3, and 3.5 (see

Figure 2).

2.3 Data collection procedures

All procedures were approved by the Stanford University Panel on Human Subjects in Medical Research

and they were conducted in accordance with its guidelines and regulations. Written informed consent and

assent were obtained from parents and children, respectively, after complete description of the study to

the participants. Neuroimaging data were collected using a 3T GE-Signa HDxt scanner (GE Healthcare)

with a quadrature head coil at the Lucas Center for Imaging at Stanford University. High-resolution T1-

weighted anatomical images were acquired with a matrix size of 256*256 and voxel size of 0.86 x 0.86 x

1.2 mm; 156 axial slices; TR = 8.5 ms, TE = 3.4 ms, inversion time = 400 ms; flip angle = 15°; FOV = 22

cm. Images were visually inspected for scanner artefacts and anatomical anomalies.

2.4 Segmentation

The T1 images were processed with FreeSurfer's (version 7.1) structural sub-millimeter pipeline (`recon-`

`all`) (Fischl et al., 2004; Zaretskaya et al., 2018), consisting of motion correction, intensity normalization,

skull stripping, and reconstruction of the volume's voxels into white and pial surfaces. FreeSurfer's output

was visually inspected for segmentation errors, which were caused by excessive motion. We then

performed a detailed segmentation of the auditory cortices using an automated toolbox (TASH; Dalboni

da Rocha et al., 2020). For morphometric measures, we used `TASH_complete`, which segments and

quantifies the cortical structure of all TTG(s) that are identified (Dalboni da Rocha et al., 2023). We

performed a visual selection of the gyri segmented by `TASH_complete`, retaining for the analysis only gyri

that had a similar orientation as the first TTG (i.e. which we will henceforth refer to as 'HG') and excluding

---

[1] Note that normative data of the TOWRE and RAN/RAS subtests are only available for ages 6 to 24, and 5 to 18, respectively. Therefore, the standard scores for the available oldest age range reference were used for adult participants.

gyri along the portion of the superior temporal plane that curved vertically (i.e., within the parietal extension,

Honeycutt et al., 2000), when present. From the resulting labels, we exported measures (in native space)

of cortical (i.e. grey matter) volume (in mm^3), surface area (in mm^2) and mean thickness (in mm). To derive

measures of the shape of TTG(s), we used another toolbox, the Multivariate Concavity Amplitude Index

(Dalboni da Rocha et al., 2023), which calculates the degree of concavity of each of the gyri segmented

by TASH. Following Dalboni da Rocha et al. (2023), lateral concavity values were used in the analysis. We

used both an index reflecting the shape of HG alone, and one reflecting the overall shape of all identified

TTG(s) ('lateral multiplication index', Dalboni da Rocha et al., 2023). The latter was derived by counting all

gyri identified by `TASH_complete`, and adding the number of gyri to the lateral concavity values summed

across all present TTGs in the respective hemisphere. Higher values for both measures (on HG and on all

TTGs) indicate a more complex shape of the TTG(s), e.g., more duplications/multiplications of the TTG (see

examples on Figure 1).

2.5 Data analysis

In the statistical analyses, we used two sets of neuroanatomical measures: (1) ones describing only the

HG and (2) ones describing all identified TTG(s). Examples of these measures from two participants are

shown in Figure 1 below. In addition to the measures of volume, surface area, average thickness and shape

of HG and of the TTG(s) per hemisphere, we also computed their lateralization indices (LI). These were

calculated using the formula `[left - right]/[left + right]`, with positive values indicating

leftward asymmetry. When computing lateralization across all TTG(s), we used the following hemispheric

summary measures: (a) for volume and area, we computed the sum of volume and area across all TTG(s)

per hemisphere; (b) for thickness, we took the average thickness across the gyri per hemisphere, and (c)

for shape, we used the lateral multiplication index, as described above (Section 2.3).

Linear mixed models were used to determine:

(1) the relationships between children's reading scores and parents' reading scores (Section 3.2);

(2) the relationships between TTG(s) anatomy and reading scores (Section 3.3);

(3) the familial similarity and intergenerational transmission of TTG anatomy (Section 3.4).

In all analyses, we included covariates of no-interest: age, sex, and SES. Analyses with all brain structural

indices as dependent variables included additionally participants' handedness and a quadratic term for

age. We accounted for different brain sizes in two ways: total intracranial volume was included as a

covariate of no-interest in analyses where TTG volume, area and thickness were either an independent or

dependent variable; in analyses of intergenerational similarity effects (i.e., where children's anatomical

measures were modeled as a functions of measures of their parents), we normalized the anatomical

measures for the corresponding whole hemisphere measures (hemispheric volume, surface area and mean
hemispheric thickness) so as to be more comparable between parents and children. To account for sibling
relationships (i.e., non-independent observations within the data) family index was included as random
intercept. In models testing for intergenerational transmission, likelihood ratio tests were used to compare
models that included parental measures with reduced models with covariates of no-interest only. Analyses
were conducted in R (R Development Core Team, 2015). All significant intergenerational similarity effects
were followed up with permutation tests to confirm that they were due to true familial relationships, rather
than to spurious, generic correlations in our population. We computed the same models on 5,000
permutations of the family index, shuffling only the behavioral or anatomical variables of the parents with
respect to those of the children, while avoiding random pairings across siblings.

In addition, a *random forest* classifier was used to determine:

(4) which, if any, of the parents' anatomical measures could predict children's reading (Section 3.5).

Here, we used the `cforest()` function in the `party` package (Hothorn et al., 2006) in R (R Development
Core Team, 2015) with 1,000 trees (i.e., `ntree = 1,000`) and four randomly selected predictors considered
at each split (i.e., `mtry = 4`). Next, the party's `varimp()` function (Strobl et al., 2008) was used to compute
the conditional permutation importance measures for the predictors, which reflect the impact of each
predictor variable on the dependent variable. If the significance (i.e. the p-value) of the association between
a given predictor and the variable of interest was lower than 0.05, the relevant covariate was included in
the predictor's conditioning scheme. Similarly, we set the `mincriterion` of $p = .05$ to include splits in
the calculation of importance.

A summary of all conducted analyses can be found in Section 7.4 of the Supplementary Materials.

3 Results

3.1 Descriptive statistics

We present the distribution of scores as well as the sample's means and standard deviations of the collected reading and reading-related test scores in Figure 2 below. Apart from for RN-LTR on which adults scored higher than children (t(103.44) = -3.84, p = .0002), children's standard scores were on average higher than those of adults. Significant differences between the groups were noted for T-SWE (t(124.97) = -2.55, p = .01) and T-PDE (t(122.7) = -4.21, p < .0001), but not for the WRMT-WA (t(116.45) = -1.73, p = .09).

Figure 2. Distributions of standard scores on reading and reading-related tasks for adults (in green) and children (in dark gray): T-SWE = Sight Word Efficiency, T-PDE = Phonemic Decoding Efficiency, RN-LTR = Rapid Automized Naming Letters, WRMT-WID = Word Identification, WRMT-WA = Word Attack. Each panel lists means and standard deviations (in parentheses) of the scores for adults (top rows, in green) and children (bottom rows, in black) separately.

The number of TTG multiplications was highly variable in the sample. In the left hemisphere, we noted only 2 cases of a single gyrus (2 adults); 63 participants had 2 gyri (28 adults, 35 children), 55 had 3 gyri (28 adults, 27 children), and 8 had 4 (1 adult, 7 children). In the right hemisphere, 23 cases of a single gyrus were observed (10 adults, 13 children), 79 participants had 2 gyri (37 adults, 42 children), 24 had 3 gyri (11 adults, 13 children), and 2 had 4 (1 adult, 1 child). Further descriptive data for adults' and children's neuroanatomical measures describing the TTG (HG and all TTG(s) separately) can be found in Table 1. Note that all volume, surface and shape measures were significantly left-lateralized (according to a one-sample t-test against 0 on the lateralization indices), with the exception of the shape of the HG in adult participants. None of the average thickness values showed significant lateralization.

	adults				children			
	HG		all TTG(s)		HG		all TTG(s)	
	left	right	left	right	left	right	left	right
Cortical volume:								
mean	1227.52	990.06	2016.92	1509.10	1413.97	1109.65	2522.26	1738.75
(SD)	(319.65)	(347.39)	(495.65)	(351.55)	(473.54)	(348.69)	(717.74)	(403.37)
LI [mean (SD)]	0.11* (0.18)		0.14* (0.13)		0.11* (0.2)		0.17* (0.14)	
Surface area:								
mean	389.25	302.35	623.24	453.54	410.22	319.26	703.67	488.74
(SD)	(108.74)	(95.89)	(158.41)	(99.87)	(133.12)	(89.70)	(175.07)	(100.99)
LI [mean (SD)]	0.12* (0.18)		0.15* (0.13)		0.12* (0.19)		0.17* (0.13)	
Average thickness:								
mean	2.48	2.52	2.58	2.61	2.69	2.70	2.80	2.78
(SD)	(0.23)	(0.21)	(0.17)	(0.18)	(0.23)	(0.21)	(0.20)	(0.20)
LI [mean (SD)]	-0.009 (0.04)		-0.006 (0.03)		-0.001 (0.04)		0.004 (0.03)	
Shape:								
mean	0.06	0.11	2.62	2.19	0.09	0.07	2.77	2.15
(SD)	(0.07)	(0.13)	(0.61)	(0.59)	(0.10)	(0.11)	(0.70)	(0.66)
LI [mean (SD)]	-0.02 (0.68)		0.09* (0.18)		0.24* (0.70)		0.13* (0.21)	

Table 1. Descriptive data for adults' and children's measures of HG (HG) and all TTG(s) (volume, area and average thickness) in native space, and their lateralization indices (LI), where positive values indicate leftward lateralization. (*) denotes significantly lateralized measures (according to a one sample t-test against 0). Note that children's values are consistently higher than adults', which is in line with normative trajectories of gray matter volume, surface area and cortical thickness reported by e.g., Bethlehem and colleagues (2022).

3.2 Intergenerational transmission of reading skills

We first tested whether children's performance on the five reading and reading-related tests (T-SWE, T-PDE, RN-LTR, WRMT-WID and WRMT-WA) was related to parents' performance on the same tests above and beyond children's demographic variables (age, sex and SES). To establish a relationship between children's and parents' reading skills, a series of linear mixed models were fitted to the children's test scores. To account for sibling relationships (i.e., non-independent observations within the data) family index was included as a random intercept in all models. For each model, a likelihood ratio test was performed to compare models that included parents' measures with reduced models that did not include them. FDR correction for multiple comparisons was applied to the resulting *p*-values (five from model comparisons including mothers' data and five with fathers' data). Figure 3A presents the results of these analyses. Children's scores were significantly related to mothers' scores on the same tests for T-PDE ($\beta = 0.63$, $t = 4.76$, $p < .001$), WRMT-WID ($\beta = 0.59$, $t = 2.88$, $p = .007$) and WRMT-WA ($\beta = 0.36$, $t = 2.34$, $p = .02$), and to fathers' scores on the same tests for T-PDE ($\beta = 0.53$, $t = 2.65$, $p = .01$), and WRMT-WID ($\beta = 0.74$, $t = 3.25$, $p = .004$). To assess if the above intergenerational similarity effects of reading scores were due to actual familial relationships and not to spurious correlations across our population, we assessed the specificity of these relationships by computing the same models on 5000 permutations of the family index. The analysis confirmed all above significant relations established in the linear mixed models between child's scores and same test scores for: (1) mother's T-PDE ($p = 0.0004$), WRMT-WID ($p = 0.01$), WRMT-WA ($p = 0.04$), and (2) father's T-PDE ($p = 0.003$), and father's WRMT-WID ($p = 0.001$). In sum,

both mother-child and father-child similarities in reading skills were observed, but were more reliable for

mother-child pairs. Interestingly, they were observed for reading subtests which target phonological

decoding skills rather than lexico-semantic, or sight-word reading.

Figure 3. (A) Relationship between parents' and children's reading scores on five tests: T-SWE, T-PDE, RN-LTR, WRMT-WID and WRMT-WA.
The intensity of the color represents increase in R^2 values between a model with demographic variables only and a model additionally including
mothers' or fathers' corresponding test scores; p-values were obtained from likelihood ratio tests, used to compare the models. (B) Relationship
between TTG(s) anatomy and performance on the reading tests across the whole sample, obtained from comparing models with demographic
variables to models that additionally included the neuroimaging data; R^2 values and p-values were derived and represented similarly as in panel A.
(C) Significant relationships between reading tests and neuroanatomical variables. In all plots, darker dots and regression lines refer to adults, and
lighter ones refer to children. (D) Familiar similarity of the anatomical measures describing the TTG(s) established by comparing models with
demographic variables to models additionally including the family index; R^2 values and p-values derived and represented as in panel A.

279 3.3 TTG(s) and reading measures

To explore whether and how the structure of the TTG(s) was related to reading measures in the whole

sample, we extracted neuroanatomical measures (volume, surface area, average thickness and shape) of

the HG and for all identified TTG(s), from both hemispheres, as well as their lateralization indices. To

establish the relationship between the neuroanatomical measures and reading scores, for each reading

score, we fitted a linear mixed model predicting this score in a model including a neuroanatomical measure

of interest, and a reduced model without it. We compared the two nested models using a likelihood ratio

test. In all models, covariates of no-interest (age [linear and quadratic term], sex, SES, handedness and

total intracranial volume) were included. FDR correction for multiple comparisons was applied to the

resulting *p*-values (120 *p*-values in total: from 24 neuroanatomical measures x 5 reading measures).

Significant associations between reading scores and TTG anatomy were observed only in the left

hemisphere, for anatomical features in all TTG(s), after correction for multiple comparison (Figure 3B, see

also Figure S1 for uncorrected results). Specifically, the models predicting the T-PDE scores by the left all

TTG(s) volume ($\Delta R^2_{\text{Marg.}}$ = 8.15, p_{FDR} = .02) and surface area ($\Delta R^2_{\text{Marg.}}$ = 9.26, p_{FDR} = .02), showed a

significantly better fit than the models without the TTG(s) indices (after FDR correction). Larger left TTG(s)

were associated with better phonemic decoding (volume: β = 0.34, *t* = 2.38, *p* = .001, area: β = 0.34,

t = 3.37, *p* = .001). Furthermore, in the model predicting T-PDE from left all TTG(s) volume, we observed

a significant interaction between left all TTG(s) volume and age (β = -0.24, *t* = -2.26, *p* = .026); the left all

TTG(s) area model showed a significant two-way interaction between left all TTG(s) area and age (β = -

0.26, *t* = -2.71, *p* = .008), and a significant three-way interaction between left all TTG(s) area, age and sex

(β = 0.29, *t* = 2.13, *p* = .036). The significant two-way interactions with age are shown in Figure 3C, where

steeper slopes of the relationship between T-PDE and left all TTG(s) measures can be observed for children

than for adults. The three-way interaction between left all TTG(s) area, age and sex is driven by the fact

that the slope of the relationship between T-PDE and left all TTG(s) area differs between girls, boys, mothers

and fathers and is the steepest for girls.

Further, Figure S1 presents the results of the analysis of the relationship between the structure of the TTG(s)

and reading measures without FDR correction for multiple comparisons. Here, both the volume of the HG

as well as the shape of all TTG(s) show significant relationships with reading measures.

We then ran a follow-up, complementary analysis on the planum temporale (PT). Indeed, additional TTGs

by definition belong to the planum, and the anatomy of the PT has been repeatedly linked to reading skills

(Altarelli et al., 2014; Blockmans et al., 2023; Serrallach et al., 2016). Therefore, using linear mixed models

with age [linear and quadratic term], sex, SES, handedness and total intracranial volume as covariates of

no-interest (with family index as a random intercept), we associated the reading measures which were

significantly related to 'all TTG(s)' volume and area in the previous analysis to the anatomical measures

describing the whole PT. We used the PT label as delineated by the Destrieux segmentation (Destrieux et

al., 2010), implemented in FreeSurfer. Both the volume (β = 0.16, *t* = 2.07, *p* = .04) of the left PT and its

surface area (β = 0.18, *t* = 2.37, *p* = .02) were significantly related to T-PDE. Directly comparing the models

predicting the T-PDE scores from 'all TTG(s)' volume and area *versus* PT volume and area revealed that

'all TTG(s)' measures were better at explaining reading skills, with $\Delta R^2_{\text{Marg.}}$ of 4.39%

(ΔAIC = -7.78) and 3.55% (ΔAIC = -6.61) for volume and area respectively. Thus, reading skills may relate

more specifically to the anatomy of the gyri encompassed within the HG/PT region of the Sylvian fissure,

than to the PT per se. Finally, a two-way interaction between left PT area and sex was observed (β = -

0.32, t = -2.36, p = .02), with a steeper slope of the relationship between T-PDE and left PT area for

females than for males.

323   3.4   Familial similarity in the structure of the TTG(s)

Next, we explored familial similarity and intergenerational transmission of TTG anatomy. A series of linear

models were fitted to the participants' volume, surface area, average thickness and shape of the left and

right HG, of all identified left and right TTG(s), and to their lateralization indices. In each of these models,

we: (1) modelled the extracted anatomical measures as a function of covariates of no-interest only

(participants' age [linear and quadratic term], sex, handedness, and total intracranial volume), and (2)

additionally modelled an index of whether individuals were related or not (i.e., 'family index'). To establish

the familial similarity effect for the four measures, for each we used a likelihood ratio test to compare the

model including the family index to the reduced model without it (i.e., with co-variates of no-interest only);

FDR correction for multiple comparisons was applied to the resulting p-values (24 p-values in total). The

results of these analyses are summarized in Figure 3D. All models including the family index explained at

least 20% more variance than models without (min. ΔR^2 = 20.44%, max. ΔR^2 = 43.96%). Including family

information significantly improved model fit for the volume (ΔR^2 = 40.66%, p_{FDR} = .024), area (ΔR^2 =

39.02%, p_{FDR} = .044), and thickness (ΔR^2 = 35.88%, p_{FDR} = .024) of the right HG, thickness of all TTG(s) in

the left (ΔR^2 = 27.66%, p_{FDR} = .044), and right (ΔR^2 = 31.40%, p_{FDR} = .044) hemispheres, shape of the HG

in the left (ΔR^2 = 41.43%, p_{FDR} = .024) and right (ΔR^2 = 39.11%, p_{FDR} = .024) hemispheres, as well as

lateralization of the volume (ΔR^2 = 40.21%, p_{FDR} = .044) and surface area of all identified TTG(s) (ΔR^2 =

43.96%, p_{FDR} = .024), and lateralization of the area of the HG (ΔR^2 = 39.71%, p_{FDR} = .050). Familial similarity

in TTG anatomy was notably stronger for surface area than thickness, in line with previous reports of higher

genetic contributions to shaping surface area (Grasby et al., 2020). Interestingly, across the regions

considered, familial similarity was strongest for the right HG, with significant familial contributions in all

anatomical features. For all anatomical features of the TTG(s) showing significant familial similarities, we

next investigated the intergenerational origin of such similarities by testing parent-of-origin effects.

346   3.4.1   Intergenerational transmission of the volume, area, and thickness of the 1st right TTG (HG)

To gain further insight into the nature of the familial similarity of the structure of the 1st right TTG, we next

examined the relationship between parent and child measures of right HG volume, surface area, and

average thickness. For this, we employed linear mixed models, treating the children's volume, surface and

thickness of the right HG as dependent variables, and treating the mothers' or fathers' corresponding

measures the independent ones, controlling for children's age [linear and quadratic term], sex, SES, and

handedness, and including family index as a random intercept to account for sibling relationships. In total,

6 models were fit (for three different anatomical measures, for the mothers' and fathers' data separately).

Prior to the analysis, the anatomical measures were normalized for the corresponding whole hemisphere

measures (hemispheric volume, surface area and mean hemispheric thickness) so as to be more

comparable between parents and children. We found the volume, area, and thickness of children's 1st right

TTG to be significantly related to mothers' 1st right TTG volume, area, and thickness respectively (volume:

$\beta = 0.38$, $t = 2.84$, $p = .009$; area: $\beta = 0.30$, $t = 2.33$, $p = .03$; thickness: $\beta = 0.38$, $t = 2.87$, $p = .008$), as

well as children's thickness to fathers' thickness values ($\beta = 0.28$, $t = 2.05$, $p = .05$), but we did not find

relationships between children's and father's respective volume and surface area values (volume: $\beta = 0.15$,

$t = 1.10$, $p = .28$; area: $\beta = 0.16$, $t = 1.14$, $p = .26$), see Table 2 and Figure 4.

		Mother			Father		
		Volume	Area	Thickness	Volume	Area	Thickness
(Intercept)	β	0.12	0.12	0.16	0.22	0.24	0.19
	SE	(0.18)	(0.18)	(0.19)	(0.21)	(0.21)	(0.20)
Parental Volume/Area/Thickness	β	**0.38****	**0.30***	**0.38****	0.16	0.16	**0.28***
	SE	**(0.13)**	**(0.13)**	**(0.13)**	(0.14)	(0.14)	**(0.13)**
Handedness	β	-0.10	-0.12	0.01	-0.12	-0.10	-0.25+
	SE	(0.10)	(0.10)	(0.11)	(0.12)	(0.13)	(0.13)
Age	β	-0.20+	-0.19	-0.20	-0.16	-0.14	-0.23
	SE	(0.11)	(0.12)	(0.13)	(0.12)	(0.13)	(0.14)
Age^2	β	-0.05	-0.06	-0.03	-0.08	-0.06	-0.03
	SE	(0.08)	(0.08)	(0.09)	(0.07)	(0.08)	(0.09)
Sex	β	-0.25	-0.26	-0.25	-0.38	-0.47+	-0.14
	SE	(0.22)	(0.22)	(0.24)	(0.24)	(0.25)	(0.26)
SES	β	-0.03	-0.02	-0.15	-0.17	-0.18	-0.09
	SE	(0.14)	(0.14)	(0.14)	(0.18)	(0.17)	(0.14)
SD (Intercept Family index)		0.53	0.47	0.45	0.63	0.53	0.35
SD (Observations)		0.70	0.75	0.83	0.71	0.80	0.86
Num.Obs.		65	65	65	60	60	60
R^2 Marg.		0.237	0.201	0.196	0.133	0.131	0.188
R^2 Cond.		0.515	0.425	0.375	0.509	0.399	0.301
AIC		190.6	192.2	201.6	183.0	187.5	187.0
BIC		210.2	211.8	221.2	201.8	206.3	205.8
ICC		0.4	0.3	0.2	0.4	0.3	0.1
RMSE		0.58	0.63	0.72	0.57	0.66	0.76

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

362
363 Table 2. Linear mixed model parameters for the volume, surface area and average thickness values of children's right HG as a function of the mothers' and fathers' respective HG volume, surface area and average thickness values.

Figure 4. (A) Relationships between right HG volume, surface area and average thickness values of parents and children plotted separately for mothers (top row) and fathers (bottom row); in all plots, darker dots and regression lines refer to girls, lighter dots refer to boys. (B) Distribution of the beta estimates after family permutation. Dotted lines indicate the best estimate of the right HG values of mothers and fathers computed with the correct family labels.

To assess if the above intergenerational similarity effects in the right HG were due to actual familial relationships and not to spurious correlations across our population, we assessed the specificity of these relationships by computing the same models on 5,000 permutations of the family index. The analysis confirmed all significant relations established in the linear mixed models (p = 0.0006, p = 0.013 and p = 0.004, for mother's volume, area and thickness, and p = 0.04 for father's thickness respectively); see Figure 4B.

To further explore the maternal transmission in the right HG volume and surface area, we checked the potential sex-specificity of this effect, e.g., whether the mother-child similarities were higher for daughters than for sons. Here, we fit further linear mixed models to children's HG volume and area (with the mothers' corresponding measures as independent variables, controlling for children's age [linear and quadratic term], sex, SES, and handedness, and with family index as a random intercept), additionally modelling an interaction between children's sex and mothers' HG volume or area. We compared these models with the models without the interaction terms using likelihood ratio tests. These showed that only the interaction model for volume offered a significantly better fit to the data (X^2 = 5.14, p = 0.023), and not the one for area (X^2 = 2.16, p = 0.14). While it seems that intergenerational similarity of right HG is stronger for mother-daughter pairs, at least for volume, the female specificity of this effect does not seem unequivocal since a

visualization of the result for girls and boys separately (Figure 4A) shows similar, positive slopes of the effect

for both boys and girls.

3.4.2   Intergenerational transmission of left and right HG shape

To further explain the familial similarity effect of TTG(s) shape reported in section 3.4, we fit linear mixed

models to the right and left HG lateral concavity values as determined by MCAI (Dalboni da Rocha et al.,

2023), with mothers' and fathers' HG shape indices as independent variables, controlling for children's age

[linear and quadratic term], sex, SES and handedness, with family index as a random intercept. We found

the shape of children's HG to be significantly related to mothers' HG shape in the left hemisphere (β =

0.31, $t = 2.12$, $p = .04$), but not for the right hemisphere ($\beta = 0.24$, $t = 1.60$, $p = .12$). The relationships

between children's and father's HG concavity values were not significant (left: $\beta = 0.24$, $t = 1.68$ $p = .11$;

right: $\beta = 0.17$, $t = 1.20$, $p = .24$), see Table 3 and Figure 5.

		Mother		Father	
		Left	Right	Left	Right
(Intercept)	β	0.01	0.11	-0.02	0.18
	SE	(0.21)	(0.20)	(0.20)	(0.20)
Parent concavity	β	**0.31***	0.24	0.24+	0.18
	SE	**(0.15)**	(0.15)	(0.14)	(0.15)
Handedness	β	-0.14	0.10	-0.03	-0.02
	SE	(0.12)	(0.10)	(0.13)	(0.12)
Age	β	-0.06	-0.01	-0.06	0.06
	SE	(0.14)	(0.11)	(0.13)	(0.13)
Age^2	β	0.13	-0.19*	0.10	-0.09
	SE	(0.09)	(0.07)	(0.08)	(0.08)
Sex	β	-0.20	-0.09	-0.31	-0.35
	SE	(0.26)	(0.22)	(0.24)	(0.24)
SES	β	0.04	-0.03	-0.01	-0.22
	SE	(0.15)	(0.16)	(0.15)	(0.17)
SD (Intercept Family index)		0.42	0.70	0.48	0.52
SD (Observations)		0.90	0.66	0.77	0.75
Num.Obs.		65	65	60	60
R^2 Marg.		0.143	0.133	0.108	0.098
R^2 Cond.		0.294	0.597	0.356	0.393
AIC		208.0	194.3	181.9	181.8
BIC		227.6	213.9	200.7	200.7
ICC		0.2	0.5	0.3	0.3
RMSE		0.79	0.51	0.65	0.62

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

396 Table 3. Multiple regression model parameters for the concavity values of children's HG as a function of mothers' and fathers' HG concavity values.

Figure 5. Familial similarity of Heschl's gyrus shape. (A) Relationship between HG lateral concavity values of parents and children (plotted separately for mothers and fathers) in the left and right hemisphere. In all plots, darker dots and regression lines refer to girls, lighter dots refer to boys. (B) Distribution of the beta estimates after family permutation. Dotted line indicates the best estimate of the HG concavity of mothers computed with the correct family labels. (C) An example of left HG shape similarity between family members.

Similarly to the analysis of HG volume, area and average thickness values, we computed 5,000 permutations of the family index to assess the specificity of the familial relationship of left HG shape. The analysis confirmed that the concavity of children's HG was significantly related to their mothers' HG concavity in the left hemisphere (p = 0.014), see Figure 5B.

Last, we explored the female-specificity of HG similarity by fitting a further linear mixed model to children's HG shape values (with the mothers' corresponding measure as independent variable, controlling for children's age [linear and quadratic term], sex, SES, and handedness, and with family index as a random intercept), with additionally modelling an interaction between children's sex and mothers' HG shape values. We compared this model with the model without the interaction term using a likelihood ratio test, which showed that the interaction model did not offer a significantly better fit to the data (X^2 = 0.37, p = 0.54). Therefore, the left HG shape intergenerational transmission did not appear sex-specific.

3.4.3 Intergenerational transmission of TTG lateralization

We further explored the familial similarity of the lateralization of all TTG(s) (see Figure 6A) by fitting further linear models to the children's lateralization indices of TTG(s) volume and surface area, with mothers' or fathers' corresponding lateralization values as independent variables, controlling for children's age [linear and quadratic term], sex, SES and handedness, see Table 4. We found the degree of lateralization of the surface area of children's all TTG(s) to be significantly and positively related to the lateralization of the

fathers' all TTG(s) surface area (β = 0.42, t = 3.00, p = .005), but not to that of the mothers (β = 0.04,

t = 0.26, p = .80), see Figure 6B. We found no significant effects in the follow-up analyses on lateralization

of volume (mother: β = 0.03, t = 0.20, p = .84; father: β = 0.26, t = 1.74, p = .09). Permutation analyses

confirmed the specificity of this effect to parent-child pairs: the lateralization of the surface area of all TTG(s)

of the fathers was significantly related to that of their children (p = 0.0006), see Figure 6C. Nested linear

mixed models comparison with a likelihood ratio test revealed that a model including an interaction term

between fathers' lateralization index of all TTG(s) area and children's sex (in addition to demographic

variables (age [linear and quadratic term], sex, SES and handedness), random intercept for family index

and fathers' lateralization index of all TTG(s) area) did not have a better fit to children's all TTG(s) area

lateralization (X^2 = 0.25, p = 0.61). The male-specificity of familial similarity of the lateralization of all TTG(s)

area could therefore not be confirmed.

		Mother		Father	
		LI Volume	LI Area	LI Volume	LI Area
(Intercept)	β	-0.06	-0.17	-0.17	-0.27
	SE	(0.20)	(0.20)	(0.22)	(0.21)
Parent LI	β	0.03	0.04	0.26+	**0.42****
	SE	(0.14)	(0.14)	(0.15)	**(0.14)**
Handedness	β	0.07	0.04	-0.02	-0.04
	SE	(0.11)	(0.11)	(0.13)	(0.13)
Age	β	-0.28*	-0.31*	-0.21	-0.21
	SE	(0.13)	(0.12)	(0.14)	(0.13)
Age^2	β	0.03	0.08	0.00	0.04
	SE	(0.08)	(0.08)	(0.08)	(0.08)
Sex	β	0.16	0.24	0.30	0.39
	SE	(0.24)	(0.24)	(0.26)	(0.25)
SES	β	-0.13	-0.12	0.01	0.05
	SE	(0.16)	(0.16)	(0.17)	(0.16)
SD (Intercept Family index)		0.62	0.61	0.62	0.58
SD (Observations)		0.77	0.78	0.76	0.81
Num.Obs.		65	65	65	60
R^2 Marg.		0.074	0.086	0.106	0.113
R^2 Cond.		0.439	0.431	0.459	0.417
AIC		195.5	203.7	202.0	190.9
BIC		210.8	223.3	221.5	209.7
ICC		0.4	0.4	0.4	0.3
RMSE		0.63	0.64	0.62	0.66

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

430
431 Table 4. Multiple regression model parameters for the lateralization indices of children's surface area of all TTG(s) as a function of mothers' and fathers' surface area of all TTG(s) lateralization indices.

TTG Lateralization

Figure 6. Familial similarity of TTG(s) lateralization. (A) Lateralization indices of TTG(s) (*x*-axis) for 37 different families (*y*-axis); each family is plotted in a different color. (B) Relationship between lateralization index of TTG(s) surface area of parents and children, plotted separately for mothers and fathers. In both plots, darker dots and regression lines refer to girls, lighter dots refer to boys. (C) Distribution of the beta estimates after family permutation. Dotted line indicates the best estimate of lateralization of TTG(s) area of fathers computed with the correct family labels.

3.4.4   Lack of intergenerational transmission of other features showing familial similarity

We also performed follow-up analyses on the remaining results that showed significant familial similarity effects in the first exploratory analysis reported in Section 3.4. These included the thickness of the left and right all TTG(s), and the lateralization of the area of the HG. Here, again, linear mixed models were used to relate these measures in children to those of their mothers and fathers. With respect to the measure of average thickness of all left TTG(s), neither the mothers' (β = 0.25, *t* = 1.78, *p* = .09) nor the fathers' (β = 0.24, *t* = 1.72, *p* = .10) measures were significantly related to those of the children. This was also the case for all right TTG(s) (mothers: β = 0.19, *t* = 1.77, *p* = .08; fathers: β = 0.28, *t* = 1.98, *p* = .053). Similarly, neither the mothers' (β = 0.22, *t* = 1.53, *p* = .14) nor the fathers' (β = 0.21, *t* = 1.47, *p* = .15) lateralization of the area of the HG were significantly related to the children's lateralization of the area of the HG. Therefore, the effect of "family" for the above neuroanatomical measures reported in Section 3.4 might have been driven by anatomical similarities between the siblings, rather than between parents and children.

3.5 Intergenerational effects on reading ability: is children's reading related to parents' neuroanatomy?

As a last step in our analyses, we aimed to determine whether mothers' or fathers' neuroanatomical measures could predict children's reading. Given the large number of neuroanatomical variables describing the TTG(s), and given that several different brain measures were either related to reading performance (Section 3.3) or showed neuroanatomical concordance between parents and children (Section 3.4), we opted for an exploratory approach using a *random forests* classifier (Breiman, 2001). Here, we used children's T-PDE scores as a dependent variable, given their association with TTG(s) anatomy established in Section 3.3. We grew two random forests modelling children's T-PDE in terms of all available (1) mothers' and (2) fathers' anatomical measures describing the HG and all TTG(s). Prior to the analysis, the anatomical measures were normalized for the corresponding whole hemisphere measures (hemispheric volume, surface area and mean hemispheric thickness). Conditional permutation importance scores were computed for all models, and are presented in Figure 7A. The dotted vertical lines demarcate the permutation scores that can differ from zero due to randomness alone; variables having importance scores to the left of this line can be considered not to be relevant predictors.

Figure 7. Results of the random forests analysis. (A) Conditional permutation importance for mothers' and fathers' neuroanatomical TTG measures in predicting children's reading skills. (B) Children's reading skills as a function of fathers' anatomical variables based on the follow-up analyses (additional plots for father's volume variables can be found in Supplementary Materials, Figure S2).

The model with mothers' variables did not point to any features of their TTG(s) to be relevant predictors of

children's reading ability, but several variables in the model with fathers' data did. The variables with the

highest conditional permutation importance in case of fathers were volume and area of the left HG, as well

as HG lateralization indices of volume and area; all had conditional permutation importance significantly

different from zero, see Figure 7A. To validate these effects and to determine their direction, we fitted

separate linear mixed models to children's T-PDE scores, each with one of the relevant predictors as

independent variables (controlling for children's sex, age and SES), and with family index as random

intercept. Fathers' left volume and area were both negatively related to children's reading (volume:

β = -0.40, t = -2.91, p = .008; area: β = -0.46, t = -3.45, p = .002), indicating that the smaller the father's

left HG, the better the child's reading scores. Fathers' lateralization of HG (both for volume and area) was

not significantly related to the children's reading (volume: β = -0.29, t = -1.27, p = .21; area: β = -0.20,

t = -1.32, p = .19), see Figure7B and Table 5.

To explore sex specificity of these effects, we performed a series of nested model comparisons with

likelihood ratio tests for models including (1) left HG Volume, (2) left HG Area, (3) LI HG Volume, or (4) LI

HG Area, additionally including an interaction term between fathers' measures and child's sex. The results

did not point to sex specificity of any of the effects (left HG Volume: X^2 = 2.25, p = 0.13, left HG Area:

X^2 = 0.88, p = 0.35, LI HG Volume: X^2 = 0.17, p = 0.68, LI HG Area: X^2 = 0.60, p = 0.44) Furthermore, we

again tested for familial specificity of the effect of father's left HG volume and area on children's reading

with permutation analyses. These revealed that the relationship between fathers' anatomical measures

(volume and surface area of the HG) was significantly more related to their own children's reading scores

than to non-related children's reading scores (p = 0.002 and p = 0.0006 for fathers' volume and surface

area, respectively).

		Father left HG Volume	Father left HG Area	Father LI HG Volume	Father LI HG Area
(Intercept)	β	-0.06	-0.01	-0.07	-0.07
	SE	(0.19)	(0.19)	(0.21)	(0.21)
Father's measure	β	**-0.40****	**-0.46****	-0.19	-0.20
	SE	**(0.14)**	**(0.13)**	(0.15)	(0.15)
Child's age	β	-0.20+	-0.20+	-0.19	-0.19
	SE	(0.12)	(0.12)	(0.12)	(0.12)
Child's sex	β	0.25	0.15	0.30	0.31
	SE	(0.26)	(0.26)	(0.27)	(0.27)
SES	β	0.15	0.16	0.12	0.12
	SE	(0.14)	(0.14)	(0.17)	(0.17)
SD (Intercept Family index)		0.37	0.31	0.56	0.56
SD (Observations)		0.84	0.84	0.82	0.82
Num.Obs.		58	58	58	58
R^2 Marg.		0.259	0.303	0.130	0.134
R^2 Cond.		0.382	0.385	0.409	0.411
AIC		172.8	170.3	178.4	178.2
BIC		187.2	184.8	192.8	192.6
ICC		0.2	0.1	0.3	0.3
RMSE		0.75	0.76	0.69	0.69

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Table 5. Multiple regression model parameters for children's reading ability as a function of the mothers' and fathers' anatomical variables having shown the highest conditional permutation importance in the random forests analysis.

Given the surprising direction of the intergenerational effects on children's reading ability, we set out to further understand why fathers' small left HG would be associated with better reading skills in children. One possibility is that fathers' small left HG goes hand in hand with more additional TTGs, or with a larger PT (region exclusively or mostly including non-primary auditory areas). Using linear models, we therefore tested (1) whether a smaller left HG in fathers was associated with a greater likelihood of having additional TTGs (reflected by a higher lateral multiplication index for left all TTG(s)), and (2) if there was a negative relationship between the volume and/or area of fathers' left HG and those of their PT. The lateral multiplication index for left all TTG(s) was not significantly associated with either the volume (β = 0.08, $t = 0.40$, $p = .69$), or the surface area (β = -0.05, $t = 0.19$, $p = .78$) of the left HG in fathers. Surface area of fathers' left HG was, however, significantly positively related to the surface area of their PT (β = 0.18, $t = 2.036$, $p = .05$), but volume of HG was not related to PT volume (β = 0.20, $t = 1.08$, $p = .29$).

We therefore further evaluated the relationships between children's reading and the characteristics of their fathers' auditory cortex regions using linear mixed models (with children's age, sex and SES as covariates of no-interest and family index as a random intercept). Here, in separate models, we included both Heschl's gyrus and planum temporale as independent variables, as well as how much space both structures occupy relatively to each other (i.e., HG:PT ratio) (see Serrallach et al., 2016 for findings of a decreased left HG:PT ratio in dyslexia). We concentrate on the surface area here, but see Supplementary Materials for convergent results for the volume. First, while fathers' PT area alone did not significantly predict children's T-PDE scores (β = 0.11, $t = 0.64$, $p = .53$), fathers' HG:PT ratio did indeed significantly and strongly predict children's reading (β = -0.48, $t = -3.57$, $p = .002$), negatively. The HG:PT ratio effect was not sex-specific:

a nested model comparison with a likelihood ratio test did not show a better fit for a model additionally

including an interaction term between fathers' HG:PT ratio and child's sex ($X^2 = 0.44$, $p = 0.51$).

Conversely, however, we noted a significantly better fit for a model with an interaction term between

fathers' PT area and child's sex over a model with fathers' PT area only ($X^2 = 5.34$, $p = 0.02$): while fathers'

PT area did not significantly predict children's T-PDE overall (as reported above), and was not significantly

related to girls' reading ($\beta = -0.12$, $t = -0.65$, $p = .53$), it was so for boys only ($\beta = 0.66$, $t = 2.40$, $p = .03$).

Permutation analyses confirmed familial specificity of the effect of fathers' HG:PT ratio on children's reading

($p < .0001$), and fathers' PT area on boys' reading ($p = .005$). Directly comparing the models predicting

children's T-PDE with either father's left HG surface area, or fathers' HG:PT ratio as independent variables

showed that the model with HG:PT ratio offered a better fit to the data with $\Delta R^2_{Marg.}$ of 0.9% (ΔAIC = -0.81).

For boys, the model with the lowest AIC was still the HG:PT ratio model (94.70 *versus* 98.99 for father's

left HG surface area, and 96.77 for fathers' left PT), despite the significant effect of fathers' left PT surface

area on boys' reading.

In addition, to confirm father-specific transmission, we ran the same three models but with mothers' HG,

PT and HG:PT ratio surface area instead of fathers'. We found no significant relationships between

mothers' neuroanatomical measures and children's reading (HG: $\beta = 0.11$, $t = 0.78$, $p = .44$; PT: $\beta = 0.13$,

$t = 0.92$, $p = .37$; HG:PT ratio: $\beta = 0.02$, $t = 0.12$, $p = .91$). Associating children's own neuroanatomical

measures (surface area/volume of left HG, left PT and left HG:PT ratio) with their reading (controlling for

age, sex, SES, handedness and total intracranial volume, and with family index as a random intercept) in

linear mixed models pointed only to the left HG as a significant predictor of reading (area: $\beta = 0.26$, $t = 2.28$,

$p = .03$; volume: $\beta = 0.31$, $t = 2.60$, $p = .01$). Neither left PT (area: $\beta = 0.16$, $t = 1.43$, $p = .16$; volume: $\beta =$

0.10, $t = 0.88$, $p = .39$), nor HG:PT ratio (area: $\beta = 0.09$, $t = 0.68$, $p = .50$; volume: $\beta = 0.19$, $t = 1.44$,

$p = .16$) were significantly associated with children's reading. Lastly, of note, there was no significant

relationship between the father's left HG, PT, or HG:PT ratio and their own T-PDE scores (left HG: $\beta = -$

0.32, $t = -1.64$, $p = .11$; left PT: $\beta = -0.08$, $t = -0.36$, $p = .72$; left HG:PT: $\beta = -0.30$, $t = -1.33$, $p = .19$),

according to linear models controlling for age, handedness, SES (and total intracranial volume for left HG

and PT models).

In sum, we observed strong intergenerational, father-specific effects on offspring's reading, manifested by

the relative sizes of left primary (HG) and secondary (PT) auditory areas: a small left HG:PT ratio in fathers

is related to worse reading in children. Moreover, fathers' PT area was positively associated with reading

in boys only.

4   Discussion

The purpose of the present study was to explore familial similarity in reading skills, to establish relationships between auditory cortex anatomy and individual differences in reading, to examine familial and intergenerational similarities in auditory cortex structure, and to examine intergenerational effects of parental brain structure on children's reading. Our main findings are summarized in Figure 8, and as follows.

(1) children's reading skills were related to those of parents;
(2) the size of all left TTG(s) was related to reading skill across the whole sample;
(3) mothers' HG was similar to those of children, and all TTG(s) were similarly lateralized across fathers and children;
(4) the relative sizes of primary (HG) and secondary (PT) auditory areas in fathers were related to children's reading.

Figure 8. A schematic representation of the most important findings, as discussed in the text.

First, (1) we found that children's reading skills were significantly related to those of parents, more so for mother-child pairs than for father-child pairs. Interestingly, they were observed for reading subtests which target phonological decoding skills rather than lexico-semantic, or sight-word reading. This finding is in line with twin studies showing higher heritability of phonological aspect of reading rather than lexically-mediated ones (Gay and Olson, 2003), and a recent genome-wide association study in which a higher proportion of non-word reading variability was accounted for in comparison to word-reading (Eising et al., 2022).

Furthermore, given that most of the mothers participating in the study were homemakers (21 out of 33,
see Table S1), this finding might also reflect the amount of time the mothers spent with their children, and
thus suggest an environmental (rather than genetic) transmission of reading skill. Alternatively, it could
reflect genetically-driven parent-of-origin effects, or genomic imprinting (Lawson et al., 2013), which we
discuss in more detail below (see Section 4.3).

Next, using newly developed toolboxes (Dalboni da Rocha et al., 2023, 2020), we performed detailed
segmentations of gyri in the superior temporal plane, i.e., the Heschl's gyrus and further TTG(s), when
present, and found that:

(2) across the whole sample, volume and surface area of all TTG(s) in the left hemisphere correlated
565   with individual differences in speeded phonemic decoding;

566   (3) there were structural brain similarities for parent-child pairs in the 1st TTG (Heschl's gyrus, HG),
567   and in the lateralization of all TTG(s) for father-child pairs;

568   (4) the relative sizes of HG (including primary) and PT (consisting exclusively or mostly of
569   secondary) auditory areas in fathers were associated with offspring reading ability.

570 Each of these points is discussed in more detail below.

4.1 TTG(s) and reading abilities

Across our whole sample, individual differences in performance on speeded phonemic decoding (T-PDE)
(Torgesen et al., 1999) was related to the structure of the auditory cortex when a stricter statistical
correction was applied; three other tests, of speeded sight-word decoding (T-SWE) (Torgesen et al., 1999),
untimed sight-word decoding (WRMT-WID) (Woodcock, 1998), and untimed phonemic decoding (WRMT-
WA) (Woodcock, 1998), showed relationships with the same neuroanatomical measures when uncorrected
statistics were used. Thus, we observed that both word (WRMT-WID, T-SWE) and pseudo-word reading
(T-PDE, WRMT-WA) were related to some degree to TTG(s) anatomy. However, the task that assessed
how well participants could pronounce syllables and pseudo-words and use their phonetic decoding skills
and knowledge of grapheme–phoneme correspondence (T-PDE), was more strongly related to TTG(s)
anatomy (see Figure 3B). This relationship was stronger for children than for adults, possibly pointing to
developmental changes of the nature of reading throughout development: the reliance on auditorily-
mediated processes is stronger in beginning stages of reading (i.e., during early school years).

Notably, rapid automatized naming (RAN) skills, as measured by the RN-LTR task (Wolf and Denckla,
2005), showed no relationships with any neuroanatomical measures of the TTG(s), even at the uncorrected
threshold. This dissociation between TTG(s) anatomical correlates of phonological processing skills *versus*
RAN skills is consistent with the double-deficit hypothesis of dyslexia (Wolf and Bowers, 1999). According

to this hypothesis, reading difficulties may be caused by impairments in either naming speed or

phonological processing, with individuals with a "double-deficit" having more reading problems than those

with a single deficit. Neuroanatomical evidence for such a dissociation has been provided by both structural

(Leonard et al., 2006) and functional (Norton et al., 2014) data (although Norton and colleagues did not

observe the functionality of TTG(s) to be related to the degree of phonological impairment), and our data

provide further evidence for the dissociation.

In terms of the precise characteristics of the TTG(s) that were related to the variability in phonological skills

in our sample, we observed positive relationships between the size (volume and surface area) of all

identified TTG(s) in the left hemisphere and the relevant reading tasks. Furthermore, inspection of the results

without FDR correction for multiple comparisons (Figure S 1) showed that both variables describing the

1st left gyrus (volume and area of the left HG), as well as the overall shape (which includes information on

the number of identified gyri as indexed by the variable 'left all TTG(s) shape') contributed to the 'left all

TTG(s) volume' and 'left all TTG(s) area' results. Thus, our data suggest that larger and more numerous

gyri in the left superior temporal plane are associated with better reading skills. This finding replicates recent

results showing that in pre-readers, the surface area of left HG as well as the duplication patterns of the

left TTG(s) positively predicted later word reading (Blockmans et al., 2023). The finding is also consistent

with higher incidence of duplicated TTGs in relation to individual differences in non-native speech sound

learning (Golestani et al., 2007a), and in expert phoneticians (Golestani et al., 2011), i.e., it extends those

results to reading, at the same time underscoring the importance of phonetic processing in reading.

Notably, in the present results, the relationship between TTG(s) anatomy with reading was stronger than

that of PT anatomy (which includes additional TTGs) with reading, indicating the specific importance of gyri

within the PT for skilled reading. As discussed elsewhere (Kepinska et al., 2023), the properties of gyri

relative to sulci, such as greater neuronal density, cytoarchitectural differences or connectivity properties,

may make them more likely to be associated with better auditory or higher-level linguistic skills than

neighboring sulcal cortex.

Of note, none of the neuroanatomical variables describing the lateralization of TTG(s) anatomy were

significantly associated with the reading measures collected, nor did the indices of the right hemisphere

gyri survive correction for multiple comparisons. While a negative relationship between the shape of the

right TTG(s) and reading ability might have been expected based on the results of Altarelli et al. (2014) and

Serrallach et al. (2016), such an association was not present in our data. This may be due to the limited

number of (very) poor readers in our sample, and as such, does not contradict the previous findings: a

positive association with the overall size and gyrification of the TTG(s) in the left hemisphere and reading in

our sample of predominantly typical readers may be a 'flip side' of the right hemisphere correlates of
impaired reading.

4.2   Familial similarity of the TTG

Regarding TTG(s) similarity between family members, we observed intergenerational effects in HG and in
the lateralization of all identified TTG(s). Intergenerational similarity between parents and children in
measures of volume, surface area, and thickness were restricted to right HG, whereas shape was only
significant in left HG. Both the HG and TTG(s) lateralization findings were confirmed to be significantly more
likely for parent-child pairs than for random adult-child pairs, confirming that the observed effects were not
due to general similarities of the brain regions studied across individuals.

4.2.1   Right HG – volume, area and thickness

The structure of the right HG was similar in terms of volume, area, and thickness between mothers and
children, and only in terms of thickness between fathers and children. HG houses the primary auditory
cortex, but its size and shape have also been shown to be related to higher-level cognitive functions, such
as musicianship (Schneider et al., 2002) and phonetic learning skill and expertise (Golestani et al., 2011,
2007a). Research has suggested that the right HG uniquely underlies processing of pitch direction
(Johnsrude et al., 2000), and that its larger volume is an anatomical marker of absolute pitch (Wengenroth
et al., 2014). The thickness of the right HG has been specifically associated with the severity of auditory
verbal hallucinations in schizophrenia patients (Chen et al., 2015). At the same time, data from our
laboratory suggest that right HG surface area is positively related to individual differences in foreign
language aptitude (Ramoser et al. *under review*). In the context of reading, Ma and colleagues (2015)
showed a relationship between the thickness of the right HG and of surrounding areas and dyslexia, and
dyslexia risk was significantly associated with polygenic risk score of bilateral HG thickness (Gialluisi et al.,
2021). Whether and how individual differences in musicality, pitch perception, and language aptitude may
be mediated by parent-child concordance in HG morphology requires further research. So far, however,
our data do not suggest that variability in children's reading ability can be related to the morphology of their
own right HG, despite the apparent parent-offspring similarities (see also Section 3.5 on intergenerational
effects on reading skills and the discussion of these findings below).

4.2.2   Shape of the left HG

We also observed that the shape of mothers' left HG was similar to that of children's left HG. The shape
of the left (but not the right) HG has previously been related to phonetic learning ability, with good phonetic
learners having a higher probability of having a duplicated TTG (Golestani et al., 2007b), as well with
phonetic expertise (Golestani et al., 2011). Phonetic learning might be an ability intergenerationally

transmitted by mothers through mother-child concordance of the shape of the left HG. To confirm this,

future research should employ phenotypic testing of auditory learning abilities in both parents and offspring.

Incidentally, gyrification patterns of the HG have been related to schizophrenia symptoms and suggested

to be associated with susceptibility to psychopathology (Takahashi et al., 2021). Schizophrenia is a partly

genetically transmitted condition (Sullivan et al., 2003), with the mother's disorder determining the high-

risk status of the offspring (Niemi et al., 2005). Mother-child similarities in HG shape may provide novel

neurobiological support for maternal transmission patterns in schizophrenia, but the status of HG shape

as a potential endophenotype of schizophrenia requires further research with affected individuals.

4.2.3   Lateralization effects

Leftward asymmetry of the temporal speech region has been well established in the literature ever since

the work of Geschwind and Levitsky (1968) (Marie et al., 2015; Penhune et al., 1996), and has been more

recently confirmed by large-scale neuroimaging efforts (Kong et al., 2018). In our data, both adults and

children showed a leftward asymmetry in the volume and area of HG and TTG(s), whereas the thickness

of HG and TTG(s) was symmetrically distributed across the hemispheres (see Table 1). This is partially

consistent with previous asymmetry investigations of auditory cortex: while Meyer and colleagues (2014)

reported leftward-lateralization in terms of volume and surface area, in line with our current findings, they

found a clear rightward asymmetry for thickness of the auditory regions (see also Kong et al., 2018 for

similar results). This discrepancy may be due to different segmentation approaches (atlas-based *versus*

individual anatomy-based), or to exclusively investigating HG (as opposed to all TTGs). We found the

degree of lateralization of the surface area of TTG(s) to be similar within families, and driven by the similarity

between the TTG(s) lateralization indices of fathers and their children. Lack of (i.e. symmetrical) or even

reserved asymmetries of the temporal speech region have previously been reported as male-specific neural

markers of dyslexia (Altarelli et al., 2014). Our results of intergenerational similarity of TTG(s) lateralization

(Section 3.4.3) suggest that these dyslexia-related individual differences in structural asymmetries may be

transmitted from fathers to offspring. Genetic influences on the structural lateralization of the temporal

speech regions have been proposed in the literature (e.g., Eckert et al., 2002), but despite the findings that

individual differences in cerebral lateralization are highly heritable (Francks, 2015), Bishop (2013) proposes

that the degree of lateralization might also be driven by environmental factors. Because intergenerational

similarities reflect both genetic and environmental factors common to families, our results may reflect both.

4.3   Parent-of-origin effects

Most of the intergenerational effects reported here were specifically patrilineal or matrilineal (with the

exception of the parent-offspring similarities in average thickness of the right HG). Their basis could be

environmental in nature (see, for example, Feldman (2023) for a discussion of paternal contributions to
development), or an unintended by-product of our particular sample. They could also potentially be due to
genetic parent-of-origin effects, which refer to phenotypic effects of an allele depending on whether it is
inherited from the mother or father, and which play an important role in the genetic architecture of complex
traits (Lawson et al., 2013). Such parent-of-origin effects are common in humans (Mozaffari et al., 2019),
and indeed Goos et al. (2006), based on animal studies, clinical data and intergenerational analyses of
behavioral data, suggest that they are "influential in brain development, with the maternal genome playing
a disproportionate role in the development of the cortex" (p. 19). Our data suggest that the structure of the
auditory cortex manifests such a parent-of-origin effect to some extent, and that maternal influences may
indeed be more important in the development of auditory regions than paternal influences (for a further
discussion of the molecular basis of the parent-of-origin effect see e.g., Hager et al., 2008; Lawson et al.,
2013).

4.4 Intergenerational effects on reading ability

Given that none of the intergenerational TTG(s) similarity effects reported in Section 3.4 were also found to
be related to reading ability (Section 3.3), we took an exploratory approach to determining whether (and
how) neuroanatomical characteristics of mothers' or fathers' TTG(s) could predict children's reading. The
structure of the left HG was found to be the best predictor of children's reading for fathers' data (no
neuroanatomical markers of mothers were found to be related to children's reading). Surprisingly, fathers
with a statistically smaller left HG had children who were better readers. This finding is counterintuitive
because, to our knowledge, no studies have reported a negative association between left HG size and
reading ability. Our follow-up analyses revealed that, in fact, it was the relative size of fathers' HG (including
primary) to secondary (PT) auditory areas that explained most variance in children's reading (small HG:PT
ratio was related to better reading). Moreover, in boys specifically, fathers' large PT was related to better
reading. These results may be related to the sex differences in the auditory cortex often observed in the
literature. As mentioned above, Altarelli and colleagues (2014) found that an increased number of right TTG
duplications and altered asymmetry of the PT are associated with dyslexia but only in boys; Sutherland et
al. (2012) found that the sex-specificity of the association between gray matter probability (likely reflecting
volume) of left Heschl's gyrus and auditory processing differs across development (i.e., is only apparent
during early adolescence).

The direction of our HG:PT ratio finding does not align with that observed in developmental disorders
reported by Serrallach and colleagues (2016): diminished HG:PT ratios were reported for adults with ADHD
and ADD (2022), and for children with ADHD, ADD and – crucially – with dyslexia. One could expect that
an association between parental brain structure and children's reading should follow the same direction

(i.e., that a small HG:PT ratio should be related to worse reading outcomes), but our data show the
opposite pattern. This might have to do with the specific, indirect nature of intergenerational effects
investigated here. We related children's behavior to parents' brain measures, and these associations might
show different patters due to, for example, neuroplasticity mechanisms operating throughout the lifespan.
The inconsistent direction of our and of previous results may also arise from a non-linear relationship
between reading level and HG:PT ratio; it could for example be that there is an inversed u-shaped function
between HG:PT surface area (of fathers) and reading level. In turn, this could arise from the fact that
intergenerational anatomical underpinnings of dyslexia may be different than those underlying healthy
variability in reading skill (in other words, similar anatomical features may exist, but these regions may
function differently, or display in functional and/or structural connectivity). Our finding of a father-specific
association between parental left auditory cortex anatomy and offspring reading ability may, furthermore,
add to the body of knowledge showing differential patterns of association between auditory cortex anatomy
and cognitive abilities across sexes. However, the exact mechanisms by which paternal HG and PT are
related to offspring reading, and whether there may be third variables that explain this relationship, should
be explored in future research.

732 4.5 Limitations and conclusions

Given the limited prior research on parent-offspring relationships with regards to reading and
neuroanatomy, the precise mechanisms underlying our findings remain uncertain; however, it is reasonable
to hypothesize that a convergence of genetic, prenatal, and postnatal environmental factors contribute to
the observed effects. Given that our study was limited to genetically-related families, future work will benefit
from including *in vitro* fertilization and adoptive families to isolate distinct genetic, prenatal, and postnatal
environmental influences on auditory cortex structure and related (reading) phenotypes. In addition, the
relatively small sample size and exploratory nature of the study underscore the need for further research
with a larger pool of participants. Nevertheless, we provide novel insights into the neural underpinnings of
reading ability in the auditory cortex, children's skills in relation to parental reading, and neuroanatomy.

742 **5 Acknowledgments**

Data collection was funded by the National Institutes of Health (NIH) under grant K23HD054720 to FH.
The authors gratefully acknowledge support by the NCCR Evolving Language, Swiss National Science
Foundation (SNSF) Agreement #51NF40_180888, and by the SNSF grant #100014_18238. OK was
funded by the Marie Jahoda Stipendium from University of Vienna. FB was successively supported by NIH
R01HD094834, the Institute of Convergence ILCB (supported by grants from France 2030 ANR-16-

CONV-0002 and the Excellence Initiative of Aix-Marseille University A*MIDEX), and the French Foundation

for Medical Research (*Fondation Médicale pour la Recherche*, FRM ARF202209015734).

6   References

Ahtam, B., Turesky, T.K., Zollei, L., Standish, J., Grant, P.E., Gaab, N., Im, K., 2021. Intergenerational

Transmission of Cortical Sulcal Patterns from Mothers to their Children. Cerebral Cortex 31, 1888–

1897. https://doi.org/10.1093/cercor/bhaa328

Altarelli, I., Leroy, F., Monzalvo, K., Fluss, J., Billard, C., Dehaene-Lambertz, G., Galaburda, A.M., Ramus,

F., 2014. Planum temporale asymmetry in developmental dyslexia: Revisiting an old question. Hum

Brain Mapp 35, 5717–5735. https://doi.org/10.1002/HBM.22579

Benner, J., Wengenroth, M., Reinhardt, J., Stippich, C., Schneider, P., Blatow, M., 2017. Prevalence and

function of Heschl's gyrus morphotypes in musicians. Brain Struct Funct 222, 3587–3603.

https://doi.org/10.1007/s00429-017-1419-x

Bethlehem, R.A.I., Seidlitz, J., White, S.R., Vogel, J.W., Anderson, K.M., Adamson, C., Adler, S.,

Alexopoulos, G.S., Anagnostou, E., Areces-Gonzalez, A., Astle, D.E., Auyeung, B., Ayub, M., Bae,

J., Ball, G., Baron-Cohen, S., Beare, R., Bedford, S.A., Benegal, V., Beyer, F., Blangero, J., Blesa

Cábez, M., Boardman, J.P., Borzage, M., Bosch-Bayard, J.F., Bourke, N., Calhoun, V.D.,

Chakravarty, M.M., Chen, C., Chertavian, C., Chetelat, G., Chong, Y.S., Cole, J.H., Corvin, A.,

Costantino, M., Courchesne, E., Crivello, F., Cropley, V.L., Crosbie, J., Crossley, N., Delarue, M.,

Delorme, R., Desrivieres, S., Devenyi, G.A., Di Biase, M.A., Dolan, R., Donald, K.A., Donohoe, G.,

Dunlop, K., Edwards, A.D., Elison, J.T., Ellis, C.T., Elman, J.A., Eyler, L., Fair, D.A., Feczko, E.,

Fletcher, P.C., Fonagy, P., Franz, C.E., Galan-Garcia, L., Gholipour, A., Giedd, J., Gilmore, J.H.,

Glahn, D.C., Goodyer, I.M., Grant, P.E., Groenewold, N.A., Gunning, F.M., Gur, R.E., Gur, R.C.,

Hammill, C.F., Hansson, O., Hedden, T., Heinz, A., Henson, R.N., Heuer, K., Hoare, J., Holla, B.,

Holmes, A.J., Holt, R., Huang, H., Im, K., Ipser, J., Jack, C.R., Jackowski, A.P., Jia, T., Johnson,

K.A., Jones, P.B., Jones, D.T., Kahn, R.S., Karlsson, H., Karlsson, L., Kawashima, R., Kelley, E.A.,

Kern, S., Kim, K.W., Kitzbichler, M.G., Kremen, W.S., Lalonde, F., Landeau, B., Lee, S., Lerch, J.,

Lewis, J.D., Li, J., Liao, W., Liston, C., Lombardo, M. V., Lv, J., Lynch, C., Mallard, T.T., Marcelis, M.,

Markello, R.D., Mathias, S.R., Mazoyer, B., McGuire, P., Meaney, M.J., Mechelli, A., Medic, N., Misic,

B., Morgan, S.E., Mothersill, D., Nigg, J., Ong, M.Q.W., Ortinau, C., Ossenkoppele, R., Ouyang, M.,

Palaniyappan, L., Paly, L., Pan, P.M., Pantelis, C., Park, M.M., Paus, T., Pausova, Z., Paz-Linares,

D., Pichet Binette, A., Pierce, K., Qian, X., Qiu, J., Qiu, A., Raznahan, A., Rittman, T., Rodrigue, A.,

Rollins, C.K., Romero-Garcia, R., Ronan, L., Rosenberg, M.D., Rowitch, D.H., Salum, G.A.,

Satterthwaite, T.D., Schaare, H.L., Schachar, R.J., Schultz, A.P., Schumann, G., Schöll, M., Sharp,
781   D., Shinohara, R.T., Skoog, I., Smyser, C.D., Sperling, R.A., Stein, D.J., Stolicyn, A., Suckling, J.,
782   Sullivan, G., Taki, Y., Thyreau, B., Toro, R., Traut, N., Tsvetanov, K.A., Turk-Browne, N.B., Tuulari,
783   J.J., Tzourio, C., Vachon-Presseau, Valdes-Sosa, M.J., Valdes-Sosa, P.A., Valk, S.L., van
784   Amelsvoort, T., Vandekar, S.N., Vasung, L., Victoria, L.W., Villeneuve, S., Villringer, A., Vértes, P.E.,
785   Wagstyl, K., Wang, Y.S., Warfield, S.K., Warrier, V., Westman, E., Westwater, M.L., Whalley, H.C.,
786   Witte, A. V., Yang, N., Yeo, B., Yun, H., Zalesky, A., Zar, H.J., Zettergren, A., Zhou, J.H., Ziauddeen,
787   H., Zugman, A., Zuo, X.N., Rowe, C., Frisoni, G.B., Binette, A.P., Bullmore, E.T., Alexander-Bloch,
788   A.F., 2022. Brain charts for the human lifespan. Nature 604, 525–533.
789   https://doi.org/10.1038/s41586-022-04554-y

790 Bishop, D.V.M., 2013. Cerebral asymmetry and language development: Cause, correlate, or
791   consequence? Science (1979) 340. https://doi.org/10.1126/science.1230531

792 Blockmans, L., Golestani, N., Dalboni, J.L., Wouters, J., Ghesquière, P., Vandermosten, M., 2023. Pre-
793   reading auditory and neurostructural predictors of later reading outcome: the mediating role of family
794   risk. Neurobiology of Language 1–69. https://doi.org/10.1162/NOL_A_00111

795 Breiman, L., 2001. Random Forests. Mach Learn 45, 5–32. https://doi.org/10.1023/A:1010933404324

796 Chen, X., Liang, S., Pu, W., Song, Y., Mwansisya, T.E., Yang, Q., Liu, H., Liu, Z., Shan, B., Xue, Z., 2015.
797   Reduced cortical thickness in right Heschl's gyrus associated with auditory verbal hallucinations
798   severity in first-episode schizophrenia. BMC Psychiatry 15, 152. https://doi.org/10.1186/s12888-
799   015-0546-2

800 Clark, K.A., Helland, T., Specht, K., Narr, K.L., Manis, F.R., Toga, A.W., Hugdahl, K., 2014.
801   Neuroanatomical precursors of dyslexia identified from pre-reading through to age 11. Brain 137,
802   3136–3141. https://doi.org/10.1093/brain/awu229

803 Dalboni da Rocha, J.L., Kepinska, O., Schneider, P., Benner, J., Degano, G., Schneider, L., Golestani, N.,
804   2023. Multivariate Concavity Amplitude Index (MCAI) for characterizing Heschl's gyrus shape.
805   Neuroimage 272, 120052. https://doi.org/10.1016/j.neuroimage.2023.120052

806 Dalboni da Rocha, J.L., Schneider, P., Benner, J., Santoro, R., Atanasova, T., Van De Ville, D., Golestani,
807   N., 2020. TASH: Toolbox for the Automated Segmentation of Heschl's gyrus. Sci Rep 10, 3887.
808   https://doi.org/10.1038/s41598-020-60609-y

Destrieux, C., Fischl, B., Dale, A., Halgren, E., 2010. Automatic parcellation of human cortical gyri and sulci
using standard anatomical nomenclature. Neuroimage 53, 1–15.
https://doi.org/10.1016/j.neuroimage.2010.06.010

Dimanova, P., Borbás, R., Raschle, N.M., 2023. From mother to child: How intergenerational transfer is
reflected in similarity of corticolimbic brain structure and mental health. Dev Cogn Neurosci 64.
https://doi.org/10.1016/j.dcn.2023.101324

Eckert, M.A., Leonard, C.M., Molloy, E.A., Blumenthal, J., Zijdenbos, A., Giedd, J.N., 2002. The epigenesis
of planum temporale asymmetry in twins. Cereb Cortex 12, 749–755.

Eising, E., Mirza-Schreiber, N., de Zeeuw, E.L., Wang, C.A., Truong, D.T., Allegrini, A.G., Shapland, C.Y.,
Zhu, G., Wigg, K.G., Gerritse, M.L., Molz, B., Alagoz, G., Gialluisi, A., Abbondanza, F., Rimfeld, K.,
van Donkelaar, M., Liao, Z., Jansen, P.R., Andlauer, T.F.M., Bates, T.C., Bernard, M., Blokland, K.,
Bonte, M., Børglum, A.D., Bourgeron, T., Brandeis, D., Ceroni, F., Csepe, V., Dale, P.S., de Jong,
P.F., DeFries, J.C., Demonet, J.F., Demontis, D., Feng, Y., Gordon, S.D., Guger, S.L., Hayiou-
Thomas, M.E., Hernandez-Cabrera, J.A., Hottenga, J.J., Hulme, C., Kere, J., Kerr, E.N., Koomar, T.,
Landerl, K., Leonard, G.T., Lovett, M.W., Lyytinen, H., Martin, N.G., Martinelli, A., Maurer, U.,
Michaelson, J.J., Moll, K., Monaco, A.P., Morgan, A.T., Nothen, M.M., Pausova, Z., Pennell, C.E.,
Pennington, B.F., Price, K.M., Rajagopal, V.M., Ramus, F., Richer, L., Simpson, N.H., Smith, S.D.,
Snowling, M.J., Stein, J., Strug, L.J., Talcott, J.B., Tiemeier, H., van der Schroeff, M.P., Verhoef, E.,
Watkins, K.E., Wilkinson, M., Wright, M.J., Barr, C.L., Boomsma, D.I., Carreiras, M., Franken, M.C.J.,
Gruen, J.R., Luciano, M., Muller-Myhsok, B., Newbury, D.F., Olson, R.K., Paracchini, S., Paus, T.,
Plomin, R., Reilly, S., Schulte-Korne, G., Bruce Tomblin, J., van Bergen, E., Whitehouse, A.J.O.,
Willcutt, E.G., Pourcain, B.S., Francks, C., Fisher, S.E., 2022. Genome-wide analyses of individual
differences in quantitatively assessed reading- and language-related skills in up to 34,000 people.
Proc Natl Acad Sci U S A 119. https://doi.org/10.1073/PNAS.2202764119/-/DCSUPPLEMENTAL

Eyler, L.T., Chen, C.H., Panizzon, M.S., Fennema-Notestine, C., Neale, M.C., Jak, A., Jernigan, T.L., Fischl,
B., Franz, C.E., Lyons, M.J., Grant, M., Prom-Wormley, E., Seidman, L.J., Tsuang, M.T., Fiecas,
M.J.A., Dale, A.M., Kremen, W.S., 2012. A comparison of heritability maps of cortical surface area
and thickness and the influence of adjustment for whole brain measures: A magnetic resonance
imaging twin study. Twin Research and Human Genetics 15, 304–314.
https://doi.org/10.1017/thg.2012.3

Fehlbaum, L. V., Peters, L., Dimanova, P., Roell, M., Borbás, R., Ansari, D., Raschle, N.M., 2022. Mother-
child similarity in brain morphology: A comparison of structural characteristics of the brain's reading
network. Dev Cogn Neurosci 53, 101058. https://doi.org/10.1016/j.dcn.2022.101058

Feldman,   R.,   2023.   Father   contribution   to   human   resilience.   Dev   Psychopathol   1–18.
https://doi.org/10.1017/S0954579423000354

Fischl, B., Van Der Kouwe, A., Destrieux, C., Halgren, E., Ségonne, F., Salat, D.H., Busa, E., Seidman,
L.J., Goldstein, J., Kennedy, D., Caviness, V., Makris, N., Rosen, B., Dale, A.M., 2004. Automatically
Parcellating   the   Human   Cerebral   Cortex.   Cerebral   Cortex   14,   11–22.
https://doi.org/10.1093/cercor/bhg087

Francks, C., 2015. Exploring human brain lateralization with molecular genetics and genomics. Ann N Y
Acad Sci 1359, 1–13. https://doi.org/10.1111/nyas.12770

Gay, J., Olson, R.K., 2003. Genetic and environmental influences on individual differences in printed word
recognition.   J.   Experimental   Child   Psychology   84,   97–123.   https://doi.org/10.1016/S0022-
0965(02)00181-9

Geschwind, N., Levitsky, W., 1968. Human brain: left-right asymmetries in temporal speech region.
Science 161, 186–187. https://doi.org/10.1126/SCIENCE.161.3837.186

Gialluisi, A., Andlauer, T.F.M., Mirza-Schreiber, N., Moll, K., Becker, J., Hoffmann, P., Ludwig, K.U.,
Czamara, D., Pourcain, B.S., Honbolygó, F., Tóth, D., Csépe, V., Huguet, G., Chaix, Y., Iannuzzi, S.,
Demonet, J.F., Morris, A.P., Hulslander, J., Willcutt, E.G., DeFries, J.C., Olson, R.K., Smith, S.D.,
Pennington, B.F., Vaessen, A., Maurer, U., Lyytinen, H., Peyrard-Janvid, M., Leppänen, P.H.T.,
Brandeis, D., Bonte, M., Stein, J.F., Talcott, J.B., Fauchereau, F., Wilcke, A., Kirsten, H., Müller, B.,
Francks, C., Bourgeron, T., Monaco, A.P., Ramus, F., Landerl, K., Kere, J., Scerri, T.S., Paracchini,
S., Fisher, S.E., Schumacher, J., Nöthen, M.M., Müller-Myhsok, B., Schulte-Körne, G., 2021.
Genome-wide association study reveals new insights into the heritability and genetic correlates of
developmental dyslexia. Mol Psychiatry 26, 3004–3017. https://doi.org/10.1038/s41380-020-
00898-x

Golestani, N., Molko, N., Dehaene, S., LeBihan, D., Pallier, C., 2007a. Brain structure predicts the learning
of foreign speech sounds. Cerebral Cortex 17, 575–582.

Golestani, N., Molko, N., Dehaene, S., LeBihan, D., Pallier, C., 2007b. Brain structure predicts the learning
of foreign speech sounds. Cerebral Cortex 17, 575–582.

Golestani, N., Paus, T., Zatorre, R.J., 2002. Anatomical correlates of learning novel speech sounds. Neuron

35, 997–1010. https://doi.org/10.1016/S0896-6273(02)00862-0

Golestani, N., Price, C.J., Scott, S.K., 2011. Born with an ear for dialects? Structural plasticity in the expert

phonetician brain. Journal of Neuroscience 31, 4213–4220.

https://doi.org/10.1523/JNEUROSCI.3891-10.2011

Goos, L.M., Silverman, I., 2006. The inheritance of cognitive skills: does genomic imprinting play a role? J

Neurogenet 20, 19–40. https://doi.org/10.1080/01677060600685840

Grasby, K.L., Jahanshad, N., Painter, J.N., Colodro-Conde, L., Bralten, J., Hibar, D.P., Lind, P.A.,

Pizzagalli, F., Ching, C.R.K., McMahon, M.A.B., Shatokhina, N., Zsembik, L.C.P., Thomopoulos, S.I.,

Zhu, A.H., Strike, L.T., Agartz, I., Alhusaini, S., Almeida, M.A.A., Alnæs, D., Amlien, I.K., Andersson,

M., Ard, T., Armstrong, N.J., Ashley-Koch, A., Atkins, J.R., Bernard, M., Brouwer, R.M., Buimer,

E.E.L., Bülow, R., Bürger, C., Cannon, D.M., Chakravarty, M., Chen, Q., Cheung, J.W., Couvy-

Duchesne, B., Dale, A.M., Dalvie, S., de Araujo, T.K., de Zubicaray, G.I., de Zwarte, S.M.C., den

Braber, A., Doan, N.T., Dohm, K., Ehrlich, S., Engelbrecht, H.-R., Erk, S., Fan, C.C., Fedko, I.O.,

Foley, S.F., Ford, J.M., Fukunaga, M., Garrett, M.E., Ge, T., Giddaluru, S., Goldman, A.L., Green,

M.J., Groenewold, N.A., Grotegerd, D., Gurholt, T.P., Gutman, B.A., Hansell, N.K., Harris, M.A.,

Harrison, M.B., Haswell, C.C., Hauser, M., Herms, S., Heslenfeld, D.J., Ho, N.F., Hoehn, D.,

Hoffmann, P., Holleran, L., Hoogman, M., Hottenga, J.-J., Ikeda, M., Janowitz, D., Jansen, I.E., Jia,

T., Jockwitz, C., Kanai, R., Karama, S., Kasperaviciute, D., Kaufmann, T., Kelly, S., Kikuchi, M., Klein,

M., Knapp, M., Knodt, A.R., Krämer, B., Lam, M., Lancaster, T.M., Lee, P.H., Lett, T.A., Lewis, L.B.,

Lopes-Cendes, I., Luciano, M., Macciardi, F., Marquand, A.F., Mathias, S.R., Melzer, T.R., Milaneschi,

Y., Mirza-Schreiber, N., Moreira, J.C. V., Mühleisen, T.W., Müller-Myhsok, B., Najt, P., Nakahara, S.,

Nho, K., Olde Loohuis, L.M., Orfanos, D.P., Pearson, J.F., Pitcher, T.L., Pütz, B., Quidé, Y.,

Ragothaman, A., Rashid, F.M., Reay, W.R., Redlich, R., Reinbold, C.S., Repple, J., Richard, G.,

Riedel, B.C., Risacher, S.L., Rocha, C.S., Mota, N.R., Salminen, L., Saremi, A., Saykin, A.J., Schlag,

F., Schmaal, L., Schofield, P.R., Secolin, R., Shapland, C.Y., Shen, L., Shin, J., Shumskaya, E.,

Sønderby, I.E., Sprooten, E., Tansey, K.E., Teumer, A., Thalamuthu, A., Tordesillas-Gutiérrez, D.,

Turner, J.A., Uhlmann, A., Vallerga, C.L., van der Meer, D., van Donkelaar, M.M.J., van Eijk, L., van

Erp, T.G.M., van Haren, N.E.M., van Rooij, D., van Tol, M.-J., Veldink, J.H., Verhoef, E., Walton, E.,

Wang, M., Wang, Y., Wardlaw, J.M., Wen, W., Westlye, L.T., Whelan, C.D., Witt, S.H., Wittfeld, K.,

Wolf, C., Wolfers, T., Wu, J.Q., Yasuda, C.L., Zaremba, D., Zhang, Z., Zwiers, M.P., Artiges, E.,

Assareh, A.A., Ayesa-Arriola, R., Belger, A., Brandt, C.L., Brown, G.G., Cichon, S., Curran, J.E.,

Davies, G.E., Degenhardt, F., Dennis, M.F., Dietsche, B., Djurovic, S., Doherty, C.P., Espiritu, R.,

Garijo, D., Gil, Y., Gowland, P.A., Green, R.C., Häusler, A.N., Heindel, W., Ho, B.-C., Hoffmann, W.U., Holsboer, F., Homuth, G., Hosten, N., Jack, C.R., Jang, M., Jansen, A., Kimbrel, N.A., Kolskår, K., Koops, S., Krug, A., Lim, K.O., Luykx, J.J., Mathalon, D.H., Mather, K.A., Mattay, V.S., Matthews, S., Mayoral Van Son, J., McEwen, S.C., Melle, I., Morris, D.W., Mueller, B.A., Nauck, M., Nordvik, J.E., Nöthen, M.M., O'Leary, D.S., Opel, N., Martinot, M.-L.P., Pike, G.B., Preda, A., Quinlan, E.B., Rasser, P.E., Ratnakar, V., Reppermund, S., Steen, V.M., Tooney, P.A., Torres, F.R., Veltman, D.J., Voyvodic, J.T., Whelan, R., White, T., Yamamori, H., Adams, H.H.H., Bis, J.C., Debette, S., Decarli, C., Fornage, M., Gudnason, V., Hofer, E., Ikram, M.A., Launer, L., Longstreth, W.T., Lopez, O.L., Mazoyer, B., Mosley, T.H., Roshchupkin, G. V., Satizabal, C.L., Schmidt, R., Seshadri, S., Yang, Q., Alvim, M.K.M., Ames, D., Anderson, T.J., Andreassen, O.A., Arias-Vasquez, A., Bastin, M.E., Baune, B.T., Beckham, J.C., Blangero, J., Boomsma, D.I., Brodaty, H., Brunner, H.G., Buckner, R.L., Buitelaar, J.K., Bustillo, J.R., Cahn, W., Cairns, M.J., Calhoun, V., Carr, V.J., Caseras, X., Caspers, S., Cavalleri, G.L., Cendes, F., Corvin, A., Crespo-Facorro, B., Dalrymple-Alford, J.C., Dannlowski, U., de Geus, E.J.C., Deary, I.J., Delanty, N., Depondt, C., Desrivières, S., Donohoe, G., Espeseth, T., Fernández, G., Fisher, S.E., Flor, H., Forstner, A.J., Francks, C., Franke, B., Glahn, D.C., Gollub, R.L., Grabe, H.J., Gruber, O., Håberg, A.K., Hariri, A.R., Hartman, C.A., Hashimoto, R., Heinz, A., Henskens, F.A., Hillegers, M.H.J., Hoekstra, P.J., Holmes, A.J., Hong, L.E., Hopkins, W.D., Hulshoff Pol, H.E., Jernigan, T.L., Jönsson, E.G., Kahn, R.S., Kennedy, M.A., Kircher, T.T.J., Kochunov, P., Kwok, J.B.J., Le Hellard, S., Loughland, C.M., Martin, N.G., Martinot, J.-L., McDonald, C., McMahon, K.L., Meyer-Lindenberg, A., Michie, P.T., Morey, R.A., Mowry, B., Nyberg, L., Oosterlaan, J., Ophoff, R.A., Pantelis, C., Paus, T., Pausova, Z., Penninx, B.W.J.H., Polderman, T.J.C., Posthuma, D., Rietschel, M., Roffman, J.L., Rowland, L.M., Sachdev, P.S., Sämann, P.G., Schall, U., Schumann, G., Scott, R.J., Sim, K., Sisodiya, S.M., Smoller, J.W., Sommer, I.E., St Pourcain, B., Stein, D.J., Toga, A.W., Trollor, J.N., Van der Wee, N.J.A., van 't Ent, D., Völzke, H., Walter, H., Weber, B., Weinberger, D.R., Wright, M.J., Zhou, J., Stein, J.L., Thompson, P.M., Medland, S.E., 2020. The genetic architecture of the human cerebral cortex. Science (1979) 367. https://doi.org/10.1126/science.aay6690

Hager, R., Cheverud, J.M., Wolf, J.B., 2008. Maternal Effects as the Cause of Parent-of-Origin Effects That Mimic Genomic Imprinting. Genetics 178, 1755–1762. https://doi.org/10.1534/genetics.107.080697

Ho, T.C., Sanders, S.J., Gotlib, I.H., Hoeft, F., 2016. Intergenerational Neuroimaging of Human Brain Circuitry. Trends Neurosci 39, 644–648. https://doi.org/10.1016/j.tins.2016.08.008

Honeycutt, N.A., Musick, A., Barta, P.E., Pearlson, G.D., 2000. Measurement of the planum temporale (PT) on magnetic resonance imaging scans: Temporal PT alone and with parietal extension. Psychiatry Res Neuroimaging 98, 103–116. https://doi.org/10.1016/S0925-4927(00)00043-3

Hothorn, T., Hornik, K., Zeileis, A., 2006. party: A Laboratory for Recursive Part (y) itioning. R package version 1.3-13 URL http://CRAN. R-project. org.

Johnsrude, I.S., Penhune, V.B., Zatorre, R.J., 2000. Functional specificity in the right human auditory cortex for perceiving pitch direction 155–163.

Kepinska, O., Dalboni da Rocha, J.L., Tuerk, C., Hervais-Adelman, A., Bouhali, F., Green, D., Price, C.J., Golestani, N., 2023. Auditory cortex anatomy reflects multilingual phonological experience. Elife 12, 2023.06.16.545298. https://doi.org/10.7554/elife.90269.1

Kong, X.Z., Mathias, S.R., Guadalupe, T., Abé, C., Agartz, I., Akudjedu, T.N., Aleman, A., Alhusaini, S., Allen, N.B., Ames, D., Andreassen, O.A., Vasquez, A.A., Armstrong, N.J., Bergo, F., Bastin, M.E., Batalla, A., Bauer, J., Baune, B.T., Baur-Streubel, R., Biederman, J., Blaine, S.K., Boedhoe, P., Bøen, E., Bose, A., Bralten, J., Brandeis, D., Brem, S., Brodaty, H., Yüksel, D., Brooks, S.J., Buitelaar, J., Bürger, C., Bülow, R., Calhoun, V., Calvo, A., Canales-Rodríguez, E.J., Canive, J.M., Cannon, D.M., Caparelli, E.C., Castellanos, F.X., Cavalleri, G.L., Cendes, F., Chaim-Avancini, T.M., Chantiluke, K., Chen, Q.L., Chen, X., Cheng, Y., Christakou, A., Clark, V.P., Coghill, D., Connolly, C.G., Conzelmann, A., Córdova-Palomera, A., Cousijn, J., Crow, T., Cubillo, A., Dale, A., Dannlowski, U., De Bruttopilo, S.A., De Zeeuw, P., Deary, I.J., Delanty, N., Demeter, D. V., Di Martino, A., Dickie, E.W., Dietsche, B., Doan, N.T., Doherty, C.P., Doyle, A., Durston, S., Earl, E., Ehrlich, S., Ekman, C.J., Elvsåshagen, T., Epstein, J.N., Fair, D.A., Faraone, S. V., Fernández, G., Filho, G.B., Förster, K., Fouche, J.P., Foxe, J.J., Frodl, T., Fuentes-Claramonte, P., Fullerton, J., Garavan, H., Garcia, D.D.S., Gotlib, I.H., Goudriaan, A.E., Grabe, H.J., Groenewold, N.A., Grotegerd, D., Gruber, O., Gurholt, T., Haavik, J., Hahn, T., Hansell, N.K., Harris, M.A., Hartman, C.A., Hernández, M.D.C.V., Heslenfeld, D., Hester, R., Hibar, D.P., Ho, B.C., Ho, T.C., Hoekstra, P.J., Van Holst, R.J., Hoogman, M., Høvik, M.F., Howells, F.M., Hugdahl, K., Huyser, C., Ingvar, M., Irwin, L., Ishikawa, A., James, A., Jahanshad, N., Jernigan, T.L., Jönsson, E.G., Kähler, C., Kaleda, V., Kelly, C., Kerich, M., Keshavan, M.S., Khadka, S., Kircher, T., Kohls, G., Konrad, K., Korucuoglu, O., Krämer, B., Krug, A., Kwon, J.S., Lambregts-Rommelse, N., Landén, M., Lázaro, L., Lebedeva, I., Lenroot, R., Lesch, K.P., Li, Q., Lim, K.O., Liu, J., Lochner, C., London, E.D., Lonning, V., Lorenzetti, V., Luciano, M., Luijten, M., Lundervold, A.J., MacKey, S., MacMaster, F.P., Maingault, S., Malpas, C.B., Malt, U.F., Mataix-Cols, D., Martin-Santos, R., Mayer, A.R., McCarthy, H., Mitchell, P.B., Mueller, B.A., Maniega, S.M., Mazoyer, B., McDonald, C., McLellan, Q., McMahon, K.L., McPhilemy, G., Momenan, R., Morales, A.M., Narayanaswamy, J.C., Moreira, J.C.V., Nerland, S., Nestor, L., Newman, E., Nigg, J.T., Nordvik, J.E., Novotny, S., Weiss, E.O., O'Gorman, R.L., Oosterlaan, J., Oranje, B., Orr, C., Overs, B., Pauli, P., Paulus, M., Plessen, K.J., Von Polier, G.G., Pomarol-Clotet, E., Portella, M.J., Qiu, J., Radua, J., Ramos-Quiroga,

J.A., Reddy, Y.C.J., Reif, A., Roberts, G., Rosa, P., Rubia, K., Sacchet, M.D., Sachdev, P.S.,
Salvador, R., Schmaal, L., Schulte-Rüther, M., Schweren, L., Seidman, L., Seitz, J., Serpa, M.H.,
Shaw, P., Shumskaya, E., Silk, T.J., Simmons, A.N., Simulionyte, E., Sinha, R., Sjoerds, Z., Smelror,
R.E., Soliva, J.C., Solowij, N., Souza-Duran, F.L., Sponheim, S.R., Stein, D.J., Stein, E.A., Stevens,
M., Strike, L.T., Sudre, G., Sui, J., Tamm, L., Temmingh, H.S., Thoma, R.J., Tomyshev, A., Tronchin,
G., Turner, J., Uhlmann, A., Van Erp, T.G.M., Van Den Heuvel, O.A., Van Der Meer, D., Van Eijk, L.,
Vance, A., Veer, I.M., Veltman, D.J., Venkatasubramanian, G., Vilarroya, O., Vives-Gilabert, Y.,
Voineskos, A.N., Völzke, H., Vuletic, D., Walitza, S., Walter, H., Walton, E., Wardlaw, J.M., Wen, W.,
Westlye, L.T., Whelan, C.D., White, T., Wiers, R.W., Wright, M.J., Wittfeld, K., Yang, T.T., Yasuda,
C.L., Yoncheva, Y., Yücel, M., Yun, J.Y., Zanetti, M.V., Zhen, Z., Zhu, X.X., Ziegler, G.C., Zierhut, K.,
De Zubicaray, G.I., Zwiers, M., Glahn, D.C., Franke, B., Crivello, F., Tzourio-Mazoyer, N., Fisher, S.E.,
Thompson, P.M., Francks, C., Farde, L., Flyckt, L., Engberg, G., Erhardt, S., Fatouros-Bergman, H.,
Cervenka, S., Schwieler, L., Piehl, F., Collste, K., Victorsson, P., Malmqvist, A., Hedberg, M., Orhan,
F., 2018. Mapping cortical brain asymmetry in 17,141 healthy individuals worldwide via the ENIGMA
consortium. Proc Natl Acad Sci U S A 115, E5154–E5163.
https://doi.org/10.1073/pnas.1718418115

Kuhl, U., Neef, N.E., Kraft, I., Schaadt, G., Dörr, L., Brauer, J., Czepezauer, I., Müller, B., Wilcke, A., Kirsten,
H., Emmrich, F., Boltze, J., Friederici, A.D., Skeide, M.A., 2020. The emergence of dyslexia in the
developing brain. Neuroimage 211. https://doi.org/10.1016/j.neuroimage.2020.116633

Lawson, H.A., Cheverud, J.M., Wolf, J.B., 2013. Genomic imprinting and parent-of-origin effects on
complex traits. Nat Rev Genet 14, 609–617. https://doi.org/10.1038/nrg3543

Leonard, C., Eckert, M., Given, B., Virginia, B., Eden, G., 2006. Individual differences in anatomy predict
reading and oral language impairments in children. Brain 129, 3329–3342.

Leonard, C.M., Eckert, M.A., Lombardino, L.J., Oakland, T., Kranzler, J., Mohr, C.M., King, W.M.,
Freeman, A., 2001. Anatomical risk factors for phonological dyslexia. Cerebral Cortex 11, 148–157.
https://doi.org/10.1093/cercor/11.2.148

Linkersdörfer, J., Lonnemann, J., Lindberg, S., Hasselhorn, M., Fiebach, C.J., 2012. Grey matter
alterations co-localize with functional abnormalities in developmental dyslexia: An ALE meta-analysis.
PLoS One 7. https://doi.org/10.1371/journal.pone.0043122

Ma, Y., Koyama, M.S., Milham, M.P., Castellanos, F.X., Quinn, B.T., Pardoe, H., Wang, X., Kuzniecky, R.,
Devinsky, O., Thesen, T., Blackmon, K., 2015. Cortical thickness abnormalities associated with

dyslexia, independent of remediation status. Neuroimage Clin 7, 177–186.
https://doi.org/10.1016/j.nicl.2014.11.005

Marie, D., Jobard, G., Crivello, F., Perchey, G., Petit, L., Mellet, E., Joliot, M., Zago, L., Mazoyer, B.,
Tzourio-Mazoyer, N., 2015. Descriptive anatomy of Heschl's gyri in 430 healthy volunteers, including
198 left-handers. Brain Struct Funct 220, 729–743. https://doi.org/10.1007/s00429-013-0680-x

Meyer, M., Liem, F., Hirsiger, S., Jäncke, L., Hänggi, J., 2014. Cortical surface area and cortical thickness
demonstrate differential structural asymmetry in auditory-related areas of the human cortex. Cerebral
Cortex 24, 2541–2552. https://doi.org/10.1093/cercor/bht094

Mozaffari, S. V., DeCara, J.M., Shah, S.J., Sidore, C., Fiorillo, E., Cucca, F., Lang, R.M., Nicolae, D.L.,
Ober, C., 2019. Parent-of-origin effects on quantitative phenotypes in a large Hutterite pedigree.
Commun Biol 2, 1–9. https://doi.org/10.1038/s42003-018-0267-4

Niemi, L.T., Suvisaari, J.M., Haukka, J.K., Lönnqvist, J.K., 2005. Childhood predictors of future psychiatric
morbidity in offspring of mothers with psychotic disorder: Results from the Helsinki high-risk study.
British Journal of Psychiatry 186, 108–114. https://doi.org/10.1192/bjp.186.2.108

Norton, E.S., Black, J.M., Stanley, L.M., Tanaka, H., Gabrieli, J.D.E., Sawyer, C., Hoeft, F., 2014.
Functional neuroanatomical evidence for the double-deficit hypothesis of developmental dyslexia.
Neuropsychologia 61, 235–246. https://doi.org/10.1016/j.neuropsychologia.2014.06.015

Penhune, V.B., Zatorre, R.J., MacDonald, J.D., Evans, A.C., 1996. Interhemispheric anatomical differences
in human primary auditory cortex: Probabilistic mapping and volume measurement from magnetic
resonance scans. Cerebral Cortex 6, 661–672. https://doi.org/10.1093/cercor/6.5.661

R Development Core Team, 2015. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing. https://doi.org/10.1007/978-3-540-74686-7

Ramoser, C., Fischer, A., Caspers, J., Schiller, N.O., Golestani, N., Kepinska, O., (*under review*) Language
Aptitude and the Morphology and Shape of the Auditory Cortex Transverse Temporal Gyri.

Schneider, P., Andermann, M., Wengenroth, M., Goebel, R., Flor, H., Rupp, A., Diesch, E., 2009. Reduced
volume of Heschl's gyrus in tinnitus. Neuroimage 45, 927–939.
https://doi.org/10.1016/j.neuroimage.2008.12.045

Schneider, P., Scherg, M., Dosch, H.G., Specht, H.J., Gutschalk, A., Rupp, A., 2002. Morphology of
Heschl's gyrus reflects enhanced activation in the auditory cortex of musicians. Nat Neurosci 5, 688–
694. https://doi.org/10.1038/nn871

Serrallach, B., Groß, C., Bernhofs, V., Engelmann, D., Benner, J., Gündert, N., Blatow, M., Wengenroth, M., Seitz, A., Brunner, M., Seither, S., Parncutt, R., Schneider, P., Seither-Preisler, A., Gündert, N., Blatow, M., Wengenroth, M., Seitz, A., Brunner, M., Seither, S., Parncutt, R., Schneider, P., Seither-Preisler, A., 2016. Neural biomarkers for dyslexia, ADHD, and ADD in the auditory cortex of children. Front Neurosci 10, 1–23. https://doi.org/10.3389/fnins.2016.00324

Serrallach, B.L., Groß, C., Christiner, M., Wildermuth, S., Schneider, P., 2022. Neuromorphological and Neurofunctional Correlates of ADHD and ADD in the Auditory Cortex of Adults. Front Neurosci 16, 850529. https://doi.org/10.3389/fnins.2022.850529

Strobl, C., Boulesteix, A.L., Kneib, T., Augustin, T., Zeileis, A., 2008. Conditional variable importance for random forests. BMC Bioinformatics 9. https://doi.org/10.1186/1471-2105-9-307

Sullivan, P.F., Kendler, K.S., Neale, M.C., 2003. Schizophrenia as a Complex Trait: Evidence from a Meta-analysis of Twin Studies. Arch Gen Psychiatry 60, 1187–1192. https://doi.org/10.1001/archpsyc.60.12.1187

Sutherland, M.E., Zatorre, R.J., Watkins, K.E., Hervé, P.Y., Leonard, G., Pike, B.G., Witton, C., Paus, T., 2012. Anatomical correlates of dynamic auditory processing: Relationship to literacy during early adolescence. Neuroimage 60, 1287–1295. https://doi.org/10.1016/j.neuroimage.2012.01.051

Takagi, Y., Okada, N., Ando, S., Yahata, N., Morita, K., Koshiyama, D., Kawakami, S., Sawada, K., Koike, S., Endo, K., Yamasaki, S., Nishida, A., Kasai, K., Tanaka, S.C., 2021. Intergenerational transmission of the patterns of functional and structural brain networks. iScience 24, 102708. https://doi.org/10.1016/j.isci.2021.102708

Takahashi, T., Sasabayashi, D., Takayanagi, Y., Furuichi, A., Kido, M., 2021. Increased Heschl's Gyrus Duplication in Schizophrenia Spectrum Disorders: A Cross-Sectional MRI Study.

Takahashi, T., Sasabayashi, D., Takayanagi, Y., Furuichi, A., Kobayashi, H., Noguchi, K., Suzuki, M., 2022. Different Heschl's Gyrus Duplication Patterns in Deficit and Non-deficit Subtypes of Schizophrenia. Front Psychiatry 13, 867461. https://doi.org/10.3389/fpsyt.2022.867461

Torgesen, J.K., Wagner, R.K., Rashotte, C.A., 1999. Test of word reading efficiency (TOWRE). Austin, TX: Pro-Ed.

Turker, S., Reiterer, S.M., Schneider, P., Seither-Preisler, A., 2019. Auditory cortex morphology predicts language learning potential in children and teenagers. Front Neurosci 13, 824. https://doi.org/10.3389/FNINS.2019.00824/BIBTEX

Turker, S., Reiterer, S.M., Seither-Preisler, A., Schneider, P., 2017. "When Music Speaks": Auditory Cortex
Morphology as a Neuroanatomical Marker of Language Aptitude and Musicality. Front Psychol 8,
2096. https://doi.org/10.3389/fpsyg.2017.02096

Turker, S., Seither-Preisler, A., Reiterer, S.M., 2021. Examining Individual Differences in Language
Learning: A Neurocognitive Model of Language Aptitude. Neurobiology of Language 1–27.
https://doi.org/10.1162/nol_a_00042

van Bergen, E., van der Leij, A., de Jong, P.F., 2014. The intergenerational multiple deficit model and the
case of dyslexia. Front Hum Neurosci 8. https://doi.org/10.3389/fnhum.2014.00346

von Economo, C., Horn, L., 1930. Über Windungsrelief, Maße und Rindenarchitektonik der
Supratemporalfläche, ihre individuellen und ihre Seitenunterschiede. Zeitschrift für die gesamte
Neurologie und Psychiatrie 130, 678–757. https://doi.org/10.1007/BF02865945

Wengenroth, M., Blatow, M., Heinecke, A., Reinhardt, J., Stippich, C., Hofmann, E., Schneider, P., 2014.
Increased volume and function of right auditory cortex as a marker for absolute pitch. Cereb Cortex
24, 1127–1137. https://doi.org/10.1093/cercor/bhs391

Wolf, M., Bowers, P.G., 1999. The double-deficit hypothesis for the developmental dyslexias. J Educ
Psychol 91, 415–438. https://doi.org/10.1037/0022-0663.91.3.415

Wolf, M., Denckla, M., 2005. Rapid Automatized Naming and Rapid Alternating Stimulus Tests: Examiner's
Manual. PRO-ED, Austin, Texas.

Wong, P.C.M., Warrier, C.M., Penhune, V.B., Roy, A.K., Sadehh, A., Parrish, T.B., Zatorre, R.J., 2008.
Volume of left Heschl's gyrus and linguistic pitch learning. Cerebral Cortex 18, 828–836.
https://doi.org/10.1093/cercor/bhm115

Woodcock, R.W., 1998. Woodcock Reading Mastery Tests—Revised/NU. American Guidance Service,
Inc., Circle Pines, MN.

Yamagata, B., Murayama, K., Black, J.M., Hancock, R., Mimura, M., Yang, T.T., Reiss, A.L., Hoeft, F.,
2016. Female-specific intergenerational transmission patterns of the human corticolimbic circuitry.
Journal of Neuroscience 36, 1254–1260. https://doi.org/10.1523/JNEUROSCI.4974-14.2016

Zaretskaya, N., Fischl, B., Reuter, M., Renvall, V., Polimeni, J.R., 2018. Advantages of cortical surface
reconstruction using submillimeter 7 T MEMPRAGE. Neuroimage 165, 11–26.
https://doi.org/10.1016/j.neuroimage.2017.09.060

1088

7 Supplementary Materials

7.1 Participants

Family ID	Family member	Age (years)	Sex	Handedness
1	Child	5.47	M	R
	Child	13.52	F	R
	Child	11.73	M	R
	Father	45.89	M	R
	Mother*	43.09	F	R
2	Child	8.14	F	R
	Child	14.43	F	R
	Child	11.31	F	R
	Father	54.40	M	L
3	Child	8.80	M	R
	Child	8.16	M	R
	Father	52.53	M	R
	Mother	39.44	F	R
4	Child	8.49	M	R
	Child	10.38	F	R
	Father	41.36	M	Both
	Mother*	40.41	F	R
5	Child	8.05	F	R
	Child	9.03	F	Both
	Mother	35.22	F	R
6	Child	7.95	F	R
	Child	9.91	F	R
	Child	13.26	M	R
	Father	39.51	M	R
	Mother*	41.70	F	R
7	Child	9.76	F	R
	Father	50.12	M	R
	Mother*	45.62	F	R
8	Child	8.78	F	R
	Father	39.59	M	R
	Mother	38.09	F	L
9	Child	9.08	M	L
	Child	7.06	M	R
	Mother*	41.24	F	R
10	Child	7.62	M	R
	Father	39.90	M	R
	Mother*	35.69	F	L
11	Child	7.99	M	L
	Child	9.05	F	R
	Mother	38.58	F	R
12	Child	7.88	F	R
	Child	10.95	F	R
	Father	45.99	M	R
	Mother	43.91	F	R
13	Child	8.41	F	R
	Child	5.81	M	R
	Father	35.90	M	R
14	Child	8.11	M	R
	Child	8.00	F	R
	Father	48.66	M	L
	Mother*	41.40	F	R
15	Child	8.65	F	L
	Child	5.76	F	L
	Father	42.23	M	R
	Mother	42.81	F	L
16	Child	8.65	M	R
	Child	10.55	M	R
	Father	42.07	M	R

	Mother*	43.96	F	R
17	Child	7.83	M	R
	Child	7.72	M	R
	Child	5.52	M	R
	Father	39.18	M	Both
	Mother*	41.70	F	R
18	Child	8.77	F	R
	Child	12.19	M	R
	Father	41.23	M	R
	Mother*	48.82	F	R
19	Child	7.51	M	R
	Child	6.99	M	R
	Father	45.43	M	R
	Mother	41.13	F	R
20	Child	8.39	F	R
	Child	6.02	F	L
	Father	36.38	M	R
	Mother	33.95	F	R
21	Child	8.28	M	R
	Father	45.65	M	R
	Mother*	38.21	F	R
22	Child	7.98	M	R
	Father	40.29	M	R
	Mother*	41.85	F	R
23	Child	8.70	F	R
	Child	10.46	F	R
	Mother*	43.74	F	R
24	Child	8.81	M	L
	Father	49.44	M	L
	Mother*	47.15	F	R
25	Child	8.12	F	R
	Child	10.20	F	R
	Child	5.12	M	R
	Father	46.34	M	R
	Mother	43.70	F	R
26	Child	8.62	M	R
	Father	47.44	M	R
27	Child	8.18	F	R
	Father	54.33	M	R
	Mother	46.05	F	R
28	Child	8.10	M	R
	Child	12.60	F	R
	Father	48.27	M	R
	Mother*	41.44	F	R
29	Child	8.72	M	L
	Child	12.28	F	R
	Father	51.62	M	R
	Mother*	49.28	F	R
30	Child	8.08	F	L
	Child	5.92	M	N/A
	Father	44.07	M	N/A
	Mother	41.13	F	N/A
31	Child	8.40	F	R
	Child	4.88	F	N/A
	Mother*	38.11	F	L
32	Child	8.18	M	R
	Mother*	44.87	F	R
33	Child	8.88	M	R
	Child	5.70	F	R
	Child	8.64	M	R
	Father	35.04	M	R
	Mother*	34.19	F	R
34	Child	7.77	F	R
	Mother*	40.85	F	R
35	Child	7.98	M	R
	Father	42.23	M	R

	Child	8.47	M	R
	Child	5.72	M	R
	Child	14.19	M	R
36	Child	14.19	M	R
	Child	9.40	M	L
	Father	32.63	M	R
	Mother	30.87	F	R
	Child	8.16	F	R
37	Child	8.16	F	R
	Father	44.49	M	R
	Mother*	38.97	F	R

Table S 1. Participant characteristics including family ID, age, sex and handedness. (*) denotes mothers who declared to be homemakers; no father declared to be homemaker.

7.2 TTG and reading measures (uncorrected)

Figure S 1. Relationship between TTG anatomy and the performance on five reading tests (T-SWE, T-PDE, RN-LTR, WRMT-WID and WRMT-WA) across the whole sample obtained from comparing models with demographic variables to models including neuroimaging data. The intensity of the color represents increase in R^2 values between a model with demographic variables only and a model additionally including mothers' or fathers' corresponding test scores; p-values were obtained from likelihood ratio tests used to compare the models and are not corrected for multiple comparisons (compare with Figure 3B, where FDR-corrected p-values are presented).

 ## 7.3 Intergenerational effects on reading ability

1101

Figure S 2. Children's reading skills as a function of fathers' cortical volume variables (of the left HG, left PT and left HG:PT ratio).

Similar to surface area results, fathers' PT volume alone did not significantly predict children's T-PDE

scores (β = 0.16, t = 0.97, p = .34), but fathers' HG:PT volumes ratio did significantly predict children's

reading (β = -0.44, t = -3.26, p = .005). Permutation analysis confirmed familial specificity of this effect

(p = .0006). Directly comparing the models predicting children's T-PDE with either father's HG volume, or

fathers' HG:PT volumes ratio as independent variables showed that the model with HG:PT ratio offered a

better fit to the data with $\Delta R^2_{Marg.}$ of 2.88% (ΔAIC = -1.88). The effect was not sex specific: a nested model

comparison with a likelihood ratio test did not show a better fit for a model additionally including an

interaction term between fathers' HG:PT volumes ratio and child's sex (X^2 = 1.85, p = 0.17).

ANALYSIS	DEPENDANT MEASURE(S)	INDEPENDENT MEASURE(S)	COVARIATES OF NO-INTEREST	QUESTION	RESULTS
1. Intergenerational transmission of reading skills (Section 3.2)	children's performance on five reading and reading-related tests (T-SWE, T-PDE, RN-LTR, WRMT-WID and WRMT-WA)	a. mothers' performance on the respective reading tests b. fathers' performance on the respective reading tests	• age • sex • SES	Is children's reading related to parents' reading?	Yes (Figure 3A): a. children's performance on T-PDE, WRMT-WID and WRMT-WA was positively related to mothers' performance on the respective reading tests b. children's performance on T-PDE, WRMT-WID was positively related to fathers' performance on the respective reading tests
2. TTG(s) and reading measures (Section 3.3)	whole sample's performance on five reading and reading-related tests (T-SWE, T-PDE, RN-LTR, WRMT-WID and WRMT-WA)	neuroanatomical measures describing the structure of the TTG(s): a. left and right HG (volume, area, thickness, shape) b. left and right all TTG(s) (volume, area, thickness, shape)	• age [linear and quadratic term] • sex • SES • handedness • total intracranial volume	Are reading skills related to the structure of the TTG(s)?	Yes (Figure 3B): T-PDE scores were positively and significantly related to the left all TTG(s) volume ($\Delta R^2_{Marg.} = 8.15$, $p_{FDR} = .02$) and surface area ($\Delta R^2_{Marg.} = 9.26$, $p_{FDR} = .02$)
2a. Follow-up analysis	whole sample's reading measures which were significantly related to 'all TTG(s)' volume and area in Analysis 2 above (T-PDE)	volume and area of the left PT:	• age [linear and quadratic term] • sex • SES • handedness • total intracranial volume	Are reading skills related to the structure of the TTG(s) specifically, or rather to PT anatomy (which contain the additional TTG(s))?	Yes: 'all TTG(s)' measures were better at explaining reading skills than PT measures, with $\Delta R^2_{Marg.}$ of 4.39% (ΔAIC = -7.78) and 3.55% (ΔAIC = -6.61) for volume and area respectively
3. Familial similarity in the structure of the TTG(s) (Section 3.4)	neuroanatomical measures describing the structure of the TTG(s): a. left and right HG (volume, area, thickness, shape) b. left and right all TTG(s) (volume, area, thickness, shape)	family index	• age [linear and quadratic term] • sex • SES • handedness • total intracranial volume	Which features of the TTG(s) anatomy are similar among family members?	a. volume, area, and thickness of right HG b. shape of left and right HG c. lateralization of volume and surface of all TTG(s) d. thickness of the left and right all TTG(s) e. lateralization of HG area
3a. Follow-up analysis on Result (a) in Analysis 3 (Section 3.4.1)	children's volume, area, and thickness of the 1st right TTG (HG) [normalized for their corresponding whole hemisphere measures]	a. mothers' volume, area, and thickness of the right HG [normalized for their corresponding whole hemisphere measures] b. fathers' volume, area, and thickness of the right HG [normalized for their corresponding whole hemisphere measures]	children's: • age [linear and quadratic term] • sex • SES • handedness	Is there intergenerational transmission of volume, area, and thickness of right HG?	Yes (Table 2, Figure 4): a. children's volume, area, and thickness of right HG was positively and significantly related to mothers' right HG volume, area, and thickness respectively (volume: β = 0.38, t = 2.84, p = .009; area: β = 0.30, t = 2.33, p = .03; thickness: β = 0.38, t = 2.87, p = .008), b. children's right HG thickness was positively and significantly related to fathers' thickness (β = 0.28, t = 2.05, p = .05), but not to other measures
3b. Follow-up analysis on Result (b) in Analysis 3 (Section 3.4.2)	children's left and right HG shape	a. mothers' left and right HG shape b. fathers' left and right HG shape	children's: • age [linear and quadratic term] • sex • SES • handedness	Is there intergenerational transmission of left and right HG shape?	Yes (Table 3, Figure 5): a. children's left HG shape was only significantly and positively related to mothers' left HG shape (β = 0.31, t = 2.12, p = .04) b. relationships between children's and fathers' HG shape were not significant[2]
3c. Follow-up analysis on Result (c) in Analysis 3 (Section 3.4.3)	children's lateralization index (LI) of volume and area of all TTG(s)	a. mothers' LI of volume and area of all TTG(s) b. fathers' LI of volume and area of all TTG(s)	children's: • age [linear and quadratic term] • sex	Is there intergenerational transmission of lateralization of all TTG(s)?	Yes (Table 4, Figure 6): a. relationships between children's and mothers' all TTG LI were not significant

---

[2] Non-significant results in gray; significant in black.

ANALYSIS		DEPENDANT MEASURE(S)	INDEPENDENT MEASURE(S)	COVARIATES OF NO-INTEREST	QUESTION	RESULTS
				• SES • handedness		b. LI of surface area of children's all TTG(s) was positively and significantly related to fathers' all TTG(s) LI of the surface area (β = 0.42, t = 3.00, p = .005)
3d.	Follow-up analysis on Result (d) in Analysis 3 (Section 3.4.4)	children's thickness of the left and right all TTG(s)	a. mothers' thickness of the left and right all TTG(s) b. fathers' thickness of the left and right all TTG(s)	children's: • age [linear and quadratic term] • sex • SES • handedness	Is there intergenerational transmission of thickness of the left and right all TTG(s)?	No: a. mothers' measures were not significantly related to those of the children b. fathers' measures were not significantly related to those of the children
3e.	Follow-up analysis on Result (e) in Analysis 3 (Section 3.4.4)	children's lateralization of HG area	a. mothers' lateralization of HG area b. fathers' lateralization of HG area	children's: • age [linear and quadratic term] • sex • SES • handedness	Is there intergenerational transmission of lateralization of HG area?	No: a. mothers' measures were not significantly related to those of the children b. fathers' measures were not significantly related to those of the children

4.	Intergenerational effects on reading ability (Section 3.5)	children's performance on T-PDE	a. mothers' neuroanatomical measures describing the structure of the TTG(s):
			  - left and right HG (volume, area, thickness [normalized for their corresponding whole hemisphere measures], shape)
			  - left and right all TTG(s) (volume, area, thickness [normalized for their corresponding whole hemisphere measures], shape)
			b. fathers' neuroanatomical measures describing the structure of the TTG(s):
			  - left and right HG (volume, area, thickness [normalized for their corresponding whole hemisphere measures], shape)
			  - left and right all TTG(s) (volume, area, thickness [normalized for their corresponding whole hemisphere measures], shape)
				-	Is children's reading related to parents' neuroanatomy?	Yes (Figure 7A):
						a. model with mothers' variables did not point to any features of their TTG(s) to be relevant predictors of children's reading ability
						b. fathers' volume and area of the left HG and HG lateralization indices of volume and area had conditional permutation importance significantly different from zero

4a.	Follow-up analysis on Results in Analysis 4	children's performance on T-PDE	a. fathers' left HG volume and area [normalized for their corresponding whole hemisphere measures] b. fathers' lateralization of volume and area of HG	children's: • age • sex • SES	Do the above effects survive in linear mixed effects models? What is their direction?	a. fathers' left HG volume and area were both negatively related to children's reading (volume: β = -0.40, t = -2.91, p = .008; area: β = -0.46, t = -3.45, p = .002) (Figure 7B) b. fathers' lateralization of HG (both for volume and area) was not significantly related to the children's reading
4b.	Follow-up analysis on Results in Analysis 4a	fathers' left HG volume and area	a. fathers' left all TTG(s) shape b. fathers' left PT volume and area	-	Is fathers' small left HG related to more additional TTGs, or with a larger PT?	a. No: fathers' small left HG is not related to more additional TTGs b. Yes: fathers' small left HG area is related to larger PT area (β = 0.18, t = 2.036, p = .05), but volume of fathers' HG was not related to PT volume (Figure 7B)
4c.	Follow-up analysis on Results in Analysis 4b	children's performance on T-PDE	a. fathers' left HG area and volume b. fathers' left PT area and volume c. fathers' left HG:PT ratio of area and volume	children's: • age • sex • SES	Which characteristics of fathers' auditory cortex (HG, PT or HG:PT ratio) explain most variance in children's reading?	fathers' left HG:PT ratio of area was the best predictor of children's reading (Figure 7B)